Predictive Modeling of *Pseudomonas syringae* Virulence on Bean using Gradient

Boosted Decision Trees

Renan N.D. Almeida[1], Michael Greenberg[1], Cedoljub Bundalovic-Torma[1], Alexandre Martel[1],

Pauline W. Wang[1,2], Maggie Middleton[1], Syama Chatterton[3], Darrell Desveaux [1], and David S.

Guttman[1,2,*]

[1]Department of Cell & Systems Biology, University of Toronto, 25 Willcocks St., Toronto,

Ontario, Canada

[2]Centre for the Analysis of Genome Evolution & Function, University of Toronto, Toronto,

Ontario, Canada

[3]Agriculture and Agri-Food Canada, Lethbridge Research and Development Centre, Lethbridge,

Alberta, Canada

*To whom correspondence should be addressed. Email: david.guttman@utoronto.ca

Running title: Predicting *P. syringae* virulence on bean using a machine learning approach

Keywords: *Pseudomonas syringae*, *Phaseolus vulgaris*, bean, machine learning, random forest,

gradient boosted decision trees, host specificity, virulence,

ABSTRACT

Pseudomonas syringae is a genetically diverse bacterial species complex responsible for numerous agronomically important crop diseases. Individual isolates of *P. syringae* are typically assigned pathovar names based on their host of isolation and the associated disease symptoms, and these pathovar designations are often assumed to reflect host specificity. Unfortunately, this assumption has rarely been rigorously tested, which poses a challenge when trying to identify genetic factors associated with host specificity. Here we develop a rapid seed infection assays to measure the virulence of 121 diverse *P. syringae* isolates on common bean (*Phaseolus vulgaris*). This collection includes *P. syringae* phylogroup 2 bean isolates (pathovar *syringae*) that cause bacterial spot disease and *P. syringae* phylogroup 3 bean isolates (pathovar *phaseolicola*) that cause the much more serious halo blight disease. We find that phylogroup 2 strains generally show lower levels of host specificity on bean, with the average level of virulence for all strains in this phylogroup (irrespective of host of isolation) being higher than the average level for all other *P. syringae* strains. We then use gradient boosted decision trees to model the *P. syringae* virulence weights using whole genome kmers, type III secreted effector kmers, and the presence / absence of type III effectors and phytotoxins. Our machine learning model performed best using whole genome data, and we were able to predict bean virulence with high accuracy (mean absolute error as low as 0.05). Finally, we functionally validated the model by predicting virulence for 16 strains and found that 15 (94%) of the strains had virulence levels within the bounds of estimated predictions given the calculated RMSE values (±0.20). This study further illustrates that *P. syringae* phylogroup 2 strains may have evolved a different lifestyle than other *P. syringae* strains and demonstrates the power of machine learning for predicting host specific adaptation.

INTRODUCTION

Pseudomonas syringae is a genetically diverse Gammaproteobacterial species complex

responsible for numerous agronomically important crop diseases (1-4). Strains in the *P.*

syringae species complex are frequently categorized into pathovars depending on pathogenic

characteristics such as host of isolation and disease symptoms (5, 6). Strains have also been

subdivided into phylogenetic groups (i.e., phylogroups) based on multilocus sequence typing or

genomic analysis (1, 7-10). Currently, there are 13 recognized phylogroups (7), of which seven

have been termed primary phylogroups based on their genetic relatedness and the near

universal presence of the canonical *P. syringae* type III secretion system (discussed below) (1,

9).

Phylogenetic analyses of *P. syringae* isolates suggests that adaptation to specific hosts has

evolved multiple times in the evolutionary history of the species complex. For example, cherry

and plum pathogens are found in clades distributed in phylogroups 1, 2, and 3 (11, 12), hazelnut

pathogens are distributed among two distinct clades in phylogroups 1 and 2 (13), while

pathogens of common bean (including snap, green, kidney, and French bean) are found in

phylogroups 2 and 3 (9). Common bean pathogens from *P. syringae* phylogroup 3 are generally

classified as pathovar *phaseolicola* and are responsible for halo blight disease, which is

characterized by large necrotic lessons surrounded by a chlorotic zone (halo) of yellow tissue

(14-16). Bean pathogens of phylogroup 2 are generally classified as pathovar *syringae* and are

responsible for bacterial spot disease, which is characterized by brown leaf spots (17, 18).

While halo blight can cause very serious crop losses, bacterial spot disease is generally of only

minor agronomic concern. The phylogroup 3 *phaseolicola* bean isolates show a high degree of

phylogenetic clustering, with most strains sharing a relatively recent common ancestor that is

closely related to a compact sister clade of soybean pathogens. In contrast, phylogroup 2

syringae bean isolates show very little phylogenetic clustering and are frequently more closely

related to a non-bean isolates than another bean isolates (9).

Assuming that host specificity is a heritable trait, the exploitation of a common host by divergent

lineages of strains can be explained by a number of different mechanisms, including: 1)

evolution via shared, vertically transmitted host specificity factors; 2) convergent evolution via

unrelated genetic mechanisms; or 3) convergent evolution via the horizontal acquisition of host

specificity factors from divergent lineages. Another layer of complexity is that host specificity

could come about either through the gain of genetic factors that promotes growth on a new host,

or alternatively, by the loss of a factor that otherwise limits growth (e.g., by inducing a host

immune response). In fact, the most thorough study of host convergence in *P. syringae*

suggests that isolates can make use of multiple mechanisms simultaneously. For example,

diverse lineages of cherry pathogens use multiple mechanisms to cause disease and have

exchanged and lost key genes that contribute to successful infections multiple times (11, 12).

One of the most important and dynamic classes of *P. syringae* virulence and host specificity

factors are type III secreted effectors (T3SEs). T3SE are proteins secreted through the type III

secretion system and translocated directly into the eukaryotic host cell where they can directly

interfere with host immunity or disrupt cellular homeostasis to promote the disease process.

There are 70 distinct families of *P. syringae* T3SEs, and most strains carry a suite of T3SEs

consisting of 12 to 50 T3SEs, with an average of ~30 (19). Plants have responded to T3SEs by

evolving immune receptors and complexes that trigger an effector-triggered immune (ETI)

response when they detect the presence or activity of a T3SE (20, 21). Consequently, the

outcome of any particular host-microbe interaction depends to a large degree on the specific

T3SE profile of the pathogen, and the complement of immune receptors carried by the host. The

strong selective pressures imposed by the host-microbe arms race results in very dynamic

evolution of T3SEs in general, with very frequent horizontal transmission, acquisition, and loss

(9, 19, 22).

While nearly every strain in phylogroups 2 and 3 carries a substantial suite of T3SEs, the suites

vary in size, with phylogroup 3 strains carrying an average of ~27 T3SEs vs. ~19 for phylogroup

2 strains (19). Phylogroup 2 strains are also known to carry more phytotoxins, which contribute

to virulence and niche competition via a variety of mechanisms such as membrane disruption

and mimicking of hormones (3, 9, 23). In general, phylogroup 2 strains are believed to show

lower levels of host specificity and better ability to survive on leaf surfaces (i.e., epiphytic

growth) (1-3, 24).

The application of statistical genetic and machine learning approaches to genomic data has

greatly increased our power to identify genes underlying traits of interest, such as host

specificity (25). Statistical genetic approaches such as genome-wide association studies

(GWAS) are well developed for studying human traits and have more recently gained traction in

the study of bacterial traits as statistical and phylogenetic methods have been developed for

dealing with the shared evolutionary history of segregating genetic variants (i.e., population

structure) (26-31). While GWAS approaches have great power for finding genotype-phenotype

associations, they generally measure associations on a locus-by-locus basis, and therefore can

miss more complex interactions among loci that impact traits. An alternative approach for

predicting genotype-phenotype associations is to use machine learning, which generally

describes a large range of statistical approaches that create models derived from a dataset

consisting of features (e.g., genetic variants) linked to a trait or outcome (e.g., host specificity).

These models can be used to predict outcomes from new samples or to identify the feature(s)

that carry the most importance in the model. Machine learning approaches are better at

identifying suites of interacting genetic variants than GWAS, but they are more limited in their

ability to deal with complex evolutionary relationships among these variants.

Here, we implemented a rapid, high-throughput method for screening a proxy of virulence for *P.*

syringae isolates on common bean. We use this screen to assess virulence of 121 strains from

nine phylogroups on bean, and then impute the virulence for an additional set isolates based on

their core genome relationship to the screened strains. While the majority of bean pathogens

come from phylogroups 2 and 3, we find that phylogroup 3 pathogens display a much stronger

degree of host specificity in comparison to phylogroup 2 pathogens. Finally, we develop a

random forest regression model using genomic kmers and virulence factors as features to

predict the virulence of *P. syringae* isolates on bean.

MATERIALS AND METHODS

Strain collection. Three hundred and thirty-four *Pseudomonas syringae* strains were used in

this study (Table S1). Forty-six *P. syringae* isolates were collected from bean fields

approximately 80 km East of Lethbridge, Alberta, Canada, during the summer of 2012. Bean

leaves with symptoms of bacterial diseases were collected from new growth during the

vegetative growth stage. The remaining 288 isolates were previously published (9) and include

49 *P. syringae* type and pathotype strains (1, 32). A type strain is the isolate to which the

scientific name of that organism is formally attached under the rules of prokaryote

nomenclature, while a pathotype strain is similar but with the additional requirement that it has

the pathogenic characteristics of its pathovar (i.e., a pathogen of a particular host) (5). Out of

the 334, 318 strains were used for comparative analyses and model training, while 16 were

used for model functional validation. A subset of 267 non-clonal representative strains

(discussed below) were selected for the predictive modeling to avoid clonal bias. A further

subset of 121 isolates, including the type and pathotype strains, were selected for virulence

assays.

Sequencing and quality control. DNA was extracted using the Gentra Puregene Yeast and

Bacteria kit (Qiagen, Hilden, Germany). Illumina libraries with 300-400 bp inserts were

generated using the Illumina Nextera XT kit according to the manufacturer's protocol (Illumina,

CA, USA). Samples were multiplexed with the Illumina Nextera XT Index kit containing 96

indices. Samples were sequenced on the Illumina NextSeq 500 Mid Output v2 (300 cycle) kit

with 150 base PE reads. All sequencing was performed at the University of Toronto's Centre for

the Analysis of Genome Evolution and Function (CAGEF). Raw read quality was assessed with

FastQC. Trimmomatic was used to remove adapters and trim raw sequencing reads based on a

sliding window approach (window size = 4, required quality = 5).

De novo **assembly.** Paired-end reads were *de novo* assembled using the CLC Genomics

Assembly Tool (CLC Genomics Workbench). Contigs shorter than 1kb were removed from the

assemblies. Low coverage contigs with matches to non-*Pseudomonas* genera and no matches

to the *Pseudomonas* genus that had a depth of coverage less than one standard deviation from

the average assembly coverage were deemed contaminants and, therefore, removed from the

final draft.

Pangenome analysis, gene prediction, annotation, and orthologous clustering. Gene

prediction and annotation for all assemblies were performed with Prokka (33). Prokka annotates

proteins by searching for sequence similarity in the UniProtKB (34) database and HMM libraries

(35). Additionally, all predicted genes were aligned against a custom T3SE database for the

identification of potential T3SEs (19). Orthologs were inferred with PIRATE (36), which clusters

genes into orthologous groups by performing all-vs-all comparisons followed by MCL clustering

given a certain percent identify threshold. Genes present in at least 95% of the genomes were

classified as core. Core protein families were individually aligned with MUSCLE (37) and later

concatenated into a single protein alignment. We used FastTree2 (38) to infer the phylogenetic

relationships of all 318 isolates.

Identification of non-clonal representative strains. We reduced the impact of phylogenetic

bias in our predictive modeling by selecting only one representative strain from each clonal

group (i.e., very closely related strains recently derived from a common ancestor). We identified

clonal groups by calculating the pairwise core genome evolutionary distance and the Jaccard

similarity for T3SE profiles. We found the minimum pairwise core-genome evolutionary distance

for isolates with identical T3SE profiles to be 0.001 average substitutions per site. We therefore

pooled the 318 isolates if they had a core genome evolutionary distance of less than 0.001,

resulting in 209 clusters. We further supplemented these clusters by adding back any strain that

had a T3SE profile Jaccard similarity value less than 0.8, resulting in 267 non-clonal clusters. A

single representative was selected out of each of these non-clonal clusters for downstream

analyses. One exception was made to the strain selection process in order to balance our

experimental design, which was skewed due to the fact that the vast majority of phylogroup 3

bean isolates (i.e., pathovar *phaseolicola*) fall into one clonal group. Initially, our selection

criteria resulted in only six phylogroup 3 bean isolates compared to 16 phylogroup 2 bean

isolates (i.e., pathovar *syringae*). We therefore added an additional seven *phaseolicola* strains

to the screened set to better balance the number of bean isolates in phylogroups 2 and 3.

Evolutionary distances and Jaccard similarity scores were inferred with MEGA7 (39) and R

version 4.0.5 (40).

Seed infection virulence assay. *P. syringae* strains were grown overnight at 30°C in King's B

media, re-suspended in 10 mM $MgSO_4$ and diluted to an OD_{600} of 0.001. *P. vulgaris* var.

Canadian red seeds were soaked for 24 hours in the bacterial suspension, planted in Sunshine

Mix 1 soil with regular watering and grown for 14 days. Plant fresh weight and germination

frequencies were measured and normalized to a control plant treated with 10 mM $MgSO_4$ sown

on each flat.

Syringe infiltration virulence assays. *P. syringae* strains were grown overnight on appropriate

antibiotics, re-suspended in 10 mM $MgSO_4$ and diluted to OD_{600} of 0.001. Two- to three-week-

old *Phaseolus vulgaris* var. Canadian red plants were syringe infiltrated and bacterial growth

assays were carried out by harvesting eight leaf disks (1 cm^2) from each plant (two per each

primary leaf) three days after infiltration. Disks were homogenized using a bead-beater in 200 µl

sterile 10 mM $MgSO_4$, serially diluted in 96-well plates, and 5 µl from each dilution was spot

plated on KB supplemented with rifampicin for positive and negative control strains, and

rifampicin and kanamycin for strains harboring constructs. Plates were incubated for at least 24

hours at 30°C and the resulting colony counts were used to calculate the number of CFUs per

cm^2 in the leaf apoplast.

Predictive modeling of *P. syringae* virulence on bean. We used an implementation of

gradient boosted decision trees to model the effect of *P. syringae* isolates on plant weights

using: 1) whole genome kmers, 2) T3SE kmers, and 3) a presence / absence matrix of T3SEs

and phytotoxins. We split sequences into 31-mers with fsm-lite and generated a binary matrix

for kmers with identical distribution patterns using custom python scripts. Next, we used the

Scikit-learn and the XGBoost python libraries (41) to generate a regression model for the

prediction of normalized plant weights using all three datasets as input features. Given the

relatively small size of our dataset, we used a cross-validation (CV) procedure to assess the

performance of our model on 50 independent splits. For each time, we randomly split the data

into training (80%) and testing (20%) sets while maintaining the same plant weight distributions

on both sets. Hyper parameters were fine-tuned using Scikit-learn's RandomizedSearchCV

module and regression models were generated with XGBoost's XGBRFRegressor module.

RESULTS

Virulence screen development. We developed a high-throughput seed infection assay to

measure the virulence of *P. syringae* isolates on common bean. We used seed infection based

on the understanding that many bean infections are caused by contaminated seeds (42, 43)

(14, 44). These infections can reduce overall plant health, which is reflected in plant fresh

weight. For the screen, we soaked bean seeds (*P. vulgaris* var. Canadian Red) in a *P. syringae*

suspension (~5x10^5 cells / ml) for 24 hours prior to planting. Plant fresh weight was then

determined after 14 days. We first assessed if plant weight was correlated with *in planta*

bacterial load by comparing our seed infection assay to the traditional syringe infiltration

virulence assay using 24 *P. syringae* isolates from 9 out of the 13 phylogroups (Fig. 1). Well-

established bean pathogens such as *P. syringae* pv. *phaseolicola* 1448A (Pph1448A) (14, 16)

and *P. syringae* pv. *syringae* B728a (PsyB728a) (17, 45) showed the highest levels of bacterial

growth and lowest plant weights, while the other isolates from phylogroups 1-7, 11, and 13

showed a range of values. On the whole there was a significant negative association between

bacterial growth and plant weight (R^2=0.63, P+5.0e-6), supporting the use of seed infection and

plant fresh weight to assess bacterial virulence.

To determine the power of this assay, we performed initial seed infection trials with six *P.*

syringae isolates and 50 or more replicate plants. We used a rarefaction analysis of normalized

plant weights to determine the number of replicate plants required to distinguish pathogens from

non-pathogens with >95% confidence (Tukey-HSD test) and found that the test power

plateaued at ~20 replicates per treatment (Fig. S1). Therefore, we performed future seed

infection assays using 30 replicate plants per treatment.

Virulence screen. We screened 121 non-clonal representative *P. syringae* isolates from nine

phylogroups to assess the virulence potential as measured by reduced plant fresh weight in 14-

day old bean plants after seed infection. The screened strains resulted a highly skewed

distribution of normalized fresh weights (i.e., virulence), with a mean of 0.78, median of 0.94,

and standard deviation of 0.30 (Fig. 2). An examination of the 30 strains in the first quartile

revealed normalized fresh weights between 0.00 and 0.61, with 56.7% (17 strains) being bean

isolates. These 17 strains represent 58.6% of all 29 bean isolates in screened.

Significant differences in virulence, as measured by normalized fresh weight, were observed

when comparing the strain collection stratified by host of isolation and phylogroup (Table 1). The

29 bean isolates had an average virulence (normalized fresh weight) of 0.59 compared to 0.85

for the 92 non-bean isolates (p=4.2e-5, 2-tailed, heteroscedastic T-test). All of the bean isolates

are found in phylogroups 2 and 3, so we compared the virulence of bean isolates to non-bean

isolates within these two phylogroups individually and found no significant difference for

phylogroup 2 (p=0.128, 2-tailed, heteroscedastic T-test) but a strong difference for phylogroup 3

(p=8.9e-4, 2-tailed heteroscedastic T-test). Additionally, there were no significant differences

between phylogroup 2 and phylogroup 3 bean isolates (p=0.460). We then looked for

differences in virulence between strains from different phylogroups irrespective of their host of

isolation (only comparing phylogroups with at least six tested strains, using 2-tailed,

heteroscedastic T-tests, Bonferroni corrected for seven total tests), and found that strains in

phylogroup 2 were significantly more virulent on bean than strains from phylogroups 1, 3, and 4

(p= 1.33e-07, 0.012, and 0.029 respectively), but not relative to phylogroup 6. In contrast,

strains from phylogroup 3 were only significantly more virulent on bean than strains from

phylogroup 1 (p=0.006). No other significant pairwise phylogroup comparisons were observed.

Interestingly, when we asked if non-bean phylogroup 2 isolates showed higher virulence on

bean than non-bean isolates from all other phylogroups, we observed a significant difference

(p=2.87e-4) indicating that while bean isolates specifically show greater virulence on bean,

phylogroup 2 isolates in general also show greater virulence on bean. This pattern was actually

reversed in phylogroup 3 where non-bean isolates had significantly lower virulence than non-

bean isolates from all other phylogroups (p=4.52e-3).

Virulence and germination inhibition. We then assessed whether the virulence of a strain

also influenced the germination frequency of bean seeds. Pathogenic microbes are known to

interfere with the seed germination (42, 43, 46) both through the direct action of phytotoxins (23)

and the indirect action of immune activation. In fact, seedling growth inhibition is a well-

established assay for immune activation in *Arabidopsis thaliana* (47). In general, the frequency

distribution for bean germination inhibition was much less skewed than the frequency

distribution for virulence, with mean = 60.1%, media = 63.3%, and standard deviation = 24.3%

(Fig 2, Table 1). The average germination frequency for all bean isolates was 45.9% compared

to 64.6% for non-bean isolates (p=4.92e-4, 2-tailed, heteroscedastic T-test). When stratifying

the bean isolates by phylogroup, we found no significant difference in germination frequency

between phylogroup 2 bean and non-bean isolates (p=0.076), while there was for phylogroup 3

(p=0.002), and we did observe a significant difference between germination frequency for

phylogroup 2 bean isolates and phylogroup 3 bean isolates (p=0.014). Similar to what was

found for the virulence assays, phylogroup 2 strains were significantly different from strains from

phylogroups 1, 3, and 4 (p=0.010, 6.23e-5, 8.53e-11 respectively 2-tailed, heteroscedastic T-

test, Bonferroni corrected), while phylogroup three strain were also significantly different from

phylogroup 4 (p=2.23e-4). Also similar to the virulence results, non-bean phylogroup 2 isolates

result in lower germination rates than all other non-bean isolates (p=9.60e-5), while non-bean

phylogroup 3 strains had a higher germination rate than non-bean isolates from all other

phylogroups (p=0.004), although this pattern disappears when non-bean phylogroup 2 strains

were removed from the analysis.

Finally, we measured the association between virulence (i.e., normalized fresh weight) and

germination frequency and found a strong association between the two metrics for the full

dataset (R^2=0.46, p=2.2e-16). Stratifying by phylogroup and bean isolates showed a strong

association for phylogroup 2 bean isolates (linear regression; F = 28.91, df = 13, p=0.0001, R^2 =

0.68), but no significant association for phylogroup 3 bean isolates (R^2=0.62, p=0.06) (Fig 3).

Predictive modelling of *P. syringae* virulence on bean. We used a gradient boosted decision

tree regression model to predict the virulence of *P. syringae* isolates on beans. The model used

one of three input feature classes: 1) genomic kmers; 2) T3SE kmers; or 3) presence / absence

of T3SEs and phytotoxins. T3SEs and phytotoxins are well-know virulence factors, with the

former often strongly associated with host specificity. Plant weight 14 days after seed infection

was used as a continuous outcome. The goal of this model was to assess the power of machine

learning to predict disease outcomes based on genome sequences and to predict the host

specificity of new isolates based on their genome sequence.

We used two nested collections of strains to generate the model. The first collection included

the 114 of the isolates directly screened for virulence, which included 50 phylogroup 2 strains

and 42 phylogroup 3 strains. This collection is slightly smaller than the full screened set since it

does not include the additional phylogroup 3 bean isolates added to balance the experimental

design (q.v., materials and methods). The second collection was an expanded strain set in

which we imputed virulence values based on genomic similarity. The imputation process

identified strains in our collection that had a core genome evolutionary distance of less than

0.001 and a T3SE Jaccard similarity of greater than 0.8 compared to one of the screened

strains. Any strains meeting these criteria were assigned the same virulence as the

corresponding screen strain. This imputation process almost tripled the size of our sample set,

resulting in an expanded collection of 318 strain, which included 67 phylogroup 2 strains and

142 phylogroup 3 strains. We also tested phylogroup 2 and phylogroup 3 strains separately

since bean isolates from these phylogroups interact with their host very differently.

Our analysis revealed a significant linear relationship between sample size and the overall

performance of the models (Fig. 5, MAE linear regression F = 5.89, df = 10, p=0.03, R^2 = 0.3).

Hence, the data sets with the least number of samples consistently showed the poorest

performance regardless of the input data (e.g., phylogroup 2 and the screened collection).

Furthermore, our models achieved the highest predictive power when looking exclusively at

isolates from PG3, which may be explained by the strong clonality patterns displayed by bean

pathogens from this phylogroup. Perhaps not surprisingly, models built using the greatest

diversity, with respect to number of strains (i.e., the expanded strain collection) and number of

genetic feature (i.e., whole genome kmers) showed the highest predictive power, with the best

model having a mean absolute error (MAE) of 0.06.

Model functional validation. We evaluate the power of our gradient boosted decision tree

regression model on unseen strains by using the kmer profiles of *P. syringae* isolates not

previously examined to predict virulence, and comparing these predictions to actual virulence

measures obtained through the seed infection virulence assay. Importantly, none of the strains

in the functional validation set were clonal related to any screened strain - they did not share a

core genome evolutionary distance of >0.001 and a T3SE Jaccard similarity of <0.8 with any

screened strain. When comparing actual to predicted virulence on the 16 strains in the

functional validation set, we found that 15 (94%) strains had virulence levels within the bounds

of estimated predictions given the calculated RMSE values (±0.20) (Fig. 5).

DISCUSSION

This work addressed the question of whether host of isolation is a reliable predictor of host

specific virulence, and conversely, can we use whole genome sequences to predict the host

specific virulence potential of individual strains. *Pseudomonas syringae* host specificity has

been studied in one form or another for many years. Some of the earliest molecular plant

immunity work was done in *P. syringae* with the cloning of the first "avirulence" gene (i.e., an

ETI-eliciting T3SE) in 1984 (48). Since those early studies, current technologies and

methodologies now enable the genome sequencing and functional characterization of hundreds

of strains (9, 24, 49). Despite this outstanding progress, we still have very limited ability to

predict the virulence potential of individual strains, even when full genome sequences are

available. The host of isolation is an extremely widely used surrogate for host specificity, and to

a large extent the basis for the pathovar nomenclature system. Despite the importance of this

assumption, it has rarely been systematically studied, and often fails when put to the test (24,

49).

Our first aim was to determine if infection of bean seeds by *P. syringae* recapitulated virulence

responses seen in more standard syringe inoculation virulence assays of bacteria into leaves.

We found a negative association between normalized plant fresh weight after seed infection and

in planta bacterial growth after syringe infiltration into leaf tissue, showing that the alternative

infection protocol works, and suggesting that virulence can systemically disrupt growth and

development. This finding is consistent with published and anecdotal reports that infected seed

stocks are a significance source of bean disease (42, 43, 50, 51). In general, we found that

bean isolates reduced mean plant fresh weight by 30.2% and median weight by 46.2%

compared to non-bean isolates. While the phylogroup 2 leaf spot bean pathogens had a

normalized mean fresh weight of 0.56±0.293 (stddev) compared to 0.64±0.246 for the

phylogroup 3 halo blight bean pathogens, this difference was not significantly different. A similar

pattern was found when we examined seed germination frequencies, where bean isolates

reduced the average germination frequency by 28.9% and median frequency by 22.0%

compared to non-bean isolates. But in this case, the 38.8% mean germination frequency of

phylogroup 2 bean pathogens was significantly lower than the 57.0% mean germination

frequency of the phylogroup 3 bean pathogens (p=0.014).

Phylogroup 2 strains in general showed lower levels of host specificity on bean. Neither our

fresh weight virulence measure nor the germination rates were significantly different between

the bean and non-bean isolates of phylogroup 2. This comparison was quite different for

phylogroup 3 strains, where bean isolates showed significantly higher virulence and lower

germination frequency compared to non-bean isolates from the same phylogroup. These

findings are consistent with other studies that have found lower levels of host specificity among

phylogroup 2 strains (24, 49) and lends support to the hypothesis that phylogroup 2 strains may

rely as much or more on toxin than T3SEs than other *P. syringae* strains.

While many genome-wide association studies have successfully identified strong genotype to

phenotype linkages to date, we were unable to identify any loci significantly associated with

bean isolation (data not shown). Consequently, we shifted our focus to machine learning

approaches as they can not only unravel genomic signatures associated with continuous

phenotypes, but also predict the virulence potential of previously unseen isolates given their

genome sequences. Regardless of phylogroup affiliation, our model was able to predict the

virulence of individual *P. syringae* isolates with low error margins, based solely on whole

genome data. However, the fact that models trained on virulence factors alone could predict

virulence with considerable accuracy supports the notion that T3SEs and phytotoxins play

crucial roles in host adaptation processes. Nonetheless, the higher predictive power of models

trained with whole genome kmers suggests that factors other than virulence-associated genes

also play important roles on disease development and adaptation to beans.

Sample size is one of the most important contributors to accurate model generation in machine

learning. Simple models trained with large data sets can learn to distinguish label categories

with much stronger accuracy than complex models trained with small datasets. Our results also

found a correlation between sample size and model accuracy, with the screened strain

collection providing a MAE of 0.153 while the larger imputed strain collection increased model

performance to a MAE of 0.065. We find much poorer model performance for phylogroup 2

strains than phylogroup 3 strains. While this may partly reflect the differences in sample size

between these two groups, it also likely reflects the underlying biology. The majority of bean

isolates in phylogroup 3 are phylogenetically clustered, while there is little clustering of bean

isolates in phylogroup 2. Consequently, the model may perform better on phylogroup 3 since it

is essentially predicting phylogenetic structure. Another contributing factor is likely the finding

discussed above, namely, host specificity appears to be weaker in phylogroup 2. If phylogroup 2

strains are more generalists than specialists, then the host specificity signal would be weaker

and any model trying to find this signal would perform more poorly.

REFERENCES

1.   Baltrus DA, McCann HC, Guttman DS. 2017. Evolution, genomics and epidemiology of
      Pseudomonas syringae: Challenges in Bacterial Molecular Plant Pathology. Mol Plant
      Pathol 18:152-168.

2.   Morris CE, Monteil CL, Berge O. 2013. The life history of *Pseudomonas syringae*: linking
      agriculture to earth system processes. Annu Rev Phytopathol 51:85-104.

3.   Xin XF, Kvitko B, He SY. 2018. *Pseudomonas syringae*: what it takes to be a pathogen.
      Nat Rev Microbiol 16:316-328.

4.   Morris CE, Sands DC, Vinatzer BA, Glaux C, Guilbaud C, Buffiere A, Yan S, Dominguez
      H, Thompson BM. 2008. The life history of the plant pathogen *Pseudomonas syringae* is
      linked to the water cycle. ISME J 2:321-34.

5.   Bull CT, De Boer SH, Denny TP, Firrao G, Fischer-Le Saux M, Saddler GS, Scortichini
      M, Stead DE, Takikawa Y. 2008. Demystifying the nomenclature of bacterial plant
      pathogens. J Plant Pathol 90:403-417.

6.   Dye DW, Bradbury JF, Goto M, Hayward AC, Lelliott RA, Schroth MN. 1980.
      International standards for naming pathovars of phytopathogenic bacteria and a list of
      pathovar names and pathotype strains. Review of Plant Pathology 59:153-168.

7.   Berge O, Monteil CL, Bartoli C, Chandeysson C, Guilbaud C, Sands DC, Morris CE.
      2014. A user's guide to a data base of the diversity of *Pseudomonas syringae* and its
      application to classifying strains in this phylogenetic complex. PLoS One 9:e105547.

8.   Sarkar SF, Guttman DS. 2004. Evolution of the core genome of *Pseudomonas syringae*,
      a highly clonal, endemic plant pathogen. Appl Environ Microbiol 70:1999-2012.

9.   Dillon MM, Thakur S, Almeida RND, Wang PW, Weir BS, Guttman DS. 2019.
      Recombination of ecologically and evolutionarily significant loci maintains genetic
      cohesion in the *Pseudomonas syringae* species complex. Genome Biol 20:3.

10.　Hwang MS, Morgan RL, Sarkar SF, Wang PW, Guttman DS. 2005. Phylogenetic

characterization of virulence and resistance phenotypes of *Pseudomonas syringae*. Appl

Environ Microbiol 71:5182-91.

11.　Hulin MT, Armitage AD, Vicente JG, Holub EB, Baxter L, Bates HJ, Mansfield JW,

Jackson RW, Harrison RJ. 2018. Comparative genomics of *Pseudomonas syringae*

reveals convergent gene gain and loss associated with specialization onto cherry

(*Prunus avium*). New Phytol 219:672-696.

12.　Hulin MT, Mansfield JW, Brain P, Xu X, Jackson RW, Harrison RJ. 2018.

Characterization of the pathogenicity of strains of *Pseudomonas syringae* towards cherry

and plum. Plant Pathol 67:1177-1193.

13.　O'Brien HE, Thakur S, Gong Y, Fung P, Zhang J, Yuan L, Wang PW, Yong C,

Scortichini M, Guttman DS. 2012. Extensive remodeling of the *Pseudomonas syringae*

pv. *avellanae* type III secretome associated with two independent host shifts onto

hazelnut. BMC Microbiol 12:141.

14.　Arnold DL, Lovell HC, Jackson RW, Mansfield JW. 2011. Pseudomonas syringae pv.

phaseolicola: from 'has bean' to supermodel. Mol Plant Pathol 12:617-27.

15.　Tsiamis G, Mansfield JW, Hockenhull R, Jackson RW, Sesma A, Athanassopoulos E,

Bennett MA, Stevens C, Vivian A, Taylor JD, Murillo J. 2000. Cultivar-specific avirulence

and virulence functions assigned to avrPphF in *Pseudomonas syringae* pv. phaseolicola,

the cause of bean halo-blight disease. EMBO J 19:3204-14.

16.　Joardar V, Lindeberg M, Jackson RW, Selengut J, Dodson R, Brinkac LM, Daugherty

SC, Deboy R, Durkin AS, Giglio MG, Madupu R, Nelson WC, Rosovitz MJ, Sullivan S,

Crabtree J, Creasy T, Davidsen T, Haft DH, Zafar N, Zhou L, Halpin R, Holley T, Khouri

H, Feldblyum T, White O, Fraser CM, Chatterjee AK, Cartinhour S, Schneider DJ,

Mansfield J, Collmer A, Buell CR. 2005. Whole-genome sequence analysis of

485  *Pseudomonas syringae* pv. *phaseolicola* 1448A reveals divergence among pathovars in

486  genes involved in virulence and transposition. J Bacteriol 187:6488-98.

487 17. Feil H, Feil WS, Chain P, Larimer F, DiBartolo G, Copeland A, Lykidis A, Trong S, Nolan

488  M, Goltsman E, Thiel J, Malfatti S, Loper JE, Lapidus A, Detter JC, Land M, Richardson

489  PM, Kyrpides NC, Ivanova N, Lindow SE. 2005. Comparison of the complete genome

490  sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. *tomato* DC3000. Proc

491  Natl Acad Sci U S A 102:11064-9.

492 18. Marco ML, Legac J, Lindow SE. 2005. Pseudomonas syringae genes induced during

493  colonization of leaf surfaces. Environ Microbiol 7:1379-1391.

494 19. Dillon MM, Almeida RND, Laflamme B, Martel A, Weir BS, Desveaux D, Guttman DS.

495  2019. Molecular evolution of *Pseudomonas syringae* type III secreted effector proteins.

496  Front Plant Sci 10:418.

497 20. Buttner D. 2016. Behind the lines-actions of bacterial type III effector proteins in plant

498  cells. FEMS Microbiol Rev 40:894-937.

499 21. Khan M, Seto D, Subramaniam R, Desveaux D. 2018. Oh, the places they'll go! A survey

500  of phytopathogen effectors and their host targets. Plant J 93:651-663.

501 22. Martel A, Ruiz-Bedoya T, Breit-McNally C, Laflamme B, Desveaux D, Guttman DS.

502  2021. The ETS-ETI cycle: evolutionary processes and metapopulation dynamics driving

503  the diversification of pathogen effectors and host immune factors. Curr Opin Plant Biol

504  62:102011.

505 23. Bender CL, Alarcon-Chaidez F, Gross DC. 1999. *Pseudomonas syringae* phytotoxins:

506  mode of action, regulation, and biosynthesis by peptide and polyketide synthetases.

507  Microbiol Mol Biol Rev 63:266-92.

508 24. Morris CE, Moury B. 2019. Revisiting the concept of host range of plant pathogens.

509  Annu Rev Phytopathol doi:10.1146/annurev-phyto-082718-100034.

25.   Allen JP, Snitkin E, Pincus NB, Hauser AR. 2021. Forest and Trees: Exploring Bacterial

Virulence with Genome-wide Association Studies and Machine Learning. Trends

Microbiol doi:10.1016/j.tim.2020.12.002.

26.   Chen PE, Shapiro BJ. 2015. The advent of genome-wide association studies for

bacteria. Curr Opin Microbiol 25:17-24.

27.   Falush D. 2016. Bacterial genomics: Microbial GWAS coming of age. Nat Microbiol

1:16059.

28.   Falush D, Bowden R. 2006. Genome-wide association mapping in bacteria? Trends

Microbiol 14:353-5.

29.   Power RA, Parkhill J, de Oliveira T. 2017. Microbial genome-wide association studies:

lessons from human GWAS. Nat Rev Genet 18:41-50.

30.   Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA, Kelly DJ, Bentley SD, Maiden

MC, Parkhill J, Falush D. 2013. Genome-wide association study identifies vitamin B5

biosynthesis as a host specificity factor in *Campylobacter*. Proc Natl Acad Sci U S A

110:11923-7.

31.   Earle SG, Wu CH, Charlesworth J, Stoesser N, Gordon NC, Walker TM, Spencer CC,

Iqbal Z, Clifton DA, Hopkins KL, Woodford N, Smith EG, Ismail N, Llewelyn MJ, Peto TE,

Crook DW, McVean G, Walker AS, Wilson DJ. 2016. Identifying lineage effects when

controlling for population structure improves power in bacterial association studies. Nat

Microbiol 1:16041.

32.   Thakur S, Weir BS, Guttman DS. 2016. Phytopathogen Genome Announcement: draft

genome sequences of 62 *Pseudomonas syringae* type and pathotype strains. Mol Plant

Microbe Interact 29:243-6.

33.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics

30:2068-9.

34. Apweiler R, Bairoch A, Wu CH, Barker WC, Boeckmann B, Ferro S, Gasteiger E, Huang

H, Lopez R, Magrane M, Martin MJ, Natale DA, O'Donovan C, Redaschi N, Yeh LS.

2004. UniProt: the Universal Protein knowledgebase. Nucleic Acids Res 32:D115-9.

35. Mistry J, Finn RD, Eddy SR, Bateman A, Punta M. 2013. Challenges in homology

search: HMMER3 and convergent evolution of coiled-coil regions. Nucleic Acids Res

41:e121.

36. Bayliss SC, Thorpe HA, Coyle NM, Sheppard SK, Feil EJ. 2019. PIRATE: A fast and

scalable pangenomics toolbox for clustering diverged orthologues in bacteria.

Gigascience 8.

37. Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time

and space complexity. BMC Bioinformatics 5:113.

38. Price MN, Dehal PS, Arkin AP. 2010. FastTree 2--approximately maximum-likelihood

trees for large alignments. PLoS One 5:e9490.

39. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. 2018. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. Mol Biol Evol 35:1547-1549.

40. R Development Core Team. 2020. R: A language and environment for statistical

computing, R Foundation for Statistical Computing, Vienna, Austria. https://www.R-

project.org/.

41. Chen T, Guestrin C. 2016. XGBoost: A Scalable Tree Boosting System, Proceedings of

the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data

Mining doi:10.1145/2939672.2939785. ACM, San Francisco, California, USA.

42. Darrasse A, Bureau C, Samson R, Morris CE, Jacques M-A. 2007. Contamination of

bean seeds by Xanthomonas axonopodis pv. phaseoli associated with low bacterial

densities in the phyllosphere under field and greenhouse conditions. Eur J Plant Pathol

119:203-215.

43. Eyster HC. 1940. The cause of decreased germination of bean seeds soaked in water.

Am J Bot 27:652-659.

44. Hirano SS, Demars SJ, Morris CE. 1981. Survival, establishment, and dispersal of

Pseudomonas s*yringae* on snap beans (*Phaseolus vulgaris* L). Phytopathology 71:881-

881.

45. Willis DK, Hrabak EM, Rich JJ, Barta TM, Lindow SE, Panopoulos NJ. 1990. Isolation

and characterization of a *Pseudomonas syringae* pathovar *syringae* mutant deficient In

lesion formation on bean. Mol Plant Microbe Interact 3:149-156.

46. Chahtane H, Nogueira Füller T, Allard PM, Marcourt L, Ferreira Queiroz E,

Shanmugabalaji V, Falquet J, Wolfender JL, Lopez-Molina L. 2018. The plant pathogen

Pseudomonas aeruginosa triggers a DELLA-dependent seed germination arrest in

Arabidopsis. Elife 7.

47. Bredow M, Sementchoukova I, Siegel K, Monaghan J. 2019. Pattern-Triggered

Oxidative Burst and Seedling Growth Inhibition Assays in Arabidopsis thaliana. J Vis Exp

doi:10.3791/59437.

48. Staskawicz BJ, Dahlbeck D, Keen NT. 1984. Cloned avirulence gene of *Pseudomonas*

syringae pv. *glycinea* determines race-specific incompatibility on *Glycine max* (L.) Merr.

Proc Natl Acad Sci U S A 81:6024-8.

49. Morris CE, Lamichhane JR, Nikolić I, Stanković S, Moury B. 2019. The overlapping

continuum of host range among strains in the *Pseudomonas syringae* complex.

Phytopathology Research 1:4.

50. Shade A, Jacques MA, Barret M. 2017. Ecological patterns of seed microbiome

diversity, transmission, and assembly. Curr Opin Microbiol 37:15-22.

51. Upper CD, Hirano SS, Dodd KK, Clayton MK. 2003. Factors that Affect Spread of

Pseudomonas syringae in the Phyllosphere. Phytopathol 93:1082-92.

TABLES

Table 1. Virulence and Germination Assay Summary

Group [1]	N Strains	Virulence [2]			Germination Frequency (%)		
		Mean	Median	Stdv	Mean	Median	Stdv
All Strains	121	0.789	0.913	0.271	60.10	63.33	24.31
PG1	8	0.972	0.995	0.043	72.99	79.17	15.71
PG2	50	0.658	0.748	0.320	46.74	53.33	26.41
PG3	42	0.841	0.964	0.218	68.15	67.50	16.53
PG4	6	0.924	1.000	0.147	83.61	82.50	4.52
PG5	3	0.966	1.000	0.059	73.33	78.33	10.14
PG6	3	1.000	1.000	0.000	80.56	81.67	3.47
PG7	2	0.956	0.956	0.062	85.00	85.00	7.07
PG11	6	0.851	0.867	0.151	43.89	50.00	20.75
PG13	1	1.000	1.000	NA	91.67	91.67	NA
All Bean Isolates	29	0.594	0.516	0.271	45.89	53.33	23.52
All Non-Bean	92	0.850	0.960	0.244	64.58	68.33	23.05
PG2 Bean	16	0.560	0.478	0.293	36.84	32.22	26.42
PG2 Non-Bean	34	0.704	0.841	0.327	51.41	56.67	25.47
PG3 Bean	13	0.635	0.606	0.246	57.02	58.89	13.21
PG3 Non-Bean	29	0.933	1.000	0.122	73.14	73.33	15.56

[1] PG = phylogroup
[2] Normalized fresh weight 2 weeks after seed infection

589
590

Table 2. Machine Learning Model Performance

Group	Samples	Dataset	MAE	RMSE
PG3	142	Whole genome kmers	0.049	0.107
PG3	142	Toxin and T3SE binary matrix	0.054	0.101
Imputed	318	Whole genome kmers	0.067	0.140
PG3	142	T3SE kmers	0.085	0.126
Imputed	318	Toxin and T3SE binary matrix	0.092	0.153
Imputed	318	T3SE kmers	0.096	0.177
PG2	66	T3SE kmers	0.154	0.233
Screened	114	Whole genome kmers	0.155	0.204
Screened	114	T3SE kmers	0.159	0.213
Screened	114	Toxin and T3SE binary matrix	0.163	0.201
PG2	66	Whole genome kmers	0.169	0.242
PG2	66	Toxin and T3SE binary matrix	0.202	0.262

FIGURE LEGENDS

Figure 1: Bacterial growth is inversely correlated with plant weight. Bacterial density in syringe infiltrated bean leaves as a function of normalized median weight of seed infected plants. There is a strong negative correlation between bacterial density and plant weight across isolates in the *P. syringae* species complex (linear regression; F = 36.95, df = 21, p = 4.95e-06, R^2 = 0.62).

Figure 2: Frequency distribution virulence and germination assays. Distributions of data from the 121 strains screened for virulence (plant fresh weight) and germination by the seed infection assay.

Figure 3: Normalized median plant fresh weights of seed infected bean plants. (A) Boxplots show virulence (normalized plant fresh weight) of 121 *P. syringae* strains after seed infection on bean. The boxplots are color-coded by the phylogroup of each strain, and asterisks above the boxplots denote bean isolates. Plants infected with bean isolates tend to have higher virulence (i.e., lower plant fresh weight) than plants infected with strains isolated from other hosts. (B) Germination frequency of seed infected plants. (C) Core genome phylogeny of 318 *P. syringae* isolates. Leaf colors represent phylogroup affiliation. Metadata row 1 shows isolation source; metadata row 2 shows normalized median weight of screened isolates, and metadata row 3 shows normalized median weight for the expanded strain collection based on phylogenetic imputation.

Figure 4: Correlation between germination frequency and virulence of seed infected bean plants. (A) Germination frequency vs virulence (i.e., normalized fresh weight) stratified by

phylogroup (PG) 2 and phylogroup 3 bean isolates. A strong correlation is found between

germination frequency and virulence of bean strains from phylogroup 2 (linear regression; $F =$

28.91, df = 13, $p = 0.00012$, $r^2 = 0.66$). (B) Germination frequency vs. virulence stratified by

phylogroup (PG) irrespective of host.

Figure 5: Performance of supervised machine learning models on virulence predictions

with genome data. Models trained with the screed collection (yellow), imputed expanded

collection (blue), imputed expanded phylogroup (PG) 2 set (orange), and imputed expanded

phylogroup 3 set (green). Diamonds, circles, and triangles represent models trained with a

phytotoxin and T3SE binary matrix, T3SE kmers, and whole genome kmers, respectively. (A)

Mean Absolute Error (MAE) as a function of number of samples across 50 cross-validation

splits. Increasing the number of samples seen by the model significantly increases model

performance (linear regression; $F = 5.89$, df = 10, $p = 0.03$, $r^2 = 0.37$). (B) Root Mean Square

Error (RMSE) as a function of number of samples across 50 cross-validation splits. Models

trained on whole genome kmers with a higher number of samples consistently outperform other

models.

Figure 6: Correlation between predicted and observed weights of plants infected with

isolates previously unseen by the model. (A) Predicted virulence as a function of observed

virulence on seed infected bean plants. Color coding indicates phylogroups (PG). Dashed line

indicates 1:1 relationship. (B) Virulence for each isolate tested ordered by predicted virulence

and colored by phylogroup. The solid line represents virulence predictions for each isolate. The

grey box represents the error margins for the predictions based on the MAE and RMSE values

for the model.

SUPPLEMENTARY MATERIAL

Figure S1: Sample size selection via a simulated experimental setup. Bean plants were

seed infected with 6 *P. syringae* isolates with a high number of replicates (>50). Plant weights

were randomly selected according to various replicate sizes (8-32). Our ability to distinguish

pathogens from non-pathogens using Tukey-HSD tests plateaus at 20 replicates per treatment.

Figure S2: MAE and RMSE across 50 cross-validation splits. Model performance in terms of

Mean Absolute Error and Root Mean Square Error. Plant weight distributions were kept the

same across splits.

Table S1: List of *P. syringae* isolates used in this study.

Fig 1

Fig 2.

667   Fig 3.

673   Fig 4.

678　　Fig 5.

683   Fig 6.

686
687
688

689   Fig S1

693 Fig S2.

696