Predictive Modeling of *Pseudomonas syringae* Virulence on Bean using Gradient

Boosted Decision Trees

Renan N.D. Almeida[1], Michael Greenberg[1], Cedoljub Bundalovic-Torma[1], Alexandre Martel[1],

Pauline W. Wang[1,2], Maggie A. Middleton[1,2], Syama Chatterton[3], Darrell Desveaux [1], and David

S. Guttman[1,2,*]

[1]Department of Cell & Systems Biology, University of Toronto, 25 Willcocks St., Toronto,

Ontario, Canada

[2]Centre for the Analysis of Genome Evolution & Function, University of Toronto, Toronto,

Ontario, Canada

[3]Agriculture and Agri-Food Canada, Lethbridge Research and Development Centre, Lethbridge,

Alberta, Canada

*To whom correspondence should be addressed. Email: david.guttman@utoronto.ca

Short title: Predicting *P. syringae* virulence on bean using a machine learning approach

Keywords: *Pseudomonas syringae*, *Phaseolus vulgaris*, bean, machine learning, random forest,

gradient boosted decision trees, host specificity, virulence,

ABSTRACT

Pseudomonas syringae is a genetically diverse bacterial species complex responsible for numerous agronomically important crop diseases. Individual *P. syringae* isolates are typically given pathovar designations based on their host of isolation and the associated disease symptoms, and these pathovar designations are often assumed to reflect host specificity although this assumption has rarely been rigorously tested. Here we developed a rapid seed infection assay to measure the virulence of 121 diverse *P. syringae* isolates on common bean (*Phaseolus vulgaris*). This collection includes *P. syringae* phylogroup 2 (PG2) bean isolates (pathovar *syringae*) that cause bacterial spot disease and *P. syringae* phylogroup 3 (PG3) bean isolates (pathovar *phaseolicola*) that cause the more serious halo blight disease. We found that bean isolates in general were significantly more virulent on bean than non-bean isolates and observed no significant virulence difference between the PG2 and PG3 bean isolates. However, when we compared virulence within PGs we found that PG3 bean isolates were significantly more virulent than PG3 non-bean isolates, while there was no significant difference in virulence between PG2 bean and non-bean isolates. These results indicate that PG3 strains have a higher level of host specificity than PG2 strains. We then employed machine learning to investigate if we could use genomic data to predict virulence on bean. We used gradient boosted decision trees to model the virulence using whole genome kmers, type III secreted effector kmers, and the presence/absence of type III effectors and phytotoxins. Our model performed best using whole genome data and was able to predict virulence with high accuracy (mean absolute error = 0.05). Finally, we functionally validated the model by predicting virulence for 16 strains and found that 15 (94%) had virulence levels within the bounds of estimated predictions. This study demonstrates the power of machine learning for predicting host specific adaptation and strengthens the hypothesis that *P. syringae* PG2 strains have evolved a different lifestyle than other *P. syringae* strains.

AUTHOR SUMMARY

Pseudomonas syringae is a genetically diverse Gammaproteobacterial species complex

responsible for numerous agronomically important crop diseases. Strains in the *P. syringae*

species complex are frequently categorized into pathovars depending on pathogenic

characteristics such as host of isolation and disease symptoms. Common bean pathogens from

P. syringae are known to cause two major diseases: the halo blight disease, which is

characterized by large necrotic lesions surrounded by a chlorotic zone or halo of yellow tissue;

and the bacterial spot disease, which is characterized by brown leaf spots. While halo blight can

cause serious crop losses, bacterial spot disease is generally of minor agronomic concern. The

application of statistical genetic and machine learning approaches to genomic data has greatly

increased our power to identify genes underlying traits of interest, such as host specificity.

Machine learning models can be used to predict outcomes from new samples or to identify the

genetic feature(s) that carry the most importance when predicting a particular phenotype. Here,

we implemented a rapid method for screening a proxy of virulence for *P. syringae* isolates on

common bean, and used this screen to assess virulence of *P. syringae* strains on bean. We

found that halo blight pathogens display a stronger degree of host specificity compared to brown

spot pathogens, and that genomic kmers and virulence factors can be used to predict the

virulence of *P. syringae* isolates on bean using machine learning models.

INTRODUCTION

Pseudomonas syringae is a genetically diverse Gammaproteobacterial species complex responsible for numerous agronomically important crop diseases [1-4]. Strains in the *P. syringae* species complex are frequently categorized into pathovars depending on pathogenic characteristics such as host of isolation and disease symptoms [5, 6]. The species complex is also subdivided into phylogenetic groups (i.e., phylogroups, PGs) based on multilocus sequence typing or genomic analysis [1, 7-10]. Currently, there are 13 recognized PGs [7], of which seven have been termed primary PGs based on their genetic relatedness and the near universal presence of the canonical *P. syringae* type III secretion system (discussed below) [1, 9].

Phylogenetic analyses of *P. syringae* isolates suggest that adaptation to specific hosts has evolved multiple times in the evolutionary history of the species complex. For example, cherry and plum pathogens are found in clades distributed in PG1, PG2, and PG3 [11, 12], hazelnut pathogens are distributed among two distinct clades in PG1 and PG2 [13], while pathogens of common bean (including snap, green, kidney, and French bean) are found in PG2 and PG3 [9].

Common bean pathogens from *P. syringae* PG3 are generally classified as pathovar *phaseolicola* and are responsible for halo blight disease, which is characterized by large necrotic lesions surrounded by a chlorotic zone or halo of yellow tissue [14-16]. Bean pathogens of PG2 are generally classified as pathovar *syringae* and are responsible for bacterial spot disease, which is characterized by brown leaf spots [17, 18]. While halo blight can cause serious crop losses, bacterial spot disease is generally of minor agronomic concern. The PG3 *phaseolicola* bean isolates show a high degree of phylogenetic clustering, with most strains sharing a relatively recent common ancestor that is closely related to a compact sister clade of soybean pathogens [9]. In contrast, PG2 *syringae* bean isolates show very little phylogenetic

clustering and are frequently more closely related to non-bean isolates than other bean isolates

[9].

Assuming that host specificity is a heritable trait, the exploitation of a common host by divergent

lineages of strains can be explained by several different mechanisms, including: 1) evolution via

shared, vertically transmitted host specificity factors; 2) convergent evolution via unrelated

genetic mechanisms; or 3) convergent evolution via the horizontal acquisition of host specificity

factors from divergent lineages. Another layer of complexity is that host specificity could come

about either through the gain of genetic factors that promotes growth on a new host, or

alternatively, by the loss of a factor that otherwise limits growth (e.g., by inducing a host immune

response). In fact, the most thorough study of host convergence in *P. syringae* suggests that

isolates can make use of multiple mechanisms simultaneously [11, 12]. For example, diverse

lineages of cherry pathogens have exchanged and lost key genes and used multiple

mechanisms to successfully infect this host [11, 12].

One of the most important and dynamic classes of *P. syringae* virulence and host specificity

factors are type III secreted effectors (T3SEs). T3SEs are proteins translocated through the type

III secretion system directly into the eukaryotic host cell where they interfere with host immunity

or disrupt cellular homeostasis to promote the disease process. There are 70 distinct families of

P. syringae T3SEs, and most strains carry a suite of T3SEs consisting of 12 to 50 T3SEs, with

an average of ~30 [19]. Plants have responded to T3SEs by evolving immune receptors and

complexes that trigger an effector-triggered immune (ETI) response when they detect the

presence or activity of a T3SE [20, 21]. Consequently, the outcome of any particular host-

microbe interaction depends to a large degree on the specific T3SE profile of the pathogen and

the complement of immune receptors carried by the host. The strong selective pressures

imposed by the host-microbe arms race results in dynamic evolution of T3SEs in general, with

frequent horizontal transmission, acquisition, and loss [9, 19, 22].

The suites of T3SEs carried by PG2 and PG3 strains vary in size, with PG2 strains carrying an

average of ~19 T3SEs vs. ~27 for PG3 strains [19]. PG2 strains are also known to carry more

phytotoxins, which contribute to virulence and niche competition via a variety of mechanisms

such as membrane disruption and hormone mimicry [3, 9, 23]. In general, PG2 strains are

believed to show lower levels of host specificity and better ability to survive on leaf surfaces

(i.e., epiphytic growth) [1, 3, 24, 25].

The application of statistical genetic and machine learning approaches to genomic data has

greatly increased our power to identify genes underlying traits of interest, such as host

specificity [26]. Statistical genetic approaches like genome-wide association studies (GWAS)

are well developed for studying human traits and have more recently gained traction in the study

of bacterial traits as statistical and phylogenetic methods have been developed to handle the

shared evolutionary history of segregating genetic variants (i.e., population structure) [27-32].

While GWAS approaches have great power for finding genotype-phenotype associations, they

generally measure associations on a locus-by-locus basis, and therefore can miss more

complex interactions among loci that impact traits. An alternative approach for predicting

genotype-phenotype associations is to use machine learning, which generally describes a large

range of statistical approaches that create models derived from a dataset consisting of features

(e.g., genetic variants) linked to a trait or outcome (e.g., host specificity). These models can be

used to predict outcomes from new samples or to identify the feature(s) that carry the most

importance in the model. Although machine learning approaches are better suited for identifying

suites of interacting genetic variants than GWAS, they are more limited in their ability to deal

with complex evolutionary relationships among these variants [33, 34].

Here, we implemented a rapid method for screening a proxy of virulence for *P. syringae* isolates

on common bean. We used this screen to assess virulence of 121 strains from nine PGs on

bean, and then expand the strain collection by imputing the virulence for an additional set of

isolates based on their core genome relationship to the screened strains. While most of bean

pathogens come from PG2 and PG3, we found that PG3 pathogens display a stronger degree

of host specificity compared to PG2 pathogens. Finally, we developed a random forest

regression model using genomic kmers and virulence factors as features to predict the virulence

of *P. syringae* isolates on bean.

RESULTS

Genome analysis. We characterized the genomic diversity of the isolates used in this study via

phylogenetic analysis (Fig. 1) and by measuring the pairwise core genome synonymous

substitution rates (Ks) and pairwise accessory genome Jaccard distances for each isolate

combination. The 36 bean halo blight pathogens of pathovar *phaseolicola* largely cluster in one

very closely related clade in PG3 and had a core genome Ks of 0.0039 and an accessory

genome Jaccard distance of 0.28 compared to the entirety of 142 PG3 strains, which had a core

genome Ks of 0.047 and an accessory genome Jaccard distance of 0.23. In contrast the 21

bean spot disease pathogens of pathovar *syringae* are broadly distributed throughout PG2 and

had a core genome Ks of 0.1218 and an accessory genome Jaccard distance of 0.51 compared

to the entirety of 66 PG2 strains, which had a core genome Ks of 0.1223 and an accessory

genome Jaccard distance of 0.27 (Fig. 2). Due to their clonal nature, PG3 bean isolates were

found to have a considerably higher number of gene families in the hard (present in 100% of the

isolates) and soft core (present in >95% of the isolates) genomes, as well as a lower number of

singleton families (present in a single isolate) in comparison to PG 2 bean isolates, despite the

higher number of PG3 samples analyzed (Fig. 3).

Virulence screen development. We developed a high-throughput seed infection assay to

measure the virulence of *P. syringae* isolates on common bean. We used seed infection based

on the understanding that many bean infections are caused by contaminated seeds [14, 45-47].

These infections can reduce overall plant health, which is reflected in plant fresh weight. For the

screen, we soaked bean seeds (*P. vulgaris* var. Canadian Red) in a *P. syringae* suspension

(~5×10^5 cells / ml) for 24 hours prior to planting. Plant fresh weight was then determined after 14

days. We first assessed if plant weight was correlated with *in planta* bacterial load by comparing

our seed infection assay to the traditional syringe infiltration virulence assay using 24 *P.*

syringae isolates from 9 out of the 13 PGs (Fig. 4). Well-established bean pathogens such as

PG3 strain *P. syringae* pv. *phaseolicola* 1448A (Pph1448A) [14, 16] and the PG2 strains *P.*

syringae pv. *syringae* B728a (PsyB728a) [17, 48] showed the highest levels of bacterial growth

and lowest plant weights, while the other isolates from PGs 1-7, 11, and 13 showed a range of

values. Overall, there was a significant negative association between bacterial growth and plant

weight (R^2=0.63, P+5.0e-6), supporting the use of seed infection and plant fresh weight to

assess bacterial virulence.

To determine the power of this assay, we performed initial seed infection trials with six *P.*

syringae isolates and 50 or more replicate plants. We used a rarefaction analysis of normalized

plant weights to determine the number of replicate plants required to distinguish pathogens from

non-pathogens with >95% confidence (Tukey-HSD test) and found that the test power

plateaued at ~20 replicates per treatment (Fig. S1). Therefore, we performed future seed

infection assays using 30 replicate plants per treatment.

Virulence screen. We screened 121 non-clonal representative *P. syringae* isolates from nine

PGs to assess the virulence potential as measured by reduced plant fresh weight in 14-day old

bean plants after seed infection (Fig. 5). This screened set was subset of the non-clonal set

selected to maximize coverage of the species complex while focusing on a manageable number

for screening. The screened strains resulted in a highly skewed distribution of normalized fresh

weights (i.e., virulence), with a mean of 0.78, median of 0.94, and standard deviation of 0.30. An

examination of the 30 strains in the first quartile revealed normalized fresh weights between

0.00 and 0.61, with 56.7% (17 strains) being bean isolates (Fig. 6). These 17 strains represent

58.6% of all 29 bean isolates screened.

Significant differences in virulence, as measured by normalized fresh weight, were observed

when comparing the strain collection stratified by host of isolation and PG (Table 1). The 29

bean isolates had an average virulence (normalized fresh weight) of 0.59 compared to 0.85 for

the 92 non-bean isolates (p=4.2e-5, 2-tailed, heteroscedastic T-test, same for tests discussed

below). As all the bean isolates are found in PG2 and PG3, we compared the virulence of bean

isolates to non-bean isolates within these two PGs individually and found no significant

difference for PG2 (p=0.128) but a strong difference for PG3 (p=8.9e-4). Additionally, there were

no significant differences between PG2 and PG3 bean isolates (p=0.460). We then looked for

differences in virulence between strains from different PGs irrespective of their host of isolation

(only comparing PGs with at least six tested strains, using 2-tailed, heteroscedastic T-tests,

Bonferroni corrected for seven total tests), and found that strains in PG2 were significantly more

virulent on bean than strains from PG1, PG3, and PG4 (p= 1.33e-07, 0.012, and 0.029

respectively), but not relative to PG6. In contrast, strains from PG3 were only significantly more

virulent on bean than strains from PG1 (p=0.006). No other significant pairwise PG comparisons

were observed. Interestingly, we noticed that PG2 non-bean isolates showed higher virulence

on bean than non-bean isolates from other PGs (p=2.87e-4). This indicates that PG2 isolates

225 show greater virulence on bean *irrespective* of host of isolation, although the degree of virulence

226 is relatively low. This pattern was reversed in PG3 where non-bean isolates had significantly

227 lower virulence than non-bean isolates from all other PGs (p=4.52e-3).

Table 1. Virulence and Germination Assay Summary

Group [1]	N Strains	Virulence [2] Mean	Median	Stdv	Germination Frequency (%) Mean	Median	Stdv
All Strains	121	0.789	0.913	0.271	60.10	63.33	24.31
PG1	8	0.972	0.995	0.043	72.99	79.17	15.71
PG2	50	0.658	0.748	0.320	46.74	53.33	26.41
PG3	42	0.841	0.964	0.218	68.15	67.50	16.53
PG4	6	0.924	1.000	0.147	83.61	82.50	4.52
PG5	3	0.966	1.000	0.059	73.33	78.33	10.14
PG6	3	1.000	1.000	0.000	80.56	81.67	3.47
PG7	2	0.956	0.956	0.062	85.00	85.00	7.07
PG11	6	0.851	0.867	0.151	43.89	50.00	20.75
PG13	1	1.000	1.000	NA	91.67	91.67	NA
All Bean Isolates	29	0.594	0.516	0.271	45.89	53.33	23.52
All Non-Bean	92	0.850	0.960	0.244	64.58	68.33	23.05
PG2 Bean	16	0.560	0.478	0.293	36.84	32.22	26.42
PG2 Non-Bean	34	0.704	0.841	0.327	51.41	56.67	25.47
PG3 Bean	13	0.635	0.606	0.246	57.02	58.89	13.21
PG3 Non-Bean	29	0.933	1.000	0.122	73.14	73.33	15.56

[1] PG = phylogroup

[2] Normalized fresh weight 2 weeks after seed infection

Germination screen. We then assessed whether the virulence of a strain also influenced the

germination frequency of bean seeds. Pathogenic microbes are known to interfere with the seed

germination both through the direct action of phytotoxins and the indirect action of immune

activation [23, 45, 46, 49, 50]. In fact, seedling growth inhibition is a well-established assay for

immune activation in *Arabidopsis thaliana [50]*. In general, the frequency distribution for bean

germination inhibition was less skewed than the frequency distribution for virulence, with mean

= 60.1%, median = 63.3%, and standard deviation = 24.3% (Fig 7, Table 1). The average

germination frequency for all bean isolates was 45.9% compared to 64.6% for non-bean isolates

(p=4.92e-4). When stratifying the bean isolates by PG, we found no significant difference in

germination frequency between PG2 bean and non-bean isolates (p=0.076), while the

comparison was significant for PG3 (p=0.002). However, in contrast to the virulence assays we

observed a slightly significant difference between germination frequency for PG2 bean isolates

and PG3 bean isolates (p=0.014). Other inter-phylogroup comparisons were similar to what was

found for the virulence assays, PG2 strains were significantly different from strains from PG1,

PG3, and PG4 (p=0.010, 6.23e-5, 8.53e-11 respectively 2-tailed, heteroscedastic T-test,

Bonferroni corrected for seven tests), while PG3 strain were also significantly different from PG4

(p=2.23e-4). Also similar to the virulence results, non-bean PG2 isolates resulted in a lower

germination frequency than all other non-bean isolates (p=9.60e-5), while non-bean PG3 strains

had a higher germination frequency than non-bean isolates from all other PGs (p=0.004),

although this pattern is absent when non-bean PG2 strains were removed from the analysis.

Finally, we measured the association between virulence (i.e., normalized fresh weight) and

germination frequency and found a strong association between the two metrics for the full

dataset (R^2=0.46, p=2.2e-16). Stratifying by PG and bean isolates showed a strong association

for PG2 bean isolates (linear regression; F = 28.91, df = 13, p=0.0001, R^2 = 0.68), but no

significant association for PG3 bean isolates (R^2=0.62, p=0.06) (Fig 8).

Predictive modelling of *P. syringae* virulence on bean. We used a gradient boosted decision

tree regression model to predict the virulence of *P. syringae* isolates on beans. The model used

one of three input feature classes: 1) genomic kmers; 2) T3SE kmers; or 3) presence / absence

of T3SEs and phytotoxins. T3SEs and phytotoxins are well-known virulence factors, with the

former often strongly associated with host specificity. Plant weight 14 days after seed infection

was used as the continuous outcome variable in our model. We could have also used seed

germination frequency in this assay but felt that plant fresh weight more accurately reflected the

virulence concerns of bean producers. The goal of analysis was to assess the power of machine

learning to predict disease outcomes based on genome sequences and to predict the host

specificity of new isolates based on their genome sequence.

We used two nested collections of strains to generate the model. The first collection was

comprised of the 121 of the isolates directly screened for virulence, which was made of 29 bean

isolates and 92 non-bean isolates, including 50 PG2 isolates (16 bean, 34 non-bean), and 42

PG3 isolates (13 bean, 29 non-bean). This collection is slightly smaller than the full screened

set since it does not include the additional PG3 bean isolates added to balance the

experimental design (q.v., materials and methods). The second collection was an expanded

strain set in which we imputed virulence values based on genomic similarity. The imputation

process involved identifying strains in our collection belonging to the same clonal lineage as

those assayed in our virulence screen (i.e., having a core genome evolutionary distance of less

than 0.001 and a T3SE Jaccard similarity of greater than 0.8 to a screened strain). Any strains

meeting these criteria were assigned the same virulence as the corresponding screen strain.

This imputation process almost tripled the size of our sample set, resulting in an expanded

collection of 320 strains, which was made of 59 bean isolates and 261 non-bean isolates,

including 66 PG2 isolates (19 bean, 47 non-bean), and 142 PG3 isolates (39 bean, 104 non-

bean). We also trained a model on PG2 and PG3 strains separately since bean isolates from

these PGs interact with their host very differently.

Our analysis revealed a significant linear relationship between sample size and the overall

performance of the models (Fig. 9, MAE linear regression $F = 5.89$, df = 10, p=0.03, $R^2 = 0.3$).

Hence, the data sets with the least number of samples consistently showed the poorest

performance regardless of the input data (e.g., PG2 and the screened collection). Furthermore,

our models achieved the highest predictive power when trained exclusively at isolates from

PG3, which may be explained by the strong clonal separation of bean vs. non-bean pathogens

within this PG which results in the statistical reinforcement of features with improved predictive

power during model training. Perhaps not surprisingly, models built using the greatest diversity,

with respect to number of strains (i.e., the expanded strain collection) and number of genetic

feature (i.e., whole genome kmers) showed the highest predictive power, with the best model

having a mean absolute error (MAE) of 0.06 (Table 2).

Table 2. Machine Learning Model Performance

Group	Samples	Dataset	MAE	RMSE
PG3	142	Whole genome kmers	0.049	0.107
PG3	142	Toxin and T3SE binary matrix	0.054	0.101
Imputed	320	Whole genome kmers	0.067	0.140
PG3	142	T3SE kmers	0.085	0.126
Imputed	320	Toxin and T3SE binary matrix	0.092	0.153
Imputed	320	T3SE kmers	0.096	0.177
PG2	66	T3SE kmers	0.154	0.233
Screened	121	Whole genome kmers	0.155	0.204
Screened	121	T3SE kmers	0.159	0.213

Screened	121	Toxin and T3SE binary matrix	0.163	0.201
PG2	66	Whole genome kmers	0.169	0.242
PG2	66	Toxin and T3SE binary matrix	0.202	0.262

Model functional validation. Finally, we evaluated the power of our gradient boosted decision

tree regression model on unseen strains by using the kmer profiles of *P. syringae* isolates not

previously examined to predict virulence and comparing these predictions to actual virulence

measures obtained through the seed infection virulence assay. Importantly, none of the strains

in the functional validation set were clonally related to any screened strain, i.e., having a core

genome evolutionary distance >0.001 and a T3SE Jaccard similarity of <0.8 compared to the

screened strains. When comparing actual to predicted virulence of the 16 strains in the

functional validation set, we found that 15 (94%) strains had virulence levels within the bounds

of estimated predictions given the calculated RMSE values (±0.20) (Fig. 10).

DISCUSSION

In this work we addressed whether host of isolation is a reliable predictor of host specific

virulence and whether whole genome sequences can be used to predict the host specific

virulence potential of individual strains. While host of isolation is a widely used surrogate for

host specificity, this assumption has rarely been empirically tested [24, 25], and the strength of

this assumption is critical when viewed from the context of the virulence potential of emerging

pathogens. Are strains isolated from one host only virulent on that host, or do they have the

potential to move to other species? Are strains isolated from environmental sources, such as

streams or soil, limited to those environments or can they 'jump' to a new host and potentially

cause a significant outbreak?

Our first aim was to determine if infection of bean seeds by *P. syringae* recapitulated virulence

responses seen in standard syringe inoculation virulence assays. We found a negative

association between our virulence measure of normalized plant fresh weight after seed infection

and *in planta* bacterial growth after syringe infiltration into leaf tissue, showing that the seed

infection protocol effectively recapitulates standard methods. This finding is consistent with

published and anecdotal reports that infected seed stocks are a significance source of bean

disease [45, 46, 51, 52]. In general, we found that bean isolates reduced mean plant fresh

weight by 30.2% and median weight by 46.2% compared to non-bean isolates. While the PG2

bean isolates (leaf spot disease caused by pathovar *syringae*) had a normalized mean fresh

weight of 0.56±0.293 (SD) compared to 0.64±0.246 for the PG3 bean isolates (halo blight

disease caused by pathovar *phaseolicola*), this difference was not significantly different. A

similar pattern was found when we examined seed germination frequencies, where bean

isolates reduced the average germination frequency by 28.9% and median frequency by 22.0%

compared to non-bean isolates, while the 38.8% mean germination frequency of PG2 bean

pathogens was significantly lower than the 57.0% mean germination frequency of the PG3 bean

pathogens (p=0.014).

We find a striking difference when comparing bean to non-bean isolates found in the same

phylogroup. As anticipated, bean seed infection with PG3 bean isolates resulted in significantly

higher virulence and lower germination frequency than PG3 non-bean isolates, while in contrast,

PG2 bean isolates did not differ significantly from PG2 non-bean isolates. PG2 strains generally

(irrespective of host of isolation) show greater virulence on bean, indicating that strains from this

phylogroup have lower host specificity, i.e., are host generalists. This is strongly supported

when comparing non-bean isolates from PG2 to non-bean PG3 isolates (normalized fresh

weight of 0.704 and 0.933, respectively; p= 4.67E-04). These findings are consistent with other

studies that have found lower levels of host specificity among PG2 strains [24, 25] and lends

support to the hypothesis that PG2 strains may rely as much or more on toxins than T3SEs

when compared to other *P. syringae* strains.

We expected that PG3 bean isolates would have higher virulence than PG2 bean isolates since

halo blight caused by PG3 pathovar *phaseolicola* is a much more severe disease than spot

disease caused by PG2 pathovar *syringae*, but this was not the case. There are several

explanations for these data. First, is a simple experimental bias explanation driven by the fact

that nearly all the PG3 bean isolates fall into one clonal group as defined by our clonality criteria

of a core genome distance of <0.001 average substitutions per site and T3SE profile Jaccard

similarity value >0.8. We attempted to address this issue by oversampling from the *phaseolicola*

clonal group. But to ensure that we did not create another bias by adding too many very closely

related strains, we only added seven additional strains to the original group of six PG3 bean

isolates. Unfortunately, this still resulted in a small set that could easily be skewed by a few

outlying measurements. Second, some of the PG3 strains likely elicit effector-triggered immunity

in the cultivar of bean assayed, which would result in healthy plants. Given the small set of PG3

bean isolates, even a few ETI-eliciting strains will result in a large average decrease in

virulence. And third, it is possible that the most severe symptoms of halo blight are only seen

after leaf-to-leaf transmission caused by water splash rather than seed transmission [14, 53].

While many genome-wide association studies have successfully identified strong genotype-to-

phenotype linkages, we were unable to identify any loci significantly associated with bean

isolation (data not shown). Consequently, we shifted our focus to machine learning approaches

as they can not only unravel genomic signatures associated with continuous phenotypes, but

also predict the virulence potential of previously unseen isolates given their genome sequences.

Regardless of PG affiliation, our model was able to predict the virulence of individual *P. syringae*

isolates with low error margins based solely on whole genome data. The fact that models

trained on virulence factors alone could predict virulence with considerable accuracy supports

the notion that T3SEs and phytotoxins play crucial roles in host adaptation processes.

Nonetheless, the higher predictive power of models trained with whole genome kmers suggests

that factors other than canonically virulence-associated genes also play important roles on

disease development and adaptation to beans.

Sample size is one of the most important contributors to accurate model generation in machine

learning. Simple models trained with large data sets can learn to predict outcomes with much

higher accuracy than complex models trained on smaller datasets. Our results also found a

correlation between sample size and model accuracy, with the screened strain collection

providing a MAE of 0.153 while the larger imputed strain collection increased model

performance to a MAE of 0.065. We also find poorer model performance for PG2 strains than

PG3 strains. While this may partly reflect the differences in sample size between these two

groups, it also likely reflects the underlying biology. The majority of bean isolates in PG3 are

phylogenetically clustered, while there is little clustering of bean isolates in PG2. Consequently,

the model may perform better on PG3 since it is essentially predicting phylogenetic structure.

Another contributing factor is likely the finding discussed above, namely, that host specificity

appears to be weaker in PG2. If PG2 strains are more generalists than specialists, then the host

specificity signal would be weaker and any model trying to find this signal would perform more

poorly.

MATERIALS AND METHODS

Strain collection. Three hundred and thirty-four *Pseudomonas syringae* strains were used in

this study (Table S1). Forty-six *P. syringae* isolates were collected from bean fields

approximately 80 km East of Lethbridge, Alberta, Canada, during the summer of 2012. Bean

leaves with symptoms of bacterial diseases were collected from new growth during the

vegetative growth stage. The remaining 288 isolates were previously published [9] and include

49 *P. syringae* type and pathotype strains [1, 35]. A type strain is the isolate to which the

scientific name of that organism is formally attached under the rules of prokaryote

nomenclature, while a pathotype strain is similar but with the additional requirement that it has

the pathogenic characteristics of its pathovar (i.e., a pathogen of a particular host) [5]. Out of the

334, 318 strains were used for comparative analyses and model training, while 16 were used for

model functional validation. A subset of 267 non-clonal representative strains (discussed below)

were selected for the predictive modeling to avoid clonal bias. A further subset of 121 isolates,

including the type and pathotype strains, were selected for virulence assays.

Sequencing and quality control. DNA was extracted using the Gentra Puregene Yeast and

Bacteria kit (Qiagen, Hilden, Germany). Illumina libraries with 300-400 bp inserts were

generated using the Illumina Nextera XT kit according to the manufacturer's protocol (Illumina,

CA, USA). Samples were multiplexed with the Illumina Nextera XT Index kit containing 96

indices. Samples were sequenced on the Illumina NextSeq 500 Mid Output v2 (300 cycle) kit

with 150 base PE reads. All sequencing was performed at the University of Toronto's Centre for

the Analysis of Genome Evolution and Function (CAGEF). Raw read quality was assessed with

FastQC. Trimmomatic was used to remove adapters and trim raw sequencing reads based on a

sliding window approach (window size = 4, required quality = 5).

***De novo* assembly.** Paired-end reads were *de novo* assembled using the CLC Genomics

Assembly Tool (CLC Genomics Workbench). Contigs shorter than 1kb were removed from the

assemblies. Low coverage contigs with matches to non-*Pseudomonas* genera and no matches

to the *Pseudomonas* genus that had a depth of coverage less than one standard deviation from

the average assembly coverage were deemed contaminants and, therefore, removed from the

final draft.

Pangenome analysis, gene prediction, annotation, and orthologous clustering. Gene

prediction and annotation for all assemblies were performed with Prokka [36]. Prokka annotates

inferred coding sequences by searching for sequence similarity in the UniProtKB [37] database

and HMM libraries [38]. Additionally, all predicted genes were aligned against a custom T3SE

database for the identification of potential T3SEs [19]. Pangenome analysis was performed via

PIRATE [39], which iteratively clusters genes into orthologous groups by performing all-vs-all

comparisons followed by MCL clustering given a certain percent identify threshold. Genes

present in at least 95% of the genomes were classified as core. Core protein families were

individually aligned with MUSCLE [40] and later concatenated into a single protein alignment.

We used the FastTree2 approximate maximum-likelihood approach [41] to infer the

phylogenetic relationships of all 320 isolates. Core genome synonymous substitution rates were

estimated with MEGA7 (41) using the Nei-Gojobori method and Jukes-Cantor model. Jaccard

distances were computed with R version 4.0.5 (42) using a binary matrix of presence and

absence of accessory genes. Rarefaction curves were generated using a custom Python script.

Identification of non-clonal representative strains. We reduced the impact of phylogenetic

bias in our predictive modeling by selecting only one representative strain from each clonal

group (i.e., very closely related strains recently derived from a common ancestor) identified from

the *P. syringae* core-genome phylogeny. We identified clonal groups by calculating the pairwise

core genome evolutionary distance and the Jaccard similarity for T3SE profiles. We found the

minimum pairwise core-genome evolutionary distance for isolates with identical T3SE profiles to

be 0.001 average substitutions per site. We therefore pooled the 318 isolates if they had a core

genome evolutionary distance of less than 0.001, resulting in 209 clusters. We further

supplemented these clusters by adding back any strain that had a T3SE profile Jaccard

similarity value less than 0.8, resulting in 267 non-clonal clusters. A single representative was

selected out of each of these non-clonal clusters for downstream analyses. One exception was

made to the strain selection process to balance our experimental design, which was skewed

due to the fact that the vast majority of PG3 bean isolates (i.e., pathovar *phaseolicola*) fall into

one clonal group. Initially, our selection criteria resulted in only six PG3 bean isolates compared

to 16 PG2 bean isolates (i.e., pathovar *syringae*). We therefore added an additional seven

phaseolicola strains to the screened set to better balance the number of bean isolates in PG2

and PG3. Evolutionary distances and Jaccard similarity scores were inferred with MEGA7 [42]

and R version 4.0.5 [43].

Seed infection virulence assay. *P. syringae* strains were grown overnight at 30°C in King's B

media, re-suspended in 10 mM $MgSO_4$ and diluted to an OD_{600} of 0.001. *P. vulgaris* var.

Canadian Red seeds were soaked for 24 hours in the bacterial suspension, planted in Sunshine

Mix 1 soil with regular watering and grown for 14 days. Plant fresh weight and germination

frequencies were measured and normalized to a control plant treated with 10 mM $MgSO_4$ sown

on each flat. Trials were repeated three times.

Syringe infiltration virulence assays. *P. syringae* strains were grown overnight on appropriate

antibiotics, re-suspended in 10 mM $MgSO_4$ and diluted to OD_{600} of 0.001. Two- to three-week-

old *Phaseolus vulgaris* var. Canadian Red plants were syringe infiltrated and bacterial growth

assays were carried out by harvesting eight leaf disks (1 cm^2) from each plant (two per each

primary leaf) three days after infiltration. Disks were homogenized using a bead-beater in 200 μl

sterile 10 mM $MgSO_4$, serially diluted in 96-well plates, and 5 μl from each dilution was spot

plated on KB supplemented with rifampicin for positive and negative control strains, and

rifampicin and kanamycin for strains harboring constructs. Plates were incubated for at least 24

hours at 30°C and the resulting colony counts were used to calculate the number of CFUs per

cm^2 in the leaf apoplast.

Predictive modeling of *P. syringae* virulence on bean. We used an implementation of

gradient boosted decision trees to model the effect of *P. syringae* isolates on plant weights as a

proxy for strain virulence using: 1) whole genome kmers, 2) T3SE kmers, and 3) a presence /

absence matrix of T3SEs and phytotoxins. We split sequences into 31-mers with fsm-lite and

generated a binary matrix for kmers with identical distribution patterns using custom python

scripts. Next, we used the Scikit-learn and the XGBoost python libraries [44] to generate a

regression model for the prediction of normalized plant weights using all three datasets as input

features. Given the relatively small size of our dataset, we used a cross-validation (CV)

procedure to assess the performance of our model on 50 independent splits. For each time, we

randomly split the data into training (80%) and testing (20%) sets while maintaining the same

plant weight distributions on both sets. Hyper parameters were fine-tuned using Scikit-learn's

RandomizedSearchCV module and regression models were generated with XGBoost's

XGBRFRegressor module.

REFERENCES

1.   **Baltrus DA, McCann HC, Guttman DS**. Evolution, genomics and epidemiology of

Pseudomonas syringae: Challenges in Bacterial Molecular Plant Pathology. *Mol Plant Pathol*

2017;18(1):152-168.

2.   **Morris CE, Monteil CL, Berge O**. The life history of *Pseudomonas syringae*: linking

agriculture to earth system processes. *Annu Rev Phytopathol* 2013;51:85-104.

3.   **Xin XF, Kvitko B, He SY**. *Pseudomonas syringae*: what it takes to be a pathogen. *Nat*

Rev Microbiol 2018;16(5):316-328.

4.   **Morris CE, Sands DC, Vinatzer BA, Glaux C, Guilbaud C et al.** The life history of the

plant pathogen *Pseudomonas syringae* is linked to the water cycle. *ISME J* 2008;2(3):321-334.

5.   **Bull CT, De Boer SH, Denny TP, Firrao G, Fischer-Le Saux M et al.** Demystifying the

nomenclature of bacterial plant pathogens. *J Plant Pathol* 2008;90(3):403-417.

6.   **Dye DW, Bradbury JF, Goto M, Hayward AC, Lelliott RA et al.** International standards

for naming pathovars of phytopathogenic bacteria and a list of pathovar names and pathotype

strains. *Review of Plant Pathology* 1980;59(4):153-168.

7.   **Berge O, Monteil CL, Bartoli C, Chandeysson C, Guilbaud C et al.** A user's guide to

a data base of the diversity of *Pseudomonas syringae* and its application to classifying strains in

this phylogenetic complex. *PLoS One* 2014;9(9):e105547.

8.   **Sarkar SF, Guttman DS**. Evolution of the core genome of *Pseudomonas syringae*, a

highly clonal, endemic plant pathogen. *Appl Environ Microbiol* 2004;70(4):1999-2012.

9.   **Dillon MM, Thakur S, Almeida RND, Wang PW, Weir BS et al.** Recombination of

ecologically and evolutionarily significant loci maintains genetic cohesion in the *Pseudomonas*

syringae species complex. *Genome Biol* 2019;20(1):3.

10.   **Hwang MS, Morgan RL, Sarkar SF, Wang PW, Guttman DS**. Phylogenetic

characterization of virulence and resistance phenotypes of *Pseudomonas syringae*. *Appl*

Environ Microbiol 2005;71(9):5182-5191.

11.   **Hulin MT, Armitage AD, Vicente JG, Holub EB, Baxter L et al.** Comparative genomics

of *Pseudomonas syringae* reveals convergent gene gain and loss associated with specialization

onto cherry (*Prunus avium*). *New Phytol* 2018;219(2):672-696.

12.   **Hulin MT, Mansfield JW, Brain P, Xu X, Jackson RW et al.** Characterization of the

pathogenicity of strains of *Pseudomonas syringae* towards cherry and plum. *Plant Pathol*

2018;67(5):1177-1193.

13.   **O'Brien HE, Thakur S, Gong Y, Fung P, Zhang J et al.** Extensive remodeling of the

Pseudomonas syringae pv. *avellanae* type III secretome associated with two independent host

shifts onto hazelnut. *BMC Microbiol* 2012;12:141.

14.   **Arnold DL, Lovell HC, Jackson RW, Mansfield JW**. *Pseudomonas syringae* pv.

phaseolicola: from 'has bean' to supermodel. *Mol Plant Pathol*, Research Support, Non-U.S.

Gov't

Review 2011;12(7):617-627.

15.   **Tsiamis G, Mansfield JW, Hockenhull R, Jackson RW, Sesma A et al.** Cultivar-

specific avirulence and virulence functions assigned to avrPphF in *Pseudomonas syringae* pv.

phaseolicola, the cause of bean halo-blight disease. *EMBO J* 2000;19(13):3204-3214.

16.   **Joardar V, Lindeberg M, Jackson RW, Selengut J, Dodson R et al.** Whole-genome

sequence analysis of *Pseudomonas syringae* pv. *phaseolicola* 1448A reveals divergence

among pathovars in genes involved in virulence and transposition. *J Bacteriol*

2005;187(18):6488-6498.

17.   **Feil H, Feil WS, Chain P, Larimer F, DiBartolo G et al.** Comparison of the complete

genome sequences of *Pseudomonas syringae* pv. *syringae* B728a and pv. *tomato* DC3000.

Proc Natl Acad Sci U S A 2005;102(31):11064-11069.

18.   **Marco ML, Legac J, Lindow SE**. *Pseudomonas syringae* genes induced during

colonization of leaf surfaces. *Environ Microbiol* 2005;7(9):1379-1391.

19.   **Dillon MM, Almeida RND, Laflamme B, Martel A, Weir BS et al.** Molecular evolution

of *Pseudomonas syringae* type III secreted effector proteins. *Front Plant Sci* 2019;10:418.

20.   **Buttner D**. Behind the lines-actions of bacterial type III effector proteins in plant cells.

FEMS Microbiol Rev 2016;40(6):894-937.

21.   **Khan M, Seto D, Subramaniam R, Desveaux D**. Oh, the places they'll go! A survey of

phytopathogen effectors and their host targets. *Plant J* 2018;93(4):651-663.

22.   **Martel A, Ruiz-Bedoya T, Breit-McNally C, Laflamme B, Desveaux D et al.** The ETS-

ETI cycle: evolutionary processes and metapopulation dynamics driving the diversification of

pathogen effectors and host immune factors. *Curr Opin Plant Biol* 2021;62:102011.

23.   **Bender CL, Alarcon-Chaidez F, Gross DC**. *Pseudomonas syringae* phytotoxins: mode

of action, regulation, and biosynthesis by peptide and polyketide synthetases. *Microbiol Mol Biol*

Rev 1999;63(2):266-292.

24.   **Morris CE, Moury B**. Revisiting the concept of host range of plant pathogens. *Annu Rev*

Phytopathol 2019.

25.   **Morris CE, Lamichhane JR, Nikolić I, Stanković S, Moury B**. The overlapping

continuum of host range among strains in the *Pseudomonas syringae* complex. *Phytopathology*

Research, journal article 2019;1(1):4.

26.   **Allen JP, Snitkin E, Pincus NB, Hauser AR**. Forest and Trees: Exploring Bacterial

Virulence with Genome-wide Association Studies and Machine Learning. *Trends Microbiol*

2021.

27.   **Chen PE, Shapiro BJ**. The advent of genome-wide association studies for bacteria.

Curr Opin Microbiol 2015;25:17-24.

28.   **Falush D**. Bacterial genomics: Microbial GWAS coming of age. *Nat Microbiol*

2016;1:16059.

29.   **Falush D, Bowden R**. Genome-wide association mapping in bacteria? *Trends Microbiol*,

Review 2006;14(8):353-355.

30.   **Power RA, Parkhill J, de Oliveira T**. Microbial genome-wide association studies:

lessons from human GWAS. *Nat Rev Genet* 2017;18(1):41-50.

31.   **Sheppard SK, Didelot X, Meric G, Torralbo A, Jolley KA et al.** Genome-wide

association study identifies vitamin B5 biosynthesis as a host specificity factor in

Campylobacter. Proc Natl Acad Sci U S A 2013;110(29):11923-11927.

32.   **Earle SG, Wu CH, Charlesworth J, Stoesser N, Gordon NC et al.** Identifying lineage

effects when controlling for population structure improves power in bacterial association studies.

Nat Microbiol 2016;1:16041.

33.   **Nicholls HL, John CR, Watson DS, Munroe PB, Barnes MR et al.** Reaching the end-

game for GWAS: Machine learning approaches for the prioritization of complex disease loci.

Front Genet 2020;11:350.

34.   **San JE, Baichoo S, Kanzi A, Moosa Y, Lessells R et al.** Current affairs of microbial

genome-wide association studies: approaches, bottlenecks and analytical pitfalls. *Front

Microbiol* 2019;10:3119.

35.   **Thakur S, Weir BS, Guttman DS**. Phytopathogen Genome Announcement: draft

genome sequences of 62 *Pseudomonas syringae* type and pathotype strains. *Mol Plant Microbe

Interact* 2016;29(4):243-246.

36.   **Seemann T**. Prokka: rapid prokaryotic genome annotation. *Bioinformatics

2014;30(14):2068-2069.

37.   **Apweiler R, Bairoch A, Wu CH, Barker WC, Boeckmann B et al.** UniProt: the

Universal Protein knowledgebase. *Nucleic Acids Res* 2004;32(Database issue):D115-119.

38.   **Mistry J, Finn RD, Eddy SR, Bateman A, Punta M**. Challenges in homology search:

HMMER3 and convergent evolution of coiled-coil regions. *Nucleic Acids Res* 2013;41(12):e121.

39.   **Bayliss SC, Thorpe HA, Coyle NM, Sheppard SK, Feil EJ**. PIRATE: A fast and

scalable pangenomics toolbox for clustering diverged orthologues in bacteria. *Gigascience*

2019;8(10).

40.   **Edgar RC**. MUSCLE: a multiple sequence alignment method with reduced time and

space complexity. *BMC Bioinformatics* 2004;5:113.

41.   **Price MN, Dehal PS, Arkin AP**. FastTree 2--approximately maximum-likelihood trees

for large alignments. *PLoS One* 2010;5(3):e9490.

42.   **Kumar S, Stecher G, Li M, Knyaz C, Tamura K**. MEGA X: Molecular Evolutionary

Genetics Analysis across Computing Platforms. *Mol Biol Evol* 2018;35(6):1547-1549.

43.   **R Development Core Team**. R: A language and environment for statistical computing.

R Foundation for Statistical Computing, Vienna, Austria; 2020. https://www.R-project.org/.

44.   **Chen T, Guestrin C**. XGBoost: A scalable tree boosting system. *Proceedings of the*

22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining series

vol. KDD '16). San Francisco, California, USA: ACM; 2016.

45.   **Darrasse A, Bureau C, Samson R, Morris CE, Jacques M-A**. Contamination of bean

seeds by *Xanthomonas axonopodis* pv. *phaseoli* associated with low bacterial densities in the

phyllosphere under field and greenhouse conditions. *Eur J Plant Pathol* 2007;119(2):203-215.

46.   **Eyster HC**. The cause of decreased germination of bean seeds soaked in water. *Am J*

Bot 1940;27(8):652-659.

47.   **Hirano SS, Demars SJ, Morris CE**. Survival, establishment, and dispersal of

Pseudomonas s*yringae* on snap beans (*Phaseolus vulgaris* L). *Phytopathology*, Meeting

Abstract 1981;71(8):881-881.

48.   **Willis DK, Hrabak EM, Rich JJ, Barta TM, Lindow SE et al.** Isolation and

characterization of a *Pseudomonas syringae* pathovar *syringae* mutant deficient In lesion

formation on bean. *Mol Plant Microbe Interact* 1990;3(3):149-156.

49.   **Chahtane H, Nogueira Füller T, Allard PM, Marcourt L, Ferreira Queiroz E et al.** The

plant pathogen *Pseudomonas aeruginosa* triggers a DELLA-dependent seed germination arrest

in *Arabidopsis. Elife* 2018;7.

50.   **Bredow M, Sementchoukova I, Siegel K, Monaghan J**. Pattern-triggered oxidative

burst and seedling growth inhibition assays in *Arabidopsis thaliana. J Vis Exp* 2019(147).

51.   **Shade A, Jacques MA, Barret M**. Ecological patterns of seed microbiome diversity,

transmission, and assembly. *Curr Opin Microbiol* 2017;37:15-22.

52.   **Upper CD, Hirano SS, Dodd KK, Clayton MK**. Factors that affect spread of

Pseudomonas syringae in the phyllosphere. *Phytopathol* 2003;93(9):1082-1092.

53.   **Butterworth J, McCartney HA**. The dispersal of bacteria from leaf surfaces by water

splash. *Journal of Applied Bacteriology* 1991;71(6):484-496.

FIGURES AND CAPTIONS

Figure 1: Distribution of bean isolates across the *Pseudomonas syringae* core genome

phylogeny. Core genome phylogeny of 320 *P. syringae* isolates. Clade numbers represent

phylogroups. Metadata row 1 shows strains isolated from diseased bean plants.

Figure 2: Core and accessory genome diversity. Comparison of (A) core genome

synonymous substitution rate (Ks) and (B) accessory genome Jaccard distance for 21 PG2

bean isolates, 36 PG3 bean isolates, and all 66 isolates from PG2 and 142 isolates from PG3.

646   **Figure 3: Rarefaction curves for the core and accessory genomes.** Families present in 95%

647   (soft core genome) of *P. syringae* isolates exponentially decay as each new genome is added to

648   the analysis. The total number of gene families identified continues to increase indefinitely with

649   the addition of new genomes when singletons (families only present in one isolate) are included.

A

R² = 0.63, p = 4.95e-06

B

Figure 4: Correlation between bacterial load following pressure infiltration of mature

bean leaves and plant weight following seed infection. (A) Bacterial density in 14 day old

infected bean leaves (pressure infiltration 3 days post infection) as a function of normalized

median weight of seed infected plants at 14 days old. There is a strong negative correlation

between bacterial density and plant weight across 24 *P. syringae* isolates (linear regression; $F =$

36.95, df = 21, $p =$ 4.95e-06, $R^2 =$ 0.62). (B) Characteristic plant phenotypes of 14 day old plants

following seedling infection. The photos on the left are for plants infected with $MgSO_4$, while the

photos on the right shows plants infected with the bean pathogen Psy0540.

Tree scale: 0.1

Phylogroup

■ PG1
■ PG2
■ PG3
■ PG4
■ PG5
■ PG6
■ PG7
■ PG11
■ PG13

Weight

Figure 5: Core genome phylogeny of 320 *P. syringae* isolates. Leaf colors represent PG

affiliation as shown in the legend. Metadata row 1 shows bean isolates in black and non-bean

isolates in gray. Metadata row 2 shows normalized median bean weight after seed infection by

the respective isolates. Green indicates a high weight, while yellow indicates a low weight as

show in the legend. Metadata row 3 shows normalized median bean weight for the expanded

strain collection based on phylogenetic imputation.

Figure 6: Virulence stratified by host and phylogroup. (A) Boxplots showing the distribution

of virulence (i.e., normalized plant weight 14 days after seed infection) ordered by the median

virulence. Colors correspond to phylogroups as shown along the top. Bean isolates are

indicated with asterisks above their respective boxplots. (B) Frequency plot of virulence for the

121 screened *P. syringae* strains. Distribution of virulence values for (C) bean verses non-bean

isolates, (D) bean and non-bean isolates stratified by phylogroup, and (E) virulence values

stratified by phylogroup (PG) for the set of 121 screened strains.

Figure 7: Germination frequencies by host and phylogroup. (A) Boxplots showing the

germination frequencies frequency for 121 screened *P. syringae* strains with strains presented

in the same manner and order as in Fig. 6. (B) Frequency plot of germination for the 121

screened strains. Distribution of germination frequencies for (C) bean verses non-bean isolates,

(D) bean and non-bean isolates stratified by phylogroup, and (E) virulence values stratified by

phylogroup (PG) for the set of 121 screened strains.

Figure 8: Correlation between germination frequency and virulence of seed infected bean plants. (A) Germination frequency vs virulence (i.e., normalized fresh weight) stratified by PG2

and PG3 bean isolates. A strong correlation is found between germination frequency and

virulence of bean strains from PG2 (linear regression; $F = 28.91$, df $= 13$, $p = 0.00012$, $r^2 =$

0.66). (B) Germination frequency vs. virulence stratified by PG irrespective of host.

Figure 9: Performance of supervised machine learning models on virulence predictions

with genome data. Models trained with the screened collection (yellow), imputed expanded

collection (blue), imputed expanded PG2 set (orange), and imputed expanded PG3 set (green).

Diamonds, circles, and triangles represent models trained with a phytotoxin and T3SE binary

matrix, T3SE kmers, and whole genome kmers, respectively. (A) Mean Absolute Error (MAE) as

a function of number of samples across 50 cross-validation splits. Increasing the number of

samples seen by the model significantly increases model performance (linear regression; $F =$

5.89, df = 10, $p = 0.03$, $r^2 = 0.37$). (B) Root Mean Square Error (RMSE) as a function of number

of samples across 50 cross-validation splits. Models trained on whole genome kmers with a

higher number of samples consistently outperform other models.

Figure 10: Correlation between predicted and observed weights of plants infected with

isolates previously unseen by the model. (A) Predicted virulence as a function of observed

virulence on seed infected bean plants. Color coding indicates PGs. Dashed line indicates 1:1

712   relationship. (B) Virulence for each isolate tested ordered by predicted virulence and colored by

713   PG. The solid line represents virulence predictions for each isolate. The grey box represents the

714   error margins for the predictions based on the MAE and RMSE values for the model.

SUPPORTING INFORMATION CAPTIONS

Figure S1: Sample size selection via a simulated experimental setup. Bean plants were seed infected with 6 *P. syringae* isolates with a high number of replicates (>50). Plant weights were randomly selected according to various replicate sizes (8-32). Our ability to distinguish pathogens from non-pathogens using Tukey-HSD tests plateaus at 20 replicates per treatment.

Figure S2: Mean Absolute Error (MAE) and Root Mean Square Error (RMSE) across 50 cross-validation splits. Model performance in terms of (A) MAE and (B) RMSE. Plant weight distributions were kept the same across splits.

Figure S3: Distribution of virulence values stratified by host and phylogroup. Boxplots showing the distribution of virulence (i.e., normalized plant weight 14 days after seed infection) values for (A) bean verses non-bean isolates for the set of 121 screened strains, and (B) for the 320 strains in the expanded dataset that includes both screened and imputed strains. (C) Distribution of virulence values for bean and non-bean isolates stratified by phylogroup for the screened strains and (D) for the expanded dataset. (E) Distribution of virulence values stratified by phylogroup (PG) for the screened strains, and the (F) expanded strain set.

Table S1: List of *P. syringae* isolates used in this study.

737   Supplement Figure 1

741   Supplement Figure 2

745   Supplement Figure 3

Screened   **Imputed**