Deciphering signatures of natural selection via deep learning

Xinghu Qin[1*#], Charleston W. K. Chiang[2], Oscar E. Gaggiotti[1*]

[1] Centre for Biological Diversity, Sir Harold Mitchell Building, University of St Andrews, Fife, KY16 9TF, UK

[2] Center for Genetic Epidemiology, Keck School of Medicine & Department of Quantitative and Computational Biology, University of Southern California, USA

Current Address: CAS Key Laboratory of Genomics and Precision Medicine, Beijing Institute of Genomics, Chinese Academy of Sciences & China National Center for Bioinformation, Beijing, 10010, China.

* Correspondence should be addressed to Xinghu Qin (qinxh@big.ac.cn) and OE Gaggiotti (oeg@st-andrews.ac.uk)

Abstract

Identifying genomic regions influenced by natural selection provides fundamental insights

into the genetic basis of local adaptation. We propose a deep learning-based framework,

DeepGenomeScan, that can detect signatures of local adaptation. We demonstrate that

DeepGenomeScan outperformed PCA and RDA-based genome scans in identifying loci

underlying quantitative traits subject to complex spatial patterns of selection. Noticeably,

DeepGenomeScan increases statistical power by up to 47.25% under non-linear

environmental selection patterns. We applied DeepGenomeScan to a European human

genetic dataset and identified some well-known genes under selection and a substantial

number of clinically important genes that were not identified using existing methods.

Keywords: deep learning, genome scan, genome-wide association studies, signatures of

natural selection

Background

One of the main challenges of modern biology is to dissect and understand the molecular

basis for naturally occurring phenotypic variation. Addressing this challenge is of

fundamental importance not only for the field of evolutionary biology but also for a wide

variety of applied fields involving human diseases, improvement of agricultural crops and

breeds and biodiversity conservation.

The recent increases in genomic data generated by modern sequencing technologies have

advanced our understanding of how natural selection, and its interactions with other

evolutionary forces, shape the genome and phenotype of species. Such technologies have

now been applied to the study of a wide range of species, but the statistical methods used to

analyse them tend to differ depending on whether the focus is on humans and domesticated

species or on wild species. In the case of humans and domesticated species, there is a

particular interest in uncovering associations between well-defined phenotypes (traits

associated with diseases or, in the particular case of domesticated species, yield) and the

genomic variants that underlie them. The standard approach, in this case, is to use Genome

Wide Association Studies (GWAS; e.g. [1]), which linearly model the additive allelic effect

of genotypes (the explicative variable) on phenotypes (the dependent variable). All of the

recent methods can take into account population stratification, using for example linear

mixed models that incorporate the genetic relationship matrix (GRM) as a random effect (e.g.

[2]). GWAS have been used to identify variants associated with a wide range of phenotypic

traits [3] and their results are used by several studies aimed at detecting polygenic adaptation

[4-8].

In the case of evolutionary studies and in particular those focused on wild species, the most

commonly used approaches are a suite of methods known as the genome-scans [9], which

ignore phenotypic traits. Many of them are population-based and use summary statistics to

uncover genomic regions exhibiting outlier behaviour. A very popular approach is to identify

loci that exhibit unusually strong among-population genetic differentiation. This approach

assumes that, in the absence of selection, differences in allele frequencies among populations

should follow a neutral distribution driven by migration and genetic drift. Any locus that

departs from this expectation is considered a good candidate for being under the effect of

natural selection. This approach initially proposed by Lewontin & Krakauer [10] has been

implemented in popular Bayesian methods (e.g. [11, 12]). More recently, there has been an

increased interest in establishing an association between a locus outlier behaviour and

environmental variables, with the assumption that such variables could represent selective

pressures acting upon genomic regions linked to the outlier loci [13-16]. Some of these

methods use raw genotypic data instead of summary statistics (e.g. [12]), but as opposed to

GWAS, they use linear mixed models that consider the genotype as the dependent variable,

and the environmental factor as the explicative variable.

Despite this apparent methodological dichotomy between studies focused on humans and

domesticated species and those focused on wild species, it is clear that a thorough

understanding of how natural selection shapes the phenotype and genome of species requires

some combination of these two different approaches. Indeed, the selective pressures exerted

by the heterogenous abiotic and biotic environment act upon individuals' phenotypes in a

complex way and lead to changes in the spatial structuring of variation in the genomic

regions underlying them. Thus, in order to fully characterise the action of natural selection,

we need to answer three fundamental questions: (i) what are the environmental drivers of

natural selection? (ii) what are the phenotypic traits upon which selective pressures act? and

(iii) what are the genomic regions underlying those adaptive traits?

In the last few years, there has been an increasing interest in the application of machine

learning approaches in population genomics [17-19] and GWAS studies [20, 21] but no

single method applicable to both has been proposed. Here we present DeepGenomeScan, a

unified deep learning approach for genome scan and GWAS, which can be used to answer

questions (i) and (iii) above (question (ii) would require the use of a quantitative genetics

method). The rationale underlying our method is that we can use the genotype of an

individual to predict not only its phenotype but also some attribute of the habitat where they

are sampled. This framework is akin to GWAS, but in our case, the response variable can be

an individual phenotypic trait, an environmental variable associated with its habitat, the

geographic location of the individual (latitude and longitude), or a variable describing spatial

genetic structure (e.g. reduced features from a dimensionality reduction technique such as

PCA).

Our approach leverages the power of deep neural networks to approximate arbitrarily

complex functions linking dependent and explicative variables [22], as well as recent

algorithms for model optimisation [23], and for inferring the importance of each explicative

variable in predicting the dependent variable [24]. It is important to note that as opposed to

the prevalent use of neural networks, i.e. prediction and pattern recognition, here our main

interest is in estimating the features (loci) that contribute the most to the predictive power of

the neural network. To achieve this goal, we use the concept of "feature importance" [24],

which represents a proxy for the effect size of any given locus. In essence, our method

estimates the effect size of genetic variants that explain a particular trait (phenotype or

environmental variable), and identifies those with outlier values as pinpointing a QTL or a

signature of natural selection.

An important advantage of our method when applied to spatial data is that it can consider any

spatial selection pattern including the usually assumed linear environmental gradient as well

as arbitrarily complex non-linear spatial patterns and, importantly, coarse-grained

heterogeneous selection with no clear spatial pattern. This represents an important advance as

existing methods only consider linear or monotonic non-linear patterns. In this particular

application, our approach generalises Yang et al.'s [25] idea of using geographic positioning

of individuals to identify loci that exhibit particularly steep slopes of allele frequency change

associated with recent positive selection. Our generalisation is two-fold: (i) instead of only

considering monotonic spatial gradients, we can detect loci associated with arbitrary and non-

monotonic selection patterns, and (ii) the dependent variable can be the geographic location

but also a phenotypic trait or an environmental factor.

In what follows, we introduce our method and evaluate its performance focusing on genome-

scan applications under spatially complex selection scenarios, but we also present a

preliminary evaluation of DeepGenomeScan performance as a tool to carry out GWAS (see

Discussion and Supplementary Material).

Results

Underlying Rationale for the Deep Learning Approach

A neural network can be considered as a generalised regression approach used to learn

complex functions expressing the association between the input data and a response variable

[22, 26]. In our case, the input or predictor variables are the genotypes of the individuals and

the response variables are observed phenotypes, environmental values, geographic locations

or reduced features describing genetic structure. In what follows, we use the term traits to

refer to these response variables.

As opposed to standard regression, which fits a fixed function to the data, a neural network

learns the (non-linear) function most appropriate for the data at hand. Here we use a

multilayer perceptron (MLP), which is composed of up to three layers of nodes (neurons)

with connections between layers but not between nodes within a layer (Fig. 1 and

Supplemental Material). Connection edges between nodes can have different weights and the

main goal of the MLP is to learn the weights that best describe the complex function that

associates inputs and outputs [27-29] by minimising the difference between the predicted and

observed traits. Once the optimal network is found, the weights connecting each input node

with the output node can be used to devise a test to identify the loci that contribute the most

to the trait under consideration. The underlying rationale for this test is that the absolute

values of weights associated with the path linking an input node (i.e. a locus) with the output

can be used to estimate the importance of an input variable [30, 31] (node or locus), which in

turn is closely associated with the effect size of a genetic variant. Loci with extreme

importance values are considered as outliers and good candidates for being under the

influence of natural selection. Thus, the approach we implement test the null hypothesis that

the importance (effect size) of a SNP is zero. Full details of our approach are provided in

Methods and Supplementary Material.

Genotype matrix   Multilayer Perceptron   Manhattan Plot

Input layer   Hidden layer 1   Hidden layer 2   Hidden layer 3   Output layer

Fig. 1. DeepGenomeScan framework. The first (input) layer receives the genotype matrix so

that each of its input nodes contains the genotypes of all sampled individuals at a single

locus. The last (output) layer contains the predicted trait values. Between these two layers

there is one or more hidden layers containing nodes that compute a non-linear transformation

of the previous layer outputs. Thus, the first hidden layer will transform the input data and

feed a signal to the second hidden layer, which in turn will apply a transformation and feed

the resulting signal to the third hidden layer, and so on and so forth until the last hidden layer,

which will carry out a final transformation and feed the results (the predicted traits) to the

output layer. After model training, optimization and hyperparameter tuning, the weights

connecting each input node with the output are used to calculate p-values that are then used

to produce a Manhattan plot. Note that all edges of the graph have associated weights but

here we include only some of them to avoid cluttering the figure.

Simulation Study

We evaluated the performance of our method using simulated data generated by Capblancq et

al.[18] (datasets are available from the Dryad Digital Repository at:

https://datadryad.org/review?doi=doi:10.5061/dryad.1s7v5). The simulations assume a two-

dimensional stepping-stone scenario and consider three quantitative traits, each coded by a

distinct set of 10 loci (QTLs), resulting in a total of 30 causal SNPs, out of a total of 1000.

Trait values are calculated simply as the sum of genotypic values of the causal loci. Each

quantitative trait (QT) is influenced by environmental factors having a distinct spatial pattern.

QT1 is influenced by factors with a quadratic spatial gradient, QT2 is influenced by factors

with a linear gradient, and QT3 is influenced by factors with a coarse and patchy spatial

pattern. There is a total of 10 environmental factors falling into one of these three categories.

Further details of the simulation approach are provided in Supplementary. Full details are

given in ref. [18].

We compared the performance of our method with those of two recently proposed machine

learning-based approaches: pcadapt [32] and RDA [18]. Figure 2 presents these results in

terms of power (proportion of true positives), False Discovery Rate (FDR) and False Positive

Rate (FPR or type I error). Using a threshold p-value $= 10^{-8}$, as typically done in GWAS, our

method has high power to detect all QTL types with a small FDR (0.117) and FPR (0.003).

On the other hand, pcadapt and RDA have no power to detect loci associated with QTs 1 and

3, and very low power to detect those associated with QT2 (although they have very low

error rates; Figs. 2a-c). QQ plots (Supplementary Material, Fig. S3) suggest using a threshold

of 0.001 for pcadapat and RDA and a threshold of up to 10^{-10} for DeepGenomeScan. Using a

less stringent threshold for pcadapt and RDA (p-value = 0.001) but keeping p-value $= 10^{-8}$ for

DeepGenomeScan still leads to a better performance of our method in terms of power when

compared to the other two methods (Fig. 2d) and also in terms of FDR and FPR when

compared to pcadapt (Fig. 2e-f). Increasing the stringency of the test for DeepGenomeScan to

the p-value $= 10^{-10}$ lowers the FDR of our method, making it similar to that of RDA while

still having the highest power to detect QTL1 and QTL3 (Fig. 2g-i). Overall, all three

methods have high power to detect loci associated with QT2, which is influenced by a linear

selective gradient. However, DeepGenomeScan with a threshold p-value $=10^{-10}$ was the best

for detecting QT1 and QT3 loci, which are influenced by non-linear selection patterns (Fig.

2g). In all cases, DeepGenomeScan outperforms both PCA and RDA in terms of statistical

power while controlling the type I error. In particular, DeepGenomeScan increases statistical

power, on average, by up to 47.25% compared with pcadapt, and up to 18.35% compared to

RDA under non-linear environmental gradient selections (QT1 and QT3). Similar results

were obtained when we used q-value thresholds (Supplementary Material, Figs. S4-5).

Fig. 2. Performance of pcadapt, RDA and DeepGenomeScan (DNN) under different p-value thresholds. (a-c): p-value = 10^{-8} for all three methods; (d-f) p-value = 0.001 for pcadapt and RDA and p-value = 10^{-8} for DeepGenomeScan; (g-i) p-value = 0.001 for pcadapt and RDA and p-value = 10^{-10} for DeepGenomeScan. Error bars indicate the standard deviation and letters are used to indicate the statistical significance of the difference in performance based on a t-test with p-value < 0.01. Panels a, d, g, j present the power to identify loci underlying

each quantitative trait separately (QT1-3). All other panels present the overall FDR and FPR

(across all three types of QTLs). All estimates were based on 100 simulated datasets.

Figure 3 presents Manhattan plots for one simulated data set with a somewhat permissive

threshold for pcadapt and RDA (0.001) and a more stringent threshold for DeepGenomeScan

(10^{-10}). This figure illustrates the advantage of DeepGenomeScan to detect outlier loci.

Clearly, very discontinuous selection patterns (i.e., QTL3) represent a big challenge for all

methods but DeepGenomeScan could still detect several of the outliers under this difficult

scenario.

Fig. 3. Manhattan plots of the results obtained with (a) pcadapt, (b) RDA and (c)

DeepGenomeScan (DNN) for one simulated data set. Note that the scale of the y axis in c

differs from that used for a and b. The threshold for pcadapt and RDA is set to 0.001 and a

threshold for DeepGenomeScan is set to 10^{-10}.

Application to a real data set

A common and longstanding framework to study the action of natural selection is to focus on

clinal variation in phenotypic traits or allele frequencies along environmental gradients [33].

One approach to detect genomic regions under selection in these clinal variation scenarios is

to identify loci exhibiting extreme frequency gradients across geographic space [25]. The

underlying assumption of this approach is that the environmental gradient is continuous,

leading to a monotonic but not necessarily linear change in allele frequency or phenotype.

Although there may be several examples of such geographic variation, spatial patterns in

selective pressures and the associated allele frequency can be non-monotonic. For example, it

is possible that the maximum allele frequency is located in the middle of the geographic

region under study [25]. As our simulations show, our approach is capable of identifying

genomic regions subject to such non-linear selection patterns.

In principle, approaches focused on clinal spatial patterns require the geographic coordinates

of each individual sample. However, sometimes this information is not available. A potential

solution to this problem is to carry out a Principal Component Analysis on the genotype

matrix and use the first two PC axes as surrogates for geographic coordinates [34, 35]. This

approach, however, is based on a linear combination of genotypes and can lead to poor

inference of spatial locations of admixed individuals [25]. A recently developed model-based

approach can be used to overcome this issue [25] but it requires the assumption of a smooth

monotonic function to describe allele frequency behaviour as a function of geographic

location, which may not be appropriate in all cases. Our implementation of

DeepGenomeScan allows the use of a more general dimensionality reduction approach,

Kernel Local Discriminant Analysis of Principal Components (KLFDAPC [36];

Supplementary Material) when spatial coordinates are not available.

Here we use a human dataset to determine if our approach can uncover new regions of the

human genome that may be under the influence of non-linear spatial selection patterns. We

applied DeepGenomeScan to the Population Reference Sample [37] (POPRES) dataset using

the known geographic coordinates and then using the two first reduced features from a

KLFDAPC analysis. POPRES represents an excellent example of human genetic variation

aligning with geography [35]. The dataset contains a total of 3,192 European individuals

genotyped at 500,568 loci using the Affymetrix 500K SNP chip. Details about data filtering

steps are provided in Online Materials.

We calculated p-values for each SNP (see Online Methods for details) and obtained a

genomic inflation factor GIF = 1 for this data set. Using a threshold p-value = 10^{-10} for

identifying the outlier loci, the first analysis using geographic coordinates as the response

variables detected 122 outlier SNPs located within 33 known genes. The full list of genes we

identified, and their chromosome positions are provided in Supplementary Table S1.

Consistent with previous widely reported regions under selection, our method detects strong

signals at the LCT region on chromosome 2, the ADH1C region on chromosome 4, the HLA

region on chromosome 6, as well as the OCA2 and the HERC2 region on chromosome 15

(Fig. 4; Supplementary Table S1).

Besides the already known selected genes, our method also detects some disease-related

genes that exhibited extreme variation across geographic space but were not identified by

existing methods (c.f. Supplementary Table 4 in ref. [25]); these include MGAT5,

TMEM163, ACMSD, CCNT2, MAP3K19, R3HDM1, UBXN4, MCM6, DARS1, EHMT2,

and CFB (Supplementary Table 1). For example, MCM6 is a regulatory element that controls

the expression of the LCT gene. The MGAT5 gene (alpha-1,6-mannosylglycoprotein 6-beta-

N-acetylglucosaminyltransferase), is one of the most important enzymes involved in the

regulation of the biosynthesis of glycoprotein oligosaccharides and is associated with

invasive malignancies and sclerosis [38-40] as well as visceral fat in women [41]. The

TMEM163 gene (transmembrane protein 163), is associated with Parkinson's disease,

ischemic stroke and coronary artery disease [42, 43]. The ACMSD gene

(aminocarboxymuconate semialdehyde decarboxylase) is associated with Parkinson's disease

[42] and childhood obesity [44]. There are several other outlier SNPs detected by our method

on chromosome 2, 4, 6, 10, and 11 (Supplementary Table 1) that are not within known

functional genes. However, they may be linked to regulatory genes and, therefore, of interest

for human genetic studies.

Fig. 4. Spatial genetic structure of European populations and signals of selection detected by

DeepGenomeScan. (A). Spatial genetic structure of European populations inferred via

KLFDAPC with a σ = 5. (B). Manhattan plot indicating loci under spatial selection obtained

from DeepGenomeScan using geographic coordinates as the response variables. (C).

Manhattan plot indicating loci under spatial selection obtained from DeepGenomeScan using

inferred spatial genetic structure (the first two reduced features of KLFDAPC) as the

response variables. The outliers in B-C were identified with a threshold of p-value $=10^{-10}$.

The top hits are shown in panels b and c with their dbSNP Reference ID (rs ID). Country

abbreviations: AL, Albania; AT, Austria; BA, Bosnia-Herzegovina; BE, Belgium; BG,

Bulgaria; CH, Switzerland; CY, Cyprus; CZ, Czech Republic; DE, Germany; ES, Spain; FR,

France; GB, United Kingdom; GR, Greece; HR, Croatia; HU, Hungary; IE, Ireland; IT, Italy;

KS, Kosovo; MK, Macedonia; NO, Norway; NL, Netherlands; PL, Poland; PT, Portugal; RO,

Romania; RS, Serbia and Montenegro; RU, Russia; Sct, Scotland; SE, Sweden; TR, Turkey;

YG, Yugoslavia. A full list of outlier loci can be found in Supplementary Table 1-2.

In the case of the analysis for which we replaced the original geographic coordinates with the

first two reduced features obtained from KLFDAPC, we first calculated the correlation

between the first two reduced features and the original geographic coordinates. We compared

these results to those of a similar analysis using the first two PC axes obtained from a PCA of

the genotype matrix. The results confirm that KLFDAPC provides better estimates of

geographic location than PCA (see Fig. 4a and Supplementary Material Fig. S8).

The genome scan based on the first two KLFDAPC reduced features identifies a somewhat

smaller number of outlier SNPs (116 loci, in 34 known genes) than when using geographic

coordinates (Fig. 4b-c and Supplementary Table S2). However, 88% of the outliers detected

in this analysis were also identified by the analysis based on geographic coordinates (102

shared loci). This analysis also detected well-known selected genomic regions (LCT, HLA,

ADH1, HERC2; Supplementary Table S2). However, there were 6 genes that were not

identified when using latitude and longitude (Supplementary Table S3). This group include

genes associated with cancer (e.g., AK5), diabetes mellitus (e.g., HSPA1L, HCG26, BAG6,

APOM), and a gene of unknown function, LOC101928978. On the other hand, there are also

6 genes that were not identified in this analysis but were highlighted by the analysis based on

geographic coordinates. This group includes genes associated with pathogen recognition and

activation of innate immunity (e.g., TLR10), speech-language disorder 1 (e.g., FOXP2), skin

and eye color (e.g., OCA2), expression of gamma-Glutamyltransferase (SOX9-AS1), bipolar

disease or neuropsychiatric disorders (e.g., CDH7), as well as a gene of unknown function,

LOC105373760 (Supplementary Table S4).

Discussion

In this study, we present DeepGenomeScan, a new deep learning method that can be used to

study how natural selection shapes genomes of species (genome-scan application) and to

identify genomic regions associated with phenotypic traits (GWAS application). For the sake

of simplicity and brevity, we focused on the genome-scan applications to detect signatures of

natural selection. However, in the Supplementary Material, we present an evaluation of the

performance of DeepGenomeScan when applied to the detection of genomic regions

associated with quantitative traits. This was done using the same set of simulations described

for the genome-scan application but in this case, the response variable was a quantitative trait

that influenced fitness (see details in Supplementary Methods). The results showed that

DeepGenomeScan can achieve high power (80%) to identify QTLs under a wide range of

spatial scenarios (linear, non-linear and coarse-grained heterogeneous environments) while

maintaining very low error rates (FDR < 0.12; FPR < 0.001; see Figure S6-7).

The insight upon which our method relies is the idea that we can use the genotypes of

individuals to predict any associated trait, not limited to just their phenotype but also their

spatial location or the environmental attributes of the habitat they live in. Intuitively, the type

of functions that can be used to describe the association between genotypes and any of these

dependent variables is likely to be radically different depending on the "trait" under

consideration. Therefore, no single model-based approach can be used to take advantage of

the above-mentioned insight. Additionally, no single model-free statistical method can be

used to approximate very complex non-linear functions linking genotypes and these disparate

traits. In contrast, Deep Neural Networks can approximate arbitrarily complex non-linear

functions linking dependent and predictor variables [22] and our study showed that they can

be an invaluable tool for population and evolutionary genomics applications.

Our approach represents an integration of genome wide association studies and genome scan

approaches in the specific case of spatially structured populations. More precisely, it is

focused on traits that vary spatially. Thus, it is not well adapted to study global selective

sweeps unless they are at an early stage where selected variants still exhibit clinal variation.

Similarly, DeepGenomeScan is not appropriate to carry out GWAS studies of panmictic

populations. On the other hand, it is perfectly suited to the study of local adaptation. There

are three questions that need to be answered in this context: (i) what are the environmental

drivers of natural selection? (ii) what are the phenotypic traits upon which selective pressures

act? and (iii) what are the genomic regions underlying those adaptive traits? As a genome-

scan method, DeepGenomeScan can identify the environmental drivers of natural selection

while as a GWAS tool, it can identify the genomic regions underlying adaptive traits.

Therefore, it can answer questions (i) and (iii). Question (ii), however, would require the

application of a quantitative genetic method to identify phenotypic traits that are good

candidates for being involved in local adaptation to heterogenous environmental conditions.

In this regard, we note that a recent review [45] has highlighted the fact that the study of local

adaptation requires combining population genomics and GWAS methods in the context of

common garden experiments. Therefore, our unified approach provides an excellent tool to

implement such frameworks.

The results of our analyses of an existing synthetic dataset [18] generated by simulations of a

polygenic trait under various spatial patterns of selection showed that DeepGenomeScan can

successfully identify outlier loci exhibiting complex non-linear associations with the target

trait. This represents an important advance over existing methods that can only consider

monotonic associations. For example, SPA uses a logistic function to describe allele

frequency behaviour as a function of geographic location and can only detect outlier loci

exhibiting monotonic geographic gradients [25]. Moreover, it cannot be applied to study

associations with phenotypic or environmental data.

Evolutionary biologists interested in understanding the mechanisms driving local adaptation

and speciation follow a longstanding tradition focused on the analysis of variation in

phenotype or allele frequencies along linear environmental gradients [33, 46]. Here we

showed that recent machine learning methods such as pcadapt and RDA can successfully

identify genomic regions associated with those patterns (Figs. 2 & 3). However, their

performance is limited in the case of non-monotonic gradients and, moreover, they are

completely incapable of considering coarse-grained heterogeneous spatial patterns in allele

frequencies (Fig 2 & 3). Our method clearly outperforms pcadapt and RDA in uncovering the

outlier loci that contribute the most to spatial population structure, regardless of the

underlying spatial pattern in selective pressures. Nevertheless, it is clear that very coarse-

grained heterogenous selection with no clear spatial pattern represents a very difficult

scenario, which reduces the power of our method to detect genomic regions associated with

the response of a species to selection.

Although less well studied empirically, coarse-grained heterogenous selection with no clear

spatial pattern has been an important focus of theoretical studies aimed at explaining

observed patterns of genetic variation [47-52]. This scenario is particularly relevant to studies

of protected polymorphisms [48, 50, 51] and hard versus soft selection [49]. Additionally, the

importance of this type of selection pattern is particularly relevant for genome-wide

association studies focused on diseases and phenotypic characters that do not exhibit clear

spatial patterns. Deep learning genome-scan methods capable of uncovering genomic regions

associated with this type of selection pattern can, therefore, provide a more general

understanding of how prevalent these particular mechanisms are.

We also applied DeepGenomeScan to the Population Reference Sample [37] (POPRES)

dataset to determine if our method could uncover genomic regions undetected by SPA and

other popular methods such iHS [53], Fst [10], Bayenv [13]. Comparing Yang et al.'s [25]

Supplementary Table 4 with our Supplementary Table 1, reveals a large number of important

genes that were not detected by those other existing methods. Some of these genes have been

linked to complex diseases such as Parkinson, obesity or various types of cancer and may not

be associated with monotonic selection gradients.

DeepGenomeScan can be applied to spatial genetic data even when individuals' geographic

coordinates are not available. In this case, our method uses the first two reduced features

obtained from a very recent machine learning method (KLFDAPC [36]), as proxies for

latitude and longitude. A comparison of the results from analyses of the POPRES dataset

based on geographic coordinates with those based on the first two reduced features reveals

very extensive overlap in terms of the genes they uncover. The small discrepancy in genes

uncovered by the two options implemented in our software is due to differences in the type of

spatial information used. It is clear that reduced features could lead to a loss of detailed

spatial information provided by geographic location. On the other hand, the use of reduced

features could avoid biases introduced by low quality geographic information or could take

into account small scale spatial effects that are not detectable at the spatial scale used for the

geographic coordinates.

There are still several methodological challenges faced when implementing deep learning

methods, which are associated with the computational cost of hyperparameter tuning and

neural network training. However, recent adaptive resampling algorithms (c.f. ref. [23]),

which we implemented in our software, carry out an efficient exploration of hyperparameter

space, allowing the use of deep learning to analyse large population genomic data sets.

Although the application of deep learning to population genetics problems is still in its

infancy [23], there are already some examples of such applications [27, 54-57]. One of these

applications is aimed at distinguishing between hard and soft sweeps and simultaneously

incorporating the confounding effects of demographic history [27]. This application involves

choosing among two or more alternative models, an objective that is usually achieved using

Approximate Bayesian Computation (ABC) [58, 59]. Although model choice in ABC is akin

to a classification application in machine learning, neural networks require a training set that

is not readily available when the target is a demographic or selection model. To overcome

this limitation, the authors train the neural network using synthetic data generated under

predefined evolutionary models. A drawback of this strategy is that it introduces model

assumptions into a computational framework that could be completely free of them and,

therefore, of very wide applicability. This drawback is absent when the objective is to assign

individuals to geographic locations as in ref. [54] or when scanning the genome in search of

outlier loci, as in our study. Overall, DeepGenomeScan fully exploits the flexibility and

power of deep neural networks, which makes it applicable to a wide range of problems

including identification of genomic regions associated with diseases or economically

important phenotypic traits, assignment of individuals to geographic locations, or

identification of environmental factors associated with selective pressures.

Methods

In what follows we first describe the architecture of the deep neural network and its

implementation. We then describe the statistical approach to identify SNPs that are located in

regions subject to positive (local adaptation or selective breeding) or negative selection

(diseases). Finally, we explain the simulation approach used to test performance and the

dataset used to provide a practical application.

MLP Architecture

In this study, we constructed a Multilayer Perceptron Network with three hidden layers (Fig.1

and Fig. S1). To describe this MLP, we first assume that the genotypes of the n sampled

individuals at p loci are described by a n×p matrix, $\mathbf{x} = (x_{ij})$, and are coded by the number of

derived alleles, which in the case of a biallelic marker takes values $x_{ij} = 0, 1, 2$, where i = 1,

2, ..., n, and j=1, 2, ..., p. Furthermore, the trait values of all individuals are arranged in a n×1

vector $\mathbf{y} = (y_i)$. Then the input layer of the MLP consists of p nodes, each one representing a

distinct locus. The information contained in node j comprises the individual genotypes at

locus j, $\mathbf{x}_{*j} = (x_{1j}, \cdots, x_{nj})$. This information is transformed into a signal which is fed to the

first hidden layer. The signal received by each node k = 1, 2, ..., K_1 of the first hidden layer

$\hat{\mathbf{s}}^{k1} = (\hat{s}_1^{k1}, \cdots, \hat{s}_i^{k1}, \cdots, \hat{s}_n^{k1})$ represents a weighted non-linear regression of individuals' trait

values on their multilocus genotypes. Here, the superscript k1 identifies the k^{th}-node of the

first hidden layer. Each element $\hat{s}_i^{k1}$ of the signal vector received by a node k of the first

hidden layer is given by

$\hat{s}_i^{k1} = f^1(\mathbf{x}_{i*}) = G\left(\sum_{j=1}^{p} x_{ij} w_j^{k1} + b^{k1}\right) = G(\mathbf{x}_{i*}\mathbf{w^{k1}} + b^{k1}),$ (1)

where w_j^{k1} is the weight of locus j, b^{k1} is a scalar representing a bias in the signal received

by node k in the first hidden layer, and $G(\cdot)$ is the activation function used to transform the

information sent by the input layer to the first hidden layer.

In a similar way, each node k of the second hidden layer will take the output signal of all

nodes of the first hidden layer and apply a transformation before sending the signal to the

third hidden layer:

$\hat{s}_i^{k2} = f^2\left(\hat{s}_i^{k1}\right) = G\left(\sum_{m=1}^{K_1} \hat{s}_i^{k1} w_m^{k2} + b^{k2}\right) = G\left(f^1(\mathbf{x}_{i*})\mathbf{w}^{k2} + b^{k2}\right).$ (2)

If there are only two hidden layers, then the neural network is represented by $f(\mathbf{x}) =$

$f^2(f^1(\mathbf{x}))$. In general, there can be an arbitrary number L of hidden layers, each consisting of

K_l nodes and the signal generated by the last hidden layer is the vector of predicted trait

values, $\hat{\mathbf{y}} = (\hat{y}_i)$. Therefore, a MLP network can be represented by,

$f(\mathbf{x}) = f^L(f^{L-1}\left(\cdots f^2\left(f^1(\mathbf{x})\right)\right),$ (3)

where $\mathbf{f}^l(\cdot)$ is the signal received by the l☐☐ layer.

Given the input data, i.e. the genotypes, and the observed trait values, the neural network

learns the weights $\mathbf{w} = (\mathbf{w}^1, ..., \mathbf{w}^L)$ with $\mathbf{w}^l = \left(w^{1l}, \cdots, w^{K_l l}\right)$ and biases $\mathbf{b} =$

$(\mathbf{b}^1, ..., \mathbf{b}^L)$ with $\mathbf{b}^l = \left(b^{1l}, \cdots, b^{K_l l}\right)$ that best describe the relationship between the inputs and

the outputs by minimising the difference between predicted and observed trait values [60,

61]. These weights represent an essential element of our genome scan method because they

are used to identify outlier loci (see below) using a procedure supported by previous studies

[31, 62], which show that the importance of a variable (locus) can be estimated as the

combination of the absolute values of weights associated with the graph edges connecting the

predictor variable (locus) with the MLP output (predicted trait).

Implementation

DeepGenomeScan trains the deep neural network to estimate the nonlinear function mapping

genotypes to individual traits. This process involves not only learning the vectors of weights

and biases by iteratively adjusting these parameters but also the tuning of a large number of

hyperparameters associated with structural components of the model, which impact model

performance. These include those determined by the network architecture (activation

functions, number of neurons per hidden layer) and those included in the optimisation

algorithm used for training (e.g. learning rate, batch size). For optimisation, we used resilient

backpropagation [63] with weight backtracking when analysing both simulated and real

datasets. However, the tuning algorithm used depended on the size of the input data. In the

case of simulated datasets, which were of limited size, we used a full resampling strategy

corresponding to Algorithm 1 in [23] based on a repeated k-fold cross validation with k = 5

and five repetitions. In the case of the large POPRES dataset, we used an adaptive resampling

approach corresponding to Algorithm 2 in [23], which incorporates futility assessment into

the model tuning process. The resampling approach in this case was the same as in the

analysis of simulated data (five replications of a five-fold cross-validation). Detailed

information about the settings used for the optimisation and tuning algorithms is presented in

Supplementary Notes.

Identification of outlier loci

As mentioned before, the SNP importance can be estimated as combinations of the absolute

values of connection weights [31, 62]. This is done once the optimal model is found. We used

Olden and Jackson's [24] method, which is based on Garson's [64] algorithm to calculate the

relative importance for each input node but adds a randomisation step to identify non-

significant connection weights.

DeepGenomeScan carries out separate runs for each trait and generates a vector of SNP

importance for each of them. Given T traits, the position of each SNP in trait space is

described by its associated vector of importance values and, therefore, it is possible to

calculate the Mahalanobis distance between the focal locus and all other loci. Our approach

leads to an intuitive definition of outlier loci as any locus with an "extreme" Mahalanobis

distance. Since the squared Mahalanobis distance follows a chi-squared distribution with T

degrees of freedom, the p-values associated with each SNP can be obtained from a χ^2_T

distribution [65] and used for identifying 'outlier' loci. This approach relies on the

assumption that variables used to calculate Mahalanobis distance follow a normal

distribution. Therefore, we used the arcsine transformation to normalise the weights before

calculating the Mahalanobis distance.

Human dataset

We applied DeepGenomeScan to European populations from the POPRES project [37]. The

dataset contains a total of 3,192 European individuals genotyped at 500,568 loci using the

Affymetrix 500K SNP chip. The sample collections and genotyping for POPRES are

described in [37]. We removed individuals from outside of Europe and individuals whose

grandparents had different geographic origins based on the criteria used by [35]. We also

removed samples that only have one individual per country, such as Denmark, Finland,

Latvia, Ukraine, Slovakia and Slovenia. Geographic coordinates for each individual

corresponded to the central point of the geographic area of the individual's country of birth.

We removed the monomorphic SNPs and filtered out autosomal biallelic SNPs with a minor

allele frequency (MAF) below 5%, and a missing rate of 5% or more. In the end, we kept 1,

382 individuals from 32 countries carrying 283, 499 autosomal SNPs. We annotated and

updated the variant reference ID based on the comprehensive report of short human

variations from the human variation database (dbSNP, GRCh37p13, b151, release 2018,

https://ftp.ncbi.nih.gov/snp/organisms/human_9606/VCF/) using bcftools [66].

Data and code availability

The simulation data used in this study is available at

https://datadryad.org/review?doi=doi:10.5061/dryad.1s7v5.

The datasets used for the analyses described in this manuscript were obtained from dbGap at

https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000145.v4.p2

through dbGap accession number phs000145.v4.p2 (request approval #90291-1).

The DeepGenomeScan R code is hosted on GitHub at

https://github.com/xinghuq/DeepGenomeScan. The scripts used in this study are available at

https://github.com/xinghuq/DeepGenomeScan/tree/webpkg/DeepGenomeScan_simulation%2

BPOPRES. The KLFDAPC package is available at https://xinghuq.github.io/KLFDAPC/.

Ethics declarations

Ethics approval and consent to participate

Ethical approval was not needed for this study.

Consent for publication

Not applicable.

Competing Interests

The authors declare no competing interests.

Acknowledgments

535   XQ was supported by a PhD scholarship from the China Scholarship Council. We benefited

536   from discussions with Pierre de Villemereuil.

Author Contributions

538   XQ and OEG designed the study, XQ carried out the analyses and interpreted results with

539   input from OEG and CWKC. OEG and XQ wrote the article with input from CWKC.

References

1.   Bush WS, Moore JH: **Chapter 11: Genome-Wide Association Studies.** *Plos Computational Biology*
2012, **8**.

2.   Kang HM, Sul JH, Service SK, Zaitlen NA, Kong SY, Freimer NB, Sabatti C, Eskin E: **Variance component**
model to account for sample structure in genome-wide association studies. *Nature Genetics* 2010,
42:348-U110.

3.   Tam V, Patel N, Turcotte M, Bosse Y, Pare G, Meyre D: **Benefits and limitations of genome-wide**
association studies. *Nature Reviews Genetics* 2019, **20:**467-484.

4.   Edge MD, Coop G: **Reconstructing the History of Polygenic Scores Using Coalescent Trees.** *Genetics*
2019, **211:**235-262.

5.   Field Y, Boyle EA, Telis N, Gao ZY, Gaulton KJ, Golan D, Yengo L, Rocheleau G, Froguel P, McCarthy MI,
Pritchard JK: **Detection of human adaptation during the past 2000 years.** *Science* 2016, **354:**760-764.

6.   Racimo F, Berg JJ, Pickrell JK: **Detecting Polygenic Adaptation in Admixture Graphs.** *Genetics* 2018,
208:1565-1584.

7.   Speidel L, Forest M, Shi SN, Myers SR: **A method for genome-wide genealogy estimation for**
thousands of samples. *Nature Genetics* 2019, **51:**1321-+.

8.   Turchin MC, Chiang CWK, Palmer CD, Sankararaman S, Reich D, Hirschhorn JN, Genetic Invest A:
Evidence of widespread selection on standing variation in Europe at height-associated SNPs.
Nature Genetics 2012, **44:**1015-+.

9.   Oleksyk TK, Smith MW, O'Brien SJ: **Genome-wide scans for footprints of natural selection.**
Philosophical Transactions of the Royal Society B-Biological Sciences 2010, **365:**185-205.

10. Lewontin RC, Krakauer J: **Distribution of gene frequency as a test of the theory of the selective**
neutrality of polymorphisms. *Genetics* 1973, **74:**175-195.

11. Beaumont MA, Balding DJ: **Identifying adaptive genetic divergence among populations from**
genome scans. *Mol Ecol* 2004, **13:**969-980.

12. Foll M, Gaggiotti O: **A Genome-Scan Method to Identify Selected Loci Appropriate for Both**
Dominant and Codominant Markers: A Bayesian Perspective. *Genetics* 2008, **180:**977-993.

13. Coop G, Witonsky D, Di Rienzo A, Pritchard JK: **Using Environmental Correlations to Identify Loci**
Underlying Local Adaptation. *Genetics* 2010, **185:**1411-1423.

14. de Villemereuil P, Gaggiotti OE: **A new F-ST-based method to uncover local adaptation using**
environmental variables. *Methods Ecol Evol* 2015, **6:**1248-1258.

15. Frichot E, Schoville SD, Bouchard G, François O: **Testing for associations between loci and**
environmental gradients using latent factor mixed models. *Molecular biology and evolution* 2013,
30:1687-1699.

16. Gaggiotti OE, Bekkevold D, Jorgensen HBH, Foll M, Carvalho GR, Andre C, Ruzzante DE: **Disentangling**
the effects of evolutionary, demographic, and environmental factors influencing the genetic
structure of natural populations: Atlantic herring as a case study. *Evolution* 2009, **63:**2939-2951.

17. Torada L, Lorenzon L, Beddis A, Isildak U, Pattini L, Mathieson S, Fumagalli M: **ImaGene: a convolutional**
neural network to quantify natural selection from genomic data. vol. 20; 2019.

18. Capblancq T, Luu K, Blum MG, Bazin E: **Evaluation of redundancy analysis to identify signatures of**
local adaptation. *Mol Ecol Resour* 2018, **18:**1223-1233.

19. Duforet-Frebourg N, Luu K, Laval G, Bazin E, Blum MG: **Detecting genomic signatures of natural**
selection with principal component analysis: application to the 1000 genomes data. *Molecular*
biology and evolution 2016, **33:**1082-1093.

20. Yan Q, Jiang Y, Huang H, Swaroop A, Chew EY, Weeks DE, Chen W, Ding Y: **Genome-wide association**
studies-based machine learning for prediction of age-related macular degeneration risk.
Translational vision science & technology 2021, **10:**29-29.

21. Sun T, Wei Y, Chen W, Ding Y: **Genome-wide association study-based deep learning for survival**
prediction. *Statistics in medicine* 2020, **39:**4605-4620.

22. Hornik K, Stinchcombe M, White H: **Multilayer feedforward networks are universal approximators.**
Neural networks 1989, **2:**359-366.

23. Kuhn M: **Futility analysis in the cross-validation of machine learning models.** *arXiv:14056974* 2014.

24. Olden JD, Jackson DA: **Illuminating the "black box": a randomization approach for understanding**
variable contributions in artificial neural networks. *Ecological modelling* 2002, **154:**135-150.

25. Yang WY, Novembre J, Eskin E, Halperin E: **A model-based approach for analysis of spatial structure**
in genetic data. *Nature Genetics* 2012, **44:**725-U163.

26. Sharma S: **Activation functions in neural networks.** *Towards Data Science* 2017, **6**.

27. Sheehan S, Song YS: **Deep Learning for Population Genetic Inference.** *Plos Computational Biology*
2016, **12**.

28. Attali J-G, Pagés G: **Approximations of functions by a multilayer perceptron: a new approach.**
Neural networks 1997, **10:**1069-1081.

29. Pal SK, Mitra S: **Multilayer perceptron, fuzzy sets, classifiaction.** 1992.

30. Gevrey M, Dimopoulos L, Lek S: **Review and comparison of methods to study the contribution of**
variables in artificial neural network models. *Ecol Model* 2003, **160:**249-264.

31. Olden JD, Joy MK, Death RG: **An accurate comparison of methods for quantifying variable**
importance in artificial neural networks using simulated data. *Ecological modelling* 2004, **178:**389-
397.

32. Luu K, Bazin E, Blum MG: **pcadapt: an R package to perform genome scans for selection based on**
principal component analysis. *Molecular ecology resources* 2017, **17:**67-77.

33. Endler JA: *Geographic variation, speciation and clines.* Princeton, NJ: Princeton University Press; 1977.

34. Lao O, Lu TT, Nothnagel M, Junge O, Freitag-Wolf S, Caliebe A, Balascakova M, Bertranpetit J, Bindoff LA,
Comas D, et al: **Correlation between genetic and geographic structure in Europe.** *Current Biology*
2008, **18:**1241-1248.

35. Novembre J, Johnson T, Bryc K, Kutalik Z, Boyko AR, Auton A, Indap A, King KS, Bergmann S, Nelson MR:
Genes mirror geography within Europe. *Nature* 2008, **456:**98-101.

36. Qin X, Chiang CWK, Gaggiotti OE: **KLFDAPC: A Supervised Machine Learning Approach for Spatial**
Genetic Structure Analysis. *Briefings in Bioinformatics* 2021**:**2021.2005.2015.444294.

37. Nelson MR, Bryc K, King KS, Indap A, Boyko AR, Novembre J, Briley LP, Maruyama Y, Waterworth DM,
Waeber G: **The Population Reference Sample, POPRES: a resource for population, disease, and**
pharmacological genetics research. *The American Journal of Human Genetics* 2008, **83:**347-358.

38. Granovsky M, Fata J, Pawling J, Muller WJ, Khokha R, Dennis JW: **Suppression of tumor growth and**
metastasis in Mgat5-deficient mice. *Nature medicine* 2000, **6:**306-312.

39. Brynedal B, Wojcik J, Esposito F, Debailleul V, Yaouanq J, Martinelli-Boneschi F, Edan G, Comi G, Hillert J,
Abderrahim H: **MGAT5 alters the severity of multiple sclerosis.** *Journal of neuroimmunology* 2010,
220:120-124.

40. Wang R, Fan Q, Zhang J, Zhang X, Kang Y, Wang Z: **Hydrogen sulfide demonstrates promising**
antitumor efficacy in gastric carcinoma by targeting MGAT5. *Translational oncology* 2018, **11:**900-
910.

41. Fox CS, Liu Y, White CC, Feitosa M, Smith AV, Heard-Costa N, Lohman K, Johnson AD, Foster MC,
Greenawalt DM: **Genome-wide association for abdominal subcutaneous and visceral adipose**
reveals a novel locus for visceral fat in women. *PLoS Genet* 2012, **8:**e1002695.

42. Nalls MA, Pankratz N, Lill CM, Do CB, Hernandez DG, Saad M, DeStefano AL, Kara E, Bras J, Sharma M:
Large-scale meta-analysis of genome-wide association data identifies six new risk loci for
Parkinson's disease. *Nature genetics* 2014, **46:**989-993.

43. Dichgans M, Malik R, König IR, Rosand J, Clarke R, Gretarsdottir S, Thorleifsson G, Mitchell BD, Assimes
TL, Levi C: **Shared genetic susceptibility to ischemic stroke and coronary artery disease: a genome-**
wide analysis of common variants. *Stroke* 2014, **45:**24-36.

44. Comuzzie AG, Cole SA, Laston SL, Voruganti VS, Haack K, Gibbs RA, Butte NF: **Novel genetic loci**
identified for the pathophysiology of childhood obesity in the Hispanic population. *PloS one* 2012,
7:e51954.

45. de Villemereuil P, Gaggiotti OE, Mouterde M, Till-Bottraud I: **Common garden experiments in the**
genomic era: new perspectives and opportunities. *Heredity* 2016, **116:**249-254.

46. Haldane JBS: **THE THEORY OF A CLINE.** *Journal of Genetics* 1948, **48:**277-284.

47. Bulmer MG: **Multiple niche polymorphism.** *Amer Natur* 1972, **106:**254-257.

48. Levene H: **Genetic equilibrium when more than one ecological niche is available.** *Amer Natur* 1953,
87:331-333.

49. Levins R, MacArthur R: **The maintenance of genetic polymorphism in a spatially heterogeneous**
environment: variations on a theme by Howard Levene. *Americal Naturalist* 1966, **100:**585-589.

50. Prout T: **SUFFICIENT CONDITIONS FOR MULTIPLE NICHE POLYMORPHISM.** *American Naturalist*
1968, **102:**493-&.

51. Strobeck C: **Haploid selection withn alleles in m niches.** *American Naturalist* 1979, **113:**439-444.

52. Maynard S: **Genetic polymorphism in a varied environment.** *American Naturalist* 1970, **104:**487-490.

53. Sabeti PC, Reich DE, Higgins JM, Levine HZP, Richter DJ, Schaffner SF, Gabriel SB, Platko JV, Patterson NJ,
McDonald GJ, et al: **Detecting recent positive selection in the human genome from haplotype**
structure. *Nature* 2002, **419:**832-837.

54. Battey CJ, Ralph PL, Kern AD: **Predicting geographic location from genetic variation with deep**
neural networks. *Elife* 2020, **9**.

55. Flagel L, Brandvain Y, Schrider DR: **The Unreasonable Effectiveness of Convolutional Neural**
Networks in Population Genetic Inference. *Molecular Biology and Evolution* 2019, **36:**220-238.

56. Åkesson M, Singh P, Wrede F, Hellander A: **Convolutional Neural Networks as Summary Statistics for**
Approximate Bayesian Computation. *arXiv preprint arXiv:200111760* 2020.

57. Jiang B, Wu T-y, Zheng C, Wong WH: **Learning summary statistic for approximate Bayesian**
computation via deep neural network. *Statistica Sinica* 2017:1595-1618.

58. Beaumont MA, Zhang WY, Balding DJ: **Approximate Bayesian computation in population genetics.**
Genetics 2002, **162:**2025-2035.

59. Csillery K, Blum MGB, Gaggiotti OE, Francois O: **Approximate Bayesian Computation (ABC) in**
practice. *Trends Ecol Evol* 2010, **25:**410-418.

60. Jinming Y, Ahmadi M, Jullien GA, Miller WC: **Model validation and determination for neural network**
activation function modeling. 1998:548-551.

61. Goodfellow Ia: **Deep learning / Ian Goodfellow, Yoshua Bengio and Aaron Courville.** pp. 0-5.
Cambridge, Massachusetts :: The MIT Press; 2016:0-5.

62. Gevrey M, Dimopoulos I, Lek S: **Review and comparison of methods to study the contribution of**
variables in artificial neural network models. *Ecological modelling* 2003, **160:**249-264.

63. Riedmiller M: **Advanced supervised learning in multi-layer perceptrons—from backpropagation to**
adaptive learning algorithms. *Computer Standards & Interfaces* 1994, **16:**265-278.

64. Garson DG: **Interpreting neural network connection weights.** *Artificial Intelligence Expert* 1991, **6:**46-
51.

65. Filzmoser P, Gregorich M: **Multivariate Outlier Detection in Applied Data Analysis: Global, Local,**
Compositional and Cellwise Outliers. *Mathematical Geosciences* 2020, **52:**1049-1066.

66. Li H: **A statistical framework for SNP calling, mutation discovery, association mapping and**
population genetical parameter estimation from sequencing data. *Bioinformatics* 2011, **27:**2987-
2993.

683