1

Deciphering signatures of natural

selection via deep learning

Xinghu Qin[1*], Charleston W. K. Chiang[2], Oscar E. Gaggiotti[1*]

[1] Centre for Biological Diversity, Sir Harold Mitchell Building, University of St Andrews, Fife, KY16 9TF, UK

[2] Center for Genetic Epidemiology, Keck School of Medicine & Department of Quantitative and Computational Biology, University of Southern California, USA

* Correspondence should be addressed to XQ (xq5@st-andrews.ac.uk) and OE Gaggiotti (oeg@st-andrews.ac.uk)

Abstract

Identifying genomic regions influenced by natural selection provides fundamental insights

into a wide range of problems including human health, animal and plant breeding, and the

understanding of local adaptation. We propose a new method, DeepGenomeScan, that can be

used to address all these problems. It is based on the principle that the genotypes of

individuals can be used to predict any associated trait; not only their phenotype but also their

spatial location or the environmental attributes of the habitat they live in. We, therefore,

implemented a deep learning method to detect candidate regions under selection by

identifying loci that contribute the most to the predictive power of the deep neural network.

Using simulations, we show that our method can successfully identify loci underlying

quantitative traits subject to complex spatial patterns of selection. We apply

DeepGenomeScan to a European human genetic variation dataset and posit that the loci that

contribute the most to the prediction of latitude and longitude are located in genomic regions

under selection. Using this approach, we identified many SNPs located within well-known

genes, some of which were not identified using existing population genetics approaches, e.g.

MCM6, MGAT5, TMEM163.

Main

One of the main challenges of modern biology is to dissect and understand the molecular

basis for naturally occurring phenotypic variation. Addressing this challenge is of

fundamental importance not only for the field of evolutionary biology but also for a wide

variety of applied fields involving human diseases, improvement of agricultural crops and

breeds and biodiversity conservation.

The recent increases in genomic data generated by modern sequencing technologies have

advanced our understanding of how natural selection, and its interactions with other

evolutionary forces, shapes the genome and phenotype of species. Such technologies have

now been applied to the study of a wide range of species, but the statistical methods used to

analyse them tend to differ depending on whether the focus is on humans and domesticated

species or on wild species. In the case of humans and domesticated species, there is a

particular interest in uncovering associations between well-defined phenotypes (traits

associated with diseases or, in the particular case of domesticated species, yield) and the

genomic variants that underlie them. The standard approach, in this case, is to use Genome

Wide Association Studies (GWAS; e.g. ref [1]), which linearly model the additive allelic effect

of genotypes (the explicative variable) on phenotypes (the dependent variable). All of the

recent methods can take into account population stratification, using for example linear

mixed models that incorporate the genetic relationship matrix (GRM) as a random effect (e.g.

ref. [2]). GWAS have been used to identify variants associated with a wide range of phenotypic

traits[3] and their results are used by several studies aimed at detecting polygenic adaptation[4-8].

In the case of evolutionary studies and in particular those focused on wild species, the most

commonly used approaches are a suite of methods known as the genome-scans [9], which

ignore phenotypic traits. Many of them are population-based and use summary statistic to

uncover genomic regions exhibiting outlier behaviour. A very popular approach is to identify

loci that exhibit unusually strong among-population genetic differentiation. This approach

assumes that, in the absence of selection, differences in allele frequencies among populations

should follow a neutral distribution driven by migration and genetic drift. Any locus that

departs from this expectation is considered a good candidate for being under the effect of

natural selection. This approach initially proposed by Lewontin & Krakauer [10] has been

implemented in popular Bayesian methods (e.g. [11,12]). More recently, there has been an

increased interest in establishing an association between a locus outlier behaviour and

environmental variables, with the assumption that such variables could represent selective

pressures acting upon genomic regions linked to the outlier loci[13-16]. Some of these methods

use raw genotypic data instead of summary statistics (e.g. [12]) but as opposed to GWAS, they

use linear mixed models that consider the genotype as the dependent variable, and the

environmental factor as the explicative variable.

Despite this apparent methodological dichotomy between studies focused on humans and

domesticated species and those focused on wild species, it is clear that a thorough

understanding of how natural selection shapes the phenotype and genome of species requires

some combination of these two different approaches. Indeed, the selective pressures exerted

by the heterogenous abiotic and biotic environment act upon individuals' phenotypes in a

complex way and lead to changes in the spatial structuring of variation in the genomic

regions underlying them. Thus, in order to fully characterise the action of natural selection,

we need to answer three fundamental questions: (i) what are the environmental drivers of

natural selection? (ii) what are the phenotypic traits upon which selective pressures act? and

(iii) what are the genomic regions underlying those adaptive traits?

Here we present DeepGenomeScan, a deep learning approach that can be used to answer all

three questions. The rationale underlying our method is that we can use the genotype of an

individual to predict not only its phenotype but also some attribute of the habitat where they

are sampled. This framework is akin to GWAS but in our case, the response variable can be

an individual phenotypic trait, an environmental variable associated with its habitat, or the

geographic location of the individual (latitude, longitude). Our approach leverages the power

of deep neural networks to approximate arbitrarily complex functions linking dependent and

explicative variables [17], as well as recent algorithms for model optimisation [18], and for

inferring the importance of each explicative variable in predicting the dependent variable [19].

It is important to note that as opposed to the prevalent use of neural networks, i.e. prediction

and pattern recognition, here our main interest is in estimating the features (loci) that

contribute the most to the predictive power of the neural network. To achieve this goal, we

use the concept of "feature importance"[19], which represents a proxy for the effect size of any

given locus.

An important advantage of our method when applied to spatial data is that it can consider any

spatial selection pattern including the usually assumed linear environmental gradient as well

as arbitrarily complex non-linear spatial patterns and, importantly, coarse-grained

heterogeneous selection with no clear spatial pattern. This represents an important advance as

existing methods only consider linear or monotonic non-linear patterns. In this particular

application, our approach generalises Yang et al.'s[20] idea of using geographic positioning of

individuals to identify loci that exhibit particularly steep slopes of allele frequency change

associated with recent positive selection. Our generalisation is two-fold: (i) instead of only

considering monotonic spatial gradients, we can detect loci associated with arbitrary and non-

monotonic selection patterns, and (ii) the dependent variable can be the geographic location

but also a phenotypic trait or an environmental factor.

Results

Underlying Rationale for the Deep Learning Approach

A neural network can be considered as a generalised regression approach used to learn complex functions expressing the association between the input data and a response variable[20]. In our case, the input or predictor variables are the genotypes of the individuals and the response variables are observed phenotypes, environmental values, or geographic locations. In what follows, we use the term traits to refer to these response variables.

As opposed to standard regression, which fits a fixed function to the data, a neural network learns the (non-linear) function most appropriate for the data at hand. Here we use a multilayer perceptron (MLP), which is composed of up to three layers of nodes (neurons) with connections between layers but not between nodes within a layer (Figure 1 and Supplementary Notes). Connection edges between nodes can have different weights and the main goal of the MLP is to learn the weights that best describe the complex function that associates inputs and outputs[21] by minimising the difference between the predicted and observed traits. Once the optimal network is found, the weights connecting each input node with the output node can be used to devise a test to identify the loci that contribute the most to the trait under consideration. The underlying rationale for this test is that the absolute values of weights associated with the path linking an input node (i.e. a locus) with the output can be used to estimate the importance of an input variable[22,23] (node or locus). Loci with extreme importance values are considered as outliers and good candidates for being under the influence of natural selection. Full details of our approach are provided in Online Methods.

Simulation Study

We evaluated the performance of our method using simulated data generated by Capblancq et al.[24] (downloaded from the Dryad Digital Repository at:

https://datadryad.org/review?doi=doi:10.5061/dryad.1s7v5). The simulations assume a two-dimensional stepping-stone scenario and consider three quantitative traits, each coded by a distinct set of 10 loci (QTLs), resulting in a total of 30 causal SNPs, out of a total of 1000. Trait values are calculated simply as the sum of genotypic values of the causal loci. Each quantitative trait (QT) is influenced by environmental factors having a distinct spatial pattern. QT1 is influenced by factors with a quadratic spatial gradient, QT2 is influenced by factors with a linear gradient, and QT3 is influenced by factors with a coarse and patchy spatial pattern. There is a total of 10 environmental factors falling into one of these three categories. Further details of the simulation approach are provided in Supplementary. Full details are given in ref. [24].

We compared the performance of our method with those of two recently proposed machine

learning-based approaches: pcadapt [25] and RDA[24]. Figure 2 presents these results in terms of

power (proportion of true positives), False Discovery Rate (FDR) and False Positive Rate

(FPR). Using a threshold p-value = 10^{-8}, as typically done in GWAS, our method has high

power to detect all QTL types, but the FDR is too high. On the other hand, pcadapt and RDA

have no power to detect loci associated with QTs 1 and 3, and very low power to detect those

associated with QT2 (Fig. 2a). We then progressively increased the stringency of the

threshold for DeepGenomScan (from p-value 10^{-16} up to 10^{-64}) and used a less stringent

threshold (p-value = 0.001) for pcadapt and RDA (Fig. 2d-l). It is possible to decrease the

FDR of our method below that of pcadapt using a threshold of 10^{-32}, and below that of RDA

using a threshold of 10^{-64} (Fig. 2h, i, k, l). Overall, all three methods have high power to

detect loci associated with QT2, which is influenced by a linear selective gradient.

DeepGenomeScan, however, has the lowest FDR under the most stringent threshold, and still

has much higher power than pcadapt and RDA to detect QT1 and QT3 loci, which are

influenced by non-linear selection patterns (Fig. 2j). It is important to note that this trade-off

between power and FDR only applies to cases with non-linear selection patterns. Indeed, the

power to detect loci underlying QT2 remains very high regardless of how stringent the

threshold we use is (Fig. 2). Qualitatively similar results were obtained when we used q-value

thresholds (Supplementary Notes).

Figure 2: Performance of pcadapt, RDA and DeepGenomeScan (DL_MLP). (a-c): Initially, the p-value threshold used for all three methods was 10^{-8}. Subsequently, we used a threshold of 0.001 for pcadapt and RDA but we progressively increased the stringency of the threshold for DeepGenomeScan: (d-f): 10^{-16}; (g-i) 10^{-32}; (j-l) 10^{-64}. Error bars indicate the standard deviation and letters are used to indicate the statistical significance of the difference in performance based on a t-test with p-value < 0.01. Panels a, d, g, j present the power to identify loci underlying each quantitative trait separately (QT1-3). All other panels present the overall FDR and FPR (across all three types of QTLs). All estimates were based on 100 simulated datasets.

Figure 3 presents Manhattan plots for one simulated data set with a somewhat permissive

threshold for pcadapt and RDA and a much more stringent threshold for DeepGenomeScan.

This figure illustrates the much higher power of DeepGenomeScan to detect outlier loci.

Indeed, in the case of the linear selection gradient, all the outlier loci fall off the y-axis scale

and only two are missed under quadratic selection gradients. Clearly, very discontinuous

selection patterns (i.e., QTL3) represent a big challenge for all methods but

DeepGenomeScan could still detect several of the outliers under this difficult scenario.

Figure 3: Manhattan plots of the results obtained with (a) pcadapt, (b) RDA and (c) DeepGenomeScan (DL_MLP) for one
simulated data set. Note that in c, all QT2 loci and some QT1 loci fall out of the range of the y axis. Note also that the scale
of the y axis in c differs from that used for a and b.

Application to a real data set

A common and longstanding framework to study the action of natural selection is to focus on

clinal variation in phenotypic traits or allele frequencies along environmental gradients [26].

One approach to detect genomic regions under selection in these clinal variation scenarios is

to identify loci exhibiting extreme frequency gradients across geographic space [20]. The

underlying assumption of this approach is that the environmental gradient is continuous,

leading to a monotonic but not necessarily linear change in allele frequency or phenotype.

Although there may be several examples of such geographic variation, spatial patterns in

selective pressures and the associated allele frequency can be non-monotonic. For example, it

is possible that the maximum allele frequency is located in the middle of the geographic

region under study [20]. As our simulations show, our approach is capable of identifying

genomic regions subject to such non-linear selection patterns.

In principle, approaches focused on clinal spatial patterns require the geographic coordinates

of each individual sample. However, sometimes this information is not available. A potential

solution to this problem is to carry out a Principal Component Analysis on the genotype

matrix and use the first two PC axes as surrogates for geographic coordinates [27,28]. This

approach, however, is based on a linear combination of genotypes and can lead to poor

inference of spatial location of admixed individuals [20]. A recently developed model-based

approach can be used to overcome this issue [20] but it requires the assumption of a smooth

monotonic function to describe allele frequency behaviour as a function of geographic

location, which may not be appropriate in all cases. Our implementation of

DeepGenomeScan allows the use of a more general dimensionality reduction approach,

Kernel Local Discriminant Analysis of Principal Components (KLFDAPC[29]; Supplementary

Material) when spatial coordinates are not available.

Here we use a human dataset to determine if our approach can uncover new regions of the

human genome that may be under the influence of non-linear spatial selection patterns. We

applied DeepGenomeScan to the Population Reference Sample [30] (POPRES) dataset using

the known geographic coordinates and then using the two first reduced features from a

KLFDAPC analysis (Online Methods). POPRES represents an excellent example of human

genetic variation aligning with geography [28]. The dataset contains a total of 3,192 European

individuals genotyped at 500,568 loci using the Affymetrix 500K SNP chip. Details about

data filtering steps are provided in Online Materials.

We calculated p-values for each SNP (see Online Methods for details) and obtained a

genomic inflation factor GIF = 1 for this data set. Using a threshold p-value = 1e-100 for

identifying the outlier loci, the first analysis using geographic coordinates as the response

variables detected 245 outlier SNPs located within 84 known genes. The full list of genes we

identified, and their chromosome positions is provided in Supplementary Table S1.

Consistent with previous widely reported regions under selection, our method detects strong

signals at the LCT region on chromosome 2, the ADH1C region on chromosome 4, the HLA

region on chromosome 6, as well as the OCA2 and the HERC2 region on chromosome 15

(Fig. 4; Supplementary Table S1).

Besides the already known selected genes, our method also detects some disease related

genes that exhibited extreme variation across geographic space but were not identified by

existing methods (c.f. Supplementary Table 4 in ref. [20]); these include MGAT5, TMEM163,

ACMSD, CCNT2, MAP3K19, R3HDM1, UBXN4, MCM6, DARS1, EHMT2, and CFB

(Supplementary Table 1). For example, MCM6 is a regulatory element that controls the

expression of the LCT gene. The MGAT5 gene (alpha-1,6-mannosylglycoprotein 6-beta-N-

acetylglucosaminyltransferase), is one of the most important enzymes involved in the

regulation of the biosynthesis of glycoprotein oligosaccharides and is associated with

invasive malignancies and sclerosis [31-33] as well as visceral fat in women [34]. The TMEM163

gene (transmembrane protein 163), is associated with Parkinson's disease, ischemic stroke

and coronary artery disease [35,36]. The ACMSD gene (aminocarboxymuconate semialdehyde

decarboxylase) is associated with Parkinson's disease [35] and childhood obesity [37]. There are

many other genes with important functions that our method highlighted as contributing very

strongly to the estimation of geographic origin (Supplementary Table S1). There are also

several other outlier SNPs detected by our method on chromosome 2, 4, 6, 8, 10, 11, 12, 15,

16, 18 (Supplementary Table 1) that are not within known functional genes. However, they

may be linked to regulatory genes and, therefore, of interest for human genetic studies.

Figure 4: Spatial genetic structure of European populations and signals of selection detected by DeepGenomeScan. (A).
Spatial genetic structure of European populations inferred via KLFDAPC with a σ = 5. (B). Manhattan plot indicating loci
under spatial selection obtained from DeepGenomeScan using geographic coordinates as the response variables. (C).
Manhattan plot indicating loci under spatial selection obtained from DeepGenomeScan using inferred spatial genetic
structure (the first two reduced features of KLFDAPC) as the response variables. The outliers in B-C were identified with a
threshold of p-value =1e-100. p-value lower than the threshold are highlighted in light green. Country abbreviations: AL,
Albania; AT, Austria; BA, Bosnia-Herzegovina; BE, Belgium; BG, Bulgaria; CH, Switzerland; CY, Cyprus; CZ, Czech
Republic; DE, Germany; ES, Spain; FR, France; GB, United Kingdom; GR, Greece; HR, Croatia; HU, Hungary; IE,
Ireland; IT, Italy; KS, Kosovo; MK, Macedonia; NO, Norway; NL, Netherlands; PL, Poland; PT, Portugal; RO, Romania;
RS, Serbia and Montenegro; RU, Russia; Sct, Scotland; SE, Sweden; TR, Turkey; YG, Yugoslavia. Note that in B and C,
some of the outlier loci on Chromosome 2 and 6 fall out of the range of the y axis. A full list of outlier loci can be found in
Supplementary Data Table 1-2.

In the case of the analysis for which we replaced the original geographic coordinates with the

first two reduced features obtained from KLFDAPC, we first calculated the correlation

between the first two reduced features and the original geographic coordinates. We compared

these results to those of a similar analysis using the first two PC axes obtained from a PCA of

the genotype matrix. The results confirm that KLFDAPC provides better estimates of

geographic location than PCA (see Fig. 4a and Supplementary Notes Fig. S6).

The genome scan based on the first two reduced features identifies a somewhat smaller

number of outlier SNPs (229, in 77 known genes) than when using geographic coordinates

(Fig. 4c and Supplementary Table S2). However, 80% of the outliers detected in this analysis

were also identified by the analysis based on geographic coordinates (185 shared loci). This

analysis also detected well-known selected genomic regions (LCT, HLA, ADH1, CHERC2;

Supplementary Table S2). However, there were 28 genes that were not identified when using

latitude and longitude (Supplementary Table S3). Genes in this group include some

associated with cancer (SLC44A5, AK5, TNXB) and other diseases such as Behcet's disease

(IL12AS1-AS1) as well as obesity and diabetes mellitus (APOM). On the other hand, there

are also 29 genes that were not identified in this analysis but were highlighted by the analysis

based on geographic coordinates. This group includes genes associated with cognitive

disorders (e.g. TMEM132D), psychiatric disorders (e.g. TRIM26) but also five others that

have unknown function or associated phenotype (Supplementary Table S4).

Discussion

In this study, we present DeepGenomeScan, a new deep learning method to study how

natural selection shapes the phenotypes and genomes of species. Our approach represents an

integration of genome wide association studies and genome scan approaches and can be used

to address three fundamental evolutionary questions: (i) what are the environmental drivers

of natural selection? (ii) what are the phenotypic traits upon which selective pressures act?

and (iii) what are the genomic regions underlying those adaptive traits.

The insight upon which our method relies is the idea that we can use the genotypes of

individuals to predict any associated trait, not limited to just their phenotype but also their

spatial location or the environmental attributes of the habitat they live in. Intuitively, the type

of functions that can be used to describe the association between genotypes and any of these

dependent variables is likely to be radically different depending on the "trait" under

consideration. Therefore, no single model-based approach can be used to take advantage of

the above-mentioned insight. Additionally, no single model-free statistical method can be

used to approximate very complex non-linear functions linking genotypes and these disparate

traits. In contrast, Deep Neural Networks can approximate arbitrarily complex non-linear

functions linking dependent and predictor variables[17] and our study showed that they can be

an invaluable tool for population and evolutionary genomics applications.

The results of our analyses of an existing synthetic dataset[24] generated by simulations of a

polygenic trait under various spatial patterns of selection showed that DeepGenomeScan can

successfully identify outlier loci exhibiting complex non-linear associations with the target

trait. This represents an important advance over existing methods that can only consider

monotonic associations. For example, SPA uses a logistic function to describe allele

frequency behaviour as a function of geographic location and can only detect outlier loci

exhibiting monotonic geographic gradients [20]. Moreover, it cannot be applied to study

associations with phenotypic or environmental data.

Evolutionary biologists interested in understanding the mechanisms driving local adaptation

and speciation follow a longstanding tradition focused on the analysis of variation in

phenotype or allele frequencies along linear environmental gradients [26,38]. Here we showed

that recent machine learning methods such as pcadapt and RDA can successfully identify

genomic regions associated with those patterns (Figs. 2 & 3). However, their performance is

limited in the case of non-monotonic gradients and, moreover, they are completely incapable

of considering coarse-grained heterogeneous spatial patterns in allele frequencies (Fig 2 & 3).

Our method clearly outperforms pcadapt and RDA in uncovering the outlier loci that

contribute the most to spatial population structure, regardless of the underlying spatial pattern

in selective pressures. Nevertheless, it is clear that very coarse-grained heterogenous

selection with no clear spatial pattern represents a very difficult scenario, which reduces the

power of our method to detect genomic regions associated with the response of a species to

selection.

Although less well studied empirically, coarse-grained heterogenous selection with no clear

spatial pattern has been an important focus of theoretical studies aimed at explaining

observed patterns of genetic variation [39-44]. This scenario is particularly relevant to studies of

protected polymorphisms [40,42,44] and hard versus soft selection [41]. Additionally, the

importance of this type of selection pattern is particularly relevant for genome-wide

association studies focused on diseases and phenotypic characters that do not exhibit clear

spatial patterns. Deep learning genome-scan methods capable of uncovering genomic regions

associated with this type of selection pattern can, therefore, provide a more general

understanding of how prevalent these particular mechanisms are.

We also applied DeepGenomeScan to the Population Reference Sample [30] (POPRES) dataset

to determine if our method could uncover genomic regions undetected by SPA and other

popular methods such iHS [45], Fst [10], Bayenv [13]. Comparing Yang et al.'s [20] Supplementary

Table 4 with our Supplementary Table 1, reveals a large number of important genes that were

not detected by those other existing methods. Some of these genes have been linked to

complex diseases such as Parkinson, obesity or various types of cancer and may not be

associated with monotonic selection gradients.

DeepGenomeScan can be applied to spatial genetic data even when individuals' geographic

coordinates are not available. In this case, our method uses the first two reduced features

obtained from a very recent machine learning method (KLFDAPC[29]), as proxies for latitude

and longitude. A comparison of the results from analyses of the POPRES dataset based on

geographic coordinates with those based on the first two reduced features reveals very

extensive overlap in terms of the genes they uncover. The small discrepancy in genes

uncovered by the two options implemented in our software is due to differences in the type of

spatial information used. It is clear that reduced features could lead to a loss of detailed

spatial information provided by geographic location. On the other hand, the use of reduced

features could avoid biases introduced by low quality geographic information or could take

into account small scale spatial effects that are not detectable at the spatial scale used for the

geographic coordinates.

There are still several methodological challenges faced when implementing deep learning

methods, which are associated with the computational cost of hyperparameter tuning and

neural network training. However, very recent adaptive resampling algorithms (c.f. ref. [18]),

which we implemented in our software, carry out an efficient exploration of hyperparameter

space, allowing the use of deep learning to analyse large population genomic data sets.

Although the application of deep learning to population genetics problems is still in its

infancy [18], there are already some examples of such applications [21,46,47]. One of these

applications is aimed at distinguishing between hard and soft sweeps and simultaneously

incorporating the confounding effects of demographic history [21]. This application involves

choosing among two or more alternative models, an objective that is usually achieved using

Approximate Bayesian Computation (ABC) [48,49]. Although model choice in ABC is akin to a

classification application in machine learning, neural networks require a training set that is

not readily available when the target is a demographic or selection model. To overcome this

limitation, the authors train the neural network using synthetic data generated under

predefined evolutionary models. A drawback of this strategy is that it introduces model

assumptions into a computational framework that could be completely free of them and,

therefore, of very wide applicability. Additionally, instead of using the raw genotype matrix,

this framework requires the pre-processing of genetic data to generate summary statistics, as

input data, leading to a loss of information. These drawbacks are absent when the objective is

to assign individuals to geographic locations as in ref. [46] or when scanning the genome in

search of outlier loci, as in our study. Overall, DeepGenomeScan fully exploits the flexibility

and power of deep neural networks, which makes it applicable to a wide range of problems

including identification of genomic regions associated with diseases or economically

important phenotypic traits, assignment of individuals to geographic locations, or

identification of environmental factors associated with selective pressures.

Methods

Methods

MLP Architecture

In this study, we constructed a Multilayer Perceptron Network with three hidden layers

(Figure 1 and Supplementary Notes Figure S1). To describe this MLP, we first assume that

the genotypes of the n sampled individuals at p loci are described by a n×p matrix, $\mathbf{x} = (x_{ij})$,

and are coded by the number of derived alleles, which in the case of a biallelic marker takes

values $x_{ij} = 0, 1, 2$, where i = 1, 2, ..., n, and j=1, 2, ..., p. Furthermore, the trait values of all

individuals are arranged in a n×1 vector $\mathbf{y} = (y_i)$. Then the input layer of the MLP consists

of p nodes, each one representing a distinct locus. The information contained in node j

comprises the individual genotypes at locus j, $\mathbf{x}_{*j} = (x_{1j}, \cdots, x_{nj})$. This information is

transformed into a signal which is fed to the first hidden layer. The signal received by each

node k = 1, 2, …, K_1 of the first hidden layer $\hat{\mathbf{s}}^{k1} = (\hat{s}_1^{k1}, \cdots, \hat{s}_i^{k1}, \cdots, \hat{s}_n^{k1})$ represents a

weighted non-linear regression of individuals' trait values on their multilocus genotypes.

Here, the superscript k1 identifies the k^{th}-node of the first hidden layer. Each element $\hat{s}_i^{k1}$ of

the signal vector received by a node k of the first hidden layer is given by

$\hat{s}_i^{k1} = f^1(\mathbf{x}_{i*}) = G\left(\sum_{j=1}^p x_{ij} w_j^{k1} + b^{k1}\right) = G(\mathbf{x}_{i*}\mathbf{w}^{k1} + b^{k1}),$ (1)

where w_j^{k1} is the weight of locus j, b^{k1} is a scalar representing a bias in the signal received

by node k in the first hidden layer, and $G(\cdot)$ is the activation function used to transform the

information sent by the input layer to the first hidden layer.

In a similar way, each node k of the second hidden layer will take the output signal of all

nodes of the first hidden layer and apply a transformation before sending the signal to the

third hidden layer:

$\hat{s}_i^{k2} = f^2(\hat{\mathbf{s}}_i^{k1}) = G\left(\sum_{m=1}^{K_1} \hat{s}_i^{k1} w_m^{k2} + b^{k2}\right) = G(f^1(\mathbf{x}_{i*})\mathbf{w}^{k2} + b^{k2}).$ (2)

If there are only two hidden layers, then the neural network is represented by $f(\mathbf{x}) =$

$f^2(f^1(\mathbf{x}))$. In general, there can be an arbitrary number L of hidden layers, each consisting of

K_l nodes and the signal generated by the last hidden layer is the vector of predicted trait

values, $\hat{\mathbf{y}} = (\hat{y}_i)$. Therefore, a MLP network can be represented by,

$f(\mathbf{x}) = f^L(f^{L-1}(\cdots f^2(f^1(\mathbf{x})))),$ (3)

where $\mathbf{f^l}(\cdot)$ is the signal received by the l^{th} layer.

Given the input data, i.e. the genotypes, and the observed trait values, the neural network

learns the weights $\mathbf{w} = (\mathbf{w}^1, ..., \mathbf{w}^L)$ with $\mathbf{w}^l = (w^{1l}, \cdots, w^{K_l l})$ and biases $\mathbf{b} =$

$(\mathbf{b}^1, ..., \mathbf{b}^L)$ with $\mathbf{b}^l = (b^{1l}, \cdots, b^{K_l l})$ that best describe the relationship between the inputs and

the outputs by minimising the difference between predicted and observed trait values [50,51].

These weights represent an essential element of our genome scan method because they are

used to identify outlier loci (see below) using a procedure supported by previous studies [23,52],

which show that the importance of a variable (locus) can be estimated as the combination of

the absolute values of weights associated with the graph edges connecting the predictor

variable (locus) with the MLP output (predicted trait).

Implementation

DeepGenomeScan trains the deep neural network to estimate the nonlinear function mapping

genotypes to individual traits. This process involves not only learning the vectors of weights

and biases by iteratively adjusting these parameters but also the tuning of a large number of

hyperparameters associated with structural components of the model, which impact model

performance. These include those determined by the network architecture (activation

functions, number of neurons per hidden layer) and those included in the optimisation

algorithm used for training (e.g. learning rate, batch size). For optimisation, we used resilient

backpropagation [53] with weight backtracking when analysing both simulated and real

datasets. However, the tuning algorithm used depended on the size of the input data. In the

case of simulated datasets, which were of limited size, we used a full resampling strategy

corresponding to Algorithm 1 in ref. [18] based on a repeated k-fold cross validation with k = 5

and five repetitions. In the case of the large POPRES dataset, we used an adaptive resampling

approach corresponding to Algorithm 2 in ref. [18], which incorporates futility assessment into

the model tuning process. The resampling approach in this case was the same as in the

analysis of simulated data (five replications of a five-fold cross-validation). Detailed

information about the settings used for the optimisation and tuning algorithms is presented in

Supplementary Notes.

Identification of outlier loci

As mentioned before, the SNP importance can be estimated as combinations of the absolute values of connection weights (**w**) [23,52]. This is done once the optimal model is found. We used Olden and Jackson's [19] method, which is based on Garson's[54] algorithm to calculate the relative importance for each input node but adds a randomisation step to identify non-significant connection weights.

DeepGenomeScan carries out separate runs for each trait and generates a vector of SNP importance for each of them. Given T traits, the position of each SNP in trait space is described by its associated vector of importance values and, therefore, it is possible to calculate the Mahalanobis distance between the focal locus and all other loci. This leads to an intuitive definition of outlier loci as any locus with an "extreme" Mahalanobis distance. Since the squared Mahalanobis distance follows a chi-squared distribution with T degrees of freedom, the p-values associated with each SNP can be obtained from a χ_T^2 distribution [55] and used for identifying 'outlier' loci.

Human dataset

We applied DeepGenomeScan to European populations from the POPRES project [30]. The dataset contains a total of 3,192 European individuals genotyped at 500,568 loci using the Affymetrix 500K SNP chip. The sample collections and genotyping for POPRES are described in [30]. We removed individuals from outside of Europe and individuals whose grandparents had different geographic origins based on the criteria used by [28]. We also

removed samples that only have one individual per country, such as Denmark, Finland,

Latvia, Ukraine, Slovakia and Slovenia. Geographic coordinates for each individual

corresponded to the central point of the geographic area of the individual's country of birth.

We removed the monomorphic SNPs and filtered out autosomal biallelic SNPs with a minor

allele frequency (MAF) below 5%, and a missing rate of 5% or more. In the end, we kept 1,

382 individuals from 32 countries carrying 283, 499 autosomal SNPs. We annotated and

updated the variant reference ID based on the comprehensive report of short human

variations from the human variation database (dbSNP, GRCh37p13, b151, release 2018,

https://ftp.ncbi.nih.gov/snp/organisms/human_9606/VCF/) using bcftools [56].

Data and code availability

The simulation data used in this study is available at

https://datadryad.org/review?doi=doi:10.5061/dryad.1s7v5.

The datasets used for the analyses described in this manuscript were obtained from dbGap at

https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000145.v4.p2

through dbGap accession number phs000145.v4.p2 (request approval #90291-1).

The DeepGenomeScan R code is hosted on GitHub at

https://github.com/xinghuq/DeepGenomeScan. The scripts used in this study are available at

https://github.com/xinghuq/DeepGenomeScan/tree/webpkg/DeepGenomeScan_simulation%2

BPOPRES. The KLFDAPC package is available at https://xinghuq.github.io/KLFDAPC/.

Acknowledgments

XQ was supported by a PhD scholarship from the China Scholarship Council.

Author Contributions

XQ and OEG designed the study, XQ carried out the analyses and interpreted results with input from OEG and CWKC. OEG and XQ wrote the article with input from CWKC.

Competing Interests

The authors declare no competing financial interests.

References

1. Bush, W.S. & Moore, J.H. Chapter 11: Genome-Wide Association Studies. *Plos Computational Biology* **8**(2012).

2. Kang, H.M. *et al.* Variance component model to account for sample structure in genome-wide association studies. *Nature Genetics* **42**, 348-U110 (2010).

3. Tam, V. *et al.* Benefits and limitations of genome-wide association studies. *Nature Reviews Genetics* **20**, 467-484 (2019).

4. Edge, M.D. & Coop, G. Reconstructing the History of Polygenic Scores Using Coalescent Trees. *Genetics* **211**, 235-262 (2019).

5. Field, Y. *et al.* Detection of human adaptation during the past 2000 years. *Science* **354**, 760-764 (2016).

6. Racimo, F., Berg, J.J. & Pickrell, J.K. Detecting Polygenic Adaptation in Admixture Graphs. *Genetics* **208**, 1565-1584 (2018).

7.   Speidel, L., Forest, M., Shi, S.N. & Myers, S.R. A method for genome-wide
genealogy estimation for thousands of samples. *Nature Genetics* **51**, 1321-+ (2019).

8.   Turchin, M.C. *et al.* Evidence of widespread selection on standing variation in Europe
at height-associated SNPs. *Nature Genetics* **44**, 1015-+ (2012).

9.   Oleksyk, T.K., Smith, M.W. & O'Brien, S.J. Genome-wide scans for footprints of
natural selection. *Philosophical Transactions of the Royal Society B-Biological*
Sciences **365**, 185-205 (2010).

10.   Lewontin, R.C. & Krakauer, J. Distribution of gene frequency as a test of the theory
of the selective neutrality of polymorphisms. *Genetics* **74**, 175-95 (1973).

11.   Beaumont, M.A. & Balding, D.J. Identifying adaptive genetic divergence among
populations from genome scans. *Molecular Ecology* **13**, 969-980 (2004).

12.   Foll, M. & Gaggiotti, O. A Genome-Scan Method to Identify Selected Loci
Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective.
Genetics **180**, 977-993 (2008).

13.   Coop, G., Witonsky, D., Di Rienzo, A. & Pritchard, J.K. Using Environmental
Correlations to Identify Loci Underlying Local Adaptation. *Genetics* **185**, 1411-1423
(2010).

14.   de Villemereuil, P. & Gaggiotti, O.E. A new F-ST-based method to uncover local
adaptation using environmental variables. *Methods in Ecology and Evolution* **6**, 1248-
1258 (2015).

15.   Frichot, E., Schoville, S.D., Bouchard, G. & François, O. Testing for associations
between loci and environmental gradients using latent factor mixed models.
Molecular biology and evolution **30**, 1687-1699 (2013).

16.   Gaggiotti, O.E. *et al.* Disentangling the effects of evolutionary, demographic, and
environmental factors influencing the genetic structure of natural populations:
Atlantic herring as a case study. *Evolution* **63**, 2939-2951 (2009).

17.   Hornik, K., Stinchcombe, M. & White, H. Multilayer feedforward networks are
universal approximators. *Neural networks* **2**, 359-366 (1989).

18.   Kuhn, M. Futility analysis in the cross-validation of machine learning models.
arXiv:1405.6974 (2014).

19.   Olden, J.D. & Jackson, D.A. Illuminating the "black box": a randomization approach
for understanding variable contributions in artificial neural networks. *Ecological*
modelling **154**, 135-150 (2002).

20.   Yang, W.Y., Novembre, J., Eskin, E. & Halperin, E. A model-based approach for
analysis of spatial structure in genetic data. *Nature Genetics* **44**, 725-U163 (2012).

21.   Sheehan, S. & Song, Y.S. Deep Learning for Population Genetic Inference. *Plos*
Computational Biology **12**(2016).

22.   Gevrey, M., Dimopoulos, L. & Lek, S. Review and comparison of methods to study
the contribution of variables in artificial neural network models. *Ecological Modelling*
160, 249-264 (2003).

23.   Olden, J.D., Joy, M.K. & Death, R.G. An accurate comparison of methods for
quantifying variable importance in artificial neural networks using simulated data.
Ecological modelling **178**, 389-397 (2004).

24.   Capblancq, T., Luu, K., Blum, M.G. & Bazin, E. Evaluation of redundancy analysis
to identify signatures of local adaptation. *Molecular Ecology Resources* **18**, 1223-
1233 (2018).

25.   Luu, K., Bazin, E. & Blum, M.G. pcadapt: an R package to perform genome scans for
selection based on principal component analysis. *Molecular ecology resources* **17**, 67-
77 (2017).

26.   Endler, J.A. *Geographic variation, speciation and clines*, (Princeton University Press,
Princeton, NJ, 1977).

27.   Lao, O. *et al.* Correlation between genetic and geographic structure in Europe.
Current Biology **18**, 1241-1248 (2008).

28.   Novembre, J. *et al.* Genes mirror geography within Europe. *Nature* **456**, 98-101
(2008).

29.   Qin, X., Chiang, C.W.K. & Gaggiotti, O.E. Kernel Local Fisher Discriminant
Analysis of Principal Components (KLFDAPC) significantly improves the accuracy
of predicting geographic origin of individuals. *bioRxiv, 2021.2005.2015.444294*
(2021).

30.   Nelson, M.R. *et al.* The Population Reference Sample, POPRES: a resource for
population, disease, and pharmacological genetics research. *The American Journal of*
Human Genetics **83**, 347-358 (2008).

31.   Granovsky, M. *et al.* Suppression of tumor growth and metastasis in Mgat5-deficient
mice. *Nature medicine* **6**, 306-312 (2000).

32.   Brynedal, B. *et al.* MGAT5 alters the severity of multiple sclerosis. *Journal of*
neuroimmunology **220**, 120-124 (2010).

33.   Wang, R. *et al.* Hydrogen sulfide demonstrates promising antitumor efficacy in
gastric carcinoma by targeting MGAT5. *Translational oncology* **11**, 900-910 (2018).

34.   Fox, C.S. *et al.* Genome-wide association for abdominal subcutaneous and visceral
adipose reveals a novel locus for visceral fat in women. *PLoS Genet* **8**, e1002695
(2012).

35. Nalls, M.A. *et al.* Large-scale meta-analysis of genome-wide association data
identifies six new risk loci for Parkinson's disease. *Nature genetics* **46**, 989-993
(2014).

36. Dichgans, M. *et al.* Shared genetic susceptibility to ischemic stroke and coronary
artery disease: a genome-wide analysis of common variants. *Stroke* **45**, 24-36 (2014).

37. Comuzzie, A.G. *et al.* Novel genetic loci identified for the pathophysiology of
childhood obesity in the Hispanic population. *PloS one* **7**, e51954 (2012).

38. Haldane, J.B.S. THE THEORY OF A CLINE. *Journal of Genetics* **48**, 277-284
(1948).

39. Bulmer, M.G. Multiple niche polymorphism. *Amer. Natur.* **106**, 254-257 (1972).

40. Levene, H. Genetic equilibrium when more than one ecological niche is available.
Amer. Natur. **87**, 331-333 (1953).

41. Levins, R. & MacArthur, R. The maintenance of genetic polymorphism in a spatially
heterogeneous environment: variations on a theme by Howard Levene. *Americal
Naturalist* **100**, 585-589 (1966).

42. Prout, T. SUFFICIENT CONDITIONS FOR MULTIPLE NICHE
POLYMORPHISM. *American Naturalist* **102**, 493-& (1968).

43. Smith, J.M. GENETIC POLYMORPHISM IN A VARIED ENVIRONMENT.
American Naturalist **104**, 487-& (1970).

44. Strobeck, C. Haploid selection withn alleles in m niches. *American Naturalist* **113**,
439-444 (1979).

45. Sabeti, P.C. *et al.* Detecting recent positive selection in the human genome from
haplotype structure. *Nature* **419**, 832-837 (2002).

46. Battey, C.J., Ralph, P.L. & Kern, A.D. Predicting geographic location from genetic
variation with deep neural networks. *Elife* **9**(2020).

47. Flagel, L., Brandvain, Y. & Schrider, D.R. The Unreasonable Effectiveness of
Convolutional Neural Networks in Population Genetic Inference. *Molecular Biology
and Evolution* **36**, 220-238 (2019).

48. Beaumont, M.A., Zhang, W.Y. & Balding, D.J. Approximate Bayesian computation
in population genetics. *Genetics* **162**, 2025-2035 (2002).

49. Csillery, K., Blum, M.G.B., Gaggiotti, O.E. & Francois, O. Approximate Bayesian
Computation (ABC) in practice. *Trends in Ecology & Evolution* **25**, 410-418 (2010).

50. Jinming, Y., Ahmadi, M., Jullien, G.A. & Miller, W.C. Model validation and
determination for neural network activation function modeling. 548-551 (1998).

51. Goodfellow, I.a. Deep learning / Ian Goodfellow, Yoshua Bengio and Aaron
Courville. 0-5 (The MIT Press, Cambridge, Massachusetts :, 2016).

52. Gevrey, M., Dimopoulos, I. & Lek, S. Review and comparison of methods to study
the contribution of variables in artificial neural network models. *Ecological modelling*
160, 249-264 (2003).

53. Riedmiller, M. Advanced supervised learning in multi-layer perceptrons—from
backpropagation to adaptive learning algorithms. *Computer Standards & Interfaces*
16, 265-278 (1994).

54. Garson, D.G. Interpreting neural network connection weights. *Artificial Intelligence*
Expert **6**, 46-51 (1991).

55. Filzmoser, P. & Gregorich, M. Multivariate Outlier Detection in Applied Data
Analysis: Global, Local, Compositional and Cellwise Outliers. *Mathematical*
Geosciences **52**, 1049-1066 (2020).

56. Li, H. A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data.
Bioinformatics **27**, 2987-2993 (2011).