TCAB1 prevents nucleolar accumulation of the telomerase RNA to

promote telomerase assembly

Basma S. Al-Masraf [1,2,3], Gloria I. Perez [1], Kate Adams-Boone [1], Scott B. Cohen [4], Li

Han [5], Kefei Yu [5], Jens C. Schmidt [1,6,*]

[1] Institute for Quantitative Health Sciences and Engineering, Michigan State University,

East Lansing, MI, U.S.A.

[2] College of Osteopathic Medicine, Michigan State University, East Lansing, MI, U.S.A.

[3] Cellular and Molecular Biology Graduate Program, College of Natural Sciences,

Michigan State University, East Lansing, MI, U.S.A.

[4] Children's Medical Research Institute and University of Sydney, Westmead, NSW

2145, Australia.

[5] Department of Microbiology and Molecular Genetics, Michigan State University, East

Lansing, MI, U.S.A.

[6] Department of Obstetrics, Gynecology, and Reproductive Biology.

[*] Correspondence: schmi706@msu.edu

Abstract

Localization of a wide variety of RNAs to non-membrane bound cellular compartments

such as nucleoli, Cajal bodies, and stress-granules is critical for their function and

stability. The molecular mechanisms that underly the recruitment and exclusion of specific

RNAs from these phase-separated organelles is poorly understood. Telomerase is a

ribonucleoprotein (RNP), that is composed of the reverse transcriptase protein TERT, the

telomerase RNA (TR), and several auxiliary proteins that associate with TR, including

TCAB1. Here we show that, that in the absence of TCAB1, TR is trapped in the nucleolus,

while TERT localizes to the nucleoplasm and is excluded from the nucleolus, which

prevents telomerase assembly. Thus, nucleolar phase separation constitutes a barrier

that counteracts telomerase assembly and TCAB1 is required to exclude the telomerase

RNA from the nucleolus. Our work provides general insight into the mechanism and

functional consequences of RNA recruitment to non-membrane bound organelles and

proposes a new model explaining the critical role of TCAB1 in telomerase function.

Introduction

Human cells contain a number of non-membrane bound organelles that carry out critical cellular functions. For instance, nucleoli and Cajal bodies are phase-separated nuclear organelles that play important roles in the biogenesis and maturation of many cellular RNAs (Hyman et al., 2014; Mitrea and Kriwacki, 2016). Nucleoli and Cajal bodies contain a wide range of small nucleolar and small Cajal body-specific RNAs (snoRNAs and scaRNAs, respectively). A subset of these snoRNAs and scaRNAs are bound by the H/ACA complex, which contains NOP10, NHP2, GAR1, and the pseudouridylase dyskerin, which modifies ribosomal and spliceosomal RNA precursors and other RNAs (Angrisani et al., 2014). A key difference between snoRNAs and scaRNAs is the presence of the Cajal-body box (CAB-box) motif in scaRNAs that directly associates with the telomerase Cajal body protein 1 (TCAB1, also known as WRAP53) (Jády et al., 2004; Schmidt and Cech, 2015; Venteicher et al., 2009). TCAB1 is required for the recruitment of scaRNAs to Cajal bodies and in its the absence scaRNAs localize to the nucleolus (Venteicher et al., 2009). Therefore, TCAB1 controls which phase-separated nuclear organelle scaRNAs associate with. Importantly, the molecular mechanism by which TCAB1 drives exclusion of scaRNAs from the nucleolus and facilitates their recruitment to Cajal bodies is unknown. In addition, it is unclear whether miss-localization of scaRNAs to the nucleolus has functional consequences.

The telomerase RNA (TR) is a scaRNA and like other scaRNAs its association with nucleoli and Cajal bodies is controlled by TCAB1 (Schmidt and Cech, 2015). Telomere maintenance by telomerase is essential for continuous proliferation of stem cell

populations in the human body and most cancers require telomerase activity for their

survival (Stewart and Weinberg, 2006). To compensate for the incomplete replication of

chromosome ends, telomerase appends TTAGGG repeats to the telomeric single-

stranded overhang (Schmidt and Cech, 2015). Telomerase-mediated telomere

maintenance requires three critical steps: Telomerase assembly, telomerase recruitment

to telomeres, and telomeric repeat synthesis (Schmidt and Cech, 2015). Mutations in

several genes have been identified that cause deficiencies in one of these critical steps

and lead to a variety of diseases known as telomere syndromes, characterized by

premature depletion of stem cell populations (Armanios and Blackburn, 2012). In addition,

telomerase is inappropriately activated in >85% of cancers (Stewart and Weinberg,

2006). While telomerase recruitment to telomeres (Nandakumar and Cech, 2013) and

telomerase catalysis (Wu et al., 2017) have been studied extensively, much less is known

about telomerase assembly. Importantly, telomerase assembly could be targeted to

reduce telomerase activity in cancer cells, or to increase telomerase function in patients

affected by genetically defined telomerase deficiency syndromes (Nagpal et al., 2020;

Shukla et al., 2020).

Telomerase is a complex ribonucleoprotein (RNP). The core components of telomerase

are the telomerase reverse transcriptase (TERT) protein, TR, the H/ACA complex, and

TCAB1 (Schmidt and Cech, 2015). The primary function of the H/ACA complex is to

stabilize TR, by directly binding to its 3'-end, preventing the exonucleolytic degradation of

TR (Stuart et al., 2015; Tummala et al., 2015). The 3'-end formation of TR is tightly

regulated by the competing activities of the poly-(A) polymerase PAPD5 and the nuclease

PARN (Shukla et al., 2016; Tseng et al., 2015). Loss of TCAB1 function leads to telomere

attrition in a variety of cell lines (Chen et al., 2018; Venteicher et al., 2009; Vogan et al.,

2016; Zhong et al., 2011). In addition, multiple mutations in TCAB1 have been identified

that cause misfolding of TCAB1 and lead to dyskeratosis congenita, a telomere syndrome

(Freund et al., 2014; Zhong et al., 2011). While these observations highlight that TCAB1

is necessary for telomere maintenance, the underlying molecular mechanism is unclear.

Initially, it was proposed that TCAB1 is required for telomerase recruitment to telomeres

(Stern et al., 2012; Venteicher et al., 2009). A more recent study suggested that TCAB1

is required for the correct folding of TR, and that its absence causes a reduction in

telomerase activity (Chen et al., 2018). Importantly, all previous studies have come to the

conclusion that TCAB1 is not required for telomerase assembly.

Here we analyze telomerase assembly in intact cells and by purification of the telomerase

RNP and demonstrate that, contrary to previous findings, TCAB1 promotes telomerase

assembly *in vivo.* Using a combination of genetic perturbations, biochemical analysis of

telomerase function, cell biological approaches, and single-molecule live cell imaging, we

show that in the absence of TCAB1, TR is sequestered in the nucleolus while TERT is

excluded from the nucleolus. The spatial separation of TERT and TR that we observe in

our experiments is inconsistent with proper telomerase assembly. Furthermore, we show

that the limited amount of telomerase that can assemble in the absence of TCAB1 is fully

active and can localize to telomeres, suggesting that TCAB1 is not required for the

enzymatic function of telomerase or its recruitment to telomeres. In addition, we

demonstrate that the diffusion properties of TERT in cells that lack TCAB1 closely

resemble those of TERT that is not bound to TR. We conclude that the sequestration of

TR in the nucleolus, when TCAB1 is absent, prevents its association with TERT, and that

this lack of telomerase assembly is the molecular mechanism underlying the critical role

of TCAB1 in telomere maintenance. Our results also demonstrate that the nucleolar

phase separation constitutes a barrier for telomerase assembly and suggest that

incompletely assembled RNPs are tightly associated with the nucleolus and do not readily

enter the nucleoplasm.

Results

Loss of TCAB1 leads to nucleolar accumulation of TR

To confirm that TR is sequestered in the nucleolus in the absence of TCAB1, we attempted to knock-out TCAB1 in parental HeLa cells and HeLa cells expressing 3xFLAG-HaloTag-TERT and mEos3.2-TRF2 from the endogenous *TERT* and *TRF2* loci (referred to as Halo-TERT from here on) using Cas9 and a single guide RNA targeting the first exon of the *TCAB1* locus (Figure 1 – figure supplement 1). We isolated clonal cell lines that lacked a wild-type *TCAB1* allele (Figure 1 – figure supplement 1) and analyzed TCAB1 expression using Western blot. While a TCAB1 antibody targeting the N-terminus of TCAB1 failed to detect TCAB1 in our knock-out clones (Figure 1 – figure supplement 1), antibodies raised against more C-terminal TCAB1 fragments detected a truncated form of TCAB1 (Figure 1 – figure supplement 1). The size of the truncated TCAB1 is consistent with the use of methionine 129 as an alternative start codon. Importantly, the truncated form of TCAB1 localizes to Cajal bodies (Figure 1 – figure supplement 1) and scaRNA localization is unaffected in its presence (Figure 1 – figure supplement 1), indicating that it is at least partially functional. To avoid expression of the truncated TCAB1 variant, we knocked out TCAB1 using Cas9 with two guide RNAs to delete exons 2 and 3 from the TCAB1 gene, which removes the coding sequence for residues 144-214 of TCAB1 and results in a frame shift (Figure 1 – figure supplement 2). TCAB1 knock-out was validated by Southern blot, PCR, Western blot, and immunofluorescence imaging (IF, Fig. 1A-C, Figure 1 – figure supplement 2). Fluorescence *in situ* hybridization (FISH) demonstrated that TR accumulates in the nucleolus in cells that lack TCAB1, as indicated by co-localization of TR and nucleolar dyskerin signals. Importantly, expression of GFP-

134 TCAB1 in TCAB1 knock-out cells rescued TR localization to Cajal bodies (Figure 1 –

135 figure supplement 2), confirming that the miss-localization of TR to nucleoli is caused by

136 absence of TCAB1. These observations confirm that TCAB1 is required to prevent TR

137 accumulation in the nucleolus (Fig. 1C).

Figure 1. TR is localized to nucleoli in TCAB1 knock-out cells. (A-B) Western blot demonstrating the absence of TCAB1 protein in TCAB1 knock-out cell lines generated from **(A)** HeLa and **(B)** Halo-TERT parental cell lines (probed with Proteintech TCAB1 antibody). **(C)** Immuno-fluorescence with anti-dyskerin and anti-TCAB1 antibodies coupled to fluorescence in-situ hybridization with probes against TR, demonstrating the absence of TCAB1 and TR localization to nucleoli in TCAB1 knock-out cells (scale bar = 5 µm).

TERT is excluded from nucleoli

Our previous observations demonstrated that TERT does not enter nucleoli in human

cancer cells (Schmidt et al., 2016). To confirm these results, we performed single-

molecule imaging of 3xFLAG-HaloTag-TERT in living HeLa cells. Consistent with our

previous results, 3xFLAG-HaloTag-TERT was not observed entering or overlapping with

nucleoli, which are discernible as circular shapes in the nucleus under transmitted light

Figure 2. TERT is excluded from nucleoli in control and TCAB1 knock-out cells. (A) Maximum intensity projections of 2000 frames of 3xFLAG-HaloTag (JF646) TERT movies (bottom), demonstrating that the TERT signal does not overlap with the nucleolus detected as circular shape in the transmitted light image in control and TCAB1 knock-out cells (top, red dashed line, scale bar = 2 µm). **(B)** TERT particle trajectories from the cells shown in Fig. 2A, demonstrating that TERT molecules move parallel to or away from the nucleolus when located at the interface between the nucleolus and nucleoplasm.

illumination (Fig. 2A,B, Figure 2 – figure supplement 1, Movie 1,2). TERT trajectories in

close proximity to nucleoli revealed that their movement is biased away from the

nucleolus, suggesting that TERT is repelled by the nucleolus (Fig. 2B). To exclude the

possibility that nucleolar exclusion is a consequence of the 3xFLAG-HaloTag on the N-

terminus of TERT used in our experiments, we transiently expressed the 3xFLAG-

HaloTag fused to a nuclear localization sequence (NLS) in HeLa cells. Single-molecule

imaging demonstrated that the nuclear 3xFLAG-HaloTag signals overlapped with the

nucleolus (Figure 2 – figure supplement 1, Movie 3). Similar to the 3xFLAG-HaloTag

alone, 3xFLAG-HaloTag-dyskerin also localized to the nucleolus (Figure 2 – figure

supplement 1). These results demonstrate that 3xFLAG-HaloTag-TERT is excluded from

the nucleolus and that this exclusion is not caused by the 3xFLAG-HaloTag but instead

is an intrinsic property of the TERT protein.

TERT and TR localize to distinct nuclear compartments in cells lacking TCAB1

It is well established that telomerase assembly is incomplete in human cancer cells, which

leads to substantial pools of TERT and TR that are not assembled into telomerase RNPs

(Xi and Cech, 2014). Together with our observation that TERT molecules rarely enter the

nucleolus (Schmidt et al., 2016) this suggests that both free TERT and assembled

telomerase RNPs do not enter the nucleolus. To test whether, like TR, TERT accumulates

in the nucleolus in cells lacking TCAB1, we carried out single-molecule imaging of

3xFLAG-HaloTag-TERT. Strikingly, TERT localization is unchanged in cells lacking

TCAB1 (Fig. 2A,B). This suggests that unlike TR, TERT does not accumulate in nucleoli

when TCAB1 is absent.

The experiments described so far demonstrate that TCAB1 is enriched in nucleoli

and TERT is excluded from nucleoli in cells lacking TCAB1, but they do not

simultaneously detect TERT and TR in the same cell. To overcome this limitation, we

over-expressed mCherry-TERT and TR in TCAB1 knock-out cells. In controls, TERT and

TR colocalized with dyskerin at telomeres marked by TRF2 (Fig. 3A). In addition, analysis

of the TERT signal across the nucleus and nucleolus revealed that TERT was depleted

from the nucleolus (Fig. 3B). Importantly, mCherry-dyskerin localized to the nucleolus,

demonstrating that the mCherry-tag does not lead to nucleolar exclusion of its fusion

partner (Figure 3 – figure supplement 1). In cells lacking TCAB1, mCherry-TERT was

diffusely localized in the nucleoplasm, localized to a subset of telomeres, and was

depleted from the nucleolus (Fig. 3A,C). Similar to endogenous TR, overexpressed TR

180 was enriched in the nucleolus in TCAB1 knock-out cells (Fig. 3A). In addition, TR was

181 frequently co-localized

Figure 3. TERT and TR localize to distinct nuclear compartments in TCAB1 knock-out cells. (A) IF-FISH images of control and TCAB1 knock-out HeLa cells (3xFLAG-Halo-TERT, mEOS3.2-TRF2) overexpressing mCherry-TERT and TR (scale bar = 5 μm). Cells were probed with antibodies against dyskerin and mCherry, and FISH probes specific to TR. The intrinsic fluorescence of mEOS3.2-TRF2 was used to detect telomeres. TR and TERT are co-localized at telomeres in control cells. In TCAB1 knock-out cells, TR is enriched in nucleoli, while TERT is depleted from nucleoli in both control and TCAB1 knock-out cells. Both TERT and TR also localize to telomeres in TCAB1 knock-out cells. **(B-C)** Line scans of **(B)** control and **(C)** TCAB1 knock-out cells along the dashed white lines in Fig. 3A, demonstrating the enrichment of TR (green) in nucleoli (blue) in TCAB1 knock-out cells and the depletion of TERT (magenta) from nucleoli in both control and TCAB1 knock-out cells. **(D)** Quantification of the fraction of cells showing TR localization exclusively to telomeres, to telomeres and nucleoli, or only to nucleoli in cells over-expressing mCherry-TERT and TR (3 independent experiments, >100 cells per experiments, mean ± standard deviation).

with telomeres when mCherry-TERT and TR were overexpressed in cells that lack

TCAB1 (Fig. 3A,C-D). Similar TR localization patterns were found when overexpressing

untagged TERT and TR (Figure 3 – figure supplement 1) and untagged TERT was

excluded from nucleoli, confirming that nucleolar exclusion is an intrinsic property of

TERT (Figure 3 – figure supplement 1). Together these results demonstrate that, when

overexpressed in TCAB1 knock-out cells, TR is enriched in nucleoli and TERT is depleted

from nucleoli, consistent with a failure of nucleolar TR to assemble with TERT. In addition,

our observations suggest that overexpression can partially overcome the sequestration

of TR in the nucleolus when TCAB1 is absent, allowing a fraction of TR to bind to TERT

and localize to telomeres.

TCAB1 promotes telomerase RNP assembly

Previous studies by other laboratories have concluded that telomerase assembly is

unaffected by the absence of TCAB1 and whether TCAB1 is required for telomerase

activity is controversial (Chen et al., 2018; Venteicher et al., 2009; Vogan et al., 2016).

To assess the role of TCAB1 in telomerase assembly, we overexpressed TERT and TR

in parental HeLa and TCAB1 knock-out cells and immuno-purified TERT using a well-

established anti-TERT antibody (Cohen et al., 2007). To quantify telomerase assembly,

we measured TERT levels by Western blot and determined the amount of TR co-purified

using Northern blot (Fig. 4A,B). The amount of TR associated with TERT purified from

TCAB1 knock-out cells was significantly reduced compared to parental cells, for both

HeLa (31 $\pm$ 9%, mean $\pm$ SEM, p = 0.001) and Halo-TERT (33 $\pm$ 4%, mean $\pm$ SEM, p =

0.001) cell lines (Fig. 4D). As a second measure of telomerase RNP assembly we

quantified the amount of dyskerin associated with TERT (Fig. 4C). Since TR bridges

TERT and dyskerin, dyskerin co-purified with TERT directly reports on the presence of

TR. Consistent with the reduction in TR, the amount of dyskerin bound to TERT was also

significantly reduced when purified from cells lacking TCAB1 compared to parental

controls for both HeLa (22 $\pm$ 8%, mean $\pm$ SEM, p = 0.002) and Halo-TERT (32 $\pm$ 9%,

mean $\pm$ SEM, p = 0.004) cell lines (Fig. 4C,E). Importantly, we also confirmed that TCAB1

Figure 4. Telomerase Assembly is reduced in the absence of TCAB1. (A) Western blots analyzing TERT immuno-purification (using a sheep anti-TERT antibody) probed with a rabbit anti-TERT antibody (Abcam) and a TCAB1 antibody. **(B)** Northern blot of RNA extracted from input and purified TERT samples probed with three radiolabeled DNA oligonucleotides complementary to TR. Standards are *in vitro* transcribed full-length TR and truncated TR_{28-324}. TR_{28-324} was added to samples prior to RNA extraction as loading and recovery control. **(C)** Western blots to analyze immuno-purified telomerase RNP composition. A single membrane was cut into two pieces that were probed with TERT and dyskerin antibodies, respectively. **(D-F)** Quantification of the amount of **(D)** TR, **(E)** dyskerin, and **(F)** the ratio of TR to TERT in TERT purifications from TCAB1 knock-out cells compared to parental controls (n = 4-5, mean). The dashed lines indicate the level in telomerase purified from wild-type TCAB1 control cells which was normalized to 1.0.

is absent from telomerase purified from TCAB1 knock-out cells (Fig. 4A). As a final

measure of telomerase assembly, we determined the ratio between TR and TERT (Fig.

4A,B,F), which was also significantly reduced when TERT was purified from TCAB1

knock-out cells (HeLa: 18 $\pm$ 3%, mean $\pm$ SEM, p < 0.001, Halo-TERT: 17 $\pm$ 5%, mean $\pm$

SEM, p < 0.001). Altogether, these results demonstrate that telomerase assembly is

significantly reduced in the absence of TCAB1 and that overexpression of TERT and TR

is not sufficient to overcome this defect in telomerase RNP formation.

TCAB1 is not required for telomerase catalytic activity

Previous work by others has suggested that the defects in telomere maintenance

Figure 5. The specific activity of telomerase is unchanged in the absence of TCAB1. (A-B) Direct telomerase extension assay of telomerase immuno-purified from parental (WT) and TCAB1 knock-out (TKO) **(A)** HeLa and **(B)** Halo-TERT cell lines. LC1 and LC2, radiolabeled DNA oligonucleotide loading controls. In -Ab samples the TERT antibody was omitted during the immuno-purification. **(C)** Quantification of telomerase activity in samples from TCAB1 knock-out cells relative to parental controls (n = 4, mean). **(D)** Specific activity of telomerase purified from TCAB1 knock-out cells relative to parental controls (n = 4, mean). Specific activity was calculated by dividing the relative activity (see Fig. 5C) by the relative amount of TR present in immuno-purified TERT samples (see Fig. 4D). The dashed lines indicate the activity level in telomerase purified from wild-type TCAB1 control cells which was normalized to 1.0.

observed in TCAB1 knock-out cells are the consequence of reduced telomerase activity

caused by the miss-folding of the telomerase RNA rather than impaired telomerase

assembly (Chen et al., 2018). This TR folding model predicts that the specific activity of

telomerase is reduced. To determine the specific activity of telomerase purified from

TCAB1 knock-out cells, we measured telomerase activity using the direct telomerase

extension assays (Fig. 5A,B, Figure 5 – figure supplement 1), and divided the activity by

the amount of TR in the respective telomerase sample (Fig. 4B,D), which is a direct

measure of the quantity of telomerase RNP present. Total activity of telomerase purified

from TCAB1 knock-out cells was significantly reduced relative to telomerase purified from

parental controls telomerase (Fig. 5C, HeLa: $24 \pm 6\%$, mean $\pm$ SEM, $p < 0.001$, Halo-

TERT: $34 \pm 9\%$, mean $\pm$ SEM, $p = 0.005$), consistent with observations by others (Chen

et al., 2018). In contrast, the specific activity of telomerase derived from cells that lack

TCAB1 was not significantly reduced compared to controls (Fig. 5D, HeLa: $90 \pm 9\%$, mean

$\pm$ SEM, $p = 0.35$, Halo-TERT: $76 \pm 10\%$, mean $\pm$ SEM, $p = 0.1$). Together these

observations suggest that TCAB1 is required for telomerase assembly and that the limited

amount of telomerase RNP that can form in the absence of TCAB1 is fully active.

TCAB1 is required for telomerase assembly in living cells

The experiments presented thus far demonstrate that telomerase assembly is

significantly reduced in the absence of TCAB1 but were carried out in fixed cells or cell

lysates. To analyze telomerase assembly in intact cells, we carried out live cell single-

molecule imaging of 3xFLAG-HaloTag-TERT and determined the diffusion coefficient of

TERT particles (Fig. 6A, Movie 4, Figure 6 – figure supplement 1). The diffusion co-

efficient is a measure of the rate of movement of a molecule and depends on the size of

the complex it is part of and reports on molecular interactions formed with other sub-

cellular structures. Analysis of the diffusion coefficients of TERT trajectories in control

cells revealed three distinct populations of TERT particles (Fig. 6B-D). A static population

(D_S = 0.03 µm^2/s, 12%) which likely represents assembled telomerase RNPs bound to

telomeres or Cajal bodies, a slowly diffusion population (D_{F1} = 0.35 µm^2/s, 28%) and a

rapidly diffusing population (D_{F2} = 1.54 µm^2/s, 60%). The slowly diffusing population likely

includes assembled telomerase RNPs, while the rapidly diffusing particles represents

TERT molecules, which are not assembled with TR (Fig. 6B). Importantly, these diffusion

coefficients are largely consistent with our previous results using a distinct method to

determine their values (Schmidt et al., 2016). In the absence of TCAB1, the diffusion

coefficient of the freely diffusing TERT population was increased (D_{F2} = 1.78 $\pm$ 0.04 µm^2/s,

mean $\pm$ SEM, p = 0.04, Fig. 6C) and the fraction of the TERT populations that includes

assembled telomerase RNPs was significantly reduced ($F_{Slow+Static}$ = 29 $\pm$ 1%, mean $\pm$

SEM, p = 0.02, Fig. 6D). This observation is consistent with our model that in the absence

of TCAB1, telomerase assembly is defective. To determine the degree to which

telomerase assembly is affected in cells lacking TCAB1, we knocked out TR, completely

abolishing telomerase assembly. TR knock-out was confirmed by PCR and Sanger

sequencing, FISH, and qPCR (Figure 6 – figure supplement 1). Similar to control cells

and TCAB1 knock-out cells, TERT was also excluded from nucleoli in cells lacking TR

(Fig. S6D). Strikingly, the diffusion coefficients and the fraction of slow and static TERT

particles in cells lacking TR closely resembled those of TCAB1 knockout cells (Fig. 6A-

D, Figure 6 – figure supplement 1, Movie 4-5). To analyze the

Figure 6. Telomerase assembly is reduced in TCAB1 knock-out cells. (A) TERT particle trajectories from control, TCAB1 knock-out, and TR knock-out cells expressing 3xFLAG-HaloTag TERT (JF646, scale bar = 2 μm). (B) Diagram of distinct populations of TERT particles detected in control cells. (C) Diffusion coefficient of the rapidly diffusing TERT population in control, TCAB1 knock-out, and TR knock-out cells (3 independent experiments, >15 cells per experiment per cell line, mean ± standard deviation, complete data in Fig. S6D). (D) Fraction of slow plus static TERT particles in control, TCAB1 knock-out, and TR knock-out cells expressing 3xFLAG-HaloTag TERT (3 independent experiments, >15 cells per experiment per cell line, mean ± standard deviation, complete data in Fig. S6D). (D) Model for the regulation of telomerase assembly by TCAB1. In the absence of TCAB1, TR is sequestered in the dense fibrillar component (DFC) of the nucleolus, which is separated from the nucleoplasm by the granular component (GC) of the nucleolus.

interaction of TERT with telomeres, we filtered out TERT trajectories that came into

proximity with telomeres marked by mEOS3.2-TRF2, as previously described (Schmidt

et al., 2016). To assess the interaction of TERT with telomeres, we plotted the step-size

vs. the distance from the closest telomere for each step of these trajectories (Figure 6 –

figure supplement 2). In control cells, we observed an enrichment of smaller step sizes

and particles in close proximity to telomeres, consistent with TERT interactions with the

telomere (Figure 6 – figure supplement 2). In contrast, TERT trajectories from TCAB1

knock-out cells lacked this enrichment, and the step size vs. distance from the closest

telomere plots were identical to those from TR knock-out cells (Figure 6 – figure

supplement 2). In addition, diffusion analysis using SpotOn revealed that the fraction of

static TERT particles at telomeres was reduced from 12% in control cells to 4-5% in

TCAB1 and TR knock-out cells (Figure 6 – figure supplement 2). These observations

indicate that in the absence of either TCAB1 or TR, stable interactions of telomerase with

telomeres occur at a lower frequency because they require base pairing of TR with the

chromosome end (Schmidt et al., 2018). Together these results demonstrate that

telomerase assembly is strongly reduced in the absence of TCAB1.

Discussion

The experiments described in this study demonstrate that TCAB1 promotes telomerase

assembly. In the absence of TCAB1 the telomerase RNA is targeted to the nucleolus via

its association with dyskerin and other components of the H/ACA complex. In contrast to

TR, TERT cannot enter the nucleolus, preventing its association with TR in cells that lack

TCAB1. This demonstrates that nucleolar phase separation counteracts telomerase

assembly. In addition, we demonstrate that sequestration of TR in the phase-separated

nucleolus in the absence of TCAB1 can be partially overcome by telomerase

overexpression, leading to telomerase assembly and localization to telomeres. This

suggests that the nucleolus has a limited capacity to accommodate cellular RNAs and

that TCAB1 is not required for telomerase recruitment to telomeres. Finally, we

demonstrate that while telomerase assembly is limited the specific activity of telomerase

is unchanged in the absence of TCAB1, which excludes a role of TCAB1 in telomerase

catalytic function. Altogether our work completely reshapes our understanding of the role

of TCAB1 in telomerase function in human cells and provides insight into the role phase-

separated organelles play in RNP assembly and function.

TCAB1 promotes telomerase assembly

The importance of TCAB1 for telomere maintenance is undisputed (Chen et al., 2018;

Venteicher et al., 2009). Knock-out or depletion of TCAB1 results in telomere shortening

(Chen et al., 2018; Venteicher et al., 2009; Vogan et al., 2016). All previous work also

concluded that TR is enriched in the nucleolus in the absence of TCAB1 (Chen et al.,

2018; Stern et al., 2012; Venteicher et al., 2009; Vogan et al., 2016; Zhong et al., 2011).

Finally, all prior studies propose that TCAB1 is not required for telomerase assembly but

instead plays a role telomerase trafficking to Cajal bodies and telomeres or is required for

telomerase catalysis (Chen et al., 2018; Stern et al., 2012; Venteicher et al., 2009; Vogan

et al., 2016; Zhong et al., 2011). In contrast, our results demonstrate that in the absence

of TCAB1, TERT and TR are localized to the distinct sub-cellular compartments, the

nucleoplasm and the nucleolus, respectively. This spatial separation strongly reduces

telomerase assembly, which leads to reduced number of telomerase RNPs per cell and

in turn telomere shortening. Our conclusions are based on a variety of quantitative

approaches analyzing telomerase assembly by purification of the telomerase RNP and in

intact cells. Analysis of TERT localization in living and fixed cells clearly demonstrates

that TERT is excluded from the nucleolus. Importantly, we confirmed that the nucleolar

exclusion of TERT is not a consequence of the protein-tags used to visualize TERT in

these experiments. Purification of telomerase from TCAB1 knock-out cells further

confirmed that TCAB1 is required for telomerase assembly *in vivo*. The amount of TR and

dyskerin associated with TERT is substantially reduced in the absence of TCAB1. Finally,

we used live cell single-molecule imaging to demonstrate that the diffusion pattern of

TERT molecules and their interactions with telomeres are significantly different in TCAB1

knock-out cells compared to control cells. More importantly, TERT dynamics in the

absence of TCAB1 are comparable to those in TR knock-out cells where telomerase

assembly is impossible. Altogether, these observations support our model that TCAB1

promotes telomerase assembly by extracting TR from the nucleolus to facilitate its

assembly with TERT. Our model is further supported by work from Vogan *et al.*, which

demonstrated that truncated TR (hTRmin) that lacks the H/ACA region and therefore

cannot bind dyskerin, accumulates in the nucleoplasm and is excluded from nucleoli

(Vogan et al., 2016). Importantly, in cells that express hTRmin, TCAB1 is not required for

telomere maintenance (Vogan et al., 2016), consistent with TCAB1 promoting telomerase

assembly by extracting TR from the nucleolus.

TCAB1 is not required for telomerase catalysis

Previous work by others has reported conflicting results regarding the role of TCAB1 in

telomerase catalysis, ranging from full enzymatic activity in initial reports to substantial

activity defects in the most recent study (Chen et al., 2018; Venteicher et al., 2009; Vogan

et al., 2016). Importantly, both our work and the only other study that analyzed the role of

TCAB1 in telomerase activity using the "gold-standard" direct telomerase extension assay

concluded that telomerase activity is reduced to approximately 20-30% in the absence of

TCAB1. In both cases, telomerase was purified from TCAB1 knock-out cells using an

antibody raised against TERT prior to measuring telomerase activity. While both studies

concur on the degree to which telomerase activity is reduced in the absence of TCAB1,

the proposed underlying molecular mechanisms differ. Chen *et al.* propose that TCAB1

is required for proper folding of the CR4/CR5 region of the telomerase RNA, which directly

associates with TERT, without affecting telomerase assembly (Chen et al., 2018). Recent

structural analysis of the telomerase RNP from human cells revealed that TCAB1 is

located far away from the CR4/CR5 region of TR (Figure 6 – figure supplement 3)(Ghanim

et al., 2021). TCAB1 resides within the H/ACA lobe of while CR4/CR5 is located within

the catalytic lobe of the telomerase RNP. The two lobes are connected by the P1 and

P4.2 helices which can adopt multiple conformations. Importantly, TCAB1 was not located

near the CR4/CR5 region of TR in any of the observed telomerase RNP conformations

(Ghanim et al., 2021). Although it is possible that telomerase can adopt additional

conformations, based on the currently available structural information it is difficult to

rationalize a molecular mechanism by which TCAB1 could specifically promote CR4/CR5

folding. In addition, due to the miss-folding of TR telomerase was proposed to adopt a

low activity state in the absence of TCAB1. Experimentally such a low activity state would

manifest itself as a reduction in the specific activity of telomerase (telomerase activity per

assembled telomerase RNP). Our experiments strongly suggest that, while telomerase

assembly is reduced in the absence of TCAB1, the limited amount of telomerase that can

assemble is fully active (i.e. does not have reduced specific activity). Both, our enzymatic

analysis and the positioning of TCAB1 within the telomerase RNP do not support a role

of TCAB1 in TR folding and telomerase catalysis but are fully consistent with TCAB1

promoting telomerase assembly.

TCAB1 is not necessary for telomerase recruitment to telomeres

TCAB1 is necessary for the localization of scaRNAs to Cajal bodies and previous work

suggested that it is also required for telomerase recruitment to telomeres (Stern et al.,

2012; Venteicher et al., 2009). Our observations demonstrate that when TERT and TR

are overexpressed in TCAB1 knockout cells a fraction of TERT can assemble with TR an

localize to telomeres. Telomerase recruitment to telomeres requires a direct interaction

between TERT and TPP1 (Nandakumar et al., 2012; Schmidt et al., 2014; Zhong et al.,

2012), therefore TR that localizes to telomeres must be assembled with TERT. These

results are fully consistent with our model that the spatial separation of TR and TERT in

the absence of TCAB1 prevents telomerase assembly. When TR is overexpressed the

capacity of the nucleolus to sequester TR may be saturated and excess TR can assemble

with TERT and localize to telomeres. Importantly, our observations also suggests that

TCAB1 is not required for telomerase localization to telomeres.

Molecular mechanism of telomerase RNP assembly

The results presented in this study demonstrate that, in the absence of TCAB1,

telomerase assembly is reduced but not completely abolished. It is likely that TR bound

by the H/ACA complex is in an equilibrium between localizing to nucleoplasm and the

nucleolus. In the presence of TCAB1, this equilibrium is shifted towards the nucleoplasm

allowing TR to associate with TERT (Fig. 6E). In contrast, in the absence of TCAB1, the

majority of TR is trapped in the nucleolus effectively reducing the amount of TR that is

available for assembly with TERT. In addition, the nucleolus disassembles during mitosis

and does not reform until early G1-phase of the cell cycle (Hernandez-Verdun, 2014). In

cells lacking TCAB1, nucleolar disassembly should lead to the release of TR, which could

transiently allow telomerase assembly during cell division or prior to reformation of the

nucleolus in G1-phase. Both, a reduction of TR available for assembly and transient

assembly during cell division are plausible models to explain the limited amount of

telomerase that can form in the absence of TCAB1.

The sub-cellular location and order in which telomerase RNP components

associate with TR in human cells is largely unknown (Fig. 6E). Our results demonstrate

that telomerase does not assemble in the nucleolus. In addition, it is unlikely that

telomerase assembly requires intact Cajal bodies. Multiple studies have shown that

knock-out of coilin, which eliminates Cajal bodies, does not impact telomerase activity or

telomere maintenance (Chen et al., 2018; Vogan et al., 2016). Our single-molecule live

cell imaging of TERT has demonstrated that TERT is almost exclusively localized to the

nucleus, it is therefore likely that human telomerase assembles in the nucleoplasm. But

it cannot be ruled out that in human cells TR is transiently exported to the cytoplasm to

assemble with nascent TERT protein, as is the case in *S. cerevisiae* (Gallardo et al.,

2008).

The H/ACA complex has been shown to associate with scaRNAs at the site of their

transcription (Darzacq et al., 2006). While co-transcriptional association of the H/ACA

complex with TR has not been demonstrated, binding of the H/ACA complex proteins to

TR is likely the first step in telomerase RNP assembly, because of their critical role in TR

stability and 3'-end formation (Stuart et al., 2015; Tummala et al., 2015). Once TR is

bound to dyskerin, TR will be targeted to the nucleolus, unless it associates with TCAB1

(Fig. 6E). Like TERT, TCAB1 is excluded from nucleoli (Fig. 1, Fig. S2D), and dyskerin

an TR are localized to the dense fibrillar component (DFC, Fig. 6E) of the nucleolus (Yao

et al., 2019), which is separated from the nucleoplasm by the granular component (GC,

Fig. 6E) (Lafontaine et al., 2021). Thus, it is unlikely that TCAB1 can bind to TR localized

to the DFC of the nucleolus. We therefore believe that TCAB1 association with TR

prevents the entry of TR into the nucleolus rather than extracting TR which is already

localized to the DFC of the nucleolus. The association of TR with either TERT or TCAB1

should prevent its sequestration in the nucleolus, but the order in which TERT and TCAB1

bind to TR cannot be inferred from our experiments, or observation by others. Since

TCAB1 is substantially more abundant than TERT it is likely that TCAB1 encounters TR

before TERT does.

Regulation of RNP assembly by nucleolar phase-separation

In addition to the mechanistic insight into the role of TCAB1 in telomerase function, our

results also demonstrate that nucleolar phase separation can effectively regulate

telomerase RNP assembly in the nucleus of human cells. How RNA molecules are

specifically recruited into, excluded from, or expelled from non-membrane bound

organelles is a key unanswered question. One model suggests that gradual replacement

of non-specific, multivalent interactions of pre-ribosomal RNAs with nucleolar proteins

such as NPM1 and fibrillarin, with specific, high-affinity interactions with ribosomal

proteins leads to the ejection of mature ribosomal subunits from the nucleolus (Riback et

al., 2020). In this model a key driving force for the retention of RNA in the nucleolus is

regions of RNA not yet bound by ribosomal proteins, that are available to interact with

nucleolar proteins (Riback et al., 2020). By analogy, this model would explain why TR

bound by the H/ACA complex but not TERT would be sequestered in the nucleolus. In

addition to the interactions formed by the H/ACA complex with nucleolar proteins and

RNA, the regions of TR that are bound by TERT in the context of telomerase (i.e. the

pseudoknot, template, and CR4/CR5) would be available to form non-specific, multivalent

interactions with nucleolar proteins to strengthen the association of TR with the nucleolus

and prevent its release.

TR is a unique among the scaRNAs because it contains the additional domains

that associate with TERT. Most other box H/ACA scaRNAs are substantially shorter (<150

nucleotides) than TR (451 nucleotides), and do not contain large regions that are not

bound by proteins and could form non-specific interactions with nucleolar proteins. It is

therefore possible, that in cells lacking TCAB1, TR is strongly retained in the nucleolus

while other scaRNAs are less tightly bound, because they lack additional interaction sites

with nucleolar proteins. This provides a potential explanation for the phenotypes observed

in patients with TCAB1 mutations that suffer from dyskeratosis congenita. The patients

have a clear deficiency in telomerase function but no defects in splicing have been

reported (Zhong et al., 2011), which would be the consequence of complete loss of

scaRNA function and their critical role in spliceosome maturation.

How TCAB1 binding leads to the exclusion of TR and other scaRNA from the

nucleolus remains a key unanswered question. TCAB1 interacts with a very short

sequence motif in TR, which is far removed from the TR regions that associate with TERT

(Fig. S8) (Ghanim et al., 2021). It is therefore unlikely that TCAB1 binding leads to

expulsion of TR from the nucleolus by reducing the number of non-specific, multivalent

interactions TR can form with nucleolar proteins. As outlined above, we believe that

TCAB1 prevents localization of scaRNAs to the nucleolus, rather than extracting

scaRNAs that are already localized to the DFC. One potential explanation is that TCAB1

counteracts scaRNA recruitment to the nucleolus by inhibiting the nucleolar localization

signals within dyskerin (Heiss et al., 1999). Dissecting the molecular mechanism by which

TCAB1 leads to exclusion of TR from the nucleolus in future studies will undoubtably shed

light on the fundamental principles RNP recruitment to non-membrane bound organelles

and its physiological role in cell biology.

Acknowledgements

We would like to thank members of the Schmidt lab and J. Nandakumar for discussions

and critical reading of the manuscript and Luke Lavis (HHMI Janelia Research Campus)

for providing HaloTag dyes. This work was supported by a grant from the NIH (R00

GM120386) to J.C.S. J.C.S. is a Damon Runyon Dale F. Frey Scientist supported (in part)

by the Damon Runyon Cancer Research Foundation (DFS-24-17). S.B.C. acknowledges

sustained support from the Ernest & Piroska Major Foundation.

Author contributions

B.S.A.-M. carried out IF-FISH experiments, telomerase purifications, analyzed telomerase assembly, generated mCherry- and 3xFLAG-HaloTag-dyskerin plasmids, determined their sub-cellular localization, and edited the manuscript. G.I.P. maintained cell lines, established TCAB1 and TR knockout cell lines and carried out IF-FISH experiments. K.A.-B. assisted in establishing the TR knockout cell line and carried out characterization of the TR knockout cells. S.B.C. purified and characterized the anti-TERT sheep antibody. L.H. and K.Y. characterized TCAB1 knock-out clones using Southern blots. J.C.S. carried out all other experiments, designed the research, analyzed data, and wrote the manuscript.

Competing interests

The authors declare no competing interests.

Materials and Methods

Cell Lines and Tissue Culture

All cell lines were based on HeLa-EM2-11ht (Weidenfeld et al., 2009) and were cultured in Dulbecco's Modified Eagle Medium including L-glutamine (Gibco) supplemented with 10% fetal bovine serum, 100 units/ml penicillin and 100 µg/ml streptomycin at 37°C with 5% CO_2. Live cell imaging was carried out using CO_2 independent media supplemented with 2 mM GlutaMAX (Life Technologies), 10% fetal bovine serum, 100 units/ml penicillin and 100 µg/ml streptomycin at 37°C with 5% CO_2. For single-molecule imaging of HaloTag-TERT cell were cultured in homemade imaging dishes made by gluing 22x22 mm Nexterion coverslips (170 ± 5 µm, Schott) onto the bottom of plastic 3.5 x 1.0 cm cell culture dishes with a hole in the middle using an epoxy adhesive. Prior to chamber assembly the coverslips were washed with 1 M KOH and 100% for 30 min each in a sonicating water bath. To enrich for cells in S-phase for live cell imaging experiments, cultures we treated with complete media including 2 mM thymidine for a minimum of 16 hours. Cells were released 2 hours prior to imaging by replacing the thymidine containing media with fresh media without thymidine. Puromycin selection was carried out at a concentration of 1 µg/ml.

Plasmid Construction and Genome Editing

All plasmids were generated by Gibson assembly (NEB) using standard protocols or by inverse PCR. All plasmids will be made available on Addgene. All Cas9 and sgRNA expression plasmids were based on pX330 (Cong et al., 2013). The homologous recombination donor for the TR knockout was generated by assembling the genomic

sequences immediately upstream and downstream (~500 bp each) of the TR sequence

flanking a puromycin resistance cassette into HpaI linearized pFastBac. The 3xFLAG-

HaloTag-NLS plasmids was generated by adding a 3xFLAG-tag to a previously described

HaloTag-NLS plasmid (a kind gift from X. Darzacq and A. Hansen) (Hansen et al., 2018).

The 3xFLAG-HaloTag-dyskerin plasmid was generated by replacing TERT in our

previously described 3xFLAG-HaloTag-TERT expression plasmid with the dyskerin

coding sequence (Schmidt et al., 2016). The mCherry-dyskerin plasmid was generated

by replacing TERT in our previously described mCherry-TERT expression plasmid with

the dyskerin coding sequence (Schmidt et al., 2014). All transfections were carried out

using Lipfectamine 2000 (Invitrogen) using the manufacturer's instructions. TCAB1 was

knocked-out using a single sgRNA or two separate sgRNA and Cas9 encoding plasmids

that were transfected alongside a GFP-expression plasmid. 24 hours after transfection

single-cell clones were sorted using the GFP signal. TCAB1 knock-out clones were

screened by PCR and confirmed by western blot, Southern Blotting of the *TCAB1* locus

and immunofluorescence imaging. TR was knocked out by transfecting two sgRNA

plasmids and a homologous recombination donor plasmid. 48 hours after transfection

puromycin selection was initiated and 1 week after the initiation of selection single-cell

clones were generated by dilution into 96-well plates. TR knock-out was confirmed using

PCR and Sanger sequencing, fluorescence *in situ* hybridization, and RT-qPCR.

Immunofluorescence and Fluorescence *In Situ* Hybridization Imaging

Fixed cell immunofluorescence imaging and fluorescence *in situ* hybridization was carried

out as previously described (Schmidt et al., 2014). Briefly, cells grown on coverslips were

fixed in PBS supplemented with 4% formaldehyde. When using the HaloTag for

fluorescence detection cells were incubated with 100 nM of JF646 HaloTag-ligand for 30

min prior to fixation. Unincorporated ligand was removed by 3 washes with complete

media followed by placing the cells back in the incubator for 10 min to let additional dye

leak out of the cells. mEOS3.2-TRF2 was detected using the intrinsic fluorescence of

green form of mEOS3.2. After removing the fixation solution using 2 PBS washes,

coverslips were transferred into aluminum foil covered humidity chambers with a parafilm

layer and rinsed with 1 ml of PBS with 0.2% Triton X-100. Cells were than incubated in

blocking buffer (PBS, 0.2% Triton X-100, 3% BSA) for 30 minutes, followed by incubation

with primary antibodies diluted in blocking buffer for 1 hour. All primary antibodies (see

key resource table) were used at a concentration of 1 µg/ml. After three washes with PBS

+ 0.2% Triton X-100, coverslips were incubated with secondary antibodies diluted in PBS

+ 0.2% Triton X-100 for 1 hour. All secondary antibodies (see key resource table) were

used at a concentration of 4 µg/ml. Cells were washed three times PBS + 0.2% Triton X-

100 prior to a second fixation with PBS + 4% formaldehyde. In cases where nuclear

staining was used the first of the three washing steps also included 0.1 µg/ml HOECHST.

After the second fixation steps coverslips were dehydrates in three steps with ethanol

(70%, 95%, 100%), re-hydrated in 2xSSC + 50% formamide, blocked for 1 hour in

hybridization buffer (100 mg/ml dextran sulfate, 0.125 mg/ml *E. coli* tRNA, 1 mg/ml

nuclease free BSA, 0.5 mg/ml salmon sperm DNA, 1 mM vanadyl ribonucleoside

complexes, 50% formamide, 2xSSC) at 37°C, before incubating the coverslips in

hybridization buffer supplemented with three TR probes (30 ng per coverslip) over night

at 37°C. Probe sequences were previously described (Tomlinson et al., 2006). After

hybridization coverslips were washes twice for 30 minutes in 2xSSC + 50% formamide

and then mounted on slides using ProLong Antifade Diamond mounting media (Life

Technologies). Microscopy was carried out using a DeltaVision Elite microscope using a

60x PlanApo objective (1.42 NA) and a pco.edge sCMOS camera. We acquired 20 Z-

sections spaced by 0.2 µm, followed by image deconvolution and maximum intensity

projection of the sections using the DeltaVision Softworx software.

Single-Molecule Live Cell Imaging

Live cell single-molecule imaging was carried out on a Olympus IX83 inverted microscope

equipped with a 4-line cellTIRF illuminator (405 nm, 488 nm, 561 nm, 640 nm lasers), a

Olympus UAPO 100x TIRF objective (1.49 NA), a CAIRN TwinCam beamsplitter, 2 Andor

iXon 897 Ultra EMCCD cameras, and a blacked-out environmental control enclosure. The

microscope was operated using the Olympus cellSense software. 3xFLAG-HaloTag-

TERT was labeled for 2 min in complete media supplemented with 100 nM JF646-

HaloTag ligand (Grimm et al., 2015). After removing the HaloTag-ligand with three

washes in complete media, cells were placed back in the incubator for 10 min to allow

unincorporated dye to leak out of the cells. Cells were then transferred into CO_2

independent media and put on the microscope which was heated to 37°C. Single-

molecule imaging was carried out at 50 or 100 frames per second using highly inclined

laminated optical sheet illumination (Tokunaga et al., 2008). Movies were typically 20

seconds in length (2000 frames) and were followed by a transmitted light acquisition to

visualize overall cell morphology.

RT-qPCR

RNA samples for RT-qPCR analysis were generated by using RNA mini prep kits (Zymogen) using ~2 million cells as starting material. Reverse transcription was carried out using random hexamer primers and SuperScript III reverse transcriptase (Invitrogen) according to the manufacturer's instructions. qPCR was carried out using the Maxima SYBR Green qPCR master mix (Thermo Scientific) using primers for GAPDH and TR according to the manufacturer's instructions. All qPCR reactions were carried out in triplicates and three independent biological replicates were analyzed.

Southern Blotting

Southern blotting was carried out using standard protocols (Southern, 2006). Briefly, genomic DNA generated by phenol-chloroform extraction after cell lysis using TE supplemented with 0.5% SDS and 0.1 mg/ml Proteinase K, was digested with BamHI (generating a 1394 bp fragment spanning exons 1-3 of the *TCAB1* locus) and separated on a 0.8% agarose gel. The DNA was then transferred on a Hybond-N+ nylon membrane using capillary transfer. The *TCAB1* locus was detected using radioactive probes (alpha-^{32}P-dCTP) generated by randomly primed DNA synthesis using an 800 bp PCR product overlapping with the 1394 bp restriction fragment as a template and Klenow polymerase (NEB).

Western Blotting

Mini-PROTEAN TGX stain-free gels (Bio-Rad) were used for SDS-PAGE. Total protein was detected using a ChemiDoc MP (Bio-Rad) after a 45 second UV activation. Western

transfer was carried out using the Trans-Blot Turbo transfer system (Bio-Rad) according

to the manufacturer's instructions using the mixed molecular weight transfer setting.

Immuno-blotting was carried out using standard protocols. The TCAB1 antibody

(Proteintech) was used at a 1:2000 dilution, the TERT antibody (Abcam) was used at a

1:4000 dilution, and the dyskerin antibody (Santa Cruz Biotech) was used at a 1:200

dilution. Secondary antibodies were used at a 1:10000 dilution.

Northen Blotting

RNA was extracted from cell lysates and purified telomerase samples supplemented with

10 ng of a loading and recovery control (*in vitro* transcribed TR 34-328) using the RNeasy

Mini kit (Qiagen) and eluted in 30 ul of RNase free water. 15 ul of eluted RNA was mixed

with 15 ul of 2x formamide loading buffer (0.1XTBE, 25 mM EDTA, 0.1% bromophenol

blue, 0.1% xylene cyanol, 93% formamide) and heated to 60 °C for 5 min. Samples were

separated on a 6% TBE, 7M Urea, polyacrylamide gel (Life Technologies), and

transferred to a Hybond N+ membrane (Cytiva) using a wet-blotting apparatus in 1x TBE

for 2 hours at 0.5 A of constant current in the cold room. After transfer, membranes were

UV-crosslinked, and pre-hybridized in Church buffer for 2 hours at 50 °C. Three DNA

oligos complementary to TR were radioactively labeled using T4 PNK (NEB) and ~10×10^6

cpm of each probe were added to the membrane. Hybridization was carried out at 50 °C

overnight. Membranes were washed three times with 2xSSC, 0.1% SDS prior to exposure

to a storage phosphorescence screen (Cytiva) which was then imaged on an Amersham

Typhoon IP phosphoimager (Cytiva).

Telomerase Expression and Purification

Cell lines were transfected in 15 cm tissue culture plates at ~90% confluency (~25-30x10⁶

cells) using 7.5 µg of TERT plasmid, 30 µg of TR plasmid and 75 µl of Lipofectamine 2000

in 1875 ul of Opti-MEM (Cristofari and Lingner, 2006). Transfected cells were split to three

15 cm dishes 24 hours after transfection. 48 hours after transfection cells were counted,

harvested, and snap frozen in liquid nitrogen. Cells were lysed in 1 ml of CHAPS lysis

buffer supplemented with 5 µl of RiboLock RNase inhibitor (10 mM TRIS pH 7.5, 1 mM

$MgCl_2$, 1 mM EGTA pH 8.0, 0.5% CHAPS, 10% glycerol) per 100x10⁶ cells and rotated

at 4 °C for 30 min. Lysates were cleared in a table-top centrifuge at 21,000xg for 15 min

at 4 °C. Identical cell equivalents were used for all samples. 45 µg of anti-TERT antibody

was added per ml of cleared lysate and samples were rotated for 1 hour at 4 °C. Lysates

were then added to 100 µl of protein G agarose and rotated for 1 hour at 4 °C. The resin

was spun down at 1000xg and washed four times with 1 ml of Buffer W (20 mM HEPES

pH 7.9, 300 mM KCl, 2 mM $MgCl_2$, 1 mM EDTA, 1 mM DTT, 1 mM PMSF, 0.1% Triton X-

100, 10% glycerol). TERT was eluted in 100 µl of Buffer W supplemented with 5 µl of 1

mM TERT peptide by rotating for 30 min at room temperature.

Telomerase Activity Assays

Telomerase assays were carried out in 20 µl of reaction buffer (50 mM TRIS pH 8.0, 150

mM KCl, 1 mM MgCl2, 2 mM DTT, 100 nM TTAGGGTTAGGGTTAGG oligo, 10 µM dATP,

10 µM dGTP, 10 µM dTTP, 0.165 µM dGTP [α-32P] 3000 Ci/mmol) including 2 µl of

purified telomerase for 1 hour at 30 °C. Telomerase was incubated with the substrate

oligo for 15 min at room temperature, prior to initiating the reaction by addition of dNTPs.

Reactions were stopped by adding 100 μl of 3.6 M of ammonium acetate supplemented

with 20 μg of glycogen and 32P 5'-end labeled loading control oligos

(TTAGGGTTAGGGTTAGGG, TTAGGGTTAGGGTTAG). Reaction products were

precipitated using 500 μl of ice-cold ethanol and stored at -20 °C over-night. Reaction

products were spun down in a table-top centrifuge at max speed for 30 min at 4 °C,

washed with 500 μl of 70% ethanol, and spun down again speed for 30 min at 4 °C. The

70% ethanol was decanted, and the reaction products were dried in an Eppendorf

vacuum concentrator at 45 °C. Reaction products were resuspended in 20 μl of loading

buffer (0.05XTBE, 25 mM EDTA, 0.05% bromophenol blue, 0.05% xylene cyanol, 46.5%

formamide) and incubated at 95 °C for 5 min. 10 μl of each sample was separated on a

12% polyacrylamide, 7 M urea sequencing gel pre-run for 45 min at 90W. Gels were dried

and exposed to a storage phosphorescence screen (Cytiva) and imaged on an

Amersham Typhoon IP phosphoimager (Cytiva).

Single-Particle Tracking

Single-particle tracking was carried out in MATLAB 2019a using a batch parallel-

processing version of SLIMfast modified to allow the input of TIFF files (kindly provided

by Xavier Darzacq and Anders Hansen) (Hansen et al., 2018), an implementation of the

Multiple-Target-Tracing algorithm (Sergé et al., 2008), with the following settings:

Exposure Time = 10 ms, NA = 1.49, Pixel Size = 0.16 μm, Emission Wavelength = 664

nm, D_{max} = 5 μm^2/s, Number of gaps allowed = 2, Localization Error = 10^{-5}, Deflation

Loops = 0. Diffusion coefficients and the fraction of molecules in each distinct particle

population were determined using the MATLAB version of the Spot-On tool (kindly

provided by Xavier Darzacq and Anders Hansen) (Hansen et al., 2018) with the following

settings: TimeGap = 10 ms or 20 ms, dZ = 0.700 μm, GapsAllowed = 2, TimePoints = 8,

JumpsToConsider = 4, BinWidth = 0.01 μm, PDF-fitting, D_Free1_3State = [1 25],

D_Free2_3State = [0.1 1], D_Bound_3State = [0.0001 0.1]. For all experiments we carried

out 3 independent biological replicates with at least 15 cells for each cell line. The

complete data set can be found in supplemental figure S6D. The statistical significance

of differences in particle fractions and diffusion coefficients were assessed using a two-

tailed T-Test.

Quantification of Fixed Cell Imaging

For the quantification of cellular TR distribution in control and TCAB1 knockout cells we

assigned cells into one of three categories: Cells with TR only at telomeres, Cells with TR

only in nucleoli, Cells with TR at telomeres and in nucleoli. We carried out 3 independent

biological replicates and counted a minimum of 100 cells for control and TCAB1 knockout

cells.

Quantification of RT-qPCR data

RT-qPCR experiments were carried out in triplicate and the TR Ct value was normalized

to the GAPDG Ct value. The mean ΔCt (Ct of TR – Ct of GAPDH) value from three

independent experiments and the corresponding standard deviation is shown in figure

S3B.

Quantification of Western Blots, Northern Blots, and Telomerase Activity Assays

Gel images from Western Blots, Northern Blots, and Telomerase Activity Assays were

analyzed using ImageQuant TL 8.2. To quantify TR levels in Northern blots the TR band

intensity was normalized to the loading and recovery control signal added to the RNA

sample prior to RNA purification. To quantify telomerase activity the whole lane intensity

starting at repeat 1 was determined and divided by the sum of the loading control signals.

Telomerase processivity was calculated by dividing product intensity > 7 repeats by the

total signal in the respective lane. The statistical significance of the observed differences

was calculated using a two-tailed T-test using a minimum of three biological replicates.

Each biological replicate (independent telomerase expression and purification) was

analyzed in technical triplicate.

References

Angrisani A, Vicidomini R, Turano M, Furia M. 2014. Human dyskerin: beyond
    telomeres. *Biol Chem* **395**:593–610. doi:10.1515/hsz-2013-0287

Armanios M, Blackburn EH. 2012. The telomere syndromes. *Nat Rev Genet* **13**:693
    704. doi:10.1038/nrg3246

Chen L, Roake CM, Freund A, Batista PJ, Tian S, Yin YA, Gajera CR, Lin S, Lee B,
    Pech MF, Venteicher AS, Das R, Chang HY, Artandi SE. 2018. An Activity Switch in
    Human Telomerase Based on RNA Conformation and Shaped by TCAB1. *Cell* **174**.
    doi:10.1016/j.cell.2018.04.039

Cohen SB, Graham ME, Lovrecz GO, Bache N, Robinson PJ, Reddel RR. 2007. Protein
    Composition of Catalytically Active Human Telomerase from Immortal Cells. *Science*
    315:1850–1853. doi:10.1126/science.1138596

Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini
    LA, Zhang F. 2013. Multiplex Genome Engineering Using CRISPR/Cas Systems.
    Science **339**:819–823. doi:10.1126/science.1231143

Cristofari G, Lingner J. 2006. Telomere length homeostasis requires that telomerase
    levels are limiting. *Embo J* **25**:565 574. doi:10.1038/sj.emboj.7600952

Darzacq X, Kittur N, Roy S, Shav-Tal Y, Singer RH, Meier UT. 2006. Stepwise RNP
    assembly at the site of H/ACA RNA transcription in human cells. *J Cell Biology*
    173:207–218. doi:10.1083/jcb.200601105

Freund A, Zhong FL, Venteicher AS, Meng Z, Veenstra TD, Frydman J, Artandi SE.
    2014. Proteostatic Control of Telomerase Function through TRiC-Mediated Folding of
    TCAB1. *Cell* **159**:1389 1403. doi:10.1016/j.cell.2014.10.059

Gallardo F, Olivier C, Dandjinou AT, Wellinger RJ, Chartrand P. 2008. TLC1 RNA
    nucleo-cytoplasmic trafficking links telomerase biogenesis to its recruitment to
    telomeres. *Embo J* **27**:748 757. doi:10.1038/emboj.2008.21

Ghanim GE, Fountain AJ, Roon A-MM van, Rangan R, Das R, Collins K, Nguyen THD.
    2021. Structure of human telomerase holoenzyme with bound telomeric DNA. *Nature*
    1–5. doi:10.1038/s41586-021-03415-4

Grimm JB, English BP, Chen J, Slaughter JP, Zhang Z, Revyakin A, Patel R, Macklin
    JJ, Normanno D, Singer RH, Lionnet T, Lavis LD. 2015. A general method to improve
    fluorophores for live-cell and single-molecule microscopy. *Nat Methods* **12**:244 50–3
    p following 250. doi:10.1038/nmeth.3256

Hansen AS, Woringer M, Grimm JB, Lavis LD, Tjian R, Darzacq X. 2018. Robust
model-based analysis of single-particle tracking experiments with Spot-On. *Elife* **7**.
doi:10.7554/elife.33125

Heiss NS, Girod A, Salowsky R, Wiemann S, Pepperkok R, Poustka A. 1999. Dyskerin
Localizes to the Nucleolus and Its Mislocalization Is Unlikely to Play a Role in the
Pathogenesis of Dyskeratosis Congenita. *Hum Mol Genet* **8**:2515–2524.
doi:10.1093/hmg/8.13.2515

Hernandez-Verdun D. 2014. Assembly and disassembly of the nucleolus during the cell
cycle. *Nucleus* **2**:189–194. doi:10.4161/nucl.2.3.16246

Hyman AA, Weber CA, Jülicher F. 2014. Liquid-Liquid Phase Separation in Biology.
Annu Rev Cell Dev Bi **30**:39–58. doi:10.1146/annurev-cellbio-100913-013325

Jády BE, Bertrand E, Kiss T. 2004. Human telomerase RNA and box H/ACA scaRNAs
share a common Cajal body–specific localization signal. *J Cell Biology* **164**:647–652.
doi:10.1083/jcb.200310138

Lafontaine DLJ, Riback JA, Bascetin R, Brangwynne CP. 2021. The nucleolus as a
multiphase liquid condensate. *Nat Rev Mol Cell Bio* **22**:165–182.
doi:10.1038/s41580-020-0272-6

Mitrea DM, Kriwacki RW. 2016. Phase separation in biology; functional organization of a
higher order. *Cell Commun Signal Ccs* **14**:1. doi:10.1186/s12964-015-0125-7

Nagpal N, Wang J, Zeng J, Lo E, Moon DH, Luk K, Braun RO, Burroughs LM, Keel SB,
Reilly C, Lindsley RC, Wolfe SA, Tai AK, Cahan P, Bauer DE, Fong YW, Agarwal S.
2020. Small-Molecule PAPD5 Inhibitors Restore Telomerase Activity in Patient Stem
Cells. *Cell Stem Cell* **26**:896-909.e8. doi:10.1016/j.stem.2020.03.016

Nandakumar J, Bell CF, Weidenfeld I, Zaug AJ, Leinwand LA, Cech TR. 2012. The TEL
patch of telomere protein TPP1 mediates telomerase recruitment and processivity.
Nature **492**:285 289. doi:10.1038/nature11648

Nandakumar J, Cech TR. 2013. Finding the end: recruitment of telomerase to
telomeres. *Nat Rev Mol Cell Bio* **14**. doi:10.1038/nrm3505

Riback JA, Zhu L, Ferrolino MC, Tolbert M, Mitrea DM, Sanders DW, Wei M-T, Kriwacki
RW, Brangwynne CP. 2020. Composition-dependent thermodynamics of intracellular
phase separation. *Nature* **581**:209–214. doi:10.1038/s41586-020-2256-2

Schmidt JC, Cech TR. 2015. Human telomerase: biogenesis, trafficking, recruitment,
and activation. *Gene Dev* **29**:1095 1105. doi:10.1101/gad.263863.115

Schmidt JC, Dalby AB, Cech TR. 2014. Identification of human TERT elements
necessary for telomerase recruitment to telomeres. *Elife* **3**:e03563.
doi:10.7554/elife.03563

Schmidt JC, Zaug AJ, Cech TR. 2016. Live Cell Imaging Reveals the Dynamics of
Telomerase Recruitment to Telomeres. *Cell* **166**:1188 1197.e9.
doi:10.1016/j.cell.2016.07.033

Schmidt JC, Zaug AJ, Kufer R, Cech TR. 2018. Dynamics of human telomerase
recruitment depend on template-telomere base pairing. *Mol Biol Cell* **29**:869 880.
doi:10.1091/mbc.e17-11-0637

Sergé A, Bertaux N, Rigneault H, Marguet D. 2008. Dynamic multiple-target tracing to
probe spatiotemporal cartography of cell membranes. *Nat Methods* **5**:687 694.
doi:10.1038/nmeth.1233

Shukla S, Jeong H-C, Sturgeon CM, Parker R, Batista LFZ. 2020. Chemical inhibition of
PAPD5/7 rescues telomerase function and hematopoiesis in dyskeratosis congenita.
Blood Adv **4**:2717–2722. doi:10.1182/bloodadvances.2020001848

Shukla S, Schmidt JC, Goldfarb KC, Cech TR, Parker R. 2016. Inhibition of telomerase
RNA decay rescues telomerase deficiency caused by dyskerin or PARN defects. *Nat
Struct Mol Biol* **23**:286 292. doi:10.1038/nsmb.3184

Southern E. 2006. Southern blotting. *Nat Protoc* **1**:518–525. doi:10.1038/nprot.2006.73

Stern JL, Zyner KG, Pickett HA, Cohen SB, Bryan TM. 2012. Telomerase Recruitment
Requires both TCAB1 and Cajal Bodies Independently. *Mol Cell Biol* **32**:2384–2395.
doi:10.1128/mcb.00379-12

Stewart SA, Weinberg RA. 2006. Telomeres: Cancer to Human Aging. *Annu Rev Cell
Dev Bi* **22**:531–557. doi:10.1146/annurev.cellbio.22.010305.104518

Stuart BD, Choi J, Zaidi S, Xing C, Holohan B, Chen R, Choi M, Dharwadkar P, Torres
F, Girod CE, Weissler J, Fitzgerald J, Kershaw C, Klesney-Tait J, Mageto Y, Shay
JW, Ji W, Bilguvar K, Mane S, Lifton RP, Garcia CK. 2015. Exome sequencing links
mutations in PARN and RTEL1 with familial pulmonary fibrosis and telomere
shortening. *Nat Genet* **47**:512 517. doi:10.1038/ng.3278

Tokunaga M, Imamoto N, Sakata-Sogawa K. 2008. Highly inclined thin illumination
enables clear single-molecule imaging in cells. *Nat Methods* **5**:159 161.
doi:10.1038/nmeth1171

Tomlinson RL, Ziegler TD, Supakorndej T, Terns RM, Terns MP. 2006. Cell cycle-
regulated trafficking of human telomerase to telomeres. *Mol Biol Cell* **17**:955 965.
doi:10.1091/mbc.e05-09-0903

Tseng C-K, Wang H-F, Burns AM, Schroeder MR, Gaspari M, Baumann P. 2015.
Human Telomerase RNA Processing and Quality Control. *Cell Reports* **13**:2232–
2243. doi:10.1016/j.celrep.2015.10.075

Tummala H, Walne A, Collopy L, Cardoso S, Fuente J de la, Lawson S, Powell J,
Cooper N, Foster A, Mohammed S, Plagnol V, Vulliamy T, Dokal I. 2015. Poly(A)-
specific ribonuclease deficiency impacts telomere biology and causes dyskeratosis
congenita. *J Clin Invest* **125**:2151–2160. doi:10.1172/jci78963

Venteicher AS, Abreu EB, Meng Z, McCann KE, Terns RM, Veenstra TD, Terns MP,
Artandi SE. 2009. A human telomerase holoenzyme protein required for Cajal body
localization and telomere synthesis. *Science* **323**:644 648.
doi:10.1126/science.1165357

Vogan JM, Zhang X, Youmans DT, Regalado SG, Johnson JZ, Hockemeyer D, Collins
K. 2016. Minimized human telomerase maintains telomeres and resolves
endogenous roles of H/ACA proteins, TCAB1, and Cajal bodies. *Elife* **5**:693.
doi:10.7554/elife.18221

Weidenfeld I, Gossen M, Löw R, Kentner D, Berger S, Görlich D, Bartsch D, Bujard H,
Schönig K. 2009. Inducible expression of coding and inhibitory RNAs from
retargetable genomic loci. *Nucleic Acids Res* **37**:e50. doi:10.1093/nar/gkp108

Wu RA, Upton HE, Vogan JM, Collins K. 2017. Telomerase Mechanism of Telomere
Synthesis. *Annu Rev Biochem* **86**:439 460. doi:10.1146/annurev-biochem-061516-
045019

Xi L, Cech TR. 2014. Inventory of telomerase components in human cells reveals
multiple subpopulations of hTR and hTERT. *Nucleic Acids Res* **42**:8565–8577.
doi:10.1093/nar/gku560

Yao R-W, Xu G, Wang Ying, Shan L, Luan P-F, Wang Yang, Wu M, Yang L-Z, Xing Y-
H, Yang L, Chen L-L. 2019. Nascent Pre-rRNA Sorting via Phase Separation Drives
the Assembly of Dense Fibrillar Components in the Human Nucleolus. *Mol Cell*
76:767-783.e11. doi:10.1016/j.molcel.2019.08.014

Zhong F, Savage SA, Shkreli M, Giri N, Jessop L, Myers T, Chen R, Alter BP, Artandi
SE. 2011. Disruption of telomerase trafficking by TCAB1 mutation causes
dyskeratosis congenita. *Gene Dev* **25**:11–16. doi:10.1101/gad.2006411

Zhong FL, Batista LFZ, Freund A, Pech MF, Venteicher AS, Artandi SE. 2012. TPP1
OB-Fold Domain Controls Telomere Maintenance by Recruiting Telomerase to
Chromosome Ends. *Cell* **150**:481 494. doi:10.1016/j.cell.2012.07.012

 Supplementary Information

Figure 1 – Figure Supplement 1. (A) Strategy to knock-out TCAB1 with a single Cas9 and a single sgRNA
targeting exon 1 of the *TCAB1* gene. **(B)** PCR of TCAB1 knock-out clones using the primers indicated in
(A) demonstrating the absence of a WT *TCAB1* allele. **(C)** Sanger sequencing of the PCR product derived
from Halo-TERT TKO1 **(B)** demonstrating deletion causing a frame shift. **(D)** Model of TCAB1 protein
indicating the antigens used to raise the indicated TCAB1 antibodies. **(E-G)** Western blots of parental cells
and TCAB1 knock-out clones using **(E)** the Novus Biological, **(F)** Abcam, and **(G)** Proteintech TCAB1
antibodies, demonstrating the presence of a truncated TCAB1 protein in the putative TCAB1 knock-out
cells. **(H)** IF using antibodies against coilin and TCAB1 (Abcam) demonstrating that truncated TCAB1 can
be detected in Cajal bodies (white arrows) in putative TCAB1 knock-out clones. **(I)** IF using coilin and
TCAB1 (Novus Biologicals) antibodies coupled to FISH with probes against the scaRNA ACA57,
demonstrating that the Novus Biologicals antibody fails to detect the truncated TCAB1 and scaRNA
localization is unchanged in putative TCAB1 knock-out cells.

862
Figure 1 – Figure Supplement 2. (A) Strategy to knock-out TCAB1 using Cas9 and two sgRNAs
targeting introns 1 and 3. **(B)** Southern blot of genomic DNA digested with BamHI from parental cells and
TCAB1 knock-out clones using a probes generated from a PCR product of the TCAB1 gene indicated in
(A) demonstrating the expected truncation of the *TCAB1* gene in Halo-TERT TCAB1 KO 2. HeLa TCAB1

KO 2 carries larger deletions completely removing exons 1 and 2 from the *TCAB1* gene. **(C)** PCR using
primers indicated in **(A)** of genomic DNA from parental cells and TCAB1 knock-out clones confirming the
deletion of critical regions of the *TCAB1* gene show in **(B)**. **(D)** Immuno-fluorescence with anti-dyskerin
and anti-TCAB1 antibodies coupled to fluorescence in-situ hybridization with probes against TR,
demonstrating that expression of GFP-TCAB1 in TCAB1 knockout cells rescues TR localization to Cajal
bodies (scale bar = 5 μm).

Figure 2 – Figure Supplement 1. (A) Images of HeLa cells transiently expressing GFP-nucleolin to mark
nucleoli. The GFP-nucleolin signal overlaps with circular shapes visible under transmitted light illumination
(scale bar = 2 μm). **(B)** Images of HeLa cells transiently expressing GFP-nucleolin and 3xFLAG-HaloTag-
NLS labeled with JF646. The 3xFLAG-HaloTag-NLS signal (maximum intensity projection of 1000 frames
of a single-molecule imaging movie) clearly overlaps with the GFP-nucleolin signal (red dashed outline),
demonstrating that 3xFLAG-HaloTag-NLS can enter the nucleolus (scale bar = 2 μm). **(C)** Images of HeLa
cells transiently expressing 3xFLAG-HaloTag-dyskerin labeled with JF646 and probed with an antibody
against dyskerin, demonstrating that 3xFLAG-HaloTag-dyskerin localizes to the nucleolus (scale bar = 5
μm).

Figure 3 – Figure Supplement 1. (A) Images of HeLa cells transiently expressing mCherry-dyskerin
probed with antibodies agains mCherry and dyskerin, demonstrating that mCherry dyskerin localizes to the
nucleolus (scale bar = 5 µm). **(B)** Images of control and TCAB1 knock-out HeLa cells (3xFLAG-Halo-TERT,
mEOS3.2-TRF2) overexpressing untagged TERT and TR. TR was detected using FISH and TRF2 was
visualized using the fluorescence signal from mEOS3.2-TRF2. In control cells TR co-localizes with
telomeres, while it is enriched in nucleoli and localized to telomeres in TCAB1 KO cells (scale bar = 5 µm).
(C) Quantification of the fraction of cells showing TR localization exclusively to telomeres, to telomeres and
nucleoli, or only to nucleoli (2 independent experiments, >100 cells per experiments, mean ± standard
deviation). **(D)** Images of HeLa cells and TCAB1 knockout cells transiently expressing untagged TERT and
TR, probed with TERT (Abcam) and dyskerin antibodies and FISH for TR demonstrating that untagged
TERT is excluded from nucleoli marked by dyskerin in parental and TCAB1 knock-out cells.

Figure 5 – Figure Supplement 1. Direct telomerase extension assay of telomerase immuno-purified from parental (WT) and TCAB1 knock-out (TKO) HeLa and Halo-TERT cell lines. LC1 and LC2, radiolabeled DNA oligonucleotide loading controls.

897
898
899
900

Figure 6 – Figure Supplement 1. (A) PCR analysis of the TR locus in parental and TR knock-out clones.
Both TR knock-out clones show PCR products with reduced length that were confirmed to be knock-outs
by Sanger sequencing. **(B)** Images of control and TR knock-out cells probed with an antibody against
coilin and FISH probes specific for TR, demonstrating the lack of TR signal in TR knock-out cells (scale
bar = 5 µm). **(C)** Determination of TR levels in control, TCAB1 knock-out, and TR knock-out cells, using
RT-qPCR with primers specific to TR normalized to GAPDH (3 independent biological replicates, 3
technical replicates for each biological replicate, mean ± standard deviation). **(D)** Maximum intensity
projection of 2000 frames of a 3xFLAG-HaloTag (JF646) TERT movie (bottom), demonstrating that the
TERT signal does not overlap with the nucleolus detected as circular shape in the transmitted light image
(top, red dashed line). **(E)** Fitting of single-particle tracking data of TERT from control, TCAB1 knock-out,
and TR knock-out cells expressing 3xFLAG-HaloTag-TERT using the Spot-On tool.
915
916

Figure 6 – Figure Supplement 2. (A) Analysis if the step size of telomeric TERT particles relative to the
distance of the particle to the closest telomere (pooled results from 3 independent biological replicates with
19-30 cells analyzed per replicate). TERT molecules bound to the telomere are expected to have small
step sizes and a short distance to the closest telomere, which is apparent in the enrichment of events in
the lower quadrants in the WT control. This enrichment is not observed in TCAB1 and TR knock-out cells.
(B) Spot-On analysis of telomeric TERT particles (pooled results from 3 independent biological replicates
with 19-30 cells analyzed per replicate). The fraction of bound TERT particles in TCAB1 and TR knock-out
cells is 4-5%, compared to 12% in the WT control cells.
926

927
928 **Figure 6 – Figure Supplement 3. (A)** Structure of the telomerase RNP showing the distance between
929 TCAB1 and CR4/CR5 (13.4 nm). TR in blue, CR4/CR5 in red, and TCAB1 in green (Ghanim et al., 2021).
930
931

Movie Legends

Movie 1. Single-particle tracking of 3xFLAG-HaloTag-TERT labeled with JF646 in a control cell acquired at 100 frames per second. Trajectories with a minimum of 5 localizations are displayed. 150x150 pixels with a pixel size of 0.16 µm.

Movie 2. Single-particle tracking of 3xFLAG-HaloTag-TERT labeled with JF646 in a TCAB1 knockout cell acquired at 100 frames per second. Trajectories with a minimum of 5 localizations are displayed. 150x150 pixels with a pixel size of 0.16 µm.

Movie 3. Movie of cell expressing GFP-nucleolin (red) and 3xFLAG-HaloTag-NLS (green) labeled with JF646 acquired at 100 frames per second, showing overlap of 3xFLAG-HaloTag-NLS with nucleoli. 140x140 pixels with a pixel size of 0.16 µm.

Movie 4. Movie of 3xFLAG-HaloTag-TERT labeled with JF646 in a control (left), TCAB1 knockout (middle), and TR knockout (right) cell acquired at 100 frames per second. Each panel is 150x150 pixels in size with a pixel size of 0.16 µm.

Movie 5. Single-particle tracking of 3xFLAG-HaloTag-TERT labeled with JF646 in a TR knockout cells acquired at 100 frames per second. Trajectories with a minimum of 5 localizations are displayed. 150x150 pixels with a pixel size of 0.16 µm.