1   Title page
2

EEG and behavioral correlates of attentional processing while walking and navigating naturalistic environments

Authors: Magnus Liebherr[1,2*], Andrew W. Corcoran[3,4*], Phillip M. Alday[3], Scott Coussens[3], Valeria Bellan[3,5], Caitlin A. Howlett[3,5], Maarten A. Immink[3,6], Mark Kohler[3,7], Matthias Schlesewsky[3], Ina Bornkessel-Schlesewsky[3]

Affiliation: [1]Department of Neuroscience, Karolinska Institutet, Stockholm, Sweden; [2]Department of General Psychology: Cognition, University Duisburg-Essen, Germany; [3]Cognitive and Systems Neuroscience Research Hub, University of South Australia, Adelaide, Australia; [4]Cognition and Philosophy Laboratory, Monash University, Melbourne, Australia; [5]Innovation, Implementation & Clinical Translation (IIMPACT) in Health, University of South Australia, Adelaide, Australia; [6]Sport, Health, Activity, Performance and Exercise Research Centre, Flinders University, Adelaide, Australia; [7]School of Psychology, University of Adelaide, Adelaide, Australia

* Both authors contributed equally to the publication
Corresponding authors: Magnus Liebherr; Andrew W. Corcoran

Emails: magnus.liebherr@gmail.com; andrew.corcoran1@monash.edu

ORCid:
ML: 0000-0001-8580-2464
AWC: 0000-0002-0449-4883
PMA: 0000-0002-9984-5745
SC: 0000-0002-3564-5859
VB: 0000-0002-8512-5288
CAH: 0000-0002-4584-8641
MAI: 0000-0002-4652-0090
MK: 0000-0001-7265-6242
MS: 0000-0003-4131-8077
IBS: 0000-0002-3238-6492

Abstract

The capacity to regulate one's attention in accordance with fluctuating task demands and environmental contexts is an essential feature of adaptive behavior. Although the electrophysiological correlates of attentional processing have been extensively studied in the laboratory, relatively little is known about the way they unfold under more variable, ecologically-valid conditions. Accordingly, this study employed a 'real-world' EEG design to investigate how attentional processing varies under increasing cognitive, motor, and environmental demands. Forty-four participants were exposed to an auditory oddball task while (1) sitting in a quiet room inside the lab, (2) walking around a sports field, and (3) wayfinding across a university campus. In each condition, participants were instructed to either count or ignore oddball stimuli. While behavioral performance was similar across the lab and field conditions, oddball count accuracy was significantly reduced in the campus condition. Moreover, event-related potential components (mismatch negativity and P3) elicited in both 'real-world' settings differed significantly from those obtained under laboratory conditions. These findings demonstrate the impact of environmental factors on attentional processing during simultaneously-performed motor and cognitive tasks, highlighting the value of incorporating dynamic and unpredictable contexts within naturalistic designs.

Keywords: Mobile EEG, ERP, attention, real-world, cognitive load

Introduction

Attention is a finite resource that must be strategically managed throughout daily life. The hyper-connected nature of the modern world, and in particular, the ubiquitous devices, apps, and media that constantly vie for our attention, expose us to an abundance of information that must be selectively sampled, navigated, and at times, held in abeyance. Arguably, it has never been more important to understand the impact of competing attentional demands on cognitive function and motor action. The present study builds on recent developments in mobile electroencephalography (EEG) by investigating the psychophysiological effects of cognitive task load, motor demands, and environmental complexity on attention in real-world scenarios.

A great deal of knowledge about the nature of attention and the influence of distracting stimuli has been derived from carefully controlled laboratory studies [1]. By contrast, much less is known about the characteristics of attention under real-world situations. However, recent technological developments in neurophysiological measures (e.g., EEG and eye-tracking) have enabled researchers to address such topics outside the laboratory (e.g., while driving) [2]. The present study exploits this technical progress to examine how event-related potentials (ERPs) relating to sensory and attentional processing differ across laboratory and more ecologically-valid situations. Here, we focus on the P3 component, which has been most frequently investigated in previous real-world studies [3], and the mismatch negativity (MMN), which constitutes an index of early sensory processing [4].

The P3 constitutes the third positive deflection after stimulus presentation, typically occurring at a latency ranging 250 to 500 ms and most dominantly in centro-parietal areas [5,6]. This component has been related to updating and categorization processes [7], the intensity of performance [8,9], as well as the amount of resources dedicated to task performance [10,11]. Furthermore, the P3 can serve as an index of the distribution of attention in competing concurrent tasks [12] and has been shown to be sensitive to alterations in task difficulty [13].

The MMN is elicited in the presence of any discriminable change in some repetitive aspect of stimulation. Peaking within 100 to 250 ms from change onset, the MMN exhibits the strongest intensity in temporal and frontal areas of topographic scalp maps, and can be detected even when the participant pays no attention to the task [14–16]. Although the MMN has been most intensively studied in the auditory modality [17], previous studies have also identified the component in visual settings [18].

Traditionally, the P3 and MMN components are most frequently studied using the oddball
paradigm. Within this paradigm, participants are typically presented with an infrequently
occurring target stimulus (e.g., a high-pitch tone) embedded within sequences of frequently
presented distractor stimuli (e.g., low-pitch tones). While both the P3 and the MMN are
elicited by such target stimuli, the MMN can also be detected when the individual is not
actively attending to the tone sequence [19].

From pioneers to contemporary studies, the oddball paradigm has also been frequently
used to investigate the influence of motor demands (e.g., varying gait parameters) on ERPs,
where participants are asked to perform the task while walking on a treadmill with varying
velocity (slow walking vs. fast walking) [20]. Here, findings suggest that the amplitude, latency
and topography of event-related potentials (N1, N2, P3) vary depending on the difficulty of
both the motor and the cognitive task [20,21]. When occurring in parallel with a cognitive task,
motor demands (e.g., during walking) can thus be viewed as inducing a dual-task setting that
calls for appropriate allocation of attention between cognitive and motor systems [22].

In recent years, studies have begun to measure ERPs during dual-task scenarios performed
under more naturalistic conditions, such as cycling [23,24], skateboarding [25], or walking [26]
outside the lab (e.g., on empty corridors, parking places, or the lawn on the outside premises).
Such findings demonstrate a relation between attentional processing and the cognitive load
induced by each situation. For example, cognitive task performance, parietal P3 amplitude,
and frontal midline theta power have been found to decrease with progressively increased
situational load (i.e., standing vs. walking on a treadmill vs. walking along a hallway) [27].
Furthermore, these findings show a greater impact of environmental factors on
electrophysiological markers of attention compared to motor demands (i.e., being wheeled
along a hallway vs. walking on a treadmill). To date, however, most real-world studies on
walking involve fairly undemanding environmental settings, such as an empty straight
corridor [27] or the lawn of an institute's grounds [3].

Within a more naturalistic context, Aspinall and colleagues described EEG changes in
terms of increased engagement and arousal in more busy environmental settings (i.e., urban
shopping street and busy commercial district) compared to green spaces [28]. However, the
interpretation of these findings is complicated by the authors' reliance on proprietary EEG
metrics that are not commonly reported in the literature. More recently, Pizzamiglio and

colleagues reported increased theta and beta power when participants were walking across a

campus while conversing, as compared to standing [29]. However, studies that have analyzed

attention-related ERP responses under such ecologically valid conditions are rare and often

limited to virtual reality [22].

In sum, previous studies on walking in naturalistic environments have either focused on the

effect of motor demands [3], used relatively monotonous scenarios (e.g., empty corridors) [27], or

did not analyze attention-related ERPs [29–31]. Here, we sought to fill an important gap in this

literature by investigating the effects of motor demands, environmental complexity, and

cognitive task load on such ERPs. We report findings from an auditory oddball paradigm in

which participants either attended to or ignored experimental stimuli while engaging in three

situational contexts: (1) ordinary laboratory conditions, (2) walking around a sports field, and

(3) navigating a route through a university campus. Behavioral data and ERPs pertaining to

the MMN and P3 were analyzed to investigate the effect of each experimental manipulation

on task performance and attentional processing.

Based on previous findings from traditional laboratory studies and studies in increasingly

naturalistic environments, we hypothesized a decrease in P3 and MMN amplitude as well as

task-performance with increased environmental and motor complexity, due to the reduction in

attentional resources available for cognitive task performance. In addition, we predicted that

P3 and MMN effects would be more pronounced under conditions in which specific emphasis

is placed on the cognitive task, compared to when the tone stimuli are ignored.

Methods

Participants

Healthy, right-handed, native English-speaking adults aged 18-40 years were eligible to

participate in this study. Of the 44 recruited participants, one was excluded due to left-

handedness, and a further two were excluded due to missing EEG data. The remaining sample

consisted of 26 females and 15 males aged 18–39 years (M = 23.1 years, SD = 4.8). All

participants reported normal (or corrected-to-normal) vision and audition, no history of

neurological or psychological disorders, and no diagnosed intellectual or language disorder.

The study was performed in accordance with the ethical standards laid down in the

Declaration of Helsinki and approved by the local ethics committee at the University of South

Australia (ID: 0000034991). All participants provided written informed consent prior to the
experiment and were informed that they could end participation at any time without reprisal.

Materials

The auditory oddball task comprised two distinct tones, one presented frequently
(standard) and the other infrequently (deviant). These stimuli were a French horn tone
(Dominant frequency of 1046.502Hz, High C or C6) and a piano tone (Dominant frequency
of 523.251Hz, Middle C or C5). We used complex tones because they constitute more
naturalistic stimuli than pure tones [32].

Tones were presented binaurally via conventional in-ear headphones (Anko) for 150 ms,
with an interstimulus interval of 350 ms. One-hundred and fifty stimuli were presented per
block, of which 13-30% were deviants ($M = 20\%$). Tones were randomly assigned as the
standard/deviant. Stimulus presentation was driven by OpenSesame v3.0.7 [33] running on a
commercial compact laptop in conjunction with BrainVision StimTrak (Gain = 10, Trigger
Level = 1V). The volume of the oddball stimuli was kept the same for all participants and
conditions (approximately 60 dB).

Procedure

After providing written, informed consent, participants completed a demographic
questionnaire and the Edinburgh Handedness Inventory [34], followed by the EEG setup. Before
starting the task, the two tones were presented to familiarize participants with the stimuli.
Participants performed 5 blocks of the oddball task in each of the three environments (Lab,
Field, Campus) while counting the number of deviant stimuli (Count), and again in each of
the three environments while ignoring task stimuli (Ignore). The order of the task
manipulation was randomized, as was the order of environmental conditions nested within
either task condition. We chose counting rather than button press responses as our measure of
attentional engagement because previous studies have reported that P3 topography is
confounded by motor-evoked potentials [35].

Within the Lab condition, participants performed the oddball task while sitting alone in an
interview room. The room was sparsely furnished with a large table, six chairs and a
sideboard, surrounded by three white walls without pictures, and one frosted glass wall.
Participants sat on a chair with armrests, with the laptop in front of them (supported by a

harness strapped to their torso). The participant's gaze was directed towards a white wall. The glass wall, which separated the interview room from a corridor, was on their right.

The Field condition involved walking around a grass sports field at the Magill campus of the University of South Australia, Adelaide. The sports field is surrounded by a grandstand, a car park for members of the university, trees and fencing (see Figure 1a). The Campus condition involved navigating a pre-defined route through the university campus adjacent to the sports field. Participants were provided a map with a cross-marked destination (red point) to which they should navigate via several waypoints (yellow points) (see Figure 1b). The route included both outdoor and indoor sections. Participants had to use stairs, open doors, and interact with other people on the campus.

Figure 1 a) Grass sports field at the Magill campus at the University of South Australia (from *Google Maps*, by Google, https://bit.ly/2R0XXjl), b) Reproduction of the map provided to the participants in the "campus condition", showing the route across the Magill campus. Participants began at the blue dot that represents the starting point and had to follow the yellow dots until they reached the red dot that represents the finish point.

During both walking conditions, participants carried the laptop in front of them (supported by the harness) while carrying the StimTrak in a plastic pocket (attached with a lanyard) over one shoulder as they made their way along the route. Participants were permitted to walk at a comfortable, self-determined pace. The entire testing session lasted between 2.5 and 3 hours, with sufficient breaks between conditions to avoid fatigue. Data collection occurred during the summer months, with temperatures typically ranging between 25 and 30ºC. Sessions were not performed when it was raining or in extreme temperatures.

EEG acquisition and preprocessing

The EEG was continuously recorded from 32 Ag/AgCl actiCAP mounted active electrodes using a Brain Vision LiveAmp 32 amplifier (Brain Products GmbH, Gilching, Germany).

Electrode positions conformed to the extended 10–20 system: Fp1, Fz, F3, F7, FT9, FC5, FC1, C3, T7, TP9, CP5, CP1, Pz, P3, P7, O1, Oz, O2, P4, P8, TP10, CP6, CP2, Cz, C4, T8, FT10, FC6, FC2, F4, F8, Fp2. All channels were digitized at a sampling rate of 500 Hz. AFz served as the ground and FCz as the online reference. Electrode impedances were below 20 kΩ at the beginning of the experiment. EEG data were stored on a micro SD card inside the amplifier; the amplifier was kept inside a mesh pocket on the back of the EEG cap during the experiment. After the experiment, the data were transferred from the SD card to a hard drive for offline processing and analysis.

EEG data were preprocessed in MATLAB R2019b (v9.7.0.1319299; The MathWorks, Natick, MA) using custom-built scripts incorporating functions from the EEGLAB (v2019.1;[36]) and ERPLAB (v8.10;[37]) toolboxes. Marker files were first manually edited to remove artefactual triggers caused by feedback from the auditory equipment. Edited data were then imported into EEGLAB and downsampled to 250 Hz. Accelerometer channels and non-experimental recording segments were excluded. The online reference was then reintroduced into the dataset, and the data re-referenced to the average of TP9 and TP10 in order to approximate linked mastoids. The data were then high-pass filtered (passband edge = 0.5 Hz, transition width = 0.5 Hz) using a zero-phase, hamming-windowed, finite impulse response (FIR) sinc filter (implemented in the *pop_eegfiltnew* function from the *firfilt* plugin; v2.4).

Filtered data were processed with artifact subspace reconstruction (implemented with the *pop_clean_artifacts* function from the *clean_rawdata* plugin; v2.3). This routine was used to remove flat or noisy channels (flatline criterion = 5 s; channel correlation criterion = 0.80; line noise criterion = 4), and to correct for large amplitude artifacts (burst criterion = 10, [38,39]). Data were then low-pass FIR filtered (passband edge = 40 Hz, transition width = 10 Hz) prior to independent component analysis (*binica* implementation of the extended logistic infomax ICA algorithm,[40,41]). Resultant decompositions were automatically classified using the ICLabel plugin (v1.3, [42]). Components classed as either ocular or cardiac in origin with >90% probability were subtracted (median = 2, range = 1-3).

Missing channels (median = 1, range = 0-4) were interpolated for the purposes of ERP visualization but excluded from statistical analysis. All EEG event markers were systematically shifted -200 ms relative to their recorded latencies (*pop_erplabShiftEventCodes*) to account for the lag in signal transmission introduced by the

StimTrak. The EEG was then epoched [-200, 800 ms] relative to the corrected event markers.

Extracted epochs were mean-centered and subjected to a moving window artefact rejection

routine (*pop_artmwppth*; voltage threshold = 125 µV, time period = [-200, 600 ms]; window

size = 100 ms, window step = 50 ms). Grand average ERP waveforms and difference waves,

along with their accompanying standard error estimates, were computed using the

pop_gaverager function (weighted averages based on number of available trials per

participant). Finally, mean voltages were extracted from three time-windows per trial epoch

(Prestimulus = [-200, 0 ms]; MMN = [100, 250 ms]; P3 = [250, 500 ms]) and exported for

statistical analysis.

Data analysis

All statistical analysis was conducted in *R* (v3.6.2 [43]) using the *RStudio Desktop IDE*

(v1.2.5033 [44]). Behavioral and EEG data were analyzed using linear mixed-effects models fit

using the *lmer()* function from *lme4* (v1.1-23 [45]). All mixed-effects models included a by-

participant random intercept term to account for non-independence of (within-subject)

observations. All categorical independent variables were entered as unordered factors using

sum-to-zero contrast coding (reference category coded -1).

For the behavioral model, deviant counts for each block of the Count task were included as

the dependent variable, while number of deviant stimuli and environmental condition were

included as independent variables. Note that including the number of presented deviants as a

covariate controlled for differences in the ground truth count across blocks, enabling us to

implicitly model count accuracy as a function of environmental condition.

Separate EEG models were fitted for the MMN and P3 time-windows, with single-trial

mean voltage estimates serving as the dependent variable. Environmental condition, stimulus

type, and task manipulation were entered as independent variables and permitted to interact

with one another. Models additionally included mean prestimulus voltage as a scaled

covariate in order to address trial-by-trial variation in prestimulus activity (essentially

functioning as a form of baseline correction; see [46]), and by-channel random intercepts to

account for topographic differences in voltage estimates (see supplementary materials for

models that specify midline channel location as a fixed effect). By-participant random slopes

of environment were included in the MMN model; by-participant random slopes of

environment and stimulus were included in the P3 model (more complex random effects

structures failed to converge).

Main effects and interaction terms were assessed using Wald tests from Type-II ANOVA

tables obtained from the *car* package (v3.0-10 [47]) Model summary tables were produced using

sjPlot (v2.8.7 [48]). P-values for model terms were derived using Satterthwaite's method for

approximating degrees of freedom. Post-hoc contrasts were obtained via the *emmeans*

package (v1.5.1 [49]). Pairwise contrasts were corrected using the Tukey method; interaction

contrasts were corrected using the Sidak method. Plotted model predictions are estimated

marginal means (also obtained from the *emmeans* package) and visualised with the aid of

tidyverse (v1.3.0 [50]). Model predictions are plotted with 84% confidence intervals, where non-

overlapping intervals indicate a significant difference at $\alpha = .05$ [51].

Data Availability Statement

Raw and preprocessed datasets, along with all code necessary to reproduce the analysis

pipeline reported in this paper, are openly available from the OpenNeuro database [Dataset:

doi: 10.18112/openneuro.ds003620.v1.0.2]. Data were archived in accordance with the Brain

Imaging Data Structure (BIDS) standard for EEG [52] with the assistance of functions from the

bids-matlab-tools plugin (v5.2; https://github.com/sccn/bids-matlab-tools).

Results

Behavioral data

Two participants were excluded from analysis due to numerous highly inaccurate count

estimates (more than double the number of presented deviants on the majority of blocks). Due

to a programming error, count estimates from the final block of trials were missing for 21

participants. A further 16 count estimates were excluded as outliers (> 3 *SD* from the sample

mean after estimates had been normalized by the number of presented deviants). The included

39 participants contributed a median 14 counts to the model.

Numerically, mean count estimates were similar across Lab and Field conditions, but

reduced in the Campus condition. Linear mixed-effects modelling revealed a significant effect

of Environment ($\chi^2(2)=12.08$, *p*=.002) after controlling for the number of presented deviants.

Post-hoc contrasts revealed that count estimates were significantly reduced during the
Campus condition compared to both the Lab ($p_{adj}=.005$) and Field ($p_{adj}=.014$) conditions. No
significant difference was found between Lab and Field conditions ($p_{adj}=.930$).

Figure 2. Estimated marginal mean counts of deviant stimuli per block across environmental
conditions (Lab, Field, Campus). Error bars indicate 84% confidence intervals. Dashed line
indicates perfect performance.

Event related potentials

A further three participants were excluded from the ERP analysis due to excessively noisy
recordings (>50% epochs rejected due to artifact). Epoch rejection rates for the remaining 36
participants ranged from 0 to 38.6% ($M=7.2$, $SD=9.9$).

Grand average ERPs and difference waveforms from electrodes Fz and Cz are displayed in
Figure 3 (see supplementary materials for additional ERP plots). In the Lab condition, both
channels revealed a marked negative-going waveform in the early time-window for deviant
stimuli, consistent with a MMN component. This effect was evident irrespective of task
manipulation, although scalp topographies displaying the mean voltage difference between
stimuli indicate that the effect was more pronounced over centro-parietal sensors during the

Count task (see Figure 3). Conversely, deviants elicited a strong positive deflection in the first

half of the later time-window, consistent with a P3 response. This effect was also strongest

over centro-parietal channels in the Count task.

Field and Campus conditions rendered qualitatively similar ERP waveforms. Deflections

in the MMN time-window were positive-going for standard tones, and negative-going for

deviants, rendering a negative difference wave that was most pronounced over central

electrode sites. This pattern of activity was again reversed in the P3 time-window, resulting in

a centrally distributed positivity that peaked approximately 300 ms post stimulus onset. This

effect was less pronounced in the Campus compared to the Field condition. Comparing across

task manipulations, the difference between deviant and standard waveforms in the MMN

time-window was slightly accentuated, and more frontally distributed, in the Ignore task,

while the P3 window revealed slight accentuation over centro-parietal regions in the Count

task.

Taken together, the qualitative pattern of both the early and late ERP components elicited

during the walking conditions was generally similar to those elicited in the Lab condition, but

of reduced magnitude.

Figure 3. Grand average event-related potentials (ERPs; solid lines), difference waveforms (broken lines) and topographies for Ignore (top panel) and Count (bottom panel) tasks across Lab (left column), Field (middle column), and Campus (right column) conditions. Light blue waveforms depict ERPs evoked by deviant stimuli; purple waveforms depict ERPs evoked by standard stimuli. Broken lines indicate the difference in mean voltage (μV) between deviants and standards. Topographies depict mean voltage differences (μV) averaged across the two time-windows of interest (MMN = 100 – 250 ms; P3 = 250 – 500 ms). Yellow dots indicate electrode locations for plotted ERPs (upper = Fz, lower = Cz).

MMN time-window

Type II Wald tests revealed a significant three-way interaction between stimulus, task, and environmental condition for the MMN model (χ^2 (2) = 7.67, p = .022; see Table 1 for full decomposition). This interaction is resolved in Figure 4 (see Table 2 for the full model summary). This plot reveals that mean voltage was significantly more negative for all pairwise contrasts of stimulus type within each factorial combination of Task and Environment (all p_{adj} <.001), consistent with the elicitation of the MMN. In line with the ERP waveforms depicted in Figure 3, the magnitude of this difference was greatest in the Lab condition, and reduced in the other two conditions. This diminution was predominantly driven by the modulation of cortical responses to standard stimuli, which were less positive-going in the Field and Campus conditions compared to the corresponding Lab condition, irrespective of task instructions (all p_{adj} <.001).

The difference between stimulus responses across environmental conditions was qualitatively similar irrespective of whether participants counted or ignored task stimuli. Nonetheless, sequential interaction contrasts revealed that counting accentuated the difference between standard and deviant responses in the Lab condition relative to the Field condition (p_{adj} =.028). A similar effect was noted for the Campus relative to the Field condition, on account of the increased difference between stimulus responses when participants counted deviants (p_{adj} =.038).

400　Table 1. Analysis of deviance tables for linear mixed-effects models evaluating voltage
401　changes in the MMN (100-250 ms) and P3 (250-500 ms) time-windows.

	MMN time-window			P3 time-window		
	χ^2	d.f.	p-value	χ^2	d.f.	p-value
stimulus	5176.37	1	**<.001**	36.39	1	**<.001**
task	0.87	1	.350	76.78	1	**<.001**
environ	27.66	2	**<.001**	8.73	2	**.013**
prestim.s	246102.39	1	**<.001**	575055.15	1	**<.001**
stimulus:task	0.07	1	.786	221.33	1	**<.001**
stimulus:environ	799.65	2	**<.001**	1775.21	2	**<.001**
task:environ	105.34	2	**<.001**	492.80	2	**<.001**
stimulus:task: environ	7.67	2	**.022**	33.40	2	**<.001**

402　*Note*. χ^2 = Type II Wald χ^2 test statistic, *d.f.* = degrees of freedom, *stimulus* = oddball task
403　stimulus (Deviant, Standard), *task* = cognitive task condition (Count, Ignore), *environ* =
404　environmental context (Lab, Field, Campus), *prestim.s* = prestimulus EEG voltage (scaled).
405

406
Figure 4. Estimated marginal mean voltage amplitudes for MMN (left column) and P3 (right
column) time-windows during the Ignore (top row) and Count (bottom row) conditions of the
oddball task. Responses to standard tone stimuli are depicted by purple circles; responses to
deviant tone stimuli are depicted by blue diamonds. Estimates for each environmental
condition (Lab, Field, Campus) are displayed along the x-axis of each subplot. Error bars
indicate 84% confidence intervals.

Table 2. Summary of linear mixed-effects models for the MMN and P3 time-windows.

Predictors	MMN time-window				P3 time-window			
	Estimate	S.E.	t-stat	p-value	Estimate	S.E.	t-stat	p-value
(Intercept)	0.21	0.07	3.00	**.004**	-0.16	0.07	-2.24	**.028**
stimulus[S.Standard]	0.22	0.00	71.38	**<.001**	-0.18	0.03	-5.94	**<.001**
task[S.Ignore]	0.00	0.00	0.82	.411	-0.04	0.00	-15.69	**<.001**
environ[S.Lab]	0.17	0.05	3.69	**.001**	-0.06	0.05	-1.22	.229
environ[S.Field]	-0.07	0.03	-2.53	**.016**	0.03	0.03	1.17	.250
prestim.s	-1.22	0.00	-496.09	**<.001**	-1.41	0.00	-758.32	**<.001**
stimulus[S.Standard] : task[S.Ignore]	-0.00	0.00	-0.24	.814	0.03	0.00	14.80	**<.001**
stimulus[S.Standard] : environ[S.Lab]	0.12	0.00	28.07	**<.001**	-0.13	0.00	-40.79	**<.001**
stimulus[S.Standard] : environ[S.Field]	-0.05	0.00	-11.21	**<.001**	0.04	0.00	11.64	**<.001**
task[S.Ignore] : environ[S.Lab]	0.03	0.00	6.39	**<.001**	-0.06	0.00	-17.78	**<.001**
task[S.Ignore] : environ[S.Field]	-0.04	0.00	-9.69	**<.001**	-0.00	0.00	-1.19	.232
stimulus[S.Standard] : task[S.Ignore] : environ[S.Lab]	-0.01	0.00	-1.48	0.138	0.02	0.00	5.15	**<.001**
stimulus[S.Standard] : task[S.Ignore] : environ[S.Field]	0.01	0.00	2.77	**.006**	-0.00	0.00	-0.29	.773

Random Effects

σ^2	26.45				15.08			
τ_{00}	0.13 subject_id				0.12 subject_id			
	0.04 channel_id				0.06 channel_id			
τ_{11}	0.08 subject_id.environ[S.Lab]				0.07 subject_id.environ[S.Lab]			
	0.02 subject_id.environ[S.Field]				0.03 subject_id.environ[S.Field]			
					0.03 subject_id.stimulus[S.Standard]			
ρ_{01}	0.06 subject_id.environ[S.Lab]				0.19 subject_id.environ[S.Lab]			
	0.05 subject_id.environ[S.Field]				-0.06 subject_id.environ[S.Field]			
					-0.28 subject_id.stimulus[S.Standard]			

N	36 subject_id	36 subject_id
	31 channel_id	31 channel_id
Observations	4400681	4400681

Note. Factor variables sum-to-zero contrast-coded. *Estimate* = beta coefficient, *S.E.* = standard error, *stimulus* = oddball task stimulus (Deviant, Standard), *task* = cognitive task condition (Count, Ignore), *environ* = environmental context (Lab, Field, Campus), *prestim.s* = prestimulus EEG voltage (scaled), σ^2 = residual variance, τ_{00} = random intercept variance, τ_{11} = random slope variance, ρ_{01} = random effect correlation coefficient, *N* = number of levels per grouping variable.

P3 time-window

Type II Wald tests revealed a significant three-way interaction between stimulus, task, and environment for the P3 model (χ^2 (2) = 33.40, p < .001; see Table 1 for full decomposition). This interaction is resolved in Figure 4 (see Table 2 for the full model summary).

Pairwise comparisons of the difference in responses to standards and deviants were significant across all factorial combinations of Task and Environment (all p_{adj} <.038), with the exception of the Ignore task in the Campus condition (p_{adj} =.588). Sequential interaction contrasts revealed that the difference between stimulus responses became progressively smaller from Lab, to Field, to Campus conditions in both task manipulations (all p_{adj} <.001). Counting served to increase the difference between stimulus responses (i.e. amplify the magnitude of the P3 effect) in each environmental condition (all p_{adj} <.001).

In contrast to our analysis of the MMN time-window, differences between the Count and Ignore tasks in the P3 time-window were primarily driven by modulations of deviant tone processing. Deviant stimuli elicited significantly stronger positivities in both the Lab and Field conditions when being counted as opposed to being ignored (all p_{adj} <.001). Counted and ignored deviants did not significantly differ from one another in the Campus condition (p_{adj} = .999).

Discussion

The present study addressed the extent to which attentional processing is influenced by the situational context and its associated task demands. This involved comparing electrophysiological correlates of attentional processing between a traditional lab-based setting and two more naturalistic, dynamic scenarios involving walking and navigation. The manipulation of task instruction (i.e., to count or ignore oddball stimuli) across these three conditions enabled us to further compare the interaction of different levels of motor and environmental demands on top-down attentional engagement with task stimuli.

While participants were generally able to perform the oddball counting task with a high degree of accuracy across all three environmental conditions, linear mixed-effects regression analysis revealed that oddball counts were significantly less accurate in the Campus condition. This suggests the additional complexity entailed by this condition – both in terms of the more variable and unpredictable sensory stimuli it availed, and the more varied patterns of motor activity it solicited – were sufficiently distracting to degrade attentional focus on this simple cognitive task. By contrast, the additional motor demands and sensory complexity induced by walking around a sports field did not significantly affect task performance compared to sitting in a quiet room.

By contrast, the main findings from our ERP analyses revealed substantial differences in both the MMN and P3 components under 'real-world' versus laboratory conditions. This finding suggests that more ecologically-valid conditions involving motor co-ordination may modulate attentional processing despite ostensibly good behavioral task performance. Differences in ERPs between the Field and the Campus condition were subtle, which is perhaps surprising given the superior behavioural performance and relative simplicity of the Field condition. Differences between these conditions were limited to the Count task, and were more pronounced for the P3 response than the MMN.

With respect to previous studies of MMN and P3 components, findings from our Lab condition confirm the validity of the present results [17,53,54]. Specifically, the pre-dominant distribution of activity across fronto-central and centro-parietal regions of the scalp, respectively, was in accordance with findings from dual-task walking studies [55,56], as well as basic studies of the MMN [14,57] and P3 [58].

Increasing evidence from previous laboratory as well as real-world studies point to the fact

that attention on a given task depends on aspects such as task-difficulty, concurrent motor

demands, as well as environmental conditions [27]. However, especially in real-world scenarios

findings are rare and partly controversial. The present findings tap into this gap – for example

– by identifying distinctions in the effects of different real-world scenarios on attentional

resources allocated to cognitive task-performance, by varying cognitive task difficulty. In

contrast, previous studies with similar cognitive tasks reported no differences in P3 amplitude

between different real-world scenarios [3]. This apparent discrepancy can be explained by the

varied characteristics of the real-world scenarios included within these studies. While simple

conditions are comparable to the present sports field condition (e.g., walking on a circuit on

the lawn in front of the university), more complex scenarios differ significantly from walking

across the campus (e.g., obstacle course on an empty area or corridor).

Together with previous findings, we suggest that commonly-encountered everyday

situations (e.g., wayfinding, interacting with other people, etc.) may place attentional

resources under higher demand than physically-demanding situations (e.g., obstacle course)

within a controlled, unchanging environment. In accordance with previous real-world EEG

findings [23,25,27], evidence from laboratory studies, which used similar cognitive tasks, also

identified no significant effects of increased motor demands (e.g., slow walking vs. fast

walking on a treadmill) [20]. However, studies with more difficult cognitive tasks (higher

number of target-stimuli and increased complexity) report differences in amplitude, latency

and topography of ERP components with increasing motor demands [21]. It follows that the

amount of attention deployed during a given cognitive task is influenced by the complexity of

real-world situations, but more importantly, that the influence of environmental complexity is

also modulated by cognitive task difficulty.

In contrast to previous studies, the present findings revealed significant interactions

between stimulus, task, and condition for both ERP time-windows [3]. The difference between

standard and deviant stimulus responses diminished from lab to 'real-world' conditions. For

the MMN, this effect was predominantly driven by a reduction in the positivity evoked by

standard stimuli while walking; deviants, by contrast, evoked similar voltage levels

irrespective of environmental or task condition. Analogously, standard stimuli evoked less-

negative potentials in the P3 time-window during the walking conditions compared to the lab.

This pattern of results is broadly consistent with a predictive coding-style account of perceptual learning [59,60] and its attentional modulation via the mechanism of synaptic gain-control [61,62]. Under the assumption that frequent, regular presentations of the standard tone enable the brain to construct a precise model of stimulus features (e.g., pitch, duration), the sensory signals generated by the repeated stimuli are efficiently cancelled-out (or 'silenced') by descending model predictions [16,63]. The present findings suggest this ability may be reduced when moving through (and actively engaging with) more complex, dynamic environments. A possible explanation for this is that the additional, competing demands imposed on attention by more naturalistic scenarios may limit the brain's capacity to accurately and precisely model recurrent stimuli, such that they are subject to more elaborate cortical processing than they would ordinarily be afforded under laboratory conditions. The functional upshot of this account is that standard stimuli may essentially be treated more like deviants when occasioned in the context of more complex, variable, and uncertain conditions than those typically encountered in the lab.

Currently, we do not fully understand the extent to which environmental stimulation and movement-related information recruit attention resources. However, based on the findings presented here and in other recently published naturalistic studies [3,27] it is becoming increasingly evident that factors such as gross motor activity (as opposed to sitting or standing), cognitive task difficulty, and the degree of experimental control over ambient conditions (i.e. laboratory vs. real-world settings), seem to be highly relevant determinants of attention and cognitive resource allocation. Such results highlight potential limitations in the generalizability of standard laboratory experiments to the more complex, dynamic conditions encountered in everyday life. However, it is also important to note some of the limitations and ongoing challenges faced by real-world designs, which we consider below.

The present study focused on EEG signals in order to avoid potential complications or confounds introduced by encumbering the participant with multiple measurement devices. One consequence of this decision was that we were unable to augment our analysis of cognitive task performance with information concerning gait parameters. Based on previous findings that identified a maintenance of cognitive performance in dual-task walking conditions by reducing walking speed [64], it can only be speculated that gait parameters in the present study may have changed with increased complexity of the situation (across the

campus vs. around the sports field) and thereby, the amount of attention focused on the

cognitive task could be maintained; however, this must remain a question for future research.

Moreover, while advancing technological innovations are revolutionizing the study of

cognition under ecologically-valid conditions, mobile EEG designs still face a number of

challenges. In addition to the concern that performing tasks and activities while

wearing/carrying recording devices may perturb normal functioning, the amount and quality

of the data collected under such conditions is often reduced compared to standard laboratory

conditions. Although we attempted to mitigate this potential issue through the application of

sophisticated artifact rejection techniques, epoch rejection rates varied considerably amongst

individuals. The problem of noisy data collection might also have been compounded by the

measurement imprecision introduced by the delayed (and in some cases, repeated)

transmission of event markers. While we took care to correct these errors without introducing

bias (i.e., applying a uniform adjustment designed to ameliorate event marker lag across all

conditions prior to statistical analysis), the additional variance introduced into the data may

have limited our sensitivity to fine-grained differences between conditions.

Despite these cautionary remarks, the rapid development of mobile EEG systems and

software solutions suggests that many of the challenges mentioned above can be overcome in

future studies. Methodological innovations such as the development of ear-EEG systems

promise to enable the acquisition of high-quality data while participants move about freely

[65,66]. Sophisticated algorithms for capturing and characterizing gait-related artefacts are also

being developed (see, e.g., [67]). Such advancements will prove critical for enhancing the

quality and robustness of mobile EEG data collected under complex and variable

environmental conditions.

Existing findings from naturalistic EEG studies provide valuable insights concerning the

nature of well-established markers of cognitive processing such as the MMN and P3

components. The present results align well with recently published mobile EEG studies and

fill an important gap in this literature about the effects of environmental complexity, motor

demands, and cognitive task load on attention within naturalistic scenarios. Furthermore, our

findings highlight the necessity of extending future cognitive research programs beyond the

confines of the laboratory. Ideally, such studies will combine complementary measurement

devices as they become increasingly portable (e.g., ear-EEG & eye-tracking), and address a
greater variety of everyday situations.
581

Acknowledgements

AWC and CAH are supported by Australian Government Research Training Program (RTP)
scholarships. IBS acknowledges the support of an Australian Research Council Future
Fellowship (FT160100437). This research was funded by the University of South Australia
under the Research Themes Investment Scheme. We would also like to thank the
undergraduate students in course BEHL 3021 ("Cognitive Neuroscience") at the University of
South Australia who participated in this study and helped with data collection as part of their
class project.

References

1. Schröger, E. & Wolff, C. Behavioral and electrophysiological effects of task-irrelevant
sound change: A new distraction paradigm. *Cogn. Brain Res.* **7**, 71–87 (1998).
2. Di Flumeri, G. *et al.* EEG-based mental workload neurometric to evaluate the impact
of different traffic and road conditions in real driving settings. *Front. Hum. Neurosci.*
12, 509 (2018).
3. Reiser, J. E., Wascher, E. & Arnau, S. Recording mobile EEG in an outdoor
environment reveals cognitive-motor interference dependent on movement complexity.
Sci. Rep. **9**, 1–14 (2019).
4. Näätänen, R. & Alho, K. Mismatch negativity-a unique measure of sensory processing
in audition. *Int. J. Neurosci.* **80**, 317–337 (1995).
5. Polich, J. *et al.* P300 topography of amplitude/latency correlations. *Brain Topogr.* **9**,
275–282 (1997).
6. Smith, M. E. *et al.* The intracranial topography of the P3 event-related potential elicited
during auditory oddball. *Electroencephalogr. Clin. Neurophysiol.* **76**, 235–248 (1990).
7. Polich, J. Updating P300: an integrative theory of P3a and P3b. *Clin. Neurophysiol.*
118, 2128–2148 (2007).
8. Kok, A. Internal and external control: a two-factor model of amplitude change of
event-related potentials. *Acta Psychol.* **74**, 213–236 (1990).
9. Polich, J. & Kok, A. Cognitive and biological determinants of P300: an integrative
review. *Biol. Psychol.* **41**, 103–146 (1995).

10. Kramer, A. F. & Strayer, D. L. Assessing the development of automatic processing: an

application of dual-task and event-related brain potential methodologies. *Biol. Psychol.*

26, 231–267 (1988).

11. Wickens, C., Kramer, A., Vanasse, L. & Donchin, E. Performance of concurrent tasks:

a psychophysiological analysis of the reciprocity of information-processing resources.

Science. **221**, 1080–1082 (1983).

12. Nash, A. J. & Fernandez, M. P300 and allocation of attention in dual-tasks. *Int. J.*

Psychophysiol. **23**, 171–180 (1996).

13. Allison, B. Z. & Polich, J. Workload assessment of computer gaming using a single-

stimulus event-related potential paradigm. *Biol. Psychol.* **77**, 277–283 (2008).

14. Duncan, C. C. *et al.* Event-related potentials in clinical research: guidelines for

eliciting, recording, and quantifying mismatch negativity, P300, and N400. *Clin.*

Neurophysiol. **120**, 1883–1908 (2009).

15. Näätänen, R., Kujala, T. & Winkler, I. Auditory processing that leads to conscious

perception: a unique window to central auditory processing opened by the mismatch

negativity and related responses. *Psychophysiology* **48**, 4–22 (2011).

16. Garrido, M. I., Kilner, J. M., Stephan, K. E. & Friston, K. J. The mismatch negativity: a

review of underlying mechanisms. *Clin. Neurophysiol.* **120**, 453–463 (2009).

17. Näätänen, R., Paavilainen, P., Rinne, T. & Alho, K. The mismatch negativity (MMN)

in basic research of central auditory processing: a review. *Clin. Neurophysiol.* **118**,

2544–2590 (2007).

18. Pazo-Alvarez, P., Cadaveira, F. & Amenedo, E. MMN in the visual modality: a review.

Biol. Psychol. **63**, 199–236 (2003).

19. Näätänen, R., Gaillard, A. W. K. & Mäntysalo, S. Early selective-attention effect on

evoked potential reinterpreted. *Acta Psychol.* **42**, 313–329 (1978).

20. Gramann, K., Gwin, J. T., Bigdely-Shamlo, N., Ferris, D. P. & Makeig, S. Visual

evoked responses during standing and walking. *Front. Hum. Neurosci.* **4**, 1-12 (2010).

21. De Sanctis, P., Butler, J. S., Malcolm, B. R. & Foxe, J. J. Recalibration of inhibitory

control systems during walking-related dual-task interference: a mobile brain-body

imaging (MOBI) study. *Neuroimage* **94**, 55–64 (2014).

22. Nenna, F., Do, C. T., Protzak, J. & Gramann, K. Alteration of brain dynamics during

dual-task overground walking. *Eur. J. Neurosci.* **00**, 1–17 (2020).

23. Zink, R., Hunyadi, B., Van Huffel, S. & De Vos, M. Mobile EEG on the bike:

disentangling attentional and physical contributions to auditory attention tasks. *J.*

Neural Eng. **13**, 1-11 (2016).

24.   Scanlon, J. E. M., Townsend, K. A., Cormier, D. L., Kuziek, J. W. P. & Mathewson, K.

649   E. Taking off the training wheels: Measuring auditory P3 during outdoor cycling using

650   an active wet EEG system. *Brain Res.* **1716**, 50–61 (2019).

651   25.   Robles, D. *et al.* EEG in motion: Using an oddball task to explore motor interference in

652   active skateboarding. *Eur. J. Neurosci.* (2021).

653   26.   De Vos, M., Gandras, K. & Debener, S. Towards a truly mobile auditory brain–

654   computer interface: exploring the P300 to take away. *Int. J. Psychophysiol.* **91**, 46–53

655   (2014).

656   27.   Ladouce, S., Donaldson, D. I., Dudchenko, P. A. & Ietswaart, M. Mobile EEG

657   identifies the re-allocation of attention during real-world activity. *Sci. Rep.* **9**, 1–10

658   (2019).

659   28.   Aspinall, P., Mavros, P., Coyne, R. & Roe, J. The urban brain: analysing outdoor

660   physical activity with mobile EEG. *Br J Sport. Med.* **49**, 272–276 (2015).

661   29.   Pizzamiglio, S., Naeem, U., Abdalla, H. & Turner, D. L. Neural correlates of single-

662   and dual-task walking in the real world. *Front. Hum. Neurosci.* **11**, 1-12 (2017).

663   30.   Park, J. L. & Donaldson, D. I. Detecting the neural correlates of episodic memory with

664   mobile EEG: Recollecting objects in the real world. *Neuroimage* **193**, 1–9 (2019).

665   31.   Piñeyro Salvidegoitia, M. *et al.* Out and about: subsequent memory effect captured in a

666   natural outdoor environment with smartphone EEG. *Psychophysiology* **56**, 1-15 (2019).

667   32.   Brilmayer, I., Sassenhagen, J., Bornkessel-Schlesewsky, I. & Schlesewsky, M.

668   Domain-general neural correlates of dependency formation: Using complex tones to

669   simulate language. *Cortex* **93**, 50–67 (2017).

670   33.   Mathôt, S., Schreij, D. & Theeuwes, J. OpenSesame: An open-source, graphical

671   experiment builder for the social sciences. *Behav. Res. Methods* **44**, 314–324 (2012).

672   34.   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh inventory.

673   *Neuropsychologia* **9**, 97–113 (1971).

674   35.   Salisbury, D. F., Rutherford, B., Shenton, M. E. & McCarley, R. W. Button-pressing

675   affects P300 amplitude and scalp topography. *Clin. Neurophysiol.* **112**, 1676–1684

676   (2001).

677   36.   Delorme, A. & Makeig, S. EEGLAB: an open source toolbox for analysis of single-

678   trial EEG dynamics including independent component analysis. *J. Neurosci. Methods*

679   **134**, 9–21 (2004).

37.   Lopez-Calderon, J. & Luck, S. J. ERPLAB: an open-source toolbox for the analysis of

event-related potentials. *Front. Hum. Neurosci.* **8**, 1-14 (2014).

38.   Chang, C.-Y., Hsu, S.-H., Pion-Tonachini, L. & Jung, T.-P. Evaluation of artifact
subspace reconstruction for automatic EEG artifact removal. in *40th Annual*
International Conference of the IEEE Engineering in Medicine and Biology Society
(EMBC) 1242–1245 (IEEE, 2018).

39.   Mullen, T. R. *et al.* Real-time neuroimaging and cognitive monitoring using wearable
dry EEG. *IEEE Trans. Biomed. Eng.* **62**, 2553–2567 (2015).

40.   Bell, A. J. & Sejnowski, T. J. An information-maximization approach to blind
separation and blind deconvolution. *Neural Comput.* **7**, 1129–1159 (1995).

41.   Lee, T.-W., Girolami, M. & Sejnowski, T. J. Independent component analysis using an
extended infomax algorithm for mixed subgaussian and supergaussian sources. *Neural*
Comput. **11**, 417–441 (1999).

42.   Pion-Tonachini, L., Kreutz-Delgado, K. & Makeig, S. ICLabel: An automated
electroencephalographic independent component classifier, dataset, and website.
Neuroimage **198**, 181–197 (2019).

43.   R Core Team. A language and environment for statistical computing. https://www.r-
project.org/ (2019).

44.   R Studio Team. RStudio: Integrated Development for R. RStudio.
http://www.rstudio.com/ (2015).

45.   Bates, D. *et al.* Package 'lme4'. *Convergence* **12** (2015).

46.   Alday, P. M. How much baseline correction do we need in ERP research? Extended
GLM model can replace baseline correction while lifting its limits. *Psychophysiology*
56, 1-14 (2019).

47.   Fox, J. & Weisberg, S. *An R companion to applied regression*. (Sage publications,
2018).

48.   Lüdecke, D. sjPlot: Data Visualization for Statistics in Social Science.
https://doi.org/10.5281/zenodo.1308157 (2020).

49.   Lenth, R. emmeans: Estimated marginal means, aka least-squares means (Version
1.3.0)[R package]. (2020).

50.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1-6 (2019).

51.   MacGregor-Fors, I. & Payton, M. E. Contrasting diversity values: statistical inferences
based on overlapping confidence intervals. *PLoS One* **8**, e56794 (2013).

52.   Pernet, C. R. *et al.* EEG-BIDS, an extension to the brain imaging data structure for
electroencephalography. *Sci. data* **6**, 1–5 (2019).

53.   Dariusz, A. *et al.* The response relevance of visual stimuli modulates the P3 component
and the underlying sensorimotor network. *Sci. Rep.* **10**, 1-20 (2020).

54.   Näätänen, R. Mismatch negativity (MMN): perspectives for application. *Int. J.*
Psychophysiol. **37**, 3–10 (2000).

55.   Holtzer, R. *et al.* Online fronto-cortical control of simple and attention-demanding
locomotion in humans. *Neuroimage* **112**, 152–159 (2015).

56.   Mirelman, A. *et al.* Increased frontal brain activation during walking while dual
tasking: an fNIRS study in healthy young adults. *J. Neuroeng. Rehabil.* **11**, 1-7 (2014).

57.   Näätänen, R., Pakarinen, S., Rinne, T. & Takegata, R. The mismatch negativity
(MMN): towards the optimal paradigm. *Clin. Neurophysiol.* **115**, 140–144 (2004).

58.   Debener, S., Makeig, S., Delorme, A. & Engel, A. K. What is novel in the novelty
oddball paradigm? Functional significance of the novelty P3 event-related potential as
revealed by independent component analysis. *Cogn. Brain Res.* **22**, 309–321 (2005).

59.   Friston, K. A theory of cortical responses. *Philos. Trans. R. Soc. B Biol. Sci.* **360**, 815–
836 (2005).

60.   Rao, R. P. N. & Ballard, D. H. Predictive coding in the visual cortex: a functional
interpretation of some extra-classical receptive-field effects. *Nat. Neurosci.* **2**, 79–87
(1999).

61.   Feldman, H. & Friston, K. Attention, uncertainty, and free-energy. *Front. Hum.*
Neurosci. **4**, 1-23 (2010).

62.   Kok, P., Rahnev, D., Jehee, J. F. M., Lau, H. C. & De Lange, F. P. Attention reverses
the effect of prediction in silencing sensory signals. *Cereb. cortex* **22**, 2197–2206
(2012).

63.   Todd, J., Finch, B., Smith, E., Budd, T. W. & Schall, U. Temporal processing ability is
related to ear-asymmetry for detecting time cues in sound: A mismatch negativity
(MMN) study. *Neuropsychologia* **49**, 69–82 (2011).

64.   Patel, P., Lamar, M. & Bhatt, T. Effect of type of cognitive task and walking speed on
cognitive-motor interference during dual-task walking. *Neuroscience* **260**, 140–148
(2014).

65.   Bleichner, M. G. & Debener, S. Concealed, unobtrusive ear-centered EEG acquisition:
cEEGrids for transparent EEG. *Front. Hum. Neurosci.* **11**, 1-14 (2017).

66.   Hölle, D., Meekes, J. & Bleichner, M. G. Mobile ear-EEG to study auditory attention
in everyday life. *Behav. Res. Methods* 1–12 (2021).

67.   Jacobsen, N. S. J., Blum, S., Witt, K. & Debener, S. A walk in the park? Characterizing

gait-related artifacts in mobile EEG recordings. *Eur. J. Neurosci.* 1-20 (2020).