No evidence of human genome integration of SARS-CoV-2 found by long-read DNA sequencing

Nathan Smits[1,10], Jay Rasmussen[2,10], Gabriela O. Bodea[1,2,10], Alberto A. Amarilla[3,10], Patricia Gerdes[1], Francisco J. Sanchez-Luque[4,5], Prabha Ajjikuttira[2], Naphak Modhiran[3], Benjamin Liang[3], Jamila Faivre[6], Ira W. Deveson[7,8], Alexander A. Khromykh[3,9], Daniel Watterson[3,9], Adam D. Ewing[1], Geoffrey J. Faulkner[1,2,11,*]

[1]Mater Research Institute - University of Queensland, TRI Building, Woolloongabba QLD 4102, Australia.

[2]Queensland Brain Institute, University of Queensland, Brisbane QLD 4072, Australia.

[3]School of Chemistry and Molecular Biosciences, University of Queensland, Brisbane QLD 4072, Australia.

[4]GENYO, Pfizer-University of Granada-Andalusian Government Centre for Genomics and Oncological Research, PTS Granada 18016, Spain.

[5]MRC Human Genetics Unit, Institute of Genetics and Cancer (IGC), University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, United Kingdom.

[6]INSERM, U1193, Paul-Brousse University Hospital, Hepatobiliary Centre, Villejuif 94800, France.

[7]Kinghorn Centre for Clinical Genomics, Garvan Institute of Medical Research, Sydney NSW 2010, Australia.

[8]St Vincent's Clinical School, Faculty of Medicine, University of New South Wales, Sydney NSW 2052, Australia.

[9]Australian Infectious Diseases Research Centre, Global Virus Network Centre of Excellence, Brisbane QLD 4072, Australia.

[10]These authors contributed equally.

[11]Lead contact.

*Correspondence should be addressed to G.J.F. (faulknergj@gmail.com).

SUMMARY

A recent study proposed severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)

hijacks the LINE-1 (L1) retrotransposition machinery to integrate into the DNA of infected

cells. If confirmed, this finding could have significant clinical implications. Here, we applied

deep (>50×) long-read Oxford Nanopore Technologies (ONT) sequencing to HEK293T cells

infected with SARS-CoV-2, and did not find the virus integrated into the genome. By

examining ONT data from separate HEK293T cultivars, we completely resolved 78 L1

insertions arising *in vitro* in the absence of L1 overexpression systems. ONT sequencing

applied to hepatitis B virus (HBV) positive liver cancer tissues located a single HBV

insertion. These experiments demonstrate reliable resolution of retrotransposon and

exogenous virus insertions via ONT sequencing. That we found no evidence of SARS-CoV-2

integration suggests such events are, at most, extremely rare *in vivo*, and therefore are

unlikely to drive oncogenesis or explain post-recovery detection of the virus.

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is a positive-sense single-stranded ~30kbp polyadenylated RNA betacoronavirus (V'kovski et al., 2020; Wu et al., 2020). SARS-CoV-2 does not encode a reverse transcriptase (RT) and therefore is not expected to integrate into genomic DNA as part of its life cycle. This assumption is of fundamental importance to the accurate diagnosis and potential long-term clinical consequences of SARS-CoV-2 infection, as demonstrated by other viruses known to incorporate into genomic DNA, such as human immunodeficiency virus 1 (HIV-1) and hepatitis B virus (HBV) (Bill and Summers, 2004; Fujimoto et al., 2012; Jiang et al., 2012; Nagaya et al., 1987). Notably, a recent work by Zhang *et al.* reported potential evidence of SARS-CoV-2 integration into the genome of infected human cells (Zhang et al., 2021). Prior analyses of mammalian genome sequences, as well as *in vivo* and *in vitro* experimental data, indicate single-stranded RNA viruses can act as templates for endogenous RTs (Belyi et al., 2010; Feschotte and Gilbert, 2012; Geuking et al., 2009; Horie et al., 2010; Kawasaki et al., 2021; Klenerman et al., 1997). These studies provide a conceptual basis to further investigate genomic integration of SARS-CoV-2, as pursued by Zhang *et al.*.

LINE-1 (L1) retrotransposons reside in all mammalian genomes (Kazazian and Moran, 2017). In humans, L1 transcribes a bicistronic mRNA encoding two proteins, ORF1p and ORF2p, essential to L1 mobility (Moran et al., 1996). ORF2p possesses endonuclease (EN) and RT activities, and exhibits strong *cis* preference for reverse transcription of L1 mRNA (Doucet et al., 2015; Garcia-Perez et al., 2007; Kulpa and Moran, 2006; Monot et al., 2013; Moran et al., 1996; Wei et al., 2001). Nonetheless, the L1 protein machinery can *trans* mobilise polyadenylated cellular RNAs, especially those produced by *Alu* and SINE-VNTR-*Alu* (SVA) retrotransposons, but also including protein-coding gene mRNAs (Dewannieux et al., 2003; Esnault et al., 2000; Garcia-Perez et al., 2007; Hancks et al., 2011; Raiz et al., 2012). Somatic L1 mobilisation *in cis* is observed in embryonic cells, the neuronal lineage, and various cancers (Evrony et al., 2015; Feusier et al., 2019; Rodriguez-Martin et al., 2020; Sanchez-Luque et al., 2019; Schauer et al., 2018; Scott et al., 2016). By contrast, somatic L1-mediated *trans* mobilisation is apparently rare *in vivo* (Evrony et al., 2015; Rodriguez-Martin et al., 2020; Sanchez-Luque et al., 2019) and is likely repressed by various mechanisms (Ahl et al., 2015; Deniz et al., 2019; Doucet et al., 2015; Ewing et al., 2020; Sanchez-Luque et al., 2019). While *Alu* and, to a lesser extent, SVA are readily mobilised in cultured cell assays by L1 proteins, the same machinery produces less than one non-retrotransposon cellular RNA insertion for every 2000 L1 insertions (Dewannieux et al., 2003; Hancks et al., 2011; Wei et

al., 2001). Both *cis* and *trans* L1-mediated insertions incorporate target site duplications (TSDs) and a 3′ polyA tract, and integrate at the degenerate L1 EN motif 5′-TTTT/AA (Dewannieux et al., 2003; Esnault et al., 2000; Garcia-Perez et al., 2007; Gilbert et al., 2005; Hancks et al., 2009; Jurka, 1997; Moran et al., 1996; Raiz et al., 2012). These sequence hallmarks can together discriminate artifacts from genuine insertions (Faulkner and Billon, 2018).

In their work, Zhang *et al.* overexpressed L1 in HEK293T cells, infected these with SARS-CoV-2, and identified DNA fragments of the virus through PCR amplification. These results, alongside other less direct (Kazachenka and Kassiotis, 2021; Yan et al., 2021) analyses, were interpreted as evidence of SARS-CoV-2 genomic integration. Crucially, Zhang *et al.* then detected 63 putative SARS-CoV-2 integrants by Oxford Nanopore Technologies (ONT) long-read sequencing. Of these, only a single integrant on chromosome X was spanned by an ONT read aligned to one locus, and was flanked by potential TSDs (**Figure 1A**). However, this SARS-CoV-2 integrant did not incorporate a 3′ polyA tract, as is expected for an L1-mediated insertion, and involved an unusual 28kb internal deletion of the SARS-CoV-2 sequence. Overall, the SARS-CoV-2 integrants reported by Zhang *et al.* were 26-fold enriched in exons, despite the L1 EN showing no preference for these regions (Flasch et al., 2019; Sultana et al., 2019). Zhang *et al.* also used Illumina short-read sequencing to map putative SARS-CoV-2 integration junctions in HEK293T cells without L1 overexpression. A lack of spanning reads and the tendency of Illumina library preparation to produce artefacts (Treiber and Waddell, 2017) leave this analysis open to interpretation.

The application of ONT sequencing to HEK293T cells nonetheless held conceptual merit. ONT reads can span germline and somatic retrotransposition events end-to-end, and resolve the sequence hallmarks of L1-mediated integration (Ewing et al., 2020; Siudeja et al., 2021). Through this approach, we previously found two somatic L1 insertions in the liver tumour sample of an individual positive for hepatitis C virus (HCV), a ~10kbp positive-sense single-stranded non-polyadenylated RNA virus (Lauer and Walker, 2001), including one PCR-validated L1 insertion spanned by a single ONT read (Ewing et al., 2020; Shukla et al., 2013). Moreover, HEK293T cells are arguably a context favourable to L1 activity. They express L1 ORF1p (Philippe et al., 2016), accommodate engineered L1-mediated retrotransposition *in cis* and *in trans* (Hancks et al., 2011; Kubo et al., 2006; Niewiadomska et al., 2007; Sanchez-Luque et al., 2019), and support SARS-CoV-2 viral replication (**Figure S1**). Endogenous L1-mediated insertions can be detected in cell culture by genomic analysis of separate cultivars derived from a common population (Klawitter et al., 2016; Nguyen et

Figure 1. Key potential SARS-CoV-2 insertions reported by Zhang *et al.*. (A) A cartoon summarising the features of a putative SARS-CoV-2 integrant on chromosome X. Numerals underneath the SARS-CoV-2 sequence represent positions relative to the QLD02 virus isolate. Potential TSDs are shown as red triangles, and motifs resembling potential pre-integration L1 EN recognition sites are highlighted, with question marks in labels intended to flag uncertain L1 involvement. No 3′ polyA tract was found. Homologous regions at sequence junctions are marked. One spanning ONT read is positioned underneath the cartoon and its identifier is displayed. **(B)** As for (A), except showing an ONT read spanning two SARS-CoV-2 insertions, on chromosome 22 and chromosome 1. The alignments to chromosome 22 were flagged as supplementary by the minimap2 aligner. 3′ polyA tracts are represented as green rectangles. Note: these chromosome 22 and chromosome X instances are the key examples reported by Zhang *et al.* in support of SARS-CoV-2 genomic integration. Neither example has a complete set of retrotransposition hallmarks (TSD, 3′ polyA tract, L1 EN motif) and the support of a uniquely aligned ONT read.

al., 2018). Based on this experimental rationale, we sought to replicate the central findings of

Zhang *et al.* and, after deeply ONT sequencing SARS-CoV-2-infected HEK293T cells, did

not detect SARS-CoV-2 genomic integration.

RESULTS

We ONT sequenced (~54× genome-wide depth, read length N50 ~ 39kbp) genomic DNA

harvested from HEK293T cells infected with SARS-CoV-2 at a multiplicity of infection

(MOI) of 1.0, as well as mock infected cells (~28× depth, N50 ~ 47kbp) (**Figures 2A** and **S1**,

and **Table S1**). As positive controls, we ONT sequenced the tumour and non-tumour liver

tissue of a HBV-positive hepatocellular carcinoma patient (Shukla et al., 2013). HBV is a

DNA virus known to be integrated into sites of genomic damage via DNA double-strand

break repair (Bill and Summers, 2004). As negative controls, we used the aforementioned

HCV-positive hepatocellular carcinoma samples, and a normal liver sample (Ewing et al.,

2020) (**Table S1**). We viewed HCV-infected samples as a suitable negative control because

HCV and SARS-CoV-2 are both positive-sense single-stranded RNA viruses, yet HCV is not

polyadenylated and is therefore unlikely to attract the L1 machinery, and has not been found

to integrate into infected hepatocytes or liver tumour genomes (Fujimoto et al., 2012; Lauer

and Walker, 2001). To these data, we added those of Zhang *et al*., and then used the

Transposons from Long DNA Reads (TLDR) (Ewing et al., 2020) software to call SARS-

CoV-2, HBV, HCV and non-reference retrotransposon insertions spanned by at least one

uniquely aligned ONT read. TLDR detected no SARS-CoV-2, HBV or HCV insertions.

In total, TLDR identified 575 non-reference human-specific L1 (L1HS) insertions,

which were typically flanked by TSDs with a median length of 14bp (**Figure 2B** and **Table**

S2). No tumour-specific L1 insertions were found, apart from the two previously detected in

the HCV-infected liver tumour (Ewing et al., 2020; Shukla et al., 2013). Seventy-eight L1

insertions were found only in our SARS-CoV-2 infected HEK293T cells (66) or the mock

infected control (12) and produced TSDs with a median length of 14bp (**Figure 2B**). Of the

78 events, 69 (88.5%) were detected by a single spanning read and 13 carried a 3′

transduction (Holmes et al., 1994; Moran et al., 1999) (**Table S2**). After random

downsampling, the more deeply sequenced SARS-CoV-2 infected HEK293T cells still had

more than 2-fold more putative cultivar-specific L1 insertions than the mock infected

HEK293T cells. Next, we chose at random 6/69 L1 insertions detected by one spanning read

for manual curation and PCR validation. All 6 L1 insertions bore a TSD and a 3′ polyA tract,

and integrated at a degenerate L1 EN motif (**Figures 2D, 2E** and **S2A-S2D**). Three were 5′

Figure 2. Detection of endogenous L1-mediated retrotransposition in human cells. (A) Experimental design. HEK-293T cells were divided into two populations (cultivars), which were then either SARS-CoV-2 infected or mock infected. DNA was extracted from each cultivar, as well as from hepatocellular carcinoma patient samples, and subjected to ONT sequencing. ONT reads were used to call non-reference L1 and virus insertions with TLDR, which also resolves TSDs and other retrotransposition hallmarks. TSDs: red triangles; polyA tract: green rectangle; ONT read: blue rectangle. Some illustrations are adapted from a previous study (Ewing et al., 2020). **(B)** TSD size distribution for non-reference L1 insertions, as annotated by TLDR, as well as cultivar-specific L1 insertions found only in either our HEK293T cells infected with SARS-CoV-2 or our mock infected cells. Pie charts indicate the percentages of exonic, intronic and intergenic insertions, annotated by RefSeq coordinates. **(C)** As for (B), except showing data for *Alu* insertions. **(D)** Detailed characterisation of an L1 insertion detected in SARS-CoV-2 infected HEK293T cells by a single spanning ONT read aligned to chromosome 14. Nucleotides highlighted in red correspond to the integration site TSD. Underlined nucleotides correspond to the L1 EN motif. The cartoon indicates a full-length L1HS insertion flanked by TSDs (red triangles), and a 3′ polyA tract (green), with the underneath numeral representing the 5′ end position relative to the mobile L1HS sequence L1.3 (Dombroski et al., 1993). The relevant spanning ONT read, with identifier, is also positioned underneath the cartoon. Symbols (α, β, δ, γ) represent the approximate position of primers used for empty/filled site and L1-genome junction PCR validation reactions. These are displayed in gel images if successful. Ladder band sizes are as indicated, NTC; non-template control. Red triangles indicate L1 amplicon expected sizes (empty triangle: no product; filled triangle: capillary sequenced on-target product). Blue triangles indicate expected empty site sizes. **(E)** As for (D), except for a 5′ inverted/deleted L1HS located on chromosome 18. Please see Figures S1 and S2, and Tables S1 and S2 for further information.

inverted (Kazazian et al., 1988; Ostertag and Kazazian, 2001) (**Figures 2E, S2C** and **S2D**)

and one carried a 3′ transduction (Holmes et al., 1994) traced to a mobile (Rodriguez-Martin

et al., 2020) full-length non-reference L1HS (**Figure S2C**). Three PCR amplified in the

SARS-CoV-2 and mock infected samples (**Figures 2D**, **2E** and **S2A**) and three did not

amplify in either sample (**Figures S2B-S2D**). The 6 integration sites were on average

spanned by 86 reads not containing the L1 insertion (**Figure S2E**), a ratio (1:86) suggesting

the L1s were absent from most cells. An additional analysis revealed 293 putative *Alu* (291)

and SVA (2) insertions specific to either one of the HEK293T populations, with 290 of these

found in the SARS-CoV-2 infected cells and 275 (93.9%) detected by a single spanning read

(**Table S2**). The median TSD size for this cohort was 13bp (**Figure 2C**). Altogether, these

and earlier (Ewing et al., 2020; Siudeja et al., 2021) experiments show that lone spanning

ONT reads can recover *bona fide* retrotransposition events, and highlight endogenous L1

activity in HEK293T cells lacking L1 overexpression systems.

We next tested whether our computational analysis parameters excluded genuine

HBV, HCV or SARS-CoV-2 insertions. We directly aligned our ONT reads to the genome of

the SARS-CoV-2 isolate (QLD002, GISAID EPI_ISL_407896) used here, as well as to a

geographically diverse set of HBV and HCV genomes (**Table S1**), and a highly mobile L1HS

sequence (Dombroski et al., 1993). In total, 3.6% of our ONT sequence bases aligned to

L1HS, whereas no alignments to the SARS-CoV-2 or HCV genomes were observed (**Figure**

3A). One read from the HBV-infected non-tumour liver sample aligned to 2,770bp of a HBV

genotype B isolate, and the remaining 2,901bp aligned to an intergenic region of

chromosome 2 (**Figure 3B** and **Table S2**). To validate this HBV insertion, we PCR amplified

and capillary sequenced its 3′ junction (**Figure 3B**). The HBV sequence was linearised and

rearranged (**Figure 3B**) as per prior reports (Fujimoto et al., 2012; Jiang et al., 2012; Nagaya

et al., 1987). Direct inspection of ONT read alignments thus recovered a HBV integrant,

which are found in ~1 per 10^1-10^4 infected hepatocytes (Mason et al., 2016; Tu et al., 2018),

yet did not reveal reads alignable to the SARS-CoV-2 genome in our ONT datasets.

Reanalysing the ONT data generated by Zhang *et al.*, we found 555 reads (out of ~12

million) that generated an alignment of ≥100bp to the SARS-CoV-2 genome (**Figure 3A**).

These reads (median length 924bp) were however 65.6% shorter than the overall dataset

(2,686kbp) and were comprised of a much higher average proportion of SARS-CoV-2

sequence (52.3%) than the proportion of L1HS sequence found in reads aligned to L1HS

(17.1%). Of the 555 reads, 79 generated an alignment of ≥100bp to the human genome,

including one matching the aforementioned integrant on chromosome X that lacked a 3′

Figure 3. ONT reads occasionally align to viral genome sequences. (A) Percentages of total ONT sequence alignable to L1HS (left), SARS-CoV-2 (middle) and HBV (right) isolate genomes. Read counts for SARS-CoV-2 and HBV are provided above histogram columns. No reads were aligned to the HCV isolate genomes. HEK293T data were generated here (SARS-CoV-2, mock) or by Zhang *et al.*. HCC tumour/non-tumour liver pairs were sequenced here (HCC32; HBV-positive) or previously (Ewing et al., 2020) (HCC33; HCV-positive). Normal liver ONT sequencing from our prior work (Ewing et al., 2020) was included as an additional control. **(B)** A HBV insertion detected in non-tumour liver. In this example, an ONT read from the non-tumour liver of HCC32 spanned the 3′ junction of a HBV integrant located on chromosome 2. Of the HBV isolate genomes considered here, this read aligned best to a representative of genotype B (Genbank accession AB602818). The HBV sequence was rearranged consistent with its linearisation prior to integration (Fujimoto et al., 2012; Jiang et al., 2012; Nagaya et al., 1987). Numerals indicate positions relative to AB602818. Symbols (β, δ) represent the approximate position of primers used to PCR validate the HBV insertion. The gel image at right shows the 3′ junction PCR results. Ladder band sizes are as indicated. The red filled triangle indicates an on-target product confirmed by capillary sequencing. Repeated attempts to PCR amplify the 5′ junction of the HBV integrant did not return an on-target product, perhaps due to a genomic deletion at the insertion site. **(C)** Percentages of exonic (black), intronic (grey) and intergenic (blue) genomic alignment breakpoints for ONT reads also aligned to the SARS-CoV-2 genome, and also for the non-reference L1-mediated insertions reported here. Genomic features were annotated according to RefSeq coordinates. Please see Tables S1 and S2 for further information.

polyA tract (**Figure 1A**). An analysis of the corresponding 79 human genome alignment

breakpoints, which could be interpreted as putative SARS-CoV-2 insertion points, as per

Zhang *et al.*, indicated 36.2% were exonic (**Figure 3C**). By comparison, 1.1% of all non-

reference L1-mediated insertions reported here were exonic (**Figure 3C**), as were 1.3% and

1.7% of cultivar-specific L1 and *Alu* insertions, respectively (**Figures 2B** and **2C**). Finally,

we investigated why TLDR called neither of the two SARS-CoV-2 insertions highlighted by

Zhang *et al.* (**Figure 1**), and found ambiguity in the ONT read alignments supporting these

examples led to their exclusion. Specifically, the putative chromosome X insertion (**Figure**

1A) was filtered due to its short 5′ genomic flank alignment, whereas the chromosome 22

example was filtered because the corresponding ONT read alignment was marked as

supplementary to another alignment on chromosome 1 (**Figure 1B**). These analyses

confirmed SARS-CoV-2 alignable reads were present in the Zhang *et al.* ONT dataset, yet

these reads were unusually short and could include molecular artifacts interpreted by Zhang

et al. as SARS-CoV-2 integrants.

DISCUSSION

We did not observe L1-mediated SARS-CoV-2 genomic integration in HEK293T cells,

despite availability of the L1 machinery (Hancks et al., 2011; Kubo et al., 2006;

Niewiadomska et al., 2007; Philippe et al., 2016; Sanchez-Luque et al., 2019) and detected

L1, *Alu* and SVA retrotransposition events. The higher number of L1 and *Alu* insertions

found in our SARS-CoV-2-infected HEK293T cells is of potential interest given viral

infection can repress host factors limiting L1 activity (Hrecka et al., 2011; Laguette et al.,

2011; Zhao et al., 2013). This preliminary finding perhaps indicates SARS-CoV-2 infection

could increase L1 or *Alu* retrotransposition *in vitro*, a possibility requiring experimental

confirmation. The comparative rarity of SVA insertions, and absence of SARS-CoV-2

insertions, is however congruent with the relative frequencies of L1, *Alu*, SVA and non-

retrotransposon cellular RNA insertions driven by L1 proteins in prior cultured cell assays

(Dewannieux et al., 2003; Hancks et al., 2011; Wei et al., 2001).

178   Our approach has several notable differences and caveats when compared to that of

Zhang *et al.*. Each study used different SARS-CoV-2 isolates, and here the multiplicity of

infection (MOI 1.0) was double that of Zhang *et al.* (MOI 0.5). The high molecular weight

DNA extraction method, ONT library preparation kit and depth and quality of sequencing

applied to HEK293T cells by Zhang *et al.* (standard isopropanol precipitation, SQK-LSK109

kit, ~21× depth, N50 ~ 11kbp) and here (Nanobind kit, SQK-LSK110 kit, ~54× depth, N50 ~

39kbp) differed. Nevertheless, the DNA extraction protocols of each study would limit

retention of extrachromosomal SARS-CoV-2 DNA potentially generated by ectopic L1

reverse transcription (Dhellin et al., 1997). The origins of the ONT reads aligned to the

SARS-CoV-2 genome reported by Zhang *et al.* are therefore unclear in our view. Zhang *et al.*

only ONT sequenced HEK293T cells transfected with an L1 expression plasmid, which

human cells would not carry *in vivo*. We did not analyse SARS-CoV-2 patient samples

although, arguably, HEK293T cells present an environment far more conducive to L1 activity

than those cells accessed *in vivo* by SARS-CoV-2 (Sungnak et al., 2020; Wiersinga et al.,

2020). Widespread cell death post-infection also reduces the probability SARS-CoV-2

integrants would persist in the body (Karki et al., 2021; Varga et al., 2020). This view aligns

with a very recent report of negligible SARS-CoV-2 DNA being detected by PCR in COVID-

19 patient nasal swabs (Briggs et al., 2021).

Finally, the incredible enrichment reported by Zhang *et al.* for putative SARS-CoV-2

insertions in exons, which this and prior studies (Flasch et al., 2019; Sultana et al., 2019)

have shown are not preferred by the L1 EN, contradicts the involvement of L1 in the events

interpreted by Zhang *et al.* as SARS-CoV-2 genomic integrants. We conclude L1 *cis*

preference strongly disfavours SARS-CoV-2 retrotransposition, making the phenomenon

mechanistically plausible but likely very rare, as for other polyadenylated non-

retrotransposon cellular RNAs (Dewannieux et al., 2003; Doucet et al., 2015; Esnault et al.,

2000; Garcia-Perez et al., 2007; Hancks et al., 2011; Kulpa and Moran, 2006; Monot et al.,

2013; Moran et al., 1996; Wei et al., 2001).

ACKNOWLEDGEMENTS

The authors thank the human subjects of this study who donated tissues to the Centre

Hépatobiliaire, Paul-Brousse Hospital. We thank S. Richardson and R. Shukla for helpful

discussions, K. Chappell and P. Young for project support, and Queensland Health for

providing the SARS-CoV-2 virus isolate QLD02. This study was funded by an Australian

Government Research Training Program (RTP) Scholarships (N.S.), an NHMRC-ARC

Dementia Research Development Fellowship (GNT1108258, G.O.B.), seed funding provided

by the Australian Infectious Disease Research Centre to establish SARS-CoV-2 research at

the University of Queensland (A.A.K), an Australian Department of Health Medical Research

Future Fund (MRFF) Novel Coronavirus Vaccine Development Grant (APP1202445-2020,

D.W.), an MRFF Investigator Grant (MRF1175457, A.D.E.), an NHMRC Investigator Grant

(GNT1173711, G.J.F.), a CSL Centenary Fellowship (G.J.F.), and the Mater Foundation.

AUTHOR CONTRIBUTIONS

N.S., J.R., G.O.B., A.A.A., P.G., F.J.S-L., P.A., N.M. and B.L. performed experiments and

analysed data. A.D.E. and G.J.F. performed bioinformatic analysis. J.F., I.W.D., A.A.K.,

D.W. and G.J.F. provided resources. G.J.F. designed the project and wrote the manuscript.

DECLARATION OF INTERESTS

The authors declare no competing interests.

REFERENCES

Ahl, V., Keller, H., Schmidt, S., and Weichenrieder, O. (2015). Retrotransposition and Crystal Structure of an Alu RNP in the Ribosome-Stalling Conformation. Mol. Cell *60*, 715–727.

Amarilla, A.A., Modhiran, N., Setoh, Y.X., Peng, N.Y.G., Sng, J.D.J., Liang, B., McMillan, C.L.D., Freney, M.E., Cheung, S.T.M., Chappell, K.J., et al. (2021). An optimized high-throughput immuno-plaque assay for SARS-CoV-2. Front. Microbiol. *12*, 625136.

Belyi, V.A., Levine, A.J., and Skalka, A.M. (2010). Unexpected inheritance: multiple integrations of ancient bornavirus and ebolavirus/marburgvirus sequences in vertebrate genomes. PLoS Pathog. *6*, e1001030.

Bill, C.A., and Summers, J. (2004). Genomic DNA double-strand breaks are targets for hepadnaviral DNA integration. Proc. Natl. Acad. Sci. U. S. A. *101*, 11135–11140.

Briggs, E., Ward, W., Rey, S., Law, D., Nelson, K., Bois, M., Ostrov, N., Lee, H.H., Laurent, J.M., and Mita, P. (2021). Assessment of potential SARS-CoV-2 virus N gene integration into human genome reveals no significant impact on RT-qPCR COVID-19 diagnostic testing (medRxiv).

Deniz, Ö., Frost, J.M., and Branco, M.R. (2019). Regulation of transposable elements by DNA modifications. Nat. Rev. Genet. *20*, 417–431.

Dewannieux, M., Esnault, C., and Heidmann, T. (2003). LINE-mediated retrotransposition of marked Alu sequences. Nat. Genet. *35*, 41–48.

Dhellin, O., Maestre, J., and Heidmann, T. (1997). Functional differences between the human LINE retrotransposon and retroviral reverse transcriptases for in vivo mRNA reverse transcription. EMBO J. *16*, 6590–6602.

Dombroski, B.A., Scott, A.F., and Kazazian, H.H., Jr (1993). Two additional potential retrotransposons isolated from a human L1 subfamily that contains an active retrotransposable element. Proc. Natl. Acad. Sci. U. S. A. *90*, 6513–6517.

Doucet, A.J., Wilusz, J.E., Miyoshi, T., Liu, Y., and Moran, J.V. (2015). A 3' Poly(A) Tract Is Required for LINE-1 Retrotransposition. Mol. Cell *60*, 728–741.

Esnault, C., Maestre, J., and Heidmann, T. (2000). Human LINE retrotransposons generate
processed pseudogenes. Nat. Genet. *24*, 363–367.

Evrony, G.D., Lee, E., Mehta, B.K., Benjamini, Y., Johnson, R.M., Cai, X., Yang, L.,
Haseley, P., Lehmann, H.S., Park, P.J., et al. (2015). Cell lineage analysis in human brain
using endogenous retroelements. Neuron *85*, 49–59.

Ewing, A.D., Smits, N., Sanchez-Luque, F.J., Faivre, J., Brennan, P.M., Richardson, S.R.,
Cheetham, S.W., and Faulkner, G.J. (2020). Nanopore Sequencing Enables Comprehensive
Transposable Element Epigenomic Profiling. Mol. Cell *80*, 915–928.

Faulkner, G.J., and Billon, V. (2018). L1 retrotransposition in the soma: a field jumping
ahead. Mob. DNA *9*, 22.

Feschotte, C., and Gilbert, C. (2012). Endogenous viruses: insights into viral evolution and
impact on host biology. Nat. Rev. Genet. *13*, 283–296.

Feusier, J., Watkins, W.S., Thomas, J., Farrell, A., Witherspoon, D.J., Baird, L., Ha, H.,
Xing, J., and Jorde, L.B. (2019). Pedigree-based estimation of human mobile element
retrotransposition rates. Genome Res. *29*, 1567–1577.

Flasch, D.A., Macia, Á., Sánchez, L., Ljungman, M., Heras, S.R., García-Pérez, J.L., Wilson,
T.E., and Moran, J.V. (2019). Genome-wide de novo L1 Retrotransposition Connects
Endonuclease Activity with Replication. Cell *177*, 837–851.

Fujimoto, A., Totoki, Y., Abe, T., Boroevich, K.A., Hosoda, F., Nguyen, H.H., Aoki, M.,
Hosono, N., Kubo, M., Miya, F., et al. (2012). Whole-genome sequencing of liver cancers
identifies etiological influences on mutation patterns and recurrent mutations in chromatin
regulators. Nat. Genet. *44*, 760–764.

Garcia-Perez, J.L., Doucet, A.J., Bucheton, A., Moran, J.V., and Gilbert, N. (2007). Distinct
mechanisms for trans-mediated mobilization of cellular RNAs by the LINE-1 reverse
transcriptase. Genome Res. *17*, 602–611.

Geuking, M.B., Weber, J., Dewannieux, M., Gorelik, E., Heidmann, T., Hengartner, H.,
Zinkernagel, R.M., and Hangartner, L. (2009). Recombination of retrotransposon and
exogenous RNA virus results in nonretroviral cDNA integration. Science *323*, 393–396.

Gilbert, N., Lutz, S., Morrish, T.A., and Moran, J.V. (2005). Multiple fates of L1
retrotransposition intermediates in cultured human cells. Mol. Cell. Biol. *25*, 7780–7795.

Hancks, D.C., Ewing, A.D., Chen, J.E., Tokunaga, K., and Kazazian, H.H., Jr (2009). Exon-
trapping mediated by the human retrotransposon SVA. Genome Res. *19*, 1983–1991.

Hancks, D.C., Goodier, J.L., Mandal, P.K., Cheung, L.E., and Kazazian, H.H., Jr (2011).
Retrotransposition of marked SVA elements by human L1s in cultured cells. Hum. Mol.
Genet. *20*, 3386–3400.

Holmes, S.E., Dombroski, B.A., Krebs, C.M., Boehm, C.D., and Kazazian, H.H., Jr (1994). A
new retrotransposable human L1 element from the LRE2 locus on chromosome 1q produces
a chimaeric insertion. Nat. Genet. *7*, 143–148.

Horie, M., Honda, T., Suzuki, Y., Kobayashi, Y., Daito, T., Oshida, T., Ikuta, K., Jern, P.,
Gojobori, T., Coffin, J.M., et al. (2010). Endogenous non-retroviral RNA virus elements in
mammalian genomes. Nature *463*, 84–87.

Hrecka, K., Hao, C., Gierszewska, M., Swanson, S.K., Kesik-Brodacka, M., Srivastava, S.,
Florens, L., Washburn, M.P., and Skowronski, J. (2011). Vpx relieves inhibition of HIV-1
infection of macrophages mediated by the SAMHD1 protein. Nature *474*, 658–661.

Jiang, Z., Jhunjhunwala, S., Liu, J., Haverty, P.M., Kennemer, M.I., Guan, Y., Lee, W.,
Carnevali, P., Stinson, J., Johnson, S., et al. (2012). The effects of hepatitis B virus
integration into the genomes of hepatocellular carcinoma patients. Genome Res. *22*, 593–601.

Jurka, J. (1997). Sequence patterns indicate an enzymatic involvement in integration of
mammalian retroposons. Proc. Natl. Acad. Sci. U. S. A. *94*, 1872–1877.

Karki, R., Sharma, B.R., Tuladhar, S., Williams, E.P., Zalduondo, L., Samir, P., Zheng, M.,
Sundaram, B., Banoth, B., Malireddi, R.K.S., et al. (2021). Synergism of TNF-α and IFN-γ
Triggers Inflammatory Cell Death, Tissue Damage, and Mortality in SARS-CoV-2 Infection
and Cytokine Shock Syndromes. Cell *184*, 149–168.

Kawasaki, J., Kojima, S., Mukai, Y., Tomonaga, K., and Horie, M. (2021). 100-My history of
bornavirus infections hidden in vertebrate genomes. Proc. Natl. Acad. Sci. U. S. A. *118*.

Kazachenka, A., and Kassiotis, G. (2021). SARS-CoV-2-Host Chimeric RNA-Sequencing
Reads Do Not Necessarily Arise From Virus Integration Into the Host DNA. Front.
Microbiol. *12*, 676693.

Kazazian, H.H., Jr, and Moran, J.V. (2017). Mobile DNA in Health and Disease. N. Engl. J.
Med. *377*, 361–370.

Kazazian, H.H., Jr, Wong, C., Youssoufian, H., Scott, A.F., Phillips, D.G., and Antonarakis,
S.E. (1988). Haemophilia A resulting from de novo insertion of L1 sequences represents a
novel mechanism for mutation in man. Nature *332*, 164–166.

Klawitter, S., Fuchs, N.V., Upton, K.R., Muñoz-Lopez, M., Shukla, R., Wang, J., Garcia-
Cañadas, M., Lopez-Ruiz, C., Gerhardt, D.J., Sebe, A., et al. (2016). Reprogramming triggers
endogenous L1 and Alu retrotransposition in human induced pluripotent stem cells. Nat.
Commun. *7*, 10286.

Klenerman, P., Hengartner, H., and Zinkernagel, R.M. (1997). A non-retroviral RNA virus
persists in DNA form. Nature *390*, 298–301.

Kubo, S., Seleme, M.D.C., Soifer, H.S., Perez, J.L.G., Moran, J.V., Kazazian, H.H., Jr, and
Kasahara, N. (2006). L1 retrotransposition in nondividing and primary human somatic cells.
Proc. Natl. Acad. Sci. U. S. A. *103*, 8036–8041.

Kulpa, D.A., and Moran, J.V. (2006). Cis-preferential LINE-1 reverse transcriptase activity
in ribonucleoprotein particles. Nat. Struct. Mol. Biol. *13*, 655–660.

Laguette, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Ségéral, E., Yatim,
A., Emiliani, S., Schwartz, O., and Benkirane, M. (2011). SAMHD1 is the dendritic- and
myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. Nature *474*, 654–657.

Lauer, G.M., and Walker, B.D. (2001). Hepatitis C virus infection. N. Engl. J. Med. *345*, 41–
52.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics *34*,
3094–3100.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
G., Durbin, R., and 1000 Genome Project Data Processing Subgroup (2009). The Sequence
Alignment/Map format and SAMtools. Bioinformatics *25*, 2078–2079.

Mason, W.S., Gill, U.S., Litwin, S., Zhou, Y., Peri, S., Pop, O., Hong, M.L.W., Naik, S.,
Quaglia, A., Bertoletti, A., et al. (2016). HBV DNA Integration and Clonal Hepatocyte
Expansion in Chronic Hepatitis B Patients Considered Immune Tolerant. Gastroenterology
151, 986–998.

Monot, C., Kuciak, M., Viollet, S., Mir, A.A., Gabus, C., Darlix, J.-L., and Cristofari, G.
(2013). The specificity and flexibility of l1 reverse transcription priming at imperfect T-
tracts. PLoS Genet. *9*, e1003499.

Moran, J.V., Holmes, S.E., Naas, T.P., DeBerardinis, R.J., Boeke, J.D., and Kazazian, H.H.,
Jr (1996). High frequency retrotransposition in cultured mammalian cells. Cell *87*, 917–927.

Moran, J.V., DeBerardinis, R.J., and Kazazian, H.H., Jr (1999). Exon shuffling by L1
retrotransposition. Science *283*, 1530–1534.

Nagaya, T., Nakamura, T., Tokino, T., Tsurimoto, T., Imai, M., Mayumi, T., Kamino, K.,
Yamamura, K., and Matsubara, K. (1987). The mode of hepatitis B virus DNA integration in
chromosomes of human hepatocellular carcinoma. Genes Dev. *1*, 773–782.

Nguyen, T.H.M., Carreira, P.E., Sanchez-Luque, F.J., Schauer, S.N., Fagg, A.C., Richardson,
S.R., Davies, C.M., Jesuadian, J.S., Kempen, M.-J.H.C., Troskie, R.-L., et al. (2018). L1
Retrotransposon Heterogeneity in Ovarian Tumor Cell Evolution. Cell Rep. *23*, 3730–3740.

Niewiadomska, A.M., Tian, C., Tan, L., Wang, T., Sarkis, P.T.N., and Yu, X.-F. (2007).
Differential inhibition of long interspersed element 1 by APOBEC3 does not correlate with
high-molecular-mass-complex formation or P-body association. J. Virol. *81*, 9577–9583.

Ostertag, E.M., and Kazazian, H.H., Jr (2001). Twin priming: a proposed mechanism for the
creation of inversions in L1 retrotransposition. Genome Res. *11*, 2059–2065.

Philippe, C., Vargas-Landin, D.B., Doucet, A.J., van Essen, D., Vera-Otarola, J., Kuciak, M.,
Corbin, A., Nigumann, P., and Cristofari, G. (2016). Activation of individual L1
retrotransposon instances is restricted to cell-type dependent permissive loci. Elife *5*, e13926.

Raiz, J., Damert, A., Chira, S., Held, U., Klawitter, S., Hamdorf, M., Löwer, J., Strätling,
W.H., Löwer, R., and Schumann, G.G. (2012). The non-autonomous retrotransposon SVA is
trans-mobilized by the human LINE-1 protein machinery. Nucleic Acids Res. *40*, 1666–1683.

Robinson, J.T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E.S., Getz, G., and
Mesirov, J.P. (2011). Integrative genomics viewer. Nat. Biotechnol. *29*, 24–26.

Rodriguez-Martin, B., Alvarez, E.G., Baez-Ortega, A., Zamora, J., Supek, F.,

Demeulemeester, J., Santamarina, M., Ju, Y.S., Temes, J., Garcia-Souto, D., et al. (2020).
Pan-cancer analysis of whole genomes identifies driver rearrangements promoted by LINE-1
retrotransposition. Nat. Genet. *52*, 306–319.

Sanchez-Luque, F.J., Kempen, M.-J.H.C., Gerdes, P., Vargas-Landin, D.B., Richardson, S.R.,
Troskie, R.-L., Jesuadian, J.S., Cheetham, S.W., Carreira, P.E., Salvador-Palomeque, C., et
al. (2019). LINE-1 Evasion of Epigenetic Repression in Humans. Mol. Cell *75*, 590–604.

Schauer, S.N., Carreira, P.E., Shukla, R., Gerhardt, D.J., Gerdes, P., Sanchez-Luque, F.J.,
Nicoli, P., Kindlova, M., Ghisletti, S., Santos, A.D., et al. (2018). L1 retrotransposition is a
common feature of mammalian hepatocarcinogenesis. Genome Res. *28*, 639–653.

Scott, E.C., Gardner, E.J., Masood, A., Chuang, N.T., Vertino, P.M., and Devine, S.E.
(2016). A hot L1 retrotransposon evades somatic repression and initiates human colorectal
cancer. Genome Res. *26*, 745–755.

Shukla, R., Upton, K.R., Muñoz-Lopez, M., Gerhardt, D.J., Fisher, M.E., Nguyen, T.,
Brennan, P.M., Baillie, J.K., Collino, A., Ghisletti, S., et al. (2013). Endogenous
retrotransposition activates oncogenic pathways in hepatocellular carcinoma. Cell *153*, 101–
111.

Siudeja, K., van den Beek, M., Riddiford, N., Boumard, B., Wurmser, A., Stefanutti, M.,
Lameiras, S., and Bardin, A.J. (2021). Unraveling the features of somatic transposition in the
Drosophila intestine. EMBO J. *40*, e106388.

Sultana, T., van Essen, D., Siol, O., Bailly-Bechet, M., Philippe, C., Zine El Aabidine, A.,
Pioger, L., Nigumann, P., Saccani, S., Andrau, J.-C., et al. (2019). The Landscape of L1
Retrotransposons in the Human Genome Is Shaped by Pre-insertion Sequence Biases and
Post-insertion Selection. Mol. Cell *74*, 555–570.

Sungnak, W., Huang, N., Bécavin, C., Berg, M., Queen, R., Litvinukova, M., Talavera-
López, C., Maatz, H., Reichart, D., Sampaziotis, F., et al. (2020). SARS-CoV-2 entry factors
are highly expressed in nasal epithelial cells together with innate immune genes. Nat. Med.
26, 681–687.

Treiber, C.D., and Waddell, S. (2017). Resolving the prevalence of somatic transposition in
Drosophila. Elife *6*, e28297.

Tu, T., Budzinska, M.A., Vondran, F.W.R., Shackel, N.A., and Urban, S. (2018). Hepatitis B
virus DNA integration occurs early in the viral life cycle in an in vitro infection model via
sodium taurocholate cotransporting polypeptide-dependent uptake of enveloped virus
particles. J. Virol. *92*, e02007–e02017.

Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B.C., Remm, M., and Rozen,
S.G. (2012). Primer3—new capabilities and interfaces. Nucleic Acids Res. *40*, e115–e115.

Varga, Z., Flammer, A.J., Steiger, P., Haberecker, M., Andermatt, R., Zinkernagel, A.S.,
Mehra, M.R., Schuepbach, R.A., Ruschitzka, F., and Moch, H. (2020). Endothelial cell
infection and endotheliitis in COVID-19. Lancet *395*, 1417–1418.

V'kovski, P., Kratzel, A., Steiner, S., Stalder, H., and Thiel, V. (2020). Coronavirus biology
and replication: implications for SARS-CoV-2. Nat. Rev. Microbiol. *19*, 155–170.

Wei, W., Gilbert, N., Ooi, S.L., Lawler, J.F., Ostertag, E.M., Kazazian, H.H., Boeke, J.D.,
and Moran, J.V. (2001). Human L1 retrotransposition: cispreference versus trans
complementation. Mol. Cell. Biol. *21*, 1429–1439.

Wiersinga, W.J., Rhodes, A., Cheng, A.C., Peacock, S.J., and Prescott, H.C. (2020).
Pathophysiology, Transmission, Diagnosis, and Treatment of Coronavirus Disease 2019
(COVID-19): A Review. JAMA *324*, 782–793.

Wu, F., Zhao, S., Yu, B., Chen, Y.-M., Wang, W., Song, Z.-G., Hu, Y., Tao, Z.-W., Tian, J.-
H., Pei, Y.-Y., et al. (2020). A new coronavirus associated with human respiratory disease in
China. Nature *579*, 265–269.

Yan, B., Chakravorty, S., Mirabelli, C., Wang, L., Trujillo-Ochoa, J.L., Chauss, D., Kumar,
D., Lionakis, M.S., Olson, M.R., Wobus, C.E., et al. (2021). Host-Virus Chimeric Events in
SARS-CoV-2-Infected Cells Are Infrequent and Artifactual. J. Virol. *95*, e0029421.

Zhang, L., Richards, A., Barrasa, M.I., Hughes, S.H., Young, R.A., and Jaenisch, R. (2021).
Reverse-transcribed SARS-CoV-2 RNA can integrate into the genome of cultured human
cells and can be expressed in patient-derived tissues. Proc. Natl. Acad. Sci. U. S. A. *118*,
e2105968118.

Zhao, K., Du, J., Han, X., Goodier, J.L., Li, P., Zhou, X., Wei, W., Evans, S.L., Li, L., Zhang,
W., et al. (2013). Modulation of LINE-1 and Alu/SVA retrotransposition by Aicardi-
Goutières syndrome-related SAMHD1. Cell Rep. *4*, 1108–1115.

FIGURE LEGENDS

Figure 1. Key potential SARS-CoV-2 insertions reported by Zhang *et al.*.

(**A**) A cartoon summarising the features of a putative SARS-CoV-2 integrant on chromosome

X. Numerals underneath the SARS-CoV-2 sequence represent positions relative to the QLD02

virus isolate. Potential TSDs are shown as red triangles, and motifs resembling potential pre-

integration L1 EN recognition sites are highlighted, with question marks in labels intended to

flag uncertain L1 involvement. No 3′ polyA tract was found. Homologous regions at sequence

junctions are marked. One spanning ONT read is positioned underneath the cartoon and its

identifier is displayed.

(**B**) As for (A), except showing an ONT read spanning two SARS-CoV-2 insertions, on

chromosome 22 and chromosome 1. The alignments to chromosome 22 were flagged as

supplementary by the minimap2 aligner. 3′ polyA tracts are represented as green rectangles.

Note: these chromosome 22 and chromosome X instances are the key examples reported by

Zhang *et al.* in support of SARS-CoV-2 genomic integration. Neither example has a complete

set of retrotransposition hallmarks (TSD, 3′ polyA tract, L1 EN motif) *and* the support of a

uniquely aligned ONT read.

Figure 2. Detection of endogenous L1-mediated retrotransposition in human cells.

(**A**) Experimental design. HEK293T cells were divided into two populations (cultivars), which were then either SARS-CoV-2 infected or mock infected. DNA was extracted from each cultivar, as well as from hepatocellular carcinoma patient samples, and subjected to ONT sequencing. ONT reads were used to call non-reference L1 and virus insertions with TLDR, which also resolves TSDs and other retrotransposition hallmarks. TSDs: red triangles; polyA tract: green rectangle; ONT read: blue rectangle. Some illustrations are adapted from a previous study (Ewing et al., 2020).

(**B**) TSD size distribution for non-reference L1 insertions, as annotated by TLDR, as well as cultivar-specific L1 insertions found only in either our HEK293T cells infected with SARS-CoV-2 or our mock infected cells. Pie charts indicate the percentages of exonic, intronic and intergenic insertions, annotated by RefSeq coordinates.

(**C**) As for (B), except showing data for *Alu* insertions.

(**D**) Detailed characterisation of an L1 insertion detected in SARS-CoV-2 infected HEK293T cells by a single spanning ONT read aligned to chromosome 14. Nucleotides highlighted in red correspond to the integration site TSD. Underlined nucleotides correspond to the L1 EN motif. The cartoon indicates a full-length L1HS insertion flanked by TSDs (red triangles), and a 3′ polyA tract (green), with the underneath numeral representing the 5′ end position relative to the mobile L1HS sequence L1.3 (Dombroski et al., 1993). The relevant spanning ONT read, with identifier, is also positioned underneath the cartoon. Symbols (α, β, δ, γ) represent the approximate position of primers used for empty/filled site and L1-genome junction PCR validation reactions. These are displayed in gel images if successful. Ladder band sizes are as indicated, NTC; non-template control. Red triangles indicate L1 amplicon expected sizes (empty triangle: no product; filled triangle: capillary sequenced on-target product). Blue triangles indicate expected empty site sizes.

(**E**) As for (D), except for a 5′ inverted/deleted L1HS located on chromosome 18.

Please see Figures S1 and S2, and Tables S1 and S2 for further information.

Figure 3. ONT reads occasionally align to viral genome sequences.

(**A**) Percentages of total ONT sequence alignable to L1HS (left), SARS-CoV-2 (middle) and HBV (right) isolate genomes. Read counts for SARS-CoV-2 and HBV are provided above histogram columns. No reads were aligned to the HCV isolate genomes. HEK293T data were generated here (SARS-CoV-2, mock) or by Zhang *et al.*. HCC tumour/non-tumour liver pairs

were sequenced here (HCC32; confirmed HBV-positive) or previously (Ewing et al., 2020)

(HCC33; HCV-positive). Normal liver ONT sequencing from our prior work (Ewing et al.,

2020) was included as an additional control.

(B) A HBV insertion detected in non-tumour liver. In this example, an ONT read from the

non-tumour liver of HCC32 spanned the 3′ junction of a HBV integrant located on

chromosome 2. Of the HBV isolate genomes considered here, this read aligned best to a

representative of genotype B (Genbank accession AB602818). The HBV sequence was

rearranged consistent with its linearisation prior to integration (Fujimoto et al., 2012; Jiang et

al., 2012; Nagaya et al., 1987). Numerals indicate positions relative to AB602818. Symbols

(β, δ) represent the approximate position of primers used to PCR validate the HBV insertion.

The gel image at right shows the 3′ junction PCR results. Ladder band sizes are as indicated.

The red filled triangle indicates an on-target product confirmed by capillary sequencing.

Repeated attempts to PCR amplify the 5′ junction of the HBV integrant did not return an on-

target product, perhaps due to a genomic deletion at the insertion site.

(C) Percentages of exonic (black), intronic (grey) and intergenic (blue) genomic alignment

breakpoints for ONT reads also aligned to the SARS-CoV-2 genome, and also for the non-

reference L1-mediated insertions reported here. Genomic features were annotated according

to RefSeq coordinates.

Please see Tables S1 and S2 for further information.

STAR METHODS

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Geoffrey J. Faulkner (faulknergj@gmail.com).

Materials availability

This study did not generate new unique reagents.

Data and code availability

Oxford Nanopore Technologies sequencing data generated by this study were deposited in the

European Nucleotide Archive (ENA) under project PRJEB44816. TLDR and instructions for

its use and application are available at https://github.com/adamewing/TLDR.

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Liver tumour and non-tumour tissue were previously obtained from a HBV-positive patient (HCC32, male, 73yrs) who underwent surgical resection at the Centre Hepatobiliaire, Paul-Brousse Hospital, and made available for research purposes with approval from the French Institute of Medical Research and Health (Reference: 11-047). Further ethics approvals were provided by the Mater Health Services Human Research Ethics Committee (Reference: HREC-15-MHS-52) and the University of Queensland Medical Research Review Committee (Reference: 2014000221). HEK293T and Vero E6 cells were obtained from the American Type Culture Collection (ATCC).

METHOD DETAILS

SARS-CoV-2 infection of HEK293T cells

HEK293T cells and African green monkey kidney cells (Vero E6) were maintained in standard Dulbecco's Modified Eagle Medium (DMEM). Culture media were supplemented with sodium pyruvate (11mg/L), penicillin (100U/mL), streptomycin (100μg/mL) (P/S) and 10% foetal calf serum (FCS) (Bovogen, USA). Cells were maintained at 37 °C with 5% CO_2.

An early Australian SARS-CoV-2 isolate (hCoV-19/Australia/QLD02/2020; GISAID Accession EPI_ISL_407896) was sampled from patient nasopharyngeal aspirates by Queensland Health Forensic and Scientific Services and used to inoculate Vero E6 African green monkey kidney cells (passage 2). A viral stock (passage 3) was then generated on Vero E6 cells and stored at -80°C. Viral titration was determined by immuno-plaque assay (iPA), as previously described (Amarilla et al., 2021). To verify viral replication in HEK293T cells, a growth kinetic was assessed using a multiplicity of infection (MOI) of 0.01, 0.1 or 1.0, and showed efficient SARS-CoV-2 replication (**Figure S1**).

HEK293T viral infection was undertaken as follows: 3×10^6 HEK293T cells were seeded onto 6-well plates pre-coated with polylysine one day before infection. Cells were infected at MOI of 1 in 200μL of DMEM (2% FCS and P/S) and incubated for 30min at 37°C. Plates were rocked every 5min to ensure the monolayer remained covered with inoculum. The inoculum was then removed, and the monolayer washed five times with 1mL of additive-free DMEM. Finally, cells were maintained with 3mL of DMEM (supplemented with 2% foetal bovine serum and P/S) and incubated at 37°C with 5% CO_2. Cell supernatant was harvested 0, 1, 2 and 3 days post-infection. The mock infected control differed only in that virus was not added to the inoculum media.

Genomic DNA was extracted from mock and SARS-CoV-2 infected (MOI 1.0)

HEK293T cells sampled 2 days post-infection, using a Nanobind CBB Big DNA Kit

(Circulomics) following the manufacturer's instructions for high molecular weight (HMW)

DNA extraction. DNA was eluted in elution buffer (10mM Tris-Cl, pH 8.5) and concentration

measured by Qubit dsDNA High-Sensitivity Assay Kit on a Qubit Fluorometer (Life

Technologies).

Hepatocellular carcinoma sample processing

DNA was extracted from the HCC32 tissues in our earlier study (Shukla et al., 2013) with a

DNeasy Blood and Tissue Kit (QIAGEN, Germany) and stored at -80ºC. To enrich for HMW

DNA, 4.5μg of DNA from the patient HCC32 tumour and non-tumour liver samples was

diluted to 75ng/μL in a 1.5mL Eppendorf DNA LoBind tube and processed with a Short Read

Eliminator XS Kit (Circulomics) following the manufacturer's instructions.

ONT sequencing

DNA libraries were prepared at the Kinghorn Centre for Clinical Genomics (KCCG) using 3-

4μg HMW input DNA, without shearing, and a SQK-LSK110 ligation sequencing kit. 350-

500ng of each prepared library was sequenced separately on one PromethION (Oxford

Nanopore Technologies) flow cell (FLO-PRO002, R9.4.1 chemistry) (**Table S1**). SARS-CoV-

2 infected HEK293T DNA was sequenced on two flow cells. Flow cells were washed (nuclease

flush) and reloaded at 24hr and 48hr with 350-500ng of additional library to maximise output.

Bases were called with guppy 4.0.11 (Oxford Nanopore Technologies).

ONT bioinformatic analyses

To call non-reference insertions with TLDR (Ewing et al., 2020), ONT reads generated here,

by Zhang *et al*., and by our previous ONT study of human tissues (Ewing et al., 2020) (**Table**

S1) were aligned to the human reference genome build hg38 using minimap2 (Li, 2018)

version 2.17 (index parameter: -x map-ont; alignment parameters: -ax map-ont -L -t 32) and

SAMtools (Li et al., 2009) version 1.12. BAM files were then processed as a group with

TLDR (Ewing et al., 2020) version 1.1 (parameters -e virus.fa -p 128 -m 1 --max_te_len

40000 --max_cluster_size 100 --min_te_len 100 --keep_pickles -n

nonref.collection.hg38.chr.bed.gz). The file virus.fa was composed of: representative HBV

and HCV isolate genomes (**Table S1**), the SARS-CoV-2 isolate used here (GISAID

Accession EPI_ISL_407896), the L1HS sequence L1.3 (Dombroski et al., 1993) (Genbank

Accession L19088), several *Alu* and SVA subfamily consensus sequences, and a consensus

sequence for human endogenous retrovirus K (HERVK), the youngest human long terminal

repeat (LTR) retrotransposon family. The file nonref.collection.hg38.chr.bed.gz is a

collection of known non-reference retrotransposon insertions available from

github.com/adamewing/tldr/. The TLDR output table was further processed to remove calls

not passing all TLDR filters, representing homopolymer insertions, where MedianMapQ < 50

or family = "NA" or remappable = "FALSE" or UnmapCover < 0.75 or LengthIns < 100 or

EndTE-StartTE < 100 or strand = "None" or SpanReads < 1. As 3′ truncation is rarely

encountered for L1-mediated insertions, calls where EndTE was more than 10bp less than the

consensus length were filtered, as were *Alu* insertions 5′ truncated by more than 1bp. The

filtered TLDR output table is provided as **Table S2**. Insertions detected in only our mock or

SARS-CoV-2 infected HEK293T datasets, but not in both experiments, and not matching a

known non-reference insertion, were designated as putative cultivar-specific insertions

(**Table S2**). Many if not most of these insertions were likely to have occurred in cell culture

prior to the cultivars being separated.

To identify L1HS and viral sequences, we directly aligned all reads to the virus.fa file

with minimap2 (index parameter: -x map-ont; alignment parameters: -ax map-ont -L -t 32).

Reads containing alignments of ≥100bp to a sequence present in virus.fa were counted with

SAMtools idxstats. Alignments to HBV, HCV or SARS-CoV-2 were excluded if they

overlapped by ≥10bp with a genomic alignment of ≥100bp. Read alignments were visualised

with SAMtools view and the Integrative Genomics Viewer (Robinson et al., 2011) version

2.8.6.

PCR validation

We used Primer3 (Untergasser et al., 2012) to design PCR primers for 6 L1 insertions found

by a single spanning ONT read, using the reference genome and L1HS sequences as inputs

(**Table S2**). These validation experiments were conducted in three phases. Firstly, we

performed an "empty/filled site" PCR using primers positioned on either side of the L1,

where the filled site is the L1 allele, and the empty site is the remaining allele(s). Each

empty/filled reaction was performed using a DNA Engine Tetrad 2 Thermal Cycler (Bio-

Rad) and Expand Long Range Enzyme Mix, with 1X Expand Long Range Buffer with

MgCl$_2$, 50pmol of each primer, 0.5mM dNTPs, 5% DMSO, 100ng of template DNA and

1.75U of enzyme, in a 25μL final volume. PCR cycling conditions were as follows: (92˚C,

3min)×1; (92˚C, 30sec; 54-57˚C, 30sec; 68˚C, 7min)×10; (92˚C, 30sec; 52-55˚C, 30sec;

68˚C, 7min + 20sec/cycle)×30; (68˚C, 10min; 4˚C, hold)×1. Amplicons were visualised on a

1% agarose gel stained with SYBR SAFE (Invitrogen). GeneRuler™ 1kb plus (Thermo

Scientific) was used as the ladder. Secondly, we combined each empty/filled primer with a

primer positioned within the L1 sequence, to amplify the 5′ and 3′ L1-genome junctions.

These reactions were undertaken on a DNA Engine Tetrad 2 Thermal Cycler (Bio-Rad), with

MyTaq HS DNA polymerase, 1X MyTaq Reaction Buffer, 10pmol of each primer, 10ng of

template DNA, and 2.5U of enzyme, in a 25µL final volume. PCR cycling conditions were as

follows: (95˚C, 1min)×1; (95˚C, 15sec; 53-55˚C, 15sec; 72˚C, 15sec)×35; (72˚C, 5min; 4˚C,

hold)×1. Amplicons were visualised on a 1.5% agarose gel stained with SYBR SAFE

(Invitrogen). Thirdly, we repeated the 5′ L1-genome junction-specific PCR using 200ng

template DNA. All PCRs were performed with non-template control, as well as DNA

extracted from the same HEK293T cells (SARS-CoV-2 and mock) subjected to genomic

analysis. Notably, L1 insertions that did not amplify in either cultivar were still likely to be

genuine events as they carried all of the relevant sequence hallmarks of L1-mediated

retrotransposition.

PCR primers for the HBV insertion 3′ junction (**Figure 3B** and **Table S2**) were

designed with Primer3 using the reference genome and closest match HBV sequence

(Genbank accession AB602818) as inputs. PCR amplification and capillary sequencing was

conducted as per the L1 insertions, except using Expand Long Range polymerase (Roche)

with 1X Expand Long Range buffer with $MgCl_2$, 10pmol of each primer, 100ng of template

DNA, 500µM of PCR Nucleotide Mix, and 3.5U of enzyme, in a 25µL final volume. PCR

cycling conditions were as follows: (92˚C, 2min)×1; (92˚C, 15sec; 65˚C, 15sec; 68˚C,

7:30min)×10; (92˚C, 15sec; 65˚C, 15sec; 68˚C, 7min+ 20sec per cycle)×35 (68˚C, 10min;

4˚C, hold)×1. Amplicons were visualized on a 1.2% agarose gel.

Amplicons in each experiment were visualised using a GelDoc (Bio-Rad) and, if of

the correct size, gel extracted using a Qiagen MinElute Gel Extraction Kit and capillary

sequenced by the Australian Genomics Research Facility (Brisbane).

QUANTIFICATION AND STATISTICAL ANALYSIS

Error bars and replicate values are defined in figure legends, where appropriate. No statistical

tests for significance were conducted.

KEY RESOURCES TABLE

REAGENT OR RESOURCE	SOURCE	IDENTIFIER
Bacterial and Virus Strains		
SARS-CoV-2 isolate hCoV-19/Australia/QLD02/2020	Queensland Health Forensic and Scientific Services	EPI_ISL_407896
Biological Samples		
Snap frozen hepatocellular carcinoma and matched non-tumour liver tissue	Centre Hépatobiliaire, Paul-Brousse Hospital	HCC32
Chemicals, Peptides, and Recombinant Proteins		
SYBR Safe DNA Gel Stain	Invitrogen	S33102
Agarose	Bioline	BIO-41026
Carboxymethylcellulose	Sigma-Aldrich	C4888
KPL Milk Diluent/Blocking Solution Concentrate	SeraCare	5140-0011
IRDye 800CW Goat anti-Mouse IgG Antibody	LI-COR	926-32210
Anti-SARS-CoV-2-spike antibody	Amarilla et al., 2021	CR3022
Critical Commercial Assays		
Ligation Sequencing Kit	Oxford Nanopore Technologies	SQK-LSK110
Qubit dsDNA HS Assay Kit	Invitrogen	Q32851
Nanobind CBB Big DNA Kit	Circulomics	NB-900-001-01
Short Read Eliminator XS Kit	Circulomics	SS-100-121-01
Expand Long Template PCR System	Merck	11681834001
MyTaq DNA Polymerase	Bioline	BIO-21105
Deposited Data		
Nanopore WGS of mock and SARS-CoV-2 infected HEK293T cells, and HBV infected samples from liver cancer patient HCC32	This paper	PRJEB44816
Nanopore WGS of SARS-CoV-2 infected HEK293T cells overexpressing L1	Zhang et al., 2021	PRJNA721333
Nanopore WGS of normal liver and HCV infected samples from liver cancer patient HCC33	Ewing et al., 2020	PRJNA629858
Experimental Models: Cell Lines		
Human embryonic kidney 293T (HEK293T) cells	ATCC	CRL-1568
Cercopithecus aethiops Vero E6 cells	ATCC	CRL-3216
Oligonucleotides		
Oligo sequences are shown in Table S2.	Integrated DNA Technologies	N/A
Software and Algorithms		
TLDR	https://github.com/adamewing/tldr	Ewing et al., 2020
Minimap2	https://github.com/lh3/minimap2	Li et al., 2018
SAMtools	https://github.com/samtools/	Li et al., 2009
IGV	https://software.broadinstitute.org/software/igv/	Robinson et al., 2011
Primer3	https://bioinfo.ut.ee/primer3-0.4.0/	Untergasser et al., 2012

Figure S1. SARS-CoV-2 is replication competent in HEK293T cells, related to Figure 2. HEK293T cells were infected with SARS-CoV-2 isolate QLD02 at an MOI of 0.01, 0.1 and 1.0. Inoculum was removed after infection and cells were washed before the addition of growth media. Supernatant was collected at the indicated time points and viral titres were quantified as focus-forming units (FFU) per mL by immuno-plaque assay (iPA) (Amarilla et al., 2021) with a limit of detection (LOD) as indicated. Data are represented as the mean ± standard deviation of three replicates.

Figure S2. Additional L1HS insertions detected by ONT sequencing in HEK293T cells, related to Figure 2.
(A) A near full-length L1. **(B)** A 5′ truncated L1. **(C)** A 5′ inverted/deleted L1 carrying a 3′ transduction (purple rectangle) traced to a non-reference source L1. **(D)** A 5′ inverted/deleted L1. **(E)** Integrative Genomics Viewer (Robinson et al., 2011) visualisation of read alignments spanning the L1 integration site displayed in Figure 2D. The L1 is coloured purple. Note: panels (A-D) show the genomic coordinates of an L1 insertion, as well as the sequence at the insertion site. Nucleotides highlighted in red correspond to the integration site TSD. Underlined nucleotides correspond to the L1 EN motif. Cartoons summarise the features of each L1, with the underneath numerals representing the 5′ end position relative to the mobile L1HS sequence L1.3 (Dombroski et al., 1993), TSDs shown as red triangles, and 3′ polyA tracts coloured as green rectangles. One spanning ONT read with its identifier is positioned underneath each cartoon. Symbols (α, β, δ, γ) represent the approximate position of primers used for empty/filled and L1-genome junction PCR validation reactions. The results of the L1-genome 3′ junction PCR are shown for panel (A). Ladder band sizes are as indicated, NTC; non-template control. The red filled triangle indicates an on-target product confirmed by capillary sequencing. No on-target products were observed for the corresponding 5′ junction PCR or the examples shown in panels (B-D).