Title:

Seasonal variations of ^{137}Cs concentration in freshwater charr through uptake and

metabolism in 1–2 years after the Fukushima accident

Authors:

Kengo Okada[1], Masaru Sakai[2,†], Takashi Gomi[1], Aimu Iwamoto[1], Junjiro N. Negishi[3],

Masanori Nunokawa[4]

Affiliations:

[1] Graduate School of Agriculture, Tokyo University of Agriculture and Technology,

Tokyo, Japan

[2] Fukushima Regional Collaborative Research Center, National Institute for

Environmental Studies, Japan, Fukushima, 963-7700, Japan

[3] Faculty of Environmental Earth Science, Hokkaido University, Hokkaido, Japan

[4] Civil Engineering Research Institute for Cold Region, Hokkaido, Japan

† ORCID (0000-0001-5361-0978)

Correspondence

Masaru Sakai, Fukushima Regional Collaborative Research Center, National Institute

for Environmental Studies, Japan, 10-2 Fukasaku, Miharu, Tamura District, Fukushima,

963-7700, Japan

Email: sakai.masaru@nies.go.jp

Abstract

Understanding the factors influencing ^{137}Cs concentrations in freshwater salmon is

crucial for reviving inland fisheries in polluted regions. We studied seasonal variations

of ^{137}Cs concentration in charr (*Salvelinus leucomaenis*) through uptake and metabolism

in forested headwaters at highly (HCS) and moderately contaminated sites (MCS).

Charr consumed both terrestrial and aquatic animals, and terrestrial prey was predated

more in summer at both sites. The ^{137}Cs concentrations in litter, which is a dominant

basal food resource of both forest and stream ecosystems, differed between forest and

stream due to ^{137}Cs leaching effect on litter submerged in streams. The difference was

greater at HCS than at MCS. The ^{137}Cs concentration in terrestrial preys was also much

higher than aquatic preys at HCS compared to MCS. The estimated prey ^{137}Cs

concentration at HCS peaked in summer when terrestrial preys are most available,

whereas it remained relatively constant at MCS because of the small difference of ^{137}Cs

concentrations in between terrestrial and aquatic preys. The specific metabolic rate of

charr was commonly changed with stream water temperature, greatest in summer and

lowest in winter at both sites. Because both prey ^{137}Cs concentrations and specific

metabolic rates peaked in summer, the combination of uptake and metabolism at HCS

largely negated seasonal ^{137}Cs fluctuations in charr, whereas specific metabolic rate

fluctuations could be the major determinant of charr ^{137}Cs concentrations at MCS. Our

results suggested the importance of balance between ^{137}Cs uptake and metabolism in the

management of ^{137}Cs in salmonids inhabiting forested headwaters.

KEYWORDS

freshwater fish; leaching; radiocesium; resource subsidy; riparian forest

1 | INTRODUCTION

The Fukushima nuclear accident caused in March 2011 have resulted a massive contamination of both terrestrial and aquatic environments around Fukushima Prefecture (MEXT, 2012). Among the emitted radionuclides, ^{137}Cs is one of the most dominant ones, and its long half-life (30 years) prolongs environmental radiation contamination. The Japanese government has carried out decontamination practices primarily in spheres of human livelihoods (e.g., residential and arable lands) whereas forested land, which is the most dominant landscape of the contaminated region, is abandoned due to the vast area (Onda et al., 2020). Thus, ^{137}Cs is still deposited and circulating within the forest landscapes trough biological activities, and parts of edible plants and animals exceed the food standards of 100 Bq/kg wet weight even now (MAFF, 2021).

Forested headwater streams harbor freshwater salmonids including *Salvelinus* and *Oncorhynchus* species. Although these species are popular for local freshwater fisheries, the ^{137}Cs contamination has caused serious ^{137}Cs accumulations in the fishes due to the heavy burden of bioavailable ^{137}Cs provided by surrounding forests (Wada et al., 2019). Freshwater fishes take up cations, including radiocesium, largely through feeding and have lower cation excretion rates during osmoregulation than marine fishes (Wada et al., 2019). Thus, the composition of food items and their radiocesium concentrations are important determinants of radiocesium concentrations in headwater salmonids (Haque et al., 2017).

Headwater salmonids consume both terrestrial and aquatic prey, depending on seasonal availability (Nakano et al., 1999a; Nakano & Murakami, 2001). Terrestrial and aquatic ecosystems in forested headwaters generally form typical detrital food webs

originating from plant litter (Wallace et al., 1997; Gomi et al., 2002; Sakai et al., 2016a).

Thus, the unique food webs may promote ^{137}Cs transfers from contaminated litter to

charr populations through potential detritivorous preys (Negishi et al., 2018).

Previous studies found that ^{137}Cs attached to litter greatly leach after falling

into streams (Sakai et al., 2015; 2016b; Gomi et al., 2018). The efficiency of the ^{137}Cs

leaching is significantly affected by initial ^{137}Cs concentration in litter, and relative

difference of ^{137}Cs concentration in litter between terrestrial and aquatic environments is

larger in more contaminated sites (Sakai et al., 2016b). Moreover, Sakai et al. (2016a)

indicated that the concentration difference can be reflected in Cs levels measured in

animals. The suite of these studies implies that terrestrial prey for salmonids may be

more contaminated compared to aquatic prey in headwaters which suffered heavier

initial ^{137}Cs fallout.

In addition to contaminant uptake from food, the whole-body burden of

contaminants in fish is influenced by their metabolism (Rowan et al., 1997; Peles et al.,

2000; Hessen et al., 2002; Doi et al., 2012). For example, the metabolic rate of bluegills

(*Lepomis macrochirus*) is negatively correlated with ^{137}Cs concentration (Kolehmainen,

1972). Variations in radiocesium levels in mosquitofish (*Gambusia affinis*) also

suggested that their metabolic rate is among primary factors determining their ^{137}Cs

concentrations (Holloman et al., 1997). The results of the previous studies imply that

^{137}Cs accumulation in salmonids in forested headwaters needs to be examined by

considering both processes involved in uptake and metabolism.

Because metabolisms of fish become more active in hotter season (Brown,

2004), ^{137}Cs elimination through metabolism will be higher in summer and lower in

winter, independent of initial ^{137}Cs fallout volume. Meanwhile, terrestrial preys for

charrs are more available in summer than in winter (Nakano et al., 1999a; Baxter et al.,

2005). Therefore, overall ^{137}Cs concentration in preys may peak in summer in heavily

contaminated sites where effective ^{137}Cs leaching from litter induces apparent difference

of ^{137}Cs concentrations in between terrestrial and aquatic preys. In contrast, overall

^{137}Cs concentration in preys may not show clear seasonality in less contaminated sites

because concentration difference between terrestrial and aquatic litter is small (Sakai et

al., 2016b). These hypotheses suggest that ^{137}Cs leaching efficiency of aquatic litter,

which varies with degree of initial ^{137}Cs fallout, influences seasonality of ^{137}Cs uptake

and consequently ^{137}Cs concentrations in charr, as a result of seasonal changes in

balance between ^{137}Cs uptake and elimination.

Although numerous studies have reported ^{137}Cs accumulation in fish, a paucity

of knowledge remains over variations in ^{137}Cs accumulation among populations by

season. Thus, the main objective of this study was to examine how different initial

fallout volumes influence seasonal changes in ^{137}Cs concentrations in *S. leucomaenis*

based on data of thoroughly measured temporal dynamics in uptake and metabolism.

Understanding these mechanisms would provide useful information for the management

of ^{137}Cs accumulation in freshwater salmonids.

2 | MATERIALS AND METHODS

2.1 | Study site

This study was conducted in headwater streams and surrounding riparian zones in

Fukushima and Gunma Prefectures, Japan. The Fukushima site was located

approximately 45 km west of the Fukushima Daiichi nuclear power plants (FDNPPs)

(37°36'N, 140°37'E), and the Gunma site was located 180 km southwest of the

FDNPPs (36°33'N, 139°21'E). Based on airborne observations conducted in June 2012,

^{137}Cs deposited on the ground surface was 100–300 kBq/m^2 at the Fukushima site and

30–60 kBq/m^2 at the Gunma site (MEXT, 2012). The Fukushima and Gunma sites have

drainage basins of 170 and 93 ha, respectively. Based on AMeDAS automated weather

station data collected nearby, the Fukushima site had mean annual precipitation and air

temperature of 1248.4 mm and 10.9°C. Based on another nearby weather station, the

Gunma site had mean annual precipitation and air temperature of 1323.3 mm and

14.5°C. The channel reaches had typical step-pool morphologies (Montgomery &

Buffington, 1997). The riparian zones of both study sites were covered with evergreen

conifer (*Cryptomeria japonica*) plantation that is a dominant forest type of Japan. We

selected 50-m study areas along the stream channels and 20-m-wide areas in the riparian

zones on both sides of the channels. The wetted widths of the study areas at both sites

were similar (1–5 m). *S. leucomaenis* was the only fish species inhabiting in both study

streams, with densities at the Fukushima and Gunma sites of 0.19 and 0.28

individuals/m^2, respectively, estimated based on the three-pass removal method (Inoue

et al., 2013).

2.2 | Field investigations

We sampled charrs four times, in August and November 2012, and February and May in

2013, representing summer, autumn, winter, and spring, respectively. Charrs were

collected using a backpack electrofisher (LR-24; Smith-Root, Inc., Vancouver, WA,

USA), measured their wet weight and standard body length, and collected stomach

content samples from each fish with stomach pumps. The samples were collected in the

early morning and at dusk, when prey was potentially most available (Nakano et al.,

1999b). The stomach content samples were immediately preserved in 99.5% ethanol. To

avoid local extinctions of the charr populations, we only sacrificed randomly selected

3–8 individuals at both sites and from all seasons for measurements of ^{137}Cs

concentrations in charr (20.5% and 11.5% of total catch at the Fukushima and Gunma

sites, respectively, Figure 1). The charr samples were stored at -20°C prior to laboratory

analyses.

To determine potential aquatic and terrestrial prey, we sampled stream channels

with driftnets (25 × 25 cm; mesh size, 250 μm) and riparian zones with pan traps (1.05

× 0.60 × 0.21 m). Five driftnets were placed longitudinally in each stream at 10-m

intervals for 1 h in the early morning and at dusk in each season. The volume of water

flowing through the driftnets was estimated from the water depth and current velocity

measured using an electromagnetic current meter (VE20; Kenek Co., Tokyo, Japan) at

the openings of the driftnets. Five pan traps filled with several centimeters of water and

a few drops of surfactant were placed along the study reaches at 10-m intervals for 7

days. During the sampling periods in autumn and winter, antifreeze liquid was added to

the pan traps to prevent the water from freezing. The samples obtained from the

driftnets and pan traps were preserved in 99.5% ethanol immediately after collection.

Measuring ^{137}Cs concentrations in individual invertebrates such as insects is

typically difficult due to their small individual body mass. Therefore, it was necessary to

collect many individuals and pool samples by taxon to provide sufficient sample

volumes for the radionuclide measurements ($\gtrsim$ several grams in dry weight) besides the

above-mentioned investigations using stomach pumps, driftnets and pan traps. Benthic

macroinvertebrates representing aquatic prey were collected from the study sites using

D-frame nets. The collected macroinvertebrates were handled with tweezers and stored

in glass vials grouped by taxon visually identified to the lowest possible taxonomic rank.

The all collections were done after fish investigations to avoid anthropogenic

disturbance which may affect foraging of charr. Invertebrates inhabiting the forest floor

representing terrestrial prey were collected with 14 baited traps (diameter, 8 cm; depth,

11 cm) located 5–10 m from the sides of the streams. We filled the traps with an

attractant consisting of a 1:1 admixture of beer and lactic acid beverage (Sakai et al.,

2016a) and left them on the forest floor for 36 h. The collected samples were gently

washed with distilled water, visually identified to the lowest possible taxonomic rank,

and stored in plastic bags. In addition, spiders, frogs, and crabs were sampled by hand

within the study riparian zones during each sampling period. All the organism samples

were stored at -20°C prior to laboratory analyses.

To estimate the ^{137}Cs concentrations of the basal food resource of the streams

and riparian zones, we collected Japanese cedar litter deposited on the forest floors and

in the streams of the study areas. Previous carbon and nitrogen stable isotope analyses

had identified that terrestrial and aquatic food webs in Japanese cedar plantations

originate mainly from Japanese cedar litter because of its persistence and abundance

(Sakai et al., 2016a; 2016c); therefore, we assumed that the basal food resource of our

study ecosystems was Japanese cedar litter. To avoid variations in ^{137}Cs concentration

due to decomposition stage, we collected intact litter both in the riparian forests and

streams. Stream water temperature was monitored in both stream channels every 10 min

using temperature loggers (TidbiT V2; Onset Computer Co., Bourne, MA, USA) to

calculate specific metabolic rates of charr (see 2.5).

2.3 | Laboratory analysis

Muscle tissues of the charr samples were extracted and dried at 60°C for 2 days. The

terrestrial and aquatic prey samples for radioactivity measurements were dried at 60°C

for more than 2 days. Japanese cedar litter was dried in a similar manner. All dried

samples were ground into powder using either an agate mortar and pestle or electrical

mill (Osaka Chemical Co., Ltd., FM-1). The samples were packed into 100-mL

polystyrene containers and sealed until the radioactivity measurements.

Prey sampled from stomach pump, pan traps, and driftnets were identified to

the order (excepting terrestrial Oligochaeta and Diplopoda) and separated into terrestrial

or aquatic organisms. Then, we considered adult aquatic insects to be aquatic animals

due to the dominance of aquatic environments in their life history (Merritt et al., 2008),

and frog species were included as terrestrial animals because they mainly predate

terrestrial invertebrates (Maeda & Matsui, 1999). The samples were dried at 60°C for

more than 2 days, and the dry weights were measured within 0.01 mg. From this, we

estimated the total, terrestrial, and aquatic prey biomass of charr. Prey availability was

expressed as biomass per volume of water flow (mg/m^3) for the driftnet samples,

biomass per area per day ($mg/m^2/day$) for the pan trap samples, and biomass per

individual (mg/individual) for the stomach pump samples.

2.4 | Radioactivity measurements

^{137}Cs radioactivity was analyzed in charr, prey, and litter using gamma ray spectroscopy.

Gamma ray emissions at 661.6keV were measured using a high-purity germanium

coaxial detector system (GEM20-70; Ortec, Oak Ridge, TN, USA) coupled with a

multi-channel analyzer (DSPECjr2.0; Ortec). The energy and efficiency calibrations

were performed using standard and blank (background) samples. The geometry was

held constant when counting all samples for ^{137}Cs concentrations. For the ^{137}Cs activity

analysis, all samples had counting errors <10%. All activities were corrected for decay

based on the sampling date, and ^{137}Cs concentrations below the detection limit were

designated as 0 Bq/kg for the statistical analysis. The detector system was assessed

using proficiency test samples from the International Atomic Energy Agency

(IAEA-TEL-2012-03).

To estimate representative values for ^{137}Cs concentrations in terrestrial and

aquatic preys, we calculated the median ^{137}Cs concentrations for them. Then, sum of

multiplication of the median concentrations and proportions of biomass of terrestrial

and aquatic prey in the stomach contents was used as "estimated ^{137}Cs concentrations in

prey" for individual charr. Because ^{137}Cs concentrations in animals can differ between

terrestrial and aquatic environments due to ^{137}Cs leaching from litter (Sakai et al.,

2016a), the relative proportions of terrestrial and aquatic prey could substantially

influence ^{137}Cs uptake in charr.

2.5 | Specific metabolic rate

Specific metabolic rate is an important parameter for estimating whole-body ^{137}Cs

concentrations in fish (Ugedal et al., 1992) that was used in this study as an indicator of

^{137}Cs elimination from charr. The specific metabolic rate (B) (W/kg) of charr was

calculated based on the equation below (Brown 2004):

$B = b_0 M^{-1/4} e^{-E/kT}$,

where b_0 is a normalization constant independent of body size and temperature, M is the

wet weight of charr (kg), E is the activation energy, k is Boltzmann's constant, and T is

the absolute water temperature (K). The wet weights of each charr were substituted into

M, and *T* was obtained from the average of the water temperatures recorded over 1 week

at the start of the field investigation.

2.6 | Statistical analysis

One-way analyses of variance (ANOVAs) with permutation test (Legendre, 2007) were

used for testing differences among seasons for (1) standard body lengths of charr in both

samples for estimating ^{137}Cs concentrations in preys and specific metabolic rates and for

measuring ^{137}Cs concentrations in charr, (2) prey biomasses from terrestrial and aquatic

environments estimated with driftnets, pan traps, and stomach pumps, (3) estimated

^{137}Cs concentrations in prey, (4) specific metabolic rates of charr, and (5) ^{137}Cs

concentrations in charr for each site. When significant, we used the same permutation

ANOVA test for post hoc comparisons among the pairs of seasons. The permutation test

allowed us to correctly handle the unbalanced data set of the study. We adjusted p

values for multiple tests using the false discovery rate correction to avoid inflated type I

error rates.

One-way ANOVAs with permutation test was also used to test differences of

standard body length of charr between the samples for estimating ^{137}Cs concentrations

in preys and specific metabolic rates and for measuring ^{137}Cs concentrations in charr.

Further, we tested the difference in ^{137}Cs concentrations in litter and prey between

habitats (i.e., terrestrial and aquatic) using the ANOVAs. All statistical procedures were

conducted using R 3.6.3 (R Core Team, 2020) with anova.1way function (Legendre,

2007) and rcompanion package (Mangiafico, 2020).

3 | RESULTS

3.1 | Body size distribution of charr

The comparisons among seasons indicated that standard body lengths of charr were similar in both the samples for estimating specific metabolic rate and ^{137}Cs concentrations in prey and for measuring ^{137}Cs concentrations in charr (Table 1 and Figure 1). Also, the size distribution was similar between the two sample categories. Although positive correlations between ^{137}Cs concentration and body size were reported in fishes (Wada et al., 2019; Ishii et al., 2020), we assumed that such body size effect is negligible for further discussion in this study.

3.1 | Charr prey

The driftnet and pan trap samples indicated that prey tended to be more abundant in summer and spring than in the other seasons at both the Fukushima and Gunma sites (Table 2). Seasonal variations in terrestrial and aquatic prey biomass were similar and peaked in summer and spring, though the statistical significances were detected only in some pan trap data (Tables 2 and 3). Based on the stomach content samples, charr consumed terrestrial prey more in summer and aquatic prey more in autumn and spring (Table 2). Charr at both the Fukushima and Gunma sites consistently consumed greater amounts of prey in summer, autumn, and spring and less in winter, though there was no statistically significant difference among seasons (Table 2). The three most common terrestrial taxa found in the stomach content samples from fish at the Fukushima site were Anurans (36.3%), Orthopterans (22.6%), and Oligochaetans (15.3%), whereas those at the Gunma site were Orthopterans (40.2%), Coleopterans (25.7%), and Anurans (16.3%) (Table 4). The three main aquatic taxa in the stomachs of the fish at the Fukushima site were Isopods (33.9%), Decapods (28.3%), and Plecopterans (10.4%),

whereas those at the Gunma site were Trichopterans (26.7%), Dipterans (24.8%), and

Decapods (14.7%) (Table 4).

3.2 | ^{137}Cs concentrations in litter and prey

The ^{137}Cs concentrations in terrestrial and aquatic litter from the Fukushima site were

14,700–36,000 and 3,960–8,500 Bq/kg, respectively, whereas those from the Gunma

site were 2,620–8,830 and 1,100–7,320 Bq/kg, respectively. ^{137}Cs concentrations in

terrestrial litter were significantly higher than those in aquatic litter at both sites, and the

difference in ^{137}Cs concentrations between aquatic and terrestrial litter was greater at the

Fukushima site than at the Gunma site (Figure 2).

We measured ^{137}Cs concentrations in four orders of terrestrial organisms and

eight orders of aquatic organisms from the Fukushima site, and four orders of terrestrial

organisms and six orders of aquatic organisms from the Gunma site. For both sites,

these measurements included the three most common taxa found in the stomach

contents of charr, except terrestrial Oligochaeta at the Fukushima site. In total, the Cs

measurements of terrestrial and aquatic prey included 88.1% of the stomach contents of

charr at the Fukushima site and 87.6% of those at the Gunma site (Table 4).

The ^{137}Cs concentrations in prey were significantly higher in terrestrial than

aquatic environments at the Fukushima site but not at the Gunma site (Figure 2). The

median ^{137}Cs concentrations in terrestrial and aquatic prey were 4,000 and 853 Bq/kg,

respectively, at the Fukushima site and 393 and 274 Bq/kg at the Gunma site. The

estimated ^{137}Cs concentrations in prey at the Fukushima site changed seasonally (Table

5 and Figure 3) by seasonal shifts of the relative contribution of terrestrial and aquatic

prey (Table 2) whereas those remained relatively constant across seasons at the Gunma

site due to the greater similarity in ^{137}Cs concentrations in terrestrial and aquatic prey

(Figure 3).

3.3 | Charr metabolism

Because of the structure of the equation of specific metabolic rate, the specific

metabolic rates of charr increased with stream water temperature and were thus highest

in summer and lowest in winter at both sites (Table 5 and Figure 3). The ranges of the

specific metabolic rates of charr were: 0.0008–0.0041 and 0.0006–0.0070 W/kg at the

Fukushima and Gunma sites, respectively.

3.4 | ^{137}Cs concentration in charr

The concentrations of ^{137}Cs in charr ranged from 1,190 to 6,080 and from 118 to 617

Bq/kg at the Fukushima and Gunma sites, respectively. The ANOVA models showed

that the ^{137}Cs concentrations in charr did not differ among seasons at the Fukushima

sites whereas those at the Gunma site was relatively higher in winter (Table 5 and

Figure 3).

4 | DISCUSSION

The charr at our study sites consumed both terrestrial and aquatic organisms, and the

consumption amounts of terrestrial and aquatic preys were greater in summer and spring,

respectively. This pattern was in agreement with seasonal dependencies in terrestrial and

aquatic food resources for salmonids in other temperate forested streams (Nakano et al.,

1999a; Baxter et al., 2005). The seasonal dependencies were basically reported from

broad-leaved deciduous forests (e.g., Cloe & Garman, 1996; Kawaguchi & Nakano,

2001; Nakano & Murakami, 2001), but our results confirmed that the similar seasonal

patterns can occur also in evergreen conifer plantations.

As previous studies reported ^{137}Cs leaching from contaminated litter in streams

(Sakai et al., 2015; Gomi et al., 2018), the ^{137}Cs concentrations in Japanese cedar litter

were apparently different between terrestrial and aquatic environments. Sakai et al.

(2015) found that ^{137}Cs concentrations in litter were 0.56 times the initial concentration

after 30 days of submersion in water. Meanwhile, our results at the Fukushima site

found a greater leaching effect on ^{137}Cs concentration in litter (0.26 times the original

concentration) compared to Sakai et al. (2015). This is attributable to longer periods of

water submersion in the field, as Gomi et al. (2018) showed that 240 days of

submersion in streams decreased ^{137}Cs concentrations by 90% in Japanese cedar litter.

In general, Japanese cedar litter is more persistent than broad-leaved deciduous litter is

(Sakai et al., 2013); therefore, Cs leaching from litter may be prolonged, resulting in

remarkably lower ^{137}Cs concentrations in aquatic litter.

Besides temporal differences in the amount of Cs leaching, our results

indicated that the difference in ^{137}Cs concentration in litter between terrestrial and

aquatic environments was smaller at the Gunma site (0.67×) than at the Fukushima site,

despite the similar decomposition stages of the litter samples. This indicates that Cs

leaching at the Fukushima site was more effective than at the Gunma site. Because

cesium leaches well from litter when differences in the concentrations between litter and

water are greater (Sheppard & Evenden, 1990), the difference of ^{137}Cs concentrations in

litter between terrestrial and aquatic environments becomes linearly greater in more

polluted sites (Sakai et al., 2016b). This is primarily associated with positive

relationships between ^{137}Cs concentrations in litter and initial fallout volumes, and

ubiquitously very low ^{137}Cs concentrations in stream water (Ueda et al., 2013; Sakai et

al., 2015; Iwagami et al., 2017). Actually, FFPRI (2012) indicated that stream water of

our Fukushima site hardly contains ^{137}Cs, probably because dissolved ^{137}Cs is unstable

and adsorbed to clay minerals immediately (Sakai et al., 2015).

The ^{137}Cs concentrations in terrestrial and aquatic prey reflected the

contamination levels of the basal food resource (Japanese cedar litter) between

terrestrial and aquatic ecosystems, in agreement with Sakai et al. (2016a) and Haque et

al. (2018). Interestingly, the difference of ^{137}Cs concentration between terrestrial and

aquatic preys was greater at the Fukushima site than at the Gunma site, corresponding to

the ^{137}Cs concentration difference between terrestrial and aquatic litter (Sakai et al.,

2016b). These results suggest that the underlying relationship between initial fallout

volume and ^{137}Cs leaching efficiency of litter is tightly linked to ^{137}Cs concentration

difference between terrestrial and aquatic preys in forested headwaters where detrital

food webs predominate.

Food resource is the most dominant source of ^{137}Cs uptake in freshwater fishes

(Hewett & Jefferies, 1978), and thus contamination of charr muscle tissues would be

influenced by seasonal variations in ^{137}Cs concentration in preys. In mammal studies, it

was reported that ^{137}Cs concentration in deer becomes higher in autumn when actively

foraging mushrooms that contain much larger amount of ^{137}Cs than other foods (Karlén

et al., 1991; Avila et al., 1999; Zibold et al., 2001). At the Fukushima site, terrestrial

preys were highly contaminated, supplied more in summer and less in winter, and the

overall ^{137}Cs concentration in preys was estimated higher in summer and lower in winter.

In contrast, the estimated ^{137}Cs concentrations in prey at the Gunma site were relatively

consistent throughout the season because ^{137}Cs concentration was similar between

terrestrial and aquatic preys. Meanwhile, seasonal change in metabolism of charr was

commonly high in summer and low in winter at the sites because that is basically

controlled by water temperature (Brown, 2004). Based on the inconsistency of ^{137}Cs

uptake and consistency of metabolism between the study sites, the ^{137}Cs concentrations

in charr was consistent at the Fukushima site, whereas that was negatively relevant to

specific metabolic rate at the Gunma site. At the Fukushima site, because both ^{137}Cs

concentrations in prey and specific metabolic rates of charr peaked in summer, ^{137}Cs

uptake and elimination in charr could have cancelled out any seasonal ^{137}Cs fluctuations,

and thus ^{137}Cs concentration in charr population was stable. On the other hand,

fluctuations in specific metabolic rate were more directly related to the ^{137}Cs

concentrations in charr at the Gunma site because ^{137}Cs concentration in preys was

consistent. These results suggest a possible threshold at which uptake or elimination

influences ^{137}Cs concentrations in charr depending on initial ^{137}Cs fallout volumes.

Although previous studies have indicated that importance of both uptake (Avila et al.,

1999) and metabolism (Rowan & Rasmussen, 1995; Peles et al., 2000), our findings

indicate that the relative influence of the two effects on ^{137}Cs concentrations in charr can

vary depending on the degree of initial ^{137}Cs fallout.

This study was conducted in an early stage of the radiation contamination (one

to two years after the accident). Currently, most litter is less contaminated compared to

those in the early stage because leaves and needles contaminated by the ^{137}Cs fallout in

2011 had already been abscised. In particular, ^{137}Cs concentrations in Japanese cedar

litter was remarkably decreased with time, whereas those in deciduous broad-leaved

litter remained at higher levels through ^{137}Cs root uptake (Onda et al., 2020). Therefore,

difference of ^{137}Cs concentrations in litter between forest and stream ecosystems may

become smaller especially in coniferous plantations of heavily contaminated areas, and

current seasonal fluctuations in ^{137}Cs concentration in headwater salmonids may be

influenced primarily by their metabolism. These characteristics of temporal changes in

^{137}Cs concentration in litter should be crucial when assess ^{137}Cs uptake to fishes from

preys that exhibit variable ^{137}Cs concentrations through the leaching effects on detrital

food webs between forest and stream ecosystems.

In this study, both metabolism and ^{137}Cs concentrations in prey items, whose

seasonality are varied due to initial ^{137}Cs fallout volume, were expected to be important

determinants for ^{137}Cs concentrations in charr. Meanwhile, the difficulty on

radioactivity measurements for small-volume samples has largely hindered quantitative

(not concentration) assessments on ^{137}Cs flows in predator-prey interactions. Our results

also could not estimate quantitative input and output of ^{137}Cs in fishes, but the plausible

processes proposed here suggest the importance of the input/output balance. At least,

our results suggested that less ^{137}Cs-concentrated freshwater salmons might be more

obtainable in hotter season, when specific metabolic rate is higher, in moderately

contaminated sites. Meanwhile in highly contaminated sites, reducing terrestrial preys

belonging to heavily contaminated detrital food webs is important to acquire less

contaminated fishes. Further studies on radiocesium transfers and dynamics in fishes

would help develop better contamination managements in freshwater fisheries. Then,

temporal changes in ^{137}Cs concentrations in litter is especially crucial to assess

headwater salmonids that belong to detrital food webs tangled in forest and stream

ecosystems.

ACKNOWLEDGMENTS

We thank the members of the Watershed Hydrology and Ecosystem Management

Laboratory at the Tokyo University of Agriculture and Technology for their support in

conducting the fieldwork. In addition, we extend our appreciation to the members of the

Watershed Conservation and Management Laboratory at Hokkaido University. This

study was supported by the Environmental Research Fund (ZD-1202) of the Ministry of

the Environment, Japan, and the JSPS KAKENHI Grant Numbers 24248058 and

15H04511.

REFERENCES

Avila, R., Johanson, K. J., & Bergström, R. (1999). Model of the seasonal variations of

fungi ingestion and [137]Cs activity concentrations in roe deer. Journal of

Environmental Radioactivity, 46, 99-112.

Baxter, C. V., Fausch, K. D., & Saunders, W. C. (2005). Tangled webs: reciprocal flows

of invertebrate prey link streams and riparian zones. Freshwater Biology, 50,

201-220.

Brown, J. H. (2004). Toward a metabolic theory of ecology. Ecology, 85, 1771-1789.

Cloe, W., & Garman, G. (1996). The energetic importance of terrestrial arthropod inputs

to three warm-water streams. Freshwater Biology, 36, 104-114.

Doi, H., Takahara, T. & Tanaka, K. (2012). Trophic position and metabolic rate predict

the long-term decay process of radioactive cesium in fish: A meta-analysis. Plos

One, 7, e29295.

Forestry and Forest Products Research Institute (FFPRI). (2012). Monitoring results of

radionuclide concentrations in stream water during Baiu season,

http://www.ffpri.affrc.go.jp/press/2012/20120921/documents/20120921.pdf

(accessed: May 2021). (in Japanese)

Gomi, T., Sidle, R. C. & Richardson, J. S. (2002). Understanding processes and

downstream linkages of headwater systems. BioScience, 52, 905-916.

Gomi, T., Sakai, M., Haque, M. E., Hosoda, K., & Toda, H. (2018). Evaluating [137]Cs

detachment from coniferous needle litter in a headwater stream: a litter bag field

experiment. Landscape and Ecological Engineering, 14, 17-27.

Haque, M. E., Gomi, T., Sakai, M., & Negishi, J. N. (2017). Developing a food

web-based transfer factor of radiocesium for fish, whitespotted char (*Salvelinus*

leucomaenis) in headwater streams. Journal of Environmental Radioactivity, 172,

191-200.

Haque, M. E., Gomi, T., Sakai, M., & Negishi, J. N. (2018). Seasonal variation in food

web-based transfer factors of radiocesium in white-spotted char (*Salvelinus*

leucomaenis) from headwater streams. Landscape and Ecological Engineering, 14,

45-53.

Hessen, D. O., Skurdal, J., Hegge, O., & Hesthagen, T. (2002). Radiocesium decay in

populations of brown trout and Arctic char in the alpine Atna area, south-eastern

Norway. Hydrobiologia, 489, 55-62.

Hewett, C. J., & Jeffries, D. F. (1978). The accumulation of radioactive caesium from

food by the plaice (*Pleuronectes platessa*) and the brown trout (*Salmo trutta*).

Journal of Fish Biology, 13, 143-153.

Holloman, K. A., Dallas, C. E., Brisbin Jr., I. L., & Jagoe, C. H. (1997). Spatial and

temporal patterns of radiocesium contamination in mosquitofish, *Gambusia*

holbrooki (Girard, 1859), inhabiting a nuclear reactor cooling reservoir. Journal of

Environmental Radioactivity, 35, 243-359.

Inoue, M., Sakamoto, S., & Kikuchi, S. (2013). Terrestrial prey inputs to streams

bordered by deciduous broadleaved forests, conifer plantations and clear-cut sites

in southwestern Japan: effects on the abundance of red-spotted masu salmon.

Ecology of Freshwater Fish, 22, 335-347.

Ishii, Y., Matsuzaki, S. S., Hayashi, S. (2020). Different factors determine ^{137}Cs

concentration factors of freshwater fish and aquatic organisms in lake and river

ecosystems. Journal of Environmental Radioactivity, 213, 106102.

Iwagami, S., Tsujimura, M., Onda, Y., Nishino, M., Konuma, R., Abe, Y., Hada, M., Pun,

I., Sakaguchi, A., Kondo, H., Yamamoto, M., Miyata, Y., & Igarashi, Y. (2017).

Temporal changes in dissolved ^{137}Cs concentrations in groundwater and stream

water in Fukushima after the Fukushima Dai-ichi Nuclear Power Plant accident.

Journal of Environmental Radioactivity, 166, 458-465.

Karlén, G., Johanson, K. J., & Bergström, R. (1991). Seasonal variation in the activity

concentration of ^{137}Cs in Swedish Roe-deer and in their daily intake. Journal of

Environmental Radioactivity, 14, 91-103.

Kawaguchi, Y., & Nakano, S. (2001). Contribution of terrestrial invertebrates to the

annual resource budget for salmonids in forest and grassland reaches of a

headwater stream. Freshwater Biology, 46, 303-316.

Kolehmainen, S. E. (1972). The balance of ^{137}Cs , stable cesium and potassium of

bluegill (*Lepomis macrochirus* Raf.) and other fish in white Oak Lake. Health

Physics, 23, 301-315.

Legendre P (2007) One-way anova with permutation test,

http://adn.biol.umontreal.ca/~numericalecology/Rcode/ (accessed: May 2021).

Maeda, N., & Matsui, M. (1999). Frogs and Toads of Japan, Revised Edition. Tokyo:

Bun-ichi Co.,Ltd.

Mangiafico S (2020) Package 'rcompanion',

https://cran.r-project.org/web/packages/rcompanion/ (accessed: May 2021).

Merritt, R. W., Cummins, K. W., & Berg, M. B. (2008). An introduction to the aquatic

insects of North America, fourth edition. Iowa: Kendall/Hunt Publishing

Company.

Ministry of Agriculture, Forestry and Fisheries, Japan (MAFF). (2021). Results on

measurement of radiocesium activity concentrations in agricultural products,

https://www.maff.go.jp/j/kanbo/joho/saigai/s_chosa/index.html (accessed: May

2021). (in Japanese)

Ministry of Education, Culture, Sports, Science and Technology, Japan (MEXT), (2012).

Results of the fifth aircraft monitoring on air dose rates and [137]Cs depositions,

http://radioactivity.nsr.go.jp/ja/contents/7000/6289/24/203_0928.pdf (accessed:

May 2021). (in Japanese)

Montgomery, D. R., & Buffington, J. M. (1997). Channel-reach morphology in

mountain drainage basins. Geological Society of America Bulletin, 109, 596-611.

Nakano, S., Miyasaka, H., & Kuhara, N. (1999a). Terrestrial-aquatic linkages: riparian

arthropod inputs alter trophic cascade in a stream food web. Ecology, 80,

2435-2441.

Nakano, S., Kawaguchi, Y., Taniguchi, Y., Miyasaka, H., Shibata, Y., Urabe, H., &

Kuhara, N. (1999b). Selective foraging on terrestrial invertebrates by rainbow

trout in a forested headwater stream in northern Japan. Ecological Research, 14,

351-360.

Nakano, S., & Murakami, M. (2001). Reciprocal subsidies: dynamic interdependence

between terrestrial and aquatic food webs. Proceedings of National Academy of

Sciences of the United States of America, 98, 166-170.

Negishi, J. N., Sakai, M., Okada, K., Iwamoto, A., Gomi, T., Miura, K., Nunokawa, M.,

& Ohhira, M. (2018). Cesium-137 contamination of river food webs in a gradient

of initial fallout deposition in Fukushima, Japan. Landscape and Ecological

Engineering, 14, 55-66.

Onda, Y., Taniguchi, K., Yoshimura, K., Kato, H., Takahashi, J., Wakiyama, Y., Coppin,

F., Smith, H. (2020). Radionuclides from the Fukushima Daiichi Nuclear Power

Plant in terrestrial systems. Nature Reviews Earth & Environment, 1, 644-660.

Peles, J. D., Philippi, T., Smith, M. H., Brisbin, Jr. I. L., & Gibbons, J. W. (2000).

Seasonal variation in radiocesium levels of largemouth bass (*Micropterus*

salmoides): implications for humans and sensitive wildlife species.

Environmental Toxicology and Chemistry, 19, 1830-1836.

R Core Team (2020). R: a language and environment for statistical computing,

https://www.R-project.org/ (accessed: May 2021).

Rowan, D. J., & Rasmussen, J. B. (1995). The elimination of radiocaesium from fish.

Journal of Applied Ecology, 32, 739-744.

Rowan, D. J., Rasmussen, J. B., & Chant, L. (1997). A bioenergetics approach to

modeling seasonal patterns in the lation of radiocesium. In G. Desmet, R. J. Blust,

R. N. J. Comans, J. A. Fernandez, J. Hilton, & A. de Bettencourt (Eds.),

Freshwater and Estuarine Radioecology (pp. 387-393). Amsterdam: Elsevier.

Sakai, M., Natuhara, Y., Fukushima, K., Imanishi, A., Imai, K., & Kato, M. (2013).

Ecological functions of persistent Japanese cedar litter in structuring stream

macroinvertebrate assemblages. Journal of Forest Research, 18, 190-199.

Sakai, M., Gomi, T., Naito, R. S., Negishi, J. N., Sasaki, M., Toda, H., Nunokawa, M.,

& Murase, K. (2015). Radiocesium leaching from contaminated litter in forest

streams. Journal of Environmental Radioactivity, 144, 15-20.

Sakai, M., Gomi, T., Negishi, J. N., Iwamoto, A., & Okada, K. (2016a). Different

cesium-137 transfers to forest and stream ecosystems. Environmental Pollution,

209, 46-52.

Sakai, M., Gomi, T., & Negishi, J. N. (2016b). Fallout volume and litter type affect

^{137}Cs concentration difference in litter between forest and stream environments.

Journal of Environmental Radioactivity, 164, 169-173.

Sakai, M., Fukushima, K., Naito, R. S., Natuhara, Y., & Kato, M. (2016c). Coniferous

needle litter acts as a stable food resource for stream detritivores. Hydrobiologia,

779, 161-171.

Sheppard, S. C., & Evenden, W. G. (1990). Leaching of radionuclides from decaying

blueberry leaves: relative rate independence of concentration. Journal of

Environmental Quality, 19, 464-469.

Ueda, S., Hasegawa, H., Kakiuchi, H., Akata, N., & Ohtsuka, Y. (2013). Fluvial

discharges of radiocaesium from watersheds contaminated by the Fukushima

Dai-ichi Nuclear Power Plant accident, Japan. Journal of Environmental

Radioactivity, 118, 96-104.

Ugedal, O., Jonsson, B., Njåstad, O., & Næumann, R. (1992). Effects of temperature

and body size on radiocaesium retention in brown trout, *Salmo trutta*. Freshwater

Biology, 28, 165-171.

Wada, T., Konoplev, A., Wakiyama, Y., Watanabe, K., Furuta, Y., Morishita, D., Kawata,

G., Nanba, K. (2019). Strong contrast of cesium radioactivity between marine and

freshwater fish in Fukushima. Journal of Environmental Radioactivity, 204,

132-142.

Wallace, J. B., Eggert, S. L., Meyer, J. L., & Webster, J. R. (1997). Multiple trophic

levels of a forest stream linked to terrestrial litter inputs. Science, 277, 102-104.

Zibold, G., Drissner, J., Kaminski, S., Klemt, E., & Miller, R. (2001). Time-dependence

of the radiocaesium contamination of roe deer: measurement and modelling.

Journal of Environmental Radioactivity, 55, 5-27.

Figure legends

FIGURE 1 Standard body lengths of charr samples of this study. The left graphs show

the standard body lengths of the charr samples used for estimating ^{137}Cs concentrations

in prey and specific metabolic rates at the Fukushima and Gunma sites. The right graphs

show those used for measuring ^{137}Cs concentrations in charr of the Fukushima and

Gunma sites. The sample sizes are shown at tops of each bar.

FIGURE 2 ^{137}Cs concentrations in Japanese cedar litter and preys collected from both

terrestrial and aquatic habitats in the Fukushima (upper) and Gunma (lower) sites.

Different alphabetical letters indicate statistically significant difference (permutation

analyses of variance, $P < 0.05$) and n.s. indicates insignificant difference.

FIGURE 3 Seasonal variations in the estimated ^{137}Cs concentrations in prey, specific

metabolic rates of charr and ^{137}Cs concentrations in charr at the Fukushima (left panels)

and Gunma sites (right panels). Different alphabetical letters and circles indicate

statistically significant difference (post hoc comparisons, $P < 0.05$ and < 0.1,

respectively) and n.s. indicates insignificant difference.

TABLE 1 Results of permutation analyses of variance testing for effect of season on the

standard body lengths of charr (sample 1: the samples for estimating specific metabolic

rate and ^{137}Cs concentrations in preys, sample 2: the samples for measuring ^{137}Cs

concentrations in charr) and comparing standard body lengths of charr between the

samples 1 and 2.

Factor	Fukushima		Gunma	
	F	P	F	P
Sample 1:				
Season	2.27	0.08	0.21	0.90
Sample 2:				
Season	0.38	0.78	0.69	0.54
Sample (1 vs 2)	0.03	0.86	1.07	0.33

			Fukushima								Gunma								
			Total prey			Terrestrial prey			Aquatic prey				Total prey			Terrestrial prey			Aquatic prey		
Method	Season	n	Mean	±	SE	Mean	±	SE	Mean	±	SE	n	Mean	±	SE	Mean	±	SE	Mean	±	SE
Driftnet	Summer	10	**8.51**	±	2.46	0.15	±	0.06	**8.36**	±	2.45	10	8.84	±	2.49	0.17	±	0.11	8.66	±	2.39
	Autumn	10	6.3	±	1.76	0.89	±	0.57	5.41	±	1.3	10	2.05	±	0.57	0.13	±	0.04	1.92	±	0.55
	Winter	10	5.17	±	1.22	0.05	±	0.03	5.13	±	1.22	10	2.41	±	0.57	0.09	±	0.03	2.32	±	0.57
	Spring	10	7.91	±	1.32	**0.96**	±	0.8	6.95	±	1.27	10	**12.21**	±	7.38	**0.64**	±	0.26	**11.58**	±	7.39
Pan trap	Summer	5	**888.96**	±	674.84[ab]	**810.12**	±	661.81[ab]	78.84	±	21.34	5	**216.96**	±	77.02[ab]	**197.01**	±	72.98	**19.95**	±	7.39
	Autumn	5	58.9	±	9.03[a]	41.78	±	5.18[a]	17.12	±	3.87	5	30.38	±	5.24[a]	20.06	±	5.14	10.32	±	2.8
	Winter	5	0.98	±	0.46[b]	0.51	±	0.25[b]	0.47	±	0.25	5	1.01	±	0.27[b]	0.34	±	0.18	0.66	±	0.25
	Spring	5	94.46	±	18.91[a]	54.98	±	12.04[a]	39.48	±	13.77	5	54.58	±	13.99[a]	35.3	±	13.94	19.28	±	4.99
Stomach pump	Summer	24	12.58	±	5.7	**9.44**	±	5.45	3.13	±	1.47	37	12.13	±	5.52	**11.22**	±	5.48	0.92	±	0.33[b]
	Autumn	34	**21.55**	±	7.67	6.39	±	3.26	15.17	±	6.68	29	**12.26**	±	5.25	7.51	±	4.13	4.75	±	1.35[a]
	Winter	26	4.18	±	2.38	0.42	±	0.29	3.76	±	2.35	36	1.94	±	0.96	0	±	0	1.94	±	0.96[ab]
	Spring	28	16.88	±	5.6	1.33	±	0.7	**15.55**	±	5.41	37	10.01	±	3.08	2.9	±	0.92	**7.11**	±	2.65[ab]

619 **TABLE 3** Results of permutation analyses of variance testing for effect of season on

620 biomass of total, terrestrial, and aquatic preys estimated by driftnet, pan trap, and

621 stomach pump in each site. Bold characters indicate statistically significant differences

622 ($P < 0.05$).

Method	Prey type	Fukushima		Gunma	
		F	*P*	*F*	*P*
Driftnet	Total prey	0.75	0.54	1.62	0.12
	Terrestrial prey	0.96	0.42	3.17	**0.01**
	Aquatic prey	0.83	0.46	1.50	0.15
Pan trap	Total prey	1.55	**<0.01**	7.88	**<0.01**
	Terrestrial prey	1.39	**<0.01**	7.58	**<0.01**
	Aquatic prey	6.97	**<0.01**	4.48	**0.02**
Stomach pump	Total prey	1.52	0.22	1.47	0.22
	Terrestrial prey	1.78	0.15	2.18	0.07
	Aquatic prey	1.83	0.14	3.18	**<0.01**

624 **TABLE 4** Total biomass (mg) of terrestrial and aquatic prey taxa estimated from the

625 stomach contents of charr.

Fukushima				Gunma			
Terrestrial		Aquatic		Terrestrial		Aquatic	
Anura	178.70	Isopoda	380.44	Orthoptera	297.63	Trichoptera	134.93
Orthoptera	111.24	Decapoda	318.09	Coleoptera	190.25	Diptera	125.08
Oligochaeta	75.40	Plecoptera	117.10	Anura	120.70	Decapoda	74.30
Diplopoda	35.00	Diptera	112.77	Lepidoptera	43.20	Ephemeroptera	70.72
Coleoptera	21.50	Trichoptera	87.91	Diplopoda	32.90	Salmoniformes	29.10
Unknown	20.77	Odonata	44.30	Araneae	26.53	Odonata	26.32
Hemiptera	18.92	Ephemeroptera	43.58	Hymenoptera	11.83	Plecoptera	24.55
Hymenoptera	15.56	Coleoptera	19.58	Oligochaeta	7.20	Coleoptera	10.88
Araneae	8.53	Neuroptera	0.01	Unknown	5.41	Isopoda	8.70
Gastropoda	5.90			Hemiptera	4.30	Neuroptera	0.17
Collembola	0.54			Collembola	0.23		
Acari	0.03			Acari	0.05		

629　**TABLE 5** Results of permutation analyses of variance testing for effect of season on

630　estimated ^{137}Cs concentrations in prey, specific metabolic rate and ^{137}Cs concentrations

631　in charr. Bold characters indicate statistically significant differences ($P < 0.05$).

	Fukushima		Gunma	
	F	*P*	*F*	*P*
Estimated ^{137}Cs concentrations in prey	5.32	**<0.01**	6.41	**<0.01**
Specific metabolic rate	128.89	**<0.01**	151.63	**<0.01**
^{137}Cs concentrations in charr	0.02	1.00	4.35	**0.02**