Research article

Partitioning the temporal changes in abundance-based beta diversity into loss and gain components

Shinichi Tatsumi[1,2,*], Ryosuke Iritani[3], and Marc W. Cadotte[1,4]

[1] Department of Biological Sciences, University of Toronto Scarborough, Toronto, ON, Canada

[2] Hokkaido Research Center, Forestry and Forest Products Research Institute, Hokkaido, Japan

[3] RIKEN Interdisciplinary Theoretical and Mathematical Sciences (iTHEMS), Saitama, Japan

[4] Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, ON, Canada

*Corresponding author: Shinichi Tatsumi

Address: Hokkaido Research Center, Forestry and Forest Products Research Institute, Hitsujigaoka 7, Toyohira-ku, Sapporo, Hokkaido 062-8516, Japan

Email: community.ecologist@gmail.com

ORCID

Shinichi Tatsumi: 0000-0002-1789-1685

Ryosuke Iritani: 0000-0002-2396-1109

Marc W. Cadotte: 0000-0002-5816-7693

Running headline: Partitioning temporal changes in beta diversity

ABSTRACT

1. Ecologists have long recognized that the losses and gains in local species abundances can either decrease or increase spatial beta diversity, phenomena often referred to as biotic homogenization and differentiation, respectively. However, quantifying such dynamic impacts of species abundances on beta diversity has remained a methodological challenge.

2. Here, we develop a numerical method to additively partition the temporal changes in beta diversity into distinct components that reflect the losses and gains in local species abundances. Our method is based on Ružička and Bray–Curtis indices and the normalized abundance-based Whittaker's beta diversity. The temporal changes in these measures are partitioned into components that represent biotic homogenization and differentiation driven by abundance losses and gains at both species and community levels.

3. Application of the method to a Swedish fish community dataset revealed decreases in beta diversity between 1990 and 2018. The homogenization of fish communities was explained by gains, but not losses, in species abundances across sites. Species-level partitioning further showed that the homogenization was largely caused by the increased population sizes of a particular species in sites where it was already present.

4. The results highlight that our partitioning method effectively identifies local population and community processes embedded in regional biodiversity patterns. We believe that explicit analyses of the losses and gains in species abundances should bring deeper insights into the dynamics of beta diversity.

Keywords: Biotic homogenization, Community assembly, Compositional dissimilarity, Ecological dynamics, Metacommunity, Spatial heterogeneity, Species turnover, Time series

1 | INTRODUCTION

Beta diversity, the variation in the identities and abundances of species among sites, is a fundamental facet of biodiversity (Whittaker, 1960; Koleff et al., 2003; Anderson et al., 2011). Beta diversity can be quantified in two ways, namely by using incidence-based (i.e., presence–absence-based) and abundances-based approaches (Chao et al., 2005; Legendre & Legendre, 2012; Baselga, 2013). The two approaches weigh rare and dominant species differently, and thus offer complementary insights into community structure (Anderson et al., 2011; Legendre & Legendre, 2012; Li et al., 2016). Extensions of existing methods for analyzing incidence-based beta diversity to account for abundance can provide a more comprehensive understanding of biodiversity (Baselga, 2013; Chao et al., 2014).

While the replacement of endemic species by cosmopolitan nonnative species has been a global concern (McKinney & Lockwood, 1999), we still have mixed evidence for the consequent changes in beta diversity over time (Olden et al., 2018). The temporal decreases and increases in beta diversity, referred to as biotic homogenization and differentiation, respectively, subsume complex processes of local population dynamics (McKinney & Lockwood, 1999; Olden & Poff, 2003; Socolar et al., 2016; Rosenblad & Sax, 2017; Tatsumi et al., 2020). Empirical and simulation studies have shown that incidence- and abundance-based approaches can result in contrasting signs and magnitudes of temporal changes in beta diversity (Cassey et al., 2008; Li et al., 2016; Petersen et al., 2021). To build a more rigorous evidence base for biotic homogenization and differentiation, we need a tool to disentangle the processes underlying beta diversity changes.

Species extinctions and colonizations (i.e., changes in presence–absence status) can alter beta diversity in multiple ways (Olden & Poff, 2003; Rosenblad & Sax, 2017; Tatsumi et al., 2020, 2021). Specifically, extinctions lead to biotic homogenization when rare, infrequent species become regionally extinct, but otherwise result in differentiation (Socolar et al., 2016; Rosenblad & Sax, 2017; Tatsumi et al., 2020). Colonizations cause homogenization when species new to the region become widespread or existing species increase their regional dominance, but drive differentiation when new species colonize a small number of sites (Socolar et al., 2016; Rosenblad & Sax, 2017; Tatsumi et al., 2020). Extinctions and colonizations can also mask each other when one increases beta diversity and the other decreases it (Tatsumi et al., 2020, 2021). In our previous study (Tatsumi et al., 2021), we proposed a numerical method to additively partition such impacts of extinction and

colonization on spatial beta diversity as quantified by incidence-based measures, namely

Jaccard and Sørensen indices and Whittaker's beta diversity.

Here, we develop a new method to additively partition the impacts of abundance losses

and gains on spatial beta diversity by extending our previous incidence-based method.

Similar to species extinctions and colonizations (i.e., binary changes between presence and

absence), quantitative decreases and increases in local species abundances can drive either

homogenization or differentiation (Socolar et al., 2016). The new method that we propose

here allows one to partition such temporal changes in spatial variation ($\Delta\beta = \beta' - \beta$, where β

and β' are the values at $t = 1$ and 2, respectively) into distinct terms that reflect abundance

losses and gains (Figure 1). Our method helps to resolve the local population dynamics and

metacommunity processes embedded in regional biodiversity patterns using both incidence

and abundance data.

Figure 1. Temporal change in spatial beta diversity and its additive components. The components represent biotic homogenization or differentiation driven by losses or gains in species abundances.

2 | METHODS

The partitioning method described below can be implemented using the ecopart R package

available from GihHub: remotes::install_github("communityecologist/ecopart").

2.1 | Partitioning equations

We describe the additive partitioning of temporal changes in pairwise dissimilarity

measures, namely using Ružička ($\beta_{\text{Ruž}}$) and Bray–Curtis indices (β_{BC}) (Bray & Curtis, 1957;

Ružička, 1958). These measures are defined as $\beta_{\text{Ruž}} = \sum_{i=1}^{S} u_i / \sum_{i=1}^{S} (c_i + u_i)$ and $\beta_{\text{BC}} =$

$\sum_{i=1}^{S} u_i / \sum_{i=1}^{S}(2c_i + u_i)$, where c_i is the component of species abundance common to both

sites, u_i is the component of species abundance unique to either site, i is species identity, and

S is the number of species. Here, let $\beta^{(\lambda)} = \sum_{i=1}^{S} u_i / \sum_{i=1}^{S}(\lambda c_i + u_i)$, then $\beta_{\text{Ruž}} = \beta^{(1)}$ and $\beta_{\text{BC}} =$

$\beta^{(2)}$. Ružička and Bray–Curtis indices are abundance-based extensions of Jaccard and

Sørensen indices, respectively, for which the partitioning methods have already been

described (Tatsumi et al., 2021). Although the Ružička and Bray–Curtis indices have

different statistical properties from other common dissimilarity measures (e.g., Shannon

entropy; Jost, 2007), we use the two indices here on account of their mathematical simplicity

and wide uses in ecology.

Previous studies categorized species extinction and colonization into six types based on

their impacts on spatial dissimilarity (Rosenblad & Sax, 2017; Tatsumi et al., 2020, 2021). We

here extend these definitions to account for species abundances (Figure 2). The first type is

the reduction in u_i (type 1 in Figure 2); that is, for the abundance of a given species i in site k

(a_{ik}), a component that was unique to either site (u_i) at time $t = 1$ becomes lost at time $t = 2$.

Type 2 is the case where a component of a_{ik} that was common to both sites (c_i) becomes lost

by an equal amount in both sites. Type 3 refers to the loss in c_i in the site where a_{ik} was

smaller than, or equal to, that in the other site, turning c_i into u_i. Type 4 refers to the gain in

u_i in the site where a_{ik} was larger than, or equal to, that in the other site. Type 5 is the case

where c_i increases by an equal amount in both sites. Type 6 refers to the gain in c_i in the site

where a_{ik} was smaller than the other site, turning u_i into c_i. Types 1, 5, and 6 decrease $\beta^{(\lambda)}$,

leading to homogenization, whereas types 2, 3, and 4 increase $\beta^{(\lambda)}$, leading to differentiation.

We write $l_i^{\mathbb{L}}, l_i^{\mathbb{E}}, l_i^{\mathbb{S}}, g_i^{\mathbb{L}}, g_i^{\mathbb{E}}$, and $g_i^{\mathbb{S}}$ for the amount of changes in abundance that correspond to

types 1, 2, 3, 4, 5, and 6, respectively (Appendix S1).

It is possible for the abundance of a given species i to change differently in the two sites

within the same time interval. Namely, if the abundance a_{ik} decreases in one site (e.g., $k = 1$)

where it was larger and increases in the other site (e.g., $k = 2$) where it was smaller, then

beta diversity can potentially show no net change (see the bottom case in Figure 2). We refer

to such offsetting replacements in species abundances as hidden dynamics (Tatsumi et al.

2021). We write d_i for the changes in abundance that fall under this definition. In our

partitioning method, we explicitly describe d_i as a distinct form of abundance losses and

gains. In total, there are 32 possible ways a_{ik} can decrease and/or increase, including the

hidden dynamics (Appendix S1). Further, beta diversity can be much more dynamic than

Figure 2. Schematic representation of six types of changes in beta diversity and hidden dynamics. The numbers in the boxes indicate species abundances. The variable u denotes the components of abundance unique to either site and c denotes the components of abundance common to both sites. For example, at $t = 1$, the u of species A is 0 and species B is 5. The c of species A is 10 and species B is 5. Thus, taking Ružička index as an example, $\beta_{\text{Ruž}} = (0 + 5)/(10 + 5 + 0 + 5) = 1/4$. In type 1 at $t = 2$, the u of species B becomes 0 as a result of an abundance loss in one of the two sites. Consequently, $\beta_{\text{Ruž}}$ decreases to 0. In type 2, the c of species A is reduced to 5 and thus $\beta_{\text{Ruž}} = (0 + 5)/(5 + 5 + 0 + 5) = 1/3$. We can see from these examples that, while types 1 and 2 are both associated with abundance losses, the changes in $\beta_{\text{Ruž}}$ can take either negative or positive values. It is such distinct types of changes in beta diversity our method allows one to partition.

any index portrays since abundances can change multiple times and in multiple ways

between sampling intervals, and so here c_i and u_i refer to the net change between $t = 1$ and 2.

For brevity, we write the sum of a given variable across all species (1, 2, ..., S) by using

the uppercase letters (e.g., $\sum_{i=1}^{S} u_i = U$, $\sum_{i=1}^{S} d_i = D$, and $\sum_{i=1}^{S} l_i^{\mathbb{L}} = L^{\mathbb{L}}$). The temporal

changes in C and U can then be written as $\Delta C = C' - C = -L^{\mathbb{E}} - L^{\mathbb{S}} + G^{\mathbb{E}} + G^{\mathbb{S}}$ and $\Delta U = U'$

$- U = -L^{\mathbb{L}} + L^{\mathbb{S}} + G^{\mathbb{L}} - G^{\mathbb{S}}$. We can additively partition the temporal changes in pairwise

dissimilarity ($\Delta\beta^{(\lambda)} = \beta'_{\text{Ružž}} - \beta_{\text{Ruž}}$ when $\lambda = 1$ and $\Delta\beta^{(\lambda)} = \beta'_{\text{BC}} - \beta_{\text{BC}}$ when $\lambda = 2$) into six

terms that correspond to the six types of abundance changes:

$\Delta\beta^{(\lambda)} = \beta^{(\lambda)'} - \beta^{(\lambda)}$

$$
119 \qquad = \underbrace{-\frac{p}{U}\left(L^{\mathbb{L}} + D\right)}_{\Delta\beta_1} + \underbrace{\frac{p}{C}L^{\mathbb{E}}}_{\Delta\beta_2} + \underbrace{\left(\frac{p}{U} + \frac{p}{C}\right)L^{\mathbb{S}}}_{\Delta\beta_3}
$$

$$
120 \qquad + \underbrace{\frac{p}{U}\left(G^{\mathbb{L}} + D\right)}_{\Delta\beta_4} + \underbrace{\frac{-p}{C}G^{\mathbb{E}}}_{\Delta\beta_5} + \underbrace{\left(-\frac{p}{U} - \frac{p}{C}\right)G^{\mathbb{S}}}_{\Delta\beta_6} \qquad (1)
$$

where $p = \frac{\lambda CU}{(\lambda C + U)(\lambda C' + U')}$ and C and U are both non-zero. See Appendix S1 for derivation of

the equation.

The variable D denotes the hidden dynamics. The quantities $-\frac{p}{U}D$ in $\Delta\beta_1$ and $-\frac{p}{U}D$ in

$\Delta\beta_4$ cancel each other out by summing up to zero. These two quantities thus allow us,

without causing any effect on $\Delta\beta$, to explicitly account for D in a manner comparable to

$-\frac{p}{U}L^{\mathbb{L}}$ and $-\frac{p}{U}G^{\mathbb{L}}$.

The first three terms are associated with gains, while the last three are associated with

losses. The six terms respectively correspond to the six types of abundance changes described

in Figure 2. The first term, which is always negative, represents homogenization by

abundance losses ($\Delta\beta_{\text{L}-}$). The second and third terms, which are always positive, indicate

differentiation by abundance losses ($\Delta\beta_{\text{L}+}$). Similarly, the fourth term represents

differentiation by abundance gains ($\Delta\beta_{\text{G}+}$) and the fifth and sixth terms indicate

homogenization by abundance gains ($\Delta\beta_{\text{G}-}$). Depending on the ecological question at hand,

one can sum the terms as $\Delta\beta_{\text{L}} = \Delta\beta_{\text{L}-} + \Delta\beta_{\text{L}+}$ and $\Delta\beta_{\text{G}} = \Delta\beta_{\text{G}-} + \Delta\beta_{\text{G}+}$ to assess the total

impact of abundance losses and gains on $\Delta\beta$, respectively (Figure 1).

See Appendix S2 for the responses of $\Delta\beta$ components ($\Delta\beta_{\text{L}-}$, $\Delta\beta_{\text{L}+}$, $\Delta\beta_{\text{G}-}$, and $\Delta\beta_{\text{G}+}$) to

the abundance losses and gains ($L^{\mathbb{L}}$, $L^{\mathbb{E}}$, $L^{\mathbb{S}}$, $G^{\mathbb{L}}$, $G^{\mathbb{E}}$, $G^{\mathbb{S}}$, and D).

2.2 | Multisite variation

Our partitioning method is applicable to multisite measures of beta diversity. Multisite

measures are used to quantify variation among more than two sites (Baselga, 2010).

Averaging pairwise dissimilarities (such as $\beta_{Ruž}$ or β_{BC}) across pairs of sites is a suboptimal

approach due to their lack of statistical independence (Baselga, 2010, 2017).

In Appendix S3, we demonstrate the partitioning of multisite beta diversity by taking,

as an example, the normalized abundance-based Whittaker's beta diversity (β_W) (cf. Baselga,

2017). We chose β_W here on account of its simple mathematical structure. Note, however,

that future works are needed for partitioning other multisite indices, typically beta diversity

based on Hill numbers (see *Discussion* for detail).

2.3 | Species-level impacts on beta diversity

We can further use the partitioning equations to quantify the response of beta diversity to

the losses and gains in the abundance of each species independently. For example, consider

a case where the abundance of a species that had existed in one site was completely lost (i.e.,

the species went locally extinct). This loss will add $-\frac{p}{U}L^{\mathbb{L}}$ to $\Delta\beta$ in Eqn. 1. Thus, the added

value can be interpreted as the consequence of the focal species' abundance loss on beta

diversity. In this way, $\Delta\beta$ can be additively partitioned into components that reflect the

decreases and increases in the population size of individual species. Note that it is possible

for the population size of a given species to decrease in some sites while increase in other

sites within the same time interval, generating both the loss and gain components (i.e., $\Delta\beta_{L-}$,

$\Delta\beta_{L+}$, $\Delta\beta_{G-}$, and $\Delta\beta_{G+}$).

3 | APPLICATIONS

We applied the partitioning method to riverine fish community data retrieved from the

Swedish Electrofishing Register database (Sers, 2013) via RivFishTIME (Comte et al., 2021).

We used data collected in 65 waterbodies consisting of a total of 181 permanent sampling

sites (2–10 sites per waterbody) across Sweden in 1990 and 2018 (see Appendix S4 for site

IDs and selection criteria). The abundance of each fish species in each site was recorded as

the number of individuals per 100 m^2. We quantified the compositional variation among

sites within each waterbody based on the normalized abundance-based Whittaker's beta

diversity using either incidence (presence–absence) or abundance data. The incidence data

was obtained by transforming the abundance values larger than zero to one. We calculated the temporal changes in beta diversity ($\Delta\beta$) and their additive components between 1990 and 2018. The components representing species extinctions and colonizations (i.e., changes from presence to absence and vice versa) are denoted as $\Delta\beta_E$ and $\Delta\beta_C$. Those that represent abundance losses and gains are denoted as $\Delta\beta_L$ and $\Delta\beta_G$.

The incidence- and abundance-based approaches provided complementary insights into the changes in beta diversity (Figure 3). While the incidence-based beta diversity showed no temporal trend (Figure 3a), the abundance-based beta diversity significantly decreased from 1990 and 2018, as indicated by $\Delta\beta$ less than zero (Figure 3b). The loss and gain components ($\Delta\beta_L$ and $\Delta\beta_G$) revealed that this homogenization of fish communities was explained by gains, but not losses, in species abundances (Figure 3b). Partitioning $\Delta\beta$ into species-level components further showed that the homogenization was largely caused by brown trout (*Salmo trutta*) (Figure 3d; see Appendix S4 for the results of all species). The fact that the colonization component $\Delta\beta_C$ of brown trout was not significant (Figure 3c) indicates that the homogenization did not result from colonizations of brown trout to new sites. Rather, it was caused by the increased sizes of brown trout populations (potentially associated with fishing restrictions and stocking; Almesjö & Limén, 2009) in sites where they were already present but in low abundances, leading to a more spatially-uniform abundance distribution.

4 | Discussion

We developed a new method to partition the temporal changes in beta diversity into distinct components that reflect the losses and gains in species abundances (Figure 1). The method provides a unified approach to analyze biotic homogenization and differentiation using both incidence and abundance data. Application of our method to an empirical dataset revealed different trends in incidence- and abundance-based beta diversity ($\Delta\beta_E$, $\Delta\beta_C$, $\Delta\beta_L$, and $\Delta\beta_G$) (Figure 3). The two approaches collectively showed that gains in abundances, but not colonization, of particular species made communities more similar over time (Figure 3). The results highlight that our partitioning method effectively identifies local population and community processes embedded in regional biodiversity patterns.

Moving forward, further generalizations of our temporal partitioning method are needed. In this study, we partitioned Ružička and Bray–Curtis indices and the normalized

Figure 3. Temporal change in beta diversity and its components of riverine fish communities in 65 waterbodies across Sweden between 1990 and 2018. Beta diversity was defined as the compositional variation among multiple sampling sites within each waterbody. (a) Temporal changes in beta diversity based on species presence–absence and its extinction and colonization components. (b) Temporal changes in beta diversity based on species abundance and its loss and gain components. (c) Impacts of local extinctions and colonizations (i.e., changes from presence to absence and vice versa) of the six most abundant species (arranged in descending order) on beta diversity. (d) Impacts of abundance losses and gains of the six species on beta diversity. In the upper panels, the plus signs, horizontal lines, boxes, and circles indicate the means, medians, interquartile ranges (IQR), and outliers (values outside 1.5×IQR), respectively. In the bottom panels, bars and lines show the means ± standard errors. Asterisks indicate that the mean value was significantly different from zero (one-sample *t*-test; $P < 0.05$).

abundance-based Whittaker's beta diversity. A promising next step will be to extend the method to beta diversity based on Hill numbers (Hill, 1973; Jost, 2007; Chao & Chiu, 2016). Hill numbers link different lines of beta diversity research together and unify multiple dissimilarity measures into a common expression (Jost, 2007; Chao & Chiu, 2016; Chao et al., 2019). Temporal partitioning of Hill-number-based beta diversity could thus give us a synthetic understanding of community dynamics. Exploring the potential connections between our method and other partitioning methods that are based on Hill numbers (Godsoe et al., 2021, 2022) would also be an important step forward.

We expect our partitioning method to serve as a useful tool in both basic and applied ecology. Specifically, the capability of our method to quantify species-level processes could help conservation practitioners to assess the impacts of particular species on regional biodiversity (e.g., increased abundance of an invasive nonnative species and consequent decreases in endangered species). Empirical ecologists could use the homogenization and differentiation components of beta diversity to infer metacommunity processes and regional coexistence mechanisms. We believe that explicit analyses of the losses and gains in species abundances bring deeper insights into ecological community structure across space and time.

AUTHORS' CONTRIBUTIONS

ST conceived the study, derived the partitioning equations, and analyzed the data. ST wrote the manuscript with inputs from RI and MWC. All authors contributed to manuscript revision.

DATA ACCESSIBILITY

The R package ecopart ('Ecological COmmunity PARTitioning' or 'Extinction and COlonization PARTitioning') and an R script for extracting the Swedish fish community data from RivFishTIME (Comte et al., 2021) are available at GitHub (https://github.com/communityecologist). The ecopart package can be installed using the remotes package (Csardi et al., 2021): remotes::install_github("communityecologist/ecopart").

ACKNOWLEDGEMENTS

We thank William Godsoe and an anonymous reviewer for constructive comments and

people who contributed to the Swedish Electrofishing Register database (Sers, 2013). ST was supported by a JSPS Overseas Research Fellowship (No. 201860500) and a JSPS grant (No. 21K14880) from the Japan Society for the Promotion of Science. RI was supported by JSPS grants (No. 19K22457, 19K23768, and 20K15882).

REFERENCES

Almesjö, L., & Limén, H. (2009). Fish populations in Swedish waters: How are they influenced by fishing, eutrophication and contaminants? The Riksdag Printing Office.

Anderson, M. J., Crist, T. O., Chase, J. M., Vellend, M., Inouye, B. D., Freestone, A. L., … Swenson, N. G. (2011). Navigating the multiple meanings of beta diversity: A roadmap for the practicing ecologist. *Ecology Letters*, *14*, 19–28.

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology and Biogeography*, *19*, 134–143.

Baselga, A. (2013). Separating the two components of abundance-based dissimilarity: Balanced changes in abundance vs. abundance gradients. *Methods in Ecology and Evolution*, *4*, 552–557.

Baselga, A. (2017). Partitioning abundance-based multiple-site dissimilarity into components: Balanced variation in abundance and abundance gradients. *Methods in Ecology and Evolution*, *8*, 799–808.

Bray, J. R., & Curtis, J. T. (1957). An ordination of the upland forest communities of southern Wisconsin. *Ecological Monographs*, *27*, 325–349.

Cassey, P., Lockwood, J. L., Olden, J. D., & Blackburn, T. M. (2008). The varying role of population abundance in structuring indices of biotic homogenization. *Journal of Biogeography*, *35*, 884–892.

Chao, A., Chazdon, R. L., & Shen, T. J. (2005). A new statistical approach for assessing similarity of species composition with incidence and abundance data. *Ecology Letters*, *8*, 148–159.

Chao, A., & Chiu, C. H. (2016). Bridging the variance and diversity decomposition approaches to beta diversity via similarity and differentiation measures. *Methods in Ecology and Evolution*, *7*, 919–928.

Chao, A., Chiu, C., Wu, S., Huang, C., & Lin, Y. (2019). Comparing two classes of alpha diversities and their corresponding beta and (dis)similarity measures, with an

application to the Formosan sika deer *Cervus nippon taiouanus* reintroduction

programme. *Methods in Ecology and Evolution*, *10*, 1286–1297.

Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., & Ellison, A. M.

(2014). Rarefaction and extrapolation with Hill numbers: A framework for sampling

and estimation in species diversity studies. *Ecological Monographs*, *84*, 45–67.

Comte, L., Carvajal-Quintero, J., Tedesco, P. A., Giam, X., Brose, U., Erős, T., … Olden, J. D.

(2021). RivFishTIME: A global database of fish time-series to study global change

ecology in riverine systems. *Global Ecology and Biogeography*, *30*, 38–50.

Csardi, G., Hester, J., Wickham, H., Chang, W., Morgan, M., & Tenenbaum, D. (2021).

remotes: R package installation from remote repositories, including 'GitHub'. R package

version 2.4.2.

Godsoe, W., Bellingham, P. J., & Moltchanova, E. (2022). Disentangling niche theory and

beta diversity change. *The American Naturalist*, *199*, 510–522.

Godsoe, W., Eisen, K. E., Stanton, D., & Sirianni, K. M. (2021). Selection and biodiversity

change. *Theoretical Ecology*, *14*, 367–379.

Hill, M. O. (1973). Diversity and evenness: A unifying notation and its consequences.

Ecology, *54*, 427–432.

Jost, L. (2007). Partitioning diversity into independent alpha and beta components. *Ecology*,

88, 2427–2439.

Koleff, P., Gaston, K. J., & Lennon, J. J. (2003). Measuring beta diversity for presence-

absence data. *Journal of Animal Ecology*, *72*, 367–382.

Legendre, P., & Legendre, L. (2012). Numerical Ecology (3rd ed.). Elsevier.

Li, S., Cadotte, M. W., Meiners, S. J., Pu, Z., Fukami, T., & Jiang, L. (2016). Convergence and

divergence in a long-term old-field succession: The importance of spatial scale and

species abundance. *Ecology Letters*, *19*, 1101–1109.

McKinney, M. L., & Lockwood, J. L. (1999). Biotic homogenization: A few winners replacing

many losers in the next mass extinction. *Trends in Ecology & Evolution*, *14*, 450–453.

Olden, J. D., Comte, L., & Giam, X. (2018). The Homogocene: A research prospectus for the

study of biotic homogenisation. *NeoBiota*, *37*, 23–36.

Olden, J. D., & Poff, N. L. (2003). Toward a mechanistic understanding and prediction of

biotic homogenization. *The American Naturalist*, *162*, 442–460.

Petersen, K. N., Freeman, M. C., Kirsch, J. E., McLarney, W. O., Scott, M. C., & Wenger, S. J.

(2021). Mixed evidence for biotic homogenization of Southern Appalachian fish communities. *Canadian Journal of Fisheries and Aquatic Sciences*, *78*, 1397–1406.

Rosenblad, K. C., & Sax, D. F. (2017). A new framework for investigating biotic homogenization and exploring future trajectories: Oceanic island plant and bird assemblages as a case study. *Ecography*, *40*, 1040–1049.

Ružička, M. (1958). Anwendung mathematisch-statisticher Methoden in der Geobotanik (synthetische Bearbeitung von Aufnahmen). *Biológia, Bratislava*, *13*, 647–661.

Sers, B. (2013). Swedish Electrofishing RegiSter – SERS. Swedish University of Agricultural Sciences (SLU), Department of Aquatic Resources. Retrieved from http://www.slu.se/electrofishingdatabase (December 21, 2021).

Socolar, J. B., Gilroy, J. J., Kunin, W. E., & Edwards, D. P. (2016). How should beta-diversity inform biodiversity conservation? *Trends in Ecology & Evolution*, *31*, 67–80.

Tatsumi, S., Iritani, R., & Cadotte, M. W. (2021). Temporal changes in spatial variation: partitioning the extinction and colonisation components of beta diversity. *Ecology Letters*, *24*, 1063–1072.

Tatsumi, S., Strengbom, J., Čugunovs, M., & Kouki, J. (2020). Partitioning the colonization and extinction components of beta diversity across disturbance gradients. *Ecology*, *101*, e03183.

Whittaker, R. H. (1960). Vegetation of the Siskiyou Mountains, Oregon and California. *Ecological Monographs*, *30*, 279–338.