Title: Complete genomes of clade G6 Saccharibacteria suggest a divergent ecological

niche and lifestyle

Author: Jonathon L. Baker[1,*]
[1] Genomic Medicine Group
J. Craig Venter Institute
4120 Capricorn Lane
La Jolla, CA 92037
9

*Corresponding Author: JLB: jobaker@jcvi.org

ORCID: JLB: 0000-0001-5378-322X

Running title: Highly divergent Clade G6 Saccharibacteria

Keywords: Saccharibacteria, oral microbiome, TM7, nanopore sequencing

ABSTRACT

Saccharibacteria (formerly TM7) have reduced genomes, a small size, and appear to have a

parasitic lifestyle dependent on a bacterial host. Although there are at least 6 major clades of

Saccharibacteria inhabiting the human oral cavity, cultured isolates or complete genomes of oral

Saccharibacteria have been previously limited to the G1 clade. In this study, nanopore

sequencing was used to obtain three complete genome sequences from clade G6. Phylogenetic

analysis suggested the presence of at least 3-5 distinct species within G6, with two discrete taxa

represented by the 3 complete genomes. G6 Saccharibacteria were highly divergent from the

more well-studied clade G1, and had the smallest genomes and lowest GC-content of all

Saccharibacteria.   Pangenome analysis showed that although 97% of shared pan-

Saccharibacteria core genes and 89% of G1-specific Core Genes had putative functions, only

50% of the 244 G6-specific Core Genes had putative functions, highlighting the novelty of this

group. Compared to G1, G6 encoded divergent metabolic pathways. G6 genomes lacked an

F1F0 ATPase, the pentose phosphate pathway, and several genes involved in nucleotide

metabolism, which were all core genes for G1. G6 genomes were also unique compared to G1

in that they encoded lactate dehydrogenase, adenylate cyclase, limited glycerolipid metabolism,

a homolog to a lipoarabinomannan biosynthesis enzyme, and the means to degrade starch.

These differences at key metabolic steps suggest a distinct lifestyle and ecological niche for clade

G6, possibly with alternative hosts and/or host-dependencies, which would have significant

ecological, evolutionary, and likely pathogenic, implications.

IMPORTANCE

Saccharibacteria are ultrasmall, parasitic bacteria that are common members of the oral

microbiota and have been increasingly linked to disease and inflammation. However, the lifestyle

and impact on human health of Saccharibacteria remains poorly understood, especially for the 5

clades (G2-G6) with no complete genomes or cultured isolates. Obtaining complete genomes is

of particular importance for Saccharibacteria, because they lack many of the "essential" core

genes used for determining draft genome completeness and few references exist outside of clade

G1. In this study, complete genomes of 3 G6 strains, representing two candidate species, were

obtained and analyzed. The G6 genomes were highly divergent from G1, and enigmatic, with

50% of the G6 core genes having no putative functions. The significant difference in encoded

functional pathways is suggestive of a distinct lifestyle and ecological niche, probably with

alternative hosts and/or host-dependencies, which would have major implications in ecology,

evolution, and pathogenesis.

OBSERVATION

Saccharibacteria (formerly TM7) have an ultrasmall cell size, reduced genomes, and are thought to be obligate epibionts, dependent on physically-associated host species (1-3). Common constituents of the oral microbiota, Saccharibacteria have been increasingly linked to inflammation and disease (4-6). Saccharibacteria contains at least 6 distinct clades (G1-G6)(7, 8), however all currently available human-associated complete genomes and cultured isolates belong to clade G1, leaving clades G2-G6 quite poorly understood. Several recent publications have provided the first draft genomes from clades G3, G5, and G6 (4, 8-11). Obtaining complete genomes is of particular importance for Saccharibacteria, because they lack many of the "essential" single-copy core genes that are typically used to estimate genome completion, as well as complete reference genomes outside of the G1 clade.

A recent, short-read-based oral microbiome study provided 21 Saccharibacteria draft genomes from clades G1, G3, and G6 (4), with several being high quality (high N50, relatively contiguous, low predicted contamination). Therefore, nanopore sequencing of the same saliva samples that had produced the draft genomes, followed by long-read and/or hybrid assembly, was used to improve these genomes, resulting in 3 complete, circular G6 genomes: JB001 (662,051 bp), JB002 (639,751 bp), and JB003 (663,165 bp). Table 1 is a summary of the genomes improved during this study and the Supplemental Methods contain a full description of the DNA extraction, sequencing, assembly, and analysis methods. These methods are a modified version of a previously reported protocol (Baker 2021, in-press). Although the G1 and G3 "near complete" improved genomes that were obtained are useful in their own right, they are still incomplete, and/or may contain contamination, therefore the 3 complete G6 genomes are the focus of this report, and the near complete genomes are briefly discussed in the Supplemental Methods.

Phylogenetic analysis using concatenated protein sequences was performed using Anvi'o (12), and included the 8 improved/completed genomes from this study, all 26 complete

Saccharibacteria genomes available on NCBI (as of 1 April 2021), and 90 Saccharibacteria draft

genomes from 5 recent studies (Table S1). JB001, JB002, and JB003 were indeed members of

Saccharibacteria clade G6 (Figure 1A, Figure S1), and represent the only human-associated,

complete Saccharibacteria genomes outside of clade G1. Notably, G6 had the smallest genomes

and the lowest GC-content of all Saccharibacteria (Figure 1A). Percent average nucleotide

identity (ANI) between the G6 genomes was calculated using Anvi'o and suggested that there are

at least 3-5 distinct species within the clade (Figure 1B; a cutoff of 95% ANI is frequently used to

estimate the species level (13, 14)). JB001, JB003, JCVI_1_bin.12, and G6_32_bin_33_unicycler

appear to be the same species, with an ANI of ≥95%, despite their source from different human

subjects and independent genome assembly (Figure 1B). JB002 and T-C-M-Bin-00022 were

over 98% ANI, likely representing the same distinct species, while CMJM-G6-HOT-870 and T-C-

M-Bin-00011 were ~98% ANI and formed what is likely an additional G6 species (Figure 1B).

CLC Genomics Workbench was used to perform whole genome alignment for JB001, JB002,

JB003, and the G1 reference strain, TM7x (Figure 1C). While JB001 and JB003 were completely

syntenic, and there were moderate differences between JB001/JB003 and JB002, TM7x and the

G6 Saccharibacteria have undergone many genomic re-arrangements and instances of gene

gain/loss since their last common ancestor (Figure 1C).

To examine functional and metabolic differences between the G6 clade and the more well-

understood G1 clade, pangenome analysis was performed using Anvi'o (15) on the 3 complete

G6 genomes and 4 diverse G1 complete genomes (Figure 2, Table S3). This identified 223 "pan-

Saccharibacteria Core Genes" appearing in all genomes, as well as all 94 "G1 Core Genes", and

244 "G6 Core Genes" (Figure 2A). While 97% of the pan-Saccharibacteria Core Genes and 89%

of the G1 Core Genes had known COG functions and pathways, only 50% of the G6 Core Genes

had known COG functions and pathways (Figure 2A), highlighting the enigmatic nature of this

clade. The likely reason for the lower number of G1 core genes is the larger amount of known

diversity within the G1 clade and the genomes analyzed here (8, 9), leading to less conservation

across the G1 pangenome. A larger pangenome analysis, examining all 11 G6 genomes and 14

diverse G1 genomes is available in Figure S2 and Table S4. This generated similar results, but

note that this analysis contains incomplete draft genomes which are incomplete and/or may

contain contamination. A complete metabolic network illustrating the known KEGG pathways

identified in the three sets of core genes identified in Figure 2A is shown in Figure 2B. Both G1

and G6 genomes encode partial cell wall metabolism, glycolysis (missing phosphofructokinase),

and arginine biosynthesis pathways, and do not encode fatty acid metabolism, a TCA cycle, or

amino acid metabolism (other than arginine) (Figure 2B). Notable pathways present in G6

genomes but absent in G1 include: maltase glucoamylase (to metabolize starch), fructose

bisphosphate aldolase (a glycolytic step), adenylate cyclase, lactate dehydrogenase, partial

lipoarabinomannan (LAM) biosynthesis, and partial glycerolipid metabolism. Conversely, G1

genomes encode the non-oxidative phase of the pentose phosphate pathway, an F1F0 ATPase,

alpha galactosidase, and several steps in nucleotide metabolism, which were not present in the

G6 genomes (Figure 2B). Between JB001 and JB002, most differences were genes with

unknown functions, therefore the differences in the KEGG pathways encoded were minor (Figure

S3). The G6 genomes examined did not contain predicted elements of a CRISPR system.

Although it is not known how Saccharibacteria obtain needed metabolites from the host, a type

IV pilus-like system is generally well-conserved across the group, has been proposed as a

candidate mechanism (8, 9), and was present in the G6 genomes here. The species-level clade

that included JB001 and JB003 encoded a ~10,000bp putative prophage element, which was

flanked by homologs to the PinE invertase and contained a T4SS VirD4 homolog and 4

hypothetical proteins, all with ~95% homology to a similar region in *Streptococcus salivarius.*

Taken together, these analyses indicate that Saccharibacteria clade G6 is highly divergent

from clade G1, and may have a different lifestyle, host, and host-dependencies. This is in line

with the recent hypothesis that G6 reside on the tongue (G6 are referred to as 'T2' in reference

9) and have a long history of association with animal hosts, while G1 reside in dental plaque and

were a much more recent acquisition from the environment (8, 9). Interestingly, the species-level

clade containing JB002 (the most reduced Saccharibacteria genome, with only 615 genes) was

the only Saccharibacteria group that resided both on the tongue and in dental plaque (9).

Although all cultured isolates of Saccharibacteria were epibionts of *Actinomyces* spp., they were

all G1 strains. Residing in a different environment, G6 may have distinct host species, possibly

Streptococcus, given the acquired homologous sequence. It is likely that G6 has fallen into the

'unknown' taxonomic bucket in the majority of past microbiome studies, thus the role of G6 in

human health remains to be elucidated. The high percentage of genes with unknown functions

further adds to the obscurity of this clade. Overall, this article highlights an urgent need for study

of Saccharibacteria, since almost nothing is known about the lifestyle, host, or ecological impact

of Saccharibacteria clade G6, and even less still is understood about clades G2, G3, G4, and G5.

ACKNOWLEDGEMENTS

I thank Karrie Goglin-Almeida, Jelena Jablanovic, and Kara Riggsbee for performing the library

preparation and sequencing, and Jeffrey S. McLean for helpful discussions. This research was

supported by NIH/NIDCR K99-DE029228.

DATA AVAILABILITY

The complete genome sequences of JB001, JB002, and JB003 have been deposited in GenBank

under the accession numbers: CP072208, CP076101, and CP076102. The BioProject accession

for this project is PRJNA624185. The short reads used to generate the assemblies are available

in the SRA database with the accession numbers SRX4318838, SRX4318837, and SRX4318835.

The long reads used to generate the assemblies are available in the SRA dataset with the

accession numbers SRX10387815, SRX11020560 and SRX11020561.

REFERENCES

1. Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R, Thomas BC, Banfield JF. 2016. A new view of the tree of life. Nat Microbiol 1:16048.
2. Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, Wilkins MJ, Wrighton KC, Williams KH, Banfield JF. 2015. Unusual biology across a group comprising more than 15% of domain Bacteria. Nature 523:208-11.
3. He X, McLean JS, Edlund A, Yooseph S, Hall AP, Liu SY, Dorrestein PC, Esquenazi E, Hunter RC, Cheng G, Nelson KE, Lux R, Shi W. 2015. Cultivation of a human-associated TM7 phylotype reveals a reduced genome and epibiotic parasitic lifestyle. Proc Natl Acad Sci U S A 112:244-9.
4. Baker JL, Morton JT, Dinis M, Alvarez R, Tran NC, Knight R, Edlund A. 2021. Deep metagenomics examines the oral microbiome during dental caries, revealing novel taxa and co-occurrences with host molecules. Genome Res 31:64-74.
5. Abu Fanas S, Brigi C, Varma SR, Desai V, Senok A, D'Souza J. 2021. The prevalence of novel periodontal pathogens and bacterial complexes in Stage II generalized periodontitis based on 16S rRNA next generation sequencing. J Appl Oral Sci 29:e20200787.
6. Bor B, Bedree JK, Shi W, McLean JS, He X. 2019. Saccharibacteria (TM7) in the Human Oral Microbiome. J Dent Res 98:500-509.
7. Camanocha A, Dewhirst FE. 2014. Host-associated bacterial taxa from Chlorobi, Chloroflexi, GN02, Synergistetes, SR1, TM7, and WPS-2 Phyla/candidate divisions. J Oral Microbiol 6.
8. McLean JS, Bor B, Kerns KA, Liu Q, To TT, Solden L, Hendrickson EL, Wrighton K, Shi W, He X. 2020. Acquisition and Adaptation of Ultra-small Parasitic Reduced Genome Bacteria to Mammalian Hosts. Cell Rep 32:107939.
9. Shaiber A, Willis AD, Delmont TO, Roux S, Chen LX, Schmid AC, Yousef M, Watson AR, Lolans K, Esen OC, Lee STM, Downey N, Morrison HG, Dewhirst FE, Mark Welch JL, Eren AM. 2020. Functional and genetic markers of niche partitioning among enigmatic members of the human oral microbiome. Genome Biol 21:292.
10. Cross KL, Campbell JH, Balachandran M, Campbell AG, Cooper SJ, Griffen A, Heaton M, Joshi S, Klingeman D, Leys E, Yang Z, Parks JM, Podar M. 2019. Targeted isolation and cultivation of uncultivated bacteria by reverse genomics. Nat Biotechnol 37:1314-1321.
11. Lamont EI, Gadkari A, Kerns KA, To TT, Daubert D, Kotsakis G, Bor B, He X, McLean JS. 2021. Modified SHI medium supports growth of a disease-state subgingival polymicrobial community in vitro. Mol Oral Microbiol 36:37-49.
12. Eren AM, Esen OC, Quince C, Vineis JH, Morrison HG, Sogin ML, Delmont TO. 2015. Anvi'o: an advanced analysis and visualization platform for 'omics data. PeerJ 3:e1319.
13. Pasolli E, Asnicar F, Manara S, Zolfo M, Karcher N, Armanini F, Beghini F, Manghi P, Tett A, Ghensi P, Collado MC, Rice BL, DuLong C, Morgan XC, Golden CD, Quince C, Huttenhower C, Segata N. 2019. Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000 Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. Cell 176:649-662 e20.
14. Jain C, Rodriguez RL, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun 9:5114.
15. Delmont TO, Eren AM. 2018. Linking pangenomes and metagenomes: the Prochlorococcus metapangenome. PeerJ 6:e4320.

 FIGURE LEGENDS

Figure 1: JB001, JB002, and JB003 are clade G6 Saccharibacteria representing two

distinct species. (A) Phylogenetic tree of Saccharibacteria annotated with genome data.

Phylogenetic analysis of the 123 Saccharibacteria genomes listed in Table S1. Firmicutes was

used as an outgroup. The bars in the innermost layer represent the number of singleton gene

clusters (i.e. genes appearing in only that one genome) in each genome. The bars in the second

layer represent the redundancy (likely contamination) within each genome. The bars in the third

layer represent the %GC content of each genome. The bars in the fourth layer represent the total

length in bp of each genome. The fifth layer displays the source/reference for each genome. The

sixth layer displays the genomes that are complete. The outermost layer, and the color of the

branches of the tree, illustrate which Saccharibacteria clade each genome is part of. Orange

stars indicate genomes that were used in the full pangenome analysis (Figure S2, Table S4).

Yellow stars indicate genomes that were used in the pangenome analysis of compete genomes

only (Figure 2, Table S3) as well as the full pangenome analysis (Figure S2, Table S4). A larger

version of this figure, with the name of each genome labeled, is available in Figure S1. Note that

CP025011_1_Candidatus_Saccharibacteria_bacterium_YM_S32_TM7_50_20_chromosome_c

omplete_genome and c_000000000001 (GCA_003516025.1_ASM351602v1_genomic.fa), the

only two complete genomes in clades G3 and G5, are from environmental, not oral, samples. The

raw data in the annotations of the tree is available in Table S1. **(B) Average nucleotide identity**

(%ANI) of G6 genomes. Heatmap of all-vs-all comparison of %ANI of all 11 G6 genomes. The

tree on the right is a scaled up version of the G6 portion of the phylogenetic tree in panel A. Full

percentage identity, which takes alignment length into account, is available in Table S2. **(C)**

Whole genome alignment of TM7x vs complete G6 genomes. Whole genome alignment

diagram produced by CLC Genomics Workbench. The tree on the right is based on the whole

genome alignment itself.

Figure 1

A Updated Saccharibacteria phylogeny

Complete?
Reference
Total length
GC-content
Redundancy
Singleton gene clusters

Reference
- Shaiber et al., 2020 (43)
- NCBI Complete genomes (26)
- Cross et al., 2019 (23)
- Baker et al. 2021 (16)
- McLean et al., 2020 (7)
- this study (6)
- Lamont et al., 2021 (1)

G3
G5
G6
JB002
JB001
JB003
G1
Firmicutes Outgroup
TM7x

★ Used in full pangenome
★ Used in full pangenome and complete genomes only pangenome

B Average Nucleotide Identity (%ANI) of G6 genomes

ORAL_T-A-F_MAG_00016	100.0%	85.4%	86.1%	86.1%	86.1%	93.8%	84.9%	85.8%	82.2%	72.9%	72.8%
G6_32_bin_33_unicycler	85.8%	100.0%	94.7%	94.4%	94.4%	85.1%	80.4%	80.3%	80.2%	72.4%	72.3%
FGB1_strain_JCVI_1_bin_12	86.4%	94.8%	100.0%	96.2%	96.3%	85.6%	80.2%	80.6%	80.3%	73.2%	72.9%
JB003_complete_genome	86.4%	94.6%	96.3%	100.0%	99.8%	85.8%	80.3%	81.2%	80.2%	73.0%	73.5%
GCA_015257795_JB001	86.6%	94.5%	96.4%	99.7%	100.0%	86.1%	80.4%	81.0%	80.2%	72.8%	72.9%
ORAL_T-B-F_Bin_00014	93.6%	85.3%	85.8%	85.9%	85.9%	100.0%	85.2%	85.9%	82.2%	72.4%	72.3%
ORAL_T-C-M_Bin_00022	85.8%	80.7%	80.8%	80.9%	80.9%	85.7%	100.0%	96.7%	81.7%	72.9%	73.0%
JB002_complete_genome	85.7%	80.3%	80.7%	81.2%	81.0%	85.8%	96.6%	100.0%	81.7%	72.8%	73.6%
ORAL_T-B-F_Bin_00010	82.6%	80.0%	80.5%	80.5%	80.5%	82.3%	81.4%	81.8%	100.0%	72.6%	72.5%
ORAL_T-C-M_Bin_00011	73.0%	72.1%	72.9%	73.0%	73.0%	72.2%	72.1%	72.8%	72.7%	100.0%	98.1%
TM7_CMJM_G6_1_HOT_870	72.4%	72.0%	72.6%	72.8%	72.6%	72.0%	72.2%	73.1%	72.6%	98.0%	100.0%

C Whole genome alignment of TM7x vs complete G6 genomes

TM7x
JB001
JB003
JB002

Figure 2: Pangenome analysis of complete genomes in Saccharibacteria clade G1 vs.
clade G6 identifies core genes with encoding distinct functional pathways. (A) The
pangenome of complete G1 and G6 genomes. The dendrogram in the center organizes the
2,279 gene clusters identified across in the genomes represented by the innermost 7 layers:
TM7x, BB001, HB001, PM004, JB003, JB001, and JB002. The data points within these 7 layers
indicate the presence of a gene cluster in a given genome. From inside to outside, the next 6
layers indicate known vs unknown COG category, COG function, COG pathway, KEGG class,
KEGG module, and KOfam. The next layer indicates single-copy pan-Saccharibacteria core
genes. The next 6 layers indicate the combined homogeneity index, functional homogeneity
index, geometric homogeneity index, max number of paralogs, number of genes in the gene
cluster, and the number of contributing genomes. The outermost layer highlights gene clusters
that correspond to the pan-Saccharibacteria Core Genes (found in all 7 genomes), the G1 Core
Genes (found in all G1 genomes and no G6 genomes), and the G6 Core Genes (found in all G6,
but no G1 genomes). The pie chart adjacent to each group of core genes indicates the breakdown
of COG categories of the gene clusters in the group. The 7 genome layers are ordered based on
the tree of the %ANI comparison, which is displayed with the red and white heatmap. The layers
underneath the %ANI heatmap, from top to bottom, indicate: the number of gene clusters, the
number of singleton gene clusters, the GC-content, and the total length of each genome. The
Venn diagrams in the inset show the number of overlapping and non-overlapping genes between
JB001 and JB002, and JB001 and TM7x. The number in parenthesis is the number of genes with
unknown functions (UF). **(B). KEGG pathways encoded by G1 and G6 core genes.** KEGG
metabolic map overlaid with the pathways encoded by the pan-Saccharibacteria core genes
(black), G1 Core Genes (green), and G6 Core Genes (red), as indicated by the Venn diagram
key. Enzymes of interest are labeled with text and arrows. Pathways are indicated by labeled
boxes, the cell wall metabolism pathways is labeled with the red background to distinguish it due
to the odd shape and overlap with the glycolysis pathway space.

Figure 2

A

Pangenome of complete G1 and G6 genomes

244 G6 Core Genes

94 G1 Core Genes

223 pan-Saccharibacteria Core Genes

B

KEGG pathways encoded by G1 and G6 core genes

- G1 Core Genes — green
- pan-Saccharibacteria Core Genes — black
- G6 Core Genes — red

Cell wall metabolism

Adenylate cyclase

Pyrimidine metabolism

Purine metabolism

Alpha galactosidase

Glycine hydroxymethyltransferase (GlyA)

LAM biosynthesis

Maltase glucoamylase

Glucose-6-phosphate dehydrogenase

Glycerolipid metabolism

Pentose phosphate pathway

Cysteine desulfurase (IscS)

Fructose bisphosphate aldolase

Glycolysis

Terpenoid biosynthesis

Lactate dehydrogenase (Ldh)

Glycine hydroxymethyltransferase (GlyA)

Arginine and glutamate metabolism

Succinyl-diaminopimelate desuccinylase (DapE)

Other sugar metabolism

F1F0 ATPase

Table 1. Saccharibacteria genomes improved using nanopore sequencing in this study

New MAG designation	Previous MAG name	previous # of contigs	previous size (bp)	updated # of contigs	updated size (bp)	updated longest contig (bp)	complete	near complete (longest contig > 700,000 bp or < 5 contigs)
JB001	Candidatus_Nanogingivalaceae_FGB1_strain_JCVI_27_bin.3	67	704,215	1	662,051	662,051	*	
JB002	Candidatus_Saccharimonas_sp._strain_JCVI_32_bin.49	14	620,057	1	639,737	639,737	*	
JB003	Candidatus_Nanogingivalaceae_FGB1_strain_JCVI_28_bin.11	34	719,702	1	663,171	663,171	*	
TM7c-JB	Candidatus_Nanosynbacter_TM7c_strain_JCVI_32_bin.19	7	793,808	1	793,363	793,363		*
none	Candidatus_Nanosynbacter_sp._TM7_MAG_III_A_2_strain_JCVI_32_bin.12	76	696,341	8	837,467	808,188		*
none	Candidatus_Nanosynbacter_GGB2_strain_JCVI_32_bin.57	32	1,040,784	6	1,054,499	762,750		*
G6_32_bin_33_unicycler	Candidatus_Nanogingivalaceae_FGB1_strain_JCVI_32_bin.33	97	521,278	31	594,688	77,761		
none	Candidatus_Nanosynbacteraceae_FGB1_strain_JCVI_32_bin.22	68	636,728	35	913,508	182,700		
none	Candidatus_Nanosynbacteraceae_FGB2_strain_JCVI_32_bin.44	31	725,781	15	819,428	300,554		
G3_32_bin_36_unicycler	Candidatus_Nanosyncoccus_FGB2_strain_JCVI_32_bin.36	32	667,180	4	688,219	265,262		*