Classification, Using Field Potential Images, of Pacemaker Micro-coordination in the Mouse Intestine

Naoko Iwata[1], Chiho Takai[1], Naoto Mochizuki[1], Mariko Yamauchi[1], Noriyuki Kaji[2]

Yoshiyuki Kasahara[3], Shinsuke Nakayama[1,*]

[1]Department of Cell Physiology, Nagoya University Graduate School of Medicine, Nagoya 466-8550, Japan

[2]Department of Veterinary Pharmacology, Azabu University School of Veterinary Medicine, Sagamihara 252-5201, Japan

[3]Department of Fetal and Maternal Therapeutics, Tohoku University Graduate School of Medicine, Sendai 980-8575, Japan

Corresponding author: Shinsuke Nakayama*

Department of Cell Physiology, Nagoya University Graduate School of Medicine,

65 Tsurumai-cho, Showa-ku, Nagoya 466-8550, Japan

Phone: +81 52 744 2045   Fax: +81 52 744 2048

E-mail: h44673a@nucc.cc.nagoya-u.ac.jp

ORCID: 0000-0003-4933-5429

Running title: MEA Classification of Pacemaker Micro-coordination

Abstract:

The flexible and sophisticated movement of the gastrointestinal (GI) tract implies the involvement of numerous mechanisms that coordinate excitation in micro-regions other than the neural reflex. We thus performed image analysis of pacemaker micro-coordination in the small intestine of mice since it contains typical network-forming pacemaker cells. Field potentials were recorded by a dialysis membrane-reinforced microelectrode array (MEA). The micro-coordination of pacemaker activity was classified into basic patterns despite large variations. In the developmental process, pacemaker activity was categorized as either an 'expanding' or a 'migrating' pattern which was initiated in or propagated to the MEA sensing area, respectively. The intercellular current of the volume conductor complicated the waveform of both activities. The existence of 'expanding' and 'migrating' patterns were attributable to duplicated pacemaker systems such as intracellular Ca^{2+} oscillation-activated and voltage-gated mechanisms. Additionally, from the spatio-temporal feature during the period of pacemaker events, the 'bumpy/aberrant' pattern was defined by aberrant, incoherent propagation and was associated with local impairment of excitability, while the 'colliding/converging' pattern involved the interaction of multiple activities in the MEA area. Interconversion between the four micro-coordination patterns occurred even in the same MEA area. 5-Hydroxytryptamine (5-HT) promoted 'migrating' activity, by decreasing the 'expanding' activity, implying an improvement or restoration of spatial conductivity. These results agree well with the action of 5-HT to change GI movement toward propulsion. In conclusion, our MEA method of image classification enables the quantitative assessment of spatio-temporal electric coordination underlying GI motility, and its application to small model animals is anticipated.

Keywords: Classification, dialysis membrane, gut pacemaker, microelectrode array

Author contributions:
NI: Investigation, Data curation, Visualization, Validation, Writing – review & editing
CT: Investigation, Data curation, Visualization
MY: Investigation, Data curation.
NM: Investigation, Data curation, Writing – review & editing.
NK: Investigation, Data curation.
YK: Conceptualization, Validation.
SN: Conceptualization, Methodology, Investigation, Data curation, Validation, Funding acquisition, Supervision, Writing – original draft.

Introduction

The flexible and sophisticated movement of the gastrointestinal (GI) tract implies an underlying mechanism that variously coordinates spatio-temporal excitability in microregions, simultaneously operated under the reflex of intrinsic neural circuits (Spencer et al. 1999; Bayliss & Starling, 1899). The patterns of GI movement, especially in the small intestine, are often described using simple terms such as 'segmentation' and 'peristalsis' for agitation and transportation of the luminal contents, whereas, in reality, the GI tract exhibits smooth, elaborate and highly varied movements that show subtle changes between cycles, for example abdominal cine magnetic resonance imaging (Teramoto et al. 2012). Electrical excitation underlies numerous GI motilities (Szurszewski, 1987). In the majority of the tract, spontaneous electrical activity occurs, and originates from pacemaker interstitial cells, which are recognized for their crucial role in cooperating local activities through their network-like structure (Faussone-Pellegrini, 2005; Nakayama et al. 2006; Huizinga & Lammers, 2009).

Spatio-temporal investigation into the interaction of pacemaker potentials has not progressed much. Reflecting spontaneous pacemaker activity, intracellular microelectrodes exhibit relatively simple wave forms comprised of initial rapid depolarizing potential followed by a plateau phase of variable amplitude and duration (Sanders et al. 2016), but simultaneous intracellular recording with multiple microelectrodes requires a masterful performance. On the other hand, extracellular recordings exhibit a wide variety of waveforms such as biphasic and triphasic potentials that may originate from recording and data processing techniques, or other sources, in addition to variations of active excitable cells under different conditions, for example, action potentials from smooth muscle cells in the absence of Ca^{2+} channel antagonists (Iwata et al. 2017). In terms of the interrelationship between these electric recordings, it has been pointed out that the waveform of extracellular field potential is theoretically proportional to the first-time derivative of the transmembrane voltage, but in practical measurements, this relationship is apparently more complicated (Sanders et al. 2016; Huizinga, 2017; O'Grady et al. 2017).

In this study, we provide an image analysis of pacemaker field potentials measured using a low-impedance microelectrode array (MEA) reinforced with dialysis membranes. This method enabled field potentials over a wide range of frequencies to be stably measured in microregions. In the small intestine of mice, despite huge variations, field potential images were classified into two major micro-coordination patterns, 'expanding' and 'migrating', depending on the developmental process and whether pacemaker activity was initiated in or migrated to the MEA area. Particularly in 'migrating' activities, a characteristic feature is that positive potentials precede negative potentials, providing a rationale for the complicated waveforms of extracellular recordings in the GI tract. Namely, due to its syncytal nature (Manchanda et al. 2019; Sanders, 2019), intercellular pacemaker current also participates in the depolarization of the membrane during an early phase of propagation, and flows in the opposite direction against the transmembrane pacemaker current. In addition, 5-hydroxytryptamine (5-HT, serotonin), which is known to promote propulsive movements (Bülbring and Lin, 1958; Mawe and Hoffman, 2013), exerted consistent modulatory effects on micro-coordination patterns with its function *in vivo*, suggesting that this MEA technique is beneficial for functional assessment.

Materials and Methods

Animals and preparations

Animals were treated ethically in accordance with the guidelines for the proper conduct of animal experiments published by the Science Council of Japan. All procedures were approved by the Animal Care and Use Committee of Nagoya University Graduate School of Medicine (Permissions #28316, #29121, #30312 and #31199). C57BL/6J wild-type mice (8–40 weeks) and *W/W^v* (WBB6F1/Kit-*Kit^W/Kit^{W-v}*/Slc) mice, which lack ICCs in the intestine (8–12 weeks, Japan SLC, Hamamatsu, Japan), were sacrificed by cervical dislocation after the inhalation of carbon dioxide. The whole GI tract was quickly resected, and small pieces (~20 mm) of the ileum were isolated and cut along the mesentery. The entire muscularis propria containing the myenteric plexus was isolated by removing it from the mucous membrane using forceps.

Immunohistochemistry

Small segments of the isolated ileal musculature were fixed with ice-cold acetone for 10 min, then blocked with 2% bovine serum albumin for 1 h. Samples were incubated with anti-c-kit rat monoclonal antibody (1:500 v/v, ACK2, BioLegend, San Diego, CA, USA) overnight (4ºC), and subsequently stained with donkey anti-rat IgG antibody, with Alexa Fluor 488 conjugate (1:500 v/v, A-21208, Thermo Fisher Scientific, Waltham, MA, USA). Fluorescence images were acquired from 70 µm thick slices under a confocal microscope (TCS SP5 II, Leica Microsystems, Tokyo, Japan) using a ×20 objective lens.

Solutions and drugs

The extracellular solution used for the MEA recordings was a modified Krebs solution with the following composition (in mM): NaCl 125, KCl 5.9, $MgCl_2$ 1.2, $CaCl_2$ 2.4, glucose 11, and Tris-HEPES 11.8 (pH 7.4). Nifedipine, 5-HT hydrochloride, 2Me-5HT and ondansetron were purchased from Sigma-Aldrich (St Louis, MO, USA). Nifedipine (2 µM) was applied to block L-type voltage-gated Ca^{2+} channels and thereby suppress smooth muscle contraction.

Electrical recordings

We developed the dialysis membrane-reinforced MEA recording technique to stably monitor the micro-organization of ICC pacemaker activity in the GI tract of small animals (Iwata et al. 2017; Morishita et al. 2017) because many animal disease models are developed using small animals. In addition, there is no consensus for the application of multi-electrode array used previously to small animals (Lammers et al. 2008; O'Grady, 2012; Sanders et al. 2016; Huizinga, 2017; O'Grady et al. 2017).

An 8 × 8 array of microelectrodes (P515A, Alpha MED Scientific, Ibaraki, Japan) was used to measure electric activity in the ileum. The inter-electrode distance of the MEA was 150 µm, and the sensing area was 1050 × 1050 µm². Samples of mouse ileal muscle sheets were firmly mounted onto the MEA, with the longitudinal muscle layer facing downward, using a small piece of dialysis membrane (cellulose tube, Visking, Chicago, IL, USA) and a slice anchor (SDH series, Harvard Apparatus Japan, Tokyo, Japan). A large reference electrode made of platinum was immersed in the extracellular solution (Fig. 1A, B). During the propagation of pacemaker potentials, the membrane currents (I_M) flowing through the access resistance (R_a) between the sensing

and reference electrodes changed the field potential ($I_M \times R_a$) (Fig. 1B). A set of 8×8 field potentials originating from c-Kit–positive interstitial cells (Fig. 1C, D) was simultaneously recorded using a multi-channel AC amplifier that stabilized the baseline potential with high-pass filtering at 0.1 Hz and reduced noise with low-pass filtering at 10 kHz in accordance with conventional extracellular electrode recording techniques (Astrand et al. 1988). The arrayed data were digitized and stored by a personal computer using a 14-bit A/D converter with a sampling rate of 20 kHz (50-μs intervals). The dynamic range of A/D conversion was set to ± 1 mV with a digital resolution of ~0.12 μV.

For all samples, the oral and anal ends of the muscles were oriented toward the upper and lower ends of the MEA, respectively. The dialysis membrane had a molecular weight cut-off of 12,000, thus oxygen, ions and energy metabolites could be exchanged. The MEA recording chamber (~2 mL volume) was placed on a heater maintained at ~34ºC. The samples were incubated at least 20 min before MEA recordings. Nifedipine (2 μM), a dihydropyridine Ca^{2+} channel antagonist, was added to the extracellular medium to selectively block L-type voltage-gated Ca^{2+} channels (a major Ca^{2+} influx pathway in GI smooth muscle) (Akbarali et al. 2010) and suppress smooth muscle contraction without impeding the pacemaker activity of ICCs (Kito and Suzuki, 2003). The above procedures stabilized the electric conditions to enable long-term recordings.

Each sensing microelectrode of the MEA (P515A) had a square profile with a side length of ~50 μm and was made from deposited platinum black nanoparticles, which increased the surface area by ~200-fold (equivalent to 0.5 mm^2) (Fig. 1B). From the capacitance (C_{ME}: 0.052 μF) and resistance (R_{ME}: 15 kΩ) of the sensing microelectrode, the estimated impedance was sufficiently low to follow a wide frequency range of electric signals (Taniguchi et al. 2013; Iwata et al. 2017). For instance, impedance (Z_{ME}) of the sensing electrode at 0.1 Hz [~31 MΩ = $\sqrt{1/(2\pi \times 0.1 \text{ Hz} \times 0.052 \text{ μF})^2 + (15 \text{ kΩ})^2}$] was sufficiently small compared with the input resistance of the multi-channel amplifier used in this study (100 MΩ). Thus, the efficacy of electric signal transmission (Tr) was estimated to be ~95% at 0.1 Hz [100 MΩ / $\sqrt{(100 \text{ MΩ})^2 + (31 \text{ MΩ})^2}$].

Field potential data processing
The arrayed field potential data were acquired at a 14-bit depth with a sampling interval of 50 μs. The digital resolution was reduced to 0.5–25 ms to display the time course of the pacemaker oscillations. Data processing, including digital filtering and linear spectrum analysis, was performed using commercial software (Kaiseki, Kyowa Electronic Instruments, Tokyo, Japan; Mobius, Alpha Med Scientific, Ibaraki, Japan; MATLAB software package, MathWorks, Natick, MA, USA).

For potential mapping (Fig. 1E), after reducing the digital resolution to 5 ms, field potential data were processed by a band-pass filter at 0.25–10.5 Hz. Numerous factors, including variations in R_{ME} and R_a and differences in I_M between different populations of pacemaker and modulator cells, were considered to affect the magnitude of the oscillation in field potential. Therefore, to compensate for these factors and reconstitute a smooth field potential map, the amplitude of the field potential recorded in the n[th] microelectrode region [ME(n)] was corrected by a normalizing factor [F_N(n)]:

$LS_{0.25-10.5}$(n)/sum[$LS_{0.25-10.5}$(n = 1 to 64)]/64,

where $LS_{0.25-10.5}(n)$ is the density of the linear spectrum in the frequency range between 0.25–10.5 Hz for the field potential data at ME(n). To avoid excessive compensation, $F_N(n)$ was limited to within 1.0 ± 0.5. The sum of the $LS_{0.25-10.5}$ values for all 64 MEs was used to represent the magnitude of pacemaker activity. Subsequently, the 8×8 field potentials were interpolated with 50 points using a 'cubic' protocol, and each point of the field potential map was displayed as a 'hot' map (MATLAB software package) (Morishita et al. 2017).

The series (video) of field potential images (for 54 – 90 s) were used to classify the spatio-temporal features of pacemaker potentials at activity and event levels. First, each pacemaker activity was classified into 'migrating' or 'expanding' activity from the developing process, depending on whether each activity migrated as a volume conductor from adjacent regions outside of the MEA sensing area, or was initiated in the MEA sensing area (1.05×1.05 mm^2) expanding toward adjacent microregions (Fig. 1F; Supplemental Fig. S1). It is therefore considered that the ratio of 'migrating' to 'expanding' activity ($R_{M/E}$) can approximately estimate the area covered by each initiating site of pacemaker activity (A_{iPA}): MEA sensing area $\times$ ($R_{M/E}$ + 1) (mm^2).

The subclasses of 'migrating' and 'expanding' activities were defined by additional features during the pacemaker event. Irrespective of the developing process ('migrating' or 'expanding'), the total event of pacemaker activity showing aberrant (extraordinary incoherent) propagation in the MEA area was defined as a 'bumpy/aberrant' pattern event. Likewise, when more than two activities independently occurred and interacted, the event was defined as 'colliding/convergence' pattern event. 'Migrating' or 'expanding' activities without aberrant propagation were classified as 'migrating' or 'expanding' events, respectively. Of note, any activity was categorized as 'migrating' and 'expanding' in the activity level classification because the developmental process outside (migrating) or inside (expanding) of the MEA area is alternative. On the other hand, subclass patterns may increase in a future study when characteristic coordination features are identified by using image recognition technology.

Each activity observed in 'bumpy/aberrant' and 'colliding/convergence' events was counted as a 'migrating' or an 'expanding' activity in the activity statistics. For example, in a 'colliding' event of one pacemaker activity migrating from an adjacent region and another activity developing in the MEA region, one 'migrating' activity and one 'expanding' activity were counted (Fig. 4G).

Motion tracking

To assess the mechanical influence on electrical recording, transmission images were acquired by a video camera system in an inverted microscope. After MEA measurement of pacemaker potentials were performed in the presence of nifedipine (2 μM), dialysis membrane-reinforced ileal muscle sheets were placed in a glass-bottomed chamber containing the same extracellular solution. The glass-bottomed chamber was placed on a heating glass (34℃), and transparent images were acquired. Measurement of transverse (X) and vertical (Y) shifts of a square tracking region from its initial position using motion tracking software (2D-PTV, Digimo, Osaka, Japan) showed negligible movement (Supplemental Fig. S2).

Statistics

Numerical data are expressed as the mean ± S.D. Comparisons of normally distributed datasets were made to estimate the probability (*P*) of significant differences using paired *t*-tests ($P < 0.05$). In the control condition, a comparison between the first and second half of each MEA measurement was made in 10 muscle samples (8-36-weeks-old mice). The effect of 5-HT was examined in nine mice (in 8-18-weeks old).

Results

Patterns of pacemaker micro-coordination

Our dialysis membrane-reinforced low-impedance MEA technique enabled electric activity in the GI tract of small animal models to be stably monitored. Smooth muscle sheets isolated from mouse ileum were mounted on an 8×8 MEA with an interpolar distance of 150 μm and a sensing region of ~1 mm^2. Each smooth muscle sheet was mounted beneath a piece of dialysis membrane that maintained the conditions required for electric recording, such as access resistance (R_a) between the sensing and reference electrodes and isolation resistance between the sensing electrodes, while allowing the exchange of ions and metabolites needed for spontaneous electric activity (Fig. 1A,B). A dihydropyridine Ca^{2+} channel antagonist (nifedipine) was added to suppress smooth muscle contraction (Supplemental Fig. S2). The MEA detected field potential changes generated by the pacemaker activity of network-forming ICCs (Fig. 1C,D). This was confirmed by the MEA measurement in *W/W^v* mice that lack network-forming ICCs in the ileum (Supplemental Fig. S3).

The pacemaker potentials recorded by the 8×8 MEA were reconstructed into pseudo-color potential images (Fig. 1E) in which positive and negative field potentials, reflecting outward and inward currents, were assigned dark and light colors, respectively. Visualization of the MEA data enabled us to distinguish spatio-temporal features (Fig. 1F; Supplemental Fig. S1; also see Supplemental Videos SV1–4).

In the developmental process, pacemaker activities were divided into two patterns. 'Expanding' activity was defined by the initiation of activity in the MEA sensing area and a subsequent progressive expansion of the active region while the activity of the initiating region gradually subsided. As a result, 'expanding' activity typically formed an annular shape during the late phase (Expanding in Fig. 1F; Supplemental Video SV1). On the other hand, 'migrating' activity was defined by pacemaker activity migrating from a region adjacent to the MEA sensing area, and was manifested by an elongated active (light-colored) region that propagated approximately orthogonally to its long axis over the MEA sensing area in typical activities (Migrating in Fig. 1F; Supplemental Video SV2). Another characteristic feature of typical 'migrating' activity was a dark-colored area (indicating a transient positive potential) preceding the light-colored active area (further explanation is provided below).

A subclass of the micro-coordination pattern was defined by additional features during the propagating period of pacemaker events, independent of whether activity initiated in or migrated to the MEA sensing area. A 'bumpy/aberrant' event was due to aberrant spatio-temporal propagations resulting in a series of potential images lacking regularity, suggesting spatially confined electric coupling. In the example (Bumpy/Aberrant event in Fig. 1F; Supplemental Video SV3), spontaneous activity was initiated in the upper-left region and propagated toward the lower-right region while avoiding the lower-left region. Another subclass of pacemaker events was termed the 'colliding/converging' pattern, in which multiple propagating pacemaker potentials interacted in the MEA sensing area. Namely, they collided and/or converged to either annihilate each other (Colliding in Fig. 1F; Supplemental Movie SV4) or merge and propagate in a different direction (explained later). This pattern of pacemaker event contained multiple activities which either initiated in or propagated to the MEA sensing area.

In 10 ileal muscle sheets obtained from wild-type mice, the frequency of pacemaker events ranged from 10–31 cycles per min (min^{-1}), agreeing well with the values reported for ICCs in the myenteric plexus region (Kito and Suzuki, 2003). To verify the stability of the electric recording and biological conditions, features were compared between the first and second halves of the MEA measurement period (Fig. 1G). The frequency of pacemaker events and the occurrence of two basic activities displayed little change between the two halves (pacemaker event: 23.7 ± 5.6 vs. 25.2 ± 3.3 min^{-1}, $P = 0.234$; expanding activity: 10.5 ± 9.7 vs. 12.7 ± 10.5 min^{-1}, $P = 0.307$; migrating activity: 19.3 ± 14.1 vs. 18.2 ± 15.7 min^{-1}, $P = 0.376$, $N = 10$). Of note, each pacemaker activity in 'colliding/convergence' and 'bumpy/aberrant' pacemaker events was summed as either one of the two basic activities ('expanding' or 'migrating' activity in Fig. 1H) by judging from the developmental process whether activity initiated in or migrated to the MEA sensing area.

Variations in the micro-coordination of pacemaker activity
Visualization of the MEA field potential data enabled us to realize considerably large variations of spatio-temporal pacemaker activity. Even in the same muscle sample, the initiating region of 'expanding' activity fluctuated in the MEA sensing area (Fig. 2A). In this sample, a continuous recording demonstrated that most cases of propagating activity were of the 'expanding' type (12 out of 13 cycles), including the first three activities (in field potential traces in Fig. 2A). The first event was initiated in the upper-left region and propagated to adjacent regions (Expanding-1), the second event was initiated in the lower-right region (Expanding-2), and the third event was initiated near the center (Expanding-3) before expanding in an annular pattern (Supplemental Video SV5).

'Migrating' activity exhibited variations in the direction of propagation (Fig. 2B). Notably, longitudinal propagation of activity in these two samples occurred in opposite directions, namely in the aboral (oral-to-anal, ordinally transporting) direction (Migrating-L1) and oral direction (Migrating-L2). In addition, pacemaker activity reversed in the circular (transverse) direction (Migrating-C1 and C2 in Supplemental Fig. S4), and propagated diagonally (Migrating-D1; Migrating-D2 in Supplemental Fig. S4). The finding that 'migrating' activity propagated in various directions rather than following the course of muscle layers and enteric innervation suggests a major contribution of the ICC network (Fig. 1D) to the generation of this activity. We also noted that numerous features of propagation (i.e., amplitude, rate and direction) varied within the same muscle sample, even between two successive cycles of activity (Supplemental Video SV6).

Careful comparisons of potential images and field potential traces in 'expanding' activity indicated that the pacemaker potential lacked a preceding transient positive potential in the initiating micro-region (blue arrow in Fig. 2A). On the other hand, in 'migrating' activity, a dark-colored area preceded a light-colored area when activity propagated from the adjacent area of the MEA sensing area (Fig. 2B; Supplemental Fig. 1G, H). This corresponds to a transient positive potential (red arrowhead) preceding the fast down-stroke of the pacemaker potential.

The processes of the propagating activity in the MEA sensing area were tentatively modeled as follows. In 'migrating' activity, the light- and dark-colored areas are considered to be the 'source' and 'sink' of a local circuit current, respectively (see

schematic illustrations in Fig. 2C and Supplemental Fig. S3). The intercellular pacemaker current progressively charges the membrane in the sink region until the transmembrane pacemaker current is activated, employing a voltage-gated mechanism (Goto et al. 2004). Thereby, propagation of pacemaker activity sequentially along the muscle sheet acts as a 'volume conductor' (Supplemental Fig. S5). On the other hand, a cellular model of 'expanding' activity was constructed by adding an initiating region which likely employs an internal clock mechanism such as the activation of Ca^{2+}-activated transmembrane pacemaker current in response to intracellular Ca^{2+} oscillations owing to Ca^{2+} release channels in the endoplasmic reticulum (ER) (Aoyama et al. 2004; Saeki et al. 2019). After 'expanding' activity has been initiated, it can subsequently propagate away from the initiating region by the same mechanism as 'migrating' activity. However, some of the pacemaker potentials near the initiating site did not have preceding positive potentials (Fig. 2A), implying that propagation in the immediate vicinity of the initiating region may contain pacemaker cells coincidently activated by an internal Ca^{2+} clock, but with some delay. An alternative possibility is that a small group of cells neighboring the initiating region were activated by the intercellular propagation of Ca^{2+} or intracellular messengers (e.g., IP_3) via gap junctions, which has been reported for other cell types (Halidi et al. 2011).

Interconversion of micro-coordination patterns
When pacemaker events were classified into four patterns by incorporating the features in a propagation period, in addition to the large variations within the same category of event, we also frequently observed interconversion between the patterns (eight out of ten samples during the continuous MEA measurement). The frequency of interconversion differed substantially between samples. Continuous MEA recordings from a muscle sheet with frequent interconversions exhibited eight consecutive pacemaker events: one 'migrating', four 'expanding' (with different initiating regions), two 'colliding', and one 'bumpy/aberrant' (Fig. 3A). In the 'migrating' event, activity crossed the MEA area in the diagonal direction (Fig. 3B), and in the 'expanding' event, activity was initiated in the upper-left region (Fig. 3C; the last 'expanding' event in Fig. 3A). In the 'bumpy/aberrant' event, activity initially propagated from the lower region of the image in the aboral direction, but the activity only reached the upper-left region and failed to propagate to the upper-right region (Fig. 3D). In the first of the two 'colliding' events (Colliding-1: Fig. 3E), two pacemaker activities expanded from the upper-left and lower-middle regions to collide near the center, following which the merged activity propagated toward the upper-right region. In the second 'colliding' event (Colliding-2: Fig. 3F), two pacemaker activities initially propagated similar to the first 'colliding' event, but the merged activity did not propagate toward the middle-right region showing a slowly oscillating potential (Fig. 3F). It is likely that the aberrant slow potential substantially prolonged the refractory period of pacemaker cells in this region, resulting in a 'bumpy/aberrant' event. Indeed, various types of regional aberrant electric activity occurred, which may account for 'bumpy/aberrant' events. For example, in another sample showing an interconversion of micro-coordination patterns, multiple pacemaker activities occurred independently in the MEA sensing area and were separated by a 'non-conducting' region where rapid potentials run frequently (Supplemental Fig. S6).

Modulation of micro-coordination patterns

The micro-coordination pattern of pacemaker activity was changeable (Fig. 3), but the occurrence frequency of each pattern was relatively stable in the control condition (Fig. 1). We thus attempted to assess if the frequency of the spatio-temporal coordination patterns of pacemaker activity is employed for functional assessment in a digital manner, like the 'open' and 'closed' states of ionic channels in patch clamp experiments. Since 5-HT is one of the most important signaling molecules in functional modulation of GI motility (Spiller, 2011; Mawe and Hoffman, 2013), we examined the effects of 5-HT after each control MEA measurement (Fig. 4). An example of a series of MEA measurements from a muscle sheet demonstrated frequent activities of the 'expanding' pattern: a typical cycle involved the initiation of activity in the upper-middle region in control conditions (Fig. 4A,B; Supplemental Video SV7), and those converted into a 'migrating' pattern in the presence of 5-HT (100 μM), accompanied by an increase in the frequency of pacemaker events (Fig. 4C,D; Supplemental Video SV8). Similar changes were observed in nine muscle sheets (see also Supplemental Fig. S8: another example of the consistent effects of 5-HT examined in a muscle sheet showing frequent 'bumpy/aberrant' events in the control).

Linear spectrum (LS) analysis revealed that the magnitude of the pacemaker potential imaging frequency range ($LS_{0.25–10.5Hz}$) normalized whereas that of the sub-basal frequency range ($LS_{0.05-0.5Hz}$) increased in the presence of 5-HT ($LS_{0.25-10.5}/LS_{0.05-0.5}$: 8.3 ± 4.7 in the control vs. 13.7 ± 7.5 in 5-HT, $P = 0.01$, $N = 9$; Fig. 4E). On the other hand, the frequency of pacemaker events increased significantly in the presence of 5-HT (27.3 ± 7.2 min^{-1} in 5-HT vs. 21.3 ± 8.4 min^{-1} in the control, $P = 0.0038$, $N = 9$; Fig. 4F). In addition, 5-HT increased the occurrence of 'migrating' activity from 10.4 ± 9.5 to 29.3 ± 16.5 min^{-1} ($P = 0.017$, $N = 9$; Fig. 4G), and decreased that of 'expanding' activity from 11.4 ± 8.8 to 5.4 ± 8.5 min^{-1} ($P = 0.095$, $N = 9$; Fig. 4G). In the sum of nine experiments (total $N = 9$, 165 events in the control vs 216 events in the 5-HT treatment for 54 s), the number of the four pacemaker events changed accordingly (Fig. 4H). 'Expanding' and 'bumpy/aberrant' events decreased (from 49 to 8 events and from 35 to 20 events, respectively) while 'migrating' and 'colliding/converging' events increased (from 69 to 134 events and from 12 to 54 events, respectively).

In exploratory research for relevant receptor types, the additional application of ondansetron (500 μM), a 5-HT$_3$ receptor antagonist, in the presence of 5-HT decreased both the frequency (33.5 ± 5.7 min^{-1} vs 24.1 ± 4.2 min^{-1}, $P = 0.041$, $N = 4$) and magnitude ($LS_{0.25-10.5Hz}$: 45.0 ± 29.1% of the value in 5-HT alone, $P = 0.047$, $N = 4$; Supplemental Fig. S8) of pacemaker activity. These results agreed well with the discernment that 5-HT$_3$ receptors exert an excitatory influence on a wide range of GI functions (Talley, 1992; Spiller, 2011).

Discussion

Biological significance of the two major patterns

As the width of sensing electrode becomes small, the transmission efficiency of low frequency electric signals decreases due to an increase in impedance (Taniguchi et al. 2013; Iwata et al. 2017). Pt black nanoparticles have resolved electrical contradictions by increasing the surface area of sensing microelectrodes. Thereby, the dialysis membrane-reinforced low-impedance MEA technique enabled the stable measurement of field potentials in a wide frequency range, and classification of spatio-temporal

coordination of pacemaker potentials in a sensing microregion, despite surprisingly large variations. We note that the electrical matching of the sensing microelectrode and detecting amplifier system is important to appropriately measure low-frequency field potentials, a characteristic feature of the GI tract.

To date, evidence has accumulated for the synchronization of electric potentials and intracellular Ca^{2+} oscillations in gut pacemaker activity (Torihashi et al. 2002; van Helden and Imtiaz, 2003; Park et al. 2006). Therefore, the initiation of a pacemaker potential, such as 'expanding' activity in the MEA sensing microregion, is considered to reflect the Ca^{2+} clock mechanism operated by the co-existence of ryanodine receptors and inositol 1,4,5-trisphosphate receptors (IP_3Rs) in the ER of ICCs (Aoyama et al. 2004; Saeki et al. 2019) that consequently activate transmembrane Ca^{2+}-activated ionic conductance (i.e. anoctamine-1 Cl^- channels: Dickens et al. 1999; Hwang et al. 2009). On the other hand, 'migrating' activity, defined by activity that propagates to the MEA sensing area, represents pacemaker currents employing voltage-gated mechanisms.

Typical 'migrating' activity was characterized by a dark-colored (positive potential) area at the front of the propagating waveform that was followed by a light-colored (negative potential) area. The spatiotemporal relationship between the light- and dark-colored areas implies that intercellular coupling causes the ICCs in the dark-colored region to become electrically charged by the intercellular pacemaker current that has been activated in the ICCs in the light-colored region (sink and source regions, respectively, in Fig. 2C). In other words, pacemaker activity propagates through a volume conductor due to the formation of local circuits. This feature explains the existence of voltage-gated inward current attributed to atypical non-selective cation channels (Goto et al. 2004) and TTX-resistant Na^+ channels ($Na_v1.5$) (Strege et al. 2005) in pacemaker cells of rodents and humans, respectively. Taken together, the image analysis in the present study thus validates the biological significance of duplicating pacemaker mechanisms in the small intestine, by implying the functional state of the sensing microregion (i.e. spontaneity and synchrony/continuity for the 'expanding' and 'migrating' patterns, respectively).

Complexity of the waveform in extracellular recording
The potential images explicitly demonstrate the two sources of pacemaker current via transmembrane and intercellular pathways that can account for the complex waveform of electric activity in extracellular recordings (Sanders et al. 2016). Namely, during the depolarization period, the directions of transmembrane and intercellular currents were opposite: Transmembrane current flows inwardly while intercellular current flows outwardly to charge the membrane as a volume conductor propagates [e.g. microregions near ME(3) and ME(2), respectively, in Fig. 2C]. On the other hand, intracellular electrodes only exhibit depolarization when detecting either transmembrane or intercellular currents, so the first-derivative of the (trans)membrane potential waveform differs from the extracellular field potential waveform in the electric syncytium (when extracellular field potential is measured in the microregion with a volume conductor, it is sufficiently resolved). The waveform is further complicated in microregions where electric potentials from multiple sources interact (e.g., Fig. 3F, Supplemental Fig. S6). In 'migrating' pacemaker activities in the present study, therefore, preceding positive potentials (outward current) of sizeable amplitude propagate in the MEA sensing area, verifying the electric syncytium nature of the GI tract (Tomita, 1970; Manchanda et al.

2019; Sanders, 2019), where smooth muscle cells and network-forming pacemaker cells are electrically connected (Dickens et al. 1999; Sperelakis and Daniel, 2004).

The present potential imaging revealed that the direction of 'migrating' activity varied considerably. Actually, volume conductors flexibly migrate, irrespective of the running circular and longitudinal muscle bundles (e.g. Fig. 2B; Supplemental Fig. S4), indicating that pacemaker currents conduct mainly via a network of ICCs (i.e. ICCs in the myenteric plexus region). The lack of pacemaker potentials in the ileum of W/W^v mice reinforces this hypothesis. Use of an extracellular electrode array with a large polar distance (1 mm) and recording area (10×10 mm^2) was previously reported for the existence of reentrant electric waves in the small intestine of healthy rats, but they were less frequent than in diabetic rats (Lammers et al. 2011, 2012). The circuit that leads reentrant electric waves may correspond to large variations of 'migrating' activity in the small MEA area. However, as previous measurements were performed in different conditions of extracellular medium and preparation (without using Ca^{2+} channel antagonists, and with the mucous intact), careful assessment is needed in a future study. In addition, the previous studies did not distinguish whether activity 'migrates' to or 'expands' in the recording region, because the size of the volume conductor is comparable to or smaller than the area covered by each sensing electrode (1×1 mm^2). In other words, the coupling mechanism between electric waves is unclear due to either electric conduction or an internal clock of the relaxation oscillator. The reentrant circuit, if there is any in the small intestine of healthy animals, likely consists of a mixture (i.e. patchwork) of micro-regions with 'migrating' (electrically conducting) and 'expanding' (initiating) activities.

Alteration of micro-coordination patterns related to motor function
5-HT signaling is involved in promoting the propulsion of foodstuff through the intestine (Bülbring and Lin, 1958; Mawe and Hoffman, 2013). Therefore, during 5-HT application, an increase in the ratio of 'migrating' to 'expanding' activity (Fig. 4) likely reflects a change in the microregion from a segmentation-preferable state to propulsion-preferable state. In addition, it is considered that the decrease in 'bumpy/aberrant' events and increase in 'colliding/converging' events indicate improved electric connectivity between network-forming myenteric ICCs. In turn, this allows pacemaker activity originating in areas adjacent to the MEA to reach the MEA-sensing area more frequently. Consistently, the ratio of 'migrating' to 'expanding' activity (R$_{M/E}$) implies a three-fold enlargement of the area covered by each pacemaker activity initiating site (i.e. R$_{M/E}$ + 1: ~1.9 in control vs. ~6.4 in 5-HT).

In the small intestine, aberrant slow electric oscillations generated in ICCs-DMP yield segmentation of the tract by waxing and waning the basal pacemaker activity in network-forming myenteric ICCs (Huizinga et al. 2014). It is speculated that 5-HT might act to amplify pacemaker activity by suppressing aberrant spontaneous rhythmicity of low frequency originating from other cell members (e.g., slow potentials as shown in Fig. 3F) thereby improving local electric conduction in microregions. This hypothesis agrees well with the fact that the magnitude of pacemaker waves relative to that of sub-basal waves increased in the presence of 5-HT, mainly reflecting the latter component of field potential waves (Supplemental Fig. S9). In human genetics, mutations of 5-HT-related genes are involved in patients with GI dysmotility, such as the LL variant of the serotonin-transporter-gene-linked polymorphic region (5-

HTTLPR) for constipation-predominant IBS (Zhang et al. 2014), or the HTR3E-untranslated region variant for female diarrhea-predominant IBS (Kapeller et al. 2008). We anticipate that the MEA method communicated in this study will be useful for investigating spatio-temporal properties of pacemaker potentials in small model animals with disorders related to 5-HT signaling.

Conclusion

The dialysis membrane-reinforced low-impedance MEA technique enabled the stable measurement of field potentials in a wide frequency range in the microregion. Thus, visualization of spatio-temporal coordination of pacemaker potentials easily distinguished between 'expanding' and 'migrating' activities attributable to the duplicating pacemaker current systems of Ca^{2+} oscillation-activated and voltage-gated channels, respectively. Namely, the image analysis of MEA data was advantageous to assess the spatio-temporal relationship between intercellular and transmembrane currents. Potential images nicely exhibited volume conductors as coupled dark and light regions in which positive intercellular and negative transmembrane currents both charged the plasma membrane positively. The opposite directions of the intercellular and transmembrane currents accounted for the complex waveforms of extracellular recording. In both classifications of developmental processes and total features of propagation during each period of pacemaker event, the micro-coordination pattern was changeable, but the occurrence of each pattern was relatively stable during prolonged measurements. In addition to accelerating the occurrence of pacemaker events, 5-HT modulated the occurrence of 'expanding' and 'migrating' activities, agreeing well with its action, which causes a shift in the contractile function of the GI tract from segmentation to propulsive motions. These results therefore provide evidence that our MEA method is useful for quantitative assessment of the functional state of GI microregions. Further refinement of the methodology and criteria for classifying the pattern of pacemaker events are required. Despite this, future studies are anticipated in small model animals with GI motility diseases.

Acknowledgments

We are grateful to Tatsuhiro Noda, Hirotaka Morishita and Tomoka Nomura (Department of Cell Physiology, Nagoya University Graduate School of Medicine) for their assistance with the MEA experiments.

Funding

This work was partly supported by a grant-in-aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS No. 19H03558) and a research grant from the Japan Agency for Medical Research and Development (AMED No. 16ek0210037h0003).

References

Akbarali HI, Hawkins EG, Ross GR, Kang M. 2010. *Neurogastroenterol. Motil.* **22**, 1045–1055.

Aoyama M, Yamada A, Wang J, Ohya S, Furuzono S, Goto T, Hotta S, Ito Y, Matsubara T, Shimokata K, Chen SR, Imaizumi Y, Nakayama S. 2004. Requirement of ryanodine receptors for pacemaker Ca^{2+} activity in ICC and

HEK293 cells. *J. Cell Sci.* **117**, 2813–2825.

Astrand P, Brock JA, Cunnane TC. 1988. Time course of transmitter action at the sympathetic neuroeffector junction in rodent vascular and non-vascular smooth muscle. *J. Physiol.* **401**, 657–670.

Bayliss WM, Starling EH. 1899. The movements and innervation of the small intestine. *J. Physiol.* **24**, 99–143.

Bülbring E, Lin RC. 1958. The effect of intraluminal application of 5-hydroxytryptamine and 5-hydroxytryptophan on peristalsis; the local production of 5-HT and its release in relation to intraluminal pressure and propulsive activity. *J. Physiol.* **140**, 381–407.

Dickens EJ, Hirst GD, Tomita T. 1999. Identification of rhythmically active cells in guinea-pig stomach. *J Physiol.* **514**, 515-531.

Faussone-Pellegrini MS. 2005. Interstitial cells of Cajal: once negligible players, now blazing protagonists. *Ital. J. Anat. Embryol.* **110**, 11–31.

Goto K, Matsuoka S, Noma A. 2004. Role of corticotropin-releasing hormone in irritable bowel syndrome and intestinal inflammation. *J Physiol.* **559**, 411–422.

Halidi N, Boittin FX, Bény JL. 2011. Propagation of fast and slow intercellular Ca^{2+} waves in primary cultured arterial smooth muscle cells. *Cell Calcium* **50**, 459–467.

Huizinga JD, Lammers WJ. 2009. Gut peristalsis is governed by a multitude of cooperating mechanisms. *Am J Physiol Gastrointest Liver Physiol.* **296**, G1–G8.

Huizinga JD, Chen JH, Zhu YF, Pawelka A, McGinn RJ, Bardakjian BL, Parsons SP, Kunze WA, Wu RY, Bercik P, Khoshdel A, Chen S, Yin S, Zhang Q, Yu Y, Gao Q, Li K, Hu X, Zarate N, Collins P, Pistilli M, Ma J, Zhang R, Chen D. 2014. The origin of segmentation motor activity in the intestine. *Nat Commun.* **5**, 3326.

Huizinga JD. 2017. The powerful advantages of extracellular electrical recording. *Nat Rev Gastroenterol Hepatol.* **14**, 372.

Hwang SJ, Blair PJ, Britton FC, O'Driscoll KE, Hennig G, Bayguinov YR, Rock JR, Harfe BD, Sanders KM, Ward SM. 2009. Expression of anoctamin 1/TMEM16A by interstitial cells of Cajal is fundamental for slow wave activity in gastrointestinal muscles. *J Physiol.* **587**, 4887-4904.

Iwata N, Fujimura T, Takai C, Odani K, Kawano S, Nakayama S. 2017. Dialysis membrane-enforced microelectrode array measurement of diverse gut electrical activity. *Biosens. Bioelectron*. **94**, 312–320.

Kapeller J, Houghton LA, Mönnikes H, Walstab J, Möller D, Bönisch H, Burwinkel B, Autschbach F, Funke B, Lasitschka F, Gassler N, Fischer C, Whorwell PJ, Atkinson W, Fell C, Büchner KJ, Schmidtmann M, van der Voort I, Wisser AS, Berg T, Rappold G, Niesler B. 2008. First evidence for an association of a functional variant in the microRNA-510 target site of the serotonin receptor-type 3E gene with diarrhea predominant irritable bowel syndrome. *Hum Mol Genet.* **17**, 2967–2977.

Kito Y, Suzuki H. 2003. Properties of pacemaker potentials recorded from myenteric interstitial cells of Cajal distributed in the mouse small intestine. *J. Physiol.* **553**, 803–818.

Lammers WJ, Ver Donck L, Stephen B, Smets D, Schuurkes JA. 2008. Focal activities and re-entrant propagations as mechanisms of gastric tachyarrhythmias. *Gastroenterology* **135**, 1601–1611.

Lammers WJEP, Al-Bloushi HM, Al-Eisaei SA, Al-Dhaheri FA, Stephen B, John R, Dhanasekaran S, Karam SM. 2011. Slow wave propagation and plasticity of interstitial cells of Cajal in the small intestine of diabetic rats. *Exp Physiol.* **96**, 1039–1048.

Lammers WJ, Stephen B, Karam SM. 2012. Functional reentry and circus movement arrhythmias in the small intestine of normal and diabetic rats. *Am J Physiol Gastrointest Liver Physiol.* **302**, G684–689.

Manchanda R, Appukuttan S, Padmakumar M. 2019. Electrophysiology of syncytial smooth muscle. *J Exp Neurosci.* **13**, 1179069518821917.

Mawe GW, Hoffman JM. 2013. Serotonin signalling in the gut – functions, dysfunctions and therapeutic targets. *Nat. Rev. Gastroenterol. Hepatol.* **10**, 473–486.

Morishita H, Iwata N, Takai C, Mochizuki N, Kaji N, Hori M, Kajioka S, Nakayama S. 2017. Micro-coordination of pacemaker potentials in the intestine of the mouse. *Gastroenterology* **152**, 1831–1833.e4.

Nakayama S, Shimono K, Liu HN, Jiko H, Katayama N, Tomita T, Goto K. 2006. Pacemaker phase shift in the absence of neural activity in guinea-pig stomach: a microelectrode array study. *J. Physiol.* **576**, 727–738.

O'Grady G. 2012. Gastrointestinal extracellular electrical recordings: fact or artifact? *Neurogastroenterol Motil.* **24**, 1-6.

O'Grady G, Paskaranandavadivel N, Du P, Angeli T, Erickson JC, Cheng LK. 2017. Correct techniques for extracellular recordings of electrical activity in gastrointestinal muscle. *Nat Rev Gastroenterol Hepatol.* **14**(6), 372.

Park KJ, Hennig GW, Lee HT, Spencer NJ, Ward SM, Smith TK, Sanders KM. 2006. Spatial and temporal mapping of pacemaker activity in interstitial cells of Cajal in mouse ileum in situ. *Am J Physiol Cell Physiol.* **290**, C1411-27.

Saeki T, Kimura T, Hashidume K, Murayama T, Yamamura H, Ohya S, Suzuki Y, Nakayama S, Imaizumi Y. 2019. Conversion of Ca^{2+} oscillation into propagative electrical signals by Ca^{2+}-activated ion channels and connexin as a reconstituted Ca^{2+} clock model for the pacemaker activity. *Biochem Biophys Res Commun.* **510**, 242–247.

Sanders KM, Ward SM, Hennig GW. 2016. Problems with extracellular recording of electrical activity in gastrointestinal muscle. *Nat Rev Gastroenterol Hepatol.* **13**, 731-741.

Sanders KM. 2019. Spontaneous electrical activity and rhythmicity in gastrointestinal smooth muscles. *Adv Exp Med Biol.* **1124**, 3–46.

Spencer N, Walsh M, Smith TK, 1999. Does the guinea-pig ileum obey the 'law of the intestine'? *J Physiol.* **517**, 889–898.

Sperelakis N, Daniel EE. 2004. Activation of intestinal smooth muscle cells by interstitial cells of Cajal in simulation studies. *Am J Physiol Gastrointest Liver Physiol.* **286**, G234–243.

Spiller RC. 2011. Targeting the 5-HT$_3$ receptor in the treatment of irritable bowel syndrome. *Curr Opin Pharmacol.* **11**, 68–74.

Strege PR, Bernard CE, Ou Y, Gibbons SJ, Farrugia G. 2005. Effect of mibefradil on sodium and calcium currents. *Am J Physiol Gastrointest Liver Physiol.* **289**, G249–G253.

Szurszewski JH. 1987. Electrical basis for gastrointestinal motility. In Physiology of Gastrointestinal Tract, 2nd edition (ed. Johnson, L.R.), pp. 383–422 (Raven Press, New York).

Talley NJ. 1992. Review article: 5-hydroxytryptamine agonists and antagonists in the modulation of gastrointestinal motility and sensation: clinical implications. *Aliment Pharmacol Ther.* **6**, 273–289.

Taniguchi M, Kajioka S, Shozib HB, Sawamura K, Nakayama S. 2013. Spatial analysis of slowly oscillating electric activity in the gut of mice using low impedance arrayed microelectrodes. *PLoS One* **8**, e75235.

Teramoto H, Shimizu T, Yogo H, Nishimiya Y, Hori S, Kosugi T, Nakayama S. 2012. Assessment of gastric emptying and duodenal motility upon ingestion of liquid meal using rapid magnetic resonance imaging. *Exp Physiol.* **97**, 516–524.

Tomita T. 1970. Electrical properties of mammalian smooth muscle. In: Bülbring E, Brading A, Jones A, Tomita T, eds. Smooth Muscle. London, England: Edward Arnold, pp. 197–243.

Torihashi S, Fujimoto T, Trost C, Nakayama S. 2002. Calcium oscillation linked to pacemaking of interstitial cells of Cajal: requirement of calcium influx and localization of TRP4 in caveolae. *J Biol Chem.* **277**, 19191–19197.

van Helden DF, Imtiaz MS. 2003. Ca^{2+} phase waves: a basis for cellular pacemaking and long-range synchronicity in the guinea-pig gastric pylorus. *J Physiol.* **548**, 271–296.

Zhang ZF, Duan ZJ, Wang LX, Yang D, Zhao G, Zhang L. 2014. The serotonin transporter gene polymorphism (5-HTTLPR) and irritable bowel syndrome: a meta-analysis of 25 studies. *BMC Gastroenterol.* **14**, 23.

Fig. 1. Dialysis membrane-reinforced microelectrode array (MEA) recordings enabled classification of pacemaker micro-coordination. (**A**) Photographs showing a muscle sample isolated from the mouse ileum and mounted on an 8 × 8 MEA using a piece of dialysis membrane and an anchor. (**B**) Illustration showing the arrangement of the muscle sample and dialysis membrane on the MEA plate. The muscle sheet was positioned with the longitudinal muscle layer facing downward and with the oral and anal ends located toward the upper and lower sensing electrodes, respectively. Each microelectrode (ME) sensed changes in field potential reflecting the flow of local membrane current (I_M) through the access resistance (R_a) between the ME and a reference electrode. CM: circular muscle; ICC: interstitial cells of Cajal; LM: longitudinal muscle. (**C**) An 8 × 8 plot showing the spontaneous electric activities recorded from a muscle sample in the presence of nifedipine. (**D**) Confocal images of c-kit immunofluorescence showing ICCs in the myenteric plexus region (MP: upper) and the deep muscular plexus region (DMP: lower) (representative of ≥3 experiments). (**E**) Procedures used for potential mapping. HPF: high-pass filtering; LPF: low-pass filtering. (**F**) Patterns of pacemaker potential micro-coordination: 'expanding', 'migrating', 'bumpy/aberrant' and 'colliding/converging' (see text and Supplemental Fig. S1 for definitions). (**G, H**) Graphs showing the frequency of pacemaker events (min^{-1}) and frequency of 'migrating' or 'expanding' activity (min^{-1}) during the first and second halves of the MEA measurements (54–90 s) in 10 samples.

Fig. 2. Variations of 'expanding' and 'migrating' activities (A,B), and schema showing cellular processes (C,D). (**A**) Field potentials recorded continuously by three microelectrodes [ME(25), ME(28) and ME(31)] arranged transversely on the same muscle preparation. The field potential images show three consecutive 'expanding' activities that were initiated in different regions: upper-left (Expanding-1), lower-right (Expanding-2) and near the center (Expanding-3), corresponding to E-1*, E-2* and E-3* in continuous field potential recording, respectively. M_c: maximum potential used for color assignment. Traces of field potential waves recorded by three individual MEs are shown below each series of field potential images. Recorded MEs are indicated by the same color on the MEA. The electric potential wave near the initiating region (blue arrow) lacked a transient positive potential (red arrowhead). Representatives of 'expanding' activities observed in 10 samples. (**B**) Field potential images showing three examples of 'migrating' activities propagating in different directions: longitudinally in the oro-anal direction (downward; Migrating-L1), longitudinally in the aboral direction (upward; Migrating-L2) and diagonally in the oro-anal direction (circumferentially leftward; Migrating-D1). Traces of representative field potential waves are shown along with each series of field potential images. Transient positive potentials (red arrowheads). Representatives of 'migrating' activities in 10 samples. (**C,D**) Illustrations showing the cellular processes proposed to underlie 'migrating' activity and 'expanding' activity, respectively (see text for an explanation). The square cells represent pacemaker interstitial cells along with adjacent smooth muscle cells electrically connected, thereby increasing electric capacitance.

Fig. 3. Interconversion of micro-coordination patterns. (**A**) Field potentials continuously recorded from a muscle sample by three microelectrodes [ME(10), ME(28) and ME(51)], showing interconversion between the four micro-coordination

patterns of pacemaker events. This sample is a representative of the 10 samples used in Fig. 1 (G, H). Asterisks indicate the pacemaker events shown as potential maps in **B–F**. (**B**) Potential images of the 'migrating' event (activity) shown in **A** (M*). (**C**) Potential images of the 'expanding' event (activity) shown in **A** (E*). (**D**) Potential images of the 'bumpy/aberrant' event shown in **A** (B*) together with superimposed field potential traces recorded by different microelectrodes. (**E**) Potential images of the first 'colliding' event shown in **A** (C-1*) together with superimposed field potential traces recorded by different microelectrodes. (**F**) Potential images of the second 'colliding' activity shown in **A** (C-2*) and the phase that followed. The superimposed field potentials reveal slow potentials in the middle-right region ('bumpy' spot), suggesting that this may have caused a 'bumpy/aberrant' event in the next cycle of pacemaker activity.

Fig. 4 Modulation of micro-coordination patterns by 5-HT. (**A**) Field potentials continuously recorded in a sample of ileal musculature in control medium. All activities are of the 'expanding' type. (**B**) Series of field potential images and superimposed field potential traces for the period marked with a bar in A. (**C, D**) Effects of 5-HT (100 μM) on pacemaker activity in the same sample. All activities are of the 'migrating' type. The size of the potential image is 1050×1050 μm^2. (**E, F**) Graphs showing the ratio of the normalized magnitude of pacemaker waves (E: $LS_{0.25-10.5Hz}/LS_{0.05-0.5Hz}$) and the frequency (F: min^{-1}) of pacemaker events under control conditions and in the presence of 5-HT. (**G**) Graphs showing the frequency of 'migrating' or 'expanding' activity (min^{-1}). (**H**) Graphs showing the relative contributions of 'migrating', 'expanding', 'bumpy/aberrant' and 'colliding/converging' patterns of pacemaker events. (**E-H**, $N = 9$ in nine samples).

Legends for Supplemental Figures

Fig. S1. Illustration showing the procedures of classification in micro-coordination of pacemaker potentials. Micro-coordination of potential images is classified into two levels: event and activity levels. In each pacemaker event, pacemaker activity is detected in the MEA sensing area. In the activity level classification, based on the developmental process of pacemaker activity which initiates in or migrates to the MEA sensing area, each activity is categorized into either a 'migrating' or an 'expanding' pattern, respectively. In the event level classification, when pacemaker activity propagates coherently, it is termed a 'migrating' or 'expanding' event. Additionally, a subclass of micro-coordination is defined as 'bumpy/aberrant' by the propagation feature lacking spatio-temporal regularity, whether activity develops as a 'migrating' or 'expanding' pattern. Likewise, another subclass of pacemaker event is termed a 'colliding/converging' event due to the existence of multiple pacemaker activities in the same pacemaker event, as they frequently collide and merge during the pacemaker event.

Fig. S2. Motion tracking of a muscle sample used in MEA measurement. (**A**) After the MEA measurement, video of transmission image acquired in a muscle sample mounted in a glass-bottomed chamber. The extracellular solution contained nifedipine. The chamber was warmed on a glass heater. (**B**) Plots of the X, Y-shift of the tracking square shown in **A**. (**C**) Plots of representative field potentials of the MEA. The measurement was performed before the video, verifying negligible movement.

Fig. S3. Microelectrode array (MEA) recordings of spontaneous electrical activity in a sample of ileal muscle from a _W/W^v_ mouse that lacks network-forming interstitial cells of Cajal (ICCs). After removal of the mucosa, the ileal muscle sample was mounted on an 8×8 MEA using a piece of dialysis membrane, as shown in Fig. 1A. (**A, B**) An 8×8 plot of field potentials recorded under control conditions (**A**) and after exposure to the dihydropyridine Ca^{2+} channel antagonist, nifedipine (**B**). (**C**) Confocal images of c-kit immunoreactivity in the regions of the myenteric plexus (upper) and the deep muscular plexus (lower). (**D**) Field potential images showing electrical activity in the presence of nifedipine, corresponding to the MEA measurement in (**B**). In contrast to the MEA recordings made in wild-type mice, there was negligible electric activity in the ileum of _W/W^v_ mice in the presence of nifedipine (**B**).

Fig. S4. 'Migrating' activities propagated in circular and diagonal directions. Field potential images and representative field potential waves showing three examples of 'migrating' activity, in addition to those in Fig. 2B. Migrating-C1 and -C2: Circumferential propagation (along the circular muscle) toward the right and left, respectively. Migrating-D2: Diagonal propagation in the oro-anal direction (circumferentially toward the left). M_c: maximum potential used for color assignment. Red arrowheads indicate transient positive potentials in the 'migrating' waves. The MEs for representative field potential waves are indicated by circles of the same color on the MEA.

Fig. S5. Schematic illustration showing the process underlying 'migrating' activity. (**A**) Intercellular pacemaker current propagates toward the cells near ME(1) and ME(2) and charges their plasma membranes. The cells near ME(1) and ME(2) are in a charging ('C') state, and those near the other MEs are in a resting ('R') state. (**B**) Next, voltage-gated inward current (a part of pacemaker current) is activated in the cell near ME(1) so that this cell enters an activated ('A') state. Meanwhile, the intercellular pacemaker current propagates toward the cells near ME(2) and ME(3). (**C**) The cell near ME(1) enters a slow oscillation ('S') state in which transmembrane outward K^+ current and Ca^{2+}-activated inward currents are continuously activated. Intracellular Ca^{2+} release processes likely contribute to the activation of Ca^{2+}-activated inward currents (e.g., anoctamin-1 Ca^{2+}-activated Cl^- channels: the equilibrium potential of pacemaker cells $\approx$ -20 mV, more positive than the resting membrane potential). Meanwhile, the cell near ME(2) enters the 'A' state, and the intercellular pacemaker current further propagates toward the cells near ME(3) and ME(3). Thus, the cells near ME(1) to ME(4) form a local circuit. The intercellular pacemaker current does not propagate toward cells on the left, because the membranes of these cells are already charged. Each cell in the scheme represents a group of electrically connected pacemaker cells, adjacent interstitial cells and smooth muscle cells, and this unit thereby possesses sizeable electric capacity.

Fig. S6. Pacemaker activities separated by a transient 'non-conducting (NC)' region, in a muscle sample showing an interconversion of pacemaker event patterns. (**A**) Field potentials recorded from different regions of the MEA. (**B**) Field potential images showing the two 'bumpy/aberrant' activities (B-1 and B-2) indicated in the field potential recorded in A. The B-1 and B-2 'bumpy/aberrant' events occurred with different cycles and propagated locally in different regions (upper and lower, respectively).

Fig. S7. Modulation of micro-coordination patterns by 5-HT in a sample showing frequent 'bumpy' activity. Field potential traces recorded from different microelectrodes and corresponding field potential images obtained under control conditions (**A, B**) and in the presence of 5-HT (**C, D**) (an example of muscle sample showing frequent 'bumpy/aberrant' events in the control out of $N = 9$ experiments in nine samples). The field potential images correspond to the B* ('bumpy/aberrant' in the control) and M* ('migrating' in 5-HT) events indicated in the field potential traces. During the first half of the control recording period, premature activity-like potentials (red arrowheads) occurred between the 'bumpy/aberrant' events. The size of the potential image is $1050 \times 1050 \ \mu m^2$.

Fig. S8. Effects of ondansetron on pacemaker activity. (**A, B**) Continuous measurement of MEA field potentials and potential images in the presence of 5-hydroxytryptamine, and (**C, D**) during additional application of ondansetron. The period of potential image processing is shown by the bars in field potentials (**A, C**). The size of the potential image is $1050 \times 1050 \ \mu m^2$. (**E, F**) Graphs showing changes in the magnitude (linear spectrum, 0.25–10.5 Hz) and frequency (min^{-1}) of pacemaker activity after the application of ondansetron (representative of $N = 4$ experiments in four samples).

Fig. S9. Effects of 5-HT on the magnitude of field potential waves. (**A**) Graphs showing the effect of 5-HT on the magnitude of the linear spectrum in the frequency range of pacemaker potential imaging (left: LS $_{0.25\text{-}10.5\ Hz}$, 1.08 ± 0.33 in the control, $P = 0.47$, $N = 9$) and in the frequency range of sub-basal waves (right: LS $_{0.05\text{-}0.5\ Hz}$, 0.73 ± 0.12 in the control, $P = 0.047$, $N = 9$).

Supplemental Videos:
The size of all videos is 1050×1050 μm^2. The speed of videos was slowed four times.
Supplemental Video SV1. Field potential video showing the 'expanding' pattern of a pacemaker event, corresponding to the data in Fig. 1F.
SV1: Movie S1.mp4

Supplemental Video SV2. Field potential video showing the 'migrating' pattern of a pacemaker event, corresponding to the data in Fig. 1F.
SV2: Movie S2.mp4

Supplemental Video SV3. Field potential video showing the 'bumpy/aberrant' pattern of a pacemaker event, corresponding to the data in Fig. 1F.
SV3: Movie S3.mp4

Supplemental Video SV4. Field potential video showing the 'colliding' pattern of a pacemaker event, corresponding to the data in Fig. 1F.
SV4: Movie S4.mp4

Supplemental Video SV5. Field potential video corresponding to Fig. 2A. Three 'expanding' activities were initiated from distinct regions.
SV5: Movie S5.mp4

Supplemental Video SV6. Field potential video showing two consecutive 'migrating' activities. The 'migrating' activities propagated diagonally toward the lower-right region and longitudinally toward the upper region.
SV6: Movie S6.mp4

Supplemental Video SV7. Field potential video showing consecutive pacemaker events in the control, corresponding to Fig. 4A. A variety of micro-coordination patterns, including those of the 'expanding' pattern, occurred in the control.
SV7: Movie S7.mp4

Supplemental Video SV8. Field potential video showing consecutive pacemaker events in the presence of 5-HT, corresponding to Fig. 4C. 'Migrating' activities alone propagated diagonally toward the lower-left region, accompanied by an increase in the frequency.
SV8: Movie S8.mp4

Field Potential Image Classification of Pacemaker Micro-coordination in the Intestine of the Mouse

Naoko Iwata[1], Chiho Takai[1], Naoto Mochizuki[1], Mariko Yamauchi[1],

Noriyuki Kaji[2]

Yoshiyuki Kasahara[3], Shinsuke Nakayama[1,*]

[1]Department of Cell Physiology, Nagoya University Graduate School of

Medicine,

Nagoya 466-8550, Japan

[2]Department of Veterinary Pharmacology, Azabu University School of

Veterinary Medicine, Sagamihara 252-5201, JAPAN

[3]Department of Fetal and Maternal Therapeutics, Tohoku University

Graduate School of Medicine, Sendai 980-8575, JAPAN

Figures:

Figure 1

Figure 2

Figure 3

Figure 4

Figure S1

Figure S2

Dialysis membrane-reinforced ileal muscle sheet in the presence of nifedipine

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Figure S9