Experimental demonstration of tethered gene drive systems for confined population modification or suppression

Matthew Metzloff[1,2], Emily Yang[1,2], Sumit Dhole[3], Andrew G. Clark[1,2], Philipp W. Messer[1], Jackson Champer[1,2,4*]

[1]Department of Computational Biology, Cornell University, Ithaca, NY 14853
[2]Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853
[3]Department of Entomology and Plant Pathology, North Carolina State University, Raleigh, North Carolina
[4]Current Address: Center for Bioinformatics, School of Life Sciences, Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, China 100871

*Corresponding author: JC (jc3248@cornell.edu)

ABSTRACT

Homing gene drives hold great promise for the genetic control of natural populations. However, current homing systems are capable of spreading uncontrollably between populations connected by even marginal levels of migration. This could represent a substantial sociopolitical barrier to the testing or deployment of such drives and may generally be undesirable when the objective is only local population control, such as suppression of an invasive species outside of its native range. Tethered drive systems, in which a locally confined gene drive provides the CRISPR nuclease needed for a homing drive, could provide a solution to this problem, offering the power of a homing drive and confinement of the supporting drive. Here, we demonstrate the engineering of a tethered drive system in *Drosophila*, using a TARE drive to support modification and suppression homing drives. Each drive was able to bias inheritance in its favor, and the TARE drive was shown to spread only when released above a threshold frequency in experimental cage populations. After the TARE drive had established in the population, it facilitated the spread of a subsequently released split homing modification drive (to all individuals in the cage) and of a homing suppression drive (to its equilibrium frequency). Our results show that the tethered drive strategy is a viable and easily engineered option for providing confinement of homing drives to target populations.

INTRODUCTION

Powerful homing gene drives can spread from low starting frequencies throughout an entire population[1–4]. However, this capability also renders such drives highly invasive since small levels of migration could facilitate their spread into any connected populations[5,6]. Though potentially desirable in some applications, this could also substantially increase the social and political difficulties associated with deploying a gene drive in the field.

Gene drive technology has improved remarkably over the past few years, and several different CRISPR-based gene drives have now been demonstrated in yeast[7–10], flies[11–25], mosquitoes[26–35], and mice[36]. Most of these are homing drives, which have successfully achieved the modification and suppression of laboratory populations[25,29,35]. Several promising applications have been proposed for this technology, such as the genetic modification of *Aedes* and *Anopheles* mosquito populations by introducing genes that could reduce transmission of malaria, dengue, and other diseases[1–4]. Furthermore, gene drives could be used to directly suppress populations of disease vectors or invasive species. Overcoming the challenge of confining a gene drive to a target population could represent an important step in bringing such approaches closer to deployment.

Several types of potentially confinable gene drives have been developed. One class of especially promising candidates are CRISPR toxin-antidote (TA) gene drives. These drives will only spread in the population when introduced above a threshold frequency, which is determined by the parameters of the drive. If migrants to connected populations cannot surpass this frequency, TA drives will not spread in these populations and can thereby remain confined[37,38]. Recent studies have already used TA drives to successfully modify cage populations[38,39]. Despite spreading more slowly than effective homing drives and also, by design, requiring larger releases sizes, such confined drives could ease concerns associated with less controllable homing drives.

The development of a confined suppression drive poses a greater challenge than confined modification drives due to the need for the drives to retain enough power to spread through the population and still provide the high genetic load required for effective population suppression. A spatially confined gene drive capable of suppression, or perhaps similarly spreading a cargo gene with a high fitness cost, would therefore be of considerable interest. Three novel systems have been suggested to this end: Toxin-Antidote Dominant Embryo (TADE) suppression drives[37,40], daisy homing drives[41], and tethered homing drive systems[42]. TADE suppression drives disrupt essential fertility genes and additionally target a haplolethal gene with Cas9, while the drive allele provides a rescue copy of the haplolethal gene[37,40]. These drives are capable of confined population suppression, but they require a haplolethal target gene, which can be difficult to work with. Daisy homing drives consist of a series of homing elements, with each element driving the next in the chain[41]. As the system spreads through a population, each element is lost in turn until the final element is no longer able to increase in frequency[41]. Daisy drives require multiple suitable target genes to avoid resistance, and because many driving

elements are required, construction is complicated[43]. In addition, with higher migration rates and lower cargo fitness costs, daisy drives could be more difficult to confine[43].

Tethered homing drive systems provide a potential alternative to solve some of these issues. A tethered homing drive system involves the modification of a population with a confined drive, along with the spread of a homing drive that is only capable of drive activity in the presence of the confined drive[42]. Thus, a tethered system potentially combines benefits of confinement with the power of a homing drives. Modeling suggests that such a system could be capable of confined suppression of a target population[42]. A tethered system for modification should also be able to spread cargo genes with much higher fitness costs than other confined drives.

Here, we show that a tethered gene drive system can modify cage populations of *Drosophila melanogaster*. We use a regionally confinable Toxin-Antidote Recessive Embryo (TARE) drive targeting an essential but haplosufficient gene. This drive already contains the necessary Cas9 for a tethered homing drive, making it particularly suitable for use in a tethered system. We then test two split homing drives in the TARE-modified populations: The first is a modification drive targeting a haplolethal gene with two gRNAs. The second is a suppression drive targeting a haplosufficient but essential female fertility gene with four gRNAs. The TARE drive spread successfully when released above its threshold frequency and then was able to provide a Cas9 source for the homing drives, which increased in frequency according to their expected behavior.

METHODS

Plasmid construction. The starting plasmid EGDh2 was constructed previously[38,39] (see Supplementary Information). Restriction enzymes for plasmid digestion, Q5 Hot Start DNA Polymerase for PCR, and Assembly Master Mix for Gibson assembly were acquired from New England Biolabs. Oligonucleotides were obtained from Integrated DNA Technologies. 5-α competent *Escherichia coli* from New England Biolabs and ZymoPure Midiprep kit from Zymo Research were used to transform and purify plasmids. See Supplementary Information for a list of DNA fragments, plasmids, primers, and restriction enzymes used for the cloning of each construct.

Generation of transgenic lines. Injections were conducted by Rainbow Transgenic Flies. The donor plasmid TAREhNU2G (507 ng/µL) was injected along with plasmid EGDhg2t (149 ng/µL), which provided gRNAs for transformation and was constructed previously[38], and pBS-Hsp70-Cas9 (442 ng/µL, from Melissa Harrison & Kate O'Connor-Giles & Jill Wildonger, Addgene plasmid #45945), which provided Cas9. A 10 mM Tris-HCl, 100 µM EDTA solution at pH 8.5 was used for the injection.

Genotypes and phenotypes. TARE drive carriers are indicated by expression of EGFP driven by the 3xP3 promoter, which is highly visible in the eyes of w^{1118} flies. Flies carrying either of

the homing drives are marked by DsRed driven by the 3xP3 promoter. For phenotyping, flies were frozen, and scored for red and green fluorescence in the eyes using the NIGHTSEA system.

Fly rearing. Flies were reared at 25 °C with a 14/10 h day/night cycle. Bloomington Standard medium was provided as food. The cage study used 30x30x30 cm (Bugdorm, BD43030D) enclosures. Flies homozygous for the TARE drive were allowed to lay eggs in food bottles for one day, and w^{1118} individuals were separately allowed to lay eggs in food bottles for one day. All adults were removed, and eight bottles were placed in each cage. Eleven days later, the bottles were replaced, and the adult flies were left in the cage and allowed to lay eggs for one day before being removed and frozen for phenotyping. This 12-day cycle was repeated until the TARE drive had nearly reached fixation. Flies heterozygous for each homing drive and homozygous for the TARE drive were separately allowed to lay eggs in food bottles for one day, and the adults were removed before the bottles were placed into the cages with TARE drive bottles. The same 12-day cycle was repeated, with all flies phenotyped.

To prevent accidental releases of gene drive flies, all live drive-carrying flies were quarantined at the Sarkaria Arthropod Research Laboratory at Cornell University under Arthropod Containment Level 2 protocols in accordance with USDA APHIS standards. In addition, the homing drives contained no Cas9 and were incapable of drive conversion in wild-type flies. All safety standards were approved by the Cornell University Institutional Biosafety Committee.

Phenotype data analysis. To calculate drive parameters, we pooled all offspring from the same type of cross together and used the combined counts to calculate rates. Because offspring had different parents and were raised in separate vials, this pooling approach could potentially distort rate and error estimates. To account for such batch effects, we performed an alternate analysis similar to the one used in previous studies[24,25,38]. This involved fitting a generalized linear mixed-effects model with a binomial distribution (fit by maximum likelihood, Adaptive Gauss-Hermite Quadrature, nAGQ = 25). In this model, offspring from a single vial were considered as a separate batch, even if they had the same parents as offspring from other vials. This approach allows for variance between batches and results in slight changes to rate and error estimates. The analysis was performed using the R statistical computing environment (3.6.1) with the packages lme4 (1.1-21, https://cran.r-project.org/web/packages/lme4/index.html) and emmeans (1.4.2, https://cran.r-project.org/web/packages/emmeans/index.html). The specific R script we used for this analysis is available on Github (https://github.com/MesserLab/Binomial-Analysis). The resulting rate estimates and errors were similar to the pooled analysis and are provided in Data Sets S1-S3.

Genotyping. To extract DNA, flies were frozen and ground in 30 µL of 10 mM Tris-HCl pH 8, 1mM EDTA, 25 mM NaCl, and 200 µg/mL recombinant proteinase K (Thermo Scientific). The mixture was incubated at 37˚C for 30 minutes, then at 95˚C for 5 minutes. The DNA was used as a template for PCR using Q5 Hot Start DNA Polymerase (New England Biolabs), following the manufacturer's protocol. The region of interest containing gRNA target sites was amplified using

DNA oligo primers hCut_S_F and hCut_S_R, and DNA fragments were isolated by gel electrophoresis. Sequences were obtained by Sanger sequencing and analyzed with ApE software (http://biologylabs.utah.edu/jorgensen/wayned/ape).

Estimation of drive fitness. We used a maximum likelihood approach to estimate the fitness parameters for the gene drive constructs. Details of this approach and its application to cage population data were previously described[44]. The fitness parameters for both homing constructs used here have been previously estimated[25,45]. For the TARE construct we estimated the viability, female fecundity, and male mating success parameters and the effective population size using a model that assumes a co-dominant, multiplicative fitness effect of the TARE construct (TARE-heterozygotes were assigned fitness equal to the square root of the TARE-homozygote fitness). The genomic site of TARE (the *h* locus) is haplosufficient, so heterozygotes that bear one wild-type and one disrupted *h* allele were assumed to have the same fitness as wildtype homozygotes. The germline and embryonic rates of cleavage of the wildtype allele by TARE were set at experimentally measured values. Parameter values were estimated by maximizing the likelihood across all generation transitions from the four cages combined (i.e, a single estimate per parameter is generated for all four cages). The effective population size parameter was estimated as a fraction of the census population size in each generation transition, using the average of both generations that were part of the transition. The code is available on GitHub (github.com/MesserLab/TetheredDrives).

RESULTS

Drive construct design. Our TARE drive is located at the *hairy* locus (*h*) on chromosome 3L (Figure 1) and is similar to the TARE drive used to modify a cage population in a previous study[38]. It includes a recoded *h* sequence, Cas9 expressed by the *nanos* promoter, EGFP as a marker expressed by a 3xP3 promoter and SV40 3' UTR, and two gRNAs expressed by the U6:3 promoter.

FIGURE 1. TARE Drive Construct. a) The TARE drive includes a recoded *h* sequence, Cas9, EGFP, and two gRNAs. **b)** The wild-type *h* allele, located on chromosome 3L, is cleaved in the third coding exon.

Our haplolethal homing drive was constructed previously[25]. In short, it is a population modification homing drive that targets the haplolethal gene *RpL35A* with two gRNAs and

provides a recoded *RpL35A* as rescue. Since the target gene is haplolethal, individuals with disrupted alleles will not be viable. Our homing suppression drive is described in detail in another study[45]. It targets a haplosufficient gene with four gRNAs. Both homing drives are split drives and do not contain Cas9. They are only capable of drive activity in TARE-modified individuals, which express Cas9 in the germline (Figure S1).

These homing drives have been engineered to reduce the incidence and impact of resistance alleles. Because the drives use multiplexed gRNAs, the formation of functional resistance alleles will be extremely rare[24]. Nonfunctional resistance alleles will lead to nonviable individuals in the haplolethal drive and will not outcompete drive alleles in the suppression drive, although the formation of large quantities of nonfunctional resistance alleles can slow the rate of spread of either drive.

TARE drive evaluation. We crossed flies heterozygous for the TARE drive to w^{1118} flies to determine drive inheritance. Observation of the EGFP phenotype in the eyes of offspring from this cross was used to identify the presence of drive alleles (Figure 2, Data Set S1). Progeny of drive females showed 72% inheritance, a rate significantly different than Mendelian inheritance ($p < 0.0001$, Fisher's exact test). Progeny of drive males showed 52% inheritance, which is not significantly different than Mendelian inheritance ($p = 0.1511$, Fisher's exact test). This was expected since nonviable individuals form in these crosses only as a result of embryo Cas9 activity, which only occurs in the progeny of drive-carrying females[37].

FIGURE 2. TARE Drive Inheritance. Individuals heterozygous for the TARE drive were crossed with w^{1118} individuals, and EGFP expression in progeny indicated the presence of the drive. The size of the dots is proportional to the number of adult progeny from a single drive individual. The rate estimates and standard error of the mean (SEM) are indicated.

To determine germline and embryo cleavage rates, we crossed male and female TARE drive heterozygotes (Data Set S1). Assuming that male and female germline cleavage rates are similar and that embryo cleavage only occurs in the offspring of females, the drive inheritance rate for this cross can be used with the rate for the cross between TARE females and w^{1118} males to determine the rates of embryo activity and germline activity. Using this approach, we calculated an embryo cleavage rate of 63.2% and a germline cleavage rate of 88.8% for the TARE drive. The latter is somewhat lower than other studies of gene drive in *D. melanogaster*[12,17,19,20,22–25,38,39] but is still sufficient to support effective population modification.

To further characterize and quantify the mechanism of our TARE drive, we crossed drive/wild-type heterozygotes with w^{1118} individuals. Flies were allowed to lay eggs for three days, and then transferred to new vials once per day for the next two days. In the second and third vials, eggs were counted, and pupae were counted in addition to phenotyping eclosed adults. The progeny of female drive heterozygotes had 80.7% egg-to-pupae survival, and the progeny of male drive heterozygotes had 88.2% egg-to-pupae survival (Figure 3, Data Set S1). The difference between these egg-to-pupae survival rates is significant ($p = 0.0013$, Fisher's exact test). Because the TARE drive relies on embryo Cas9 activity to create nonviable genotypes in the progeny of females, this difference indicates drive activity in females, as expected. Egg-to-pupae survival was also determined for crosses between TARE heterozygotes and haplolethal homing drive heterozygotes (Figure S2).

FIGURE 3. Egg-to-Pupae Viability. Individuals heterozygous for the TARE drive were crossed with w^{1118} individuals, and eggs and pupae were counted as well as eclosed adults. The size of the dots is proportional to the number of eggs from a single female. The rate estimates and standard error of the mean (SEM) are indicated.

Tethered homing drive evaluation. We crossed flies heterozygous for both the TARE drive and each homing drive with w^{1118} individuals to determine drive inheritance for the homing drives. Observation of the EGFP phenotype in the eyes of offspring identified the presence of the TARE drive, and the dsRed phenotype identified either the homing suppression drive (Figure 4, Data Set S2) or the haplolethal homing drive (Figure 5, Data Set S3). In both cases, inheritance rates were lower than the rates observed in the original studies using these drive constructs[25,38] This is likely because the *nanos*-Cas9 source (part of the TARE drive) led to lower levels of Cas9 expression than the previously tested source. In addition, while the TARE drive and haplolethal homing drive are located at two distant loci, they are still on the same chromosome, and *D. melanogaster* chromosomes do not undergo crossovers in males during meiosis. This could result in suppression of the TARE drive by nonfunctional resistance allele formation from the haplolethal homing drive.

FIGURE 4. Homing Suppression Drive Inheritance. Individuals heterozygous for both the TARE drive and the homing suppression drive were crossed with w^{1118} individuals. EGFP indicated the presence of the TARE drive, and dsRed indicated the presence of the homing drive. The size of the dots is proportional to the number of adult progeny from a single drive individual. Rate estimates and SEM are indicated.

FIGURE 5. Haplolethal Homing Drive Inheritance. Individuals heterozygous for both the TARE drive and the haplolethal homing drive were crossed with w^{1118} individuals. EGFP indicated the presence of the TARE drive, and dsRed indicated the presence of the homing drive. The size of the dots is proportional to the number of adult progeny from a single drive individual. Rate estimates and SEM are indicated.

Tethered system cage study. To assess the performance of the tethered homing drive system in large cage populations, we allowed individuals homozygous for the TARE drive to lay eggs in bottles for one day. Similarly, w^{1118} individuals were allowed to lay eggs in another set of bottles for one day. Flies were removed, and the bottles were placed in four population cages. Emerging adults were considered to be generation zero, and the TARE drive was allowed to spread in each population, with all adults phenotyped for EGFP (Figure 6, Data Set S4). In all cages, the population fluctuated, averaging 3,491 individuals (Figure S3).

In one cage with a lower initial release frequency, the TARE drive increased in frequency only slowly, and in another cage, the drive frequency remained constant. These two cages experienced fluctuations in frequency (possibly due to differences in the health of initially released TARE and w^{1118} individuals) before quickly stabilizing at drive carrier frequencies of 20% and 10%, suggesting that the introduction threshold of the TARE drive may be approximately 5-15%, a value that depends on the efficiency and fitness cost of the drive[37,38]. With the cleavage parameters calculated based on drive inheritance (Data Set S1), the fitness cost of the drive in homozygotes was perhaps 10-15% based on previous models[37]. A maximum-likelihood based analysis[44] indicates that drive homozygotes had a 12% fitness cost (fitness 0.879 compared to wild-type fitness of 1, with a 95% confidence interval of 0.788 - 0.977) that impacted female fecundity and male mating success, rather than offspring viability. This would result in a 10% introduction frequency threshold. The estimated effective population size was 2.0% of the census population size (with a 95% confidence interval of 1.3% - 2.9%). However, models with fitness

based on just mating success or fecundity fit the data similarly well (these would yield similar introduction thresholds).

In the two cages where the TARE drive reached 100% frequency, individuals homozygous for the TARE drive and heterozygous for one of the homing drives were released. The haplolethal homing drive was introduced at 6% frequency, and the homing suppression drive was introduced at 33% frequency (Figure 6, Data Set S4). The homing drives were allowed to spread for several generations, with all adults phenotyped for dsRed. The haplolethal homing drive eventually reached a frequency of 100%, while the homing suppression drive approached an apparent equilibrium frequency of somewhat over 50%. The rate of increase of these homing drives was lower than in other studies that used these lines[25,45], as expected from the somewhat lower drive conversion rate of these drives when combined with the TARE drive compared to the previously used *nanos*-Cas9 supporting element. The additional gRNAs in the TARE drive may also have somewhat reduced the cleavage activity at each site in the split homing drives due to saturation of Cas9, though this reduction was likely small[24].

FIGURE 6. Tethered Homing System Cage Study. Individuals homozygous for the TARE drive were introduced into a population that was wild type at the drive site. The cage populations were followed for several discrete generations, each lasting twelve days, including one day of egg-laying. After the TARE drive reached fixation in two cages, individuals that were homozygous for the TARE drive and heterozygous for one of the split homing drives were released. All individuals from each generation were phenotyped for DsRed and EGFP, with positive drive carriers having either one or two drive alleles.

DISCUSSION

In this study, we performed an experimental trial of the "tethered gene drive" system. This system was designed to allow a potentially confined drive system, such as a TARE drive or an underdominance system, to support a homing drive, thus allowing costly alleles to be spread with the powerful homing drive, while also preventing it from spreading to populations beyond the supporting confined drive[42]. Our demonstration with TARE followed by a tethered population modification drive[25] was successful, though the tethered homing suppression drive did not perform as well. However, this was most likely due to inadequate performance of the suppression drive itself[45], rather than a feature of the tethered system, which would only be expected to slightly reduce drive performance due to saturation of Cas9 with additional gRNAs in the combined system.

Since the TARE drive utilized the same rescue element, gRNA promoter, and target sites as a previous split TARE system[38], its reduced performance as indicated by reduced drive inheritance was almost certainly due to the Cas9 element. Though both of these TARE drives had identical *nanos*-Cas9 elements (with the same orientation with respect to the 3xP3 promoter of the fluorescent element), their genomic locations were substantially different. In this study, the Cas9 gene was located within the TARE drive on chromosome 3L, rather than a separate genomic site on chromosome 2R. This led to reduced cleavage rates, particularly in the early embryo, a critical parameter that can influence drive inheritance in individual crosses. In this case, such reduced performance could potentially be useful to make the drive more confined to a target region. However, for the homing suppression element, high performance would likely be needed to achieve high enough drive conversion efficiency and genetic load for population suppression[24], particularly in complex natural environments[46,47]. Higher expression rates of Cas9 within the TARE element could possibly be achieved by reorienting the Cas9 element within the drive allele. It would also be straightforward to choose a different site within the *h* gene, or more likely, a different target gene entirely. Essential but haplosufficient genes can likely be found in most genomes, and similar CRISPR toxin-antidote ClvR elements have already targeted other such genes with success[39,48].

In general, the tethered system provides a highly promising strategy for confining a homing drive to a target region. If an efficient homing drive can be made, then a suitable TARE system can likely be engineered as well. This is because a suitable homing system would have a high drive conversion rate, implying at least an equal rate of germline cleavage (which includes both the

drive conversion rate and germline resistance allele formation rate for homing drives). A TARE system would be expected to still have good efficiency without embryo cutting if the germline cut rate was high[37], allowing it to utilize the same promoter. While TARE would additionally need a rescue element, several studies have already shown that their engineering is feasible in flies[25,38,39,48] and moquitoes[35]. Furthermore, a TARE drive could be engineered to have an additional CRISPR nuclease with a different promoter and different gRNA specificity[49–51], allowing germline-restricted CRISPR cleavage for the homing element and both germline and embryo cleavage for the TARE element. For more stringent confinement, other CRISPR toxin-antidote systems with higher introduction thresholds could likely be engineered with similar ease, such as 2-locus TARE systems[40]. However, in situations where an efficient homing drive could not be engineered due to difficulty in achieving a high enough rate of drive conversion by homology-directed repair after germline cleavage, then TADE suppression systems[37,40] may present a useful alternative option for confined suppression if targeting of haplolethal genes could be successfully engineered.

This study experimentally demonstrated the feasibility of using tethered homing gene drives in cage populations based on CRISPR toxin-antidote drive systems. Such systems can allow for the power of homing drives to be limited to target populations. This could be useful and even necessary when dealing with invasive species outside their native range, when sociopolitical considerations demand a limited drive, or even during an initial testing phase of a homing gene drive system.

ACKNOWLEDGEMENTS

This study was supported by the National Institutes of Health awards R21AI130635 to JC, AGC, and PWM, award F32AI138476 to JC, and award R01GM127418 to PWM.

REFERENCES

1. Champer J, Buchman A, Akbari OS. Cheating evolution: engineering gene drives to manipulate the fate of wild populations. *Nat Rev Genet*, 17, 146–159, 2016.
2. Burt A. Heritable strategies for controlling insect vectors of disease. *Philos Trans R Soc L B Biol Sci*, 369, 20130432, 2014.
3. Alphey L. Genetic control of mosquitoes. *Annu Rev Entomol*, 59, 205–224, 2014.
4. Esvelt KM, Smidler AL, Catteruccia F, Church GM. Concerning RNA-guided gene drives for the alteration of wild populations. *Elife*, e03401, 2014.
5. Unckless RL, Messer PW, Connallon T, Clark AG. Modeling the manipulation of natural populations by the mutagenic chain reaction. *Genetics*, 201, 425–431, 2015.
6. Noble C, Adlam B, Church GM, Esvelt KM, Nowak MA. Current CRISPR gene drive systems are likely to be highly invasive in wild populations. *Elife*, 7, e33423, 2018.
7. DiCarlo JE, Chavez A, Dietz SL, Esvelt KM, Church GM. Safeguarding CRISPR-Cas9

gene drives in yeast. *Nat Biotechnol*, 33, 1250–1255, 2015.

8. Roggenkamp E, Giersch RM, Schrock MN, Turnquist E, Halloran M, Finnigan GC. Tuning CRISPR-Cas9 gene drives in Saccharomyces cerevisiae. *G3*, 8, 999–1018, 2018.

9. Basgall EM, Goetting SC, Goeckel ME, Giersch RM, Roggenkamp E, Schrock MN, Halloran M, Finnigan GC. Gene drive inhibition by the anti-CRISPR proteins AcrIIA2 and AcrIIA4 in Saccharomyces cerevisiae. *Microbiology*, 164, 464–474, 2018.

10. Shapiro RS, Chavez A, Porter CBM, Hamblin M, Kaas CS, DiCarlo JE, Zeng G, Xu X, Revtovich A V., Kirienko N V., Wang Y, Church GM, Collins JJ. A CRISPR–Cas9-based gene drive platform for genetic interaction analysis in Candida albicans. *Nat Microbiol*, 3, 73–82, 2018.

11. Oberhofer G, Ivy T, Hay BA. Behavior of homing endonuclease gene drives targeting genes required for viability or female fertility with multiplexed guide RNAs. *Proc Natl Acad Sci*, 115, E9343–E9352, 2018.

12. KaramiNejadRanjbar M, Eckermann KN, Ahmed HMM, Sánchez C. HM, Dippel S, Marshall JM, Wimmer EA. Consequences of resistance evolution in a Cas9-based sex-conversion suppression gene drive for insect pest management. *Proc Natl Acad Sci*, 201713825, 2018.

13. Chae D, Lee J, Lee N, Park K, Moon SJ, Kim HH. Chemical controllable gene drive in Drosophila. *ACS Synth Biol*, 2020.

14. López Del Amo V, Leger BS, Cox KJ, Gill S, Bishop AL, Scanlon GD, Walker JA, Gantz VM, Choudhary A. Small-Molecule Control of Super-Mendelian Inheritance in Gene Drives. *Cell Rep*, 31, 107841, 2020.

15. Kandul NP, Liu J, Buchman A, Gantz VM, Bier E, Akbari OS. Assessment of a split homing based gene drive for efficient knockout of multiple genes. *G3 Genes, Genomes, Genet*, 10, 827–837, 2020.

16. Xu X-RS, Bulger EA, Gantz VM, Klanseck C, Heimler SR, Auradkar A, Bennett JB, Miller LA, Leahy S, Juste SS, Buchman A, Akbari OS, Marshall JM, Bier E. Active Genetic Neutralizing Elements for Halting or Deleting Gene Drives. *Mol Cell*, 2020.

17. López Del Amo V, Bishop AL, Sánchez C HM, Bennett JB, Feng X, Marshall JM, Bier E, Gantz VM. A transcomplementing gene drive provides a flexible platform for laboratory investigation and potential field deployment. *Nat Commun*, 11, 1–12, 2020.

18. Gantz VM, Bier E. Genome editing. The mutagenic chain reaction: a method for converting heterozygous to homozygous mutations. *Science (80-)*, 348, 442–444, 2015.

19. Champer J, Reeves R, Oh SY, Liu C, Liu J, Clark AG, Messer PW. Novel CRISPR/Cas9 gene drive constructs reveal insights into mechanisms of resistance allele formation and drive efficiency in genetically diverse populations. *PLoS Genet*, 13, e1006796, 2017.

20. Champer J, Liu J, Oh SY, Reeves R, Luthra A, Oakes N, Clark AG, Messer PW. Reducing resistance allele formation in CRISPR gene drive. *Proc Natl Acad Sci*, 115, 5522–5527, 2018.

21. Champer J, Wen Z, Luthra A, Reeves R, Chung J, Liu C, Lee YL, Liu J, Yang E, Messer PW, Clark AG. CRISPR gene drive efficiency and resistance rate is highly heritable with no common genetic loci of large effect. *Genetics*, 2019.

22. Champer J, Chung J, Lee YL, Liu C, Yang E, Wen Z, Clark AG, Messer PW. Molecular safeguarding of CRISPR gene drive experiments. *Elife*, 8, 2019.

23. Guichard A, Haque T, Bobik M, Xu X-RS, Klanseck C, Kushwah RBS, Berni M, Kaduskar B, Gantz VM, Bier E. Efficient allelic-drive in Drosophila. *Nat Commun*, 10,

1640, 2019.

24. Champer SE, Oh SY, Liu C, Wen Z, Clark AG, Messer PW, Champer J. Computational and experimental performance of CRISPR homing gene drive strategies with multiplexed gRNAs. *Sci Adv*, 6, eaaz0525, 2020.

25. Champer J, Yang E, Lee E, Liu J, Clark AG, Messer PW. A CRISPR homing gene drive targeting a haplolethal gene removes resistance alleles and successfully spreads through a cage population. *Proc Natl Acad Sci*, 2020.

26. Hammond AM, Kyrou K, Bruttini M, North A, Galizi R, Karlsson X, Kranjc N, Carpi FM, D'Aurizio R, Crisanti A, Nolan T. The creation and selection of mutations resistant to a gene drive over multiple generations in the malaria mosquito. *PLOS Genet*, 13, e1007039, 2017.

27. Hammond A, Galizi R, Kyrou K, Simoni A, Siniscalchi C, Katsanos D, Gribble M, Baker D, Marois E, Russell S, Burt A, Windbichler N, Crisanti A, Nolan T. A CRISPR-Cas9 gene drive system targeting female reproduction in the malaria mosquito vector Anopheles gambiae. *Nat Biotechnol*, 34, 78–83, 2015.

28. Gantz VM, Jasinskiene N, Tatarenkova O, Fazekas A, Macias VM, Bier E, James AA. Highly efficient Cas9-mediated gene drive for population modification of the malaria vector mosquito Anopheles stephensi. *Proc Natl Acad Sci U S A*, 112, E6736–E6743, 2015.

29. Kyrou K, Hammond AM, Galizi R, Kranjc N, Burt A, Beaghton AK, Nolan T, Crisanti A. A CRISPR-Cas9 gene drive targeting doublesex causes complete population suppression in caged Anopheles gambiae mosquitoes. *Nat Biotechnol*, 2018.

30. Pham TB, Phong CH, Bennett JB, Hwang K, Jasinskiene N, Parker K, Stillinger D, Marshall JM, Carballar-Lejarazú R, James AA. Experimental population modification of the malaria vector mosquito, Anopheles stephensi. *PLOS Genet*, 15, e1008440, 2019.

31. Li M, Yang T, Kandul NP, Bui M, Gamez S, Raban R, Bennett J, Sánchez C HM, Lanzaro GC, Schmidt H, Lee Y, Marshall JM, Akbari OS. Development of a confinable gene drive system in the human disease vector aedes aegypti. *Elife*, 9, 2020.

32. Simoni A, Hammond AM, Beaghton AK, Galizi R, Taxiarchi C, Kyrou K, Meacci D, Gribble M, Morselli G, Burt A, Nolan T, Crisanti A. A male-biased sex-distorter gene drive for the human malaria vector Anopheles gambiae. *Nat Biotechnol*, 1–7, 2020.

33. Carballar-Lejarazú R, Ogaugwu C, Tushar T, Kelsey A, Pham TB, Murphy J, Schmidt H, Lee Y, Lanzaro GC, James AA. Next-generation gene drive for population modification of the malaria vector mosquito, Anopheles gambiae. *Proc Natl Acad Sci U S A*, 117, 22805–22814, 2020.

34. Hammond A, Karlsson X, Morianou I, Kyrou K, Beaghton A, Gribble M, Kranjc N, Galizi R, Burt A, Crisanti A, Nolan T. Regulating the expression of gene drives is key to increasing their invasive potential and the mitigation of resistance. *PLOS Genet*, 17, e1009321, 2021.

35. Adolfi A, Gantz VM, Jasinskiene N, Lee HF, Hwang K, Terradas G, Bulger EA, Ramaiah A, Bennett JB, Emerson JJ, Marshall JM, Bier E, James AA. Efficient population modification gene-drive rescue system in the malaria mosquito Anopheles stephensi. *Nat Commun*, 11, 1–13, 2020.

36. Grunwald HA, Gantz VM, Poplawski G, Xu X-RS, Bier E, Cooper KL. Super-Mendelian inheritance mediated by CRISPR–Cas9 in the female mouse germline. *Nature*, 566, 105–109, 2019.

37.   Champer J, Kim IK, Champer SE, Clark AG, Messer PW. Performance analysis of novel toxin-antidote CRISPR gene drive systems. *BMC Biol*, 18, 27, 2020.
38.   Champer J, Lee E, Yang E, Liu C, Clark AG, Messer PW. A toxin-antidote CRISPR gene drive system for regional population modification. *Nat Commun*, 11, 1082, 2020.
39.   Oberhofer G, Ivy T, Hay BA. Cleave and Rescue, a novel selfish genetic element and general strategy for gene drive. *Proc Natl Acad Sci*, 201816928, 2019.
40.   Champer J, Champer SE, Kim IK, Clark AG, Messer PW. Design and analysis of CRISPR-based underdominance toxin-antidote gene drives. *Evol Appl*, eva.13180, 2020.
41.   Noble C, Min J, Olejarz J, Buchthal J, Chavez A, Smidler AL, DeBenedictis EA, Church GM, Nowak MA, Esvelt KM. Daisy-chain gene drives for the alteration of local populations. *Proc Natl Acad Sci*, 201716358, 2019.
42.   Dhole S, Lloyd AL, Gould F. Tethered homing gene drives: a new design for spatially restricted population replacement and suppression. *Evol Appl*, eva.12827, 2019.
43.   Dhole S, Vella MR, Lloyd AL, Gould F. Invasion and migration of spatially self-limiting gene drives: A comparative analysis. *Evol Appl*, 11, 794–808, 2018.
44.   Liu J, Champer J, Langmüller AM, Liu C, Chung J, Reeves R, Luthra A, Lee YL, Vaughn AH, Clark AG, Messer PW. Maximum likelihood estimation of fitness components in experimental evolution. *Genetics*, 211, 1005–1017, 2019.
45.   Yang E, Metzloff M, Langmüller AM, Clark AG, Messer PW, Champer J. A homing suppression gene drive with multiplexed gRNAs maintains high drive conversion efficiency and avoids functional resistance alleles. *bioRxiv*, 2021.05.27.446071, 2021.
46.   North AR, Burt A, Godfray HCJ. Modelling the suppression of a malaria vector using a CRISPR-Cas9 gene drive to reduce female fertility. *BMC Biol*, 18, 98, 2020.
47.   Champer J, Kim IK, Champer SE, Clark AG, Messer PW. Suppression gene drive in continuous space can result in unstable persistence of both drive and wild-type alleles. *Mol Ecol*, 30, 1086–1101, 2021.
48.   Oberhofer G, Ivy T, Hay BA. Gene drive and resilience through renewal with next generation Cleave and Rescue selfish genetic elements. *Proc Natl Acad Sci U S A*, 117, 9013–9021, 2020.
49.   Chatterjee P, Jakimo N, Jacobson JM. Minimal PAM specificity of a highly similar SpCas9 ortholog. *Sci Adv*, 4, eaau0766, 2018.
50.   Chatterjee P, Jakimo N, Lee J, Amrani N, Rodríguez T, Koseki SRT, Tysinger E, Qing R, Hao S, Sontheimer EJ, Jacobson J. An engineered ScCas9 with broad PAM range and high specificity and activity. *Nat Biotechnol*, 1–5, 2020.
51.   Nishimasu H, Shi X, Ishiguro S, Gao L, Hirano S, Okazaki S, Noda T, Abudayyeh OO, Gootenberg JS, Mori H, Oura S, Holmes B, Tanaka M, Seki M, Hirano H, Aburatani H, Ishitani R, Ikawa M, Yachie N, Zhang F, Nureki O. Engineered CRISPR-Cas9 nuclease with expanded targeting space. *Science (80-)*, 361, 1259–1262, 2018.

SUPPLEMENTARY INFORMATION

The following table shows the DNA fragments used for Gibson Assembly of the plasmid. PCR products are shown with the oligonucleotide primer pair used, and plasmid digest is shown with the restriction enzymes used.

TAREhNU2G	*Template*	*Oligo/Enzyme 1*	*Oligo/Enzyme 2*
PCR Product	BHDaaN	SV40_U6_F	NosCas9_1_R
PCR Product	BHDaaN	Cas9_2_F	Nos3_3x_R
Plasmid Digest	EGDh2	HindIII	MluI

Construction primers
```
Cas9_2_F: AAACAGCTCAAGAGGCGCC
Nos3_3x_R: TTCAATTAGAGCTAATTCAATTAGGATCCAAGCTTTCCTTCCTGGCCCTTTTCGA
NosCas9_1_R: TCCTGTATATCGGCGCCTCTT
SV40_U6_F: GGAGCAATCACAGGTGAGCAAAAAAACGCGTTAAGATACATTGATGAGTTTGGACAAACC
```

Sequencing primers
```
EGFP_S_F: AGCGCACCATCTTCTTCAAGG
h3utr_S_F: AAGGACCTTCATCAGACGCAC
hCut_S_F: CCAAATTGGAAAAGGCCGACA
hCut_S_R: AACATGGGTTGCTGTTGTGC
```

FIGURE S1. Tethered Drive Mechanism. a) A regionally confined TARE drive is first released. Cas9 activity occurs in the germline and embryo of drive-carrying individuals, disrupting wild-type alleles. Individuals with two disrupted alleles are not viable. Over several generations, the TARE drive comes to fixation. Common events are indicated with solid arrows. **b)** A homing drive is released into the population of TARE-modified individuals. Cas9 activity leads to drive conversion by homology-directed repair or to disrupted alleles. In the haplolethal homing drive, genotypes with any disrupted alleles are not viable. The suppression drive targets a haplosufficient gene, and individuals require at least one wild-type allele to be viable. **c)** Homing drive activity cannot occur in individuals that do not express Cas9.

FIGURE S2. Egg-to-Pupae Viability for Haplolethal Homing Drive. Individuals heterozygous for the TARE drive and the haplolethal homing drive were crossed with w^{1118} individuals, and eggs and pupae were counted as well as eclosed adults. The size of the dots is proportional to the number of eggs from a single female. Rate estimates and SEM are indicated.

FIGURE S3. Population Sizes for Cage Study. The population is shown for each generation. Arrows indicate generations where homing drives were released in the first two cages.

Table S1. Combined maximum likelihood parameter estimates from cage populations

Log-likelihood shows a relative probability (higher values indicate a better model fit)
AICc: Akaike information criterion, corrected (low values indicate a better match of the model without overfitting)
Effective population size refers to the percent of census size, which varies with each generation (the average was 3491)
Fitness values are for drive homozygotes (with multiplicative fitness per allele). A value of 1 is equivalent to wild-type fitness, and "1" indicates that the parameter was fixed

Fitness cost model	Log-likelihood	AICc	Effective population size	Mating fitness	Fecundity fitness	Viability fitness
None	90.5	-178.86	1.8%	1	1	1
All	92.4	-175.85	2.0%	0.98	0.891	0.946
All with mating = fecundity	93.2	-179.78	2.0%	0.879	0.879	1.00
Mating and fecundity	93.2	-179.80	2.0%	0.847	0.915	1
Mating and viability	93.1	-179.72	2.0%	0.783	1	1.00
Fecundity and viability	93.0	-179.43	2.0%	1	0.768	1.00
Mating	93.1	-181.99	2.0%	0.783	1	1
Fecundity	93.0	-181.70	2.0%	1	0.768	1
Viability	91.5	-178.76	1.9%	1	1	0.898
Mating = fecundity	93.2	***-182.05***	2.0%	0.879	0.879	1