Molecular foundations of Precambrian uniformitarianism

Mateusz Kędzior[1], Amanda K. Garcia[1], Meng Li[2], Arnaud Taton[3], Zachary R. Adam[4], Jodi N. Young[3] and Betül Kaçar[1,4*]

[1]Department of Molecular and Cellular Biology, University of Arizona, Tucson, AZ
[2]School of Oceanography, University of Washington, Seattle, WA
[3]Division of Biological Sciences, University of California San Diego, La Jolla, CA
[4]Department of Planetary Sciences, Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ
*Corresponding author: betul@arizona.edu

Keywords: RuBisCO, carbon fixation, carbon isotope record, molecular paleobiology, paleogenetics, ancestral sequence reconstruction

Abstract

Uniformitarian assumptions underlie the oldest evidence for living organisms on Earth, the distinct isotope fractionation between inorganic and organic carbon. Aside from a handful of compelling deviations, the $^{13}C/^{12}C$ isotopic mean of preserved organic carbon ($\delta^{13}C_{org}$) has remained remarkably unchanged through time. RuBisCO is the principal carboxylase/oxygenase biomolecular component that is thought to primarily account for the generation of these distinct carbon isotopic signals. However, it is difficult to reconcile a mostly unchanging mean $\delta^{13}C_{org}$ with several known factors that can affect the isotope fractionation of RuBisCO, such as atmospheric composition and the amino acid composition of the enzyme itself, which have each changed markedly over Earth history. Here we report the resurrection and genetic incorporation of a Precambrian-age, Form IB RuBisCO in a modern cyanobacterial host. The isotopic composition of biomass relative to CO_2 (ε_p) in ancestral and control strains were much greater when grown under Precambrian CO_2 concentrations compared to modern ambient levels, but displaying values within a nominal envelope of modern-day RuBisCO IB enzyme variants. We infer that these isotopic differences derive indirectly from the decreased fitness of the AncIB strain, which includes diminished growth capacity and total cell RuBisCO activity. We argue that to answer the greatest questions of deep-time paleobiology, ancient biogeochemical signals should be reproduced in the laboratory through the synthesis of the geologic record with experimentally-derived constraints on underlying ancient molecular biology.

Significance Statement

The earliest geochemical indicators of microbes, and the enzymes that powered them, extend back almost 3.8 billion years on our planet. Paleobiologists often attempt to understand these indicators by assuming that the behaviors of modern microbes and enzymes are consistent (uniform) with those of their predecessors. This assumption seems uncomfortably at odds with the great variability of Earth's

environment and its highly adaptive microbes. Here we examine whether a uniformitarian assumption for an enzyme thought to generate these indicators, RuBisCO, can be corroborated by independently studying the history of changes recorded within RuBisCO's genetic sequences. We outline a new approach to paleobiology that informatively links molecule-level evolutionary changes with planet-level geochemical conditions in Earth's deep past.

Introduction

The history of life on Earth may be broadly subdivided into two, mutually exclusive macroevolutionary phases, the Phanerozoic and the Precambrian (1). The Phanerozoic (~542 million years ago to present) may be characterized by physiological and anatomical innovations and their resultant effects on ecosystem expansion, trophic tiering, and sociality (2-4). Hard- and soft-anatomical preservation provides a rich template for reconstructing Phanerozoic adaptive trends, correlating them with geographical and climatological changes (5, 6), and for testing observed diversity trends against possible systematic effects of preservation bias (7-9). By contrast, the Precambrian (the ~4 billion years preceding the Phanerozoic) is primarily characterized by genetic and metabolic biomolecular innovations, traded amongst microscopic organisms of uncertain phylogenetic assignment (10, 11). The Precambrian record of evolutionary change appears to be cryptic and may have been comparatively static. This may be attributable to macroevolutionary dynamics that were distinctly non-Phanerozoic, or it may merely indicate a lack of direct paleontological and geological evidence of the specific timing and extent of intermediate biomolecular adaptive steps (1).

Comparative analyses of extant organisms have traditionally been the most informative means of interpreting the scant direct evidence of Precambrian life, but such analyses inevitably face pitfalls. A reasonable null hypothesis is that evolution is largely a uniformitarian process, such that rates or tempos may change but the underlying processes or modes of evolutionary change (as established through observations of the more comprehensive Phanerozoic record) are likely invariant in deep time (12). Uniformitarian assumptions of ancient biology inferred from extant or Phanerozoic phenotypes are often employed to make sense of the Precambrian record, including body plan function (13, 14), cladistic assignment (15-18), and isotope biosignature traces (19-23). However, more recent studies complicate the use of uniformitarian assumptions, namely indicating that modes of biological variation can actually vary through time (12). A major crux of the problem is that even the simplest modern organisms, as well as the macromolecules that compose them, differ from their ancient predecessors in having been shaped by the cumulative effects of billions of years of Earth-life co-evolution and ecosystem upheaval. For this reason, Precambrian functional or sequence comparisons may be of limited utility at organismal or biomolecular levels of adaptation, undercutting interpretations made possible by uniformitarian assumptions. Novel experimental approaches may help to distinguish inferred paleobiological phenotypes from characteristically modern adaptive overprints.

The interpretation of the Precambrian carbon isotope record, comprising the oldest signatures of life on Earth, may be aided by novel experimental constraints on ancient phenotypes. Interpretations of this record is conventionally subject to uniformitarian assumptions regarding ancient biogeochemistry (19, 24). The $^{13}C/^{12}C$ isotopic mean of preserved organic carbon ($\delta^{13}C_{org} \approx -25‰$) has remained notably static over geologic time (19, 21, 24), and is leveraged as a general signature of ancient biological activity (25-

27). Given its role in the Calvin-Benson-Bassham cycle—likely the predominant mode of carbon fixation
for much of Earth history (19)—the majority of contextual information used to assess how carbon isotope
biosignatures might have been generated over Earth's deep history comes from studies of the modern
enzyme RuBisCO (Ribulose 1,5-Bisphosphate (RuBP) Carboxylase/Oxygenase, EC 4.1.1.39). RuBisCO
catalyzes the uptake of inorganic CO_2 from the environment and facilitates CO_2 reduction and
incorporation into organic biomass. The $^{13}C/^{12}C$ isotopic fractionation of modern Form I RuBisCO variants
in photosynthetic organisms consistently measures ~-25‰ (28-31), approximately the same isotopic
difference observed between inorganic and organic carbon in the Precambrian rock record. The carbon
isotope discrimination of RuBisCO has therefore been presumed to have remained constant over the
history of life. Recent data, however, demonstrate that RuBisCO can produce significantly different carbon
isotope signatures within organic matter in response to external factors, such as levels of atmospheric
CO_2 and/or O_2 (32-35) or cellular carbon concentrating mechanisms, which affect the catalytic efficiency
of RuBisCO (36-38). Internal factors may also affect fractionation, such as single-point mutations that can
alter the interaction between RuBisCO and CO_2 (39). Given the sensitivity of RuBisCO to both external and
internal variables, it seems unlikely that ancestral forms under ancient environmental conditions
generated the same isotope fractionation signal as descendent homologs in modern organisms. An
experimental assessment that combines the phylogenetic history of RuBisCO with the study of intra- and
extracellular conditions may provide a more insightful basis for comparing extant and Precambrian carbon
isotope fractionation patterns.
104
The Form I clade of the RuBisCO phylogeny (and its macroevolutionary tractability afforded through green
plant, algal and cyanobacterial fossil lineages) makes it an exemplary paleomolecular system for assessing
uniformitarian assumptions applied to Precambrian biosignatures. Here, we establish an experimental
system for the reconstruction of ancestral biomolecules with which to interpret evidence of Precambrian
biological activity. Specifically, we report the resurrection and genetic incorporation of a phylogenetically
reconstructed, ancient Form IB RuBisCO variant in a modern strain of cyanobacteria, *Synechococcus*
elongatus PCC 7942 (thereafter *S. elongatus*) (40, 41). We compared expression and activity levels of
RuBisCO variants and the resulting changes in the growth of *S. elongatus* and isotope fractionation under
ambient air as well as a CO_2 concentration that reflects Precambrian conditions.
114

Results

Construction of a *S. elongatus* strain harboring ancestral RuBisCO. To experimentally investigate the
generation of carbon isotope biosignatures in deep time, we designed a paleomolecular system to
engineer computationally inferred, ancestral RuBisCO enzymes in extant cyanobacteria. We previously
reconstructed a comprehensive phylogenetic history of RuBisCO and inferred maximum-likelihood
ancestral RuBisCO large-subunit (RbcL) protein sequences (42) (Fig. 1A). For this study, we selected the
ancestor of the Form IB RuBisCO clade (cyanobacteria, green algae, and land plants) for laboratory
resurrection, designated "AncIB." Chlorophyte and land plant RuBisCO homologs are nested among
cyanobacterial sequences within the Form IB clade. The Form IB topology therefore recapitulates a
primary plastid endosymbiotic history from cyanobacterial to Chlorophyte ancestors (43-45) and
constrains the minimum age of ancestral Form IB to older than the Archaeplastida. As a conservative
estimate, AncIB is thus likely older than ~1 Ga (the age of the oldest well-characterized, crown-group red
and green algal fossils (16, 46)) and younger than maximum age estimates of cyanobacteria (~3 Ga, as

constrained by oxidized sediments potentially indicating the early presence of oxygenic photosynthesis
(47, 48)).

The ancestral AncIB and *S. elongatus* native RbcL proteins differ at 37 sites and share 92% amino acid
identity (Fig. 2). This site variation is evenly spread across the length of the protein, except for a highly
conserved region between approximately site 170 to 285 (site numbering here and hereafter based on
aligned WT *S. elongatus* RbcL; Fig. 2A) that constitutes a portion of the catalytic C-terminal domain and is
proximal to the L-L interface and active site. Critical residues for carboxylase activity, including the Lys-
198 site that binds CO_2, are conserved in AncIB. Homology modeling of AncIB using the *S. elongatus* RbcL
template (PDB: 1RBL (49)) indicates high structural conservation without predicted disruption to
secondary structure (Fig. 2B). The nucleotide sequence for the reconstructed AncIB RbcL protein was
codon-optimized for *S. elongatus* and cloned within a copy of the *rbc* operon into pSyn02 (50) for insertion
into the *S. elongatus* chromosomal neutral site 2 (NS2). The native *rbc* operon was subsequently deleted
to create the AncIB strain. In addition, we generated the control strain Syn01 harboring WT *rbcL* at NS2
(see Materials and Methods for full genome strategy). Thus, the AncIB and Syn01 strains carry a single
ectopic copy of the *rbc* operon at NS2 and only differ in the coding sequence of the large RuBisCO subunit
(Table 1).

Ancestral RuBisCO complements photoautotrophic growth of extant *S. elongatus*. We cultured wild-
type (WT) and engineered (AncIB and Syn01) strains of *S. elongatus* in both ambient air and 2% CO_2 to
evaluate the physiological impact of ancestral RuBisCO under estimated Precambrian CO_2 concentrations
(51) (Fig. 3A). The AncIB strain was capable of photoautotrophic growth in both ambient air and 2% CO_2.
Maximum growth rates for all strains were relatively comparable under each atmospheric condition
(averaging doubling times of ~15 to 20 hours), and generally increased under 2% CO_2 relative to air ($p <$
0.001; Table 2). The AncIB strain exhibited a significantly diminished maximum cell density ($OD_{750} \approx 5$)
relative to both WT and Syn02 strains ($OD_{750} \approx 8$; $p < 0.001$). No significant difference between WT and
Syn01 growth rate or maximum cell density was observed under ambient air or 2% CO_2. In addition, a
moderate increase in the midpoint time for the AncIB strain was observed under air, indicating lag in
growth (p < 0.01). These growth parameters taken together suggest decreased fitness of the AncIB strain
relative to WT and Syn01.

AncIB RuBisCO protein produced a more modest impact on the oxygen evolution of *S. elongatus* compared
to that on growth parameters. Cell suspensions were briefly incubated in the dark and subsequently
exposed to saturated light in an electrode chamber to detect evolution of molecular oxygen (normalized
to chlorophyll *a* concentration (52)). For cells sampled from cultures grown in air, there was no statistical
difference detected between any of the WT, Syn01, or AncIB strains (Fig. 3B). We did find a modest but
significant decrease in photosynthetic activity for AncIB relative to WT for cells cultured in 2% CO_2,
generating ~200 and ~320 nmol $O_2 \cdot h^{-1} \cdot \mu g^{-1}$ chlorophyll *a*, respectively. However, a slower O_2 evolution
rate was also observed for Syn01 at 2% CO_2.

Ancestral RuBisCO is overexpressed and less catalytically active relative to WT. We assessed the impact
of ancestral RuBisCO on gene expression at both the transcript and protein levels. *S. elongatus rbcL*
transcript was measured by quantitative reverse-transcription PCR (RT-qPCR) and normalized to that of
the *secA* reference gene (53). For strains cultured in ambient air, we found that the AncIB strain produced

a ~29-fold increase in *rbcL* transcript relative to WT or the control strain Syn01 ($p < 0.001$; Fig. 4A). The
magnitude of AncIB *rbcL* overexpression was lower in 2% CO_2, with a ~4 and ~2-fold increase observed
relative to WT and the control strain, respectively ($p < 0.001$).
176
RbcL protein was quantified for all *S. elongatus* strains by immunodetection using rabbit anti-RbcL
antibody. We found that the amount of RbcL protein was also increased in the AncIB strain by ~3-fold (p
< 0.05) and ~5-fold ($p < 0.001$) relative to WT or Syn01 in air and 2% CO_2, respectively (Fig. 4B). No
difference in RbcL quantity was detected between WT and Syn01 strains at either atmospheric condition.
Finally, we confirmed assembly of the hexadecameric L_8S_8 RuBisCO complex in the AncIB strain by native
PAGE and detection by anti-RbcL antibody (Fig. S1).
183
184 The total carboxylase activity of *S. elongatus* harboring ancestral RuBisCO was measured from crude cell
185 lysates. Activity was assessed by an *in vitro* spectrophotometric coupled-enzyme assay that measures
186 NADH oxidation and is reported as the RuBP consumption rate normalized to total soluble protein content
187 (54). For two sets of assays using either 2.5 mM or 5 mM HCO_3^-, AncIB lysate generated less than half the
188 carboxylase activity of WT lysate ($p < 0.05$) (Fig. 4C).
189
S. elongatus harboring ancestral RuBisCO produces greater carbon isotopic fractionation than wild-
type. We measured the carbon isotope discrimination of *S. elongatus* strains cultured in ambient air and
2% CO_2 to evaluate how the fractionation behavior of ancestral RuBisCO might influence the
interpretation of ancient isotopic biosignatures preserved in the geologic record. The $^{13}C/^{12}C$ carbon
isotope composition of *S. elongatus* was measured for biomass ($\delta^{13}C_{biomass}$) as well as dissolved inorganic
carbon (DIC; $\delta^{13}C_{DIC}$) in the growth medium (Table S1). We calculated the carbon isotope fractionation
associated with photosynthetic CO_2 fixation (ε_p) following Freeman and Hayes (33) after estimating $\delta^{13}C_{CO2}$
from measured $\delta^{13}C_{DIC}$ (55, 56) (see Materials and Methods).
198
Overall, ε_p values were greater for *S. elongatus* strains cultured in 2% CO_2 compared to ambient air. At 2%
CO_2, ε_p ranged between 21‰ and 26‰ compared to only 8‰ and 14‰ in air. We found that *S. elongatus*
engineered with AncIB RbcL had an ε_p ~5‰ greater than both WT and Syn01 when cultured in air ($p <$
0.001) and ~2‰ to 4‰ greater than WT and Syn01 when cultured at 2% CO_2 ($p < 0.01$) (Fig. 1C), though
these differences in ε_p appear driven by the inorganic carbon pool composition rather than biomass (Table
S1). A substantially smaller increase in ε_p (~1‰; $p < 0.001$) was observed for Syn01 relative to WT under
ambient air. Conversely, under 2% CO_2, Syn01 ε_p was decreased by ~2‰ ($p < 0.001$) relative to WT.
206

Discussion

208
Form IB ancestral RuBisCO, when engineered into an extant strain of *S. elongatus*, decreased both the
organismal growth capacity and the total cell RuBisCO activity. The genetic engineering strategy for
insertion of the AncIB *rbcL* sequence in the cyanobacterial genome cannot solely account for these
physiological differences since differences between the WT and Syn01 control strains were insignificant
for most measured properties, or not comparable in magnitude to differences between the WT and AncIB
strains. Rather, the observed differences for the AncIB strain indicate that the resultant phenotype is likely
attributable to the functionality of the ancestral enzyme itself. The unique phenotype of the AncIB strain
could be a direct result of the ancestral RbcL subunit or due to impediments to the assembly and activation

of a hexadecamer RuBisCO complex containing both the ancestral RbcL and modern RbcS subunits.
Another possibility is hampered integration of ancestral RbcL given a modern suite of associated proteins
required for RuBisCO folding and assembly (57). Further, while overexpression of the ancestral RbcL
occurred at the level of transcription and translation, the ancestral strain appears to have comparable
levels of assembled hexadecamer RuBisCO, suggesting lower rates of RuBisCO assembly (or faster
degradation). Even lower rates of measured total carboxylase activity suggest that the AncIB has
decreased efficacy, which could be directly representing ancestral RuBisCO kinetics as well as the
challenges associated with hybrid enzyme activation and activity.

Overexpression of the amount of ancestral RuBisCO shown by RT-qPCR and immunodetection assays is a
common physiological response to decreased enzymatic efficacy throughout the cell (e.g., (58, 59)).
However, expression compensation is insufficient to fully restore the extant WT phenotype, as indicated
by the reduced fitness (i.e., decreased maximum cell density, oxygen evolution, and total carboxylase
activity) of the ancestral strain harboring the ancestral RuBisCO compared to WT and Syn01.

There are few *in vitro* measurements of the kinetic isotope effect of Form IB RuBisCO in modern-day
organisms, but those available range from ~22‰ to 28‰ for cyanobacteria and C_3 plants, respectively
(28-31). The ~26‰ ε_p of AncIB strain biomass grown under 2% CO_2 suggests that the ancestral RuBisCO
also fractionates within this range. It has been theorized that RuBisCO kinetics have adapted in response
to CO_2 availability, either due to increased environmental CO_2 (36) or the emergence of CCMs (e.g., C_4
photosynthesis in plants (60)). Considering the positive relationship between enzymatic fractionation and
RuBisCO's specificity to CO_2 (36), reconstructing ancient RuBisCO kinetic isotope effect could provide
insights into the co-evolution of atmospheric concentrations of CO_2 and O_2 and carbon fixation strategies
during the Precambrian, in particular the emergence of carbon concentrating mechanisms (CCMs) (34,
61-63). This is relevant as precise estimates of the magnitude of atmospheric CO_2 elevation during the
Precambrian relative to the present, as well as the emergence and effectiveness of Precambrian CCMs,
are unknown.

We did observe statistically significant differences in ε_p of the ancestral strain compared to WT *S.*
elongatus under both ambient air and 2% CO_2 atmospheric conditions. However, upon inspection it
appears that the diminished activity of the AncIB strain is influencing the composition of the DIC pool
(both $\delta^{13}C$ and concentration) in our cultures, and it is in fact the differences in DIC composition driving
the calculated differences in ε_p. Though strains were harvested at similar cell densities, small differences
in cell concentrations at high densities can strongly influence carbonate chemistry of the media (64). The
lower CO_2 availability in the air treatment is more sensitive to cellular influence, resulting in a larger
difference in ε_p compared to the 2% CO_2 treatment. These differences in DIC are unlikely to be due to
experimental setup (e.g., variations in CO_2 bubbling), as biological replicates showed similar values.
Therefore, the differences in fractionation reported here are likely implicated indirectly with the less
efficient AncIB ancestral enzyme. Further comparative biomolecular characterization of AncIB and WT *S.*
elongatus RbcL forms is needed to determine the degree to which enzymatic inefficiencies are
contributing directly to the AncIB strain phenotype.

The observed carbon isotopic fractionation values corroborate a uniformitarian assumption for applying
the maximal range of extant organism-level isotope fractionation values to interpret deep time isotopic

biosignatures. There are, however, several potentially important contextual caveats. The most obvious is
that the isotopic fractionation values of all strains (WT, Syn01, and AncIB) are increased under simulated
Precambrian conditions with elevated CO_2. The Form IB ancestor represents predecessors that are at least
1 billion years old, but it is also genetically and functionally still likely to be very different from the putative
'root' or common ancestor of all RuBisCO variants that emerged much earlier. Reconstruction of older
ancestors may further expand this maximal envelope of RuBisCO-generated carbon fractionation, or it
may indicate that the extant maximal envelope is pervasive (and perhaps characteristic) across all
functional variants of RuBisCO.

Another important caveat lies in the observation that, whereas all strains produce increased isotopic
fractionation under elevated CO_2, the comparative difference between ancestral AncIB and WT RbcL
fractionation is relatively muted under 2% CO_2 relative to ambient air. One possibility is that elevated CO_2
brings the intrinsic fractionation properties of RuBisCO into relief (35, 65-67), at least compared to
fractionation effects deriving from the overlying organismal physiology. By contrast, in present-day
conditions, RuBisCO-mediated fractionation processes may be more significantly overprinted by physical
factors that can affect RuBisCO catalytic efficiency, including cellular diffusion of O_2/CO_2 or other factors
such as the presence of carbon concentration mechanisms.

There are many fundamental attributes of extant and ancestral metabolism for which the systemic effects
on biosignature production have yet to be characterized. Disentangling these effects is critical for
interpretation of the oldest biogeochemical record. A host cyanobacterium *S. elongatus* engineered with
a Form IB RbcL ancestor confirms that organism- and enzyme-level effects on biosignature production are
not always synonymous but differ in nuanced ways. These differences are contingent upon changes to
internal (cellular, physiological) and external (environmental) conditions that have demonstrably varied
over Earth's long history. Cyanobacteria, with well-characterized genetic and morphological features (40,
41, 61-63) and a tractable paleobiological history (18, 68), are ideal hosts for investigating a range of early
Precambrian metabolic processes (68-70).

Discernible trends (or steadfast consistencies) in metabolic outputs over macroevolutionary timescales
can lead to foundational uniformitarian approaches to deep time molecular paleobiology. The available
rock record becomes vanishingly sparse with greater age, but it is arguably well-sampled across key global-
scale and biotically relevant isotopic systems at least through the early Archaean. Greater geologic
sampling will therefore likely generate diminishing returns for shedding new light on deep time
paleobiological trends. Innovative approaches that can chart a comprehensive envelope of biomolecular
variability over time are a promising new means of reconciling coarse geochemical data with the nuance
and complexity of ancient biological activity.

The engineering of ancient-modern hybrid organisms and their characterization can be used to
complement the existing array of fossil remains, biogeochemical signatures, and modern organismal and
molecular proxies to assess and contextualize plausible ranges of Precambrian carbon isotope
biosignature production. Hybrid organisms may be particularly useful to disentangle the regulatory,
physiological, and inter- and intramolecular factors that have impacted isotope fractionation, none of
which are individually expressed in the geologic record. These factors must be systematically accounted

for when interpreting bulk fractionation signals, even if only to elucidate the evolutionary molecular
underpinnings of uniformitarian phenomena over geologic time.
306

Conclusions

After engineering a cyanobacterium with an ancient RuBisCO large protein subunit and cultivating it under
conditions that mimic those prevailing through much of the Precambrian, we found the resultant carbon
isotope fractionation to be within the range of organisms utilizing modern Form IB RuBisCO. The
underlying biomolecular and organismal adjustments made by the cell to accommodate the ancestral
gene were tracked, and we conclude that the small fractionation differences observed are likely
attributable indirectly to decreased fitness of the AncIB strain, which influenced the inorganic carbonate
chemistry of the media. The consistency of isotopic signatures generated by this strain indicates that
uniformitarian assumptions based on the range of phenotypes of modern RuBisCO variants may apply for
Precambrian environmental conditions, but that further study is warranted to discern organism- and
enzyme-level trends in carbon isotope fractionation that may extend deeper into the early Precambrian.
319

Materials and Methods

Inference of ancestral AncIB RbcL protein sequence. A RuBisCO RbcL phylogeny was reconstructed as
previously described (42). Briefly, RbcL orthologs were identified from the NCBI protein database by BLAST
(sequence dataset and the tree can be found at https://github.com/kacarlab/rubisco). Phylogenetic
analysis was performed by Phylobot (71), a web portal that integrates alignment, phylogenetic
reconstruction by RAxML (72), and ancestral sequence inference by PAML (73). A maximum likelihood
phylogeny was built using a MSAProbs alignment (74) and the best-fit PROTCATWAG model (75, 76),
determined by the Akaike information criterion (77). Ancestral states were reconstructed at each amino
acid site for all phylogenetic nodes, and gap characters were inferred according to Fitch's parsimony (78).
330

Cyanobacterial growth and maintenance. *S. elongatus* PCC 7942 strains were cultured in BG-11 medium
(79) as liquid cultures or on agar plates (1.5% (w/v) agar and 1 mM $Na_2S_2O_3 \cdot 5H_2O$). Liquid cultures were
grown at 30°C, continuous shaking at 120 rpm, sparged with ambient air or 2% CO_2, and with 115 µmol
photon$\cdot m^{-2} \cdot s^{-1}$ (with the exception of cultures used to prepare samples for O_2 evolution, which were grown
with 80 µmol photon$\cdot m^{-2} \cdot s^{-1}$). The 2% CO_2 gas mix was controlled by an environment chamber (Percival,
Cat. No. I36LLVLC8) with a CO_2 tank input. For recombinant strains, liquid and solid media were
supplemented with appropriate antibiotics: 2 µg$\cdot ml^{-1}$ Spectinomycin (Sp) plus 2 µg$\cdot ml^{-1}$ Streptomycin
(Sm), 5 µg$\cdot ml^{-1}$ Kanamycin (Km). Cyanobacterial growth was measured at an optical density of 750 nm
(OD_{750}) and growth parameters were estimated using the Growthcurver package for R (80) (Growthcurver
analysis script can be found at https://github.com/kacarlab/rubisco). Cultures were sampled at the middle
exponential growth phase, i.e., at an OD_{750} of ~2.5 (AncIB) or ~4.5 (WT and Syn01) for all subsequent
experiments.
343

344 **Genetic engineering of cyanobacteria.** Recombinant strains of *S. elongatus* were constructed by natural
345 transformation using standard protocols (81) with minor modifications (50). The plasmids and strains used
346 in this study are listed in Table 1. The construction of *S. elongatus* Syn01, carrying a single ectopic copy of
347 the *rbc* operon at NS2, as well as the plasmids pSyn01 and pSyn02 used to construct strain Syn01, were
348 described previously (50). The construction of strain AncIB was performed similarly to Syn01. Briefly,

pSyn03, which carries the AncIB nucleotide sequence within the entire *rbc* operon (including flanking sequences and homologous regions for recombination at neutral site 2 (NS2) of the *S. elongatus* chromosome), was transformed in *S. elongatus*. Transformation of WT *S. elongatus* with pSyn03 generated strain Syn03 carrying a second copy of the *rbc* operon at NS2. Strain Syn03 was subsequently transformed with pSyn01 to replace the native *rbc* operon with a spectinomycin/streptomycin resistance gene as described previously (50), producing strain AncIB. Transformants of Syn03 and AncIB were screened for complete segregation by colony PCR using primers F06, R06, F07, and R07 (Fig. S2; Table S2) and the strain sequences at the deletion and insertion sites were further verified by Sanger sequencing using the primers R07, F08, R08, F15, and F16 (Table S2). To construct plasmid pSyn03, pSyn02 excluding the *rbcL* coding sequence was PCR-amplified and linearized using primers F13/R13 and assembled with the AncIB *RbcL* coding sequence codon-optimized for *S. elongatus* and synthesized by Twist Bioscience. Both DNA fragments were assembled using the GeneArt™ Seamless Cloning and Assembly Kit (Invitrogen, Cat. No. A13288).

Analysis of *rbcL* expression by RT-qPCR. Cells were pelleted by centrifugation and resuspended in TE buffer (10 mM Tris, pH 8.0, 1 mM EDTA). Total RNA was extracted using the RNeasy® Protect Bacteria Mini Kit (QIAGEN, Cat. No. 74524). DNase I-treated RNA was then used in reverse transcription (RT) performed with the SuperScript™ IV First-Strand Synthesis System (Invitrogen, Cat. No. 18091050). F09/R09, F11/R11, F14/R14 pairs of qPCR primers (Table S2) were designed with Primer3Plus (http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi). The quality of cDNA and primer specificity was assessed by PCR using cDNA templates (RT positive reactions) and RT negative controls. qPCR was performed by the real-time thermal cycler qTOWER[3] G (Analytik Jena AG) using qPCRsoft software. The relative expression of native and AncIB *rbcL* was calculated as the average fold change normalized to the *secA* reference gene (53) using the delta-delta Ct method. The experiment was carried out using three biological replicates and three technical replicates.

Immunodetection of RbcL protein. Cells were pelleted by centrifugation and resuspended in 95°C TE buffer supplemented with 1% (w/v) SDS and incubated at 95°C for 10 min. The mixture was sonicated and centrifuged to remove cell debris. Total protein concentration in the crude cell lysates was measured using the Pierce™ BCA Protein Assay Kit (Thermo Scientific, Cat. No. 23225). Lysates containing 5 µg of total protein in Laemmli sample buffer were loaded onto a 6% (v/v) polyacrylamide stacking gel. Proteins were electrophoresed in a 12% polyacrylamide resolving gel and blotted onto a nitrocellulose membrane. Detection of RbcL and total protein load was performed as previously described (50). The densitometric analysis of RbcL signal intensity, normalized to total protein load, was performed with Quantity One® software (Bio-Rad) for three to six biological replicates.

Confirmation of RuBisCO assembly. Assembly of the RuBisCO large and small subunits into a hexadecameric complex in each strain was evaluated by native gel electrophoresis and immunodetection, as previously described (50). Immunodetection of the RuBisCO complex was performed for three biological replicates with the same primary and secondary antibodies that were used to detect RbcL, as described above.

Catalytic activity of RuBisCO. The activity of RuBisCO in cyanobacterial lysates was measured using a spectrophotometric coupled-enzyme assay that links this activity with the rate of NADH oxidation (82). Cell lysis and the activity assay were carried out as previously described (50) with either 2.5 mM or 5 mM $NaHCO_3$. After 20 min at 25 °C for activation of Rubisco, the reaction was initialized with the addition of ribulose 1,5-bisphosphate (RuBP) (0.5 mM) and the absorbance at 340 nm was recorded using a Synergy

396   H1 plate reader (BioTek). RuBisCO activity was reported as the RuBP consumption rate normalized to total
397   soluble protein content. The assay was performed for three biological replicates.
398
Photosynthetic oxygen evolution rate. *S. elongatus* strain photosynthetic activity was assayed using a
Clark-type oxygen electrode chamber to measure the level of molecular oxygen produced in
cyanobacterial cultures. Cells were pelleted and resuspended in fresh BG-11 to an OD_{750} of ~1 following
De Porcelinis (83). Concentration of chlorophyll *a* (for normalization) was measured following the protocol
by Zavrel et al. (84). The remaining suspension was incubated in the dark for 20 min with gentle agitation.
Samples from each suspension were analyzed in an oxygen electrode chamber under saturated light, using
the Oxygraph+ System (Hansatech Instruments) equipped with the OxyTrace+ software. Oxygen evolution
rate was monitored for 10 min and expressed as nanomoles of molecular oxygen evolved per hour per
microgram of chlorophyll *a*. The assay was performed for three biological replicates.
408
Carbon isotope fractionation in bulk cyanobacterial biomass. Cells were pelleted by centrifugation and
washed in 10 mL of 10 mM NaCl (OD_{750} for Syn-1 ~ 4.5, OD_{750} for AncIB ~2.5). Pellets were then dried at
50°C. In parallel, the supernatant from centrifuged culture samples was sterilized through 0.2 μm filtration
for DIC isotopic analysis of growth media. Sterilized media was transferred to Exetainer vials leaving no
headspace and stored at 4°C until analysis. Isotopic analysis was performed for three biological replicates.
The carbon isotope composition of bulk biomass ($\delta^{13}C_{biomass}$) and DIC ($\delta^{13}C_{DIC}$) was determined at the UC
Davis Stable Isotope Facility. $\delta^{13}C_{biomass}$ was analyzed using a PDZ Europa ANCA-GSL elemental analyzer
interfaced to a PDZ Europa 20-20 isotope ratio mass spectrometer (Sercon Ltd.). DIC samples were
analyzed by gas evolution and composition was measured by a GasBench II system interfaced to a Delta
V Plus IRMS (Thermo Scientific). The carbon isotopic composition values were reported relative to the
Vienna PeeDee Belemnite standard (V-PDB):
420

$$\delta^{13}C_{sample} = \left(\frac{^{13}C / ^{12}C_{sample}}{^{13}C / ^{12}C_{V-PDB}} - 1 \right) \times 1000$$

421
422
423   The isotopic composition of dissolved molecular CO_2 ($\delta^{13}C_{CO2}$) was estimated from $\delta^{13}C_{DIC}$ following Rau et
424   al. (55) and Mook et al. (56):
425

$$\delta^{13}C_{CO_2} = \delta^{13}C_{DIC} + 23.644 - \frac{9701.5}{T_K}$$

426
427
428   The carbon isotope fractionation associated with photosynthetic CO_2 fixation (ε_p) was calculated relative
429   to $\delta^{13}C_{CO2}$ in the post-culture medium according to Freeman and Hayes (1992):
430

$$\varepsilon_p = \frac{\delta^{13}C_{CO_2} - \delta^{13}C_{biomass}}{1 + \delta^{13}C_{biomass} / 1000}$$

431
432
433   **Statistical analyses.** Results for experimental analyses were presented as the mean and the sample
434   standard deviation (1σ) values of at least three biological replicates. For comparisons of two groups,
435   statistical significance was analyzed by an unpaired, two-tailed *t*-test assuming equal variance. For
436   comparisons of three or more groups, significance was analyzed by one-way ANOVA and a post-hoc Tukey
437   HSD test.

Acknowledgements

We sincerely thank Emily Peñaherrera, Jenan Kharbush, Ryan Ward, Sky Dominguez, and the University of California-Davis Stable Isotope Facility for assistance. This work was supported by the National Aeronautics and Space Administration Early Career Faculty (ECF) Award No. 80NSSC19K1617 (BK), the National Science Foundation Emerging Frontiers Program Award No. 1724090 (BK), National Aeronautics and Space Administration Postdoctoral Fellowship (AKG), Simons Foundation Early Career Award No. 561645 (JNY and ML), and National Institute of General Medical Sciences of the National Institutes of Health award number R01GM118815 (AT, to James W. Golden at the University of California-San Diego).

References

1. N. J. Butterfield, Macroevolution and Macroecology through Deep Time. *Palaeontology* **50**, 41-55 (2007). https://doi.org/10.1111/j.1475-4983.2006.00613.x

2. B. Hannisdal, S. E. Peters, Phanerozoic Earth System Evolution and Marine Biodiversity. *Science* **334**, 1121-1124 (2011). https://doi.org/10.1126/science.1210695

3. R. K. Bambach, "Classes and Adaptive Variety: The Ecology of Diversification in Marine Faunas Through the Phanerozoic" in *Phanerozoic Diversity Patterns,* J. W. Valentine, Ed. (Princeton University Press, 1986), pp. 191-254.

4. D. J. Bottjer, W. I. Ausich, Phanerozoic development of tiering in soft substrata suspension-feeding communities. *Paleobiology* **12**, 400-420 (2016). https://doi.org/10.1017/s0094837300003134

5. J. J. Sepkoski, Biodiversity: Past, Present, and Future. *J Paleontol* **71**, 533-539 (2015). https://doi.org/10.1017/s0022336000040026

6. G. J. Vermeij, Adaptation, Versatility, and Evolution. *Syst Zool* **22** (1973). https://doi.org/10.2307/2412953

7. A. J. W. Hendy, "Taphonomic Overprints on Phanerozoic Trends in Biodiversity: Lithification and Other Secular Megabiases" in *Taphonomy,* P. A. Allison, D. J. Bottjer, Eds. (Springer, Dordrecht, 2010), pp. 19-77.

8. S. E. Peters, Geologic constraints on the macroevolutionary history of marine animals. *Proc Natl Acad Sci USA* **102**, 12326-12331 (2005). https://doi.org/10.1073/pnas.0502616102

9. S. E. Peters, M. Foote, Biodiversity in the Phanerozoic: a reinterpretation. *Paleobiology* **27**, 583-601 (2016). https://doi.org/10.1666/0094-8373(2001)027<0583:Bitpar>2.0.Co;2

10. A. H. Knoll, Paleobiological Perspectives on Early Microbial Evolution. *Cold Spring Harbor Perspectives in Biology* **7**, a018093 (2015). https://doi.org/10.1101/cshperspect.a018093

11. J. W. Schopf, Disparate rates, differing fates: tempo and mode of evolution changed from the Precambrian to the Phanerozoic. *Proc Natl Acad Sci USA* **91**, 6735-6742 (1994). https://doi.org/10.1073/pnas.91.15.6735

12. D. H. Erwin, Evolutionary uniformitarianism. *Dev Biol* **357**, 27-34 (2011). https://doi.org/10.1016/j.ydbio.2011.01.020

13. Z. R. Adam, M. L. Skidmore, D. W. Mogk, N. J. Butterfield, A Laurentian record of the earliest fossil eukaryotes. *Geology* **45**, 387-390 (2017). https://doi.org/10.1130/g38749.1

14. Z. Yin, M. Zhu, D. J. Bottjer, F. Zhao, P. Tafforeau, Meroblastic cleavage identifies some Ediacaran Doushantuo (China) embryo-like fossils as metazoans. *Geology* **44**, 735-738 (2016). https://doi.org/10.1130/g38262.1

15. N. J. Butterfield, *Bangiomorpha pubescens* n. gen., n. sp.: implications for the evolution of sex,
multicellularity, and the Mesoproterozoic/Neoproterozoic radiation of eukaryotes. *Paleobiology*
26, 386-404 (2000). https://doi.org/10.1666/0094-8373(2000)026<0386:Bpngns>2.0.Co;2
16. Q. Tang, K. Pang, X. Yuan, S. Xiao, A one-billion-year-old multicellular chlorophyte. *Nat Ecol Evol*
4, 543-549 (2020). https://doi.org/10.1038/s41559-020-1122-9
17. S. Golubic, L. Seong-Joo, Early cyanobacterial fossil record: preservation, palaeoenvironments
and identification. *Eur J Phycol* **34**, 339-348 (1999).
https://doi.org/10.1080/09670269910001736402
18. J. W. Schopf, The paleobiological record of photosynthesis. *Photosynth Res* **107**, 87-101 (2011).
https://doi.org/10.1007/s11120-010-9577-1
19. M. Schidlowski, Carbon isotopes as biogeochemical recorders of life over 3.8 Ga of Earth history:
evolution of a concept. *Precambrian Res* **106**, 117-134 (2001). https://doi.org/Doi
10.1016/S0301-9268(00)00128-5
20. B. L. Beard, Iron Isotope Biosignatures. *Science* **285**, 1889-1892 (1999).
https://doi.org/10.1126/science.285.5435.1889
21. A. K. Garcia, C. M. Cavanaugh, B. Kacar, The curious consistency of carbon biosignatures over
billions of years of Earth-life coevolution. *ISME J* (2021). https://doi.org/10.1038/s41396-021-
00971-5
22. A. D. Anbar, O. Rouxel, Metal Stable Isotopes in Paleoceanography. *Annu Rev Earth Pl Sc* **35**,
717-746 (2007). https://doi.org/10.1146/annurev.earth.34.031405.125029
23. J. W. Schopf *et al.*, Sulfur-cycling fossil bacteria from the 1.8-Ga Duck Creek Formation provide
promising evidence of evolution's null hypothesis. *Proc Natl Acad Sci USA* **112**, 2087-2092
(2015). https://doi.org/10.1073/pnas.1419241112
24. J. Krissansen-Totton, R. Buick, D. C. Catling, A statistical analysis of the carbon isotope record
from the Archean to Phanerozoic and implications for the rise of oxygen. *Am J Sci* **315**, 275-316
(2015). https://doi.org/10.2475/04.2015.01
25. M. Schidlowski, A 3,800-Million-Year Isotopic Record of Life from Carbon in Sedimentary-Rocks.
Nature **333**, 313-318 (1988). https://doi.org/10.1038/333313a0
26. J. W. Schopf, K. Kitajima, M. J. Spicuzza, A. B. Kudryavtsev, J. W. Valley, SIMS analyses of the
oldest known assemblage of microfossils document their taxon-correlated carbon isotope
compositions. *Proc Natl Acad Sci USA* **115**, 53-58 (2018).
https://doi.org/10.1073/pnas.1718063115
27. E. A. Bell, P. Boehnke, T. M. Harrison, W. L. Mao, Potentially biogenic carbon preserved in a 4.1
billion-year-old zircon. *Proc Natl Acad Sci USA* **112**, 14518-14521 (2015).
https://doi.org/10.1073/pnas.1517557112
28. K. M. Scott *et al.*, Kinetic isotope effect and biochemical characterization of form IA RubisCO
from the marine cyanobacterium *Prochlorococcus marinus* MIT9313. *Limnol Oceanogr* **52**, 2199-
2204 (2007).
29. R. D. Guy, M. L. Fogel, J. A. Berry, Photosynthetic fractionation of the stable isotopes of oxygen
and carbon. *Plant Physiol* **101**, 37-47 (1993). https://doi.org/10.1104/pp.101.1.37
30. S. von Caemmerer, Y. Tazoe, J. R. Evans, S. M. Whitney, Exploiting transplastomically modified
Rubisco to rapidly measure natural diversity in its carbon isotope discrimination using tuneable
diode laser spectroscopy. *J Exp Bot* **65**, 3759-3767 (2014). https://doi.org/10.1093/jxb/eru036
31. C. A. Roeske, M. H. O'Leary, Carbon isotope effects on the enzyme-catalysed carboxylation of
ribulose bisphosphate. *Biochemistry* **23**, 6275-6284 (1984).
32. M. Eichner, S. Thoms, S. A. Kranz, B. Rost, Cellular inorganic carbon fluxes in *Trichodesmium*: a
combined approach using measurements and modelling. *J Exp Bot* **66**, 749-759 (2015).
https://doi.org/10.1093/jxb/eru427

33. K. H. Freeman, J. M. Hayes, Fractionation of carbon isotopes by phytoplankton and estimates of
ancient CO_2 levels. *Global Biogeochem Cycles* **6**, 185-198 (1992).
https://doi.org/10.1029/92gb00190
34. S. J. Hurley, B. A. Wing, C. E. Jasper, N. C. Hill, J. C. Cameron, Carbon isotope evidence for the
global physiology of Proterozoic cyanobacteria. *Sci Adv* **7** (2021).
https://doi.org/10.1126/sciadv.abc8998
35. E. B. Wilkes, R. B. Y. Lee, H. L. O. McClelland, R. E. M. Rickaby, A. Pearson, Carbon isotope ratios
of coccolith–associated polysaccharides of *Emiliania huxleyi* as a function of growth rate and CO_2
concentration. *Org Geochem* **119**, 1-10 (2018).
https://doi.org/10.1016/j.orggeochem.2018.02.006
36. G. G. Tcherkez, G. D. Farquhar, T. J. Andrews, Despite slow catalysis and confused substrate
specificity, all ribulose bisphosphate carboxylases may be nearly perfectly optimized. *Proc Natl
Acad Sci USA* **103**, 7246-7251 (2006). https://doi.org/10.1073/pnas.0600605103
37. R. A. Studer, P. A. Christin, M. A. Williams, C. A. Orengo, Stability-activity tradeoffs constrain the
adaptive evolution of RubisCO. *Proc Natl Acad Sci USA* **111**, 2223-2228 (2014).
https://doi.org/10.1073/pnas.1310811111
38. A. I. Flamholz *et al.*, Revisiting trade-offs between Rubisco kinetic parameters. *Biochemistry* **58**,
3365-3376 (2019). https://doi.org/10.1021/acs.biochem.9b00237
39. D. B. McNevin *et al.*, Differences in carbon isotope discrimination of three variants of D-ribulose-
1,5-bisphosphate carboxylase/oxygenase reflect differences in their catalytic mechanisms. *J Biol
Chem* **282**, 36068-36076 (2007). https://doi.org/10.1074/jbc.M706274200
40. B. M. Berla *et al.*, Synthetic biology of cyanobacteria: unique challenges and opportunities. *Front
Microbiol* **4** (2013). https://doi.org/10.3389/fmicb.2013.00246
41. Y. Chen, C. K. Holtman, A. Taton, S. S. Golden, "Functional Analysis of the Synechococcus
elongatus PCC 7942 Genome" in *Functional Genomics and Evolution of Photosynthetic Systems,*
R. Burnap, W. Vermaas, Eds. (Springer, Dordrecht, 2012), pp. 119-137.
42. B. Kacar, V. Hanson-Smith, Z. R. Adam, N. Boekelheide, Constraining the timing of the Great
Oxidation Event within the Rubisco phylogenetic tree. *Geobiology* **15**, 628-640 (2017).
https://doi.org/10.1111/gbi.12243
43. H. S. Yoon, J. D. Hackett, C. Ciniglia, G. Pinto, D. Bhattacharya, A Molecular Timeline for the
Origin of Photosynthetic Eukaryotes. *Mol Biol Evol* **21**, 809-818 (2004).
https://doi.org/10.1093/molbev/msh075
44. S. Maruyama, E. Kim, "Evolution of Photosynthetic Eukaryotes; Current Opinion, Perplexity, and
a New Perspective" in *Symbiosis: Cellular, Molecular, Medical and Evolutionary Aspects,* M. Kloc,
Ed. (Springer, Cham, 2020), pp. 337-351.
45. P. M. Shih *et al.*, Biochemical characterization of predicted Precambrian RuBisCO. *Nat Commun*
7, 10382 (2016). https://doi.org/10.1038/ncomms10382
46. T. M. Gibson *et al.*, Precise age of *Bangiomorpha pubescens* dates the origin of eukaryotic
photosynthesis. *Geology* **46**, 135-138 (2017). https://doi.org/10.1130/g39829.1
47. S. A. Crowe *et al.*, Atmospheric oxygenation three billion years ago. *Nature* **501**, 535-538 (2013).
https://doi.org/10.1038/nature12426
48. A. D. Anbar *et al.*, A whiff of oxygen before the great oxidation event? *Science* **317**, 1903-1906
(2007). https://doi.org/10.1126/science.1140325
49. J. Newman, C. I. Brändén, T. A. Jones, Structure determination and refinement of ribulose 1,5-
bisphosphate carboxylase/oxygenase from *Synechococcus* PCC6301. *Acta Crystallogr D* **49**, 548-
560 (1993). https://doi.org/10.1107/s090744499300530x

50. A. K. Garcia *et al.*, System-level effects of CO_2 and RuBisCO concentration on carbon isotope
fractionation. bioRxiv [Preprint] (2021). https://doi.org/10.1101/2021.04.20.440233 (accessed
28 May 2021).
51. D. C. Catling, K. J. Zahnle, The Archean atmosphere. *Sci Adv* **6**, eaax1420 (2020).
https://doi.org/10.1126/sciadv.aax1420
52. T. Zavřel, M. A. Sinetova, D. Búzová, P. Literáková, J. Červený, Characterization of a model
cyanobacterium *Synechocystis* sp. PCC 6803 autotrophic growth in a flat-panel photobioreactor.
Eng Life Sci **15**, 122-132 (2015). https://doi.org/10.1002/elsc.201300165
53. E. Szekeres, C. Sicora, N. Dragos, B. Druga, Selection of proper reference genes for the
cyanobacterium Synechococcus PCC 7002 using real-time quantitative PCR. *FEMS Microbiol Lett*
359, 102-109 (2014). https://doi.org/10.1111/1574-6968.12574
54. D. S. Kubien, S. M. Whitney, P. V. Moore, L. K. Jesson, The biochemistry of Rubisco in Flaveria. *J
Exp Bot* **59**, 1767-1777 (2008). https://doi.org/10.1093/jxb/erm283
55. G. H. Rau, U. Riebesell, D. Wolf-Gladrow, A model of photosynthetic ^{13}C fractionation by marine
phytoplankton based on diffusive molecular CO_2 uptake. *Mar Ecol Prog Ser* **133**, 275-285 (1996).
https://doi.org/10.3354/meps133275
56. W. G. Mook, J. C. Bommerson, W. H. Staverman, Carbon isotope fractionation between
dissolved bicarbonate and gaseous carbon dioxide. *Earth Planet Sc Lett* **22**, 169-176 (1974).
https://doi.org/10.1016/0012-821x(74)90078-8
57. L. H. Gunn, E. Martin Avila, R. Birch, S. M. Whitney, The dependency of red Rubisco on its
cognate activase for enhancing plant photosynthesis and growth. *Proc Natl Acad Sci USA* **117**,
25890-25896 (2020). https://doi.org/10.1073/pnas.2011641117
58. B. Kacar, X. Ge, S. Sanyal, E. A. Gaucher, Experimental Evolution of Escherichia coli Harboring an
Ancient Translation Protein. *J Mol Evol* **84**, 69-84 (2017). https://doi.org/10.1007/s00239-017-
9781-0
59. M. Goldsmith, D. S. Tawfik, Potential role of phenotypic mutations in the evolution of protein
expression and stability. *Proc Natl Acad Sci USA* **106**, 6197-6202 (2009).
https://doi.org/10.1073/pnas.0809506106
60. P.-A. Christin *et al.*, Evolutionary Switch and Genetic Convergence on rbcL following the
Evolution of C4 Photosynthesis. *Mol Biol Evol* **25**, 2361-2368 (2008).
https://doi.org/10.1093/molbev/msn178
61. B. M. Long, B. Förster, S. B. Pulsford, G. D. Price, M. R. Badger, Rubisco proton production can
drive the elevation of CO_2 within condensates and carboxysomes. *Proc Natl Acad Sci USA* **118**
(2021). https://doi.org/10.1073/pnas.2014406118
62. R. Burnap, M. Hagemann, A. Kaplan, Regulation of CO_2 Concentrating Mechanism in
Cyanobacteria. *Life* **5**, 348-371 (2015). https://doi.org/10.3390/life5010348
63. Jeffrey C. Cameron, Steven C. Wilson, Susan L. Bernstein, Cheryl A. Kerfeld, Biogenesis of a
Bacterial Organelle: The Carboxysome Assembly Pathway. *Cell* **155**, 1131-1140 (2013).
https://doi.org/10.1016/j.cell.2013.10.044
64. D. Shi, Y. Xu, F. M. M. Morel, Effects of the pH/$p$$CO_2$ control method on medium chemistry and
phytoplankton growth. *Biogeosciences* **6**, 1199-1207 (2009). https://doi.org/10.5194/bg-6-1199-
2009
65. J. M. Hayes, Factors controlling 13C contents of sedimentary organic compounds: Principles and
evidence. *Mar Geol* **113**, 111-125 (1993). https://doi.org/10.1016/0025-3227(93)90153-m
66. R. R. Bidigare *et al.*, Consistent fractionation of ^{13}C in nature and in the laboratory: growth-rate
effects in some haptophyte algae. *Global Biogeochem Cycles* **11**, 279-292 (1997).
https://doi.org/10.1029/96gb03939

67. B. A. Schubert, A. H. Jahren, The effect of atmospheric CO_2 concentration on carbon isotope
fractionation in C_3 land plants. *Geochim Cosmochim Acta* **96**, 29-43 (2012).
https://doi.org/10.1016/j.gca.2012.08.003
68. B. Kacar, L. Guy, E. Smith, J. Baross, Resurrecting ancestral genes in bacteria to interpret ancient
biosignatures. *Philos Trans R Soc A* **375** (2017). https://doi.org/10.1098/rsta.2016.0352
69. A. K. Garcia, H. McShea, B. Kolaczkowski, B. Kacar, Reconstructing the evolutionary history of
nitrogenases: Evidence for ancestral molybdenum-cofactor utilization. *Geobiology* **18**, 394-411
(2020). https://doi.org/10.1111/gbi.12381
70. A. K. Garcia, B. Kacar, How to resurrect ancestral proteins as proxies for ancient
biogeochemistry. *Free Radic Biol Med* **140**, 260-269 (2019).
https://doi.org/10.1016/j.freeradbiomed.2019.03.033
71. V. Hanson-Smith, A. Johnson, PhyloBot: A Web Portal for Automated Phylogenetics, Ancestral
Sequence Reconstruction, and Exploration of Mutational Trajectories. *PLoS Comput Biol* **12**,
e1004976 (2016). https://doi.org/10.1371/journal.pcbi.1004976
72. A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of large
phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
https://doi.org/10.1093/bioinformatics/btu033
73. Z. Yang, PAML 4: phylogenetic analysis by maximum likelihood. *Mol Biol Evol* **24**, 1586-1591
(2007). https://doi.org/10.1093/molbev/msm088
74. Y. Liu, B. Schmidt, D. L. Maskell, MSAProbs: multiple sequence alignment based on pair hidden
Markov models and partition function posterior probabilities. *Bioinformatics* **26**, 1958-1964
(2010). https://doi.org/10.1093/bioinformatics/btq338
75. N. Lartillot, H. Philippe, A Bayesian mixture model for across-site heterogeneities in the amino-
acid replacement process. *Mol Biol Evol* **21**, 1095-1109 (2004).
https://doi.org/10.1093/molbev/msh112
76. S. Whelan, N. Goldman, A general empirical model of protein evolution derived from multiple
protein families using a maximum-likelihood approach. *Mol Biol Evol* **18**, 691-699 (2001).
https://doi.org/10.1093/oxfordjournals.molbev.a003851
77. F. Abascal, R. Zardoya, D. Posada, ProtTest: selection of best-fit models of protein evolution.
Bioinformatics **21**, 2104-2105 (2005). https://doi.org/10.1093/bioinformatics/bti263
78. W. M. Fitch, Toward Defining the Course of Evolution: Minimum Change for a Specific Tree
Topology. *Syst Zool* **20** (1971). https://doi.org/10.2307/2412116
79. R. Rippka, J. Deruelles, J. B. Waterbury, M. Herdman, R. Y. Stanier, Generic Assignments, Strain
Histories and Properties of Pure Cultures of Cyanobacteria. *Microbiology* **111**, 1-61 (1979).
https://doi.org/https://doi.org/10.1099/00221287-111-1-1
80. K. Sprouffske, A. Wagner, Growthcurver: an R package for obtaining interpretable metrics from
microbial growth curves. *BMC Bioinformatics* **17**, 172 (2016). https://doi.org/10.1186/s12859-
016-1016-7
81. E. M. Clerico, J. L. Ditty, S. S. Golden, Specialized techniques for site-directed mutagenesis in
cyanobacteria. *Methods Mol Biol* **362**, 155-171 (2007). https://doi.org/10.1007/978-1-59745-
257-1_11
82. D. S. Kubien, C. M. Brown, H. J. Kane, Quantifying the amount and activity of Rubisco in leaves.
Methods Mol Biol **684**, 349-362 (2011). https://doi.org/10.1007/978-1-60761-925-3_27
83. A. J. De Porcellinis *et al.*, Overexpression of bifunctional fructose-1,6-
bisphosphatase/sedoheptulose-1,7-bisphosphatase leads to enhanced photosynthesis and
global reprogramming of carbon metabolism in *Synechococcus* sp. PCC 7002. *Metabolic
Engineering* **47**, 170-183 (2018). https://doi.org/10.1016/j.ymben.2018.03.001

84.   T. Zavřel, M. Sinetova, J. Červený, Measurement of Chlorophyll *a* and Carotenoids Concentration
672   in Cyanobacteria. *Bio-Protocol* **5** (2015). https://doi.org/10.21769/BioProtoc.1467

Figures and Figure Legends

Figure 1. Reconstruction of ancestral RuBisCO biogeochemical signatures. (*A*) Maximum likelihood Form I RuBisCO RbcL phylogeny (derived from full RuBisCO phylogeny described in Kacar et al. (42)). Ancestral AncIB node and descendent Form IB clade highlighted in blue. Approximate likelihood ratio (aLR) branch support indicated by asterisks (**: >10, ****: >1000). Carbon isotope record figure adapted from Garcia et al. (21), with data from Schidlowski et al. (25) (grey) and Krissansen-Totton et al. (24) (dark grey). Approximate age range of AncIB indicated by blue field (see text for discussion). (*B*) Genetic engineering of *S. elongatus* strains. Strain Syn01 was constructed by inserting a second copy of the *rbc* operon in the chromosomal neutral site 2 (NS2). Strain AncIB was constructed by inserting the genetic sequence encoding for the ancestral AncIB *rbcL* within the NS2 *rbc* operon. The native *rbc* operon was removed in both strains Syn01 and AncIB. (*C*) Photosynthetic carbon isotope fractionation (ε_p) of *S. elongatus* strains in this study, cultured in ambient air or 2% CO_2. n = 3 for each data point and error bars indicate 1σ (error bars smaller than some datapoints).

692
Figure 2. Structure and sequence features of ancestral RuBisCO. (*A*) Amino acid sequence alignment
between ancestral AncIB and extant *S. elongatus* RbcL. Ancestral site variation relative to the *S. elongatus*
template is highlighted in red. (*B*) Modeled structure of the ancestral AncIB L_2 dimer (blue), aligned to the
active conformation of the extant *S. elongatus* L_8S_8 hexadecamer (grey; PDB: 1RBL (49)). Highlighted
residues in (*A*) are also highlighted in (*B*). Site numbering from extant *S. elongatus*. Conserved residues
are indicated by dots and secondary structure is indicated above the sequences (blue rectangle: α-helix;
blue arrow: β-sheet).
700

702
703 **Figure 3.** (*A*) Growth curves and (*B*) photosynthetic oxygen evolution of *S. elongatus* strains cultured in
704 ambient air or 2% CO_2. (*A, B*) n = 3 for each data point or bar and error bars indicate 1σ.
705
706

Figure 4. Expression and total cell activity of RuBisCO in *S. elongatus* strains. (*A*) Expression of *rbcL* in
Syn01 and AncIB detected by RT-qPCR (*secA* reference gene), relative to WT. (*B*) Immunodetection of
RbcL protein. *Top*, RbcL signal intensities normalized to those for total soluble protein load. *Bottom*,
Western blot showing RbcL protein detected by anti-RuBisCO antibody and total protein stain from
crude cell lysates (*C*) Total cell RuBisCO activity, measured with 2.5 mM and 5 mM HCO_3^- concentrations.
(*A-C*) n = 3 for each bar (except 2% CO_2 RbcL signal intensity data, n = 6) and error bars indicate 1σ.

 # Tables

719 **Table 1.** Strains and plasmids used in this study.

Strain or plasmid	Description/Genotype	Antibiotic resistance	Source/Reference
WT	Wild-type strain of *S. elongatus* PCC 7942	-	Susan S. Golden (UC San Diego)
Syn01	*S. elongatus* strain Syn02 with the native *rbc* operon removed: Syn02 and Δ(*rbcL-rbcS-purK)*::*aadA*.	Km, Sp+Sm	(50)
Syn02	*S. elongatus* PCC 7942 carrying a second copy of the *rbc* operon and flanking sequences at NS2: NS2::*aphI-rbcL-rbcS-purK-*Synpcc7942_1429-Synpcc7942_1430	Km	(50)
Syn03	*S. elongatus* PCC 7942 carrying a modified copy of the *rbc* operon (with the ancestral *rbcL*) and flanking sequences at NS2: NS2::*aphI-ancIB-rbcS-purK-*Synpcc7942_1429-Synpcc7942_1430	Km, Sp+Sm	This study
AncIB	*S. elongatus* strain Syn03 carrying the ancestral AncIB *rbcL* gene in the *rbc* operon copy at NS2 and having the native *rbc* operon replaced with a Sp/Sm resistance gene: Δ(*rbcL-rbcS-purK*)::*aadA*.	Km	This study
pSyn01	Plasmid to replace *S. elongatus'* native *rbc* operon (CP000100: 1479070-1482595) with a Sp/Sm resistance gene: Δ(*rbcL-rbcS-purK)*::*aadA*.	Sp+Sm	This study
pSyn02	Plasmid for recombination at NS2 of *S. elongatus* chromosome carrying the *rbc* operon including *rbcL*, *rbcS*, *purK,* and flanking sequences from *S. elongatus* PCC 7942 (CP000100: 1479071-1484283)	Km	(50)
pSyn03	pSyn02 in which the coding sequence of RbcL was replaced with the coding sequence of AncIB	Km	This study

720
721
722

723 **Table 2.** Growth parameters of *S. elongatus* strains. Values represent the mean of three replicates ± 1σ.
724 Asterisks indicate significance relative to WT for the same atmospheric condition, determined by one-
725 way ANOVA and post-hoc Tukey HSD tests (**: $p < 0.01$; ***: $p < 0.001$).
726

Strain	Atmosphere	Doubling time (h)	Midpoint time (h)	Maximum cell density (OD_{750})
WT	Air	18.9 ± 1.0	106.4 ± 2.8	7.7 ± 0.3
Syn01		18.6 ± 0.3	103.8 ± 1.9	7.7 ± 0.1
AncIB		20.5 ± 0.4	118.4 ± 2.1**	5.4 ± 0.2***
WT	2% CO_2	16.9 ± 1.5	110.8 ± 6.1	8.0 ± 0.3
Syn01		17.3 ± 1.0	113.9 ± 2.3	8.1 ± 0.5
AncIB		15.8 ± 0.5	117.8 ± 1.7	5.0 ± 0.4***

727
728
729

Supplemental Information

731
732

733
734   **Figure S1.** Immunodetection of assembled RuBisCO. Western blot showing the assembly of RbcL (WT and
735   AncIB) and RbcS into the L_8S_8 hexadecameric complex (520 kDa), detected by anti-RbcL antibody.
736
737

738
739   **Figure S2.** Genotyping of *S. elongatus* strains. Primers F06/R06 (Table S2) were used to confirm the *rbc*
740   operon insertion at the neutral site 2 (NS2), either with WT *rbcL* (for Syn01) or AncIB *rbcL* (*rbc* insert:
741   8,617 bp; no insert: 1,970 bp). Primers F07/R07 were used to confirm the presence or absence of the rbc
742   operon at the native site (*rbc* present at its native site: 5,140 bp; *rbc* knocked out and replaced with the
743   aadA gene: 3,342 bp).
744
745

746   **Table S1.** Isotopic composition of biomass and DIC in growth medium of *S. elongatus* cultures.

Strain	Atmosphere	$\delta^{13}C_{biomass}$ (‰)	$\delta^{13}C_{DIC}$ (‰)	$\delta^{13}C_{CO2}$ (‰)	[DIC] (mM)
WT	Air	-19.63 ± 0.07	-2.92 ± 0.02	-11.28 ± 0.02	5.14 ± 0.07
	2% CO_2	-26.41 ± 0.15	5.77 ± 0.30	-2.59 ± 0.30	7.35 ± 0.24
Syn01	Air	-20.20 ± 0.19	-2.63 ± 0.10	-11.00 ± 0.10	5.23 ± 0.02
	2% CO_2	-26.84 ± 0.10	3.29 ± 0.25	-5.07 ± 0.25	7.53 ± 0.34
AncIB	Air	-20.60 ± 0.08	1.26 ± 0.10	-7.10 ± 0.10	3.44 ± 0.05
	2% CO_2	-31.2 ± 0.34	3.15 ± 0.62	-5.21 ± 0.62	5.70 ± 0.70

749   **Table S2.** Primers used in this study.

Primer[a]	Sequence (5'-3')	Description
F06	GACAATCCTGTTCTCCGGCA	Genotyping *S. elongatus* strains to screen for the *rbc* operon insert (either with native or AncIB *rbcL*) at NS2 (PCR product size: with *rbc* insert – 8,617 bp, without *rbc* insert – 1,970 bp).
R06	ATCAACGCCGTACCCGTATC	
F07	GGAGTCAATTCTGCAAGAGC	Genotyping *S. elongatus* strains to confirm the presence or absence of the *rbc* operon at the native site (PCR product size: with the *rbc* operon – 5,140 bp, without the *rbc* operon – 3,342 bp) R07 was also used to sequence the native *rbc* operon deletion site.
R07	TCAAGCTCGGTCTACTGC	
F08	GAATGCTCCGCTGGACTTGC	Sequencing the *rbc* operon insertion site at NS2. F08 was also used to sequence the native *rbc* operon deletion site.
R08	TGTACTCGATTTGTGCAGCG	
F09	ACCACCTTGGCAAAATGGTG	qPCR analysis of the expression of *rbcL* (ID: Synpcc7942_1426) that encodes the RuBisCO large subunit.
R09	TTTGTCGCCTTCCAGTTTGC	
F11	ATTACCTGCGCGACAACATG	qPCR analysis of the expression of *secA* (reference gene, ID: Synpcc7942_0289) that encodes the preprotein translocase subunit SecA.
R11	TGCCCGCATGTATTTTTCGC	
F13	GGAGCCTCTGACTATCGCTGGGGGAG	Linearization of pSyn02 without the *rbcL* coding sequence.
R13	GTCGTCTCTCCCTAGAGATATG	
F14	AAACCGGCGAAATCAAAGGC	qPCR analysis of the expression of AncIB *rbcL* that encodes the AncIB RuBisCO large subunit.
R14	CGTGCATGATGATGGGCATG	
F15	TGTTTGAAGAAGGCAGCGTG	Sequencing the internal region of AncIB *rbcL* inserted at NS2.
F16	CCAAATGGTGCCGCGATAAC	

750   [a]F - forward, R – reverse