Longevity interventions temporally scale healthspan in *Caenorhabditis elegans*

Cyril Statzer[1§], Peter Reichert[2§], Jürg Dual[2*], and Collin Y. Ewald[1*]

1 Laboratory of Extracellular Matrix Regeneration, Institute of Translational Medicine,

Department of Health Sciences and Technology, ETH Zürich, Schwerzenbach CH-8603,

Switzerland

2 Eidgenössische Technische Hochschule Zürich, Department of Mechanical and Process

Engineering, Institute for Mechanical Systems, Zürich CH-8092, Switzerland

§ Authors contributed equally

*Corresponding authors: collin-ewald@ethz.ch (CYE) and dual@imes.mavt.ethz.ch (JD)

Keywords: acoustophoresis, microfluidics, healthy aging, maximum muscle strength,

healthspan, sickspan, aging biomarker, lifespan machine

Abstract

Longer lived individuals, such as centenarians or longevity mutants of model organisms, have later onsets and lower incidence rates of late-life morbidities or disabilities than the average population. However, whether increased lifespan is caused by a compression of the portion of life spent in a state of morbidity, i.e., "sickspan," is highly debated. It is unclear which health matrices are representative for measuring healthspan (time spent in good health); however, muscular performance is generally a good indicator for the health status in humans and a predictor for their mortality, regardless of the underlying cause. Here, we developed a novel microfluidic device that employs acoustophoretic force fields to quantify the maximum muscle strength and dynamic power in aging *C. elegans* populations. We found that longevity mutants have a delayed onset and lower declining rates in maximum muscle strength compared to wild type. Reconciling previous conflicting reports, we confirmed that certain longevity mutants exhibited a mild increase in relative sickspan measured by voluntary movement matrices, which is not the case when using our acoustophoretic force measurements, swim endurance, or other approaches. Using six different biomarkers for healthspan, we observed a time-dependent onset of morbidity, starting with a loss of stress resilience, a decline in dynamic power, and a decline in structural integrity, culminating finally in inactivity (lethargy) and a loss of mobility. We observed that a subset of aging biomarkers correlate with each other and maybe functionally interconnected. Surprisingly, we did not observe a compression of sickspan in longevity mutants but instead observed a temporal scaling of healthspan with diminishing returns for extreme lifespan extensions. Given the conservation of these longevity interventions, this raises the question of whether the healthspan of mammalian longevity interventions is temporally scaled as well.

Introduction

The continuously growing elderly population is projected to result in 1.5 billion people above the age of 65 globally by 2050 [1]. This poses a significant challenge since old age is the major risk factor for developing cancer, dementia, cardiovascular, and metabolic diseases [2], especially since people suffer for approximately 20% of their lifespan from one or multiple of these chronic illnesses, which are themselves accompanied by other late-life disabilities [2]. Current estimates indicate that delaying the onset of these chronic diseases by two years would save $7.1 trillion over the next fifty years in the US alone [3]. Therefore, major research efforts are dedicated to understanding how to increase the time spent in good health (*i.e.*, healthspan) and to postpone and compress the time spent suffering from age-related pathologies and chronic diseases (*i.e.,* sickspan) [2,4–6].

People that are more than one hundred years old, so-called centenarians, display a delayed onset and a lower incidence rate of late-life morbidities compared to people in the age bracket of 80 to 89 years [7–12]. Genome-wide association studies have shown associations between the exceptional longevity of centenarians and aging-related genes identified in model organisms [2,13,14]. Mutations in genes that promote longevity in model organisms, such as *C. elegans,* have been instrumental in identifying mechanisms that promote healthy aging [2,13–15].

A recent study has questioned this approach of using *C. elegans* longevity mutants to gain insights for promoting healthy aging or mechanisms that prolong healthspan [16]. Using four matrices (resilience to heat and oxidative stress, voluntary movement, and swimming performance) to assess the "health" status of aging *C. elegans*, they found that four commonly used longevity mutants outperformed wild type at any given timepoint at older ages, consistent with previous reports. However, compared to their maximum lifespan, longevity mutants displayed an increased sickspan-to-healthspan ratio compared to wild type [16]. Other studies have not observed an increase of sickspan in long-lived *C. elegans* mutants, except in the case of lower mobility or movement scores for the insulin/IGF-1 receptor longevity *daf-2(e1370)*

mutants [17–20]. A large part of the "prolonged sickspan" based on the motility of these *daf-*

2(e1370) mutants was attributed to behavioral preferences of food over exploration linked to

odr-10 gene expression [17] or improper dauer-like quiescence behavior [20–24]. Although all these

studies showed that sickspan is not increased in longevity mutants, the question remained about

how healthspan changes when the lifespan is extended. We hypothesized that using other health

matrices independent of voluntary or behavioral influences, such as physical properties of

muscular strength, which is one of the best predictors for all-cause mortality in humans [25], we

might be able to quantify the health trajectory of *C. elegans* longevity mutants.

Here we confirm that voluntary movement during aging declines and this fragility is not

extended in longevity mutants, except mildly in *daf-2* mutants, using high-resolution lifespan

and movement measurements on plates. Instead of assessing swimming speed, we measured

swimming endurance and found *daf-2* longevity mutants outperforming wild type during aging.

We developed a novel microfluidic device and applied acoustophoretic force fields to quantify

the maximum force and power of *C. elegans*. Using a high frequency and high power acoustic

force field, it becomes possible to set up a contactless, constant in time, and uniform force field

acting along the whole *C. elegans* body. Therefore, this force field challenges swimming *C.*

elegans in a similar way body-weight exercises do for humans in a gravity field. Furthermore,

the acoustic field showed to be suited for stimulating a swimming response of resting *C.*

elegans. All longevity mutants showed delayed onset of the decline in maximum force and

dynamic power during aging. We observed heterogeneity between individuals across all

genotypes in the onset of age-related phenotypes, several correlated phenotypes, and a time-

dependent occurrence of multiple disabilities. However, we did not find a compression of

sickspan, but rather a temporal scaling of healthspan relative to their maximal lifespan across

genotypes.

Results

Voluntary movement healthspan is proportionally increased by longevity interventions

To obtain highly quantitative data on lifespan and healthspan, we used a lifespan machine [26]. Here, we defined the "voluntary movement healthspan" as the time spent fast crawling and the "voluntary movement sickspan" as the time spent slow crawling or displaying minimal posture changes (see Materials and Methods for detailed definition). We chose *eat-2(ad1116)* as a genetic model for dietary restriction-mediated longevity, *glp-1(e2141)* as a genetic model for germ-stem-cell-less-mediated longevity, *daf-2(e1368)* and *daf-2(e1370)* as genetic models for reduced insulin/IGF-1 signaling mediated longevity. We cultured all animals at the same temperature (15°C) and in the same environment with the same food source, except *glp-1* that underwent a brief temperature upshift during development as in preparation for the lifespan assay. To avoid dauer-specific traits that occur in reduced insulin/IGF-1 signaling mutants [22], and to avoid pathogenicity from a bacterial food source [20], lifespans were run at 15°C on heat-killed bacteria. Thus, the experimental setup was designed to offer optimal conditions and was kept identical while *C. elegans* genotypes were varied.

As expected, we measured a significant increase in lifespan for these long-lived mutants compared to wild type (Fig. 1a; Supplementary Table 1, Supplementary Video 1). Interestingly, our data showed that the longer-lived the mutant was, the more prolonged was the voluntary movement healthspan (Fig. 1b). Therefore, interventions that increase lifespan also increase the time spent moving fast and actively.

Relative increase for both health- and sickspan in long-lived mutants

To better understand the rescaling of the time spent in frailty in these long-lived mutants, we analyzed the fraction of slow-moving animals per day. We observed gaussian activity distributions, which were shifted along the time axis for these longevity mutants (Fig. 1c). This delayed onset of the sickspan (Fig. 1c) is consistent with the prolonged healthspan of these

long-lived mutants (Fig. 1b). However, except for dietary restricted *eat-2* mutants, the width

and the area of the Gaussian distributions were bigger for long-lived mutants than wild type

(Fig. 1c), suggesting an overall increase of sickspan. Thus, based on voluntary movement

tracking, long-lived mutants display increased absolute health- and absolute sickspan compared

to wild type. Next, we asked whether the fraction spent in health- and sickspan during the

lifespan is altered. Wild-type animals spent 78% of their lifespan fast-moving and 22% slow-

moving (Fig. 1d). For long-lived mutants, we recorded about 70-79% of their lifespan are spent

fast-moving (Fig. 1d, Extended Fig. 1), suggesting no compression of sickspan but rather a

proportional scaling of both health- and sickspan relative to their lifespan.

Heterogeneity in the length of sickspan but a fixed onset of sickspan

Since we did not observe a compression of voluntary movement sickspan of the entire

population, we wondered whether individual animals that outlived their siblings would display

a compressed sickspan. When we compared the sickspan traces of individual *C. elegans* for

each genotype, we were surprised to measure such a vast heterogeneity (Fig. 1d), given that all

these individual animals of a population are genetically identical, consume the same food, and

are housed in the same environment. Only the *glp-1*-mediated longevity showed an overall

compression of individual sickspan traces (Fig. 1d). For comparison among these different

genotypes, we decided to use "relative age" by dividing lifespan curves into quartiles and

computing the health-to-sickspan ratio for each quartile (Fig. 1d insets; Extended Data Fig. 2).

Consistent with previous reports on wild type [27], we found that in the first quantile of the

lifespan curve, individual animals spent about 90% of their lifetime fast-moving and 10% slow-

moving, indicating that these animals die young with a compressed sickspan compared to the

last quartile wherein animals spent about 75% of their lifetime fast-moving and 25% slow-

moving (Fig. 1d insets; Extended Data Fig. 1). Remarkably, it looks like the onset of an

individual's sickspan is a fixed event starting approximately when the first 10% of the isogenic

population starts to die (Fig. 1d). This observation suggests that up to a certain time point, the

animal's physiological integrity is maintained. After this time point, there appears to be a

stochastic decay resulting in a heterogenous sickspan distribution. Viewing the data using this

alternative interpretation of a fixed onset of sickspan would explain why animals in the first

quartile of the lifespan curve die young and spend less time in poor health, while animals in the

last quartile of the lifespan curve die old and spend more time in poor health. Thus, the time

spent fast- vs. slow-moving seems to have a fixed onset in time.

Voluntary movement healthspan temporally scales with lifespan except in *daf-2* mutants

The model of a fixed onset-timepoint for frailty would suggest that longevity interventions

would simply delay the onset. To address this, we contrasted the number of days spent fast-

moving (healthspan) for each individual as a function of their time lived (lifespan in days; Fig.

1e). We found that the time lived correlated and predicted the time spent fast-moving with an

R squared of 0.7 for wild type and R squared ranging from 0.5 to 0.8 for the longevity mutants

(Fig. 1e). Furthermore, the *glp-1* with an R squared of 0.8 and *daf-2(e1368)* with an R squared

of 0.5 indicate lower or higher heterogeneity, respectively, compared to wild type (Fig. 1e).

This is also apparent in the increased or decreased spread of data points below the regression

line in Fig. 1e and by increased or decreased lengths of the individual sickspan traces in Fig.

1d, respectively. One interesting aspect to note is that individuals in quartile 2 and 3, which

expire in the middle of the lifespan curves, displayed shortened healthspan relative to their

lifespan, whereas individuals in the last quartile showed an extended healthspan relative to their

lifespan (Fig. 1e). This might be because sicker individuals simply died earlier, leading to an

enrichment of healthier-aging individuals in the last quartile (Extended Data Fig. 1, 2). Based

on the high R squared values for all genotypes, we applied a linear model to investigate the

relationship between health- and lifespan (Fig. 1e). Steeper linear regression lines compared to

wild type would indicate an increase in health- to lifespan ratio. The slopes of the linear model

were steeper for *eat-2* and *glp-1*, but less steep for the two *daf-2* mutants compared to wild type

(Fig. 1e), suggesting that *glp-1* and *eat-2* spent a larger fraction and *daf-2* mutants spent a

smaller fraction of their lifespan actively moving. Since slopes of linear models can be sensitive

to extreme values, we compared the population means of health- and lifespan across all

genotypes (Fig. 1f). When we extrapolated the mean healthspan to mean lifespan ratio of wild

type, we found that *eat-2* and *glp-1* were close to this extrapolated line, whereas the *daf-2*

mutants lacked approximately three days (i.e., 7%) of mean healthspan in respect to their mean

relative lifespan (Fig. 1f). Thus, we uncovered that the prolonged voluntary movement

healthspan temporally scales with the prolonged lifespan for each of these longevity mutants

except less stringently for *daf-2* mutants.

Exhaustion of Swimming endurance and stress resilience during aging

Thus far, our observations and interpretations on healthspan are based on the decline of

voluntary movement on culturing plates in the abundance of food. Certain genotypes like *daf-*

2(e1370) are less motivated to forage and display a more rapid decline in voluntary foraging

behavior compared to wild type leading to the interpretation of being less healthy [17]. In our

setting, this lower foraging behavior is less pronounced in *daf-2(e1370)* since they were

cultured at 15°C, an environment that leads to less dauer program activation [22]. We used a

second approach to assess health based on voluntary movement and conducted a swimming

assay in the absence of food. Placing *C. elegans* in a water-based physiological buffer (M9

buffer) stimulates active trashing. When animals were young, they swim for more than 30 hours

without a decline in activity levels (Fig. 2). In the case of older animals, this endurance capacity

declines but less rapidly in longevity mutants (Fig. 2). Similarly, exposing aging animals to the

oxidative-stress-inducing agent arsenite, we report a progressive decline in stress resilience of

wild-type animals (Fig. 2). This decline in stress resilience is shifted to chronological older

animals in long-lived strains compared to wild type (Fig. 2). For both endurance swimming and

stress resilience, aging *daf-2* mutants surpassed chronological age-matched wild-type animals

(Fig. 2). Taken together, these results suggest that the observed deficit in healthspan of *daf-2*

mutants using a food-dependent movement assay might be largely due to their preferences in

lower foraging or exploratory behavior [17] rather than a loss of capability or frailty [16]. Moreover,

the progressive decline in endurance swimming and stress resilience during aging occurs

comparably in both wild type and longevity mutants but is delayed by just 1 or 2 days of

chronological age for longevity mutants (Fig. 2). This difference between two voluntary

movement assays led us to develop an inducible, motivation-independent exercise platform for

C. elegans. Our goal is to address the following shortcomings of current methods: the

movement should be inducible with a strong stimulus and not dependent on secondary cues like

food or intrinsic motivation to thrash, it should be measurable in a short time window to assess

health in this instant, and it should directly measure a physiologically relevant parameter like

maximum muscle force or functional tissue integrity. This is especially important when

comparing different genotypes, which often respond differently to their environment.

Acoustophoretic characterization of *C. elegans* force and muscle power

In humans, one of the best predictors for all-cause mortality is the decline in muscle maximum

force and power [25,28,29]. However, a tool or device to quantify the maximum force and power

of *C. elegans* muscles did not exist. The application potential would be immense since *C.*

elegans muscle structures are strongly conserved, as in mammals, and forced maximum

strength measurements to the point of collapse would be unethical in mammalian models. We

developed a microfluidic device harnessing the power of acoustic standing waves (Fig. 3a, 3b,

Supplementary Video 2, 3, Supplementary Methods). We have recently applied ultrasonic

waves to compress, move and quantitatively characterize larval *C. elegans* [30]. We reasoned that

we could employ ultrasonic standing waves to trap and stretch out *C. elegans* in the minima of

the acoustic force fields (Fig. 3c, 3d). *C. elegans* dislike being trapped and try to escape by

applying mechanical forces (body bending) against the acoustic force field (Fig. 3c, 3d). The

further away from the acoustic force field minimum, the harder it gets to move against the force

field (Fig. 3d). If the animal is stronger than the applied acoustic force field, then it can turn

around in the microfluidic chamber (Fig. 3c), typical escaping behavior of *C. elegans* known

as omega reversals [31]. Thus, the degree of deflection of the *C. elegans* body away from the

acoustic force field minimum provides an estimate of the maximal muscle strength the animal

can master to try to escape the acoustic trap.

Muscular strength declines in aging *C. elegans*

To quantify muscular forces, we developed a model by dividing the *C. elegans* body plan into

13 rigid links connected by joints along the animal's midline (Fig. 3e). Upon applying acoustic

force fields, we measured the deflection of these 13 nodes for 30 seconds and the number of

times the animal escaped the force fields (Fig. 3f). A typical exercise round is structured in up

to ten cycles consisting of 30 seconds of ultrasonic force and a 5-second break (Fig. 3g). We

measured the muscular forces of aging wild-type *C. elegans* (Fig. 3g). After 3-5 cycles, we

observed muscle fatigue, which set in earlier the older the animals were (Fig. 3g). We observed

first an increase and then a decrease in the heterogeneity of individual *C. elegans* muscular

strengths (Fig. 3g). Similarly, we first saw an increase and then a progressive decline in muscle

power during aging (Fig. 3g). By contrast, we found that longevity mutants performed better in

terms of muscle strength and function at day 20 of adulthood (Fig. 3h). This indicates the

preservation of muscle power in aging longevity mutants.

Longevity mutants showed prolonged healthspan assessed by the strength performance

in longitudinal comparison to wild type

The overall force and power of a *C. elegans* depend on its muscle strength as well as its total

body size. In agreement with previous reports [32,33], we observed adult *C. elegans* kept growing

in body size beyond the reproductive period and then shrunk during aging (Fig. 4a, 4b). A

prolonged growing phase correlates with longevity [32]. We found that longevity mutants

prolonged their growing phase and shrunk less than wild-type animals during aging (Fig. 4a,

4b). Structural integrity declines during *C. elegans* aging, such as internal organ atrophy [34], loss

of internal pressure (Supplementary Method) [35], and disorganization of the exoskeleton cuticle

occurs [36]. We noticed that in the acoustic force field, *C. elegans* undergoes compression, and

this compressibility stays fairly constant during aging (Fig. 4d, 4e). We conclude that although

morphological changes occur during aging, the mechanical properties in regards to compression

are less affected by age. This points towards muscular strength playing a key role. On average,

young *C. elegans* can overcome the acoustic force field leading to an omega turn eight times

each cycle (Fig. 4e, Supplementary Video 4) . The ability to overcome the force field and turn

in the microfluidic chip progressively declines during aging but is preserved in longevity

mutants (Fig. 4e, Extended Data Fig. 3). Turning in the chip can be viewed as a measure of

high-intensity muscular capacity since the animal can completely overcome the force field. It

also showcases that the animal is not placid but trying to escape. Next, we assessed the overall

energy per individual *C. elegans* as an assessment of overall body volume deflected against the

force field. We found an increase of energy per individual until mid-age and then a decline (Fig.

4f) reminiscent of the longitudinal body size curve (Fig. 4a). In our measurements to determine

the overall force and power of *C. elegans*, the body size is a confounding factor. In all our

longitudinal measurements, we had included a positive control in the form of a muscle-

defective mutant (CB190) that carries a mutation in the muscle myosin class II heavy chain

(*unc-54*). These muscle-constriction-defective mutants were unable to perform omega turns in

the chip but showed similar compressibility and longitudinal growth curves, illustrating that the

rise and fall of the overall energy are confounded by the organismal growth curve. Therefore,

we decided to use the dynamic power as defined as energy expenditure relative to the previous

time point and normalized it by the volume of each animal (*i.e.,* volume; Fig. 4g). Using a

human analogy, total refers to how long a weight can be lifted, dynamic power only considers

the process of lifting the weight without holding the weight, and dynamic power takes the

weight of the person lifting the weight into account. In this way, we found that the overall force

and power of longevity mutants were preserved for almost three-quarters (55-83%) compared

to about one-third (30%) of their lifespan in wild type (Fig. 4g). Thus, muscular strength is

maintained longer in longevity mutants.

Temporal scaling of age-related pathologies in longevity mutants

Next, we asked whether other age-related pathologies or morphological changes show any

delayed onset on longevity mutants compared to wild type. We quantified 592 animals,

investigated timepoints between day 0 and day 33 (12 animals on average per strain and time

point) at the first two cycles of actuation (1183-time sequences), which comprised over 800'000

frames in total. We then manually quantified additional morphological changes such as intestine

length and diameter, pixel intensity, wrinkles in the cuticle, and pharynx diameter in a

subsampled representative subset (approx. 50'000 frames; Extended Data Fig. 4). Although not

all, many age-related phenotypes were delayed in their onset and displayed a slowed decline in

longevity mutants compared to wild type (Fig. 5, Extended Data Fig. 5). Rescaling phenotypic

trajectories of wild type by the lifespan extension observed in the long-lived strains revealed

that many closely match the trajectories observed in long-lived strains for both *daf-2* mutants

(Fig. 5C). Notably, for animals' length, diameter, volume, and intestine length, phenotypic

scaling was observed when comparing wild type to *daf-2(e1368)* and *daf-2(e1370)*. In the case

of *eat-2,* the observed lifespan extension was too limited to draw conclusions and *glp-1* never

ceased growing. The severely paralyzed myosin mutant *unc-54(e190)* displayed the opposite

phenotypic trajectories than all other genotypes (Fig. 5C). When approximating the phenotypic

trajectories as segmented linear fit reflecting the separated phases of growth and decline, we

observed that the starting values as young adults are often similar (Fig. 5D). However, the

slopes and point of decline are shifted compared to wild type.

Investigating each phenotype in isolation is hindered by the inherent noise in the measurement

as well as by the incomplete picture each phenotype provides. Furthermore, many phenotypes

like length, diameter, and volume were strongly correlated. For this reason, we subjected all

phenotypes to Principle Component Analysis (PCA) to study the overall age trajectory

(Extended Data Fig. 6A). We traced these phenotypes of all genotypes across the PCA plot as

they age (Extended Data Fig. 6B). All physiological parameters increased from young to

middle-aged and then revert again as the animal reached old age. However, muscle strength

density decreased steadily. Using the paralyzed mutant, we were able to establish the bottom

left quadrant as a reduced health area. This was only possible when using both physiological

and performance measurements and was entered only by the paralyzed strain as well as old

wild-type animals. The multi-phenotype traces are also shown for each genotype individually

(Extended Data Fig. 6C, 6D). Taken together, this suggested that many of these phenotypes

change similarly during aging, that many are temporally scaled in longevity interventions, and

that maximum muscle strength offers an orthogonal perspective on studying aging compared

to physiological features. This highlights the importance of performing high-intensity muscle

strength measurements when studying physiological aging and quantifying healthspan.

317   Longevity mutants show prolonged absolute but not relative healthspan

Our data revealed that longevity mutants stay healthier compared to wild type and experience

a slower decline in physiological integrity. Indeed, dividing the lifespan of each genotype into

three chronological fixed age categories: young (less than 7 days), middle (older than 8 but

younger than 19 days), and old age (>20 days of adulthood), showed a progressing decline of

volume-corrected work performed (Fig. 6a, 6b). Longevity mutants performed better in the

middle age group than wild type, but only *daf-2(e1368)* outperformed wild type in the old age

group (Fig. 6a, 6b). Using hierarchical clustering of temporally scaled phenotypes as a

complementary analysis, we found that longevity mutants often cluster with chronologically

younger wild-type samples (Extended Data Fig. 7).

328 Integration of voluntary movement and forced maximum muscle strength quantification

329 to yield a comprehensive understanding of *C. elegans* healthspan

Having two independent assessments of healthspan that act on very different stringency levels,

we can further divide *C. elegans* healthspan into 3 divisions: prime health (passes both

matrices), fragile health (passes 1 metric), and sickspan (failing both matrices) (Fig. 6c).

Consistent with previous observations in other species, the maximum power drops prior to the

cessation in general mobility [25,28,29]. Muscle performance is much more improved in longevity

mutants in relation to wild type compared to voluntary movement (Fig. 6c). Integrating both

matrices revealed that wild type spent around 16% of their lifespan in prime health, whereas

longevity mutants spent double the time in prime health (28-42%; Fig. 6c). From all four

longevity genotypes, *daf-2(e1368)* appears to be the healthiest strain (Fig. 6c). Thus, combining

multiple matrices of physiological and behavioral integrity is a powerful assessment of

healthspan.

Discussion

Understanding the relationship between healthspan and lifespan is an important question in aging research since geroscience aims to increase the time spent in good health and to postpone and compress the time suffering from age-related pathologies and chronic diseases [2,4,5,37]. Model organisms like *C. elegans* are used to identify longevity-promoting interventions that can then be of translational value for humans [13,15]. There is a fierce debate whether *C. elegans* longevity interventions show compression of sickspan and are of translational value for improving healthspan or healthy aging in humans [16–20]. In this study, we set out to develop a robust method to quantify maximum muscle strength as a highly interpretable healthspan metric with translational value. Lifespan measurements are well-established and allowed the study of hundreds of lifespan-extending compounds and genetic alterations, leading to ground-breaking discoveries. However, this development is not reflected in the area of healthspan extension. Numerous methods exist which often measure proxy phenotypes for healthspan that are also motivation dependent like pharyngeal pumping, thrashing, and others. With our approach, we were able to directly quantify muscle health in *C. elegans*. This is especially relevant since *C. elegans* is an ideal model system for large-scale genetic screening and both the microfluidic device as well as the image detection can be multiplexed. This approach could translate to the much-needed identification of (muscle) health-promoting interventions.

The microfluidic device operates using acoustophoresis to generate an acoustic force field to quantify the physical fitness and muscle strength of aging *C. elegans*. Using six different ways to assess healthspan in forms of voluntary movements, swimming endurance, stress resilience, muscular force, muscular fatigue, structural integrity/compressibility, and quantifying several age-related morphological changes, including cuticle/skin wrinkles, body, and internal organ sizes, we find that most of these phenotypic changes are postponed in longevity mutants. We observed a hierarchical and time-dependent succession in the occurrence of these phenotypes,

starting with a loss of stress resilience, then a loss of swimming endurance, decline in maximal

force as indicated by overcoming acoustic field (omega turns), decline in dynamic power,

seizing of body and organ growth (intestine), and then decline in voluntary movement and

becoming inactive and lethargic. Using principle component analysis, we show that many of

these phenotypes are strongly cross-correlated. The delay of all these age-related phenotypic

changes is evident when using chronological age as a reference point for comparison of

longevity mutants with wild type but disappears when using relative age as a reference. This

points to the idea of temporal scaling of the healthspan. Consistent with this idea, we find that

sickspan is neither compressed nor prolonged in longevity mutants compared to wild type.

Thus, our quantifications suggest that *C. elegans* healthspan undergoes temporal scaling in

longevity.

Aging is defined as a set of phenotypes or senescent pathologies occurring with a higher

proportion in older individuals [38]. Which senescent pathologies limit the lifespan depends on

the context and are different for different species, genotypes, and environments [38,39]. Whether

our chosen set of phenotypes assessed are directly limiting or affecting lifespan is unclear.

However, it is evident that not one single mechanism underlies all our measured age-related

phenotypes. On the other extreme, we do not observe a "one mechanism causing one age-

related pathology" mechanism. Our data shows that some phenotypes correlate and also follow

a hierarchal time-dependent order of occurrence, indicating that these senescent pathologies are

interconnected. This favors a mixed model of several causal mechanisms affecting multiple

connected and independent senescent pathologies/phenotypes, including lifespan limiting

phenotypes [38,39]. Even if we might not measure lifespan limiting phenotypes directly, the

interconnectedness of phenotypes should reveal the same picture of temporal scaling of age-

related pathologies.

The lifespan of *C. elegans* can be increased by up to ten-fold [40] and decreased by 40-

fold, but surprisingly the lifespan curves often follow the same rescaled distribution [41].

Temporal scaling was also noted when comparing expression profiles of longevity mutants

compared to wild-type *C. elegans* [42]. Aging is thought to be caused by the accumulation of

damage [14,43]. Organisms either remove, detoxify, compartmentalize, or dilute molecular

damage [43]. In dividing stem cells, yeast, or bacteria, damaged macromolecules or organelles

are kept in the "mother" cell leading to accumulation of damage [14,44,45]. Since *C. elegans*

somatic cells are post-mitotic, the main driver of aging dictating the lifespan of *C. elegans*

might be the accumulation of damage in non-dividing cells. In a perfect life-conditioned

environment such as the laboratory setting, free from ever-changing environmental fluctuations

or pathogens, the only lifespan limiting factor probably boils down to the imperfectness of

biological systems [43], leading to accumulation of damage, stochastic decay of homeostatic

mechanisms [14], and consequently to age-related pathologies and phenotypes. In light of this, it

might be less surprising that the onset of pathologies is strongly linked to lifespan and thereby,

healthspan is rather temporally scaled than prolonged in longevity.

Whether temporal scaling also occurs in mammalian longevity needs to be determined

in the future. However, certain longitudinal *C. elegans* phenotypes are comparable to human

age-related phenotypes. Analogous to the *C. elegans* volume increase to peak mid-age and then

decrease is that human BMI and waist circumference also follows this early-to-mid-life increase

reaching a peak around 65-70 years and then declining [46]. Furthermore, grip strength

progressively declines after the age of 30-40 [46], similarly to *C. elegans* muscular strength. This

raises the question of whether non-compression of sickspan observed in *C. elegans* means or

interpolates to non-compression of sickspan in humans? Since aging is universal, it is tempting

to speculate that the underlying mechanisms of aging or age-dependent phenotypes are also

universal. This might be a potentially erroneous or unproven extension of the observation that

almost all living things age [38]. Although phenotypic changes, such as the loss of *C. elegans*

muscle force, is analogous to loss of grip strength or muscle strength loss in humans, the

underlying biological mechanisms resulting in physical weakness might be different. Our study

makes no conclusion or interpolation about the compression of the sickspan in humans.

There is an accumulating body of evidence that long-lived humans are healthy during

old age. For instance, 56-83% and 15-23% of centenarians, people over the age of hundred

years, delay the onset of chronic age-dependent diseases and physical disabilities or were even

free of such co-morbidities and frailties, respectively [7,10]. Centenarians have lower incidence

rates of chronic illnesses compared to their 90- or 80-year old matched-controls [8,9,11,12]. This

also extends to family members related to centenarians compared to families without

centenarians [47,48]. Thus, centenarians have a later onset and a lower rate of incidence compared

to people in their eighties, similar to our observation when comparing longevity mutants to

wild-type *C. elegans*. However, since centenarians get to enjoy at least 20 more years, how

would this comparison look if we were to compare relative age to chronological age? Is

sickspan compressed or temporally scaled in centenarians compared to the average population?

As our life expectancy doubled in the last hundred years and we are on the course of potentially

reaching the limit of our lifespan [37], the accompanied delayed onset of disabilities already

started to decelerate in longer-lived woman compared to men [49]. On the other hand, there are

several interventions that increase healthspan without increasing lifespan per se identified in

mice [50,51] and Rhesus monkeys [52]. Thus, studying longevity is an important first step of

identifying molecular mechanisms promoting healthy aging, but our study and others [50–52] point

toward that it is crucially important for geroscience to start investigating interventions that

improve healthspan directly in future studies [6]. Initial steps in defining healthspan [4,5] and also

tools and experimental setups, including this study, are being developed to reliably quantify

healthspan [53–57].

In summary, we have demonstrated that *C. elegans* sickspan is neither compressed nor extended

in longevity mutants providing an alternative answer to an ongoing debate in the aging field.

With our measurements, we showed that previous claims that insulin/IGF-1 receptor mutants

have increased sickspan compared to wild type are correct if the voluntary movement is

measured, but not the case if the muscular function or other healthspan measurements are

considered that do not rely on the behavioral state of the animal. By adjusting the reference

system from chronological age to relative age, we provide evidence that the healthspan of

longevity mutants undergoes temporal scaling. Future studies using our acoustophoresis

approach to study the role of healthspan will reveal novel strategies to improve healthy aging.

Author contributions

All authors participated in analyzing and interpreting the data and designing the experiments.

CS performed lifespan assays. CYE and CS performed swimming and oxidative stress assays.

PR designed and performed microfluidic assays. CS and PR analyzed the datasets jointly. CS

and CYE wrote the manuscript in consultation with the other authors.

Author Information

The authors have no competing interests to declare. Correspondence should be addressed to C.

Y. E. and J.D.

Acknowledgment

We thank the Ewald lab for constructive comments on the manuscript. Some strains were

provided by the CGC, which is funded by NIH Office of Research Infrastructure Programs (P40

OD010440). Funding from the Swiss National Science Foundation PP00P3_163898 to CYE

and CS.

Figure 1: Voluntary movement quantification in aging C. elegans populations. C. elegans lifespan analysis displaying survival (A), healthspan (B), and sickspan (C) for each genotype. Error bars display the standard error between plates at 24h intervals. Healthspan refers to the timespan of fast movement, sickspan to the time spend in sedentary movement, and lifespan to the time until the animal fails to move irretrievably. The fate of each individual is displayed separately for each genotype overlayed with the population's survival (D). Here, each individual's healthspan is marked as a transparent line spanning from young adulthood to the onset of sickspan marked by a black dot and then extends further as sickspan until the individual's point of death on the population survival curve. The inset displays the overall proportion each genotype spends in their healthspan for each lifespan quantile ($Q_1 – Q_4$). The correlation between health and lifespan is shown in figure (E). Each individual is represented as a point with its lifespan on the x-axis and its corresponding healthspan on the y-axis. A linear model passing through the origin is shown as a solid line. The N2 model is superimposed on the longevity mutants as a red and white dashed line. All genotypes are compared to temporally scaled N2 with the mean population life- and healthspan and error bars indicating the standard errors (F). The extrapolated ratio of health- to the lifespan of wild type

(0.78) is displayed as a dashed black line. The distance of each population average is marked by a vertical line, and the
difference in expected healthspan is indicated.

Extended Data Fig. 1: Paradoxon - young animals die apparently in good health.
The sickspan distribution for every quartile of the lifespan distribution is displayed as boxplots with each strain in a separate

panel (A). Animals experience progressively higher sickspan ratios with each quantile, with the two middle quantiles being the most similar. The distributions of the first and fourth quartile are shown as violin plots with their median line highlighted by a horizontal segment showcasing the increase of relative sickspan and heterogeneity with age across genotypes (B). To compare the individual genotypes for each relative age cohort, the corresponding quartiles are contrasted as boxplots for each strain and each quartile in a separate panel and indicate temporal scaling of healthspan (C). The sickspan ratio is compared across genotypes and quartiles (Mann-Whitney test) and the P-values are depicted as symbols (ns > 0.05, * < 0.05, ** < 0.01, *** < 0.001, **** <0.0001). When all quartiles are displayed, the median sickspan percentage of the first and last quartile of the wild type population are indicated as dashed horizontal lines. The number of observations associated with every subpopulation is displayed at the bottom of each observation. Each strain is shown in its distinct color, and increasing age is reflected by increasingly dark coloring.

Extended Data Fig. 2: Relative healthspan within early and late dying C. elegans groups.

To address the question whether outliers drive the overall health-to-lifespan ratio, each genotype was divided into seven groups
with equally sized age bins between the first and last recorded death in each genotype's population to visualize the distribution
of death events. At the top of each bar, the number of animals observed to die in this cohort and its share in the overall
population is displayed (**A**). The healthspan distribution in each age group is shown as a violin plot (**B**). The average
percentage of each population spent in their healthspan is displayed above each group, with the population size indicated in
brackets. The health-to-lifespan diagonal is shown as a dashed line, and a loess trendline was fitted to the binned datasets with
a 95% confidence interval shaded in grey. In case the sample size falls below a threshold, the violin plot was omitted. Each
analysis is displayed separately for every C. elegans genotype, which are each shown in a separate line and in a distinct color.
The spread of the healthspan distribution increases at older ages, with some animals experiencing nearly no sickspan and
others spending most of their life in the sickspan portion.

Figure 2: Progressive movement decline with age in unchallenged swimming assay and when exposed to arsenite.
In all panels, the time after subjecting the C. elegans populations to the arsenite stress is displayed on the x-axis and the movement activity of animals on the y-axis. The animal age at the start of the experiment is shown at the right of every panel. The mean activity in 14 mM arsenite is shown as a continuous line and in M9 as dashed lines. M9 was selected to avoid issues related to osmolarity. Vertical lines reflect the standard deviation at every measurement point. Two daf-2 alleles and N2 populations were maintained at 15°C and subjected to movement quantification at three aging time points (a). The daf-2(e1370) allele was also assayed in a spe-9 background and therefore maintained at 15°C until L3 and then shifted to 25°C until the start of the assay (b). The glp-1(e2141) strain was compared against a spe-9 population and maintained at 25°C after population lysis until the time of measurement.

A camera
glass
silicon
transducer
microfludic channel

~~~ *C. elegans*
— standing wave

B transducer
fluid connectors — temperature sensor
MR — microfluidic channel

C (1) ultrasound off (2) ultrasound on (3) dorsal muscle contraction (4) turnaround
214 µm

D U_{ac} U_{Ce}
○ spheres for force estimation
→ acoustic force
⊙ *C. elegans* cross section dorsal/ventral muscles

E 1 2 3 4 5 6 7 8 9 10 11 12 13

F (1) (2)(3) (4)
U_{Ce} [J] 1e−10
Ultrasound off
C. elegans turn
W_{Ce} [J] 1e−11
2.5
0.0
−2.5
0 10 20 30
Time [s]

G Progressive exhaustion with exercise duration
0 4 7 14 21 25 28 33
$<P>^{Ce}_{tot}$ [pW]
6e−12
4e−12
2e−12
0e+00
1234567891 0
exercise round

H Progressive exhaustion with exercise duration
daf-2 (e1368) *daf-2 (e1370)* *eat-2* *glp-1* *N2*
$<P>^{Ce}_{tot}$ [pW]
6e−12
4e−12
2e−12
0e+00
old adult
1234567891 0
exercise round

526
527 *Figure 3: Experimental setup to measure maximum C. elegans muscle strength:*
528 *Top view of the schematic representation of the silicon/glass acoustofluidic chip, fluid in- and outlets, and the piezoelectric*
529 *transducer positioned at the bottom and C. elegans trapped in the standing wave (A). A photograph of the back of the chip*
530 *together with the metal clamps, the attached fluid connectors, temperature probe, and a piezoelectric transducer is shown (B,*
531 *top). The front view of the chip provides an overview of the device shape, the microfluidic channel, and the measurement*
532 *region (MR) (B, bottom). A day 4 wild type C. elegans, as seen by the camera in the measurement region, is displayed*
533 *together with the image processing output highlighting the outline and the segmented midline of the animal as well as the*
534 *channel borders and centerline (C). The effect of the acoustic field (frequency: 3.543 MHz, voltage amplitude: 76 Vpp, λ/2*
535 *mode) is shown in (C, 2), aligning the animal at the midline. The animal exercises maximum muscle power attempting to*
536 *bend its head away from the midline (C, 3) to achieve a turn (C, 4), likely as an attempted escape response. The working*
537 *model to quantify the C. elegans muscular force consists of 13 rigid links along the animal's midline, which are connected by*
538 *joints. Blue arrows reflect the muscle activity acting on each joint to generate a force against the acoustic field (E). The*
539 *acoustic force acting along the animal's body is modeled as the individual acoustic forces acting on the grey spheres aligned*
540 *along the body and represented by black arrows, with the length of each arrow being proportional to its force or moment*
541 *magnitude. The acoustic radiation potential is illustrated using a C. elegans cross-section highlighting the animal's four*
542 *body wall muscles (D). If stretched, the animal rests at the minimum (see C 2) and, upon muscle contraction, moves upwards*
543 *in its potential energy well (see C3). The four C. elegans frames (C 1-4) are put in the context of one 30 seconds actuation*
544 *cycle showing the time-resolved total energy and mechanical work quantification for this animal (F). Red areas reflect*
545 *regions of zero energy due to the acoustic field being turned off, and the blue areas indicate C. elegans turn movements*
546 *during which muscle force estimation is paused. A C. elegans maximum force assessment routine consists of multiple 30*
547 *seconds exercise rounds intermitted by 5-second breaks for a total of up to ten actuation cycles. For wild type, this exercise*
548 *regimen is displayed with the exercise rounds on the x-axis, the total power (time-averaged) on the y axis, as derived from*
549 *the respective total energy curve in (F), and the age of the population given at the top of each facet (G). Wild type C. elegans*
550 *is contrasted to long-lived C. elegans genotypes across up to ten exercise rounds focusing on the aged cohort above day 20*
551 *of adulthood (H).*

N2 vs. daf−2(e1368)

Figure 4: Time course of fitness- and structural-related phenotypes for wild type and long-lived C. elegans under acoustic stimulation.

Acoustic compression was used to quantify phenotypes directly (# turns, energy, diameter- and volume compression, and dynamic power by volume) as well as indirectly by exploiting the non-destructive linear alignment of the animals (length, volume). Randomly selected images for N2 (A) and daf-2(e1368) (B) at different ages are displayed to illustrate the positioning of the C. elegans in the microfluidic measurement channel and how all matrices are shown below were obtained. Measurements are displayed as mean ± standard error for each assessed age point and subjected to local polynomial regression fitting displayed as a full line with the confidence interval set to 95% and bounded by dashed lines. For the acoustic compression only, the fitted line is shown. The length (C) and volume (D) of the animal was quantified automatically

using the entire length and area of the animal in the channel, respectively. The number of turns was quantified manually and corresponded to the number of times an animal successfully changed its orientation in the channel by 180°. The total energy of the individual was calculated using the magnitude of the lateral deflection of the animal from the channel middle (F). The compression experienced by the animal diameter (G) and volume (H) when the field is activated was computed automatically and displayed as a full line when the field is on and a dashed line when the field is off. The dynamic power/volume (I) reflects the work the animal performs against the field to change its lateral position in the channel and is normalized by the overall volume of the animal to enable comparisons across genotypes making this the most informative health parameter. The mean value for day 0 and day 33 N2 animals are indicated in black dashed lines, and the half-activity value between these two extremes is shown as a full line which is also used to deduce the health- to sickspan transition for each genotype. Using this N2 half-activity value, the individual strains reach their sickspan at approximately 10 days for N2, 28 days for daf-2(e1368), 20 days for daf-2(e1370), 18 days for eat-2, and 27 days for glp-1 while CB190 spends its entire lifespan in its sickspan fraction. Long-lived genotypes, in general, do not experience the same linear energy density decline as wild type since their total energy decrease is slowed down, as is their growth leading to a non-linear energy density trajectory.

Extended Data Fig. 3. C. elegans movement in the acoustic exercise chamber.
*The exercise regimen was benchmarked for wild type (N2) C. elegans at different ages to quantify after how many cycles they
experience muscle fatigue (A, B). Every trace corresponds to one individual animal that was up to ten times subjected to 30
seconds maximum force exercise followed by a 5 seconds break (A). Thicker lines represent the population means at every
time point. To quantify maximum muscle strength, we selected the first two actuation cycles (B) at which no fatigue can yet
be identified. The total energy is computed over the entire body and muscle apparatus of the animal. Interestingly, the spline
position most activated in this exercise is the head and tail sections (C). Focusing on the head we quantified the relative head
movement for different strains at old age and also observed the same trend of decreasing movement with higher actuation
cycles (D). The extreme case of the paralyzed CB190 vs. N2 in young adulthood displays that while the mutant is still able to
move the head away from the midline of the field when the acoustic field is off, it is unable to move when the field is on while
N2 successfully fights against the applied forces (E).*

Figure 5.Temporal scaling of C. elegans aging phenotypes.
The measured values of selected phenotypes are shown using different scales on the y-axis, animal age on the x-axis, and facetted by strain (A). To address the temporal scaling of phenotypes hypothesis, the loess fit of the measured values for wild type is shown as a solid grey line, and it is temporally scaled values using the respective mean lifespan increase experienced by the respective strain are shown as a dashed line also in grey. The measured phenotype trajectory for each strain is shown in color, each in a separate panel. The maximum fitted value is marked using. White point and its increase relative to the maximum fitted values measured for wild type are shown for both the age and phenotype variables. The same phenotypes as in subfigure (C) are modeled using a piecewise linear relationship in (B). The breakpoint of the segmented fit is estimated by the model at the age value where the linear relationship between the measured phenotype and population age changes. The individual animals measured at each time point are displayed as mean +/- standard error.

A diameter and length (acoustic on/off)

B Assumptions & simplifications

- Rod shaped intestine
- No volume variation during measurement
- No density and compressibility changes
- «Sphere» assumption for acoustic force calculation
- Energy calculation includes no turns folds since these periods are inaccessible to image processing
- Batchwise measurement along the lifespan and the batches are assumed to be identical.

C Intestine length, diameter, pharynx size and wrinkles

- Wrinkles occur rarely and only at very high age (> 25 days)
- Intestine and pharynx are often difficult to detect
- Quantification ability is dependent on the rotation of the animal

D Phenotype interdependence

E Intensity heterogeneity across sections

F Increasing bending angles with age

G Shrinking and wrinkle occurence with age

H Acoustic field triggers escape response

Extended Data Fig. 4. Phenotypes quantified within the microfluidic setup:
Automated quantification of C. elegans diameter (D = diameter) evaluated at the middle segments colored in red and the
length (L = length) of the animal along its spine colored in light blue (A), The change in diameter and length quantification
as the field is switched (US = ultrasound) is indicated. The assumptions and necessary simplifications that were made to
analyze the acquired frames are listed (B). Excerpts of the manual quantification of C. elegans intestine length and diameter,
pharynx width and height, and the number of cuticle wrinkles are displayed (C). Annotation was performed in a self-
developed image annotator suite based on the python programming language. Principal component analysis (PCA) of the

measured phenotypes are displayed for aging wild type (N2) C. elegans individuals, with each point representing one individual. The overlaying components refer to the highly correlated features length, volume, and diameter. Tissue heterogeneity and age pigments were studied by analyzing the intensity distribution across different cross-sections along the animal's fitted spine. Representative images of tissue and cuticle weakening displaying an increased bending angle of older animals compared to younger individuals as they attempt to turn in the channel (F). Similarly, older animals are smaller and sometimes display cuticle wrinkles (G). The strong stimulation of the acoustic force in the animals is shown for a representative daf-2(e1368) animal (H). The spline midline and head position is displayed for seconds preceding (left), and after (right), the acoustic field is activated. The animal directly responds with a strong escape response.

Scaled & unscaled phenotypes

A Longevity−scaled phenotypes

B Longevity−scaled phenotype trajectories

C Unscaled phenotypes

D Unscaled phenotype trajectories

627
628 ***Extended Data Fig. 5: Rationale behind the partitioning of physical properties into scaled and unscaled phenotypes.***
629 *Phenotypes displaying an age-dependent change and at least a partial temporal shift for any long-lived strain were manually*
630 *classified as temporally scaled phenotypes (A, B), and phenotypes that do not satisfy these conditions are categorized as*
631 *unscaled phenotypes (C, D). The quantified phenotypes are fitted using a loess model in all panels. The observed values for*
632 *each individual are displayed using mean +/- standard error for each timepoint (A, C). The maximum value predicted by the*
633 *loess fit is shown, and the increase relative to the wild type maximum is displayed at the top of the graph.*

A Variables – PCA

B PCA of *C. elegans* genotypes and age groups

C PCA of *C. elegans* genotypes and age groups

D PCA of wild type (N2) *C. elegans* age groups

635
636
637 ***Extended Data Fig. 6: Principal component analysis of temporally scaled phenotypes to identify similarity between***
638 ***phenotypes and their contributions.***
639 *The subset of temporally scaled phenotypes for each individual was subjected to principal component analysis, and the first*
640 *two principle components are shown together with the variance they explain. The contribution of each phenotype to the two*
641 *first principle components is displayed as vectors with the orientation reflecting the directionality and the color and length*
642 *capturing the contribution of each phenotype (A). The overall progression of young to middle-aged to old individuals*
643 *through the phenotype landscape is schematically illustrated. The bottom left quadrant is highlighted in red to indicate its*
644 *association with animals experience a poor health status. To compare the effect of age on the clustering of the different C.*
645 *elegans genotypes all animals were grouped into 3 age categories, and the individual genotypes were shown in different*

646   *colors (B). Confidence ellipses are drawn at the level of 95% for all samples (transparent) and the sample means (non-*
647   *transparent). To compare the effect of age separately for each genotype, the three age categories are shown for each strain,*
648   *young in green, middle-aged in yellow and old animals in red (C). Complete separation of age and genotype is provided in*
649   *panel (D), highlighting the measurements for every individual animal encompassed by the 95% sample confidence ellipse.*

651
652

Extended Data Fig.7: Hierarchical clustering of temporally scaled phenotypes.
The similarity between sampled genotypes and ages are displayed using hierarchical clustering. The comparison between all samples (A), between each long-lived genotype and wild type (B, D, F, H), and within each long-lived is shown (C, E, G, I). The tree is cut in five clusters when comparing all samples and in all other cases in four clusters and colored from left to right.

664
665 **Figure 6 Integration of voluntary movement and forced maximum muscle strength quantification to yield a**
666 **comprehensive understanding of C. elegans healthspan.**
667 *The volume-corrected relative work performed by each genotype and grouped by age category (young <= 7 days, 8 < middle*
668 *< 20 days, old >= 20 days) is shown as boxplots facetted by genotype (A) and by age category (B). P-values of selected*
669 *comparisons (Mann-Whitney test) are displayed as symbols (ns > 0.05, * < 0.05, ** < 0.01, *** < 0.001, **** <0.0001).*
670 *The Population medians for young and old N2 are displayed as horizontal dashed lines across all panels.*
671 *C) The measured lifespan of each C. elegans genotype is displayed with animal age on the x-axis, the fraction of the*
672 *population that is alive on the y-axis, and the mean and standard error between plates shown as point and line range. The*
673 *voluntary movement was quantified using the active vs sedentary behavior of the unstimulated animals on plates and is*
674 *shown for each population as a bar at the top of each panel. Muscle health and the corresponding onset of sickspan due to*
675 *reduced muscle function is depicted as a bar at the bottom of each facet. The position on the lifespan curve corresponding to*
676 *the health-to-sickspan transition of either the unstimulated or stimulated healthspan quantification is marked by a white*
677 *circle. The two health assessments divide the lifespan curve into three segments with decreasing health status reflected by*
678 *increasing transparency.*

Materials and Methods

Strains

Caenorhabditis elegans strains were maintained on NGM plates and OP50 *Escherichia coli* bacteria. The wild-type strain was N2 Bristol. Mutant strains used are described at www.wormbase.org: LGII: *eat-2(ad1116)*; LGIII: *daf-2(e1368, e1370), glp-1(e2141)*.

C. elegans culturing conditions

C. elegans populations were age-synchronized by isolating eggs from gravid C. elegans adults, incubating them for 18 hours in M9 until hatching in the presence of 5 μg mL−1 cholesterol (Sigma-Aldrich). Hatched L1-larvae were grown on standard culturing NGM OP50 plates and then shifted to 50 mM FUDR plates seeded with heat-killed OP50 plates when reaching the L4 state. Animals were maintained on FUDR plates until the measurement of different ages was taken. The *glp-1* genotype was placed at 25°C after bleach treatment and shifted back to 15°C at the L4 stage and otherwise treated equally to the other strains. Except for *glp-1* during its development, all animals were always maintained and aged at 15°C.

Automated survival assays using the lifespan machine

To compare the lifespans among wild type and long-lived mutants, we raised all animals for several generations in parallel. Automated survival analysis was conducted using the lifespan machine setup described by [26]. Briefly, approximately 1000 L4 animals were resuspended in M9 and transferred to NGM plates containing 50 μM 5-Fluoro-2'deoxyuridine (FUdR) seeded with heat-killed OP50 bacteria and incubated at 15°C until day 4 of adulthood. Animals were then resuspended in M9 and transferred to fresh FUDR plates containing tight-fitting lids (BD Falcon Petri Dishes, 50x9mm), and the plates were dried with their lids open for 30 minutes after transfer. The plates were incubated for five additional days to rule out contamination and then loaded in the lifespan machine. Air-cooled Epson V800 scanners were utilized for all

719 experiments operating at a scanning frequency of one scan of 30 minutes. Temperature probes

720 (Thermoworks, Utah, US) were used to monitor the temperature on the scanner flatbed and kept

721 constant at 15°C.

723 **Voluntary movement healthspan measured by lifespan machine**

724 The time point at which the animal stops moving completely and irretrievably is classified as

725 the point of death and defines the lifespan of each individual. The health- to sickspan transition

726 is estimated by the time point when major movement ceases and exclusively head movements,

727 posture change, and minor body movements can be observed. The animal is also required to be

728 sedentary and remains in the rough vicinity of the area; it will ultimately die.

730 **Microfluidics device measuring muscle strength**

731 The detailed description of the microfluidics device development and characterization is found

732 in the Supplementary Methods file.

734 **Oxidative stress assay arsenite with automated wormtracker**

735 Voluntary movement in liquid, as well as resistance to oxidative stress, was quantified using

736 the wMicroTracker (NemaMetrix) platform. Briefly, animals were resuspended in M9 at

737 indicated time points and washed three times with M9 by centrifugation. 30 – 40 *C. elegans*

738 were pipetted into each well of a round-bottomed 96- well plate in either M9 or 14 mM

739 sodium arsenite solution and assayed for 50 hours. Each genotype was always measured both

740 in M9 as well as arsenite in parallel to enable comparisons. Activity measurements were

741 analyzed using the R programming language, and average activity, as well as the standard

742 deviation of wells, are shown.

744 **Figure generation and statistics**

745   The analysis was performed using the statistical software R. data processing and visualization

746   was performed using the tidyverse package collection, most prominently dplyr and ggplot2.

747   Furthermore, packages were used for lifespan analysis (survival, survminer), computing and

748   visualizing PCA (stats and ggfortify, factoextra), fitting loess models (stats), and segmented fits

749   (segmented), labeling (ggrepel) comparing distributions (ggpubr) and arranging figures

750   (cowplot).

References

1. Nations, D. of E. and S. A. P. D. U. World Population Ageing 2019: Highlights. 1–46 (2019).

2. Partridge, L., Deelen, J. & Slagboom, P. E. Facing up to the global challenges of ageing. *Nature* **561**, 45–56 (2018).

3. Goldman, D. P. *et al.* Substantial health and economic returns from delayed aging may warrant a new focus for medical research. *Health affairs (Project Hope)* **32**, 1698–1705 (2013).

4. Kaeberlein, M. Translational geroscience: A new paradigm for 21st century medicine. *Translational Medicine of Aging* **1**, 1–4 (2017).

5. Kennedy, B. K. *et al.* Geroscience: linking aging to chronic disease. *Cell* **159**, 709–713 (2014).

6. Olshansky, S. J. From Lifespan to Healthspan. *JAMA : the journal of the American Medical Association* **320**, 1323–1324 (2018).

7. Ailshire, J. A., Beltrán-Sánchez, H. & Crimmins, E. M. Becoming centenarians: disease and functioning trajectories of older US Adults as they survive to 100. *The journals of gerontology Series A, Biological sciences and medical sciences* **70**, 193–201 (2015).

8. Andersen, S. L., Sebastiani, P., Dworkis, D. A., Feldman, L. & Perls, T. T. Health span approximates life span among many supercentenarians: compression of morbidity at the approximate limit of life span. *The journals of gerontology Series A, Biological sciences and medical sciences* **67**, 395–405 (2012).

9. Evans, C. J. *et al.* Place and cause of death in centenarians: a population-based observational study in England, 2001 to 2010. *PLoS medicine* **11**, e1001653 (2014).

10. Evert, J., Lawler, E., Bogan, H. & Perls, T. Morbidity profiles of centenarians: survivors, delayers, and escapers. *The Journals of Gerontology: Series A* **58**, 232–237 (2003).

11. Ismail, K. *et al.* Compression of Morbidity Is Observed Across Cohorts with Exceptional Longevity. *Journal of the American Geriatrics Society* **64**, 1583–1591 (2016).

12. Kheirbek, R. E. *et al.* Characteristics and Incidence of Chronic Illness in Community-Dwelling Predominantly Male U.S. Veteran Centenarians. *Journal of the American Geriatrics Society* **65**, 2100–2106 (2017).

13. Kenyon, C. J. The genetics of ageing. *Nature* **464**, 504–512 (2010).

14. López-Otín, C., Blasco, M. A., Partridge, L., Serrano, M. & Kroemer, G. The hallmarks of aging. *Cell* **153**, 1194–1217 (2013).

788   15. Magalhães, J. P. de, Stevens, M. & Thornton, D. The Business of Anti-Aging Science.
789   *Trends in Biotechnology* **35**, 1062–1073 (2017).

790   16. Bansal, A., Zhu, L. J., Yen, K. & Tissenbaum, H. A. Uncoupling lifespan and healthspan
791   in Caenorhabditis elegans longevity mutants. *Proceedings of the National Academy of*
792   *Sciences* **112**, E277-86 (2015).

793   17. Hahm, J.-H. *et al.* C. elegans maximum velocity correlates with healthspan and is
794   maintained in worms with an insulin receptor mutation. *Nature communications* **6**, 8919
795   (2015).

796   18. Huang, C., Xiong, C. & Kornfeld, K. Measurements of age-related changes of
797   physiological processes that predict lifespan of Caenorhabditis elegans. *Proceedings of the*
798   *National Academy of Sciences of the United States of America* **101**, 8084–8089 (2004).

799   19. Stamper, B. L. N. *et al.* Movement decline across lifespan of Caenorhabditis elegans
800   mutants in the insulin/insulin-like signaling pathway. *Aging cell* **17**, e12704 (2018).

801   20. Podshivalova, K. & Kerr, R. A. How a Mutation that Slows Aging Can Also
802   Disproportionately Extend End-of-Life Decrepitude. *Cell reports* **19**, 441–450 (2017).

803   21. Ewald, C. Y., Landis, J. N., Abate, J. P., Murphy, C. T. & Blackwell, T. K. Dauer-
804   independent insulin/IGF-1-signalling implicates collagen remodelling in longevity. *Nature*
805   **519**, 97–101 (2015).

806   22. Ewald, C. Y., Castillo-Quan, J. I. & Blackwell, T. K. Untangling Longevity, Dauer, and
807   Healthspan in Caenorhabditis elegans Insulin/IGF-1-Signalling. *Gerontology* **64**, 96–104
808   (2018).

809   23. Gems, D. *et al.* Two pleiotropic classes of daf-2 mutation affect larval arrest, adult
810   behavior, reproduction and longevity in Caenorhabditis elegans. *Genetics* **150**, 129–155
811   (1998).

812   24. Hess, M., Gomariz, A., Goksel, O. & Ewald, C. Y. In-vivo quantitative image analysis of
813   age-related morphological changes of C. elegans neurons reveals a correlation between
814   neurite bending and novel neurite outgrowths. *Eneuro* **6**, ENEURO.0014-19.2019 (2019).

815   25. Leong, D. P. *et al.* Prognostic value of grip strength: findings from the Prospective Urban
816   Rural Epidemiology (PURE) study. *Lancet* **386**, 266–273 (2015).

817   26. Stroustrup, N. *et al.* The Caenorhabditis elegans Lifespan Machine. *Nature Methods* **10**,
818   665–670 (2013).

819   27. Zhang, W. B. *et al.* Extended Twilight among Isogenic C. elegans Causes a
820   Disproportionate Scaling between Lifespan and Health. *Cell Systems* **3**, 333-345.e4 (2016).

821   28. Petrella, J. K., Kim, J., Tuggle, S. C., Hall, S. R. & Bamman, M. M. Age differences in
822   knee extension power, contractile velocity, and fatigability. *J Appl Physiol* **98**, 211–220
823   (2005).

824   29. Kostka, T. Quadriceps maximal power and optimal shortening velocity in 335 men aged
825   23–88 years. *Eur J Appl Physiol* **95**, 140–145 (2005).

826   30. Baasch, T. *et al.* Acoustic Compressibility of Caenorhabditis elegans. *Biophysical journal*
827   **115**, 1817–1825 (2018).

828   31. Donnelly, J. L. *et al.* Monoaminergic Orchestration of Motor Programs in a Complex C.
829   elegans Behavior. *Plos Biol* **11**, e1001529 (2013).

830   32. Hulme, S. E. *et al.* Lifespan-on-a-chip: microfluidic chambers for performing lifelong
831   observation of C. elegans. *Lab on a chip* **10**, 589–597 (2010).

832   33. Shi, C. & Murphy, C. T. Mating induces shrinking and death in Caenorhabditis mothers.
833   *Science (New York, NY)* **343**, 536–540 (2014).

834   34. Ezcurra, M. *et al.* C. elegans Eats Its Own Intestine to Make Yolk Leading to Multiple
835   Senescent Pathologies. *Current biology : CB* **28**, 2544-2556.e5 (2018).

836   35. Gilpin, W., Uppaluri, S. & Brangwynne, C. P. Worms under Pressure: Bulk Mechanical
837   Properties of C. elegans Are Independent of the Cuticle. *Biophysical journal* **108**, 1887–1898
838   (2015).

839   36. Essmann, C. L. *et al.* Mechanical properties measured by atomic force microscopy define
840   health biomarkers in ageing C. elegans. *Nature communications* **11**, 1043 (2020).

841   37. Olshansky, S. J. & Carnes, B. A. Inconvenient Truths About Human Longevity. *The*
842   *journals of gerontology Series A, Biological sciences and medical sciences* **74**, S7–S12
843   (2019).

844   38. Freund, A. Untangling Aging Using Dynamic, Organism-Level Phenotypic Networks.
845   *Cell Systems* **8**, 172–181 (2019).

846   39. Gems, D. The aging-disease false dichotomy: understanding senescence as pathology.
847   *Frontiers in genetics* **6**, 212 (2015).

848   40. Ayyadevara, S., Alla, R., Thaden, J. J. & Reis, R. J. S. Remarkable longevity and stress
849   resistance of nematode PI3K-null mutants. *Aging cell* **7**, 13–22 (2008).

850   41. Stroustrup, N. *et al.* The temporal scaling of Caenorhabditis elegans ageing. *Nature* **530**,
851   103–107 (2016).

852   42. Tarkhov, A. E. *et al.* A universal transcriptomic signature of age reveals the temporal
853   scaling of Caenorhabditis elegans aging trajectories. *Scientific reports* **9**, 7368 (2019).

854   43. Gladyshev, V. N. The origin of aging: imperfectness-driven non-random damage defines
855   the aging process and control of lifespan. *Trends in genetics : TIG* **29**, 506–512 (2013).

856   44. Lippuner, A. D., Julou, T. & Barral, Y. Budding yeast as a model organism to study the
857   effects of age. *FEMS microbiology reviews* **38**, 300–325 (2014).

858   45. Yang, Y. *et al.* Temporal scaling of aging as an adaptive strategy of Escherichia coli.
859   *Science advances* **5**, eaaw2069 (2019).

860   46. Kuo, P. L. *et al.* A roadmap to build a phenotypic metric of ageing: insights from the
861   Baltimore Longitudinal Study of Aging. *Journal of Internal Medicine* **287**, 373–394 (2020).

862   47. Ash, A. S. *et al.* Are Members of Long-Lived Families Healthier Than Their Equally
863   Long-Lived Peers? Evidence From the Long Life Family Study. *The journals of gerontology*
864   *Series A, Biological sciences and medical sciences* **70**, 971–976 (2015).

865   48. Sebastiani, P. *et al.* Families Enriched for Exceptional Longevity also have Increased
866   Health-Span: Findings from the Long Life Family Study. *Frontiers in public health* **1**, 38
867   (2013).

868   49. Freedman, V. A., Wolf, D. A. & Spillman, B. C. Disability-Free Life Expectancy Over 30
869   Years: A Growing Female Disadvantage in the US Population. *American journal of public*
870   *health* **106**, 1079–1085 (2016).

871   50. Fischer, K. E. *et al.* A cross-sectional study of male and female C57BL/6Nia mice
872   suggests lifespan and healthspan are not necessarily correlated. *Aging* **8**, 2370–2391 (2016).

873   51. Garcia-Valles, R. *et al.* Life-long spontaneous exercise does not prolong lifespan but
874   improves health span in mice. *Longevity & healthspan* **2**, 14 (2013).

875   52. Mattison, J. A. *et al.* Impact of caloric restriction on health and survival in rhesus
876   monkeys from the NIA study. *Nature* **489**, 318–321 (2012).

877   53. Bellantuono, I. *et al.* A toolbox for the longitudinal assessment of healthspan in aging
878   mice. *Nature Protocols* **15**, 540–574 (2020).

879   54. Collins, J. J., Huang, C., Hughes, S. & Kornfeld, K. The measurement and analysis of
880   age-related changes in Caenorhabditis elegans. *WormBook : the online review of C elegans*
881   *biology* 1–21 (2008) doi:10.1895/wormbook.1.137.1.

882   55. Teuscher, A. C., Statzer, C., Pantasis, S., Bordoli, M. R. & Ewald, C. Y. Assessing
883   Collagen Deposition During Aging in Mammalian Tissue and in Caenorhabditis elegans.
884   *Methods Mol Biology Clifton N J* **1944**, 169–188 (2019).

885   56. Haefke, A. & Ewald, C. Tail Tendon Break Time for the Assessment of Aging and
886   Longitudinal Healthspan in Mice. *Bio-protocol* **10**, (2020).

887   57. Teuscher, A. C. & Ewald, C. Y. Overcoming Autofluorescence to Assess GFP Expression
888   During Normal Physiology and Aging in Caenorhabditis elegans. *BIO-PROTOCOL* **8**, (2018).