1

Loss of Stk11 in basolateral amygdalar projection neurons impairs CTA learning by altering the temporal pattern of taste response plasticity in GC

Roshan D. Nanu[2], Thomas J. Murdy[2], David Levitan[2], Sacha B. Nelson[2,3], Donald B. Katz[1,3]

Departments of Psychology[1], Biology[2], and Volen Center for Complex Systems[3], Brandeis University, Waltham, Massachusetts 02454, USA.

Author Contributions

D.L., S.N., and D.K. initiated the project. R.N., D.L. and D.K. designed the experiments. R.N. and T.M. performed experiments. D.L. performed histology. R.N. analyzed the data. R.N., D.K. and S.N. wrote the manuscript. S.N. reviewed the manuscript.

Proofs and correspondence to:

Donald B. Katz, PhD, Brandeis University

Department of Psychology, MS 062, 415 South Street, Waltham, MA 02454,

Email: dbkatz@brandeis.edu

Declaration of Interest: The authors declare no competing financial interests.

Acknowledgements: This research was supported by US National Institutes of Health grant R21-MH104318 and the National Institute on Deafness and Other Communication Disorders (NIDCD) grants DC006666 and DC007703.

Abstract
Gustatory cortex (GC) responds to tastes on the tongue with dynamic ensemble activity that
represents first the presence, then the identity, and finally the hedonic value (palatability) of
tastes. This final state of the taste response is uniquely altered by conditioned taste aversion
(CTA) – a powerful one-trial learning paradigm in which a taste becomes aversive after
association with gastric malaise – a process requiring coordination between GC and basolateral
amygdala (BLA). Previously, we found that one key requirement for learning in this circuit is
expression of the serine/threonine kinase 11 (Stk11) gene (a tumor suppression gene which has
only recently been associated with learning; Levitan et al., 2020). When Stk11 is knocked out in
BLA projection neurons (BLApn), CTA learning fails to occur. Here we have examined how
learning-related response plasticity in GC taste responses is impacted by the knockout of Stk11
in BLApn. Contrary to the commonly held assumption that a lack of learning means a lack of
such plasticity, but consistent with the fact that Stk11KO has been shown to increase the
excitability of BLApn, our data reveal that the knockout of Stk11 in BLApn does not eliminate
plasticity; rather, it shifts the impact of CTA training on GC taste responses to an earlier,
learning-inappropriate epoch. Even naïve taste representations are altered—specifically, the
pattern of similarities and differences among the different taste responses are rendered
abnormal by Stk11 KO, and these relationships fail to change with training. Finally, the latency
of behavior-related dynamic ensemble features of the GC taste response, which is also
abnormal in naïve KO mice, is rendered disorganized by CTA. Together, these results suggest
that Stk11 plays a role in governing the coordination of GC activity by BLA, and demonstrate
that alterations in the function of BLApn caused by Stk11KO inhibit learning not by inhibiting
plasticity but by changing its temporal properties.

48
Introduction
Following placement of a taste on the tongue, gustatory cortical (GC) ensemble
responses exhibit a rich temporal structure, evolving through a sequence of distinct encoding
states that signal first the presence of a taste, then the identity of the taste, and finally the
palatability or hedonic value of the taste (Katz et al., 2001; Jones et al., 2007; Piette et al., 2012;
Sadacca et al., 2012). On individual taste presentations, the onset of the palatability state is a
sudden and coherent ensemble transition, the timing of which reliably predicts and drives
behavior (Moran & Katz, 2014; Sadacca et al., 2016; Mukherjee et al., 2019).
When presentation of a particular taste is paired with induced gastric malaise such that
an animal learns a conditioned taste aversion (CTA), response plasticity is temporally-specific:
the coding content (as measured in firing rates) of the palatability phase of that taste's
response changes to reflect depreciation of the tastant's value, while pre-training activity in
prior response phases is largely preserved (Grossman et al., 2008; Moran & Katz, 2014; Lavi et
al., 2018).
This learning-relevant palatability-related information appears to be in part provided to
GC by the basolateral amygdala (BLA) (Piette et al., 2012; Lin et al., 2021). BLA is necessary for
proper CTA learning (Gallo et al., 1992; Yamamoto et al., 1995; Grossman et al., 2008; Garcia-
Delatorre et al., 2014), as are the axons of BLA→GC projection neurons (BLApn) themselves
(Yamamoto et al., 1984; Levitan et al., 2020). Of particular interest in this regard are recent
findings demonstrating that the BLApn-specific knockout of the serine/threonine kinase 11

(Stk11) gene, commonly studied as a tumor suppressor and regulator of cell growth in tissue
throughout the body (Shaw, 2009; Nakada et al., 2010; Beirowski, 2019), prevents the
acquisition of CTA (Levitan et al., 2020).
At the cellular level, the effect of losing Stk11 is an increase in the basal excitability of
BLA→GC projection neurons (Levitan et al., 2020), a fact that has intriguing implications for the
impact of the knockout on taste-related GC neural activity over CTA training. Most generally,
these results suggest the possibility that the Stk11 KO may cause an "over-drive" of GC by BLA
input; far from inhibiting plasticity, the increase in excitability might be expected to enhance GC
response plasticity while rendering it temporally inappropriate for learned behavior.
Here, we have set out to test this unusual hypothesis, investigating how the knockout of
Stk11 in BLApn disrupts learning-related changes in GC taste responses. We recorded
extracellular single neuron responses in GC of awake, freely behaving mice as these mice
underwent CTA training, and observed how taste responses in GC changed—both those of
individual neurons and of ensembles of neurons. Our results reveal subtle alterations of taste
processing in naïve Stk11 KO mouse, and show that, in these mice, CTA training-related changes
are not nonexistent but rather learning-inappropriate.
85
86 **Results**
87
We performed a selective, inducible knock out of Stk11 in BLA projection neurons (Stk11
KO, N=6 mice) using viral transfection to express Cre in Stk11$^{fl/fl}$ mice. One week later, we
implanted bundles of electrodes bilaterally in GC and inserted an intraoral canula (IOC) under
the masseter muscle so that tastes could be delivered directly onto the tongue. Ten days after
virus injection and electrode/IOC implantation, we recorded bilaterally from ensembles of GC
single neurons throughout a combined tasting/CTA training regime: mice were presented with
a battery of tastes of known hedonic values; one day later, the mice were subjected to a 24hr
CTA protocol designed to train an aversion to saccharin, a taste that is palatable to the naïve
mouse, and then given another, post-learning taste battery (Figure 1B). We also subjected a
control cohort (Control, N=10 Stk11 $^{fl/fl}$ mice) that had undergone injection of a viral vector
coding only for GFP (Figure 1A) to the same experimental regime. Given that CTA learning
occurs across <24hrs (with only one taste-illness pairing), we were able in some cases to
monitor the responses of single neurons in GC from before until after learning.
We evaluated learning by comparing nightly water bottle consumption before CTA
training to saccharin consumption after CTA training (10 min access each time). Given the
complexity of the recording protocol (see Methods), some extinction was expected to occur
before the test of learning. Furthermore, relatively mild learning is an inevitable result of
embedding the training in a context of multiple recording sessions involving presentation of
multiple tastes (see Discussion). In recognition of these factors, we set a relatively low
threshold: animals whose saccharin consumption was <80% of their mean nightly water
consumption were deemed to have learned and retained the aversion; this criterion was
reached by 5 of the 10 control mice, but by none of the Stk11 KO mice that drank water
normally (see Methods). These results are consistent with those shown previously (Levitan et
al., 2020)—Stk11 knockout in BLApn inhibits normal CTA learning.

A total of 460 well-isolated single neurons (313 from Stk11 KO mice; 147 from control
mice that learned CTA) were subjected to initial analysis (13 ± 2.1 cells per session). The basic
properties of taste responses were largely similar for neurons from the two groups of mice, but
Figure 2 (top-left) shows that the average firing rates of putative pyramidal neurons, computed
over the 1s before stimulus presentation (baseline), were higher in Stk11 KO mice than in
control mice (95% CI of difference: [0.40, 1.0] Hz). A two-way ANOVA comparing group and
training (pre- vs post-CTA) effects on pyramidal cell firing rates revealed a significant group
effect ($F_{(1,384)}=5.1$, $p=0.024$) on baseline firing rates (no other effects were significant). No
between-group differences were observed between interneuron firing rates, and whole-
response firing rates across the 1.5s after taste delivery (response, with no examination of
"temporal coding") were similar between groups for both cell types.
123
124   *The dynamics of learning-related response plasticity is abnormal in the BLA-specific Stk11 KO*
125
In GC responses, palatability-correlated firing emerges across the 1.5s after stimulus
delivery (Levitan et al., 2019; Levitan et al., 2020). The same pattern was observed here (Figure
3A) for both groups of mice. A two-way ANOVA for group and training effects performed on
these palatability correlation data revealed significant effects of both the KO ($F_{(1,1940)}=63$,
$p=3.0e-15$) and training ($F_{(1,1940)}=6.4$, $p=.012$; Figure 3B)—Stk11 KO mice demonstrated a
consistently higher degree of palatability-related correlation than Controls (95% CI of
difference: [0.029, 0.044]), and both groups showed a mild elevation in palatability correlation
after training.
In our previous paper (Levitan et al. 2020), we failed to observe significant differences in
the correlation of GC firing to palatability between Stk11 KO and Control. This small discrepancy
between these data and our prior results is likely due to our increased sample of Stk11 KO
neurons and the fact that our current taste battery lacked sucrose, the reliable high palatability
of which may have overwhelmed subtle between-group differences (see Discussion).
Regardless, the slight, general enhancement in GC palatability-related firing makes sense: since
a portion of palatability information in GC appears to come from BLA projections (Piette et al.,
2012; Lin et al., 2021), a larger than normal contribution of BLA to GC taste responses might be
expected resultant from the increased BLA excitability in the Stk11 knockout (Levitan et al.,
2020).
The fact that CTA training changed GC responses in Stk11 KO mice (mice that failed to
learn CTAs) was more surprising, as failure to learn is typically assumed to imply a lack of
training-related response plasticity. CTA learning, however, is supported not simply by response
changes, but by changes that are specific in timing and nature—changes in late-epoch firing
that leave the conditioned taste responses more similar to those caused by innately aversive
tastes (Grossman et al., 2008; Moran & Katz, 2014; Lavi et al., 2018; Mukherjee et al., 2019;
Arieli et al., 2020). We therefore hypothesized that GC response changes wrought in Stk11 KO
mice by CTA training might differ from those observed in control mice in these temporal
specifics. To test this hypothesis, we focused on evaluating stimulus-evoked response dynamics
in neurons held across the entire CTA training regimen. Firing rates relative to baseline were
compared for the 1.5s post-stimulus in 250ms non-overlapping bins (and statistically analyzed
using one-way ANOVA with Bonferroni correction).

As has been shown previously for rats (Grossman et al., 2008; Moran & Katz, 2014),
significant learning-related changes in GC neurons recorded from control animals were largely
restricted to the late epoch of the response (>750ms; Figure 4a, left). This pattern is consistent
with the generally agreed-upon suggestion that CTA changes the palatability of a taste, leaving
taste identity intact (Bures et al., 1998; Fonseca et al., 2020), and with the fact that it is the
transition into this late state that is implicated in the driving of palatability-related behavior (Li
et al., 2016; Sadacca et al., 2016; Mukherjee et al., 2019).
These findings contrast with the results for Stk11 KO mice. Training-related changes
were in fact present in these mice, but the response plasticity was abnormally centered on
early parts of the response, with firing rates returning to pre-training levels by the late phase
(Figure 4a, right). Figure 4b shows the examples in Figure 4a to be representative: training-
related response plasticity in the control group responses tended to appear late in responses,
but clustered at earlier response times in Stk11 KO mice; these distributions differed
significantly ($\chi^2(1)=13$, p=0.026, V=0.580), confirming that training-related response plasticity in
Stk11 KO mice, while robust, was abnormal.
It was important to consider the possibility that this seemingly abnormal response
plasticity observed in Stk11 KO mice was not simply a "normal" result of the fact that these
mice failed to learn. Prior work from our lab suggests that this is not the case, in that sham-CTA,
in which no aversive element is introduced, results in almost no (<5% of cells) response
plasticity (Grossman et al., 2008). Additionally, examination of GC responses in the subset of
control mice for which learning failed to reach significance (data not shown) revealed a
distribution that looked, if anything, more like those of control mice that learned ($\chi^2(1)=2.1$,
p=0.83, V=0.36) than they did Stk11 KO mice ($\chi^2(1)=14$, p=0.018, V=0.63)—a fact that likely
reflects learning rendered subthreshold by extinction and the inclusion of a taste battery (see
Discussion). Thus, we conclude that the abnormal training-related taste response plasticity
observed in our Stk11 KO mice is not the result of a normal failure or absence of learning, but
truly reflects a change in training-related plasticity wrought by the Stk11 KO in BLApn.
183
184 *Impaired ensemble representations of saccharin with Stk11 KO*
185
The above analyses demonstrate that CTA training changes the content of GC taste
responses in both control and Stk11 KO mice, but that the nature of those changes (defined by
known differences in coding in different temporal epochs of the taste responses, see Fontanini
& Katz, 2006; Jones et al., 2007; Miller & Katz, 2010; Maffei et al., 2012; Moran & Katz, 2014;
Sadacca et al., 2016) is group-dependent; the plasticity is learning-appropriate in control mice
only. These analyses beg the question of the implications of these changes for the information
content of the responses.
To investigate this aspect of response plasticity, we examined the information available
in the subset of single neurons that were held across either both pre-CTA or both post-CTA
recording sessions so that saccharin responses could be related to those of other tastants. We
embedded the population firing rates from the early (<750ms) and late (750-1500ms) epochs of
the response into separate 2-D spaces using the manifold data reduction technique multi-
dimensional scaling (MDS). By taking the Euclidean distances between saccharin responses and
the mean quinine and NaCl responses (distances D_Q and D_N, respectively), and normalizing to

the distance between the mean quinine and NaCl responses (D_{QN}), we created a dimensionless
distance metric $\left(D_S = \left[D_Q - D_N\right]/D_{QN}\right)$ that allowed us to accurately compare how similar the
ensemble saccharin representations (even those with different numbers of neurons) were to
each. Values greater than 0 signified that the saccharin representation was more similar to NaCl
(the palatable tastant) than to quinine. Note that this analysis does not necessarily reflect the
amount of palatability information embedded in the ensemble response, but instead shows the
organization of the N-dimensional saccharin representation in relation to those of tastants
whose valence remains fixed over the CTA paradigm (see Methods).
The results of this analysis are presented in Figure 5. Consistent with what is known of
CTA learning (Moran & Katz, 2014), late epoch saccharin representations in control mice shifted
toward those of quinine as a result of CTA training (see Table 1). Saccharin responses in the
Stk11 KO group, meanwhile, showed no such change. Instead, these responses were
significantly closer to quinine than those of controls before training, which then failed to
change that coding significantly; if anything, the "early epoch" responses (which were the site
of most of the training-related changes, see Figure 4b) became more "NaCl-like" with training –
an opposite change than that expected to occur with CTA learning.
In summary, Stk11 KO seems to disrupt the relationships among taste representations in
naïve mice; furthermore, while conditioning-related response changes occur, these changes
don't impact this organization in any manner that would be expected to support learning.
219
220 *Single-trial response dynamics are altered with Stk11 KO*
221
The data presented thus far are consistent with prior work from the Katz lab regarding
the sequential processing of taste identity and palatability in GC. That prior work has also
shown that palatability-related "states" arise suddenly in GC within single trials; state-to-state
transition times vary widely between trials, but the overall sequence is highly reliable (Miller &
Katz, 2010; Moran & Katz, 2014; Sadacca et al., 2016). CTA learning impacts the timing of these
transitions (at least in rats), without altering the general dynamical structure of the responses
(Moran & Katz, 2014).
To investigate the impact of Stk11 KO on these ensemble properties of GC taste
responses before and after CTA training, we considered taste-evoked ensemble spike trains as
multivariate Poisson processes segmented into discriminable states that could be modelled as a
first-order Markov chain – that is, we applied Hidden Markov Models (HMMs) to whole
ensemble responses for each taste in each session, in order to capture the underlying state
dynamics of the evoked responses. Based on previous work (see also Methods) we fit 3-state
models connected in a feed-forward fashion; these models account for the presence of a
background state and distinct early and late response states (Figure 6a).
Results showed that Stk11 KO and control mice produced taste responses with similar
basic state dynamics—both groups' GC ensemble responses featured sudden and coherent
firing rates changes across the population (Figure 6b). In control mice, these dynamics were
largely unchanged by CTA training, a fact that represents a basic difference between mice and
rats: in rats, the transition to the late (palatability) state shifts earlier in the trial with CTA
training (Moran & Katz, 2014); in our control mice, this trend was at most subtle, and failed to
reach significance (Figure 7A left, 7B left, 7C).

More pertinent to the current hypotheses, our analysis revealed multiple differences
between control and Stk11 KO transition times. For one thing, transitions were substantially
delayed in naïve Stik11 KO mice: before training, this transition occurred >300ms later in the KO
than control mice (Figure 7A right, 7C); a two-way ANOVA comparing the effects of group and
training showed this difference between groups to be significant ($F_{(1,157)}=20$, p=1.4e-5; Figure
7B).
While the average transition time for the KO moves earlier with training (by 170ms on
average), no training ($F_{(1,157)}=2.8$, p=0.10) or interaction effects ($F_{(1,157)}=1.7$, p=0.19) were
seen. This failure to identify a significant training effect, together with the fact that training also
appeared to decrease the sharpness of that transition (Figure 7C magenta), led us to more
closely examine the entire distribution of single-trial transition times in both groups (Figure 8a).
Scrutiny of these data suggests that the post-training behavior of Stk11 KO ensembles is far
from normal; further analysis (Figure 8b) reveals that whereas the distribution of transition
times into the late (palatability) state are reasonably well fit by a gaussian (pre-CTA Control
$r^2=0.90$; post-CTA Control $r^2=0.83$; pre-CTA Stk11 KO $r^2=0.74$) in typical sets of saccharin
responses, this structure breaks down in the Stk11 KO mice after training—the distribution
loses its single strong peak ($r^2=0.040$), becoming significantly different from before training (χ
$^2(1)= 41.0$, p=1.5e-5, V=0.76). Clearly, the knockout of Stk11 in BLA projection neurons did not
render the mice incapable of response plasticity, but this response plasticity was highly
abnormal, and resulted in taste responses that were highly abnormal.
264
265 **Discussion**
266
267 *Mouse CTA resembles rat CTA*
268
The gustatory neuroaxis consists of multiple brain regions acting in concert to produce
precisely timed neural codes that in GC indicate, in turn, the identity and hedonic value of a
tastant—information that is used to generate an appropriate behavioral response. This system
supports learning whereby a tastant is associated with negative gastric consequences occurring
minutes or even hours after a single ingestion session; this learning mechanism allows animals
to avoid ingesting substances that have proven harmful, and is thereby evolutionarily adaptive.
Here we take a step toward understanding how interactions between brain regions contribute
to the formation of this robust form of learning by looking into how the knockout of a single
gene in BLA projection neurons impacts ensemble activity in GC during taste processing and
over the course of CTA training.
Prior studies using rats have shown that it is primarily the late "epoch" of the taste
response that changes over the course of CTA training, as the code continues to reflect the
(changing) value of the tastant and reliably predict behavior (Grossman et al., 2008; Sadacca et
al., 2016; Mukherjee et al., 2019). This remains true in this study – in control mice (that did
learn the CTA), changes in single unit responses to saccharin primarily occurred in the late
epoch (>750ms) with CTA training (Figure 4), such that the ensemble representations of
saccharin during this epoch shifted dramatically from resembling another palatable taste (NaCl)
towards more resembling an unpalatable taste (quinine) over learning (Figure 5). In fact, the
only rat result that we were unable to clearly replicate in control mice was a significant

decrease in the onset latency of the late state over CTA learning (Moran & Katz, 2014); a trend
toward this shift was observed (Figure 7B&C, Figure 8B), but the magnitude was smaller (in our
hands insignificantly so) than that seen in rats.
It's worth noting that in our control mice, learned aversions to saccharin were mild—a
fact that might have made it difficult to observe a reduction in transition latency. This is
probably a reflection of the fact that: 1) we tested learning only after first performing post-
training electrophysiological recordings during which mice received saccharin without any
associated adverse effects; and 2) we tested responses to a whole battery of tastes both just
before and just after the training sessions. Both of these features of our preparation would be
likely to interfere with the expression of learning—by the time we assessed CTA, some
extinction (Nolan et al., 1997; Bouton, 2004; Moran & Katz, 2014) had probably occurred
because of the safe presentation of saccharin, and some diffusion (Richardson et al., 1984;
McLaren & Mackintosh, 2002; Heyer et al., 2003) or overshadowing (Hall & Symonds, 2006;
Kwok et al., 2012) of learning had probably also occurred because of the peri-training
presentation of other tastes.
303
304 *Stk11KO impacts taste processing*
305
While our prior work (Levitan et al., 2020) did not unearth significant differences in the
overall structure of taste processing caused by Stk11 KO, the more in-depth analysis performed
here reveals some interpretable abnormalities in the neural taste processing of mice lacking
Stk11 in BLApn. At the most basic level, putative pyramidal neurons in GC show slightly higher
basal firing rates in Stk11 KO than control mice—a change easily attributable to the previously-
noted increase of excitability in BLApn, and to the subsequent increase of excitatory drive into
GC from BLA. Together with the observed elevation in palatability-relatedness of firing
throughout the taste response in Stk11 KO animals (Figure 3), these data suggest that one
result of losing BLApn Stk11 is an alteration in the impact of BLA on GC, such that the
magnitude of the transfer of information from one to the other is enhanced.
This latter novel result may have been previously undetectable due to our prior study's
inclusion of sucrose in the taste battery: as sucrose has reliably high palatability (and
correspondingly large differences in response firing rates from other tastants), it is likely that its
inclusion swamps the impact of other (subtler) response differences on the correlation
computation. By only considering the responses of the NaCl, quinine and citric acid here, we
may actually obtain a clearer picture of how these subtler taste response differences lay out
with respect to taste palatability. This conclusion is consistent with the suggestion that the loss
of Stk11 enhances the contribution of BLA to GC activity (*via* increases in PN excitability).
We also observed a difference in how naïve Stk11 KO mice process saccharin. Typically,
saccharin taste responses are more similar to NaCl than to quinine, by virtue of the palatability-
relatedness of late epoch coding and the assumption that NaCl and saccharin are similarly
palatable (Moran & Katz, 2014). In Stk11 KO mice, however, the *de novo* saccharin
representation is significantly closer to quinine than controls in both epochs of the taste
response. Though separable from the actual calculation of palatability (which was evaluated in
relation to coding of quinine, NaCl, and citric acid), this does suggest a certain corruption of
information content – or the organization of information – in the GC of Stk11 KO mice. Future

work will investigate whether perhaps Stk11 KO mice naturally find saccharin less appealing
than do control mice, as has been found with other mouse strains and manipulations (Yirmiya
et al., 1988; Harkin et al., 2002; O'Brien et al., 2013; Awad et al., 2020).
Finally, the ensemble dynamics of naïve taste processing are also altered by KO of Stk11
in BLApn, in that the onset of the late state in single trials tends to be greatly delayed prior to
training (Figure 7B). With this transition being directly linked to the behavioral decision of
whether to spit or swallow the tastant (Sadacca et al., 2016; Mukherjee et al., 2019), this large
delay in the occurrence of this transition merits further investigation as it could mean a large
delay in making a potentially life-saving decision; alternatively, it could mean that these
abnormally timed and coded states no longer command the same behavioral relevance in the
KO mice. Further experiments looking at both amygdalar responses to tastants and the timings
of orofacial responses to tastants will place the changes in GC taste processing observed here
into richer context, and thereby help to explain further how the circuit is disrupted.
345
346 *CTA-related response plasticity is abnormal (but present) in Stk11KO mice*
347
Our naïve prediction was that the lack of learning in Stk11KO mice should be reflected in
a concomitant lack of response plasticity. There was ample *prima facie* reason to doubt this
simplest hypothesis, however, starting with the fact that the impact of a loss of Stk11 in BLApn
is actually to increase these neurons' basal excitability (Levitan et al., 2020) – a change
conceptually associated with increases in network plasticity (Cohen et al., 1999; Malik &
Chattarji, 2012; Yiu et al., 2014).
In fact, the initial hypothesis proved premature: we observed that GC saccharin
responses of single neurons in Stk11 KO animals did change significantly with CTA training
(Figure 4); these changes were not "learning-appropriate," however, in that they centered on a
period prior to the palatability epoch (i.e., <750ms post-stimulus delivery) that is involved in
stimulus identification (Yoshida & Katz, 2011), and left the later, normally learning-sensitive
phase of the responses, relatively untouched. In fact, the later onset of the palatability state in
these mice (at ~980ms post-stimulus delivery) places the training-related changes seen firmly
within this earlier identification epoch of the taste response. Furthermore, the GC coding
dynamics in Stk11 KO mice, which (again) are already abnormal prior to learning, are rendered
incoherent by CTA training, with the very organization of state-to-state transition times falling
apart (Figure 8B, bottom). In summary, the course of CTA training causes plastic changes in
mice lacking Stk11 in BLApn, but the nature of the changes is such that they disrupt the
processing of the training taste and fail to support the learned behavior.
If anything, the changes in Stk11 KO taste responses wrought by our training regimen
are specifically unsupportive of CTA. While single neurons exhibit earlier changes in their
responses to saccharin, the representational content (i.e., the relationship between saccharin
responses and those to tastes whose positive or negative hedonics are untouched by training)
remains mostly constant. Not only does "learning" impact the "wrong" portion of the response,
that influence fails to change coding in the right direction. These data further suggest that the
loss of Stk11 alters the timing of the impact of BLA on GC, changing the window of the taste
response impacted by the training.

Finally, it is important to consider the fact that GC, like most sensory cortices (Olshausen
& Field, 2004; Niell & Stryker, 2008; Poo & Isaacson, 2009; Sakata & Harris, 2009; Crochet et al.,
2011), normally relies on sparse firing to encode and process gustatory information. The
increased input from BLA results in increased activity in GC; baseline firing rates (in putative
pyramidal neurons) are higher, more neurons are active in response to tastes appearing on the
tongue, and palatability-related firing is enhanced. A consequence of this abnormality is the
augmentation of GC's normal sparse coding scheme with additional (but not necessarily useful)
information. It is entirely possible, given this fact, that the loss of Stk11 in BLApn alters the
perceptual properties of specific tastants (our data in fact suggests that saccharin might be
perceived differently by KO mice). If so, even if naïve mice can distinguish tastes, the "trained"
mouse might fail to identify the tastant associated with the gastric malaise as the same tastant
presented prior to training. This abnormality is also consistent with the disturbed organization
of taste representations seen in Stk11 KO mice prior to – and over the course of – CTA training
(Figure 5). In future work, we will investigate whether Stk11 KO mice can discriminate between
tastants reliably, and whether they can identify a conditioned tastant as the same stimulus that
had been presented prior to CTA training.
391
392   **Materials and Methods**
393
394   *Subjects*
395
Male Stk11$^{fl/fl}$ (B6(Cg)$^{Stk11tm1.1Sjm}$/J) mice aged P60-P90 were used for these experiments. Mice
were kept on a 12-hour light-dark cycle and given ad libitum access to food and water except
during experiments habituation and recording days, at which time water access was restricted,
while food remained available ad libitum. All procedures were approved by the Brandeis
University Institutional Animal Care and use Committee (IACUC) in accordance with NIH
guidelines.
402
403   *Surgery*
404
All animals were habituated to the testing environment prior to surgery. Mice were initially
anesthetized with an I.P. injection of a ketamine/xylazine (K.X.) mixture (50 mg/kg & 2.66
mg/kg, respectively). Their heads were shaved and then inserted into the stereotaxic frame
(David Kopf Instruments; Tujunga, CA). Anesthesia was maintained with 0.5 – 2% isoflurane.
The skull was exposed and cleaned, and craniotomies were made for BLA virus injection (-1.4
mm AP, ±3.4 mm ML) and GC electrode implantation (+1.0mm AP, ±3.4mm ML). BLA was
injected bilaterally with AAV2/5-Camk2α::Cre-GFP (Stk11 KO; N=6) or a control virus consisting
of AAV2/5-Camk2α::GFP (Control; N=10) (UNC, vector core) using sterile glass micropipettes
(10-20µm diameter) attached to a partially automated microinjection device (Nanoject III
Microinjector, Drummond Scientific; Broomall, PA). The micropipettes were slowly lowered into
BLA at 4.6 mm then 4.3 mm from the dura; At each depth, 200nL of the viral solution were
delivered via 10 pulses of 20nL every 10 seconds. The micropipette was retracted 10 minutes
after the dorsal-most injection on each side. Small burr holes were drilled anterior to lambda on
either side of the midline and M1 stainless steel screws were inserted both to act as grounding

contacts and to anchor the assembly to the skull. Two 16-wire bundles of formvar-coated 25
μm nichrome wire attached to mini-microdrives were slowly lowered to 0.5mm above ventral
GC (2.5 mm from the dura) as done previously (Levitan et al., 2019; Levitan et al., 2020) and
secured in place with C&B Metabond (Parkell; Edgewood, NY). To aid with later identification of
electrode locations, bundle tips were lightly coated with DiL stain (ThermoFisher Scientific;
Indianapolis, IN) before implantation.
Mice were given a booster I.P. injection of the original induction mix 10 minutes before
being briefly removed from the stereotaxic frame, so that an intraoral cannula (IOC; flexible
plastic tubing) could be inserted close to the tongue inside the cheek (Fontanini & Katz, 2006),
after which they were replaced in the stereotaxic frame, and stainless-steel grounding wire was
connected from the circuit board to the grounding screws. The craniotomies were sealed with
Kwik-Sil (World Precision Instruments; Sarasota, FL) and components were embedded in dental
acrylic and thereby secured to the skull.
Mice were allowed to recover on a heating mat before being returned to their home cages.
Post-operative injections (s.c.) of an analgesic (Metacam; 2 mg/kg; Boehringer Ingelheim
Animal Health USA Inc.; Duluth, GA), lactated ringer's solution (0.4 ml) and an antibiotic (Pro-
Pen-G; 150,000 U/kg; Bimeda; Oakbrook Terrace, IL) were given immediately following surgery
and at 24hr and 48hr later. Injections of lactated ringer's solution continued for several extra
days until weight stabilized.
438
439 *Behavioral Procedures*
440
After 7d of recovery from surgery, mice were habituated to the testing environment for 30 min
per day. At post-op day 10, animals were placed on a water restriction schedule for the
remainder of the experiment (6 days) and habituated to receiving ~1ml (60 pulses, 16-18 μL per
pulse, 15-25ms between pulses) of water through the IOC in the morning (they also received 10
minutes of free water access in their home cages in evening). On Post-op days 11 & 12,
electrode bundles were lowered a total of 0.5mm, using the built-in microdrive assembly;
recording started 24hrs later—4 sessions over 3 days. The first session (Tastes 1) consisted of
60 IOC deliveries of a battery of 4 taste solutions (15 deliveries per taste, ~16 μL per delivery,
15-25ms between deliveries). These solutions were water, 0.02M citric acid, 0.1M NaCl and
0.001M quinine (concentrations chosen to maximize comparability with earlier papers (Moran
& Katz, 2014; Levitan et al., 2019)); 2 tastes were delivered in pseudo-random order for the first
half of the session, and then the others. Pairings differed for each animal, but citric acid and
quinine were never paired, in order to prevent animals from receiving too many consecutive
un-palatable taste deliveries (Tastes 1).
The next day mice were given 60 deliveries of saccharin (0.02 M, 16-18μL per delivery; CTA
Train session). Due to an equipment malfunction, 2 animals were only given 45 deliveries (20-
21μL per delivery) to keep the amount of total amount of saccharin consumed during training
the same across animals. Five minutes after the final saccharin delivery, they were given an I.P.
injection of 0.15M LiCl (2% b.w.) to induce gastric malaise. This procedure allows, under normal
circumstances, a selective association (condition taste aversion, CTA) to form between the
malaise and the novel tastant. One day later, animals received 15 deliveries of saccharin (CTA
Test), and 1-2hr later underwent another session of the battery of 4 tastes (Tastes 2). The

evening after the final recording session, instead of 10 min access to water, animals received 10
min access to saccharin solution in order to evaluate strength of CTA. To assess CTA, the
amount of saccharin consumed was compared to water consumption on previous nights. Based
on prior work in the lab (and since the earlier consumption of saccharin as well as other tastes
likely resulted in a certain mildness of learning), we chose a saccharin consumption of 80% of
average nightly water consumption as a conservative threshold, below which animals were
deemed to have learned the CTA; this was an adequate criterion for our purposes because the
primary measure was difference between groups, and thus the prediction was that many
control mice would reach this criterion but that few Stk11KO mice would do so. 5 of the 10
control mice reached this learning criterion, and 5 of the 6 Stk11 KO mice did not. The one
Stk11 KO mouse whose saccharin consumption was below the threshold was excluded from
analysis because that animal's nightly water consumption steadily decreased over habituation
days, and we could not be sure if its lack of saccharin consumption was an effect of CTA
learning or some other impairment resulting in reduced intake.
477
478 *Electrophysiology*
479
During taste presentations, up to 32 extracellular potentials were simultaneously sampled and
digitized by an Intan RHD2132 headstage and sent to a DAQ @ 30kHz (Intan Technologies; Los
Angeles, CA). These time series, as well as time-stamped records of stimulus onset and offset
times, were stored for offline analysis in python using the blechpy analysis package:
discriminable action potentials with at least a 3:1 signal-to-noise ratio were isolated offline by
amplitude thresholding; 2ms snippets of each waveform were extracted, up-sampled 10x and
aligned to the wave peak. The UMAP algorithm was used to extract waveform features, which
were fit by Gaussian mixture models to cluster spiking events—clusters that were then
manually curated (by experimenters blind to taste responses and injection type) to ensure
conservatism. Only clusters clearly separated in UMAP or PCA space, and containing no 1ms ISI
violations and fewer than 1% 2ms ISI violations, were deemed to reflect single-neuron activity.
These methods and criteria are in accordance with previously published methods from our lab
(Moran & Katz, 2014; Sadacca et al., 2016; Levitan et al., 2019; Mukherjee et al., 2019).
493
494 *Histology*
495
496 Animals were deeply anesthetized with isoflurane and euthanized with a K.X. overdose. Animals
497 were then perfused transcardially with 0.9% saline solution followed by 4% formaldehyde in
498 PBS. Brains were extracted and stored in 4% formaldehyde, 30% sucrose solution. Once fully
499 fixed, 50 μm coronal sections of BLA and GC were visualized under a fluorescent microscope
500 (KEYENCE BZ-X800; Itasca, IL) to verify injection and electrode positions.
501
502 *Detection of single neurons reliably held across sessions*
503
To determine if a waveform recorded from a single electrode for multiple sessions represented
the same single neuron, we used a spike shape analysis successfully used in several previous
studies (Nicolelis et al. 2003; Grossman et al. 2008; Herry et al. 2008; Moran & Katz 2014). A

non-parametric clustering statistic (J_3) was used to quantify the separation and grouping
between 2 clusters. J_3-values were computed for all single neurons across all sessions using
spikes from the first and last thirds of each recording session to generate a distribution of intra-
session J_3 values of known single neurons. Then, for each pair of single cells on the same
electrode between 2 consecutive sessions, the J_3 was computed and compared to the intra-
session distribution. A conservative criterion was applied: only pairs of units with an inter-
session J_3 value less than the 95[th] percentile of the intra-session J_3 distribution (for the entire
experimental group) were deemed to be held across sessions. This mean that only pairs of units
from across days that are more similar to each other than at least 5% of single units are to
themselves within a recording session.
517
518 *Analysis of single neuron taste responses*
519
Single neurons were classified as regular spiking or fast-spiking based on inspection of spike
waveform shape and ISI histograms (for details see Levitan et al., 2019). Regular spiking cells
were taken as putative pyramidal neurons while fast-spiking cells were labelled as
interneurons. Baseline firing rates were computed as the average spike rates over the 1s period
prior to stimulus deliveries. Response firing rates were initially computed to be the average
spike rate over the entire 1.5s post-stimulus period. Two-way ANOVAs were used to assess the
effects of group (Stk11 KO vs Control) and training (pre- vs post- CTA) on baseline and response
firing rates for each unit type (Figure 2).
Using the data from the pre- and post-CTA taste battery sessions (Tastes 1 and Tastes 2),
we also looked at the correlation between the response firing rates and the rank-order hedonic
values of the given tastants (Quinine [1], Citric Acid [2] and NaCl [3]; Figure 3). This allowed us
to compare the decision-relevant information content of the single-neuron responses between
groups. Firing rates were computed in non-overlapping 250ms bins. Within each time bin, the
Spearman correlation coefficient (R^2) was computed for the correlation between the ranked
firing rates and the palatability values. A two-way ANOVA was used to investigate the effects of
group and training on the correlation of responses to palatability. All ANOVA analyses used
Bonferroni corrected post-hoc t-tests for within group comparisons.
537
538 *Changes in held neuron responses to Saccharin over training*
539
To understand the time course of changes to single neuron responses to Saccharin over
training, we analyzed the responses of neurons that were deemed reliably held from the CTA
Train session to the CTA Test session (Figure 4). For these, firing rates were computed in six
consecutive, non-overlapping 250ms bins post-stimulus, similar to the manner used previously
(Grossman et al., 2008). Response changes in each time bin were evaluated using a one-way
ANOVA (p-values were Bonferroni corrected), and the number of neurons showing significant
changes across training in each time bin was compared between groups using a χ^2-test.
Due our conservative criteria for held neurons, this analysis was the limiting factor for
determining sample size needed for our study. Preliminary data suggested this analysis would
have an effect size of around 0.960 (Cramer's V) which would have required a total sample of
15 neurons to reliably detect at a power 0.95 (computed using G*Power 3.1; see Faul et al.,

2009). Post-hoc analysis found that our final effect size was 0.580; with a final sample size of 38
held cells, this resulted in a final power of 0.946 for our analysis.
553
554 *Multidimensional Scaling (MDS)*
555
Multidimensional scaling allows for the reduction of dimensionality in multi-variate data while
preserving the relative two-dimensional distances between points. We used MDS, as has been
done many times previously (e.g. Chang & Scott, 1984; Geran & Travers, 2006; Moran & Katz,
2014), to analyze the similarity between saccharin taste responses and those of NaCl (palatable)
and quinine (unpalatable). Activity projections into 2D space were computed separately for the
early (0-750ms) and late (750-1500ms) epochs of the response; in each, Euclidean distances
were computed between representations of individual saccharin responses and the mean
quinine (D_Q) and NaCl representations (D_N) for each recording set. These distances were then
normalized by the distance between the NaCl and quinine means (D_{QN}) to provide a
dimensionless distance metric
566
567 $$D_S = \frac{D_Q - D_N}{D_{QN}} \qquad (1)$$
568
which quantifies the similarity of the saccharin response to the NaCl and quinine
representations: positive values indicate that the saccharin representation is more similar to
NaCl than quinine, while negative values indicate the opposite (Figure 5). The populations for
this analysis consisted only of units that were reliably held from Tastes 1 to CTA Train (pre-CTA)
or from CTA Test to Tastes 2 (post-CTA), and any recording sets that did not have at least 2 held
units across these periods were excluded from this analysis.
575
576 *Poisson Hidden Markov Models (HMMs)*
577
To ascertain the underlying temporal structure of single trial ensemble responses, we fit sets of
simultaneously-recorded spike trains to Poisson HMMs (Rabiner, 1989; Abeles et al., 1995;
Jones et al., 2007; Kemere et al., 2008; Miller & Katz, 2010; Moran & Katz, 2014; Sadacca et al.,
2016). Since our goal was to look at the timing of these state transitions (and not to have the
model classify taste responses), we fit separate models for each tastant for each recording
session. We constrained the search space for our parameters, fitting models for -250 ms to
2000 ms peri-stimulus with 2-6 states connected in a feed-forward manner. The Bayesian
Information Criteria (BIC) was used to evaluate model fit, and while we saw no statistically
significant differences in BIC relative to the number of states, 2 or 3 state models consistently
fit the data best (i.e., had the lowest BIC values). Given the extensive evidence (Katz et al.,
2001; Fontanini & Katz, 2006; Miller & Katz, 2010; Piette et al., 2012; Sadacca et al., 2012;
Moran & Katz, 2014; Sadacca et al., 2016; Levitan et al., 2019) that cortical taste responses
consist of 2 major chemosensory coding states or epochs, we settled on models containing 3
states connected in feed-forward manner (Figure 6a) so as to account for the presence of a
background state, an early state and a late state.

Spike trains were provided to the model at 1ms resolution, and initial model firing rates
were randomly selected from a normal distribution using the mean and standard deviation for
each units' firing rates across trials. Values for the initial transition matrix were drawn randomly
from a normal distribution with mean 0.05 and standard deviation of 0.01. Diagonal values (the
probability of remaining in the same state at each time step) were similarly drawn from a
normal distribution with mean of 0.99 and sigma of 0.01 to ensure a high probability of
remaining in the same state, and the elements corresponding to transitions backward or
advancing more than 1 state were set to zero. Finally, all rows were normalized to sum to 1.
For each dataset, models were initialized and fit 50 separate times; in each run, fitting
terminated when the change in log likelihood over iterations dropped below 1e-10. The model
with the highest log likelihood was deemed the best description of the data. Fitting was done
using the Baum-Welch algorithm (Rabiner, 1989), which provided the probabilities of the
ensemble being in each state at each time point for each trial (Figure 7c). The optimal
sequences of states for each trial were decoded using the Viterbi decoding algorithm (Viterbi,
1967), which provides the maximum likelihood decoding of each trial based on the fitted
model. While the 3-state models routinely provided good fits to the data (Figure 6b), for
subsequent analysis we nonetheless chose early and late states independently, designating
them to be the most probable sequence of states present in the bins 200-700ms and 750-
1500ms respectively.
Only recording sessions containing at least 3 well-isolated single cells were included in
any such analysis, to avoid allowing results to be unduly biased toward the dynamics of single
neurons. Furthermore, only the first 15 saccharin trials of CTA-Train sessions were used for this
analysis since all other sessions only had 15 trials per tastant to fit; this ensured that model
quality and hyperparameters (number of states, time window, bin size, etc.) could remain
constant across all trained models.
618
619   *Analysis of HMM Transition Times*
620
As the offset times for early states and onset times for the late states were reliably
coupled, we defined transition times as the offset times for the early states. Mean transition
times (Figure 7B) were compared using a two-way ANOVA for group (Stk11 KO vs Control) and
training (pre- vs post-CTA) effects. However, as inspection of the distributions revealed large
variability in the transition times of Stk11 KO responses post-CTA, these distributions were then
compared between pre- and post-CTA within each group using χ^2-tests, and fit to Gaussian
functions (Figure 8B) to quantify the differences in transition distributions. Only trials decoded
as having both the early and late states present post-stimulus were used in these analyses.
629
630
631   References
632

Abeles, M., Bergman, H., Gat, I., Meilijson, I., Seidemann, E., Tishby, N., & Vaadia, E. (1995).
Cortical activity flips among quasi-stationary states. *Proc Natl Acad Sci U S A, 92*(19),
8616-8620. doi:10.1073/pnas.92.19.8616

Arieli, E., Gerbi, R., Shein-Idelson, M., & Moran, A. (2020). Temporally-precise basolateral
amygdala activation is required for the formation of taste memories in gustatory cortex. *J*
Physiol, 598(23), 5505-5522. doi:10.1113/JP280213
Awad, G., Roeckel, L. A., Massotte, D., Olmstead, M. C., & Befort, K. (2020). Deletion of mu
opioid receptors reduces palatable solution intake in a mouse model of binge eating.
Behav Pharmacol, 31(2&3), 249-255. doi:10.1097/FBP.0000000000000496
Beirowski, B. (2019). The LKB1-AMPK and mTORC1 Metabolic Signaling Networks in
Schwann Cells Control Axon Integrity and Myelination: Assembling and upholding
nerves by metabolic signaling in Schwann cells. *Bioessays, 41*(1), e1800075.
doi:10.1002/bies.201800075
Bouton, M. E. (2004). Context and behavioral processes in extinction. *Learning & memory (Cold*
Spring Harbor, N.Y.), 11(5), 485-494. doi:10.1101/lm.78804
Bures, J., Bermudez-Rattoni, F., & Yamamoto, T. (1998). *Conditioned Taste Aversion: Memory*
of a Special Kind: Oxford University Press.
Chang, F. C., & Scott, T. R. (1984). Conditioned taste aversions modify neural responses in the
rat nucleus tractus solitarius. *The Journal of Neuroscience, 4*(7), 1850-1862.
doi:10.1523/jneurosci.04-07-01850.1984
Cohen, A. S., Coussens, C. M., Raymond, C. R., & Abraham, W. C. (1999). Long-lasting
increase in cellular excitability associated with the priming of LTP induction in rat
hippocampus. *J Neurophysiol, 82*(6), 3139-3148. doi:10.1152/jn.1999.82.6.3139
Crochet, S., Poulet, J. F., Kremer, Y., & Petersen, C. C. (2011). Synaptic mechanisms underlying
sparse coding of active touch. *Neuron, 69*(6), 1160-1175.
doi:10.1016/j.neuron.2011.02.022
Faul, F., Erdfelder, E., Buchner, A., & Lang, A. G. (2009). Statistical power analyses using
G*Power 3.1: tests for correlation and regression analyses. *Behav Res Methods, 41*(4),
1149-1160. doi:10.3758/BRM.41.4.1149
Fonseca, E., Sandoval-Herrera, V., Simon, S. A., & Gutierrez, R. (2020). Behavioral
Disassociation of Perceived Sweet Taste Intensity and Hedonically Positive Palatability.
eNeuro, 7(5). doi:10.1523/ENEURO.0268-20.2020
Fontanini, A., & Katz, D. B. (2006). State-dependent modulation of time-varying gustatory
responses. *J Neurophysiol, 96*(6), 3183-3193. doi:10.1152/jn.00804.2006
Gallo, M., Roldan, G., & Bures, J. (1992). Differential involvement of gustatory insular cortex
and amygdala in the acquisition and retrieval of conditioned taste aversion in rats. *Behav*
Brain Res, 52(1), 91-97. doi:10.1016/s0166-4328(05)80328-6
Garcia-Delatorre, P., Perez-Sanchez, C., Guzman-Ramos, K., & Bermudez-Rattoni, F. (2014).
Role of glutamate receptors of central and basolateral amygdala nuclei on retrieval and
reconsolidation of taste aversive memory. *Neurobiol Learn Mem, 111*, 35-40.
doi:10.1016/j.nlm.2014.03.003
Geran, L. C., & Travers, S. P. (2006). Single Neurons in the Nucleus of the Solitary Tract
Respond Selectively to Bitter Taste Stimuli. *Journal of Neurophysiology, 96*(5), 2513-
2527. doi:10.1152/jn.00607.2006
Grossman, S. E., Fontanini, A., Wieskopf, J. S., & Katz, D. B. (2008). Learning-related plasticity
of temporal coding in simultaneously recorded amygdala-cortical ensembles. *J Neurosci,*
28(11), 2864-2873. doi:10.1523/JNEUROSCI.4063-07.2008
Hall, G., & Symonds, M. (2006). Overshadowing and latent inhibition of context aversion
conditioning in the rat. *Auton Neurosci, 129*(1-2), 42-49.
doi:10.1016/j.autneu.2006.07.013

Harkin, A., Houlihan, D. D., & Kelly, J. P. (2002). Reduction in preference for saccharin by
684   repeated unpredictable stress in mice and its prevention by imipramine. *J*
685   *Psychopharmacol, 16*(2), 115-123. doi:10.1177/026988110201600201

686   Heyer, B. R., Taylor-Burds, C. C., Tran, L. H., & Delay, E. R. (2003). Monosodium glutamate
687   and sweet taste: generalization of conditioned taste aversion between glutamate and
688   sweet stimuli in rats. *Chem Senses, 28*(7), 631-641. doi:10.1093/chemse/bjg056

689   Jones, L. M., Fontanini, A., Sadacca, B. F., Miller, P., & Katz, D. B. (2007). Natural stimuli
690   evoke dynamic sequences of states in sensory cortical ensembles. *Proc Natl Acad Sci U S*
691   *A, 104*(47), 18772-18777. doi:10.1073/pnas.0705546104

692   Katz, D. B., Simon, S. A., & Nicolelis, M. A. L. (2001). Dynamic and Multimodal Responses of
693   Gustatory Cortical Neurons in Awake Rats. *The Journal of Neuroscience, 21*(12), 4478-
694   4489. doi:10.1523/jneurosci.21-12-04478.2001

695   Kemere, C., Santhanam, G., Yu, B. M., Afshar, A., Ryu, S. I., Meng, T. H., & Shenoy, K. V.
696   (2008). Detecting neural-state transitions using hidden Markov models for motor cortical
697   prostheses. *J Neurophysiol, 100*(4), 2441-2452. doi:10.1152/jn.00924.2007

698   Kwok, D. W., Livesey, E. J., & Boakes, R. A. (2012). Serial overshadowing of taste aversion
699   learning by stimuli preceding the target taste. *Learn Behav, 40*(4), 427-438.
700   doi:10.3758/s13420-011-0064-0

701   Lavi, K., Jacobson, G. A., Rosenblum, K., & Luthi, A. (2018). Encoding of Conditioned Taste
702   Aversion in Cortico-Amygdala Circuits. *Cell Rep, 24*(2), 278-283.
703   doi:10.1016/j.celrep.2018.06.053

704   Levitan, D., Lin, J. Y., Wachutka, J., Mukherjee, N., Nelson, S. B., & Katz, D. B. (2019). Single
705   and population coding of taste in the gustatory cortex of awake mice. *J Neurophysiol,*
706   *122*(4), 1342-1356. doi:10.1152/jn.00357.2019

707   Levitan, D., Liu, C., Yang, T., Shima, Y., Lin, J. Y., Wachutka, J., . . . Nelson, S. B. (2020).
708   Deletion of Stk11 and Fos in mouse BLA projection neurons alters intrinsic excitability
709   and impairs formation of long-term aversive memory. *Elife, 9*. doi:10.7554/eLife.61036

710   Li, J. X., Maier, J. X., Reid, E. E., & Katz, D. B. (2016). Sensory Cortical Activity Is Related to
711   the Selection of a Rhythmic Motor Action Pattern. *J Neurosci, 36*(20), 5596-5607.
712   doi:10.1523/JNEUROSCI.3949-15.2016

713   Lin, J. Y., Mukherjee, N., Bernstein, M. J., & Katz, D. B. (2021). Perturbation of amygdala-
714   cortical projections reduces ensemble coherence of palatability coding in gustatory
715   cortex. *Elife, 10*. doi:10.7554/eLife.65766

716   Maffei, A., Haley, M., & Fontanini, A. (2012). Neural processing of gustatory information in
717   insular circuits. *Curr Opin Neurobiol, 22*(4), 709-716. doi:10.1016/j.conb.2012.04.001

718   Malik, R., & Chattarji, S. (2012). Enhanced intrinsic excitability and EPSP-spike coupling
719   accompany enriched environment-induced facilitation of LTP in hippocampal CA1
720   pyramidal neurons. *J Neurophysiol, 107*(5), 1366-1378. doi:10.1152/jn.01009.2011

721   McLaren, I. P., & Mackintosh, N. J. (2002). Associative learning and elemental representation:
722   II. Generalization and discrimination. *Anim Learn Behav, 30*(3), 177-200.
723   doi:10.3758/bf03192828

724   Miller, P., & Katz, D. B. (2010). Stochastic transitions between neural states in taste processing
725   and decision-making. *J Neurosci, 30*(7), 2559-2570. doi:10.1523/JNEUROSCI.3047-
726   09.2010

727   Moran, A., & Katz, D. B. (2014). Sensory cortical population dynamics uniquely track behavior
728   across learning and extinction. *J Neurosci, 34*(4), 1248-1257.
729   doi:10.1523/JNEUROSCI.3331-13.2014

Mukherjee, N., Wachutka, J., & Katz, D. B. (2019). Impact of precisely-timed inhibition of gustatory cortex on taste behavior depends on single-trial ensemble dynamics. *Elife, 8*. doi:10.7554/eLife.45968

Nakada, D., Saunders, T. L., & Morrison, S. J. (2010). Lkb1 regulates cell cycle and energy metabolism in haematopoietic stem cells. *Nature, 468*(7324), 653-658. doi:10.1038/nature09571

Niell, C. M., & Stryker, M. P. (2008). Highly selective receptive fields in mouse visual cortex. *J Neurosci, 28*(30), 7520-7536. doi:10.1523/JNEUROSCI.0623-08.2008

Nolan, L. J., McCaughey, S. A., Giza, B. K., Rhinehart-Doty, J. A., Smith, J. C., & Scott, T. R. (1997). Extinction of a Conditioned Taste Aversion in Rats: I. Behavioral Effects. *Physiology & Behavior, 61*(2), 319-323. doi:10.1016/s0031-9384(96)00411-8

O'Brien, L. D., Wills, K. L., Segsworth, B., Dashney, B., Rock, E. M., Limebeer, C. L., & Parker, L. A. (2013). Effect of chronic exposure to rimonabant and phytocannabinoids on anxiety-like behavior and saccharin palatability. *Pharmacol Biochem Behav, 103*(3), 597-602. doi:10.1016/j.pbb.2012.10.008

Olshausen, B. A., & Field, D. J. (2004). Sparse coding of sensory inputs. *Curr Opin Neurobiol, 14*(4), 481-487. doi:10.1016/j.conb.2004.07.007

Piette, C. E., Baez-Santiago, M. A., Reid, E. E., Katz, D. B., & Moran, A. (2012). Inactivation of basolateral amygdala specifically eliminates palatability-related information in cortical sensory responses. *J Neurosci, 32*(29), 9981-9991. doi:10.1523/JNEUROSCI.0669-12.2012

Poo, C., & Isaacson, J. S. (2009). Odor representations in olfactory cortex: "sparse" coding, global inhibition, and oscillations. *Neuron, 62*(6), 850-861. doi:10.1016/j.neuron.2009.05.022

Rabiner, L. R. (1989). A tutorial on hidden Markov models and selected applications in speech recognition. *Proceedings of the IEEE, 77*(2), 257-286. doi:10.1109/5.18626

Richardson, R., Williams, C., & Riccio, D. C. (1984). Stimulus generalization of conditioned taste aversion in rats. *Behavioral and Neural Biology, 41*(1), 41-53. doi:10.1016/s0163-1047(84)90706-4

Sadacca, B. F., Mukherjee, N., Vladusich, T., Li, J. X., Katz, D. B., & Miller, P. (2016). The Behavioral Relevance of Cortical Neural Ensemble Responses Emerges Suddenly. *J Neurosci, 36*(3), 655-669. doi:10.1523/JNEUROSCI.2265-15.2016

Sadacca, B. F., Rothwax, J. T., & Katz, D. B. (2012). Sodium concentration coding gives way to evaluative coding in cortex and amygdala. *J Neurosci, 32*(29), 9999-10011. doi:10.1523/JNEUROSCI.6059-11.2012

Sakata, S., & Harris, K. D. (2009). Laminar structure of spontaneous and sensory-evoked population activity in auditory cortex. *Neuron, 64*(3), 404-418. doi:10.1016/j.neuron.2009.09.020

Shaw, R. J. (2009). LKB1 and AMP-activated protein kinase control of mTOR signalling and growth. *Acta Physiol (Oxf), 196*(1), 65-80. doi:10.1111/j.1748-1716.2009.01972.x

Viterbi, A. (1967). Error bounds for convolutional codes and an asymptotically optimum decoding algorithm. *IEEE Transactions on Information Theory, 13*(2), 260-269. doi:10.1109/tit.1967.1054010

Yamamoto, T., Azuma, S., & Kawamura, Y. (1984). Functional relations between the cortical gustatory area and the amygdala: electrophysiological and behavioral studies in rats. *Exp Brain Res, 56*(1), 23-31. doi:10.1007/BF00237438

Yamamoto, T., Fujimoto, Y., Shimura, T., & Sakai, N. (1995). Conditioned taste aversion in rats
with excitotoxic brain lesions. *Neuroscience Research, 22*(1), 31-49. doi:10.1016/0168-
0102(95)00875-t
Yirmiya, R., Lieblich, I., & Liebeskind, J. C. (1988). Reduced saccharin preference in CXBK
(opioid receptor-deficient) mice. *Brain Research, 438*(1-2), 339-342. doi:10.1016/0006-
8993(88)91360-1
Yiu, A. P., Mercaldo, V., Yan, C., Richards, B., Rashid, A. J., Hsiang, H. L., . . . Josselyn, S. A.
(2014). Neurons are recruited to a memory trace based on relative neuronal excitability
immediately before training. *Neuron, 83*(3), 722-735. doi:10.1016/j.neuron.2014.07.017
Yoshida, T., & Katz, D. B. (2011). Control of prestimulus activity related to improved sensory
coding within a discrimination task. *J Neurosci, 31*(11), 4101-4112.
doi:10.1523/JNEUROSCI.4380-10.2011
788

789
Figure 1. *Overview of experimental setup.* (A) Genetic constructs Cre-GFP (Stk11 KO) and GFP
(Control) under the control of a CamKIIα promoter were encapsulated in AAV2/5 viral vectors.
(B) Overview of the experimental timeline starting from the virus injection and electrode/IOC
implantation surgery through to recording days – Tastes 1, CTA Train, CTA Test and Taste 2.
Except for the 10-day recovery, the spacing between blocks indicates the passage of 1 day. (C)
Representative histological slices showing viral transfection of BLA (left; -1.2mm AP) and co-
localization of electrodes (red, DiL stain) and BLA axon terminals (green, GFP) in ventral GC
(right; +1mm AP).

798
799 **Figure 2.** *Pyramidal neurons in Stk11 KO mice exhibit elevated baseline firing rates.* Baseline
800 (top; 1s pre-stimulus) and response (bottom; 1.5s post-stimulus) firing rates for pyramidal cells
801 (left) and interneurons (right) in both groups over CTA training. Error bars show the 95%
802 confidence intervals (CI) for the means. Pyramidal cell basal firing rates were slightly (95% CI of
803 difference: [0.396, 1.03] Hz), but significantly, elevated in Stk11 KO mice. *p<0.05.

804
Figure 3. *Enhanced correlation to taste palatability in single neuron taste responses of Stk11 KO*
mice. (A) Peri-stimulus time histograms of representative single neurons from control (top) and
Stk11 KO (bottom) mice in response to tastes from the Tastes 1 (pre-CTA) session: NaCl (blue),
Citric Acid (orange) and Quinine (green); water was excluded since it's palatability could not be
properly ranked relative to the other stimuli. The shaded regions show the SEM, and the dashed
black line indicates the Spearman R^2 correlation coefficient (right axis) for the unit firing rates to
the rank-order palatabilites of the tastants; note that for both, the palatability-relatedness of the
firing increased across the 2nd half-second of the response. (B) Average correlation values across
1.5 sec post-stimulus for all cells. Error bars represent the 95% CI of the means. The palatability-
relatedness of GC firing was enhanced in Stk11 KO animals compared to controls (95% CI of
difference: [0.0286, 0.0435]); ***p<0.001.

816
Figure 4. *CTA training causes earlier changes to single-neuron saccharin responses in Stk11 KO*
mice than controls. (A) Saccharin-induced firing rates (relative to mean baseline) before and after
CTA for representative single neurons from control (left) and Stk11 KO (right) mice. Asterisks
with horizontal lines identify significant differences — the results of one-way ANOVAs conducted
on each time bin and Bonferroni corrected. (B) Percent of single neurons held across training and
testing sessions for which training significantly changed firing rates, for each time bin. For Stk11
KO mice (red), changes were found in earlier portions of the responses than was true for control
animals (green); *p<0.05, **p<0.01, ***p<0.001.

825
Figure 5. *Ensemble taste representations of saccharin are impaired* de novo *and fail to change*
with CTA training. A plot of a metric showing the relative normalized distances between the
saccharin representation and the average NaCl and quinine representations. Control animals
showed a significant shift in the saccharin representation towards greater similarity to quinine
over CTA learning in the late half of the response. Conversely, in Stk11 KO animals pre-CTA
representations of saccharin did not move significantly across training. Groups were compared
with non-parametric Kruskal-Wallis tests (see Table 1). Error bar reflect the 95% CI of the mean,
and significance stars represent results of post-hoc Games-Howell tests; ***p<0.001.

Table 1a. Summary of MDS relative distances for Saccharin trials (Figure 5)

			Mean Relative Distance	
			$(D_Q - D_N)/D_{QN}$	
Group	Learning Epoch	N (trials)	Early (0-750ms)	Late (750-1500 ms)
Control	Pre-CTA	135	0.200 ± 0.633	0.488 ± 0.578
	Post-CTA	60	-9.27E-3 ± 0.743	-0.0401 ± 0.607
Stk11-	Pre-CTA	165	-0.192 ± 0.821	0.0349 ± 0.770
	Post-CTA	60	-0.0908 ± 0.643	-0.121 ± 0.558
Kruskal Wallis	Statistic (dof)		28.2 (3)	50.1 (3)
	p-value		3.30E-06	7.66E-11

± represents the st. dev.

Table 1b. Games-Howell post-hoc results for the Early Epoch (0-750 ms)

		Mean Difference (Hedges' g)			
		Control		Stk11-	
Group	Learning Epoch	Pre-CTA	Post-CTA	Pre-CTA	Post-CTA
Control	Pre-CTA	--	--	--	--
	Post-CTA	-0.209 ± 0.110 (-0.293)	--	--	--
Stk11-	Pre-CTA	-0.391 ± 0.0840 (-0.539)***	0.183 ± 0.115 (-0.238)	--	--
	Post-CTA	-0.291 ± 0.0994 (-0.452)*	-0.0815 ± 0.127 (-0.117)	-0.101 ± 0.105 (0.145)	--

Table 1c. Games-Howell post-hoc results for the Late Epoch (750-1500 ms)

		Mean Difference (Hedges' g)			
		Control		Stk11-	
Group	Learning Epoch	Pre-CTA	Post-CTA	Pre-CTA	Post-CTA
Control	Pre-CTA	--	--	--	--
	Post-CTA	-0.528 ± 0.0928 (-0.880)***	--	--	--
Stk11-	Pre-CTA	-0.453 ± 0.0779 (-0.674)***	0.0750 ± 0.0986 (-0.114)	--	--
	Post-CTA	-0.609 ± 0.0875 (-1.08)***	-0.0805 ± 0.106 (-0.137)	-0.156 ± 0.0936 (-0.250)	--

± represents the SEM
*p<0.05
***p<0.001

834
835 **Table 1.** *Summary of statistics for comparison of MDS relative distances presented in Figure 5.*

836
Figure 6. *Fits of 3-state feed-forward Hidden Markov Models to GC ensemble taste responses.*
(A) A depiction of the hidden state sequence used to fit ensemble taste responses, with black
arrows indicating possible state transitions. (B) Representative trials with HMM solutions for each
tastant. Each raster shows ensemble spiking activity (black, one row per neuron) for a single trial
in response to a taste delivered at time 0. The colored lines indicate the probability of being in each
state at each time point (y-axis), and the background color reflects the state sequences as
determined by the Viterbi decoding algorithm.

Figure 7. *In Stk11 KO mice, state transitions are later Pre-CTA, and more variable Post-CTA.*
(A) Representative HMM fits for training (top) and testing (bottom) session responses to saccharin.
(B) Comparison of the mean transition times from the early to the late states. Transition times were
significantly later in Stk11 KO mice than in control before CTA training. Error bars represent the
95% CI of the mean. (C) Mean probability of being in the "late" state at each post-stimulus time
point for each group. In Stk11 KO mice, the rise in probability pre-CTA occurred noticeably late,
while in "post-CTA" testing sessions it is much more gradual (compared to the more sudden
transitions seen in other groups), a result suggesting greater variability in transition times on
individual trials. Shaded region depicts the 95% CI of the mean.; ***p<0.001.

Figure 8. *State transitions become less consistent in Stk11 KO mice following (unsuccessful) CTA*
training. (A) A depiction of decoded state sequences for all well-fit saccharin trials, sorted by late
state onset time. Sequences produced by Stk11 KO mice show more varied structure (i.e., more
gradual slope) after training. (B) Probability density histograms of transition times with lines
representing gaussian fits of the distributions. χ^2-tests reveal significantly different training/testing
distributions of transition times for Stk11 KO mice ($\chi^2(1)=41.0$, p=1.47e-5, w=0.755) but not
controls ($\chi^2(1)=11.6$, p=0.505, w=0.361). Furthermore, the distribution of Stk11 KO post-CTA
transitions times was more poorly fit by a gaussian ($r^2=0.0402$) than were all other groups ($r^2>0.7$)
demonstrating not just a shift in the distribution, but a loss of consistent structure in the taste
responses. Significance stars represent the comparison between distributions, not between fitted
lines; ***p<0.001.