The human language system, including its inferior frontal component in 'Broca's area',

does not support music perception

Authors

Xuanyi Chen*[1,2,3], Josef Affourtit[2,3], Rachel Ryskin[2,3,4], Tamar I. Regev[2,3], Samuel

Norman-Haignere[5], Olessia Jouravlev[2,3,6], Saima Malik-Moraleda[2,3,7], Hope Kean[2,3],

Rosemary Varley†[8], and Evelina Fedorenko*†[2,3,7]

† Co-senior authors

* Corresponding Authors

jchenxy17@gmail.com and evelina9@mit.edu; 43 Vassar Street, Room 46-

3037, Cambridge, MA, 02139

Affiliations

[1]Department of Cognitive Sciences, Rice University, TX 77005, USA

[2]Department of Brain and Cognitive Sciences, MIT, Cambridge, MA 02139, USA

[3]McGovern Institute for Brain Research, MIT, Cambridge, MA 02139, USA

[4]Department of Cognitive & Information Sciences, University of California, Merced,

Merced, CA 95343, USA

[5]Zuckerman Mind, Brain, Behavior Institute, Columbia University, New York, NY

10027, USA

[6]Department of Cognitive Science, Carleton University, Ottawa, Ontario, Canada

[7]The Program in Speech and Hearing Bioscience and Technology, Harvard University,

Cambridge, MA 02138, USA

[8]Psychology & Language Sciences, UCL, London, WCN1 1PF, UK

Abstract

Language and music are two human-unique capacities whose relationship remains debated.

Some have argued for overlap in processing mechanisms, especially for structure processing.

Such claims often concern the inferior frontal component of the language system located within

'Broca's area'. However, others have failed to find overlap. Using a robust individual-subject

fMRI approach, we examined the responses of language brain regions to music stimuli, and

probed the musical abilities of individuals with severe aphasia. Across four experiments, we

obtained a clear answer: music perception does not engage the language system, and judgments

about music structure are possible even in the presence of severe damage to the language

network. In particular, the language regions' responses to music are generally low, often below

the fixation baseline, and never exceed responses elicited by non-music auditory conditions, like

animal sounds. Further, the language regions are not sensitive to music structure: they show low

responses to intact and structure-scrambled music, and to melodies with vs. without structural

violations. Finally, in line with past patient investigations, individuals with aphasia who cannot

judge sentence grammaticality perform well on melody well-formedness judgments. Thus the

mechanisms that process structure in language do not appear to process music, including music

syntax.

Keywords: language, music, syntactic processing, fMRI, domain specificity

Introduction

To interpret language or appreciate music, we must understand how different elements—words in language, notes and chords in music—relate to each other. Parallels between the structural properties of language and music have been drawn for over a century (e.g., Riemann 1877, as cited in Swain 1995; Lindblom and Sundberg 1969; Fay 1971; Boiles 1973; Cooper 1973; Bernstein 1976; Sundberg and Lindblom 1976; Lerdahl and Jackendoff 1977, 1983; Roads and Wieneke 1979; Krumhansl and Keil 1982; Baroni et al. 1983; Swain 1995; cf. Jackendoff 2009; Temperley 2022). However, the question of whether music processing relies on the same mechanisms as those that support language processing continues to spark debate.

The empirical landscape is complex. A large number of studies have argued for overlap in structural processing based on behavioral (e.g., Fedorenko et al. 2009; Slevc et al. 2009; Hoch et al. 2011; Van de Cavey and Hartsuiker 2016; Kunert et al. 2016), ERP (e.g., Janata 1995; Patel et al. 1998; Koelsch et al. 2000), MEG (e.g., Maess et al. 2001), fMRI (e.g., Koelsch et al. 2002; Levitin and Menon 2003; Tillmann et al. 2003; Koelsch 2006; Kunert et al. 2015; Musso et al. 2015), and ECoG (e.g., Sammler et al. 2009, 2013; Rietmolen et al. 2022) evidence (see Tillman 2012; Kunert and Slevc 2015; LaCroix et al. 2016, for reviews). However, we would argue that no prior study has compellingly established reliance on shared syntactic processing mechanisms in language and music.

First, evidence from behavioral, ERP, and, to a large extent, MEG studies is indirect because

they do not allow to unambiguously determine where neural responses originate (in ERP and

MEG, this is due to the 'inverse problem'; Tarantola 2004; Baillet et al. 2014).

Second, the bulk of the evidence comes from structure-violation paradigms. In such paradigms,

responses to the critical condition—which contains an element that violates the rules of tonal

music—are contrasted with responses to the control condition, where stimuli obey the rules of

tonal music. (For language, syntactic violations, like violations of number agreement, are often

used.) Because structural violations (across domains) constitute unexpected events, a brain

region that responds more strongly to the structure-violation condition than the control (no

violation) condition *may* support structure processing in music, but it may also reflect domain-

general processes, like attention or error detection/correction (e.g., Bigand et al. 2001; Poulin-

Charronat et al. 2005; Tillmann et al. 2006; Hoch et al. 2011; Perruchet and Poulin-Charronnat

2013) or low-level sensory effects (e.g., Bigand et al. 2014; Collins et al. 2014; cf. Koelsch et al.

2007). In order to argue that a brain region that shows a *structure-violation > no violation* effect

supports structure processing in music, one would need to establish that this brain region i) is

selective for structural violations and does not respond to unexpected non-structural (but

similarly salient) events in music or other domains, and ii) responds to music stimuli even when

no violation is present. This latter point is (surprisingly) not often discussed but is deeply

important: if a brain region supports the processing of music structure, it should be engaged

whenever music is processed (similar to how language areas respond robustly to well-formed

sentences, in addition to showing sensitivity to violated linguistic expectations; e.g., Fedorenko

et al. 2020). After all, in order to detect a structural violation, a brain region needs to process the

structure of the preceding context, which implies that it should be working whenever a music

stimulus is present. No previous study has established both of the properties above—selectivity

for structural relative to non-structural violations and robust responses to music stimuli with no

violations—for the brain regions that have been argued to support structure processing in music

(and to overlap with regions that support structure processing in language). In fact, some studies

that have compared unexpected structural and non-structural events in music (e.g., a timbre

change) have reported similar neural responses in fMRI (e.g., Koelsch et al. 2002; cf. some

differences in EEG effects – e.g., Koelsch et al. 2001). Relatedly, and in support of the idea that

effects of music structure violations largely reflect domain-general attentional effects, meta-

analyses of neural responses to unexpected events across domains (e.g., Corbetta and Shulman

2002; Fouragnan et al. 2018; Corlett et al. 2021) have identified regions that grossly resemble

those reported in studies of music structure violations (see Fedorenko and Varley 2016 for

discussion).

Third, most prior fMRI (and MEG) investigations have relied on comparisons of group-level

activation maps. Such analyses suffer from low functional resolution (e.g., Nieto-Castañón and

Fedorenko 2012; Fedorenko 2021), especially in cases where the precise locations of functional

regions vary across individuals, as in the association cortex (Fischl et al. 2008; Frost and Goebel

2012; Tahmasebi et al. 2012; Vazquez-Rodriguez et al. 2019). Thus, observing activation

overlap at the group level does not unequivocally support shared mechanisms. Indeed, studies

that have used individual-subject-level analyses have reported a low or no response to music in

the language-responsive regions (Fedorenko et al. 2011; Rogalsky et al. 2011; Deen et al. 2015).

Fourth, the interpretation of some of the observed effects has relied on the so-called 'reverse

inference' (Poldrack 2006, 2011; Fedorenko 2021), where function is inferred from a coarse

anatomical location: for example, some music-structure-related effects observed in or around

'Broca's area' have been interpreted as reflecting the engagement of linguistic-structure-

processing mechanisms (e.g., Maess et al. 2001; Koelsch et al. 2002) given the long-standing

association between 'Broca's area' and language, including syntactic processing specifically

(e.g., Caramazza and Zurif 1976; Friederici et al. 2006). However, this reasoning is not valid:

Broca's area is a heterogeneous region, which houses components of at least two functionally

distinct brain networks (Fedorenko et al. 2012; Fedorenko and Blank 2020): the language-

selective network, which responds during language processing, visual or auditory, but does not

respond to diverse non-linguistic stimuli (Fedorenko et al. 2011; Monti et al. 2009, 2012; see

Fedorenko and Varley 2016 for a review) and the domain-general executive control or 'multiple

demand (MD)' network, which responds to any demanding cognitive task and is robustly

modulated by task difficulty (Duncan 2010, 2013; Fedorenko et al. 2013; Assem et al. 2020). As

a result, here and more generally, functional interpretation based on coarse anatomical

localization is not justified.

Fifth, many prior fMRI investigations have not reported the magnitudes of response to the

relevant conditions and only examined statistical significance maps for the contrast of interest

(e.g., a whole brain map showing voxels that respond reliably more strongly to melodies with vs.

without a structural violation, and to sentences with vs. without a structural violation). Response

magnitudes of experimental conditions relative to a low-level baseline and to each other are

critical for interpreting a functional profile of a brain region (see e.g., Chen et al. 2017, for

discussion). For example, a reliable *violation > no violation* effect in music (similar arguments

apply to language) could be observed when both conditions elicit above-baseline responses, and

the violation condition elicits a stronger response (**Figure 1A** left bar graph)—a reasonable

profile for a brain region that supports music processing and is sensitive to the target structural

manipulation. However, a reliable *violation > no violation* effect could also be observed when

both conditions elicit below-baseline responses, and the violation condition elicits a less negative

response (**Figure 1A** middle bar graph), or when both conditions elicit low responses—in the

presence of a strong response to stimuli in other domains—and the between-condition difference

is small (**Figure 1A** right bar graph; note that with sufficient power even very small effects can

be highly reliable, but this does not make them theoretically meaningful; e.g., Cumming 2012;

Sullivan and Feinn, 2012). The two latter profiles, where a brain region is more active during

silence than when listening to music, or when the response is overall low and the effect of

interest is minuscule, would be harder to reconcile with a role of this brain region in music

processing (see also the second point above).

Similarly, with respect to the music-language overlap question, a reliable *violation > no*

violation effect for both language and music could be observed in a brain region where sentences

and melodies with violations elicit similarly strong responses, and those without violations elicit

lower responses (**Figure 1B** left bar graph); but it could also arise in a brain region where

sentences with violations elicit a strong response, sentences without violations elicit a lower

response, but melodies elicit an overall low response, with the violation condition eliciting a

higher response than the no-violation condition (**Figure 1B** right bar graph). Whereas in the first

case, it may be reasonable to argue that the brain region in question supports some computation

that is necessary to process structure violations in both domains, such interpretation would not be

straightforward in the second case. In particular, given the large main effect of language > music,

any account of possible computations supported by such a brain region would need to explain

this difference instead of simply focusing on the presence of a reliable effect of violation in both

domains. In summary, without examining the magnitudes of response, it is not possible to

distinguish among many, potentially very different, functional profiles, without which

formulating hypotheses about a brain region's computations is precarious.

Figure 1: Illustration of the importance of examining the magnitudes of neural response to the

experimental conditions rather than only the statistical significance maps for the contrast(s) of interest. A

significant *violation > no violation* effect (A) and overlap between a significant *violation > no violation*

effect in language vs. in music (B) are each compatible with multiple distinct functional profiles, only one

of which (on the left in each case) supports the typically proposed interpretation (a region that processes

structure in some domain of interest in A, and a region that processes structure across domains, in both

language and music, in B).

Aside from the limitations above, to the best of our knowledge, all prior brain imaging studies

have used a single manipulation in one set of materials and one set of participants. To

compellingly argue that a brain region supports (some aspects of) structural processing in both

language and music, it is important to establish both the *robustness* of the key effect by

replicating it with a new set of experimental materials and/or in a new group of participants, and

its *generalizability* to other contrasts between conditions that engage the hypothesized

computation and ones that do not. For example, to argue that a brain region houses a core

syntactic mechanism needed to process hierarchical relations and/or recursion in both language

and music (e.g., Patel 2003; Fadiga et al. 2009; Roberts 2012; Koelsch et al. 2013; Fitch and

Martins 2014), one would need to demonstrate that this region i) responds robustly to diverse

structured linguistic and musical stimuli (which all invoke the hypothesized shared computation),

ii) shows replicable responses across materials and participants, and iii) is sensitive to more than

a single manipulation targeting the hypothesized computations specifically, as needed to rule out

paradigm-/task-specific accounts (e.g., structured vs. unstructured stimuli, stimuli with vs.

without structural violations, stimuli that are more vs. less structurally complex—e.g., with long-

distance vs. local dependencies, adaptation to structure vs. some other aspect of the stimulus,

etc.).

Finally, the neuropsychological patient evidence is at odds with the idea of shared mechanisms

for processing language and music. If language and music relied on the same syntactic

processing mechanism, individuals impaired in their processing of linguistic syntax should also

exhibit impairments in musical syntax. Although some prior studies report subtle musical deficits

in patients with aphasia (Patel et al. 2008a; Sammler et al. 2011), the evidence is equivocal, and

many aphasic patients appear to have little or no difficulties with music, including the processing

of music structure (Luria et al. 1965; Brust 1980; Marin 1982; Basso and Capitani 1985; Polk

and Kertesz 1993; Slevc et al. 2016; Faroqi-Shah et al. 2020; Chiapetta et al. 2022; cf. Omigie

and Samson 2014 and Sihvonen et al. 2017 for discussions of evidence that musical training may

lead to better outcomes following brain damage/resection). Similarly, children with Specific

Language Impairment (now called Developmental Language Disorder)—a developmental

disorder that affects several aspects of linguistic and cognitive processing, including syntactic

processing (e.g., Bortolini et al. 1998; Bishop and Norbury 2002)—show no impairments in

musical processing (Fancourt 2013; cf. Jentschke et al. 2008). In an attempt to reconcile the

evidence from acquired and developmental disorders with claims about structure-processing

overlap based on behavioral and neural evidence from neurotypical participants, Patel (2003,

2008, 2012; see Slevc and Okada 2015, Patel and Morgan 2017, and Asano et al. 2021 for

related proposals) put forward a hypothesis whereby the representations that mediate language

and music are stored in distinct brain areas, but the mechanisms that perform online

computations on those representations are partially overlapping. We return to this idea in the

Discussion.

To bring clarity to this ongoing debate, we conducted three fMRI experiments with neurotypical

adults, and a behavioral study with individuals with severe aphasia. For the fMRI experiments,

we took an approach where we focused on the 'language network'—a well-characterized set of

left frontal and temporal brain areas that selectively support linguistic processing (e.g.,

Fedorenko et al. 2011) and asked whether any parts of this network show responses to music and

sensitivity to music structure. In each experiment, we used an extensively validated language

'localizer' task based on the reading of sentences and nonword sequences (Fedorenko et al.

2010; see Scott et al. 2017 and Malik-Moraleda, Ayyash et al. 2022 for evidence that this

localizer is modality-independent) to identify language-responsive areas in each participant

individually. Importantly, these areas have been shown, across dozens of brain imaging studies,

to be robustly sensitive to linguistic syntactic processing demands in diverse manipulations (e.g.,

Keller et al. 2001; Röder et al. 2002; Friederici 2011; Pallier et al. 2011; Bautista and Wilson

2016, among many others)—including when defined with the same localizer as the one used here

(e.g., Fedorenko et al. 2010, 2012a, 2020; Blank et al. 2016; Mollica et al. 2020; Shain, Blank et

al. 2020; Shain et al. 2022)—and their damage leads to linguistic, including syntactic, deficits

(e.g., Caplan et al. 1996; Dick et al. 2001; Wilson and Saygin 2004; Tyler et al. 2011; Wilson et

al. 2012; Mesulam et al. 2014; Ding et al. 2020; Matchin and Hickok 2020, among many others).

To address the critical research question, we examined the responses of these language areas to

music, and their necessity for processing music structure. In Experiment 1, we included several

types of music stimuli including orchestral music, single-instrument music, synthetic drum

music, and synthetic melodies, a minimal comparison between songs and spoken lyrics, and a set

of non-music auditory control conditions. We additionally examined sensitivity to structure in

music across two structure-scrambling manipulations. In Experiment 2, we further probed

sensitivity to structure in music using the most common manipulation, contrasting responses to

well-formed melodies vs. melodies containing a note that does not obey the constraints of

Western tonal music. And in Experiment 3, we examined the ability to discriminate between

well-formed melodies and melodies containing a structural violation in three profoundly aphasic

individuals across two tasks. Finally, in Experiment 4, we examined the responses of the

language regions to yet another set of music stimuli in a new set of participants. Further, the

participants were all native speakers of Mandarin, a tonal language, which allowed us to evaluate

the hypothesis that language regions may play a greater role in music processing in individuals

with higher sensitivity to linguistic pitch (e.g., Deutsch et al. 2006, 2009; Bidelman et al. 2011;

Creel et al. 2018; Ngo et al. 2016; Liu et al. 2021).

Materials and methods

Participants

<u>Experiments 1, 2, and 4 (fMRI)</u>:

48 individuals (age 18-51, mean 24.3; 28 female, 20 male) from the Cambridge/Boston, MA

community participated for payment across three fMRI experiments (n=18 in Experiment 1;

n=20 in Experiment 2; n=18 in Experiment 4; 8 participants overlapped between Experiments 1

and 2). 33 participants were right-handed and four left-handed, as determined by the Edinburgh

handedness inventory (Oldfield 1971), or self-report (see Willems et al. 2014, for arguments for

including left-handers in cognitive neuroscience research); the handedness data for the remaining

11 participants (one in Experiment 2 and 10 in Experiment 4) were not collected. All but one

participant (with no handedness information) in Experiment 4 showed typical left-lateralized

language activations in the language localizer task described below (as assessed by numbers of

voxels falling within the language parcels in the left vs. right hemisphere (LH vs. RH), using the

following formula: (LH-RH)/(LH+RH); e.g., Jouravlev et al. 2020; individuals with values of

0.25 or greater were considered to have a left-lateralized language system). For the participant

with right-lateralized language activations (with a lateralization value at or below -0.25), we used

right-hemisphere language regions for the analyses (see **SI-3** for analyses where the LH language

regions were used for this participant and when this participant is excluded; the critical results

were not affected). Participants in Experiments 1 and 2 were native English speakers;

participants in Experiment 4 were native Mandarin speakers and proficient speakers of English

(none had any knowledge of Russian, which was used in the unfamiliar foreign-language

condition in Experiment 4). Detailed information on the participants' music background was,

unfortunately, not collected, except for ensuring that the participants were not professional

musicians. All participants gave informed written consent in accordance with the requirements of

MIT's Committee on the Use of Humans as Experimental Subjects (COUHES).

Experiment 3 (behavioral):

Individuals with aphasia. Three participants with severe and chronic aphasia were recruited to

the study (SA, PR, and PP). All participants gave informed consent in accordance with the

requirements of UCL's Institutional Review Board. Background information on each participant

is presented in **Table 1**. Anatomical scans are shown in **Figure 2A** and extensive perisylvian

damage in the left hemisphere, encompassing areas where language activity is observed in

neurotypical individuals, is illustrated in **Figure 2B**.

Patient	Sex	Age (years)	Time post-onset	Handedness	Etiology	Premorbid musical experience	Premorbid employment

		at testing	(years) at testing				
SA	M	67	21	R	Subdural empyema	Sang in choir; basic sight-reading ability. No formal training.	Police sergeant
PR	M	68	14	L	Left hemisphere stroke	Drummer in band; basic sight-reading ability. No formal training.	Retail manager
PP	M	77	10	R	Left hemisphere stroke	Childhood musical training (5 years). No adult experience.	Minerals trader

288   **Table 1.** Background information on the participants with aphasia.

A. Anatomical scans

SA   PR   PP

right   left

B. Language network overlay

PR

PP

C. Language tasks

Comprehension of spoken
reversible sentences

PR   SA PP

0.4   0.6   0.8   1.0
Proportion correct out of 80

Spoken grammaticality
judgement

SA PR PP

0.4   0.6   0.8   1.0
Proportion correct out of 48

Figure 2: **A.** Anatomical scans (T2-weighted for SA, T1-weighted for PR and PP) of the aphasic

participants (all scans were performed during the chronic phase, as can be seen from the ventricular

enlargement). Note that the right side of the image represents the left side of the brain. **B.** P.R.'s

(top) and P.P.'s (bottom) anatomical scans (blue-tinted) shown with the probabilistic activation

overlap map for the fronto-temporal language network overlaid (SA's raw anatomical data were

not available). The map was created by overlaying thresholded individual activation maps (red-

tinted) for the *sentences > nonwords* contrast (Fedorenko et al. 2010) in 220 neurotypical

participants (none of whom were participants in any experiments in the current study). As the images show, the language network falls largely within the lesioned tissue in the left hemisphere. **C.** Performance of the control participants and participants with aphasia on two measures of linguistic syntax processing (see Design, materials, and procedure – Experiment 3): the comprehension of spoken reversible sentences (top), and the spoken grammaticality judgments (bottom). The densities show the distribution of proportion correct scores in the control participants and the boxplot shows the quartiles of the control population (the whiskers show 1.5x interquartile range and points represent outliers). The dots show individual participants (for the individuals with aphasia, the initials indicate the specific participant). Dashed grey lines indicate chance performance.

Control participants. We used Amazon.com's Mechanical Turk platform to recruit normative samples for the music tasks and a subset of the language tasks that are most critical to linguistic syntactic comprehension. Ample evidence now shows that online experiments yield data that closely mirror the data patterns in experiments conducted in a lab setting (e.g., Crump et al. 2013). Data from participants with IP addresses in the US who self-reported being native English speakers were included in the analyses. 50 participants performed the critical music task, and the Scale task from the MBEA (Peretz et al. 2003), as detailed below. Data from participants who responded incorrectly to the catch trial in the MBEA Scale task (n=5) were excluded from the analyses, for a final sample of 45 control participants for the music tasks. A separate sample of 50 participants performed the *Comprehension of spoken reversible sentences* task. Data from one participant who completed fewer than 75% of the questions and another participant who did not report being a native English speaker were excluded for a final sample of 48 control participants.

Finally, a third sample of 50 participants performed the *Spoken grammaticality judgment* task.

Data from one participant who did not report being a native English speaker were excluded for a

final sample of 49 control participants.

Design, materials, and procedure

<u>Experiments 1, 2, and 4 (fMRI)</u>:

Each participant completed a language localizer task (Fedorenko et al. 2010) and one or more of

the critical music perception experiments, along with one or more tasks for unrelated studies.

The scanning sessions lasted approximately two hours.

Language localizer. This task is described in detail in Fedorenko et al. (2010) and subsequent

studies from the Fedorenko lab (e.g., Fedorenko et al. 2011; Blank et al. 2014; Blank et al. 2016;

Pritchett et al. 2018; Paunov et al. 2019; Fedorenko et al. 2020; Shain, Blank et al. 2020, among

others) and is available for download from https://evlab.mit.edu/funcloc/). Briefly, participants

read sentences and lists of unconnected, pronounceable nonwords in a blocked design. Stimuli

were presented one word/nonword at a time at the rate of 450ms per word/nonword. Participants

read the materials passively and performed a simple button-press task at the end of each trial

(included in order to help participants remain alert). Each participant completed two ~6-minute

runs. This localizer task has been extensively validated and shown to be robust to changes in the

materials, modality of presentation (visual vs. auditory), and task (e.g., Fedorenko et al. 2010;

Fedorenko 2014; Scott et al. 2017; Diachek, Blank, Siegelman et al. 2020; Malik-Moraleda,

Ayyash et al. 2022; Lipkin et al. 2022; see the results of Experiments 1 and 4 for additional

replications of modality robustness). Further, a network that corresponds closely to the localizer

contrast (*sentences > nonwords*) emerges robustly from whole-brain task-free data—voxel

fluctuations during rest (e.g., Braga et al. 2020; see Braga 2021 for a general discussion of how

well-validated localizers tend to show tight correspondence with intrinsic networks recovered in

a data-driven way). The fact that different regions of the language network show strong

correlations in their activity during naturalistic cognition (see also Blank et al. 2014; Paunov et

al. 2019; Malik-Moraleda, Ayyash et al. 2022) provides support for the idea that this network

constitutes a 'natural kind' in the brain (a subset of the brain that is strongly internally integrated

and robustly dissociable from the rest of the brain) and thus a meaningful unit of analysis.

However, we also examine individual regions of this network, to paint a more complete picture,

given that many past claims about language-music overlap have concerned the inferior frontal

component of the language network.

Experiment 1. Participants passively listened to diverse stimuli across 18 conditions in a long-

event-related design. The materials for this and all other experiments are available at OSF:

https://osf.io/68y7c/. All stimuli were 9 s in length. The conditions were selected to probe

responses to music, to examine sensitivity to structure scrambling in music, to compare

responses to songs vs. spoken lyrics, and to compare responses to music stimuli vs. other

auditory stimuli.

The four non-vocal music conditions (all Western tonal music) included orchestral music, single-

instrument music, synthetic drum music, and synthetic melodies (see **SI-5** for a summary of the

acoustic properties of these and other conditions, as quantified with the MIR toolbox; Lartillot

and Toiviainen 2007; Lartillot and Grandjean 2019). The orchestral music condition consisted of

12 stimuli (**SI-Table 4a**) selected from classical orchestras or jazz bands. The single-instrument

music condition consisted of 12 stimuli (**SI-Table 4b**) that were played on one of the following

instruments: cello (n=1), flute (n=1), guitar (n=4), piano (n=4), sax (n=1), or violin (n=1). The

synthetic drum music condition consisted of 12 stimuli synthesized using percussion patches

from MIDI files taken from freely available online collections. The stimuli were synthesized

using the MIDI toolbox for MATLAB (writemidi). The synthetic melodies condition consisted

of 12 stimuli transcribed from folk tunes obtained from freely available online collections. Each

melody was defined by a sequence of notes with corresponding pitches and durations. Each note

was composed of harmonics 1 through 10 of the fundamental presented in equal amplitude, with

no gap in-between notes. Phase discontinuities between notes were avoided by ensuring that the

starting phase of the next note was equal to the ending phase of the previous note.

The synthetic drum music and the synthetic melodies conditions had scrambled counterparts to

probe sensitivity to music structure. This intact > scrambled contrast has been used in some past

studies of structure processing in music (e.g., Levitin and Menon 2003) and is parallel to the

sentences > word-list contrast in language, which has been often used to probe sensitivity for

combinatorial processing (e.g., Fedorenko et al. 2010). The scrambled drum music condition was

created by jittering the inter-note-interval (INI). The amount of jitter was sampled from a

uniform distribution (from -0.5 to 0.5 beats). The scrambled INIs were truncated to be no smaller

than 5% of the distribution of INIs from the intact drum track. The total distribution of INIs was

then scaled up or down to ensure that the total duration remained unchanged. The scrambled

melodies condition was created by scrambling both pitch and rhythm information. Pitch

information was scrambled by randomly re-ordering the sequence of pitches and then adding

jitter to disrupt the key. The amount of jitter for each note was sampled from a uniform

distribution centered on the note's pitch after shuffling (from -3 to +3 semitones). The duration of

each note was also jittered (from -0.2 to 0.2 beats). To ensure the total duration was unaffected

by jitter, N/2 positive jitter values were sampled, where N is the number of notes, and then a

negative jitter was added with the same magnitude for each of the positive samples, such that the

sum of all jitters equaled 0. To ensure the duration of each note remained positive, the smallest

jitters were added to the notes with the smallest durations. Specifically, the note durations and

sampled jitters were sorted by their magnitude, summed, and then the jittered durations were

randomly re-ordered.

To allow for a direct comparison between music and linguistic conditions within the same

experiment, we included auditory sentences and auditory nonword sequences. The sentence

condition consisted of 24 lab-constructed stimuli (half recorded by a male, and half by a female).

Each stimulus consisted of a short story (each three sentences long) describing common,

everyday events. Any given participant heard 12 of the stimuli (6 male, 6 female). The nonword

sequence condition consisted of 12 stimuli (recorded by a male).

We also included two other linguistic conditions: songs and spoken lyrics. These conditions were

included to test whether the addition of a melodic contour to speech (in songs) would increase

the responses of the language regions. Such a pattern might be expected of a brain region that

responds to both linguistic content and music structure. The songs and the lyrics conditions each

consisted of 24 stimuli. We selected songs with a tune that was easy to sing without

accompaniment. These materials were recorded by four male singers: each recorded between 2

and 11 song-lyrics pairs. The singers were actively performing musicians (e.g., in *a cappella*

groups) but were not professionals. Any given participant heard either the song or the lyrics

version of an item for 12 stimuli in each condition.

Finally, to assess the specificity of potential responses to music, we included three non-music

conditions: animal sounds and two kinds of environmental sounds (pitched and unpitched),

which all share some low-level acoustic properties with music (see SI-5). The animal sounds

condition and the environmental sounds conditions each consisted of 12 stimuli taken from in-

lab collections. If individual recordings were shorter than 9 s, then several recordings of the same

type of sound were concatenated together (100 ms gap in between). We included the pitch

manipulation in order to test for general responsiveness to pitch—a key component of music—in

the language regions.

(The remaining five conditions were not directly relevant to the current study or redundant with

other conditions for our research questions and therefore not included in the analyses. These

included three distorted speech conditions—lowpass-filtered speech, speech with a flattened

pitch contour, and lowpass-filtered speech with a flattened pitch contour—and two additional

low-level controls for the synthetic melody stimuli. The speech conditions were included to

probe sensitivity to linguistic prosody for an unrelated study. The additional synthetic music

control conditions were included to allow for a more rigorous interpretation of the intact >

scrambled synthetic melodies effect had we observed such an effect. For completeness, on the

OSF page, https://osf.io/68y7c/, we provide a data table that includes responses to these five conditions.)

For each participant, stimuli were randomly divided into six sets (corresponding to runs) with each set containing two stimuli from each condition. The order of the conditions for each run was selected from four predefined palindromic orders, which were constructed so that conditions targeting similar mental processes (e.g., orchestral music and single-instrument music) were separated by other conditions (e.g., speech or animal sounds). Each run contained three 10 s fixation periods: at the beginning, in the middle, and at the end. Otherwise, the stimuli were separated by 3 s fixation periods, for a total run duration of 456 s (7 min 36 s). All but two of the 18 participants completed all six runs (and thus got a total of 12 experimental events per condition); the remaining two completed four runs (and thus got 8 events per condition).

Because, as noted above, we have previously established that the language localizer is robust to presentation modality, we used the visual localizer to define the language regions. However, in SI-2 we show that the critical results are similar when auditory contrasts (*sentences > nonwords* in Experiment 1, or *Mandarin sentences > foreign* in Experiment 4) are instead used to define the language regions.

Experiment 2. Participants listened to well-formed melodies (adapted and expanded from Fedorenko et al. 2009) and melodies with a structural violation in a long-event-related design and judged the well-formedness of the melodies. As discussed in the Introduction, this type of manipulation is commonly used to probe sensitivity to music structure, including in studies

examining language-music overlap (e.g., Patel et al. 1998; Koelsch et al. 2000, 2002; Maess et al. 2001; Tillmann et al. 2003; Fedorenko et al. 2009; Slevc et al. 2009; Kunert et al. 2015; Musso et al. 2015). The melodies were between 11 and 14 notes. The well-formed condition consisted of 90 melodies, which were tonal and ended in a tonic note with an authentic cadence in the implied harmony. All melodies were isochronous, consisting of quarter notes except for the final half note. The first five notes established a strong sense of key. Each melody was then altered to create a version with a "sour" note: the pitch of one note (from among the last four notes in a melody) was altered up or down by one or two semitones, so as to result in a non-diatonic note while keeping the melodic contour (the up-down pattern) the same. The structural position of the note that underwent this change varied among the tonic, the fifth, and the major third. The full set of 180 melodies was distributed across two lists following a Latin Square design. Any given participant heard stimuli from one list.

For each participant, stimuli were randomly divided into two sets (corresponding to runs) with each set containing 45 melodies (22 or 23 per condition). The order of the conditions, and the distribution of inter-trial fixation periods, was determined by the optseq2 algorithm (Dale et al. 1999). The order was selected from among four predefined orders, with no more than four trials of the same condition in a row. In each trial, participants were presented with a melody for three seconds followed by a question, presented visually on the screen, about the well-formedness of the melody ("Is the melody well-formed?"). To respond, participants had to press one of two buttons on a button box within two seconds. When participants answered, the question was replaced by a blank screen for the remainder of the two-second window; if no response was made within the two-second window, the experiment advanced to the next trial. Responses received within one

second after the end of the previous trial were still recorded to account for the possible slow

responses. The screen was blank during the presentation of the melodies. Each run contained 151

s of fixation interleaved among the trials, for a total run duration of 376 s (6 min 16 s). Fourteen

of the 20 participants completed both runs, four participants completed one run, and the two

remaining participants completed two runs but we only included their first run because, due to

experimenter error, the second run came from a different experimental list and thus included some

of the melodies from the first run in the other condition (the data pattern was qualitatively and

quantitatively the same if both runs were included for these participants). Finally, due to a script

error, participants only heard the first 12 notes of each melody during the three seconds of stimulus

presentation. Therefore, we only analyzed the 80 of the 90 pairs (160 of the 180 total melodies)

where the contrastive note appeared within the first 12 notes.

Experiment 4. Participants passively listened to single-instrument music, environmental sounds,

sentences in their native language (Mandarin), and sentences in an unfamiliar foreign language

(Russian) in a blocked design. All stimuli were 5-5.95s in length. The conditions were selected to

probe responses to music, and to compare responses to music stimuli vs. other auditory stimuli.

The critical music condition consisted of 60 stimuli selected from classical pieces by J.S. Bach

played on cello, flute, or violin (n=15 each) and jazz music played on saxophone (n=15). The

environmental sounds condition consisted of 60 stimuli selected from in-lab collections and

included both pitched and unpitched stimuli. The foreign language condition consisted of 60

stimuli selected from Russian audiobooks (short stories by Paustovsky and "Fathers and Sons" by

Turgenev). The foreign language condition was included because creating a 'nonwords' condition

(the baseline condition we typically use for defining the language regions; Fedorenko et al. 2010)

is challenging in Mandarin given that most words are monosyllabic, thus most syllables carry some

meaning. As a result, sequences of syllables are more akin to lists of words. Therefore, we included

the unfamiliar foreign language condition, which also works well as a baseline for language

processing (Malik-Moraleda, Ayyash et al. 2022). The Mandarin sentence condition consisted of

120 lab-constructed sentences, each recorded by a male and a female native speaker. (The

experiment also included five conditions that were not relevant to the current study and therefore

not included in the analyses. These included three conditions probing responses to the participants'

second language (English) and two control conditions for Mandarin sentences. For completeness,

on the OSF page, https://osf.io/68y7c/, we provide a data table that includes responses to these five

conditions.)

Stimuli were grouped into blocks with each block consisting of three stimuli and lasting 18s

(stimuli were padded with silence to make each trial exactly six seconds long). For each participant,

blocks were divided into 10 sets (corresponding to runs), with each set containing two blocks from

each condition. The order of the conditions for each run was selected from eight predefined

palindromic orders. Each run contained three 14 s fixation periods: at the beginning, in the middle,

and at the end, for a total run duration of 366 s (6 min 6 s). Five participants completed eight of

the 10 runs (and thus got 16 blocks per condition; the remaining thirteen completed six runs (and

thus got 12 blocks per condition). (We had created enough materials for 10 runs, but based on

observing robust effects for several key contrasts in the first few participants who completed six

to eight runs, we administered 6-8 runs to the remaining participants.)

Because we have previously found that an English localizer works well in native speakers of

diverse languages, including Mandarin, as long as they are proficient in English (Malik-Moraleda,

Ayyash et al. 2022), we used the same localizer in Experiment 4 as the one used in Experiments 1

and 2, for consistency. However, in SI-2 (**SI-Figure 2c**, **SI-Table 2c**) we show that the critical

results are similar when the *Mandarin sentences > foreign* contrast is instead used to define the

language regions.

<u>Experiment 3 (behavioral):</u>

Language assessments. Participants with aphasia were assessed for the integrity of lexical

processing using word-to-picture matching tasks in both spoken and written modalities (ADA

Spoken and Written Word-Picture Matching; Franklin et al. 1992). Productive vocabulary was

assessed through picture naming. In the spoken modality, the Boston Naming Test was employed

(Kaplan et al. 2001), and in writing, the PALPA Written Picture Naming subtest (Kay et al.

1992). Sentence processing was evaluated in both spoken and written modalities through

comprehension (sentence-to-picture matching) of reversible sentences in active and passive

voice. In a reversible sentence, the heads of both noun phrases are plausible agents, and

therefore, word order, function words, and functional morphology are the only cues to who is

doing what to whom. Participants also completed spoken and written grammaticality judgment

tasks, where they made a yes/no decision as to the grammaticality of a word string. The task

employed a subset of sentences from Linebarger et al. (1983).

All three participants exhibited severe language impairments that disrupted both comprehension

and production (**Table 2**). For lexical-semantic tasks, all three participants displayed residual

comprehension ability for high imageability/picturable vocabulary, although more difficulty was

evident on the synonym matching test, which included abstract words. They were all severely

anomic in speech and writing. Sentence production was severely impaired with output limited to

single words, social speech (expressions, like "How are you?"), and other formulaic expressions

(e.g., "and so forth"). Critically, all three performed at or close to chance level on spoken and

written comprehension of reversible sentences and grammaticality judgments; each patient's

scores were lower than all of the healthy controls (**Table 2** and **Figure 2C**).

Participant	SA	PR	PP	Controls
Lexical-semantic assessments				
ADA Spoken Word-Picture Matching (chance = 16.5)	60/66	61/66	64/66	N/A
ADA Written Word-Picture Matching (chance = 16.5)	62/66	66/66	58/66	N/A
ADA spoken synonym matching (chance = 80)	123/160	121/160	135/160	N/A
ADA written synonym matching (chance = 80)	121/160	145/160	143/160	N/A
Boston Naming Test (NB: accepting both spoken and written responses)	4/60	4/60	11/60	N/A

PALPA 54 Written Picture Naming	24/60	2/60	1/60	N/A
Syntactic assessments				
Comprehension of spoken reversible sentences (chance = 40)	49/80	38/80	52/80	Mean = 79.5/80 SD = 1.03 Min = 74/80 Max = 80/80 N=48
Comprehension of written reversible sentences (chance = 40)	42/80	49/80	51/80	N/A
Spoken grammaticality judgments (chance = 24)	33/48	34/48	35/48	Mean = 45.5/48 SD = 2.52 Min = 36/48 Max = 48/48 N=49
Written grammaticality judgments (chance = 24)	29/48	24/48	29/48	N/A

Table 2. Results of language assessments for participants with aphasia and healthy controls. For

each test, we show number of correctly answered questions out of the total number of questions.

Critical music task. Participants judged the well-formedness of the melodies from Experiment 2.

Judgments were intended to reflect the detection of the key violation in the sour versions of the

melodies. The full set of 180 melodies was distributed across two lists following a Latin Square

design. All participants heard all 180 melodies. The control participants heard the melodies from

one list, followed by the melodies from the other list, with the order of lists counter-balanced

across participants. For the participants with aphasia, each list was further divided in half, and

each participant was tested across four sessions, with 45 melodies per session, to minimize

fatigue.

Montreal Battery for the Evaluation of Amusia. To obtain another measure of music

competence/sensitivity to music structure, we administered the Montreal Battery for the

Evaluation of Amusia (MBEA) (Peretz et al. 2003). The battery consists of six tasks that assess

musical processing components described by Peretz and Coltheart (2003): three target melodic

processing, two target rhythmic processing, and one assesses memory for melodies. Each task

consists of 30 experimental trials (and uses the same set of 30 base melodies) and is preceded by

practice examples. Some of the tasks additionally include a catch trial, as described below. For

the purposes of the current investigation, the critical task is the "Scale" task. Participants are

presented with pairs of melodies that they have to judge as identical or not. On half of the trials,

one of the melodies is altered by modifying the pitch of one of the tones to be out of scale. Like

our critical music task, this task aims to test participants' ability to represent and use tonal

structure in Western music, except that instead of making judgments on each individual melody,

participants compare two melodies on each trial. This task thus serves as a conceptual replication

(Schmidt 2009). One trial contains stimuli designed to be easy, intended as a catch trial to ensure

that participants are paying attention. In this trial, the comparison melody has all its pitches set at

random. This trial is excluded when computing the scores.

Control participants performed just the Scale task. Participants with aphasia performed all six

tasks, distributed across three testing sessions to minimize fatigue.

fMRI data acquisition, preprocessing, and first-level modeling (for Experiments 1, 2, and 4)

Data acquisition. Whole-brain structural and functional data were collected on a whole-body 3

Tesla Siemens Trio scanner with a 32-channel head coil at the Athinoula A. Martinos Imaging

Center at the McGovern Institute for Brain Research at MIT. T1-weighted structural images were

collected in 176 axial slices with 1 mm isotropic voxels (repetition time (TR) = 2,530 ms; echo

time (TE) = 3.48 ms). Functional, blood oxygenation level-dependent (BOLD) data were

acquired using an EPI sequence with a 90° flip angle and using GRAPPA with an acceleration

factor of 2; the following parameters were used: thirty-one 4.4 mm thick near-axial slices

acquired in an interleaved order (with 10% distance factor), with an in-plane resolution of 2.1

mm × 2.1 mm, FoV in the phase encoding (A >> P) direction 200 mm and matrix size 96 × 96

voxels, TR = 2000 ms and TE = 30 ms. The first 10 s of each run were excluded to allow for

steady state magnetization (see OSF https://osf.io/68y7c/ for the pdf of the scanning protocols).

(Note that we opted to use a regular, continuous, scanning sequence in spite of investigating

responses to auditory conditions. However, effects of scanner noise are unlikely to be

detrimental given that all the stimuli are clearly perceptible, as also confirmed by examining

responses in the auditory areas.)

Preprocessing. fMRI data were analyzed using SPM12 (release 7487), CONN EvLab module

(release 19b), and other custom MATLAB scripts. Each participant's functional and structural

data were converted from DICOM to NIFTI format. All functional scans were coregistered and

resampled using B-spline interpolation to the first scan of the first session (Friston et al. 1995).

Potential outlier scans were identified from the resulting subject-motion estimates as well as

from BOLD signal indicators using default thresholds in CONN preprocessing pipeline (5

standard deviations above the mean in global BOLD signal change, or framewise displacement

values above 0.9 mm; Nieto-Castañón 2020). Functional and structural data were independently

normalized into a common space (the Montreal Neurological Institute [MNI] template;

IXI549Space) using SPM12 unified segmentation and normalization procedure (Ashburner and

Friston 2005) with a reference functional image computed as the mean functional data after

realignment across all timepoints omitting outlier scans. The output data were resampled to a

common bounding box between MNI-space coordinates (-90, -126, -72) and (90, 90, 108), using

2mm isotropic voxels and 4th order spline interpolation for the functional data, and 1mm

isotropic voxels and trilinear interpolation for the structural data. Last, the functional data were

smoothed spatially using spatial convolution with a 4 mm FWHM Gaussian kernel.

First-level modeling. For both the language localizer task and the critical experiments, effects

were estimated using a General Linear Model (GLM) in which each experimental condition was

modeled with a boxcar function convolved with the canonical hemodynamic response function

(HRF) (fixation was modeled implicitly, such that all timepoints that did not correspond to one

of the conditions were assumed to correspond to a fixation period). Temporal autocorrelations in

the BOLD signal timeseries were accounted for by a combination of high-pass filtering with a

128 seconds cutoff, and whitening using an AR(0.2) model (first-order autoregressive model

linearized around the coefficient a=0.2) to approximate the observed covariance of the functional

data in the context of Restricted Maximum Likelihood estimation (ReML). In addition to

experimental condition effects, the GLM design included first-order temporal derivatives for

each condition (included to model variability in the HRF delays), as well as nuisance regressors

to control for the effect of slow linear drifts, subject-motion parameters, and potential outlier

scans on the BOLD signal.

Definition of the language functional regions of interest (fROIs) (for Experiments 1, 2, and 4)

For each critical experiment, we defined a set of language functional regions of interest (fROIs)

using group-constrained, subject-specific localization (Fedorenko et al. 2010). In particular, each

individual map for the *sentences > nonwords* contrast from the language localizer was

intersected with a set of five binary masks. These masks (**Figure 3**; available at OSF:

https://osf.io/68y7c/) were derived from a probabilistic activation overlap map for the same

contrast in a large independent set of participants (n=220) using watershed parcellation, as

described in Fedorenko et al. (2010) for a smaller set of participants. These masks covered the

fronto-temporal language network in the left hemisphere. Within each mask, a participant-

specific language fROI was defined as the top 10% of voxels with the highest *t*-values for the

localizer contrast.

Validation of the language fROIs

To ensure that the language fROIs behave as expected (i.e., show a reliably greater response to

the sentences condition compared to the nonwords condition), we used an across-runs cross-

validation procedure (e.g., Nieto-Castañón and Fedorenko 2012). In this analysis, the first run of

the localizer was used to define the fROIs, and the second run to estimate the responses (in

percent BOLD signal change, PSC) to the localizer conditions, ensuring independence (e.g.,

Kriegeskorte et al. 2009); then the second run was used to define the fROIs, and the first run to

estimate the responses; finally, the extracted magnitudes were averaged across the two runs to

derive a single response magnitude for each of the localizer conditions. Statistical analyses were

performed on these extracted PSC values. Consistent with much previous work (e.g., Fedorenko

et al. 2010; Mahowald and Fedorenko 2016; Diachek, Blank, Siegelman et al. 2020), each of the

language fROIs showed a robust *sentences > nonwords* effect (all ps < 0.001).

Statistical analyses for the fMRI experiments

All analyses were performed with linear mixed-effects models using the "lme4" package in R

with p-value approximation performed by the "lmerTest" package (Bates et al. 2015; Kuznetsova

et al. 2017). Effect size (Cohen's d) was calculated using the method from Westfall et al. (2014)

and Brysbaert and Stevens (2018).

Sanity check analyses and results

To estimate the responses in the language fROIs to the conditions of the critical experiments here

and in the critical analyses, the data from all the runs of the language localizer were used to

define the fROIs, and the responses to each condition were then estimated in these regions.

Statistical analyses were then performed on these extracted PSC values. (For Experiments 1 and

4, we repeated the analyses using alternative language localizer contrasts to define the language

fROIs (auditory *sentences > nonwords* in Experiment 1, and *Mandarin sentences > foreign* in

Experiment 4), which yielded quantitatively and qualitatively similar responses (see **SI-2**).)

We conducted two sets of sanity check analyses. First, to ensure that auditory conditions that

contain meaningful linguistic content elicit strong responses in the language regions relative to

perceptually similar conditions with no discernible linguistic content, we compared the auditory

sentences condition with the auditory nonwords condition (Experiment 1) or with the foreign

language condition (Experiment 4). Indeed, as expected, the auditory sentence condition elicited

a stronger response than the auditory nonwords condition (Experiment 1) or the foreign language

condition (Experiment 4). These effects were robust at the network level (ps < 0.001; **SI-Table**

1a). Further, the *sentences > nonwords* effect was significant in all but one language fROI in

Experiment 1, and the *sentences > foreign* effect was significant in all language fROIs in

Experiment 4 (ps < 0.05; **SI-Table 1a**).

And second, to ensure that the music conditions elicit strong responses in auditory cortex, we

extracted the responses from a bilateral anatomically defined auditory cortical region (area Te1.2

from the Morosan et al. 2001 cytoarchitectonic probabilistic atlas) to the six critical music

conditions: orchestral music, single instrument music, synthetic drum music, and synthetic

melodies in Experiment 1; well-formed melodies in Experiment 2; and the music condition in

Experiment 4. Statistical analyses, comparing each condition to the fixation baseline, were

performed on these extracted PSC values. As expected, all music conditions elicited strong

responses in a primary auditory area bilaterally (all ps ≅ 0.001; **SI-Table 1b**; **SI-Figure 1**).

Critical analyses

To characterize the responses in the language network to music perception, we asked three

questions. First, we asked whether music conditions elicit strong responses in the language

regions. Second, we investigated whether the language network is sensitive to structure in music,

as would be evidenced by stronger responses to intact than scrambled music, and stronger

responses to melodies with structural violations compared to the no-violation control condition.

And third, we asked whether music conditions elicit strong responses in the language regions of

individuals with high sensitivity to linguistic pitch—native speakers of a tonal language

(Mandarin).

For each contrast (the contrasts relevant to the three research questions are detailed below), we

used two types of linear mixed-effect regression models:

i) the language network model, which examined the language network as a whole; and

ii) the individual language fROI models, which examined each language fROI separately.

As alluded to in the Introduction, treating the language network as an integrated system is

reasonable given that the regions of this network a) show similar functional profiles, both with

respect to selectivity for language over non-linguistic processes (e.g., Fedorenko et al. 2011;

Pritchett et al. 2018; Jouravlev et al. 2019; Ivanova et al. 2020, 2021) and with respect to their

role in lexico-semantic and syntactic processing (e.g., Fedorenko et al. 2012b; Blank et al. 2016;

Fedorenko et al. 2020); and b) exhibit strong inter-region correlations in both their activity

during naturalistic cognition paradigms (e.g., Blank et al. 2014; Braga et al. 2020; Paunov et al.

2019; Malik-Moraleda, Ayyash et al. 2022) and key functional markers, like the strength or

extent of activation in response to language stimuli (e.g., Mahowald and Fedorenko 2016;

Mineroff, Blank et al. 2018). However, to allow for the possibility that language regions differ in

their response to music and to examine the region on which most claims about language-music

overlap have focused (the region that falls within 'Broca's area'), we supplement the network-

wise analyses with the analyses of the five language fROIs separately.

For each network-wise analysis, we fit a linear mixed-effect regression model predicting the

level of BOLD response in the language fROIs in the contrasted conditions. The model included

a fixed effect for condition (the relevant contrasts are detailed below for each analysis) and

random intercepts for fROIs and participants. Here and elsewhere, the *p*-value was estimated by

applying the Satterthwaite's method-of-moment approximation to obtain the degrees of freedom

(Giesbrecht and Burns 1985; Fai and Cornelius 1996; as described in Kuznetsova et al. 2017).

For the comparison against the fixation baseline, the random intercept for participants was

removed because it is no longer applicable.

743   *Effect size ~ condition + (1 | fROI) + (1 | Participant)*

For each fROI-wise analysis, we fit a linear mixed-effect regression model predicting the level of

BOLD response in each of the five language fROIs in the contrasted conditions. The model

included a fixed effect for condition and random intercepts for participants. For each analysis,

the results were FDR-corrected for the five fROIs. For the comparison against the fixation

baseline, the random intercept for participants was removed because it is no longer applicable.

Effect size ~ condition + (1 | Participant)

Results

Does music elicit a response in the language network?

As discussed in the Introduction, a brain region that supports (some aspect of) music processing,

including structure processing, should show a strong response to music stimuli. To test whether

language regions respond to music, we used four contrasts using data from Experiments 1 and 2.

First, we compared the responses to each of the music conditions (orchestral music, single

instrument music, synthetic drum music, and synthetic melodies in Experiment 1; well-formed

melodies in Experiment 2) against the fixation baseline—the most liberal baseline. Second, we

compared the responses to the music conditions against the response to the nonword strings

condition—an unstructured and meaningless linguistic stimulus (in Experiment 1, we used the

auditory nonwords condition, and in Experiment 2, we used the visual nonwords condition from

the language localizer). Third, in Experiment 1, we additionally compared the responses to the

music conditions against the response to non-linguistic, non-music stimuli (animal and

environmental sounds). A brain region that supports music processing should elicit a strong

positive response relative to the fixation baseline and the nonwords condition (our baseline for

the language regions); further, if the response is selective, it should be stronger than the response

elicited by non-music auditory stimuli. Finally, in Experiment 1, we also directly compared the

responses to songs vs. lyrics. A brain region that responds to music should respond more

strongly to songs given that they contain a melodic contour in addition to the linguistic content.

None of the music conditions elicited a strong response in the language network (**Figure 3**;

Table 3). The responses to music (i) fell at or below the fixation baseline (except for the well-

formed melodies condition in Experiment 2, which elicited a small positive response in some

regions), (ii) were lower than the response elicited by auditory nonwords (except for the LMFG

language fROI, where the responses to music and nonwords were similarly low), and (iii) did not

significantly differ from the responses elicited by non-linguistic, non-music conditions. Finally,

the response to songs, which contain both linguistic content and a melodic contour, was not

significantly higher than the response elicited by the linguistic content alone (lyrics); in fact, at

the network level, the response to songs was reliably lower than to lyrics.

Figure 3. Responses of the language fROIs (pooling across the network – top, and for each fROI individually – bottom) to the language localizer conditions (in grey), to the four auditory conditions containing speech in Experiment 1 (red shades), to the five music conditions in Experiments 1 and 2 (blue shades), and to the three non-linguistic/non-music auditory conditions (green shades) in Experiment 1. Here and elsewhere, the error bars represent standard errors of the mean by participants. For the language localizer results, we include here all participants in Experiments 1 and 2. The responses to the music

792 conditions cluster around the fixation baseline, are much lower than the responses to sentences, and are

793 not higher than the responses to non-music sounds.

Contrast	Language network	LIFGorb	LIFG	LMFG	LAnt Temp	LPost Temp
music > fixation						
orchestral music >fixation	β=0.028 se=0.135 df=17.383 d=0.042 t=0.209 p=0.837	β=-0.129 se=0.193 df=nan d=nan t=-0.667 p=1.000	β=0.082 se=0.162 df=nan d=nan t=0.506 p=1.000	β=0.117 se=0.165 df=nan d=nan t=0.711 p=1.000	β=0.040 se=0.130 df=nan d=nan t=0.310 p=1.000	β=0.030 se=0.143 df=nan d=nan t=0.210 p=1.000
single-instrument music >fixation	β=-0.294 se=0.163 df=18.616 d=-0.378 t=-1.809 p=0.087	β=-0.552 se=0.223 df=nan d=nan t=-2.471 p=0.120	β=-0.141 se=0.155 df=nan d=nan t=-0.906 p=1.000	β=-0.243 se=0.217 df=nan d=nan t=-1.122 p=1.000	β=-0.272 se=0.152 df=nan d=nan t=-1.794 p=0.455	β=-0.264 se=0.164 df=nan d=nan t=-1.611 p=0.630
synthetic drum music >fixation	β=-0.255 se=0.112 df=18.000 d=-0.432 t=-2.281 p=0.035*	β=-0.258 se=0.155 df=nan d=nan t=-1.667 p=0.570	β=-0.306 se=0.172 df=nan d=nan t=-1.780 p=0.465	β=-0.168 se=0.162 df=nan d=nan t=-1.040 p=1.000	β=-0.319 se=0.106 df=nan d=nan t=-3.020 p=0.040*	β=-0.226 se=0.103 df=nan d=nan t=-2.189 p=0.215
synthetic melodies	β=-0.243 se=0.100	β=-0.286 se=0.154	β=-0.299 se=0.120	β=-0.108 se=0.177	β=-0.247 se=0.103 df=nan	β=-0.276 se=0.089 df=nan

>fixation	df=18.000 d=-0.441 t=-2.423 p=0.026*	df=nan d=nan t=-1.856 p=0.405	df=nan d=nan t=-2.485 p=0.120	df=nan d=nan t=-0.611 p=1.000	d=nan t=-2.395 p=0.140	d=nan t=-3.093 p=0.035*
well-formed melodies (Expt 2) >fixation	β=0.201 se=0.135 df=17.483 d=0.281 t=1.488 p=0.155	β=0.139 se=0.166 df=nan d=nan t=0.836 p=1.000	β=0.396 se=0.181 df=nan d=nan t=2.182 p=0.210	β=0.371 se=0.197 df=nan d=nan t=1.885 p=0.375	β=-0.008 se=0.136 df=nan d=nan t=-0.056 p=1.000	β=0.107 se=0.096 df=nan d=nan t=1.109 p=1.000
music > nonwords						
orchestral music >nonwords	β=-0.746 se=0.092 df=157.707 d=-0.978 t=-8.097 p<0.001***	β=-0.811 se=0.276 df=36.000 d=-0.981 t=-2.945 p=0.030*	β=-0.569 se=0.142 df=18.000 d=-0.779 t=-4.015 p=0.005*	β=-0.210 se=0.221 df=18.000 d=-0.276 t=-0.954 p=1.000	β=-1.187 se=0.147 df=18.000 d=-1.884 t=-8.101 p<0.001***	β=-0.950 se=0.205 df=18.000 d=-1.427 t=-4.646 p<0.001***
single-instrument music >nonwords	β=-1.068 se=0.100 df=157.689 d=-1.314 t=-10.714 p<0.001***	β=-1.234 se=0.296 df=36.000 d=-1.388 t=-4.167 p<0.001***	β=-0.791 se=0.222 df=18.000 d=-1.101 t=-3.567 p=0.010*	β=-0.571 se=0.235 df=18.000 d=-0.661 t=-2.431 p=0.130	β=-1.500 se=0.196 df=18.000 d=-2.236 t=-7.648 p<0.001***	β=-1.244 se=0.234 df=17.998 d=-1.765 t=-5.315 p<0.001***
synthetic drum music >nonwords	β=-1.029 se=0.087 df=157.720 d=-1.408	β=-0.940 se=0.212 df=18.000 d=-1.246	β=-0.956 se=0.182 df=18.000 d=-1.275	β=-0.496 se=0.245 df=18.000 d=-0.658	β=-1.546 se=0.187 df=18.000 d=-2.621	β=-1.207 se=0.177 df=18.000 d=-2.012

	t=-11.839	t=-4.430	t=-5.252	t=-2.026	t=-8.262	t=-6.817
	p<0.001***	p<0.001***	p<0.001***	p=0.290	p<0.001***	p<0.001***
synthetic melodies -nonwords	β=-1.017	β=-0.969	β=-0.949	β=-0.435	β=-1.474	β=-1.256
	se=0.087	se=0.209	se=0.153	se=0.252	se=0.195	se=0.176
	df=157.683	df=18.000	df=18.000	df=18.000	df=36.000	df=18.000
	d=-1.421	d=-1.286	d=-1.441	d=-0.556	d=-2.513	d=-2.164
	t=-11.623	t=-4.642	t=-6.223	t=-1.727	t=-7.541	t=-7.136
	p<0.001***	p<0.001***	p<0.001***	p=0.505	p<0.001***	p<0.001***
well-formed melodies (Expt 2) >nonwords (visual)	β=-0.449	β=-0.490	β=-0.403	β=-0.686	β=-0.242	β=-0.375
	se=0.090	se=0.226	se=0.208	se=0.250	se=0.134	se=0.123
	df=179.063	df=20.989	df=20.056	df=20.173	df=20.737	df=20.455
	d=-0.562	d=-0.611	d=-0.444	d=-0.705	d=-0.470	d=-0.792
	t=-4.998	t=-2.164	t=-1.938	t=-2.748	t=-1.812	t=-3.056
	p<0.001***	p=0.210	p=0.335	p=0.060	p=0.420	p=0.030*

music > non-linguistic, non-music auditory conditions

music (combined) >animal sounds	β=-0.114	β=-0.306	β=-0.295	β=0.080	β=-0.002	β=-0.048
	se=0.060	se=0.148	se=0.146	se=0.151	se=0.090	se=0.094
	df=427.876	df=72.000	df=72.000	df=72.000	df=72.000	df=72.000
	d=-0.177	d=-0.422	d=-0.451	d=0.111	d=-0.004	d=-0.088
	t=-1.915	t=-2.069	t=-2.021	t=0.528	t=-0.023	t=-0.513
	p=0.056	p=0.210	p=0.235	p=1.000	p=1.000	p=1.000
music (combined) >environmental (pitched)	β=0.019	β=0.005	β=-0.104	β=0.055	β=0.092	β=0.045
	se=0.060	se=0.144	se=0.133	se=0.159	se=0.094	se=0.094
	df=427.902	df=72.000	df=72.000	df=72.000	df=72.000	df=72.000
	d=0.028	d=0.006	d=-0.156	d=0.071	d=0.171	d=0.081
	t=0.307	t=0.033	t=-0.781	t=0.347	t=0.975	t=0.475
	p=0.759	p=1.000	p=1.000	p=1.000	p=1.000	p=1.000

music (combined) >environmental (unpitched)	β=-0.052 se=0.063 df=427.856 d=-0.079 t=-0.823 p=0.411	β=-0.109 se=0.163 df=72.000 d=-0.140 t=-0.666 p=1.000	β=-0.118 se=0.152 df=72.000 d=-0.182 t=-0.778 p=1.000	β=-0.030 se=0.151 df=72.000 d=-0.040 t=-0.198 p=1.000	β=0.042 se=0.097 df=72.000 d=0.083 t=0.429 p=1.000	β=-0.043 se=0.100 df=72.000 d=-0.082 t=-0.426 p=1.000
(melodic contour + linguistic content) > linguistic content						
songs >lyrics	β=-0.408 se=0.102 df=157.896 d=-0.377 t=-4.014 p<0.001***	β=-0.705 se=0.287 df=18.000 d=-0.569 t=-2.454 p=0.125	β=-0.394 se=0.195 df=18.000 d=-0.400 t=-2.025 p=0.290	β=-0.243 se=0.219 df=18.000 d=-0.226 t=-1.107 p=1.000	β=-0.313 se=0.163 df=18.000 d=-0.356 t=-1.925 p=0.350	β=-0.384 se=0.171 df=18.000 d=-0.392 t=-2.246 p=0.185

Table 3. Statistical results (two-sided) for the contrasts between music conditions and three kinds of baselines (fixation, nonwords, and non-linguistic non-music auditory conditions—animal sounds and environmental sounds) in Experiments 1 and 2, and for the contrast between songs and lyrics in Experiment 1. Abbreviations: b=the beta estimate for the effect; se=standard error of the mean by participants; df=degrees of freedom; d=Cohen's d (Westfall et al. 2014; Brysbaert and Stevens 2018); t=the t statistic; p=the significance value (for the individual fROIs, these values have been FDR-corrected for the number of fROIs (n=5)). In light grey, we highlight the results that are *not consistent* with the role of the language regions in music perception: of the 84 tests performed, 1 showed an effect predicted by language-music overlap accounts: a small and statistically weak (only emerging at the network level but not in any individual fROI) positive response, relative to the weakest baseline (fixation), to one of the five music conditions examined; and this response was still ~4 times lower than the response to an unstructured linguistic condition (nonwords).

Is the language network sensitive to structure in music?

Experiments 1 and 2 (fMRI): Because most prior claims about the overlap between language and music concern the processing of *structure*—given the parallels that can be drawn between the syntactic structure of language and the tonal and rhythmic structure in music (e.g., Lerdahl and Jackendoff 1977, 1983; cf. Jackendoff 2009)—we used three contrasts to test whether language regions are sensitive to music structure. First and second, in Experiment 1, we compared the responses to synthetic melodies vs. their scrambled counterparts, and to synthetic drum music vs. the scrambled drum music condition. The former targets both tonal and rhythmic structure, and the latter selectively targets rhythmic structure. The reason to examine rhythmic structure is that some patient studies have argued that pitch contour processing relies on the right hemisphere, and rhythm processing draws on the left hemisphere (e.g., Zatorre 1984; Peretz 1990; Alcock et al. 2000; cf. Boebinger 2021 for fMRI evidence of bilateral responses in high-level auditory areas to both tonal and rhythmic structure processing and for lack of spatial segregation between the two), so although most prior work examining the language-music relationship has focused on tonal structure, rhythmic structure may *a priori* be more likely to overlap with linguistic syntactic structure given their alleged co-lateralization based on the patient literature. And third, in Experiment 2, we compared the responses to well-formed melodies vs. melodies with a sour note. A brain region that responds to structure in music should respond more strongly to intact than scrambled music (similar to how language regions respond more strongly to sentences than lists of words; e.g., Fedorenko et al. 2010; Diachek, Blank, Siegelman et al. 2020), and also exhibit sensitivity to structure violations (similar to how language regions respond more strongly to sentences that contain grammatical errors: e.g., Embick et al. 2000; Newman et al. 2001;

Kuperberg et al. 2003; Cooke et al. 2006; Friederici et al. 2010; Herrmann et al. 2012; Fedorenko

et al. 2020). Note that given the lack of a strong and consistent response to music in the language

regions (**Figure 3** and **Table 3**), the answer to this narrower question is somewhat of a foregone

conclusion: even if one or more of the language regions showed a reliable effect in these music-

structure-probing contrasts, such effects would be difficult to interpret as reflecting music

structure processing given that structured music stimuli elicit a response approximately at the

level of the fixation baseline in the language areas. Nevertheless, we report the results for these

three contrasts for completeness, and because most prior studies have focused on such contrasts.

The language regions did not show consistent sensitivity to structural manipulations in music

(**Figure 4**; **Table 4**). In Experiment 1, the responses to synthetic melodies did not significantly

differ from (or were weaker than) the responses to the scrambled counterparts, and the responses

to synthetic drum music did not significantly differ from the responses to scrambled drum music.

In Experiment 2, at the network level, we observed a small and weakly significant ($p<0.05$)

effect of *sour-note > well-formed melodies*. This effect was not significant in any of the five

individual fROIs (even prior to the FDR correction).

Figure 4. Responses of the language fROIs (pooling across the network – top, and for each fROI

individually – bottom) to the language localizer conditions (in grey), and to the three sets of conditions

that target structure in music (in blue). The error bars represent standard error of the mean by participants.

For the language localizer results, we include here participants in Experiments 1 and 2. The responses to

the music conditions cluster around the fixation baseline, and are much lower than the response to

sentences. One of the three critical contrasts (*sour-note > well-formed* melodies) elicits a small and

weakly reliable effect at the network level, but it is not individually significant in any of the five fROIs.

Contrast	Language network	LIFGorb	LIFG	LMFG	LAnt Temp	LPost Temp
synthetic drum music >scrambled drum music	β=0.099 se=0.073 df=157.823 d=0.140 t=1.358 p=0.176	β=0.252 se=0.191 df=18.000 d=0.288 t=1.322 p=1.000	β=0.027 se=0.176 df=18.000 d=0.034 t=0.156 p=1.000	β=0.014 se=0.186 df=18.000 d=0.018 t=0.073 p=1.000	β=0.124 se=0.103 df=18.000 d=0.247 t=1.210 p=1.000	β=0.079 se=0.110 df=18.000 d=0.165 t=0.718 p=1.000
synthetic melodies >scrambled synthetic melodies	β=-0.124 se=0.061 df=157.720 d=-0.238 t=-2.015 p=0.046*	β=-0.147 se=0.130 df=18.000 d=-0.245 t=-1.133 p=1.000	β=-0.009 se=0.153 df=18.000 d=-0.017 t=-0.057 p=1.000	β=-0.143 se=0.202 df=18.000 d=-0.216 t=-0.708 p=1.000	β=-0.199 se=0.101 df=18.000 d=-0.572 t=-1.971 p=0.320	β=-0.121 se=0.106 df=18.000 d=-0.365 t=-1.142 p=1.000
sour-note melodies >well-formed melodies	β=0.145 se=0.069 df=175.884 d=0.196 t=2.102 p=0.037*	β=0.195 se=0.098 df=20.000 d=0.245 t=1.985 p=0.305	β=0.150 se=0.105 df=20.000 d=0.180 t=1.431 p=0.840	β=0.212 se=0.090 df=20.000 d=0.252 t=2.363 p=0.140	β=0.065 se=0.051 df=20.000 d=0.114 t=1.280 p=1.000	β=0.104 se=0.056 df=20.000 d=0.248 t=1.856 p=0.390

Table 4. Statistical results (two-sided) for the contrasts between the synthetic drum music and scrambled

drum music, synthetic melodies and scrambled synthetic melodies, and sour-note and well-formed

melodies contrasts in Experiments 1 and 2. Abbreviations: b=the beta estimate for the effect; se=standard

error of the mean by participants; df=degrees of freedom; d=Cohen's d (Westfall et al. 2014; Brysbaert

and Stevens 2018); t=the t statistic; p=the significance value (for the individual fROIs, these values have

been FDR-corrected for the number of fROIs (n=5)). In light grey, we highlight the results that are ***not***

consistent with the role of the language regions in the processing of music structure: of the 18 tests

performed, 1 showed an effect predicted by language-music overlap accounts: a small and statistically

weak response to one of the three structure-targeting contrasts (in the presence of an overall very weak

response to music relative to fixation; see Figure 3 and Table 3).

Experiment 3 (behavioral): In Experiment 3, we further asked whether individuals with severe

deficits in processing linguistic syntax also exhibit difficulties in processing music structure. To

do so, we assessed participants' ability to discriminate well-formed ("good") melodies from

melodies with a sour note ("bad"), while controlling for their response bias (how likely they are

overall to say that something is well-formed) by computing *d'* for each participant (Green and

Swets 1966), in addition to proportion correct. We then compared the *d'* values of each

individual with aphasia to the distribution of *d'* values of healthy control participants using a

Bayesian test for single case assessment (Crawford and Garthwaite 2007) as implemented in the

psycho package in R (Makowski 2018). (Note that for the linguistic syntax tasks, it was not

necessary to conduct statistical tests comparing the performance of each individual with aphasia

to the control distribution because the performance of each individual with aphasia was lower

than 100% of the control participants' performances.) We similarly compared the proportion

correct on the MBEA scale task of each individual with aphasia to the distribution of accuracies

of healthy controls. If linguistic and music syntax draw on the same resources, then individuals

with linguistic syntactic impairments should also exhibit deficits on tasks requiring the

processing of music syntax.

In the critical music task, where participants were asked to judge the well-formedness of musical

structure, neurotypical control participants responded correctly, on average, on 87.1% of trials,

suggesting that the task was sufficiently difficult to preclude ceiling effects. Patients with severe

aphasia showed intact sensitivity to music structure. The three patients had accuracies of 89.4%

(PR), 94.4% (SA), and 97.8% (PP), falling on the higher end of the controls' performance range

(**Figure 5**; **Table 5**). Crucially, none of the three aphasic participants' *d'* scores were lower than

the average control participants' *d'* scores (M = 2.75, SD = 0.75). In fact, the patients' *d'* scores

were high: SA's *d'* was 3.51, higher than 83.91% (95% Credible Interval (CI) [75.20, 92.03]) of

the control population, PR's *d'* was 3.09, higher than 67.26% (95% CI [56.60, 78.03]) of the

control population, and PP's *d'* was 3.99, higher than 94.55% (95% CI [89.40, 98.57]) of the

control population. None of the three aphasic participants' bias/criterion c scores (Green and

Swets 1966) differed reliably from the control participants' c scores (M = -0.40, SD = 0.40).

SA's c was -0.53, lower than 62.34% (95% CI [50.40, 71.67]) of the control population, PR's c

was -0.74, lower than 79.48% (95% CI [69.58, 88.44]) of the control population, and PP's c was

-0.29, higher than 60.88% (95% CI [50.08, 70.04]) of the control population. In the Scale task

from the Montreal Battery for the Evaluation of Aphasia, the control participants' performance

showed a similar distribution to that reported in Peretz et al. (2003). All participants with aphasia

performed within the normal range, with two participants making no errors. PR and PP's score

was higher than 85.24% (95% CI [76.94, 93.06]) of the control population, providing a

conceptual replication of the results from the well-formed/sour-note melody discrimination task.

SA's score was higher than 30.57% (95% CI [20.00, 41.50]) of the control population.

Participant	SA	PR	PP	Controls

Critical Music Task	170/180	161/180	176/180	M = 156.5/180 SD = 15.8 Min = 109/180 Max = 177/180 N=45
Montreal Battery for the Evaluation of Amusia				
(Critical for this study) Task 1 (Scale)	27/30	30/30	30/30	M = 28/30 SD = 1.89 Min = 23/30 Max = 30/30 N = 45
Task 2 (Interval; "Same Contour" on MBEA CD)	26/30	22/30	18/30	
Task 3 (Contour; "Different Contour" on MBEA CD)	22/30	23/30	18/30	
Task 4 (Rhythm; "Rhythmic Contour" on MBEA CD)	25/30	25/30	22/30	
Task 5 (Meter; "Metric" on MBEA CD)	28/30	22/30	24/30	
Task 6 (Incidental Memory)	28/30	28/30	22/30	

Table 5. Results for participants with aphasia and control participants on the critical music task and the

Scale task of the MBEA (Peretz et al., 2003). For participants with aphasia, we report the results from all

six MBEA tasks, for completeness.

Figure 5. Performance of the control and aphasic participants on two measures of music syntax processing:

the critical music task (left), the Scale task of the MBEA (right). The densities show the distribution of

proportion correct scores in the control participants and the boxplot shows the quartiles of the control

population (the whiskers show 1.5x interquartile range and points represent outliers). The dots show

individual participants (for the aphasic individuals, the initials indicate the specific participant). Dashed

grey lines indicate chance performance.

Does music elicit a response in the language network of native speakers of a tonal language?

The above analyses focus on the language network's responses to music stimuli and its

sensitivity to music structure in English native speakers. However, some have argued that

responses to music may differ in speakers of languages that use pitch to make lexical or

grammatical distinctions (e.g., Deutsch et al. 2006, 2009; Bidelman et al. 2011; Creel et al. 2018;

Ngo et al. 2016, Liu et al. 2021). In Experiment 4, we therefore tested whether language regions

of Mandarin native speakers respond to music. Similar to Experiment 1, we compared the

response to the music condition against a) the fixation baseline, b) the foreign language

condition, and c) a non-linguistic, non-music condition (environmental sounds). A brain region

that supports music processing should respond more strongly to music than the fixation baseline

and the foreign condition; if the response is further selective, it should be stronger than the

response elicited by environmental sounds.

Results from Mandarin native speakers replicated the results from Experiment 1: the music

condition did not elicit a strong response in the language network (**Figure 6**; **Table 6**). Although

the response to music was above the fixation baseline at the network level and in some fROIs,

the response did not differ from (or was lower than) the responses elicited by an unfamiliar

foreign language (Russian) and environmental sounds.

Figure 6. Responses of the language fROIs (pooling across the network – top, and for each fROI

individually – bottom) to the language localizer conditions (in grey), to the two auditory conditions

containing speech (red shades), to the music condition (blue), and to the non-linguistic/non-music

auditory condition (green) in Experiment 4. The error bars represent standard error of the mean by

participants. The response to the music condition is much lower than the response to sentences, and is not

higher than the response to foreign language and environmental sounds.

Contrast	Language network	LIFGorb	LIFG	LMFG	LAnt Temp	LPost Temp
music >fixation	β=0.454 se=0.177 df=17.646 d=0.517 t=2.565 p=0.020*	β=0.299 se=0.228 df=nan d=nan t=1.308 p=1.000	β=0.761 se=0.207 df=nan d=nan t=3.683 p=0.010*	β=0.480 se=0.260 df=nan d=nan t=1.848 p=0.410	β=0.268 se=0.171 df=nan d=nan t=1.568 p=0.675	β=0.462 se=0.156 df=nan d=nan t=2.962 p=0.045*
music >foreign	β=-0.359 se=0.141 df=162.000 d=-0.308 t=-2.547 p=0.012*	β=-0.360 se=0.416 df=18.000 d=-0.258 t=-0.865 p=1.000	β=0.123 se=0.309 df=18.000 d=0.124 t=0.398 p=1.000	β=-0.219 se=0.473 df=18.000 d=-0.149 t=-0.463 p=1.000	β=-0.703 se=0.240 df=18.000 d=-0.870 t=-2.926 p=0.045*	β=-0.638 se=0.254 df=18.000 d=-0.686 t=-2.511 p=0.110
music >environmental sounds	β=-0.141 se=0.108 df=157.749 d=-0.154 t=-1.299 p=0.196	β=-0.249 se=0.187 df=18.000 d=-0.280 t=-1.328 p=1.000	β=-0.240 se=0.193 df=18.000 d=-0.302 t=-1.248 p=1.000	β=0.038 se=0.304 df=18.000 d=0.030 t=0.125 p=1.000	β=-0.042 se=0.147 df=18.000 d=-0.065 t=-0.285 p=1.000	β=-0.210 se=0.179 df=18.000 d=-0.310 t=-1.171 p=1.000

Table 6. Statistical results (two-sided) for the contrasts between the music condition and fixation, foreign language, and environmental sounds in Experiment 4. Abbreviations: b=the beta estimate for the effect; se=standard error of the mean by participants; df=degrees of freedom; d=Cohen's d (Westfall et al. 2014; Brysbaert and Stevens 2018); t=the t statistic; p=the significance value (for the individual fROIs, these values have been FDR-corrected for the number of fROIs (n=5)). In light grey, we highlight the results that are ***not consistent*** with the role of the language regions in music perception: of the 18 tests performed, 3 showed an effect predicted by language-music overlap accounts: a small positive response

to the music condition relative to the weakest baseline (fixation) at the network level and in two fROIs

individually; this response was still ~2 lower than the unfamiliar foreign language condition and was

numerically lower than the environmental sounds condition.

Discussion

We here tackled a much investigated but still debated question: do the brain regions of the

language network support the processing of music, especially music structure? Across three

fMRI experiments, we obtained a clear answer: the brain regions of the language network, which

support the processing of linguistic syntax (e.g., Fedorenko et al. 2010, 2020; Pallier et al. 2011;

Bautista and Wilson 2016; Blank et al. 2016), do not support music processing (see **Table 7** for a

summary of the results). We found overall low responses to music (including orchestral pieces,

solo pieces played on different instruments, synthetic music, and vocal music) in the language

brain regions (**Figure 3**; see Sueoka et al. 2022, for complementary evidence from the inter-

subject correlation approach applied to a rich naturalistic music stimulus), including in speakers

of a tonal language (**Figure 6**), and no consistent sensitivity to manipulations of music structure

(**Figure 4**). We further found that the ability to make well-formedness judgments about the tonal

structure of music was preserved in patients with severe aphasia who cannot make

grammaticality judgments for sentences (**Figure 5**), although we acknowledge the possibility

that general ability to detect unexpected events may have contributed to performance on the

critical music-structure tasks (e.g., Bigand et al. 2014; Collins et al. 2014) and that additional

controls would be needed to conclusively determine whether these patients have preserved

music-structure processing abilities. Nevertheless, given the brain imaging results (summarized

in **Table 7**), a critical role of the language system in music structure processing is unlikely.

Our findings align with a) prior neuropsychological patient evidence of language/music

dissociations (e.g., Luria et al. 1965; Brust 1980; Marin 1982; Basso and Capitani 1985; Polk

and Kertesz 1993; Peretz et al. 1994, 1997; Piccirilli et al. 2000; Peretz and Coltheart 2003;

Slevc et al. 2016; Faroqi-Shah et al. 2020; Chiapetta et al. 2022) and with b) prior evidence that

music is processed by music-selective areas in the auditory cortex (Norman-Haignere et al.

(2015; see also Boebinger et al. 2021; see Peretz et al. 2015, for review and discussion). The

latter, music-selective areas are strongly sensitive to the scrambling of music structure in stimuli

like those used here in Experiment 1 (see also Fedorenko et al. 2012c, Boebinger 2021; see Mehr

et al. 2019 for *a priori* reasons to expect the effects of tonal structure manipulations in music-

selective brain regions). (We provide the responses of music-responsive areas to the conditions

of Experiments 1 and 2 at: https://osf.io/68y7c/).) In contrast, our findings stand in sharp contrast

to numerous reports arguing for shared structure processing mechanisms in the two domains,

including specifically in the inferior frontal cortex, within 'Broca's area' (e.g., Patel et al. 1998;

Koelsch et al. 2000; Maess et al. 2001; Koelsch et al. 2002; Levitin and Menon 2003; see Kunert

and Slevc 2015; LaCroix et al. 2016; Vuust et al. 2022 for reviews).

	Contrast	Experiment 1	Experiment 2	Experiment 4
Basic sensitivity to music stimuli	Music > fixation (6 different music conditions tested:	**No**	No (except for the network level)	Yes

	4 in Expt1, 1 in Expt2, and 1 in Expt4)			
	Music > nonwords/unfamiliar foreign language	**No**		**No**
	Music > non-linguistic, non-music auditory conditions	**No**		**No**
	Songs (melodic contour + linguistic content) > Lyrics (linguistic content)	**No**		
Sensitivity to manipulations of music structure	Intact music > scrambled music (synthetic melodies)	**No**		
	Intact music > scrambled music (synthetic drums)	**No**		
	Sour-note melodies > well-formed melodies		No (except for the network level)	

Table 7. A summary of the results for the tests of the language network's sensitivity to music in general

and to music structure specifically. This pattern of results constitutes strong evidence against the role of

the language system—or any of its components—in music perception, including the processing of music

structure. With respect to sensitivity to music stimuli: 4 of the 6 conditions failed to elicit a response

above the low-level (fixation) baseline anywhere in the language network; 1 condition (in Experiment 2)

elicited a small and weakly significant above-fixation response at the network level only (not in any

individual fROIs); and 1 condition (in Experiment 4) elicited a small above-fixation response (including

in two individual fROIs) but this response was not higher than that elicited by other auditory conditions

like environmental sounds. With respect to sensitivity to music structure: 2 of the 3 manipulations failed

to elicit a response anywhere in the language network, and the remaining manipulation elicited a small

and weakly significant effect at the network level, which was not reliable in any individual ROI.

1008

1009

1010   Below, we discuss several issues that are relevant for interpreting the current results and/or that

1011   these results inform, and outline some limitations of scope of our study.

1012

1013   *1. Theoretical considerations about the language-music relationship.*

1014

Why might we *a priori* think that the language network, or some of its components, may be

important for processing music in general, or for processing music structure specifically?

Similarities between language and music have long been noted and discussed. For example, as

summarized in Jackendoff (2009; see also Patel 2008), both capacities are human-specific,

involve the production of sound (though this is not always the case for language: cf. sign

languages, or written language in literate societies), and have multiple culture-specific variants.

Furthermore, language and music are intertwined in songs, which appear to be a cultural

universal (e.g., Brown 1991; Nettl 2015; see Mehr et al. 2019 for empirical support; see Norman-

Haignere et al. 2021 for evidence of neural selectivity for songs in the auditory cortex).

However, Jackendoff (2009) notes that i) most cognitive capacities / mechanisms that have been

argued to be common to language and music are not *uniquely* shared by language and music, and

ii) language and music differ in several critical ways, and these differences are important to

consider alongside potential similarities when theorizing about possible shared representations

and computations.

1029

To elaborate on the first point: the cognitive capacity that has perhaps received the most attention

in discussions of cognitive and neural mechanisms that may be shared by language and music is

the combinatorial capacity of the two domains (e.g., Riemann 1877, as cited in Swain 1995;

Lindblom and Sundberg 1969; Fay 1971; Sundberg and Lindblom 1976; Lerdahl and Jackendoff

1977, 1983; Roads 1979; Krumhansl and Keil 1982). In particular, in language, words can be

combined into complex hierarchical structures to form novel phrases and sentences, and in

music, notes and chords can similarly be combined to form novel melodies. Further, in both

domains, the combinatorial process is constrained by a set of conventions. However, this

capacity can be observed, in some form, in many other domains, from visual processing, to math,

to social cognition, to motor planning, to general reasoning. Similarly, other cognitive capacities

that are necessary to process language and music—including a large long-term memory store for

previously encountered elements and patterns, a working memory capacity needed to integrate

information as it comes in, an ability to form expectations about upcoming elements, and an

ability to engage in joint action—are important for information processing in other domains. An

observation that some mental capacity is necessary for multiple domains is compatible with at

least two architectures: one where the relevant capacity is implemented (perhaps in a similar

way) in each relevant set of domain-specific circuits, and another where the relevant capacity is

implemented in a centralized mechanism that all domains draw on (e.g., Fedorenko and Shain

2021). Those arguing for overlap between language and music processing advocate a version of

the latter. Critically, any shared mechanism that language and music would draw on should also

support information processing in other domains that require the relevant computation (see

Section 3 below for arguments against this kind of architecture). (A possible exception,

according to Jackendoff (2009), may be the fine-scale vocal motor control that is needed for

speech and vocal music production (cf. sign language or instrumental music), but not any other

behaviors, but this kind of ability is implemented outside of the core high-level language system,

in the network of brain areas that support articulation (e.g., Basilakos et al. 2015; Guenter

2016).)

1057

More importantly, aside from the similarities that have been noted between language and music,

numerous differences characterize the two domains. Most notable are their different functions.

Language enables humans to express propositional meanings, and thus to share thoughts with

one another. The function of music has long been debated (e.g., Darwin 1871; Pinker 1994; see

e.g., McDermott 2008 and Mehr et al. 2020, for a summary of key ideas), but most proposed

functions have to do with emotional or affective processing, often with a social component[1]

(Jackendoff 2009; Savage et al. 2020). If function drives the organization of the brain (and

biological systems more generally; e.g., Rueffler et al. 2012) by imposing particular

computational demands on each domain (e.g., Mehr et al. 2020), these fundamentally different

functions of language and music provide a theoretical reason to expect cognitive and neural

separation between them. Besides, even the components of language and music that appear

similar on the surface (e.g., combinatorial processing) differ in deep and important ways (e.g.,

Patel 2008; Jackendoff 2009; Slevc 2009; Temperley 2022).

---

[1] Although some have discussed the notions of 'meaning' in music (e.g., Meyer 1961; Raffman 1993; Cross and Tolbert 2009; Koelsch 2001), it is uncontroversial that music cannot be used to express propositional thought (for discussion, see Patel 2008; Jackendoff 2009; Slevc 2009).

1071

1072   *2. Functional selectivity of the language network.*

1073

The current results add to the growing body of evidence that the left-lateralized fronto-temporal

brain network that supports language processing is highly selective for linguistic input (e.g.,

Fedorenko et al. 2011; Monti et al. 2009, 2012; Deen et al. 2015; Pritchett et al. 2018; Jouravlev

et al. 2019; Ivanova et al. 2020, 2021; Benn, Ivanova et al. 2021; Liu et al. 2020; Deen and

Freiwald 2021; Paunov et al. 2022; Sueoka et al. 2022; see Fedorenko and Blank 2020 for a

review) and not critically needed for many forms of complex cognition (e.g., Lecours and

Joanette 1980; Varley and Siegal 2000; Varley et al. 2005; Apperly et al. 2006; Woolgar et al.

2018; Ivanova et al. 2021; see Fedorenko and Varley 2016 for a review). Importantly, this

selectivity holds across all components of the language network, including the parts that fall

within 'Broca's area' in the left inferior frontal gyrus. As discussed in the introduction, many

claims about shared structure processing in language and music have focused specifically on

Broca's area (e.g., Patel 2003; Fadiga et al. 2009; Fitch and Martins 2014). The evidence

presented here shows that the language-responsive parts of Broca's area, which are robustly

sensitive to linguistic syntactic manipulations (e.g., Just et al. 1996; Stromswold et al. 1996;

Ben-Shachar et al. 2003; Caplan et al. 2008; Peelle et al. 2010; Blank et al. 2016; see e.g.,

Friederici 2011 and Hagoort and Indefrey 2014 for meta-analyses), do not respond when we

listen to music and are not sensitive to structure in music. These results rule out the hypothesis

that language and music processing rely on the same mechanism housed in Broca's area.

1092

It is also worth noting that the very *premise* of the latter hypothesis—of a special relationship

between Broca's area and the processing of linguistic syntax (e.g., Caramazza and Zurif 1976;

Friederici 2018)—has been questioned and overturned. *First*, syntactic processing does not

appear to be carried out focally, but is instead distributed across the entire language network,

with all of its regions showing sensitivity to syntactic manipulations (e.g., Fedorenko et al. 2010,

2020; Pallier et al. 2011; Blank et al. 2016; Shain, Blank et al. 2020; Shain et al. 2022), and with

damage to different components leading to similar syntactic comprehension deficits (e.g., Caplan

et al. 1996; Dick et al. 2001; Wilson and Saygin 2004; Mesulam et al. 2014; Mesulam et al.

2015). And *second*, the language-responsive part of Broca's area, like other parts of the language

network, is sensitive to both syntactic processing and word meanings, and even sub-lexical

structure (Fedorenko et al. 2010, 2012b, 2020; Regev et al. 2021; Shain et al. 2021). The lack of

segregation between syntactic and lexico-semantic processing is in line with the idea of

'lexicalized syntax' where the conventions for how words can combine with one another are

highly dependent on the particular lexical items (e.g., Goldberg 2002; Jackendoff 2002, 2007;

Sag et al. 2003; Levin and Rappaport-Hovav 2005; Bybee 2010; Jackendoff and Audring 2020),

and is contra the idea of combinatorial rules that are blind to the content/meaning of the to-be-

combined elements (e.g., Chomsky 1965, 1995; Fodor 1983; Pinker and Prince 1988; Pinker

1991, 1999; Pallier et al. 2011).

1111

1112   *3. Overlap in structure processing in language and music outside of the core language*

1113   *network?*

1114

We have here focused on the core fronto-temporal language network. Could structure processing

in language and music draw on shared resources elsewhere in the brain? The prime candidate is

the domain-general executive control, or Multiple Demand (MD), network (e.g., Duncan and

Owen 2000; Duncan 2001, 2010; Assem et al. 2020), which supports functions like working

memory and inhibitory control. Indeed, according to Patel's Shared Structural Integration

Resource Hypothesis (SSIRH; 2003, 2008, 2012), language and music draw on separate

representations, stored in distinct cortical areas, but rely on the same working memory store to

integrate incoming elements into evolving structures. Relatedly, Slevc et al. (2013; see Asano et

al. 2021 for a related proposal) have argued that another executive resource—inhibitory

control—may be required for structure processing in both language and music. Although it is

certainly possible that some aspects of linguistic and/or musical processing would require

domain-general executive resources, based on the available evidence from the domain of

language, we would argue that any such engagement does not reflect the engagement of

computations like syntactic structure building. In particular, Blank and Fedorenko (2017) found

that activity in the brain regions of the domain-general MD network does not closely 'track'

linguistic stimuli, as evidenced by low inter-subject correlations during the processing of

linguistic input (see Paunov et al. 2021 and Sueoka et al. 2022 for replications). Further,

Diachek, Blank, Siegelman et al. (2020) showed in a large-scale fMRI investigation that the MD

network is not engaged during language processing in the absence of secondary task demands

(cf. the core language network, which is relatively insensitive to task demands and responds

robustly even during passive listening/reading). And Shain, Blank et al. (2020; also, Shain et al.

2022) have shown that the language network, but not the MD network, is sensitive to linguistic

surprisal and working-memory integration costs (see also Wehbe et al. 2021 for evidence that

activity in the language, but not the MD, network reflects general incremental processing

difficulty).

1140

In tandem, this evidence argues against the role of executive resources in core linguistic

computations like those related to lexical access and combinatorial processing, including

syntactic parsing and semantic composition (see also Hasson et al. 2015 and Dasgupta and

Gershman 2021 for general arguments against the separation between memory and computation

in the brain). Thus, although the contribution of executive resources to music processing

deserves further investigation (cf. https://osf.io/68y7c/ for evidence of low responses of the MD

network to the music conditions in the current study), any overlap within the executive system

between linguistic and music processing cannot reflect core linguistic computations, as those

seem to be carried out by the language network (see Fedorenko and Shain 2021, for a review).

Functionally identifying the MD network in individual participants (e.g., Fedorenko et al. 2013;

Shashidhara et al. 2019) is a powerful way to help interpret the observed effects of music

manipulations as reflecting general executive demands (see Saxe et al. 2006, Blank et al. 2017

and Fedorenko 2021, for general discussions of greater interpretability of fMRI results obtained

from the functional localization approach). Importantly, given the ubiquitous sensitivity of the

MD network to cognitive demands, it is / will be important to rule out task demands, rather than

stimulus processing, as the source of overlap between music and language processing in

interpreting past studies and designing future ones.

1158

1159 *4. Overlap between music processing and other aspects of speech / language.*

1160

The current study investigated the role of the language network—which supports 'high-level'

comprehension and production—in music processing. As a result, the claims we make are

restricted to those aspects of language that are supported by this network. These include the

processing of word meanings and combinatorial (syntactic and semantic) processing, but exclude

speech perception, prosodic processing, higher-level discourse structure building, and at least

some aspects of pragmatic reasoning. Some of these components of language (e.g., pragmatic

reasoning) seem *a priori* unlikely to share resources with music. Others (e.g., speech perception)

have been shown to robustly dissociate from music (Norman-Haignere et al. 2015; Overath et al.

2015; Kell et al. 2018; Boebinger et al. 2021). However, some components of speech and

language may, and some do, draw on the same resources as aspects of music. For example,

aspects of pitch perception have been argued to overlap between speech and music based on

behavioral and neuropsychological evidence (e.g., Wong and Perrachione 2007; Perrachione et

al. 2013; Patel et al. 2008b). Indeed, brain regions that selectively respond to different kinds of

pitched sounds have been previously reported (Patterson et al. 2002; Penagos et al. 2004;

Norman-Haignere et al. 2013, 2015). Some studies have also suggested that music training may

improve general rapid auditory processing and pitch encoding that are important for speech

perception and language comprehension (e.g., Overy 2003; Tallal and Gaab 2006; Wong et al.

2007), although at least some of these effects likely originate in the brainstem and subcortical

auditory regions (e.g., Wong et al. 2007). Other aspects of high-level auditory perception,

including aspects of rhythm, may turn out to overlap as well, and deserve further investigation

(see Patel 2008, for a review).

1182

We also have focused on Western tonal instrumental music here. In the future, it would be useful

to extend these findings to more diverse kinds of music. That said, given that individuals are

most sensitive to structure in music with which they have experience (e.g., Cuddy et al. 1981;

Cohen 1982; Curtis and Barucha 2009), it seems unlikely that music from less familiar traditions

would elicit a strong response in the language areas (see Boebinger 2021, for evidence that

music-selective areas of the auditory cortex respond to culturally diverse music styles). Further,

given that evolutionarily early forms of music were likely vocal (e.g., Trehub 2003; Mehr 2017),

it would be useful to examine the responses of the language regions to vocal music without

linguistic content, like humming or whistling. Based on preliminary unpublished data from our

lab (available upon request), responses to such stimuli in the language areas appear low.

1193

In conclusion, we have here provided extensive evidence against the role of the language

network in music perception, including the processing of music structure. Although the

relationship between music and aspects of speech and language will likely continue to generate

interest in the research community, and aspects of speech and language other than those

implemented in the core fronto-temporal language-selective network (Fedorenko et al. 2011;

Fedorenko and Thompson-Schill 2014) may indeed share some processing resources with

(aspects of) music, we hope that the current study helps bring clarity to the debate about structure

processing in language and music.

1202

1203

1204

1205

Data availability

The datasets generated during and/or analyzed during the current study are available in the OSF

repository: https://osf.io/68y7c/.

Code availability

Scripts for statistical analysis are available at: https://osf.io/68y7c/.

Acknowledgements

We would like to acknowledge the Athinoula A. Martinos Imaging Center at the McGovern

Institute for Brain Research at MIT, and its support team (Steve Shannon and Atsushi

Takahashi). We thank former and current EvLab members for their help with fMRI data

collection (especially Meilin Zhan for help with Experiment 4). We thank Josh McDermott for

input on many aspects of this work, Jason Rosenberg for composing the melodies used in

Experiments 2 and 3, and Zuzanna Balewski for help with creating the final materials used in

Experiments 2 and 3. For Experiment 3, we thank Vitor Zimmerer for help with creating the

grammaticality judgment task, Ted Gibson for help with collecting the control data, and Anya

Ivanova for help with Figure 2. For Experiment 4, we thank Anne Cutler, Peter Graff, Morris

Alper, Xiaoming Wang, Taibo Li, Terri Scott, Jeanne Gallée, and Lauren Clemens for help with

constructing and/or recording and/or editing the language materials, and Fatemeh Khalilifar,

Caitlyn Hoeflin, and Walid Bendris for help with selecting the music materials and with the experimental script. Finally, we thank the audience at the Society for Neuroscience conference (2014), the Neurobiology of Language conference (virtual edition, 2020), Ray Jackendoff, Dana Boebinger, and members of the Fedorenko and Gibson labs for helpful comments and discussions. RR was supported by NIH award F32-DC-015163. SNH was supported by a graduate NSF award, as well as postdoctoral awards from the HHMI / Life Sciences research foundation and a K99/R00 award from the NIH (1K99DC018051-01A1). SMM was supported by a La Caixa fellowship LCF/BQ/AA17/11610043. RV was supported by Alzheimer's Society and The Stroke Association. EF was supported by the R00 award HD057522, R01 awards DC016607, DC016950, and NS121471, by the Paul and Lilah Newton Brain Science Award, and funds from the Brain and Cognitive Sciences department, the McGovern Institute for Brain Research, and the Simons Center for the Social Brain.

Author contributions:

	XC	JA	RR	TR	SNH	OJ	SMM	HK	RV†	EF†
Conceptualization									☑	☑
Methodology	☑			☑	☑				☑	☑
Software	☑	☑								
Investigation	☑	☑	☑			☑	☑	☑	☑	☑
Investigation: fMRI data collection	☑	☑				☑				☑

Investigation: fMRI data preprocessing and analysis	☑	☑				☑	☑	☑		
Investigation: Behavioral data collection	☑		☑						☑	
Investigation: Behavioral data analysis			☑						☑	☑
Formal statistical analysis	☑		☑							☑
Validation	☑		☑							
Visualization	☑		☑	☑						
Writing: Original draft	☑		☑						☑	☑
Writing: Editing + comments		☑		☑	☑	☑	☑	☑		
Resources									☑	☑
Project administration; overall supervision										☑

1245

Conflict of interest

1246

1247 The authors declare no competing financial interests.

References

Alcock KJ, Wade D, Anslow P, Passingham RE. 2000. Pitch and timing abilities in adult left-hemisphere-dysphasic and right-hemisphere-damaged subjects. *Brain Lang.* 75(1):47-65.

Amalric M, Dehaene S. 2018. Cortical circuits for mathematical knowledge: evidence for a major subdivision within the brain's semantic networks. *Philos Trans R Soc Lond B Biol Sci.* 373(1740):20160515.

Apperly IA, Samson D, Carroll N, Hussain S, Humphreys G. 2006. Intact first-and second-order false belief reasoning in a patient with severely impaired grammar. *Soc Neurosci.* 1(3-4):334-348.

Asano R, Boeckx C, Seifert U. 2021. Hierarchical control as a shared neurocognitive mechanism for language and music. *Cognition. 216*:104847.

Assem M, Glasser MF, Van Essen DC, Duncan J. 2020. A domain-general cognitive core defined in multimodally parcellated human cortex. *Cereb Cortex.* 30(8):4361-4380.

Baillet S. 2014. Forward and Inverse Problems of MEG/EEG. In: Jaeger D, Jung R, editors. Encyclopedia of Computational Neuroscience. New York (NY): Springer. p.1-8.

Baroni M, Maguire S, Drabkin W. 1983. The concept of musical grammar. *Music Anal*, 2(2):175-208.

Basilakos A, Rorden C, Bonilha L, Moser D, Fridriksson J. 2015. Patterns of poststroke brain damage that predict speech production errors in apraxia of speech and aphasia dissociate. Stroke. 46(6):1561-1566.

Basso A, Capitani E. 1985. Spared musical abilities in a conductor with global aphasia and ideomotor apraxia. *J Neurol Neurosurg Psychiatry*. 48(5):407-412.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *J Stat Softw*. 67(1):1–48.

Bautista A, Wilson SM. 2016. Neural responses to grammatically and lexically degraded speech. *Lang Cogn Neurosci*. 31(4):567-574.

Ben-Shachar M, Hendler T, Kahn I, Ben-Bashat D, Grodzinsky Y. 2003. The neural reality of syntactic transformations: Evidence from functional magnetic resonance imaging. *Psychol Sci*. 14(5):433-440.

Benn Y*, Ivanova A*, Clark O, Mineroff Z, Seikus C, Santos Silva J, Varley R^, Fedorenko E^. 2021. No evidence for a special role of language in feature-based categorization. *bioRxiv*.

Bernstein L. 1976. The unanswered question: Six talks at Harvard. Cambridge (MA): Harvard University Press.

1295   Bidelman GM, Gandour JT, Krishnan A. 2011. Musicians and tone-language speakers share

1296   enhanced brainstem encoding but not perceptual benefits for musical pitch. *Brain Cogn.* 77(1):1-

1297   10.

1298

1299   Bigand E, Tillmann B, Poulin B, D'Adamo DA, Madurell F. 2001. The effect of harmonic

1300   context on phoneme monitoring in vocal music. *Cognition.* 81(1):B11-B20.

1301

1302   Bigand E, Delbé C, Poulin-Charronnat B, Leman M, Tillmann B. 2014. Empirical evidence for

1303   musical syntax processing? Computer simulations reveal the contribution of auditory short-term

1304   memory. *Front Syst Neurosci.* 8:94.

1305

1306   Bishop DVM, Norbury CF. 2002. Exploring the borderlands of autistic disorder and specific

1307   language impairment: a study using standardised diagnostic instruments. *J Child Psychol*

1308   *Psychiatry.* 43(7):917-929.

1309

1310   Blank I, Kanwisher N, Fedorenko E. 2014. A functional dissociation between language and

1311   multiple-demand systems revealed in patterns of BOLD signal fluctuations. *J Neurophysiol.*

1312   112(5):1105-1118.

1313

1314   Blank I, Balewski Z, Mahowald K, Fedorenko E. 2016. Syntactic processing is distributed across

1315   the language system. *Neuroimage.* 127:307-323.

1316

1317   Blank IA, Fedorenko E. 2017. Domain-general brain regions do not track linguistic input as

1318   closely as language-selective regions. *J Neurosci*. 37(41):9999-10011.

1319

1320   Blank IA, Kiran S, Fedorenko E. 2017. Can neuroimaging help aphasia researchers? Addressing

1321   generalizability, variability, and interpretability. *Cogn Neuropsychol, 34*(6):377-393.

1322

1323   Boebinger D, Norman-Haignere SV, McDermott JH, Kanwisher N. 2021. Music-selective neural

1324   populations arise without musical training. *J Neurophysiol, 125*(6):2237-2263.

1325

1326   Boilès CL. 1973. Reconstruction of proto-melody. *Anuario Interamericano de Investigacion*

1327   *Musical*, 9:45-63.

1328

1329   Bortolini U, Leonard LB, Caselli MC. 1998. Specific Language Impairment in Italian and

1330   English: evaluating alternative accounts of grammatical deficits. *Lang Cogn Process.* 13(1):1-20.

1331

1332   Braga RM, DiNicola LM, Becker HC, Buckner RL. 2020. Situating the left-lateralized language

1333   network in the broader organization of multiple specialized large-scale distributed networks. *J*

1334   *Neurophysiol.* 124(5):1415-1448.

1335

1336   Brown DR. 1991. *Human Universals*. Philadelphia (PA):Temple University Press.

1337

1338   Brust JC. 1980. Music and language: musical alexia and agraphia. *Brain.* 103(2):367-392.

1339

1340   Brysbaert M, Stevens M. 2018. Power analysis and effect size in mixed effects models: A

1341   tutorial. J Cogn. 1(1):9.

1342

1343   Bybee J. 2010. Language, usage and cognition. Cambridge (UK): Cambridge University Press.

1344

1345   Caplan D, Hildebrandt N, Makris N. 1996. Location of lesions in stroke patients with deficits in

1346   syntactic processing in sentence comprehension. *Brain.* 119(3):933-949.

1347

1348   Caplan D, Stanczak L, Waters G. 2008. Syntactic and thematic constraint effects on blood

1349   oxygenation level dependent signal correlates of comprehension of relative clauses. *J Cogn*

1350   *Neurosci.* 20(4):643-656.

1351

1352   Caramazza A, Zurif EB. 1976. Dissociation of algorithmic and heuristic processes in language

1353   comprehension: Evidence from aphasia. *Brain Lang.* 3(4):572-582.

1354

1355   Chen G, Taylor PA, Cox RW. 2017. Is the statistic value all we should care about in

1356   neuroimaging?. *Neuroimage.* 147:952-959.

1357

1358   Chiappetta B, Patel AD, Thompson CK. 2022. Musical and linguistic syntactic processing in

1359   agrammatic aphasia: An ERP study. *J Neurolinguistics.* 62:101043.

1360

1361   Chomsky N. 1965. Aspects of the Theory of Syntax. Cambridge (MA): MIT press.

1362

1363   Chomsky N. 1995. The minimalist program. Cambridge (MA): MIT Press.

1364

1365   Collins T, Tillmann B, Barrett FS, Delbé C, Janata P. 2014. A combined model of sensory and

1366   cognitive representations underlying tonal expectations in music: from audio signals to behavior.

1367   *Psychol Rev, 121*(1):33.

1368

1369   Cooke A, Grossman M, DeVita C, Gonzalez-Atavales J, Moore P, Chen W, Gee J, Detre J. 2006.

1370   Large-scale neural network for sentence processing. *Brain Lang*, 96(1):14-36.

1371

1372   Cooper R. 1973. Propositions pour un modele transformationnel de description musicale.

1373   *Musique en Jeu*. 10:70-88.

1374

1375   Corbetta M, Shulman GL. 2002. Control of goal-directed and stimulus-driven attention in the

1376   brain. *Nat Rev Neurosci.* 3(3):201-215.

1377

1378   Corlett PR, Mollick JA, Kober H. 2021. Substrates of Human Prediction Error for Incentives,

1379   Perception, Cognition, and Action. psyarxiv.

1380

1381   Crawford JR, Garthwaite PH. 2007. Comparison of a single case to a control or normative

1382   sample in neuropsychology: Development of a Bayesian approach. *Cogn Neuropsychol.* 24(4),

1383   343–372.

1384

1385   Creel SC, Weng M, Fu G, Heyman GD, Lee K. 2018. Speaking a tone language enhances

1386   musical pitch perception in 3–5-year-olds. *Dev Sci.* 21(1): e12503.

1387

1388   Crump MJ, McDonnell JV, Gureckis TM. 2013. Evaluating Amazon's Mechanical Turk as a tool

1389   for experimental behavioral research. *PLoS One.* 8(3):e57410.

1390

1391   Cohen AJ. 1982. Exploring the sensitivity to structure in music. Can Univ Music Rev. 3:15-30.

1392

1393   Cuddy LL, Cohen AI, Mewhort DJK. 1981. Perception of structure in short melodic sequences. J Exp

1394   Psychol Hum Percept Perform. 7:869-883.

1395

1396   Cumming G. 2012. *Understanding the new statistics: Effect sizes, confidence intervals, and*

1397   *meta-analysis*. New York (NY): Taylor & Francis.

1398

1399   Curtis ME, Bharucha JJ. 2009. Memory and musical expectation for tones in cultural context. Music

1400   Percept. 26:365–375.

1401

1402   Dale AM. 1999. Optimal experimental design for event-related fMRI. *Hum Brain Mapp.* 8(2-

1403   3):109-114.

1404

1405   Darwin C. 1871. *The Descent of Man, and Selection in Relation to Sex*. London (UK): John

1406   Murray.

1407

1408   Dasgupta I, Gershman SJ. 2021. Memory as a Computational Resource. *Trends Cogn Sci*,

1409   25(3):240-251.

1410

1411   Deen B, Koldewyn K, Kanwisher N, Saxe R. 2015. Functional organization of social perception

1412   and cognition in the superior temporal sulcus. *Cereb Cortex*. 25(11):4596-4609.

1413

1414   Deen B, Freiwald WA. 2021. Parallel systems for social and spatial reasoning within the cortical

1415   apex. *bioRxiv*.

1416

1417   Deutsch D, Henthorn T, Marvin E, Xu H. 2006. Absolute pitch among American and Chinese

1418   conservatory students: Prevalence differences, and evidence for a speech-related critical period. *J*

1419   *Acoust Soc Am*. 119(2):719-722.

1420

1421   Deutsch D, Dooley K, Henthorn T, Head B. 2009. Absolute pitch among students in an

1422   American music conservatory: Association with tone language fluency. *J Acoust Soc Am*.

1423   125(4):2398-2403.

1424

1425   Diachek E*, Blank I*, Siegelman M*, Affourtit J, Fedorenko E. 2020. The domain-general

1426   multiple demand (MD) network does not support core aspects of language comprehension: a

1427   large-scale fMRI investigation. *J Neurosci*. 40(23):4536–4550.

1428

1429 Dick F, Bates E, Wulfeck B, Utman JA, Dronkers N, Gernsbacher MA. 2001. Language deficits,

1430 localization, and grammar: evidence for a distributive model of language breakdown in aphasic

1431 patients and neurologically intact individuals. *Psychol Rev*, 108(4):759-788.

1432

1433 Ding J, Martin RC, Hamilton AC, Schnur TT. (2020). Dissociation between frontal and

1434 temporal-parietal contributions to connected speech in acute stroke. *Brain.* 143(3), 862-876.

1435

1436 Duncan J, Owen AM. 2000. Common regions of the human frontal lobe recruited by diverse

1437 cognitive demands. *Trends Neurosci.* 23(10):475-483.

1438

1439 Duncan J. 2001. An adaptive coding model of neural function in prefrontal cortex. *Nat Rev*

1440 *Neurosci.* 2(11):820-829.

1441

1442 Duncan J. 2010. The multiple-demand (MD) system of the primate brain: mental programs for

1443 intelligent behaviour. *Trends Cogn Sci.* 14(4):172-179.

1444

1445 Duncan J. 2013. The structure of cognition: attentional episodes in mind and brain. *Neuron.*

1446 80(1):35-50.

1447

1448 Embick D, Marantz A, Miyashita Y, O'Neil W, Sakai KL. 2000. A syntactic specialization for

1449 Broca's area. *Proc Natl Acad Sci USA*, 97(11):6150-6154.

1450

1451 Fadiga L, Craighero L, D'Ausilio A. 2009. Broca's area in language, action, and music. *Ann N Y*

1452 *Acad Sci.* 1169(1):448-458.

1453

1454 Fancourt A. 2013. Exploring musical cognition in children with Specific Language Impairment.

1455 Doctoral thesis, Goldsmiths, University of London.

1456

1457 Faroqi-Shah Y, Slevc LR, Saxena S, Fisher SJ, Pifer M. 2020. Relationship between musical and

1458 language abilities in post-stroke aphasia. *Aphasiology.* 34(7):793-819.

1459

1460 Fay T. 1971. Perceived hierarchic structure in language and music. *J Music Theory.* 15(1/2):112-

1461 137.

1462

1463 Fedorenko E, Patel A, Casasanto D, Winawer J, Gibson E. 2009. Structural integration in

1464 language and music: Evidence for a shared system. *Mem Cognit*, 37(1):1-9.

1465

1466 Fedorenko E, Hsieh P-J, Nieto-Castañon A, Whitfield-Gabrieli S, Kanwisher N. 2010. A new

1467 method for fMRI investigations of language: Defining ROIs functionally in individual subjects. *J*

1468 *Neurophysiol.* 104(2):1177-94.

1469

1470 Fedorenko E, Behr M, Kanwisher N. 2011. Functional specificity for high-level linguistic

1471 processing in the human brain. *Proc Natl Acad Sci USA.* 108(39):16428-16433.

1472

1473   Fedorenko E, Duncan J, Kanwisher N. 2012a. Language-selective and domain-general regions

1474   lie side by side within Broca's area. *Curr Biol.* 22(21):2059-2062.

1475

1476   Fedorenko E, Nieto-Castañon A, Kanwisher N. 2012b. Lexical and syntactic representations in

1477   the brain: An fMRI investigation with multi-voxel pattern analyses. *Neuropsychologia.*

1478   50(4):499-513.

1479

1480   Fedorenko E, McDermott J, Norman-Haignere S, Kanwisher N. 2012c. Sensitivity to musical

1481   structure in the human brain. *J Neurophysiol.* 108(12):3289-3300.

1482

1483   Fedorenko E, Duncan J, Kanwisher N. 2013. Broad domain-generality in focal regions of frontal

1484   and parietal cortex. *Proc Natl Acad Sci USA.* 110(41):16616-16621.

1485

1486   Fedorenko E. 2014. The role of domain-general cognitive control in language comprehension.

1487   *Front Psychol.* 5:335.

1488

1489   Fedorenko E, Thompson-Schill SL. 2014. Reworking the language network. *Trends Cogn Sci.*

1490   18(3):120-126.

1491

1492   Fedorenko E, Varley, R. 2016. Language and thought are not the same thing: Evidence from

1493   neuroimaging and neurological patients. *Ann N Y Acad Sci.* 1369(1):132-153.

1494

1495 Fedorenko E, Blank I. 2020. Broca's Area Is Not a Natural Kind. *Trends Cogn Sci*. 24(4):270-

1496 284.

1497

1498 Fedorenko E, Blank I, Siegelman M, Mineroff Z. 2020. Lack of selectivity for syntax relative to

1499 word meanings throughout the language network. *Cognition.* 203:104348.

1500

1501 Fedorenko E. 2020. The brain network that supports high-level language processing. In

1502 Gazzaniga M, Ivry RB, Mangun GR, editors. Cognitive Neuroscience: The Biology of the Mind

1503 (5th edition). New York (NY): WW Norton and Company.

1504

1505 Fedorenko E. 2021. The early origins and the growing popularity of the individual-subject

1506 analytic approach in human neuroscience. *Curr Opin Behav Sci*. 40:105-112.

1507

1508 Fedorenko E, Shain C. 2021. Similarity of computations across domains does not imply shared

1509 implementation: The case of language comprehension. *Curr Dir Psychol Sci*. 30(6):526-534.

1510

1511 Fischl B, Rajendran N, Busa E, Augustinack J, Hinds O, Yeo BT, Mohlberg H, Amunts K, Zilles

1512 K. 2008. Cortical folding patterns and predicting cytoarchitecture. *Cereb Cortex.* 18(8):1973-

1513 1980.

1514

1515 Fitch WT, Martins MD. 2014. Hierarchical processing in music, language, and action: Lashley

1516 revisited. *Ann N Y Acad Sci.* 1316(1):87-104.

1517

Fodor JD. 1983. Phrase structure parsing and the island constraints. *Linguist Philos*. 6(2):163-223.

Fouragnan E, Retzler C, Philiastides MG. 2018. Separate neural representations of prediction error valence and surprise: Evidence from an fMRI meta‑analysis. *Hum Brain Mapp*. 39(7):2887-2906.

Franklin S, Turner JE, Ellis AW. 1992. ADA Comprehension Battery. York (UK): University of York.

Friederici AD, Fiebach CJ, Schlesewsky M, Bornkessel ID, Von Cramon DY. 2006. Processing linguistic complexity and grammaticality in the left frontal cortex. *Cereb Cortex*. 16(12):1709-1717.

Friederici AD, Kotz SA, Scott SK, Obleser J. 2010. Disentangling syntax and intelligibility in auditory language comprehension. *Hum Brain Mapp*. 31(3):448-457.

Friederici AD. 2011. The brain basis of language processing: from structure to function. *Physiol Rev*. 91(4):1357-1392.

Friederici AD. 2018. The neural basis for human syntax: Broca's area and beyond. *Curr Opin Behav Sci*. 21:88-92.

1541   Frost MA, Goebel R. 2012. Measuring structural–functional correspondence: spatial variability

1542   of specialised brain regions after macro-anatomical alignment. *Neuroimage* 59(2):1369-1381.

1543

1544   Giesbrecht F, Burns J. 1985. Two-Stage Analysis Based on a Mixed Model: Large-Sample

1545   Asymptotic Theory and Small-Sample Simulation Results. *Biometrics. 41*(2):477-486.

1546

1547   Goldberg AE. 2002. Construction Grammar. In: Nadel L, editor. *Encyclopedia of Cognitive*

1548   *Science*. Stuttgart (Germany): Macmillan.

1549

1550   Green DM, Swets JA. 1966. *Signal detection theory and psychophysics*. New York (NY): Wiley.

1551

1552   Guenther FH. 2016. Neural control of speech. Cambridge (MA): MIT Press.

1553

1554   Hagoort P, Indefrey P. 2014. The neurobiology of language beyond single words. *Ann Rev*

1555   *Neurosci, 37*:347-362.

1556

1557   Hasson U, Chen J, Honey CJ. 2015. Hierarchical process memory: memory as an integral

1558   component of information processing. *Trends Cogn Sci. 19*(6):304-313.

1559

1560   Herholz SC, Zatorre RJ. 2012. Musical training as a framework for brain plasticity: behavior,

1561   function, and structure. *Neuron. 76*(3):486-502.

1562

1563   Herrmann B, Obleser J, Kalberlah C, Haynes JD, Friederici AD. 2012. Dissociable neural

1564   imprints of perception and grammar in auditory functional imaging. *Hum Brain Mapp*.

1565   33(3):584-595.

1566

1567   Hoch L, Poulin-Charronnat B, Tillmann B. 2011. The influence of task-irrelevant music on

1568   language processing: syntactic and semantic structures. *Front Psychol*. 2:112.

1569

1570   Hrong-Tai Fai A, Cornelius PL. 1996. Approximate F-tests of multiple degree of freedom

1571   hypotheses in generalized least squares analyses of unbalanced split-plot experiments. *J Stat*

1572   *Comput Simul. 54*(4):363-378.

1573

1574   Ivanova A, Srikant S, Sueoka Y, Kean H, Dhamala R, O'Reilly U-M, Bers MU, Fedorenko E.

1575   2020. Comprehension of computer code relies primarily on domain-general executive resources.

1576   *eLife*. 9:e58906.

1577

1578   Ivanova A, Mineroff Z, Zimmerer V, Kanwisher N, Varley R, Fedorenko E. 2021. The language

1579   network is recruited but not required for non-verbal semantic processing. bioRxiv.

1580

1581   Jackendoff R. 2002. English particle constructions, the lexicon, and the autonomy of syntax. In

1582   Dehé N, Jackendoff R, McIntyre A, Urban S, editors. *Verb-particle explorations*. Berlin

1583   (Germany): De Gruyter. p. 67-94.

1584

1585   Jackendoff R. 2007. A parallel architecture perspective on language processing. *Brain Res.*

1586   1146:2-22.

1587

1588   Jackendoff R. 2009. Parallels and nonparallels between language and music. *Music Percept.*

1589   26(3):195-204.

1590

1591   Jackendoff R, Audring J. 2020. The texture of the lexicon: relational morphology and the parallel

1592   architecture. Oxford (UK): Oxford University Press.

1593

1594   Janata P. 1995. ERP measures assay the degree of expectancy violation of harmonic contexts in

1595   music. *J Cogn Neurosci.* 7(2):153-164.

1596

1597   Jentschke S, Koelsch S, Sallat S, Friederici AD. 2008. Children with specific language

1598   impairment also show impairment of music-syntactic processing. *J Cogn Neurosci. 20*(11):1940-

1599   1951.

1600

1601   Jouravlev O, Zheng D, Balewski Z, Pongos A, Levan Z, Goldin-Meadow S, Fedorenko E. 2019.

1602   Speech-accompanying gestures are not processed by the language-processing mechanisms.

1603   *Neuropsychologia.* 132:107132.

1604

1605   Jouravlev O, Kell A, Mineroff Z, Haskins AJ, Ayyash D, Kanwisher N, Fedorenko E. 2020.

1606   Reduced language lateralization in autism and the broader autism phenotype as assessed with

1607   robust individual-subjects analyses. *Autism Res.* 13(10):1746-1761.

1608

1609   Just MA, Carpenter PA, Keller TA, Eddy WF, Thulborn KR. 1996. Brain activation modulated

1610   by sentence comprehension. *Science*. 274(5284):114-116.

1611

1612   Kaplan E, Goodglass H, Weintraub S. 2001. Boston Naming Test. 2nd Ed. Philadelphia (PA):

1613   Lippincott Williams & Wilkins.

1614

1615   Kay J, Lesser R, Coltheart M. 1992. Psycholinguistic Assessments of Language Processing in

1616   Aphasia (PALPA). Hove (UK): Lawrence Erlbaum.

1617

1618   Kell AJ, Yamins DL, Shook EN, Norman-Haignere SV, McDermott JH. 2018. A task-optimized

1619   neural network replicates human auditory behavior, predicts brain responses, and reveals a

1620   cortical processing hierarchy. *Neuron*. 98(3):630-644.

1621

1622   Keller TA, Carpenter PA, Just MA. 2001. The neural bases of sentence comprehension: a fMRI

1623   examination of syntactic and lexical processing. *Cereb Cortex*. 11(3):223-237.

1624

1625   Koelsch S, Gunter T, Friederici AD, Schröger E. 2000. Brain indices of music processing:

1626   "nonmusicians" are musical. *J Cogn Neurosci.* 12(3):520-541.

1627

1628   Koelsch S, Gunter TC, Schröger E, Tervaniemi M, Sammler D, Friederici AD. 2001.

1629   Differentiating ERAN and MMN: an ERP study. *NeuroReport*. 12(7):1385-1389.

1630

1631   Koelsch S, Gunter TC, von Cramon DY, Zysset S, Lohmann G, Friederici AD. 2002. Bach

1632   speaks: A cortical" language-network" serves the processing of music. *Neuroimage.* 17(2):956-

1633   966.

1634

1635   Koelsch S. 2006. Significance of Broca's area and ventral premotor cortex for music-syntactic

1636   processing. *Cortex.* 42(4):518-520.

1637

1638   Koelsch S, Jentschke S, Sammler D, Mietchen D. 2007. Untangling syntactic and sensory

1639   processing: An ERP study of music perception. *Psychophysiology.* 44(3):476-490.

1640

1641   Koelsch S, Rohrmeier M, Torrecuso R, Jentschke S. 2013. Processing of hierarchical syntactic

1642   structure in music. *Proc Natl Acad Sci USA.* 110(38):15443-15448.

1643

1644   Kriegeskorte N, Simmons WK, Bellgowan PS, Baker CI. 2009. Circular analysis in systems

1645   neuroscience: the dangers of double dipping. *Nat Neurosci.* 12(5):535.

1646

1647   Krumhansl CL, Keil FC. 1982. Acquisition of the hierarchy of tonal functions in music. *Mem*

1648   *Cognit.* 10(3):243-251.

1649

1650   Kunert R, Slevc LR. 2015. A Commentary on:"Neural overlap in processing music and speech".

1651   *Front Hum Neurosci.* 9:330.

1652

1653   Kunert R, Willems RM, Casasanto D, Patel AD, Hagoort P. 2015. Music and language syntax

1654   interact in Broca's area: An fMRI study. *PLoS One.* 10(11):e0141069.

1655

1656   Kunert R, Willems RM, Hagoort P. 2016. Language influences music harmony perception:

1657   effects of shared syntactic integration resources beyond attention. *R Soc Open Sci.* 3(2):150685.

1658

1659   Kuperberg GR, Holcomb PJ, Sitnikova T, Greve D, Dale AM, Caplan D. 2003. Distinct patterns

1660   of neural modulation during the processing of conceptual and syntactic anomalies. *J Cogn*

1661   *Neurosci.* 15(2):272-293.

1662

1663   Kuznetsova A, Brockhoff PB, Christensen RH. 2017. lmerTest package: tests in linear mixed

1664   effects models. *J Stat Softw.* 82(13):1-26.

1665

1666   LaCroix A, Diaz AF, Rogalsky C. 2015. The relationship between the neural computations for

1667   speech and music perception is context-dependent: an activation likelihood estimate study. *Front*

1668   *Psychol.* 6:1138.

1669

1670   Lartillot O, Grandjean D. 2019. Tempo and metrical analysis by tracking multiple metrical levels

1671   using autocorrelation. *Appl Sci. 9*(23):5121.

1672

1673   Lartillot O, Toiviainen P. 2007. September). A Matlab toolbox for musical feature extraction

1674   from audio. In: Proceedings of the 10th International Conference on Digital Audio Effects; 2007 Sep 10-

1675   15; Bordeaux, France. p. 244.

1676

Lecours A, Joanette Y. 1980. Linguistic and other psychological aspects of paroxysmal aphasia.

Brain Lang. 10(1):1-23.

1679

Lerdahl F, Jackendoff R. 1977. Toward a formal theory of tonal music. *J Music Theory.*

21(1):111-171.

1682

Lerdahl F, Jackendoff R. 1983. An overview of hierarchical structure in music. *Music Percept.*

1(2):229-252.

1685

Levin B, Rappaport-Hovav M. 2005. Argument realization. Cambridge (UK): Cambridge

University Press.

1688

Levitin DJ, Menon V. 2003. Musical structure is processed in "language" areas of the brain: a

possible role for Brodmann Area 47 in temporal coherence. *Neuroimage.* 20(4):2142-2152.

1691

Linebarger MC, Schwartz MF, Saffran EM. 1983. Sensitivity to grammatical structure in so-

called agrammatic aphasics. *Cognition.* 13(3):361-392.

1694

Lindblom B, Sundberg J. 1969. Towards a generative theory of melody. Speech Transmission

Laboratory. *Quarterly Progress and Status Reports.* 10:53-86.

1697

1698   Lipkin B, Tuckute G, Affourtit J, Small H, Mineroff Z, Jouravlev O, Rakocevic L, Pritchett B, et al.

1699   2022. Probabilistic atlas for the language network based on precision fMRI data from> 800

1700   individuals. *Sci Data. 9*(1):1-10.

1701

1702   Liu YF, Kim J, Wilson C, Bedny M. 2020. Computer code comprehension shares neural

1703   resources with formal logical inference in the fronto-parietal network. *Elife. 9*:e59340.

1704

1705   Liu J, Hilton CB, Bergelson E, Mehr SA. 2021. Language experience shapes music processing

1706   across 40 tonal, pitch-accented, and non-tonal languages. *bioRxiv*.

1707

1708   Luria AR, Tsvetkova LS, Futer DS. 1965. Aphasia in a composer. *J Neurol Sci.* 2(3):288-292.

1709

1710   Maess B, Koelsch S, Gunter TC, Friederici AD. 2001. Musical syntax is processed in Broca's

1711   area: an MEG study. *Nat Neurosci.* 4(5):540-545.

1712

1713   Mahowald K, Fedorenko E. 2016. Reliable individual-level neural markers of high-level

1714   language processing: A necessary precursor for relating neural variability to behavioral and

1715   genetic variability. *Neuroimage.* 139:74-93.

1716

1717   Makowski D. 2018. The psycho Package: An Efficient and Publishing-Oriented Workflow for

1718   Psychological Science. *J Open Source Softw.* 3(22):470.

1719

1720   Malik-Moraleda S, Ayyash D, Gallée J, Affourtit J, Hoffmann M, Mineroff Z, Jouravlev O,

1721   Fedorenko E. 2022. An investigation across 45 languages and 12 language families reveals a

1722   universal language network. *Nat Neurosci. 25*(8):1014-1019.

1723

1724   Marin OSM. 1982. Neurological Aspects of Music Perception and Performance. New York

1725   (NY): Academic Press.

1726

1727   Matchin W, Hickok G. 2020. The cortical organization of syntax. *Cereb Cortex.* 30(3):1481-

1728   1498.

1729

1730   Mehr SA, Krasnow MM. 2017. Parent-offspring conflict and the evolution of infant-directed

1731   song. Evol Hum Behav. 38(5):674-684.

1732

1733   Mehr SA, Singh M, Knox D, Ketter DM, Pickens-Jones D, Atwood S, Lucas C, Jacoby N, Egner

1734   AA, Hopkins EJ, et al. (2019). Universality and diversity in human song. *Science.*

1735   *366*(6468):eaax0868.

1736

1737   Mehr S, Krasnow M, Bryant G, Hagen E. 2020. Origins of music in credible signaling. *Behav*

1738   *Brain Sci.* 44:e60.

1739

1740   Mesulam MM, Rogalski EJ, Wieneke C, Hurley RS, Geula C, Bigio EH, Thompson CK,

1741   Weintraub S. 2014. Primary progressive aphasia and the evolving neurology of the language

1742   network. *Nat Rev Neurol.* 10(10):554.

1743

1744   Mesulam MM, Thompson CK, Weintraub S, Rogalski EJ. 2015. The Wernicke conundrum and

1745   the anatomy of language comprehension in primary progressive aphasia. *Brain*. 138(8):2423-

1746   2437.

1747

1748   McDermott J. 2008. The evolution of music. *Nature*. 453(7193):287-288.

1749

1750   Mineroff Z*, Blank I*, Mahowald K, Fedorenko E. 2018. A robust dissociation among the

1751   language, multiple demand, and default mode networks: evidence from inter-region correlations

1752   in effect size. *Neuropsychologia*. 119:501-511.

1753

1754   Mollica F, Shain C, Affourtit J, Kean H, Siegelman M, Fedorenko E. 2020. Another look at the

1755   constituent structure of sentences in the human brain [Poster presentation]. SNL 2020; October

1756   21-24; virtual.

1757

1758   Monti MM, Parsons LM, Osherson DN. 2009. The boundaries of language and thought in

1759   deductive inference. *Proc Natl Acad Sci USA*. 106(30):12554-12559.

1760

1761   Monti MM, Parsons LM, Osherson DN. 2012. Thought beyond language: Neural dissociation of

1762   algebra and natural language. *Psychol Sci.* 23(8):914-922.

1763

1764   Morosan P, Rademacher J, Schleicher A, Amunts K, Schormann T, Zilles K. 2001. Human

1765   primary auditory cortex: cytoarchitectonic subdivisions and mapping into a spatial reference

1766   system. *Neuroimage*. 13(4):684-701.

1767

1768   Musso M, Weiller C, Horn A, Glauche V, Umarova R, Hennig J, Schneider A, Rijntjes M. 2015.

1769   A single dual-stream framework for syntactic computations in music and language. *Neuroimage*.

1770   117:267-283.

1771

1772   Nettl B. 2015. *The study of ethnomusicology: Thirty-three discussions*. Champaign (IL):

1773   University of Illinois Press.

1774

1775   Newman AJ, Pancheva R, Ozawa K, Neville HJ, Ullman MT. 2001. An event-related fMRI

1776   study of syntactic and semantic violations. *J Psycholinguist Res*. 30(3):339-364.

1777

1778   Ngo MK, Vu KPL, Strybel TZ. 2016. Effects of music and tonal language experience on relative

1779   pitch performance. *Am J Psychol*. 129(2):125-134.

1780

1781   Nieto-Castañon A, Fedorenko E. 2012. Subject-specific functional localizers increase sensitivity

1782   and functional resolution of multi-subject analyses. *Neuroimage*. 63(3):1646-1669.

1783

1784   Norman-Haignere S, Kanwisher N, McDermott JH. 2013. Cortical pitch regions in humans

1785   respond primarily to resolved harmonics and are located in specific tonotopic regions of anterior

1786   auditory cortex. *J Neurosci*. 33(50):19451-19469.

1787

1788 Norman-Haignere S, Kanwisher NG, McDermott JH. 2015. Distinct cortical pathways for music

1789 and speech revealed by hypothesis-free voxel decomposition. *Neuron*. 88(6):1281-1296.

1790

1791 Norman-Haignere SV, Feather J, Boebinger D, Brunner P, Ritaccio A, McDermott JH, Schalk G,

1792 Kanwisher N. 2022. A neural population selective for song in human auditory cortex. *Curr Biol.*

1793 *32*(7):1470-1484.

1794

1795 Oldfield RC. 1971. The assessment and analysis of handedness: the Edinburgh inventory.

1796 *Neuropsychologia*. 9(1):97-113.

1797

1798 Omigie D, Samson S. 2014. A protective effect of musical expertise on cognitive outcome

1799 following brain damage?. *Neuropsychol Rev. 24*(4):445-460.

1800

1801 Overath T, McDermott JH, Zarate JM, Poeppel D. 2015. The cortical analysis of speech-specific

1802 temporal structure revealed by responses to sound quilts. *Nat Neurosci. 18*(6):903-911.

1803

1804 Pallier C, Devauchelle AD, Dehaene S. 2011. Cortical representation of the constituent structure

1805 of sentences. *Proc Natl Acad Sci USA*. 108(6):2522-2527.

1806

1807 Patel AD, Gibson E, Ratner J, Besson M, Holcomb PJ. 1998. Processing syntactic relations in

1808 language and music: An event-related potential study. *J Cogn Neurosci. 10*(6):717-733.

1809

1810   Patel AD. 2003. Language, music, syntax and the brain. *Nat Neurosci.* 6(7):674-681.

1811

1812   Patel AD. 2008. Music, Language, and the Brain. Oxford (UK): Oxford University Press.

1813

1814   Patel AD, Iversen JR, Wassenaar M, Hagoort P. 2008a. Musical syntactic processing in

1815   agrammatic Broca's aphasia. *Aphasiology*. 22(7-8):776-789.

1816

1817   Patel AD, Wong M, Foxton J, Lochy A, Peretz I. 2008b. Speech intonation perception deficits in

1818   musical tone deafness (congenital amusia). *Music Percept. 25*(4):357-368.

1819

1820   Patel AD. 2012. Language, music, and the brain: a resource-sharing framework. In: Rebuschat P,

1821   Rohrmeier M, Hawkins J, Cross I, editors. *Language and Music as Cognitive Systems*. Oxford

1822   (UK): Oxford University Press. p. 204-223.

1823

1824   Patel AD, Morgan E. 2017. Exploring cognitive relations between prediction in language and

1825   music. *Cogn Sci. 41*:303-320.

1826

1827   Patterson RD, Uppenkamp S, Johnsrude IS, Griffiths TD. 2002. The processing of temporal pitch

1828   and melody information in auditory cortex. *Neuron*. 36(4):767-776.

1829

1830   Paunov A, Blank IA, Fedorenko E. 2019. Functionally distinct language and Theory of Mind

1831   networks are synchronized at rest and during language comprehension. *J Neurophysiol.*

1832   121:1244-1265.

1833

1834   Paunov AM, Blank IA, Jouravlev O, Mineroff Z, Gallée J, Fedorenko E. 2022. Differential

1835   tracking of linguistic vs. mental state content in naturalistic stimuli by language and Theory of

1836   Mind (ToM) brain networks. *Neurobiol Lang. 3*(3):413-440.

1837

1838   Peelle JE, Troiani V, Wingfield A, Grossman M. 2010. Neural processing during older adults'

1839   comprehension of spoken sentences: age differences in resource allocation and connectivity.

1840   *Cereb Cortex*. 20(4):773-782.

1841

1842   Penagos H, Melcher JR, Oxenham AJ. 2004. A neural representation of pitch salience in

1843   nonprimary human auditory cortex revealed with functional magnetic resonance imaging. *J

1844   Neurosci.* 24(30):6810-6815.

1845

1846   Peretz I. 1990. Processing of local and global musical information by unilateral brain-damaged

1847   patients. *Brain*. 113(4):1185-1205.

1848

1849   Peretz I, Kolinsky R, Tramo M, Labrecque R, Hublet C, Demeurisse G, Belleville S. 1994.

1850   Functional dissociations following bilateral lesions of auditory cortex. *Brain*. 117(6):1283-1301.

1851

1852   Peretz I, Belleville S, Fontaine S. 1997. Dissociations between music and language functions

1853   after cerebral resection: a new case of amusia without aphasia. *Can J Exp Psychol* 51(4):354-368.

1854

1855   Peretz I, Champod AS, Hyde K. 2003. Varieties of musical disorders: the Montreal Battery of

1856   Evaluation of Amusia. *Ann N Y Acad Sci.* 999(1):58-75.

1857

1858   Peretz I, Coltheart M. 2003. Modularity of music processing. *Nat Neurosci.* 6(7):688-691.

1859

1860   Peretz I, Vuvan D, Lagrois MÉ, Armony JL. 2015. Neural overlap in processing music and

1861   speech. *Philos Trans R Soc Lond B Biol Sci.* 370(1664):20140090.

1862

1863   Perrachione TK, Fedorenko EG, Vinke L, Gibson E, Dilley LC. 2013. Evidence for shared

1864   cognitive processing of pitch in music and language. *PLoS One.* 8(8):e73372.

1865

1866   Perruchet P, Poulin-Charronnat B. 2013. Challenging prior evidence for a shared syntactic

1867   processor for language and music. *Psychon Bull Rev.* 20(2):310-317.

1868

1869   Piccirilli M, Sciarma T, Luzzi S. 2000. Modularity of music: evidence from a case of pure

1870   amusia. *J Neurol Neurosurg Psychiatry*. 69(4):541-545.

1871

1872   Pinker S, Prince A. 1988. On language and connectionism: Analysis of a parallel distributed

1873   processing model of language acquisition. *Cognition*. 28(1-2):73-193.

1874

1875   Pinker S. 1991. Rules of language. *Science*. 253(5019):530-535.

1876

1877   Pinker S. 1994. The Language Instinct: How the Mind Creates Language, New York (NY):

1878   Harper Collins Publishers, Inc.

1879

1880   Pinker S. 1999. Out of the minds of babes. *Science*. 283(5398):40-41.

1881

1882   Poldrack RA. 2006. Can cognitive processes be inferred from neuroimaging data?. *Trends Cogn*

1883   *Sci*. 10(2):59-63.

1884

1885   Poldrack RA. 2011. Inferring mental states from neuroimaging data: from reverse inference to

1886   large-scale decoding. *Neuron*. 72(5):692-697.

1887

1888   Polk M, Kertesz A. 1993. Music and language in degenerative disease of the brain. *Brain Cogn*.

1889   22(1):98-117.

1890

1891   Poulin-Charronnat B, Bigand E, Madurell F, Peereman R. 2005. Musical structure modulates

1892   semantic priming in vocal music. *Cognition*. 94:B67-B78.

1893

1894   Pritchett B, Hoeflin C, Koldewyn K, Dechter E, Fedorenko E. 2018. High-level language

1895   processing regions are not engaged in action observation or imitation. *J Neurophysiol*.

1896   120(5):2555-2570.

1897

1898   Riemann H. 1877. Musikalische Syntaxis: Grundriss einer harmonischen Satzbildungslehre.

1899   Leipzig: Breitkopf und Härtel.

1900

1901   Roads C, Wieneke P. 1979. Grammars as representations for music. *Comput Music J.* 3(1):48-55.

1902

1903   Roberts I. 2012. Comments and a conjecture inspired by Fabb and Halle. In: Rebuschat P,

1904   Rohrmeier M, Hawkins JA, Cross I, editors. *Language and Music as Cognitive Systems.* Oxford:

1905   Oxford University Press. p. 51-66.

1906

1907   Röder B, Stock O, Neville H, Bien S, Rösler F. 2002. Brain activation modulated by the

1908   comprehension of normal and pseudo-word sentences of different processing demands: a

1909   functional magnetic resonance imaging study. *Neuroimage.* 15(4):1003-1014.

1910

1911   Rogalsky C, Hickok G. 2011. The role of Broca's area in sentence comprehension. *J Cogn*

1912   *Neurosci.* 23(7):1664-1680.

1913

1914   Rueffler C, Hermisson J, Wagner GP. 2012. Evolution of functional specialization and division

1915   of labor. *Proc Natl Acad Sci USA.* 109(6):E326-E335.

1916

1917   Sag I, Wasow T, Bender E. 2003. Formal syntax, an introduction. CSLI publication.

1918

1919   Sammler D, Koelsch S, Ball T, Brandt A, Elger CE, Friederici AD, Grigutsch M, Huppertz H-J,

1920   Knosche TR, Wellmer J, Widman G, Schulze-Bonhaged A. 2009. Overlap of musical and

1921   linguistic syntax processing: intracranial ERP evidence. *Ann N Y Acad Sci.* 1169(1):494-498.

1922

1923 Sammler D, Koelsch S, Friederici AD. 2011. Are left fronto-temporal brain areas a prerequisite

1924 for normal music-syntactic processing?. *Cortex*. 47(6):659-673.

1925

1926 Sammler D, Koelsch S, Ball T, Brandt A, Grigutsch M, Huppertz HJ, Wellmer J, Widman G,

1927 Elger CE, Friederici AD, Schulze-Bonhaged A. 2013. Co-localizing linguistic and musical

1928 syntax with intracranial EEG. *Neuroimage*. 64:134-146.

1929

1930 Savage PE, Loui P, Tarr B, Schachner A, Glowacki L, Mithen S, Fitch WT. 2021. Music as a

1931 coevolved system for social bonding. *Behav Brain Sci*. 44(e59):1-22.

1932

1933 Schmidt S. 2009. Shall We Really Do It Again? The Powerful Concept of Replication Is

1934 Neglected in the Social Sciences. *Rev Gen Psychol*. 13(2):90-100.

1935

1936 Scott TL, Gallée J, Fedorenko E. 2017. A new fun and robust version of an fMRI localizer for

1937 the frontotemporal language system. *Cogn Neurosci*. 8(3):167-176.

1938

1939 Shain C*, Blank I*, Van Shijndel M, Schuler W, Fedorenko E. 2020. fMRI reveals language-

1940 specific predictive coding during naturalistic sentence comprehension. *Neuropsychologia*.

1941 138:107307.

1942

1943 Shain C, Kean H, Lipkin B, Affourtit J, Siegelman M, Mollica F, Fedorenko E. 2021.

1944 'Constituent length'effects in fMRI do not provide evidence for abstract syntactic processing.

1945 *bioRxiv*.

1946

1947 Shain C, Blank IA, Fedorenko E, Gibson E, Schuler W. 2022. Robust effects of working

1948 memory demand during naturalistic language comprehension in language-selective cortex. *J*

1949 *Neurosci. 42*(39):7412-7430.

1950

1951 Shashidhara S, Mitchell DJ, Erez Y, Duncan J. 2019. Progressive recruitment of the

1952 frontoparietal multiple-demand system with increased task complexity, time pressure, and

1953 reward. *J Cogn Neurosci.* 31(11):1617-1630.

1954

1955 Sihvonen AJ, Särkämö T, Leo V, Tervaniemi M, Altenmüller E, Soinila S. 2017. Music-based

1956 interventions in neurological rehabilitation. *Lancet Neurol. 16*(8):648-660.

1957

1958 Slevc LR, Rosenberg JC, Patel AD. 2009. Making psycholinguistics musical: Self-paced reading

1959 time evidence for shared processing of linguistic and musical syntax. *Psychon Bull Rev.*

1960 16(2):374-381.

1961

1962 Slevc LR, Reitman J, Okada B. 2013. Syntax in music and language: the role of cognitive

1963 control. In: Proceedings of the Annual Meeting of the Cognitive Science Society; 2013 Jul 31-Aug

1964 3; Berlin, Germany; p. 3414-3419.

1965

1966 Slevc LR, Okada BM. 2015. Processing structure in language and music: a case for shared

1967 reliance on cognitive control. *Psychon Bull Rev.* 22(3):637-652.

1968

1969   Slevc LR, Faroqi-Shah Y, Saxena S, Okada BM. 2016. Preserved processing of musical structure

1970   in a person with agrammatic aphasia. *Neurocase*. 22(6):505-511.

1971

1972   Stromswold K, Caplan D, Alpert N, Rauch S. 1996. Localization of syntactic comprehension by

1973   positron emission tomography. *Brain Lang.* 52(3):452-473.

1974

1975   Sueoka Y, Paunov A, Ivanova A, Blank IA, Fedorenko E. 2022. The language network reliably

1976   'tracks' naturalistic meaningful non-verbal stimuli. bioRxiv

1977

1978   Sullivan GM, Feinn R. 2012. Using effect size—or why the P value is not enough. *J Grad Med*

1979   *Educ. 4*(3):279-282.

1980

1981   Sundberg J, Lindblom B. 1976. Generative theories in language and music descriptions.

1982   *Cognition*. 4(1):99-122.

1983

1984   Swain JP. 1995. The concept of musical syntax. *Music Q.* 79(2):281-308.

1985

1986   Tahmasebi AM, Davis MH, Wild CJ, Rodd JM, Hakyemez H, Abolmaesumi P, Johnsrude IS.

1987   2012. Is the link between anatomical structure and function equally strong at all cognitive levels

1988   of processing?. *Cereb Cortex.* 22(7):1593-1603.

1989

1990   Tarantola A. 2005. *Inverse problem theory and methods for model parameter estimation*.

1991   Philadelphia (PA): Society for Industrial and Applied Mathematics.

1992

1993   Temperley D. 2022. Music and Language. Annu Rev Linguist. 8:153-170.

1994

1995   te Rietmolen NA, Mercier M, Trebuchon A, Morillon B, Schon D. 2022. Speech and music

1996   recruit frequency-specific distributed and overlapping cortical networks. *bioRxiv.*

1997

1998   Tillmann B, Janata P, Bharucha JJ. 2003. Activation of the inferior frontal cortex in musical

1999   priming. *Cogn Brain Res.* 16(2):145-161.

2000

2001   Tillmann B, Koelsch S, Escoffier N, Bigand E, Lalitte P, Friederici AD, von Cramon DY. 2006.

2002   Cognitive priming in sung and instrumental music: activation of inferior frontal cortex.

2003   *Neuroimage.* 31(4):1771-1782.

2004

2005   Tillmann B. 2012. Music and Language Perception: Expectations, Structural Integration, and

2006   Cognitive Sequencing. *Top Cogn Sci.* 4(4):568-584.

2007

2008   Trehub SE. 2003. The developmental origins of musicality. *Nat Neurosci. 6*(7):669-673.

2009

2010   Tyler LK, Marslen-Wilson WD, Randall B, Wright P, Devereux BJ, Zhuang J, Papoutsi M,

2011   Stamatakis EA. 2011. Left inferior frontal cortex and syntax: function, structure and behaviour in

2012   patients with left hemisphere damage. *Brain.* 134(2):415-431.

2013

Van de Cavey J, Hartsuiker RJ. 2016. Is there a domain-general cognitive structuring system? Evidence from structural priming across music, math, action descriptions, and language. *Cognition.* 146:172-184.

Varley R, Siegal M. 2000. Evidence for cognition without grammar from causal reasoning and 'theory of mind' in an agrammatic aphasic patient. *Curr Biol.* 10(12):723-726.

Varley RA, Klessinger NJ, Romanowski CA, Siegal M. 2005. Agrammatic but numerate. *Proc Natl Acad Sci USA.* 102(9):3519-3524.

Vázquez-Rodríguez B, Suárez LE, Markello RD, Shafiei G, Paquola C, Hagmann P, van den Heuvel MP, Bernhardt BC, Spreng RN, Misic B. 2019. Gradients of structure–function tethering across neocortex. *Proc Natl Acad Sci USA.* 116(42):21219-21227.

Vuust P, Heggli OA, Friston KJ, Kringelbach ML. 2022. Music in the brain. *Nat Rev Neurosci.* *23*(5):287-305.

Wehbe L, Blank I, Shain C, Futrell R, Levy R, Malsburg T, Smith N, Gibson E, Fedorenko E. 2021. Incremental language comprehension difficulty predicts activity in the language network but not the multiple demand network. Cereb Cortex. 31(9):4006-4023.

Westfall J, Kenny DA, Judd CM. 2014. Statistical power and optimal design in experiments in which samples of participants respond to samples of stimuli. J Exp Psychol. 143(5):2020-2045.

2037

2038   Willems RM, Van der Haegen L, Fisher SE, Francks C. 2014. On the other hand: including left-

2039   handers in cognitive neuroscience and neurogenetics. *Nat Rev Neurosci.* 15(3):193-201.

2040

2041   Wilson SM, Saygın AP. 2004. Grammaticality judgment in aphasia: Deficits are not specific to

2042   syntactic structures, aphasic syndromes, or lesion sites. *J Cogn Neurosci.* 16(2):238-252.

2043

2044   Wilson SM, Galantucci S, Tartaglia MC, Gorno-Tempini ML. 2012. The neural basis of

2045   syntactic deficits in primary progressive aphasia. *Brain Lang.* 122(3):190-198.

2046

2047   Wong PC, Perrachione TK. 2007. Learning pitch patterns in lexical identification by native

2048   English-speaking adults. *Appl Psycholinguist.* 28(4):565-585.

2049

2050   Woolgar A, Duncan J, Manes F, Fedorenko E. 2018. Fluid intelligence is supported by the

2051   multiple-demand system not the language system. *Nat Hum Behav.* 2(3):200-204.

2052

2053   Zatorre RJ. 1984. Musical perception and cerebral function: A critical review. *Music Percept.*

2054   2(2):196-221.