A multi-tiered map of EMT defines major transition points and identifies vulnerabilities

Indranil Paul[1], Dante Bolzan[3], Heather Hook[4], Ahmed Youssef[1], Gopal Karemore[5], Michael UJ Oliphant[6], Weiwei Lin[1], Qian Liu[7], Sadhna Phanse[1], Carl White[1], Dzmitry Padhorny[10], Sergei Kotelnikov[10], Guillaume Andrieu[8], Pingzhao Hu[7], Gerald V. Denis[9], Dima Kozakov[10], Brian Raught[12], Trevor Siggers[4], Stefan Wuchty[3,11†], Senthil Muthuswamy[6†], Andrew Emili[1,2†*]

Affiliations

[1] Department of Biochemistry, Boston University School of Medicine, Boston University, 72 East Concord Street, Boston, MA 02118, USA

[2] Department of Biology, Charles River Campus, Boston University, Life Science & Engineering (LSEB-602), 24 Cummington Mall, Boston, MA 02215 USA

[3] Department of Computer Science, University of Miami, 1365 Memorial Drive, Coral Gables, FL 33146, USA

[4] (a) Department of Biology, Boston University, 24 Cummington Mall, Boston, MA 02115, USA; (b) Biological Design Center, Boston University, 610 Commonwealth Avenue, Boston, MA 02215

[5] Advanced Analytics, Novo Nordisk A/S, 2760 Måløv, Denmark

[6] Cancer Research Institute, Department of Medicine, Beth Israel Deaconnes Medical Center, Boston MA 02115, USA

[7] Department of Biochemistry and Medical Genetics, University of Manitoba, Winnipeg, Manitoba R3E 0J9, Canada

[8] Team 2 - Institut Necker-Enfants Malades, INSERM U1151, Paris, France

[9] Boston Medical Center Cancer Center, Boston University, Boston University, 72 East Concord Street, Boston, MA 02118, USA

[10] (a) Department of Applied Mathematics and Statistics, Stony Brook University, 11794 Stony Brook, NY. (b) Laufer Center for Physical and Quantitative Biology, Stony Brook University, Stony Brook, New York, 11794, United States

[11] (a) Department of Biology, University of Miami, 1301 Memorial Drive, Coral Gables, FL 33146, USA; (b) Sylvester Comprehensive Cancer Center, 1475 NW 12th Ave, Miami, FL 33136, USA

[12] Discovery Tower (TMDT), 101 College St, Rm. 9-701A, University of Toronto, Toronto, Ontario, M5G 1L7, Canada

† Co-communicating authors

33   *Correspondence to: aemili@bu.edu

Summary

TGFβ mediated epithelial to mesenchymal transition (EMT) proceeds through hybrid 'E/M' states. A deeper understanding of these states and events which regulate entry to and exit from the E/M states is needed for therapeutic exploitation. We quantified >60,000 molecules across ten time points and twelve omic layers in mammary epithelial cells. Proteomes of whole cells, phosphoproteins, nucleus, extracellular vesicles, secretome and membrane resolved major shifts, E→E/M and E/M→M during EMT, and defined state-specific signatures. Metabolomics identified early activation of arachidonic acid pathway and an enzyme-mediated switch from Cytochrome P450 to Cyclooxygenase / Lipoxygenase branches during E→E/M. Single-cell transcriptomics identified GLIS2 as an early modulator of EMT. Integrative modeling-predicted combinatorial inhibition of AURKB, PP2A and SRC exposed vulnerabilities at E→E/M juncture. Covariance analysis revealed remarkable discordance between proteins and transcripts, and between proteomic layers, implying insufficiency of current approaches. Overall, this dataset provides an unprecedented resource on TGFβ signaling, EMT and cancer.

Short title

Multi-level time-resolved deconstruction of EMT program

Introduction

Epithelial to mesenchymal transition (EMT) regulates cell plasticity during embryonic
development, tissue homeostasis and cancer, where polarized epithelial (E) cells dedifferentiate,
transition through intermediate E/M hybrid states (E/M) and acquire mesenchymal (M)
properties (Nieto et al., 2016). These E/M share several clinically-relevant attributes of
circulating tumor cells (CTCs) and are responsible for EMT-associated stemness,
chemoresistance, immune evasion and metastasis (Dongre and Weinberg, 2019). Complete
molecular characterization of E/M states, and the mechanisms driving E→E/M and E/M→M
transitions will enable development of refined mechanistic models and discovery of new
therapeutic targets.

Approximately 150 genes are currently described as hallmarks of EMT (MSigDB)
(http://www.gsea-msigdb.org/gsea/msigdb/) that were identified from studies measuring the
expression of 'endpoint' markers (e.g. CDH1, MUC1, VIM, FN1) to track the process (Sha et al.,
2019). The prevalent view on EMT as a transcriptionally-driven program is debatable (Yang et
al., 2020). In fact, the poor correlation between genomic alterations or mRNA levels and
(sub)cellular proteoforms during EMT has been noted (Califano and Alvarez, 2017; Liu et al.,
2016). This is crucial because EMT is likely an emergent property where the shifts in cell
physiology and phenotype are orchestrated by intra- and extra-cellular signaling (Sigston and
Williams, 2017), extensive receptor-ligand crosstalk, protein relocalization (Hung and Link,
2011) and metabolic adaptations (Tabassum and Polyak, 2015; Thomson et al., 2019). Attempts
made to model EMT primarily invoke gene-regulatory networks (GRNs) controlled by key
transcription factors (TFs) and miRNAs. These models are usually based on a restricted number
of factors and thus do not capture the multi-layered architecture of signaling during EMT (Hong
et al., 2015; Zhang et al., 2014). To date, no EMT-focused studies have simultaneously measured
metabolite and gene expression changes at different functional levels (e.g., mRNA, total protein,
nucleus, secretome, etc.).

Consequentially, several aspects of EMT remain unclear. This includes dependencies between
molecular layers, secreted molecules, and specific signatures at various stages of EMT, kinetics
and scope of metabolic reprogramming, dynamics of subcellular protein localizations and ligand-
receptor mediated intercellular crosstalk. To bridge these gaps, we employed multiple high-
throughput platforms including microarray, scRNAseq and precision mass spectrometry (MS) to
quantify molecules spanning 12 distinct layers of biological information. Our ability to integrate
this information allowed us to define signatures of E/M states and identify molecular regulators
of key transition points.

RESULTS

A comprehensive resource on TGFβ-induced EMT

The human mammary epithelial cell line MCF10A has been widely used to investigate TGFβ (transforming growth factor β) induced EMT (Liu et al., 2019b; Wei et al., 2015). To generate extensive temporally resolved quantitative expression maps of the evolving EMT landscape, cells were treated with TGFβ (TGF-β1; 10 ng/mL) over 12 days (0, 4 hrs, 1–6, 8 & 12 days), spanning multiple 'omics' layers and employing complementary technology platforms (**Fig. 1A-B**, **Fig. S1A-C**). Across all time points, after stringent quality control (see **STAR Methods**), we report nanoLC-MS/MS-based quantifications (**Fig. 1C**, **Table S1**) of 6,540 whole cell (*WC*), 4,198 nuclear (*Nuc*), 2,223 plasma membrane (*Mem*), 1,209 extracellular vesicle (*EV*) and 1,133 secreted (*Sec*) proteins. Using a serial enrichment workflow (see **STAR Methods**) we also quantified the total phosphoproteome (*Phos*; 8,741 high-confidence sites on 2,254 proteins; including 6975 Ser, 962 Thr and 140 Tyr residues), N-glycoproteome (*Glyco*; 549 proteins), acetylome (*Acet*; 349 sites on 165 proteins) and peptidome (*Pep*; 547 peptides from 202 proteins). For proteomics, samples for each point were multiplexed using isobaric tandem mass-tag (TMT-10), enabling higher throughput and robust comparisons. In addition to proteins, we tracked cellular metabolism in the same samples by nanoLC-MS/MS-based untargeted metabolomics, quantifying 8,325 HMDB-indexed endogenous small molecules (*Metabol*). Furthermore, we measured 23,787 gene transcripts (mRNA) and 2,578 microRNAs (miRNA) using microarrays. To assess cellular heterogeneity, we employed scRNAseq to quantify transcriptomes of 1,913 individual cells (>200 cells per time point) undergoing EMT.

In total, we provide temporal quantifications of >60,000 proteins, phosphosites, mRNAs, miRNAs, and metabolites combined, in addition to 9,785 mRNAs in our scRNAseq dataset (**Fig. 1C**). Subcellular enrichments were performed using previously established MS-compatible protocols (see **STAR Methods**), yielding high purity determined through keyword matching against a cellular compartment annotation database (**Fig. S1D**). Quantitative reproducibility across the 3 biological replicates was excellent (**Fig. S1E**). The expression profiles in **Fig. 1D** provide a snapshot of concurrent changes of a given gene over various layers during EMT. We reproduced expected expression behavior for many established markers of EMT, including an increase of M markers VIM, CDH2 and concomitant decrease of E markers SCRIB, MUC1. Using strict criteria for differential expression (Benjamini-Hochberg adj. *p-value* < 0.05; $r^2 \geq 0.6$ & $|log2FC| \geq 1$; see **STAR Methods**) we identified >10,000 significantly regulated molecules (**Table S2**). As such, this study is the largest experimental description of EMT regulome induced by TGFβ, till date.

We found that molecular abundances were highly variable across layers and time-points (**Fig. S1F**) and all layers contributed significantly to the overall variation in the system (**Fig. 1E**). While each layer showed substantive alterations, the magnitudes (**Fig. S1G**), response profiles (**Fig. 1F**) and fraction of differentially expressed molecules (**Fig. 1G**) of each omic data set during the time course varied significantly. These findings provide direct evidence for global reorganization of the cellular architecture during EMT. For example, *Phos* and *Sec* showed the highest fractional change (36% & 40%, respectively) (**Fig. 1G**), suggesting extensive intra- and extra-cellular signaling during EMT (Scheel et al., 2011). These results add new depths to our current

133 understanding of TGFβ signaling and EMT (**Fig. 1H, I**).

Integrative multi-tiered topology of EMT reveals key transition points and identifies molecular drivers

Although E→M progression is considered a continuum, distinct stages have been reported for
MCF10A cells (Zhang et al., 2014), and cancer tissues (Liu et al., 2019a; Pastushenko et al., 2018).
Using a phylogenetic clustering approach (Hughes and Friedman, 2009) we estimated 'distances'
between time steps to understand the transition kinetics (**Fig. 2A**). We observed that up to 24
hours cells maintained their parental E phenotype, while day 2 marked a switch-like exit from
the E state, suggesting that at day 2 the E/M began and continued through Day 6, after which cells
gradually entered the M state. Cells in day 2 to 6 can be further sub-divided into E/M-1 (Day 2/3,
late E) and E/M-2 (Day 4/5/6, early M) states. These cellular reconfigurations were consistent
with principal component analysis of individual layers, such as *Mem*, *Phos* and *WC* (**Fig. S2A**).

Our observations suggest that each molecular layer has a distinct and complementary role in
shaping the transition (**Fig 2B**, **Fig 1E-G**, **Fig S1F, G**). Although, the topology was generally
correlated (**Fig. 2C**), pairwise coefficients of determination (adj. R^2) revealed a strikingly low
concordance between total proteins and other proteomic layers (mean R^2 ranging from 0.015 for
Acet to 0.299 for *Glyco*) (**Fig. 2D, Upper panel**). The discordance was even stronger between
mRNA and various proteomic layers (mean R^2 ranging from 0.011 for *Pep* to just 0.109 for *WC*)
(**Fig. 2D, Lower panel**), which increased further with EMT progression. Collectively, these
observations illustrate that at a systems level mRNA quantity is a poor proxy of protein
abundance. Total protein or mRNA quantities are also unreliable predictors of a proteins PTMs
or subcellular trafficking, which ultimately determines signaling output.

Next, to model the regulatory patterns of molecules, we employed a 3-step computational
workflow. First, we used a regression strategy for time-course measurements (Conesa et al.,
2006) to remove residual noise (time dimension) and non-reliable (replicates) components.
Second, an unsupervised machine learning approach (self-organizing maps, SOMs) (Wirth et al.,
2012) was applied on the retained molecules to generate time-point specific SOM portraits (**Fig.
2E, Fig. S2B-D**, see **STAR Methods** for details). SOMs are a powerful integration tool for large
and diverse global datasets to extract fundamental underlying patterns of co-regulation (Lu et al.,
2009; Tamayo et al., 1999). Third, we performed pathway enrichments (Ulgen et al., 2019) of
these SOMs to elucidate the functional themes at each time-step of EMT (**Fig. 2F**).

SOMs traced the temporal unfolding of the transition and provided molecular fingerprints of each
time-step. Molecules ranking high (top 1% i.e., rank ≤ 250 & log2FC ≥ 1) with SOMs for control to
day 1 (430 unique molecules) primarily included proteins characteristic of an E state (**Fig. 2E,
Table S3**), such as, ACTG1 (*Mem*; establishes cell junctions and shape) and EPHB2 (*Glyco*;
regulates angiogenesis, contact dependent adhesion and migration; tumor suppressor). SOMs for
days 6–12 (252 unique molecules) contained established M markers (e.g., CALD1, CDH2, FLNA,
FN1, FSTL1, LGALS1, NT5E, TAGLN, TPM1, TPM2, TPM4, VIM). This group included molecules
with established roles in cancer and EMT, including miR-5189 (*miR*; targets ARF6 which

internalizes CDH1), CALR (*Acet*; Lys 62; calcium homeostasis, promotes metastasis, imparts resistance to anoikis), ALCAM (*Nuc*; prognostic marker in multiple cancers linked to nuclear translocation of β-catenin and stemness), ARHGAP33 (*Sec*; regulates intracellular trafficking), RAPGEF5 (*Sec*; promotes nuclear translocation of β-catenin) and SOCS3 (*Phos*; Thr 3/13; E3 ligase, E3 ligase inhibiting TGFβ signaling). The onset of the major regulatory switch at day 2 is likely driven by molecules peaking at its corresponding SOM (95 molecules) such as miR-675-5p/H19 (*miR*; induces HIF1α, SNAIL activity), LTBP1 (*Sec*; master regulator of integrin-dependent TGFβ activation) and CD44 (*Sec*; signal transduction). Notably, many of the genes/molecules identified in the SOM, to our knowledge, have not been previously linked to TGFβ signaling or EMT, presumably because they do not show clear changes in *mRNA* or *WC* but in 'other' molecular layers, such as *EV* or *Sec* (**Fig. S2E**). This contrasts with most MSigDB hallmarks of EMT for which we captured clear transcriptional profiles (**Fig. S2F**).

Enrichment analysis of active subnetworks (Ulgen et al., 2019) identified 237 significant pathways (**Fig. 2F**, **Table S3**), discretized across sequential steps of EMT. For example, 'Beta oxidation of hexanoyl-CoA to butanoyl-CoA' declined as cells leave E and enter E/M, indicating reprogramming of mitochondrial fatty acid β-oxidation, consistent with a metastatic phenotype (Ma et al., 2018). Conversely, 'RHO GTPases mediated activation of ROCKs/PAKs/IQGAPs' increased as cells leave E/M and enter M, suggestive of their key role at this stage of EMT (Ungefroren et al., 2018). The E/M specifically were associated with migration-associated pathways such as 'anchoring fibril formation', 'ECM proteoglycans' and 'laminin interactions', consistent with their shared property with CTCs.

Overall, we catalogued complex kinetics of thousands of molecules spanning multiple molecular layers during EMT. Importantly, we predict signatures specific to each stage of EMT, e.g., the E/M, with potential clinical value.

Metabolomics reveals kinetics and predicts novel enzyme-metabolite associations

TGFβ is known to regulate Warburg effect in cancer cells, but may also regulate other metabolic pathways with implications for cancer management (Hua et al., 2019). Active subnetwork analysis of *WC* and *Phos* datasets identified 13 enriched KEGG metabolic pathways during TGFβ-induced EMT (**Fig. S3A**). Notably, significant enrichment ($p-value \le 0.05$) was only observed post day 2 for pathways such as Steroid biosynthesis (SHB), Sphingolipid metabolism (SLM) and Glycosaminoglycan degradation (GGM). To evaluate these gene-centric inferences of metabolic phenotypes, we directly profiled intra-cellular small molecules by applying an optimized untargeted metabolomics workflow to the same set of samples. Using stringent criteria (see **STAR Methods**), we quantified >8,000 putative HMDB-compounds covering a wide range of chemical classes (**Fig. 3A**).

Phylogenetic clustering (**Fig. 3B**) and SOM analysis (**Fig. 3C**) revealed kinetics of metabolic remodeling prompted by TGFβ. In particular, a distinct temporal separation from parental E cells was observed as early as 4 hours indicating rapid modulation of cellular metabolism by TGFβ, preceding changes in most other layers. To glean further insights, we performed integrative

network analysis of metabolite SOMs with differential molecules in *WC* and *Phos* using MetaboAnalyst (Chong et al., 2019) (**Fig. 3D**). This analysis reiterated several enriched pathways predicted with protein expression alone, e.g., glycerophospholipid metabolism (GPLM) and lysine degradation (LD). However, integration of metabolite and protein data within the framework of metabolite SOMs revealed pathway activities representative of key transition steps of EMT driven primarily by corresponding metabolite signatures (**Fig. S3B-D**). Indeed, we observed that arachidonic acid metabolism (AAM), GPLM and LD pathways were activated within 4 hours of TGFβ stimulation, which was not captured by gene enrichment. Consistent with observations in **Fig 2F**, processes such as fatty acid metabolism, SHB and SLM appear after day 2, as cells prepare for a metastatic phenotype (Koundouros and Poulogiannis, 2020).

Pairwise-correlation based integration of metabolite profiles with proteomics measurements from the same samples could enable mechanistic predictions and aid discovery of novel players. To explore this, we chose AAM as an example. Arachidonic acid is an omega-6 fatty acid stored as membrane phosphoglycerolipid. Its cytosolic release enables stoichiometric chain reactions and results in >100 functionally diverse compounds (Hanna and Hafez, 2018) impacting processes such as redox state, proliferation, apoptosis and chemotaxis (Tallima and El Ridi, 2018). We computed Pearson's correlations between abundance of annotated enzymes, in *WC* and *Phos*, and metabolites of AAM pathway, in *Metabol*, quantified at SOM for 4 hours to day 1 (**Fig. 3E, F**). Overall, we found that metabolites mapping to AAM either increased linearly (CYP450 branch and thromboxane) or initially decreased before rising (Lipoxygenase branch and prostaglandin) (**Fig. 3G**), suggesting fine-tuned regulation of the different branches. Encouragingly, our enzyme-metabolite association map identified PLA2G4A, the rate-limiting phospholipase of AAM pathway (Hanna and Hafez, 2018), among the top candidates (**Fig. 3F**). Interestingly, correlation and expression profiles of PLA2G15, another phospholipase, (**Fig. 3E-G**) indicated a potential enzyme-mediated switch from Cytochrome P450 to Cyclooxygenase/Lipoxygenase branches during E→E/M.

Overall, these observations provide direct evidence for the kinetics of metabolic reprogramming during TGFβ induced EMT. We found metabolites and protein signatures coordinating processes such as AAM, GPLM and LD during key stages of the transition. We also demonstrate how our enzyme-metabolite correlation map could be used as a resource to predict novel enzymes for observed metabolic changes.

scRNAseq analysis reveals heterogeneous responses to TGFβ and novel transcriptional regulators of EMT

scRNAseq studies in murine epithelial Py2T cells treated with TGFβ (Krishnaswamy et al., 2018), MCF10A cells undergoing confluence-dependent EMT (McFaline-Figueroa et al., 2019) and LPS-induced EMT in alveolar epithelial cells (Riemondy et al., 2019), provided critical insights into EMT. However, a temporal analysis of EMT to understand the transition states has not been reported.

After quality control, we retained 1,913 single cells with a combined depth of 9,785 genes (**Fig.**

S4A, Table S1). As anticipated, many of the top expressing genes (TGFB1, TPT1, KRT6A, TMSB10,
MT2A) are key players in EMT (**Fig. S4B**). Interestingly, similar to many primary human tumors
(Aiello et al., 2018; Puram et al., 2018), we did not observe an explicit loss of several classical E
markers at the transcript level (**Fig. 4A, Fig. S4C**), indicating post-transcriptional regulation. This
may be energetically economical for tumor cells by conferring invasive properties without losing
capacity for metastatic seeding (Lambert et al., 2017).

To understand the stages of cell differentiation, we took advantage of the multiple time points in
our dataset (as opposed to a 'pseudo-time'), using Monocle3 (see **STAR Methods**) and identified
20 cell clusters in 3 disjoint partitions (**Fig. 4B, Fig. S4D**). Partition P2 (12 clusters) represents
the primary EMT axis while P1/P3 predominantly expressed genes related to cell cycle (**Fig. S4E**)
and were ignored for further analysis. We observed that C3 responded strongly to TGFβ (**Fig. 4C**),
C4/6 resisted EMT, C5/C8 are the 'transition' states and C13/14 represented terminal M cells (in
terms of hallmark M markers; **Fig. 4C**, **right panel**). Examining clusters C9/18/19 which were
composed of cells from nearly all time points, suggests presence of stable M-type cells in MCF10A
populations and appears to be transcriptional impasse to TGFβ signaling.

To explore the underlying gene expression program, we used hierarchical clustering to group the
individual clusters into 6 subtypes (**Fig. 4C**) and employed SCENIC (Aibar et al., 2017) to infer
TFs and GRNs underlying these subtypes (**Fig. D-F; Table S5**). For each subtype, we identified
several unique (**Fig. 4E**) or highly active TFs (**Fig. 4F**), including both established and novel
players. Several TFs implicated in EMT (TWIST2, FOXK2, ZEB1, ID2, MSX1, ING4) were over-
represented in S4-S6, corresponding to later stages of EMT. In contrast, direct evidence of
mechanistic links of TFs enriched in early stages of EMT (i.e., S3, S4) are lacking, indicating gaps
in current EMT models. Using human TF–binding arrays (see **STAR Methods, Fig. 4G**) we
confirmed elevated activity of three S3 TFs, GLIS2, SP1 and ZNF266 upon TGFβ induction (**Fig.
4H**). In addition to providing experimental evidence to our predictions, the TF-binding array also
identified several other TFs potentially playing important roles at the early stages of EMT (**Fig.
4I**).

Overall, we provide a high-resolution temporal map of gene expression programs of individual
cells as they respond to TGFβ signaling and undergo EMT. More generally, our data suggests that
most transcriptional changes occur at early time points (also **Fig. 2D, Lower panel**), followed by
further adaptations driven predominantly by secondary or post-transcriptional mechanisms.

Spatial regulation of proteins and inter-cellular communication during EMT

Regulation of protein distribution is a crucial signaling mechanism (Ferrell, 1998), but remains
insufficiently understood in EMT. The average pairwise Pearson's correlation between proteins
quantified in various cellular compartments (CCs) ranged between $r = 0.12$ to 0.58 indicating fine-
tuned regulation of protein distributions (**Fig. 5A**). We found 3,965 proteins localizing to ≥2 CCs
(**Fig. S5A**), which we categorized into 2 classes as follows: Class I proteins (1,424) displayed a
correlated trend ($r ≥ 0.4$) consistently across all CC pairs suggesting regulation primarily at or
before the translation step (**Fig. 5B, C; Table S5**). Class II proteins (1,205) displayed anti-

correlated trends ($r \leq -0.4$) between any two CCs, implying active post-translational control of their asymmetric distributions (**Fig. 5D, E; Table S5**).

The Bromodomain and ExtraTerminal (BET) cofactors, BRD2 and BRD4, were identified as Class II proteins, with contrasting kinetic profiles (**Fig. 5F**). Using TF-binding arrays, we verified enhanced global recruitment of BRD4, but not two other cofactors p300 and LSD1, (**Fig. 5G**) indicating its importance during EMT. Indeed, treatment with a pan-BET inhibitor JQ-1 suppressed EMT (**Fig. 5H**). Another notable Class II protein was SCRIB (**Fig. 5I**), which regulates apical-basal polarity and directional migration by acting as a molecular scaffold through protein-protein interactions (Bonello and Peifer, 2019). Using an *in* vivo proximity ligation (BioID) screen of SCRIB ((**Fig. S5B**, see **STAR Methods**), we identified multiple novel interactors (**Table S6**) of which many have known roles in EMT (**Fig. 5J**). Using immunoprecipitation, we verified interactions of SCRIB with SNAP23 and ARHGEF7 (**Fig. 5K**).

Regulation of extracellular signals is important for EMT (Scheel et al., 2011). Subtypes within a cell population can differ in their capacity to send and receive signals, with implications for metastasis and drug resistance (Kim et al., 2018; Tabassum and Polyak, 2015). To map the inter-cellular communication between subgroups of cells during TGFβ-induced EMT, we integrate proteomics and scRNAseq data to perform a systems-wide survey of ligand-receptor (L-R) pair mediated crosstalk (**Fig. 5L**).

First, using a database of >2,500 curated binary L-R interactions (Ramilowski et al., 2015), we searched for pairs of L and R in our *Sec* and *Mem/Glyco* datasets, respectively, assuming that co-directional expression changes in L and/or R of a pair (FDR adj. *p-value* < 0.05 and combined L-R |log2FC| ≥1) can indicate biological role. Currently, at least two L-R pairs are implicated in TGFβ-induced EMT in several tissue models (Heldin et al., 2012). Our analysis detected 67 upregulated and 12 downregulated L-R pairs at any given time point (**Fig. 5M, Table S7**) following TGFβ treatment. Notably, none of these pairs have been directly implicated in TGFβ signaling or EMT, although individually many of the identified L or R occur frequently in the context of EMT and/or cancer. For instance, LAMC2 with its 7 receptors (CD151, COL17A1, ITGA2, ITGA3, ITGA6, ITGB1, ITGB4) exhibit significant alteration during EMT. LAMC2 is overexpressed in cancers (Garg et al., 2014) and its silencing can reverse EMT (Moon et al., 2015; Pei et al., 2019). The cognate receptors, CD151, ITGA3, ITGA6 and ITGB1 synergize with TGFβ signaling to promote metastatic behavior (Pellinen et al., 2018; Sadej et al., 2010; Shirakihara et al., 2013; Zhang et al., 2017).

Next, by systematically comparing the expression patterns of L & R (identified above) among the 16 clusters (identified in our scRNAseq dataset), we obtained cell-cell communication networks (sender → receiver) (**Fig. 5N, Fig. S5C, Table S7**). For example, C13 cells, which appeared at day 3 and showed highest M genes expression (**Fig. 4C**), produced the receptor CD44 to its cognate ligand MMP7 expressed by C17 (**Fig. S5D**). Together, this suggests that communication between cell subgroups (C17:MMP7 → C13:CD44) may exist during EMT which can have potential ramifications for tumor growth (Tabassum and Polyak, 2015). Encouragingly, several of these identified L-R pairs showed strong correlation (**Fig. 5O**) in human breast invasive carcinoma samples (Cancer Genome Atlas Network, 2012; Cerami et al., 2012).

Our study provides a comprehensive analysis of TGFβ-triggered subcellular trafficking as cells

334 undergo EMT. Such translocations, potentially driven by differential PPI, could mediate the tight
335 coordination between functional modules (e.g., SCRIB complex) and EMT phenotypes such as
336 cytoskeletal rearrangement (Huang et al., 2012). We also uncovered novel cell-cell
337 communication pathways via L-R interactions in driving EMT, representing an untapped clinical
338 opportunity.

Modeling phosphoproteome dynamics during EMT reveals kinase susceptibilities

Phosphoregulatory mechanisms are a key aspect of TGFβ signaling and EMT. We confidently
quantified 8,741 phosphosites (p-sites; 6,975 Ser, 962 Thr and 140 Tyr residues) (**Fig. S6A-C**)
over a dynamic range of 10^6 orders of magnitude (**Fig. S6D**), and phospho–STY frequencies (**Fig.**
S6E) in line with previous reports (D'Souza et al., 2014) mapping to 2,254 proteins (**Fig. S6F**). Of
all p-sites, 3,138 (35.8%) were differentially regulated in at least one time point (**Fig. S6G**). At
protein level, different patterns of regulation were noted; some proteins, such as DEK, VIM and
MISP, were regulated at ~90% of detected sites, some proteins, such as CAV1, CAMK2 and
GOLGA1, showed a ~50% mixture of regulated and unregulated sites, while others such as
AHNAK, PML and BCLAF1 showed ~2% differential sites (**Fig. 6A**). Interestingly, the fraction of
regulated p-sites in several proteins, e.g., VIM, increased with EMT progression (**Fig. 6B**).

We observed that in ~50%, p-sites dynamics were not explained ($r \leq 0.4$) by a corresponding
change at the protein level (**Fig. 6C, D**). Interestingly, for ~26%, a directionally opposite change
between phosphorylation and the corresponding protein abundance was noted ($r \leq -0.1$),
suggesting effects on protein stability. Phospho-regulated proteins were enriched for '*nucleus*',
'*cytoskeleton*' and '*focal adhesion*' annotations (**Fig. 6E**) reflecting the importance of CC
remodeling during EMT.

We also computed correlations between nuclear localization profiles (*Nuc*) and phosphorylation
kinetics (*Phos*) of individual proteins (**Fig. 6F, G**). For instance, phosphorylation of MICAL3 at
T684 and S685 regulates CSCs by promoting symmetric division (Tominaga et al., 2019). The
pattern of phosphorylation at residues T684, S685 and S687 (**Fig. 6D,** MICAL3) which are located
at the consensus NLS motif of MICAL3 suggests a role in regulating nuclear translocation and/or
retention at E/M→M transition. Indeed, the bipartite NLS motif of MICAL3 interacts with
Importin-α and the p-sites T684, S685, S687 are directly adjacent to the binding interface (**Fig.**
6H).

To analyze stage-specific kinase activities we generated a ternary model which distinguishes
active kinases into 3 broad stages (i.e., E, E/M, and M) of EMT (**Fig. 6I, Fig. S6H**). An example that
illustrates the utility of this model are AKT isoforms that have distinct and opposing roles during
cancer development (Hinz and Jücker, 2019). We predict AKT1 is strongly associated with the E-
stage, which is consistent with its role in maintaining the E phenotype (Li et al., 2016). In fact,
depletion of AKT1 in MCF10A cells promoted TGFβ-induced E→E/M transition (Iliopoulos et al.,
2009). Our model further predicts key roles for AKT2 and AKT3 at E/M→M transition. Indeed,
AKT2 and AKT3 were associated with tumor invasiveness, stemness and sensitivity to drug
treatment (Chin et al., 2014), key characteristics of the E/M populations (Dongre and Weinberg,

2019). Among several other kinases (**Fig. S6J**), our ternary model predicts critical roles for PRKCA
and AURKB at the junction of E→E/M and E/M→M transitions. PRKCA is reportedly a hub and
therapeutic target for EMT-induced breast CSCs (Tam et al., 2013). Similarly, inhibition of AURKB
was found to reverse EMT and reduce breast cancer metastasis *in vivo* (Zhang et al., 2020).

Overall, we reveal the rich intricacies of the phosphoregulome during EMT, identified functional
p-sites, predict novel kinase susceptibilities, and provide a mechanistic framework to enhance
understanding of the signaling mechanisms during EMT.

Integrative systems causal model of EMT identifies mechanistic vulnerabilities

Systems biology approaches that combine multiple molecular types (proteins, miRNAs,
metabolites) into a framework of established knowledge allow for a rich assessment of a
biological context (Hawe et al., 2019; Karczewski and Snyder, 2018). Using experimentally
validated functional priors (compiled from ENCODE, PhosphoSitePlus, SignaLink 2.0, SIGNOR 2.0,
HINT, miRTarBase and MetaBridge), we combined causal inference and PCSF (Prize Collecting
Steiner Forest) (Akhmedov et al., 2017) to construct hierarchical mechanistic models of the EMT
program (**Fig. 7A**, see **STAR Methods**). The final 'EMT network' comprised of 3,255 edges
connecting 2,217 molecules, including 723 kinase/phosphatase–substrate, 1,407 TF–target, 746
miRNA–target and 31 metabolite–gene interactions.

One of the potential applications of the EMT network is to discover signaling paths from
TGFBR1/2 to any gene(s) of interest within the network. As a demonstration, we queried several
genes (FN1, MMP7, CD44, SCRIB, TWISTNB, ZEB1, SNAI2) and recovered previously known and
unknown paths to them putatively active at multiple stages of EMT (**Fig. 7B**). Next, to identify key
factors driving EMT, we extracted 'controller' nodes (Liu et al., 2011; Vinayagam et al., 2016)
exerting a significant influence on EMT network topology (see **STAR Methods**). Survival analysis
against a large publicly available dataset of primary breast cancers with long-term patient
outcomes (Cancer Genome Atlas Network, 2012) showed a significant association between
tumors with altered expression of these controllers and shortened overall survival (**Fig. 7C**). Not
surprisingly, a few of them are established key regulators of metastasis (**Fig. 7D**), providing
independent support to our model. Unlike controllers, however, the non-controller nodes were
poorly represented in EMT literature (**Fig. 7E**), again highlighting gaps, and potentially
identifying new regulatory processes.

We queried the EMT network to identify the signaling context in which these controllers are
active at various key stages of EMT, which could also provide clues into mechanistic
vulnerabilities (**Fig. 7F**). As cells are stimulated with TGFβ, the TFs SMAD2 and SMAD3 are
activated, as expected. Another early responder was RHO GTPase RAC1, an effector of both KRAS
(Wu et al., 2014) and TGFβ signaling (Ungefroren et al., 2018), suggesting potential crosstalk. The
downstream effector of RAC1, MAPK14 (p38 MAPK), was also regulated early in EMT, suggesting
cooperation between RAC1 and MAPK pathways (Santibáñez et al., 2010). Our model suggests
SMAD3 regulates two other TF hubs, CEBPB (CCAAT/enhancer-binding protein β) and FOXA1.
Loss of CEBPB reportedly switches TGFβ signaling from growth-inhibiting to EMT-inducing

(Johansson et al., 2013), while FOXA1 is reportedly a key TF during EMT (Wang et al., 2013). We
observed that STAT3 is suppressed at later stages of EMT. A recent study in KRAS-driven lung
and pancreatic cancer found that STAT3 is required for maintaining the E state and is lost during
acquisition of M phenotypes (D'Amico et al., 2018).

The EMT network directly predicts novel avenues for blocking EMT. To assess this, we performed
a morphometry-based screening where we treated MCF10A cells with TGFβ in combination with
drugs which were predicted to inhibit several of the controller nodes active at E→E/M transition
(**Fig. 7G,** see **STAR Methods**). Our analysis using a custom-built and publicly accessible image
analysis software (Ochs et al., 2019) (**Fig. 7G**) showed noticeable efficacy of LB100+Barasertib,
LB100+PP1 and Sonidegib+Autocamtide in reverting the elongated (= 'eccentric') phenotype of
EMT-induced cells (**Fig. 7H**), thus providing direct experimental evidence to our predictions.

Overall, our EMT network recapitulates known signaling pathways, uncovers novel routes of
information flow to known regulators of EMT, identifies new signaling players and pathways,
makes substantial and provocative novel predictions and reveals cohesive time-resolved
regulatory patterns and mechanistic links between both controllers and non-controllers.

DISCUSSION

EMT and cancer are *emergent systems* (Sigston and Williams, 2017) wherein progression through
various stages is regulated by intricate networks of intra- and extra-cellular signaling within and
between cells. The key to understanding such complex biological phenomena are establishing
experimental workflows that integrate multiple tiers of biological information (Karczewski and
Snyder, 2018).

Theoretical discussions on EMT are often guided by the Waddington metaphor of a ball (=cell)
rolling over a phenotypic landscape, which is dynamically shaped by multiple parameters:
topologies of signaling networks, molecular stochasticity, extraneous cooperating and opposing
forces (e.g., EMT inducing and/or inhibitory ligands, interaction with other cells) (Li and Balazsi,
2018). By integrating several molecular layers, SOMs and 'neighbor joining' approaches revealed
the kinetics of cell-fate transitions and major phenotypic switch points driven by TGFβ. Several
studies have indicated molecular and phenotypic granularity in EMT continuum and suggested
existence of discrete metastable E/M (Sha et al., 2019). Many current EMT markers are biased
toward the later stages of EMT, when the process is approaching completion (Song et al., 2019).
Consequently, the molecular nature of E/M is still poorly understood. Our time-course integrative
SOM allowed us to trace the temporal unfolding and uncover the molecular nature of the E→M
transition.

While previous omics studies have measured either one or two molecular layers to describe EMT,
our multi-tiered datasets enabled the discovery of several new aspects of EMT which were not
captured by previous approaches. For example, the correlation between *mRNA* and proteins were
weak. Strikingly, correlations between the various *proteomics layers* were also found to be quite
poor, indicating that systems behavior cannot readily be extrapolated by any single layer (e.g.,
mRNA or total proteome), but instead needs an integrated analysis of several molecular layers.

We show that TGFβ-induced EMT is only partially driven by transcription, where several E genes are repressed only post-transcriptionally, while transcripts of M genes are upregulated but mostly during earlier time points. At later stages, post-translational mechanisms become more prominent in driving the process, suggesting that the regulatory control of EMT may be more flexible than previously appreciated. Similarly, our results also predict the mechanistic importance of protein subcellular localization during EMT. A comparison of proteins detected in *EV*, *Sec*, *Glyco*, *Mem* and *Nuc* indicated extensive regulation of protein localizations. Further, previous studies on EMT have largely focused on cell-autonomous signaling, whereas multiple inter-cellular signaling mechanisms are evident from our integrative analysis. Indeed, we found both *EV* and *Sec* to be extensively regulated during EMT. Once considered as cellular 'garbage bins', their active participation in signaling and crosstalk is increasingly recognized (H. Rashed et al., 2017). Our results demonstrating limited qualitative overlap between *EV* and *Sec* hint at a potentially overlooked mechanistic distinction and provides opportunities for new biological insights. We utilized our *Mem*, *Glyco* and *Sec* datasets to provide a repertoire of 79 putative new L-R pairs with potential roles during EMT, which should be validated in future studies. By combining these results with the scRNAseq profiles, we were able to define an extensive network of inter-cellular communications.

In conclusion, we have established a comprehensive multi-tiered molecular landscape of TGFβ-induced EMT. This study aims to provide a valuable resource which is accessible through an interactive website (https://www.bu.edu/dbin/cnsb/emtapp/) (**Fig. S7**) and will strongly complement hypothesis-driven research with direct implications for epithelial cancers.

REFERENCES

Aibar, S., González-Blas, C.B., Moerman, T., Huynh-Thu, V.A., Imrichova, H., Hulselmans, G.,
Rambow, F., Marine, J.-C., Geurts, P., Aerts, J., et al. (2017). SCENIC: Single-cell regulatory
network inference and clustering. Nat. Methods *14*, 1083–1086.

Aiello, N.M., Maddipati, R., Norgard, R.J., Balli, D., Li, J., Yuan, S., Yamazoe, T., Black, T., Sahmoud,
A., Furth, E.E., et al. (2018). EMT Subtype Influences Epithelial Plasticity and Mode of Cell
Migration. Dev. Cell *45*, 681-695.e4.

Akhmedov, M., Kedaigle, A., Chong, R.E., Montemanni, R., Bertoni, F., Fraenkel, E., and Kwee, I.
(2017). PCSF: An R-package for network-based interpretation of high-throughput data. PLOS
Comput. Biol. *13*, e1005694.

Bonello, T.T., and Peifer, M. (2019). Scribble: A master scaffold in polarity, adhesion,
synaptogenesis, and proliferation. J. Cell Biol. *218*, 742–756.

Califano, A., and Alvarez, M.J. (2017). The recurrent architecture of tumour initiation,
progression and drug sensitivity. Nat. Rev. Cancer *17*, 116–130.

Cancer Genome Atlas Network (2012). Comprehensive molecular portraits of human breast
tumours. Nature *490*, 61–70.

Cerami, E., Gao, J., Dogrusoz, U., Gross, B.E., Sumer, S.O., Aksoy, B.A., Jacobsen, A., Byrne, C.J.,
Heuer, M.L., Larsson, E., et al. (2012). The cBio Cancer Genomics Portal: An Open Platform for
Exploring Multidimensional Cancer Genomics Data: Figure 1. Cancer Discov. *2*, 401–404.

Chin, Y.R., Yoshida, T., Marusyk, A., Beck, A.H., Polyak, K., and Toker, A. (2014). Targeting Akt3
Signaling in Triple-Negative Breast Cancer. Cancer Res. *74*, 964–973.

Chong, J., Wishart, D.S., and Xia, J. (2019). Using MetaboAnalyst 4.0 for Comprehensive and
Integrative Metabolomics Data Analysis. Curr. Protoc. Bioinforma. *68*, e86.

Conesa, A., Nueda, M.J., Ferrer, A., and Talón, M. (2006). maSigPro: a method to identify
significantly differential expression profiles in time-course microarray experiments.
Bioinforma. Oxf. Engl. *22*, 1096–1102.

D'Amico, S., Shi, J., Martin, B.L., Crawford, H.C., Petrenko, O., and Reich, N.C. (2018). STAT3 is a
master regulator of epithelial identity and KRAS-driven tumorigenesis. Genes Dev. *32*, 1175–
1187.

Dongre, A., and Weinberg, R.A. (2019). New insights into the mechanisms of epithelial–
mesenchymal transition and implications for cancer. Nat. Rev. Mol. Cell Biol. *20*, 69–84.

D'Souza, R.C.J., Knittle, A.M., Nagaraj, N., Dinther, M. van, Choudhary, C., Dijke, P. ten, Mann, M.,
and Sharma, K. (2014). Time-resolved dissection of early phosphoproteome and ensuing
proteome changes in response to TGF-β. Sci. Signal. *7*, rs5–rs5.

Ferrell, J.E. (1998). How regulated protein translocation can produce switch-like responses.
Trends Biochem. Sci. *23*, 461–465.

Garg, M., Braunstein, G., and Koeffler, H.P. (2014). LAMC2 as a therapeutic target for cancers.
Expert Opin. Ther. Targets *18*, 979–982.

H. Rashed, M., Bayraktar, E., K. Helal, G., Abd-Ellah, M.F., Amero, P., Chavez-Reyes, A., and Rodriguez-Aguayo, C. (2017). Exosomes: From Garbage Bins to Promising Therapeutic Targets. Int. J. Mol. Sci. *18*.

Hanna, V.S., and Hafez, E.A.A. (2018). Synopsis of arachidonic acid metabolism: A review. J. Adv. Res. *11*, 23–32.

Hawe, J.S., Theis, F.J., and Heinig, M. (2019). Inferring Interaction Networks From Multi-Omics Data. Front. Genet. *10*.

Heldin, C.-H., Vanlandewijck, M., and Moustakas, A. (2012). Regulation of EMT by TGFβ in cancer. FEBS Lett. *586*, 1959–1970.

Hinz, N., and Jücker, M. (2019). Distinct functions of AKT isoforms in breast cancer: a comprehensive review. Cell Commun. Signal. CCS *17*.

Hong, T., Watanabe, K., Ta, C.H., Villarreal-Ponce, A., Nie, Q., and Dai, X. (2015). An Ovol2-Zeb1 Mutual Inhibitory Circuit Governs Bidirectional and Multi-step Transition between Epithelial and Mesenchymal States. PLOS Comput. Biol. *11*, e1004569.

Hua, W., ten Dijke, P., Kostidis, S., Giera, M., and Hornsveld, M. (2019). TGFβ-induced metabolic reprogramming during epithelial-to-mesenchymal transition in cancer. Cell. Mol. Life Sci.

Huang, R.Y.-J., Guilford, P., and Thiery, J.P. (2012). Early events in cell adhesion and polarity during epithelial-mesenchymal transition. J. Cell Sci. *125*, 4417–4422.

Hughes, A.L., and Friedman, R. (2009). A phylogenetic approach to gene expression data: evidence for the evolutionary origin of mammalian leukocyte phenotypes. Evol. Dev. *11*, 382–390.

Hung, M.-C., and Link, W. (2011). Protein localization in disease and therapy. J. Cell Sci. *124*, 3381–3392.

Iliopoulos, D., Polytarchou, C., Hatziapostolou, M., Kottakis, F., Maroulakou, I.G., Struhl, K., and Tsichlis, P.N. (2009). MicroRNAs differentially regulated by Akt isoforms control EMT and stem cell renewal in cancer cells. Sci. Signal. *2*, ra62.

Johansson, J., Berg, T., Kurzejamska, E., Pang, M.-F., Tabor, V., Jansson, M., Roswall, P., Pietras, K., Sund, M., Religa, P., et al. (2013). MiR-155-mediated loss of C/EBPβ shifts the TGF-β response from growth inhibition to epithelial-mesenchymal transition, invasion and metastasis in breast cancer. Oncogene *32*, 5614–5624.

Karczewski, K.J., and Snyder, M.P. (2018). Integrative omics for health and disease. Nat. Rev. Genet. *19*, 299–310.

Kim, E., Kim, J.-Y., Smith, M.A., Haura, E.B., and Anderson, A.R.A. (2018). Cell signaling heterogeneity is modulated by both cell-intrinsic and -extrinsic mechanisms: An integrated approach to understanding targeted therapy. PLoS Biol. *16*.

Koundouros, N., and Poulogiannis, G. (2020). Reprogramming of fatty acid metabolism in cancer. Br. J. Cancer *122*, 4–22.

Krishnaswamy, S., Zivanovic, N., Sharma, R., Pe'er, D., and Bodenmiller, B. (2018). Learning time-varying information flow from single-cell epithelial to mesenchymal transition data. PLoS ONE *13*.

Lambert, A.W., Pattabiraman, D.R., and Weinberg, R.A. (2017). Emerging Biological Principles of
Metastasis. Cell *168*, 670–691.

Li, C., and Balazsi, G. (2018). A landscape view on the interplay between EMT and cancer
metastasis. Npj Syst. Biol. Appl. *4*, 1–9.

Li, C.-W., Xia, W., Lim, S.-O., Hsu, J.L., Huo, L., Wu, Y., Li, L.-Y., Lai, C.-C., Chang, S.-S., Hsu, Y.-H., et
al. (2016). AKT1 Inhibits Epithelial-to-Mesenchymal Transition in Breast Cancer through
Phosphorylation-Dependent Twist1 Degradation. Cancer Res. *76*, 1451–1462.

Liu, X., Li, J., Cadilha, B.L., Markota, A., Voigt, C., Huang, Z., Lin, P.P., Wang, D.D., Dai, J., Kranz, G., et
al. (2019a). Epithelial-type systemic breast carcinoma cells with a restricted mesenchymal
transition are a major source of metastasis. Sci. Adv. *5*, eaav4275.

Liu, Y., Beyer, A., and Aebersold, R. (2016). On the Dependency of Cellular Protein Levels on
mRNA Abundance. Cell *165*, 535–550.

Liu, Y., Xue, M., Du, S., Feng, W., Zhang, K., Zhang, L., Liu, H., Jia, G., Wu, L., Hu, X., et al. (2019b).
Competitive endogenous RNA is an intrinsic component of EMT regulatory circuits and
modulates EMT. Nat. Commun. *10*, 1637.

Liu, Y.-Y., Slotine, J.-J., and Barabási, A.-L. (2011). Controllability of complex networks. Nature
473, 167–173.

Lu, R., Markowetz, F., Unwin, R.D., Leek, J.T., Airoldi, E.M., MacArthur, B.D., Lachmann, A., Rozov,
R., Ma'ayan, A., Boyer, L.A., et al. (2009). Systems-level dynamic analyses of fate change in
murine embryonic stem cells. Nature *462*, 358–362.

Ma, Y., Temkin, S.M., Hawkridge, A.M., Guo, C., Wang, W., Wang, X.-Y., and Fang, X. (2018). Fatty
acid oxidation: An emerging facet of metabolic transformation in cancer. Cancer Lett. *435*, 92–
100.

McFaline-Figueroa, J.L., Hill, A.J., Qiu, X., Jackson, D., Shendure, J., and Trapnell, C. (2019). A
pooled single-cell genetic screen identifies regulatory checkpoints in the continuum of the
epithelial-to-mesenchymal transition. Nat. Genet. *51*, 1389–1398.

Moon, Y.W., Rao, G., Kim, J.J., Shim, H.-S., Park, K.-S., An, S.S., Kim, B., Steeg, P.S., Sarfaraz, S.,
Changwoo Lee, L., et al. (2015). LAMC2 enhances the metastatic potential of lung
adenocarcinoma. Cell Death Differ. *22*, 1341–1352.

Nieto, M.A., Huang, R.Y.-J., Jackson, R.A., and Thiery, J.P. (2016). EMT: 2016. Cell *166*, 21–45.

Ochs, F., Karemore, G., Miron, E., Brown, J., Sedlackova, H., Rask, M.-B., Lampe, M., Buckle, V.,
Schermelleh, L., Lukas, J., et al. (2019). Stabilization of chromatin topology safeguards genome
integrity. Nature *574*, 571–574.

Pastushenko, I., Brisebarre, A., Sifrim, A., Fioramonti, M., Revenco, T., Boumahdi, S., Van
Keymeulen, A., Brown, D., Moers, V., Lemaire, S., et al. (2018). Identification of the tumour
transition states occurring during EMT. Nature *556*, 463–468.

Pei, Y.-F., Liu, J., Cheng, J., Wu, W.-D., and Liu, X.-Q. (2019). Silencing of LAMC2 Reverses
Epithelial-Mesenchymal Transition and Inhibits Angiogenesis in Cholangiocarcinoma via
Inactivation of the Epidermal Growth Factor Receptor Signaling Pathway. Am. J. Pathol. *189*,
1637–1653.

Pellinen, T., Blom, S., Sánchez, S., Välimäki, K., Mpindi, J.-P., Azegrouz, H., Strippoli, R., Nieto, R.,
Vitón, M., Palacios, I., et al. (2018). ITGB1-dependent upregulation of Caveolin-1 switches TGFβ
signalling from tumour-suppressive to oncogenic in prostate cancer. Sci. Rep. *8*, 1–14.

Puram, S.V., Parikh, A.S., and Tirosh, I. (2018). Single cell RNA-seq highlights a role for a partial
EMT in head and neck cancer. Mol. Cell. Oncol. *5*, e1448244.

Ramilowski, J.A., Goldberg, T., Harshbarger, J., Kloppmann, E., Lizio, M., Satagopam, V.P., Itoh, M.,
Kawaji, H., Carninci, P., Rost, B., et al. (2015). A draft network of ligand–receptor-mediated
multicellular signalling in human. Nat. Commun. *6*, 1–12.

Riemondy, K.A., Jansing, N.L., Jiang, P., Redente, E.F., Gillen, A.E., Fu, R., Miller, A.J., Spence, J.R.,
Gerber, A.N., Hesselberth, J.R., et al. (2019). Single-cell RNA sequencing identifies TGF-β as a key
regenerative cue following LPS-induced lung injury. JCI Insight *4*, e123637.

Sadej, R., Romanska, H., Kavanagh, D., Baldwin, G., Takahashi, T., Kalia, N., and Berditchevski, F.
(2010). Tetraspanin CD151 Regulates Transforming Growth Factor β Signaling: Implication in
Tumor Metastasis. Cancer Res. *70*, 6059–6070.

Santibáñez, J.F., Kocić, J., Fabra, A., Cano, A., and Quintanilla, M. (2010). Rac1 modulates TGF-β1-
mediated epithelial cell plasticity and MMP9 production in transformed keratinocytes. FEBS
Lett. *584*, 2305–2310.

Scheel, C., Eaton, E.N., Li, S.H.-J., Chaffer, C.L., Reinhardt, F., Kah, K.-J., Bell, G., Guo, W., Rubin, J.,
Richardson, A.L., et al. (2011). Paracrine and Autocrine Signals Induce and Maintain
Mesenchymal and Stem Cell States in the Breast. Cell *145*, 926–940.

Sha, Y., Haensel, D., Gutierrez, G., Du, H., Dai, X., and Nie, Q. (2019). Intermediate cell states in
epithelial-to-mesenchymal transition. Phys. Biol. *16*, 021001.

Shirakihara, T., Kawasaki, T., Fukagawa, A., Semba, K., Sakai, R., Miyazono, K., Miyazawa, K., and
Saitoh, M. (2013). Identification of integrin α3 as a molecular marker of cells undergoing
epithelial-mesenchymal transition and of cancer cells with aggressive phenotypes. Cancer Sci.
104, 1189–1197.

Sigston, E.A.W., and Williams, B.R.G. (2017). An Emergence Framework of Carcinogenesis. Front.
Oncol. *7*.

Song, J., Wang, W., Wang, Y., Qin, Y., Wang, Y., Zhou, J., Wang, X., Zhang, Y., and Wang, Q. (2019).
Epithelial-mesenchymal transition markers screened in a cell-based model and validated in lung
adenocarcinoma. BMC Cancer *19*, 680.

Tabassum, D.P., and Polyak, K. (2015). Tumorigenesis: it takes a village. Nat. Rev. Cancer *15*,
473–483.

Tallima, H., and El Ridi, R. (2018). Arachidonic acid: Physiological roles and potential health
benefits – A review. J. Adv. Res. *11*, 33–41.

Tam, W.L., Lu, H., Buikhuisen, J., Soh, B.S., Lim, E., Reinhardt, F., Wu, Z.J., Krall, J.A., Bierie, B., Guo,
W., et al. (2013). Protein Kinase C α Is a Central Signaling Node and Therapeutic Target for
Breast Cancer Stem Cells. Cancer Cell *24*, 347–364.

Tamayo, P., Slonim, D., Mesirov, J., Zhu, Q., Kitareewan, S., Dmitrovsky, E., Lander, E.S., and Golub,
T.R. (1999). Interpreting patterns of gene expression with self-organizing maps: Methods and
application to hematopoietic differentiation. Proc. Natl. Acad. Sci. *96*, 2907–2912.

Thomson, T.M., Balcells, C., and Cascante, M. (2019). Metabolic Plasticity and Epithelial-
Mesenchymal Transition. J. Clin. Med. *8*, 967.

Tominaga, K., Minato, H., Murayama, T., Sasahara, A., Nishimura, T., Kiyokawa, E., Kanauchi, H.,
Shimizu, S., Sato, A., Nishioka, K., et al. (2019). Semaphorin signaling via MICAL3 induces
symmetric cell division to expand breast cancer stem-like cells. Proc. Natl. Acad. Sci. *116*, 625–
630.

Ulgen, E., Ozisik, O., and Sezerman, O.U. (2019). pathfindR: An R Package for Comprehensive
Identification of Enriched Pathways in Omics Data Through Active Subnetworks. Front. Genet.
10.

Ungefroren, H., Witte, D., and Lehnert, H. (2018). The role of small GTPases of the Rho/Rac
family in TGF-β-induced EMT and cell motility in cancer. Dev. Dyn. *247*, 451–461.

Vinayagam, A., Gibson, T.E., Lee, H.-J., Yilmazel, B., Roesel, C., Hu, Y., Kwon, Y., Sharma, A., Liu, Y.-
Y., Perrimon, N., et al. (2016). Controllability analysis of the directed human protein interaction
network identifies disease genes and drug targets. Proc. Natl. Acad. Sci. U. S. A. *113*, 4976–4981.

Wang, H., Meyer, C.A., Fei, T., Wang, G., Zhang, F., and Liu, X.S. (2013). A systematic approach
identifies FOXA1 as a key factor in the loss of epithelial traits during the epithelial-to-
mesenchymal transition in lung cancer. BMC Genomics *14*, 680.

Wei, S.C., Fattet, L., Tsai, J.H., Guo, Y., Pai, V.H., Majeski, H.E., Chen, A.C., Sah, R.L., Taylor, S.S.,
Engler, A.J., et al. (2015). Matrix stiffness drives Epithelial-Mesenchymal Transition and tumour
metastasis through a TWIST1-G3BP2 mechanotransduction pathway. Nat. Cell Biol. *17*, 678–
688.

Wirth, H., von Bergen, M., and Binder, H. (2012). Mining SOM expression portraits: feature
selection and integrating concepts of molecular function. BioData Min. *5*, 18.

Wu, C.-Y.C., Carpenter, E.S., Takeuchi, K.K., Halbrook, C.J., Peverley, L.V., Bien, H., Hall, J.C.,
DelGiorno, K.E., Pal, D., Song, Y., et al. (2014). PI3K regulation of RAC1 is required for KRAS-
induced pancreatic tumorigenesis in mice. Gastroenterology *147*, 1405-1416.e7.

Yang, J., Antin, P., Berx, G., Blanpain, C., Brabletz, T., Bronner, M., Campbell, K., Cano, A.,
Casanova, J., Christofori, G., et al. (2020). Guidelines and definitions for research on epithelial–
mesenchymal transition. Nat. Rev. Mol. Cell Biol. 1–12.

Zhang, J., Tian, X.-J., Zhang, H., Teng, Y., Li, R., Bai, F., Elankumaran, S., and Xing, J. (2014). TGF-β–
induced epithelial-to-mesenchymal transition proceeds through stepwise activation of multiple
feedback loops. Sci Signal *7*, ra91–ra91.

Zhang, J., Lin, X., Wu, L., Huang, J.-J., Jiang, W.-Q., Kipps, T.J., and Zhang, S. (2020). Aurora B
induces epithelial–mesenchymal transition by stabilizing Snail1 to promote basal-like breast
cancer metastasis. Oncogene *39*, 2550–2567.

Zhang, K., Myllymäki, S.-M., Gao, P., Devarajan, R., Kytölä, V., Nykter, M., Wei, G.-H., and
Manninen, A. (2017). Oncogenic K-Ras upregulates ITGA6 expression via FOSL1 to induce
anoikis resistance and synergizes with αV-Class integrins to promote EMT. Oncogene *36*, 5681–
5694.

Author contributions

Indranil Paul: Conceptualization, Methodology, Software, Data curation, Writing – Original draft, Visualization, Investigation, Formal analysis

Dante Bolzan: Software, Formal analysis

Heather Joan Hook: Investigation

Ahmed Youssef: Software

Gopal Karemore: Software

Michael Uretz John Oliphant: Investigation

Weiwei Lin: Investigation

Qian Liu: Formal analysis

Sadhna Phanse: Software

Dzmitry Padhorny: Software

Sergei Kotelnikov: Software

Carl White: Software

Andrieu Guillaume: Investigation

Pingzhao Hu: Formal analysis

Gerald Denis: Investigation

Dima Kozakov: Software

Brian Raught: Investigation

Trevor Siggers: Investigation

Stefan Wuchty: Software, Formal analysis

Senthil Muthuswamy: Conceptualization, Investigation

Andrew Emili: Conceptualization, Supervision, Writing – Original draft, Resources, Funding acquisition, Formal analysis

Acknowledgements

We dedicate our sincere gratitude to Stefano Monti, Xarelabos Varelas, Valentina Perissi and Matthew Layne for providing critical feedback on the manuscript. We thank the Microarray and Sequencing Core Facility at Boston University School of Medicine for acquisition of mRNA, miRNA and scRNAseq data.

Funding

S.M., G.D., and A.E. acknowledge joint funding (UO1CA243004) from the NCI program Research Projects in Cancer Systems Biology. The CNSB has generous ongoing support from Boston University.

Competing interests

The authors declare no competing financial interest.

Main figure legends

Figure 1. A multi-dimensional resource on TGFβ-induced EMT

(A) MCF10A cells were exposed to TGFβ for the indicated time points. A major premise of the study was to characterize various cell-states and the transition points through which EMT occurs.

(B) Samples from 3 biological replicates were aliquoted and multiple technology platforms were employed to quantify various molecular layers.

(C) An overview of numbers of molecules quantified in various layers.

(D) Expression snapshots of some well-known EMT markers. Heatmaps show log2FC values (relative to Control, FDR = 0.05).

(E) Variance explained by each layer over 4 principal components.

(F) Ridge plot showing differential molecules, as % of total quantified, for each layer.

(G) Pie chart showing the overall fraction (in %) of differential molecules (yellow portion) relative to all molecules quantified in each layer.

(H) Overlap between established EMT databases (MSigDB, www.gsea-msigdb.org; dbEMT2.0, http://dbemt.bioinfo-minzhao.org) and differential proteins and miRNAs (adj. *p-value* < 0.05; $r^2 \geq 0.6$ & $|log2FC| \geq 1$) from this study.

(I) Differential molecules (proteins, miRNAs, metabolites) from this study were used to assess the number of coherent functional modules, i.e., biological interactions or network edges, by employing the Prize-Collecting Steiner Forest algorithm on an interaction network compiled from PathwayCommons, miRTarBase and STITCH.

Figure 2. The topological architecture of EMT

(A) Neighbor-joining was applied as a distance metric to reveal the similarities between timepoints, considering all bulk measurements, excluding *mRNA* (see text).

(B) Combined pseudo-eigenvalues space of all datasets, indicating the contribution of each dataset to the eigenvalue (variance).

(C) Matrix correlations between each pair of datasets.

(D) Line plots show the distribution of adjusted coefficient of determination (R^2) values between layers as a function of time points.

(E) *Left panel.* SOM-portraits characterizing expression landscape of the samples. Color gradient refers to over- or under-expression of metagenes (=cluster of genes) in each time-point compared to the mean expression level of the metagene in the pool of all time-points: red = high, yellow/green = intermediate levels and blue = low (see **STAR Methods** for details). *Middle panel*. Representative examples that appeared among the highest-ranking features (top 1%) in each time-point. *Right panel*. Barplot shows the number of features that were contributed by each layer among the highest-ranking features (top 1%) in time-point.

(F) Heatmap depicts sample-wise pathway scores, which were derived from subnetwork-based pathway enrichment analysis using the highest-ranking molecules (top 1%) for each SOM.

Figure 3. Dynamics of TGFβ-induced metabolic adaptations

    (A) Barplot showing the 'class' distribution of quantified metabolite features as defined in HMBD. Also shown are the relative proportions of differentially expressed features for each class (*p-value* ≤ 0.01, absolute log2FC ≥ 1).

    (B) Neighbor-joining tree for the *Metabol* dataset.

    (C) SOM analysis of the *Metabol* dataset.

    (D) Top 10% of metabolite features of each SOM were grouped based on clusters in `B` and used for "Network analysis" using MetaboAnalyst (https://www.metaboanalyst.ca).

    (E) Identified metabolites of AAM pathway in SOM for 4 hours to day 1 were taken and Pearson's correlation computed with known metabolic enzymes (KEGG) quantified in the *Phos* & *Prot* datasets.

    (F) The plot shows enzymes ranked according to their Pearson's correlation with metabolites of AAM pathway, as detected in this study.

    (G) Schematics of information flow from TGFβ signaling to AAM pathway, mediated by known enzymes. Heatmaps and line plots display 'standardized' expression values over the time points.

Figure 4. scRNAseq analysis reveals cellular dynamics and novel TFs for EMT

    (A) Violin plots showing expression of well-known EMT hallmarks in each time point.

    (B) UMAP of scRNAseq by Monocle3. Dots represent single cells and are colored by inferred clusters, while trajectories depict cells during EMT.

    (C) Heatmap showing the number of cells (in log2 scale) in each cluster of partition P2. Mean expression of some well-known E and M markers in each cluster are also shown.

    (D) The plot shows all TFs ranked according to their SCENIC score. TF names are shown for the 5 TFs with the highest scores and some well-known EMT associated TFs.

    (E) Tree displays unique TFs identified by SCENIC for each subtype. TFs highlighted in 'bold' are known players in EMT as of dbEMT2.0. The genes in the outer circle are representative examples used by SCENIC to infer TF activity.

    (F) Barplot shows the inferred activities of top 15 TFs based on SCENIC scores in each subtype.

    (G) Schematics of the human TF-binding array workflow (see **STAR Methods** for details).

    (H) & (I) Density plots of ∂B-scores of indicated TFs. Two-sided Kolmogorov-Smirnov test was performed to evaluate the significance of distribution differences between Control and TGFβ treated conditions.

Figure 5. Spatial regulation of proteins and intercellular communication

    (A) The schematic summarizes correlations based on Pearson's coefficients between the indicated layers. Each pie chart depicts the fraction of differential proteins (orange slice) with respect to all proteins quantified in the layer. The dotted lines indicate correlation of overlapping proteins between layers.

    (B) The plot displays top 25 Class I proteins. Each pair is represented by a different shape, as shown.

    (C) Expression profiles of top 3 Class I proteins. Each colored line represents a molecular layer, as shown.

    (D) The plot displays top 25 Class II proteins. Each pair is represented by a different shape, as shown.

(E) Expression profiles of top 3 Class II proteins. Each colored line represents a molecular
layer, as shown.
(F) Expression profiles of BRD2 and BRD4 in various layers, as indicated. Legend as in E
(G) Density plots of ∂B-scores of indicated co-factors. Two-sided Kolmogorov-Smirnov test
was performed to evaluate the significance of difference in distributions.
(H) Phase-contrast images after 6 days of TGFβ treatment in presence or absence of active JQ-
1 (100 nM) or its inactive analogue.
(I) Expression profile of SCRIB in various layers, as indicated. Legend as in E
(J) PPI network of SCRIB interactors identified using BioID.
(K) Immunoblots showing interactions of a few SCRIB partners identified using BioID.
(L) Schematics of analysis pipeline for discovering active L-R pairs (see **STAR Methods**).
(M) Heatmap showing combined log2FCs of L-R pairs in the *Sec* and *Mem/Glyco* datasets.
(N) Network plot showing L-R interactions detected between different P2 cell clusters.
(O) Scatter plot of correlations between indicated gene-pairs in Breast invasive carcinoma
samples. Regression line is shown in red.

Figure 6. Phosphoproteome dynamics during EMT

(A) Fraction of detected p-sites on a protein that are regulated during EMT.
(B) Detected p-sites on VIM and their expression during EMT. The gray lines indicate all p-
sites that are catalogued in PhosphoSitePlus database.
(C) Distribution of Pearson's correlation between expression of proteins and the p-sites
detected on them.
(D) Schematic showing two examples each where expression of proteins and p-sites showed
either low (≤–0.1; CDS2, CBX1) or high (≥0.4; MISP, MICAL3) correlation.
(E) Gene ontology enrichment of genes with at least a single regulated p-site at any time point.
(F) Distribution of Pearson's correlation between expression of proteins detected in *Nuc*
layer and the p-sites detected on them in the *Phos* layer. A few EMT hallmarks are
highlighted.
(G) Heatmaps of expression profiles of indicated molecules in *Nuc* and *Phos* layers.
(H) Structural model of MICAL3 p-sites, NLS and Importin-α.
(I) Ternary plot of kinase activity scores binned into 3 broad stages of EMT, i.e., E, ICS, and
M.
(J) *Top*. Expression of PRKCA as detected in various layers. *Bottom*. Pathway enrichment of
all differential substrates of PRKCA detected in our dataset.
(K) *Top*. Expression of AURKB as detected in various layers. *Bottom*. Pathway enrichment of
all differential substrates of AURKB detected in our dataset.

Figure 7. Integrative systems causal model of EMT

(A) Schematics of network and causal inference workflow. Non-redundant genes with most
significant expression (Hotelling's T^2 statistic) profiles were used for the analysis.
CausalPath-estimated logical networks were used to augment a custom-built confidence-
weighted scaffold interactome which was then used to solve the Steiner forest problem
using the OmicsIntegrator software. Only differential molecules (relative to Control, FDR
adj.*p-value* ≤ 0.05, |log2FC| ≥ 1) were considered as 'prizes'.
(B) A subgraph showing the signaling context of well-known EMT players 'queried' on the
EMT network.

(C) Kaplan-Meier plots comparing prognostic performance of MSigDB hallmarks and
'controllers' identified in this study, as indicated.
(D) Overlap between established EMT databases (as in Figure 1) and the 'controllers'
identified in this study.
(E) Overlap between the EMT databases and the 'non-controllers' identified in this study.
(F) A simplified schematic showing the hierarchical relationships between several bottleneck
genes at different stages of EMT, as indicated.
(G) Workflow for morphometric screening of drug combinations. The tool GENIMASEG is
available here: https://figshare.com/articles/software/GenImaSeg_Tool_zip/14371778
(H) The barplot of results of the above experiment indicates varying degrees of synergy or
antagonism between inhibitors, in influencing EMT-associated changes in cell shape
(eccentricity). Error bars indicate spread of datapoints across all quantified cells in each
condition.

Supplementary figure legends

Figure S1. Related to Figure 1. A multi-dimensional resource on TGFβ-induced EMT

(A) TGFβ treatments were staggered, at defined time periods, such that all plates were
harvested at the same time. Cells were serum starved for 16 hours before harvesting.
(B) Phase-contrast images of MCF10A cells at different time points. Scale bar = 100 μm.
(C) For maximizing efficiency while maintaining compatibility with technology-specific
protocols, after harvesting cells were collected in 4 aliquots per replicate, as shown.
Conditioned media were also collected.
(D) Upper panel – Barplots showing percentage of quantified proteins with annotation in the
'Cellular Component' category. Lower panel – Boxplots showing intensity values (log2) of
proteins (= markers) commonly used to assess sub-cellular fractionation purity. Box
edges correspond to 25th and 75th percentiles, whiskers include extreme data points.
(E) Cumulative distribution (%) of Pearson's coefficients across the samples (= time points).
Significant overlap between the 3 biological replicates, shown as 3 colors, indicates high
reproducibility.
(F) Heatmaps of molecular expression profiles of indicated layers. The mean of all quantified
molecules across the 3 replicates was used.
(G) Violin plots showing the spread of log2FCs of molecules at each time point (relative to
Control) for each layer.

Figure S2. Related to Figure 2. The topological architecture of EMT

(A) PCA of the various molecular layers. Time points are shown with different shapes and
colors. Points with similar shape/color indicate biological replicates.
(B) The population map presents the number of genes mapped to each individual metagene.
(C) The plot summarizes co-variance structure of datasets at the metagene level.
(D) The plot shows correlation of expression patterns of individual genes and the metagenes
in which they are contained.
(E) Temporal expression profiles of a few example genes identified in SOM analysis. Each
colored line represents a molecular layer, as shown.
(F) Temporal expression profiles of a few example genes enlisted as 'EMT hallmarks' in
MSigDB database. Each colored line represents a molecular layer, as shown.

Figure S3. Related to Figure 3. Dynamics of TGFβ-induced metabolic adaptations

    (A) The heatmap displays the results of active subnetwork analysis using differential molecules in *Prot* and *Phos* datasets.

    (B) – (D) Metabolite-metabolite interaction networks, using MetaboAnalyst, for indicated clusters. High connectivity indicates co-regulation of functionally related metabolites.

Figure S4. Related to Figure 4. scRNAseq analysis reveals cellular dynamics and novel TFs for EMT

    (A) An outline of the QC pipeline employed for scRNAseq data analysis.

    (B) The plot shows top 25 most expressed genes. Each row corresponds to a gene and each bar corresponds to the expression of the gene in single cells.

    (C) Expression of indicated genes in *mRNA* layer.

    (D) Developmental trajectories of MCF10A cells in response to TGFβ, inferred by Monocle3. Clusters are indicated by colors.

    (E) Heatmap showing aggregate expression of groups of genes (=Modules) with similar expression pattern across the *partitions*, by Monocle3. Modules 9/12 were highly expressed in P1/P3 and were enriched for 'Cell cycle' related GO annotations.

Figure S5. Related to Figure 5. Spatial regulation of proteins and intercellular communication

    (A) The plot shows number of common genes (intersection size, *y* axis) between layers as indicated. Only differential genes in each layer ('Set size', FDR ≤ 0.05; adj. *p-value* ≤ 0.05; $r^2 ≥ 0.6$ & |log2FC|, relative to Control, of ≥ 1) were considered for the analysis.

    (B) Schematics of the BioID experiment performed to discover novel SCRIB partners induced by TGFβ signaling.

    (C) Circos plot showing L-R interactions between the P2 clusters.

    (D) UMAP plots of scRNAseq data highlighting the co-expression patterns of MMP7 and CD44.

Figure S6. Related to Figure 6. Phosphoproteome dynamics during EMT

    (A) An outline of QC pipeline for *Phos* data analysis.

    (B) Enrichment efficiency for TiO_2 workflow employed for the study.

    (C) About 74% of all detected p-sites were reliably localized (=Class I) by MaxQuant.

    (D) Quantifications of p-sites were achieved with a dynamic range of 10^6 orders of magnitude.

    (E) Proportions of number of phosphate moieties in each detected p-site.

    (F) Number of phosphoproteins regulated over the time course.

    (G) Magnitudes of log2FC values for p-sites over the time course.

    (H) The plot shows all kinases ranked according to their KSEA enrichment. Mean of |z-score| values over all time points were taken.

Figure S7. Related to Figure 7. Integrative systems causal model of EMT

    (A) A snapshot of the companion website which can be interactively and freely accessed at (https://srphanse.shinyapps.io/EMTAPP_2/).

Figure 1

A Experimental design and premise

MCF10A cells: TGFβ (10 ng/mL)

Control 4 hrs | 1 2 3 4 5 6 8 12 days

Epithelial → Intermediate hybrids → Mesenchymal

Epithelial **E**
Hybrid **E/M**
Mesenchymal **M**
Transition points ⊚

Protein layers · Metabolites · miRNA · mRNA
Topological relations · Transition kinetics · Molecular landscape Discriminants

B Data acquisition, technologies and platforms

Single-cell RNA sequencing
Bulk Transcriptome, Metabolites & Proteome

WC proteome † · Phosphorylation · Nuclear region
N-Glycosylation * · Membrane · Secretome · Exosome · Peptidome

Barcode cells
Library preparation
ddSeq SC seq
Cells
Genes / Time points
Data matrices

mRNA miRNA · Metabolome
Microarray
Time points
Genes
Data matrix

10-plex (TMT labelling)
basic RP fractionation
nLC/Q-Exactive HF-X
Ion spectrum

C Datasets (this study, 10 time points)

Proteomics

Component	Peptides	Genes (signif. up/dn) **	PTM sites * (signif. up/dn) **
Whole cell proteome (WC)	552,716	6,540 (403/470)	--
Nuclear region (Nuc)	286,162	4,198 (298/259)	--
Membrane (Mem)	35,870	2,223 (149/131)	--
Peptidome (Pep)	735	202 (31/31)	--
Secretome (Sec)	27,136	1,133 (247/200)	--
Extracellular vesicles (EV)	20,438	1,209 (149/123)	--
Glycosylated proteins (Glyco)	6696	590 (17/26)	--
Phosphorylation (Phos)	149,469	2,254	8,741 (1,536/1,602)
Total unique proteins	--	7,632	--

Transcriptomics

Single cell	Matrix	**Total RNA**	Genes (signif. up/dn)
Number of cells ***	1,913	mRNA	23,787 (2,441/1,993)
mRNA	9,785	mi-RNA	2,574 (178/118)

Metabolomics

	Features (MS1 peaks)	HMDB indexed	KEGG indexed
Features (Level 2) ****	27,759	8,325 (610/468)	1,118 (65/59)

* Unique PTM sites remaining after QC steps; ** relative to `Control`; absolute log2FC ≥ 1, FDR ≤ 0.05, r² ≥ 0.6;
*** Number of cells & genes remaining after QC steps **** Level 2 identifications (Salek et al., 2013)

D MultiOmic profiles of EMT markers

Epithelial markers: CDH13, SCRIB, ABLIM2, MUC1
Mesenchymal markers: CDH2, CALD1, VIM, PDLIM4

EV, Sec, Glyco, Mem, Phos, Acet, Pep, WC, Nuc, mRNA
4 hours, Day 1, Day 2, Day 3, Day 4, Day 5, Day 6, Day 8, Day 12

log2FC Down — Up

E Variance contribution

Layers: EV, Mem, Metabol, miRNA, mRNA, Nuc, Pep, Phos, Prot, Sec
Variance
PC 1, PC 2, PC 3, PC 4
Principal components (PC)

F Differential molecules over time points

WC, Sec, Phos, Pep, Nuc, mRNA, miRNA, Metabol, Mem, Glyco, EV
4 hours, Day 1, Day 2, Day 3, Day 4, Day 5, Day 6, Day 8, Day 12

G Differential molecules as % of total detected

Differential / No change

Acet 7% / 93%	EV 23% / 77%	Glyco 8% / 92%	Mem 13% / 87%
Metabol 13% / 87%	miRNA 11% / 89%	mRNA 20% / 80%	Nuc 13% / 87%
Pep 19% / 81%	Phos 36% / 64%	WC 14% / 86%	Sec 40% / 60%

H Comparison to current knowledge

Proteomics
This study 1,667 — dbEMT2 703 — 213 — MSigDB 365 — 126 — 46 — 73

ACTN4, CD82, CDH1, CDH7, CLU, CTNND1, F11R, F3, FN1, FOXO1, IGFBP7, IRF6, ITGB1, ITGB4, JUN, LGALS1, LOX, LOXL2, MMP2, MUC1, NUMB, OCLN, PCDH9, PKP3, PLA2G4A, ROCK1, S100A8, SCEL, SERPINE1, SERPINF1, SKP1, SLC9A3R1, SPARC, TIMP3, TNC

microRNAs
This study 1,066 — 93 — dbEMT2 56

MIR106B, MIR130A, MIR130A, MIR130B, MIR130B, MIR132, MIR141, MIR141, MIR145, MIR149, MIR150, MIR150, MIR15A, MIR17, MIR193A, MIR193A, MIR200A, MIR200A, MIR200B, MIR20A, MIR221, MIR221, MIR222, MIR23A, MIR23B, MIR29C, MIR301A, MIR30A, MIR30B, MIR30D, MIR331, MIR331, MIR345, MIR345, MIR345, MIR424, MIR452, MIR491, MIR491, MIR506, MIR506, MIR612, MIR675, MIR675

I Enhanced discovery of coherent functional modules

Prize-Collecting Steiner Forest Number of biological interactions

Day 12, Day 8, Day 6, Day 5, Day 4, Day 3, Day 1, 4 hours

WC, Phos, Nuc, Sec, Mem, EV, Pep, Glyco

Current knowledge (MSigDB)

Figure 2

A Neighbor joining

B Relative pseudoeigengene variance

C Matrix correlations

D R² between layers

E SOMs · Features · Contribution

F Active subnetwork analysis of upregulated genes

Figure 3

Figure 4

Figure 5

Figure 6

A Fraction of regulated p−sites

B Regulation of VIM p−sites

C Correlation: Prot and Phos

D Regulation of Prot and Phos

E Enriched "Cellular components"

F Correlation: Nuc and Phos

G Nuc and Phos

H Structural model of MICAL3 p-sites, NLS and Importin-α

I EMT stage-dependent kinase activities

J Expression profiles

K Expression profiles

Figure 7

A An overview of network analysis pipeline

B PathLinks: TGFBR1 to key EMT players

C EMT network-derived signatures predict patient survival

D Overlap with 27 controllers

E Overlap with 2191 non-controllers

F A simplified schematic of links between several *controllers* during EMT

G Workflow for morphometric analysis of inhibitor screen

H Exposed combinatorial vulnerabilities

Figure S1. Related to Figure 1

Figure S2. Related to Figure 2

Figure S3. Related to Figure 3

A GeneSet enrichment of metabolic pathways using active subnetworks

B Metabolite-metabolite interaction network of metabolites identified in SOM for **4 hours to day 1**

C Metabolite-metabolite interaction network of metabolites identified in SOM for **day 2 to day 4**

D Metabolite-metabolite interaction network of metabolites identified in SOM for **day 5 to day 12**

Figure S4. Related to Figure 4

A scRNAseq QC pipeline

Data matrices ● 26,364 genes • 2125 cells

Genes not expressed in any cell ● 16,666 genes • 2125 cells

Outlier cells (MAD ≥ 3) ● 16,666 genes • 1913 cells

Cells with ≤ 200 genes ● 16,666 genes • 1913 cells

Genes detected in ≤ 4 cells ● 14,442 genes • 1913 cells

Genes with average UMI count ≤ 0.1 ● 9785 genes • 1913 cells

C Bulk mRNA expression

CDH1 EPCAM

FAP FN1

Standardized expression (log2)

B Top 25 expressing genes

Mean expression

TGFBI
TPT1
KRT6A
KRT17
TMSB10
MT2A
S100A2
KRT5
RPLP1
S100A10
KRT81
RPS14
FTH1
RPS12
SERPINE1
PERP
RPS6
CD24
RPS11
LAMC2
RPS8
RPL8
S100A6
RPS18
RPS4X

Distribution of expression across cells

D Differentiation trajectory

UMAP 2

UMAP 1

E Module and pathway enrichment of the partitions

GO Biological process

Module 12
Module 9
Module 4
Module 1
Module 7
Module 3
Module 6
Module 2
Module 10
Module 11
Module 8
Module 5

Partition 3
Partition 1
Partition 2

Mitotic cell cycle
Cell division
Microtubule cytoskeleton organization
Chromosome segregation
Mitotic nuclear division
Organelle localization
DNA metabolic process
Cellular response to DNA damage stimulus
Meiotic cell cycle
Positive regulation of transferase activity
Gamete generation
Ciliary basal body–plasma membrane docking
Anaphase-promoting complex–dependent catabolic process
Retrograde vesicle-mediated transport, Golgi to ER
Antigen processing and presentation of exogenous peptide antigen via MHC class II
Protein localization to chromosome, centromeric region
Nuclear envelope organization
Regulation of ubiquitin protein ligase activity

DNA metabolic process
Chromosome organization
DNA repair
Response to radiation
DNA replication–independent nucleosome organization
DNA replication–independent nucleosome assembly
Telomere maintenance via semi–conservative replication
Ciliary basal body–plasma membrane docking
Deoxyribonucleoside monophosphate biosynthetic process

Positive regulation of biological process
Organonitrogen compound metabolic process
System development
Immune system process
Tissue development
Locomotion
Cell proliferation
Response to cytokine
Cell adhesion
Cell migration
Leukocyte activation
Cytokine–mediated signaling pathway
Regulation of endopeptidase activity
Extracellular matrix organization
Regulation of peptidyl–tyrosine phosphorylation
Cornification
Negative regulation of viral life cycle

Overlap size (%)

Figure S5. Related to Figure 5

A

Overlap between various molecular layers

B

BioID of SCRIB

C

L–R crosstalk between cell clusters

D

Expression in scRNAseq dataset

Figure S6. Related to Figure 6

Figure S7. Related to Figure 7

Interface for querying the EMT network