Title

Phenotypic and transcriptomic analyses reveal major differences between apple and pear scab nonhost resistance

Authors

Vergne E.[1*], Chevreau E.[1*], Ravon E.[1], Gaillard S.[1], Pelletier S.[1], Bahut M.[2], Perchepied L.[1]

Affiliations

[1] Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, F-49000 Angers, France

[2] Univ Angers, SFR QUASAV, F-49000 Angers, France

*E. Vergne and E. Chevreau made equal contributions to this work.

Corresponding author

E.Vergne

Emilie.vergne@inrae.fr

Abstract

Background. Nonhost resistance is the outcome of most plant/pathogen interactions, but it has rarely been described in Rosaceous fruit species. Apple (*Malus x domestica* Borkh.) have a nonhost resistance to *Venturia pyrina*, the scab species attacking European pear (*Pyrus communis* L.). Reciprocally, *P. communis* have a nonhost resistance to *Venturia inaequalis*, the scab species attacking apple. The major objective of our study was to compare the scab nonhost resistance in apple and in European pear, at the phenotypic and transcriptomic levels.

Results. Macro- and microscopic observations after reciprocal scab inoculations indicated that, after a similar germination step, nonhost apple/*V. pyrina* interaction remained nearly symptomless,

whereas more hypersensitive reactions were observed during nonhost pear/*V. inaequalis*

interaction. Comparative transcriptomic analyses of apple and pear nonhost interactions with *V.*

pyrina and *V. inaequalis*, respectively, revealed differences. Very few differentially expressed genes

were detected during apple/*V. pyrina* interaction, preventing the inferring of underlying molecular

mechanisms. On the contrary, numerous genes were differentially expressed during pear/*V.*

inaequalis interaction, allowing a deep deciphering. Pre-invasive defense, such as stomatal closure,

could be inferred, as well as several post-invasive defense mechanisms (apoplastic reactive oxygen

species accumulation, phytoalexin production and alterations of the epidermis composition). In

addition, a comparative analysis between pear scab host and nonhost interactions indicated that,

although specificities were observed, two major defense lines seems to be shared in these

resistances: cell wall and cuticle potential modifications and phenylpropanoid pathway induction.

Conclusion. This first deciphering of the molecular mechanisms underlying a nonhost scab resistance

in pear offers new possibilities for the genetic engineering of sustainable scab resistance in this

species. Concerning nonhost scab resistance in apple, further analyses must be considered with the

aid of tools adapted to this resistance with very few cells engaged.

Keywords: apple, pear, nonhost resistance, transcriptomics

Background

Apple (*Malus x domestica* Borkh.) and European pear (*Pyrus communis* L.) are two closely

related species belonging to the *Rosaceae* family. Reclassification of the *Rosaceae* placed both *Pyrus*

and *Malus* genera in the subfamily *Spiraeoideae*, tribe *Pyreae* and subtribe *Pyrinae,* this subtribe

corresponding to the long-recognized subfamily *Maloideae* [1]. Efforts to resolve relationships within

this subtribe have frequently failed, and Campbell et al. [2] concluded that the genera of this subtribe

Pyreae have not diverged greatly genetically. The recent sequencing of the pear genome [3] allowed

a precise comparison with the apple genome [4] and led to the estimation of a divergence time

between the two genera of ≈ 5.4 – 21.5 million years ago. Furthermore, apple and pear genomes

share similar chromosome number (n=17), structure and organization.

Scab disease, caused by *Venturia* spp., affects several rosaceous fruit tree species. These

hemibiotrophic pathogens can infect only a limited host-range during their parasitic stage, but they

can overwinter as saprophytes in the leaf litter of a larger range of plant species [5]. Scab disease is

caused by *V. inaequalis* on apple, by *V. pyrina* (formerly named *V. pirina* [6]) on European pear, and

by *V. nashicola* on Japanese (*P. pyrifolia* Nakai) and Chinese (*P. ussuriensis* Maxim) pears. Cross

inoculations of *Venturia* spp. on different rosaceous fruit trees indicates that these pathogens are

highly host specific, probably indicating a close co-evolution of these pathogens with their hosts [7].

A plant species unable to be successfully infected by all isolates of a pathogen species is considered

as a nonhost for this pathogen. Nonhost interactions of *Venturia* spp. on apple and pear have rarely

been described. Microscopic observations have been made on *P. communis* / *V. nashicola* [8] as well

as *M. domestica* / *V. pyrina* and *P. communis* / *V. inaequalis* [5, 9]. In all cases, conidia germinated

and produced appressoria and runner hypheae, but failed to establish a network of stroma. No

macroscopic symptoms were visible.

Because of its durability, nonhost resistance has attracted numerous studies over the last

decade, which have uncovered its multiple and complex defense components. The underlying

mechanisms of nonhost resistance comprise pre-invasion resistance with preformed or induced cell-

wall defenses, metabolic defense with phytoanticipin or phytoalexin accumulation, pattern-triggered

immunity (PTI) as well as elicitor-triggered immunity (ETI) and various signaling pathways [10].

To our knowledge, the molecular bases of scab nonhost resistance of apple and pear have

never been investigated. We were not able to find reports on large-scale fungi nonhost resistance

analyses in pear and the few available in apple are about *Penicillium digitatum* and are conducted on

fruit [11, 12].

However, it is possible to find genome-wide molecular analyses of scab host resistance in

apple and pear. Thus, Perchepied et al. [13] performed a detailed transcriptomic analysis of the host

resistance of pear against *V. pyrina* strain VP102, deployed in a transgenic pear bearing the well-

known apple *Rvi6* resistance gene against *V. inaequalis*. They reported the modulation of expression

of 4170 genes and revealed that downstream of the pathogen recognition, the signal transduction

was triggered with calcium, G-proteins and hormonal signaling (jasmonic acid (JA) and

brassinosteroids), without involvement of salicylic acid (SA), and that this led to the induction of

defense responses such as a remodeling of primary and secondary cell wall, cutin and cuticular waxes

biosynthesis, systemic acquired resistance (SAR) signal perception in distal tissues, and the

biosynthesis of phenylpropanoids (flavonoids and lignin). Only four other transcriptomic studies

involving pear/pathogen host interactions have been published so far but are not concerning scab.

Yan et al. [14] reported the modulation of expression of 144 pear genes after fruit treatment by

Meyerozyma guilliermondii, an antagonistic yeast used for biocontrol of natural pear fruit decay.

Zhang et al. [15] similarly reported the modulation of expression of 1076 pear genes after treatment

with *Wickerhamomyces anomalus*, another biocontrol agent. Using RNA-seq, Wang et al. [16]

reported a major role of ethylene signalization during the compatible interaction between *P. pyrifolia*

and *Alternaria alternata*, a necrotrophic pathogen. Finally, Xu et al. [17] applied RNA-seq to

characterize the genes of *Penicillium expansum* activated after infection of pear fruits.

Concerning host resistance of apple against *V. inaequalis*, subtractive hybridization [18, 19]

and cDNA-AFLP [20] led to the identification of a limited set of differentially expressed genes in *Rvi6*

natural resistant 'Florina' variety (scab inoculated 'Florina' versus mock, [19]), or in *Rvi6* resistant

transgenic 'Gala' lines (*Rvi6* transgenic 'Gala' versus non-transformed 'Gala', after scab inoculation,

[18]; *Rvi6* transgenic 'Gala' before versus post scab inoculation, [20]). Recently, Perchepied et al. [13]

also performed a transcriptomic analysis of the *Rvi6* resistance in a transgenic 'Gala' line (transgenic

versus non-transformed, before and after scab inoculation). They reported the modulation of

expression of 2977 genes and revealed that downstream of the pathogen recognition, signal

transduction was triggered with calcium and interconnected hormonal signaling (auxins and

brassinosteroids), without involvement of SA, and that this led to the induction of defense responses

such as a remodeling of primary and secondary cell wall, galactolipids biosynthesis, SAR signal

generation and the biosynthesis of flavonoids. Genome-wide molecular analyses of apple scab host

resistance have also been achieved in other context than the *Rvi6* resistance. A RNA-seq analyze

identified five candidate genes putatively involved in the ontogenic scab resistance of apple [21]. In

addition, nuclear proteome analysis identified 13 proteins with differential expression patterns

among varying scab resistance 'Antonovka' accessions [22]. Recently, Masoodi et al. [23] performed

a RNA-seq analyze comparing three scab-resistant ('Florina', 'Prima', and 'White Dotted Red') and

three susceptible ('Ambri', 'Vista Bella', and 'Red Delicious') apple genotypes out to mine new scab

resistance genes. They reported the modulation of expression of 822 genes related to various

pathways, i.e., metabolic, protein processing, biosynthesis of secondary metabolites, plant hormone

signal transduction, autophagy, ubiquitin-mediated proteolysis, plant-pathogen interaction, lipid

metabolism, and protein modification pathways.

Thus, if large-scale analyses of pear and apple scab host resistance can be found, in-depth

knowledge of transcriptional patterns and gene functions involved in apple and pear scab nonhost

resistance is still needed. The objectives of our study were 1) to precisely describe nonhost resistance

symptoms in *M. domestica* / *V.pyrina* and *P. communis* / V. *inaequalis* interactions 2) to analyze the

underlying molecular mechanisms of both nonhost interactions through a transcriptomic study 3) to

compare the mechanism of host [13, 23] and nonhost scab resistance in apple and European pear.

Results

Macroscopic and microscopic symptoms analyze

Nonhost interactions were observed in a test performed on leaves of 'Gala' apple and

'Conference' pear cultivars, inoculated by a *V. pyrina* strain (VP102) and a *V. inaequalis* strain (VI

EUB05) respectively. At the macroscopic level, a total absence of sporulation was observed on all

nonhost interactions (Table 1), on the contrary to host interactions (Fig. 1 A and B). Very few pear

plants inoculated with *V. inaequalis* presented resistance symptoms such as pin points (Fig. 1C) and

chlorotic lesions (Fig. 1D), whereas the apple 'Gala' remained completely symptomless after *V. pyrina*

inoculations (Fig. 1E).

Table 1: Scab qualitative note of pear and apple lines inoculated with *V. pyrina* and *V. inaequalis*.

Percentage (number) of plants in the different classes of symptoms, 42 days after inoculation

Class of symptoms	*V. pyrina* strain VP102		*V. inaequalis* strain EUB05	
	'Conference'	'Gala'	'Conference'	'Gala'
0: absence of symptoms	0	100 (28)	90 (17)	0
1: hypersensitivity (pin points)	0	0	5 (1)	0
2: resistance (chlorotic lesions, slight necrosis, crinkled aspect)	0	0	5 (1)	0
3a: weak resistance (necrotic or chlorotic lesions with occasional very light sporulation)	0	0	0	0
3b: weak susceptibility (clearly sporulating chlorotic or necrotic lesions)	0	0	0	0
4: susceptibility (sporulation only)	100 (19)	0	0	100 (28)

At the microscopic level, three days after inoculation, there was no clear difference between

host and nonhost interactions: the conidia of *V. inaequalis* and *V. pyrina* germinated equally on both

hosts forming one or two appressoria (Fig. 1 F and G). However, 14 days after inoculation, there was

a clear reaction of the plant cells in contact with the appressoria (accumulation of red

autofluorescent compounds and enlargement of these cells), which could indicate very small scale

hypersensitive reactions (HR) (Fig. 1 H and I) in both plant species, more frequently in pear than in

apple (Fig1. J and K). No formation of subcuticular stroma and no conidiogenesis were observed in

the nonhost interactions.

Global gene expression analyze

Differentially expressed genes (DEGs) were obtained by comparing transcript abundance in

leaves between T0 and 24 hours post inoculation (hpi) and between T0 and 72 hpi, in the nonhost

interactions 'Gala' / *V. pyrina* VP102 and 'Conference' / *V. inaequalis* EUB05. These time points were

chosen in order to cover the period of establishment of the first intimate contacts between fungal

and plant cells: conidia germination and appressoria formation. For each comparison, the

experimental design is a dye switch approach [24] between the two biological repeats made by

condition (genotype x treatment x time). Each biological repeat is a pool of three leaves from three

different plants.

In total, 60 DEGs in apple and 1857 DEGs in pear were identified, which amounts to 0.19 % of

all apple genes on the apple AryANE v2.0 microarray, and 4.23 % of all pear genes on the Pyrus v1.0

microarray (Table 2). Among the 1857 pear DEGs, 80.2 % were only detected at 24 hpi and 15.4 %

only at 72 hpi, whereas 4.2 % were up-regulated or down- regulated similarly at both time points of

the experiment. Among all the pear DEGs observed at 24 and 72 hpi, the proportion of up-regulated

DEGs was higher (68.8 %) than the proportion of down-regulated DEGs (31.2 %).

Table 2. Number of DEGs identified during apple and pear nonhost response to *V. pyrina* and *V.*

inaequalis

	'Gala' / VP102		'Conference' / EUB05	
	24 hpi	72 hpi	24 hpi	72 hpi
Total # of DEGs*	49	11	1570	364
DEGs in % of all genes on the microarray**	0.16	0.03	3.58	0.83
% of up-regulated DEGs	67.3	36.4	74.5	25.5
% of down-regulated DEGs	32.7	63.6	25.5	74.5
% of DEGs without *Arabidopsis* homolog	27.1	30.4	0.70	1.09

163 *: DEGs numbers were calculated using the p-adj values ≤ 0.01 as selection threshold
164 **: 31311 genes on the apple Ariane V2 microarray, 43906 genes on the pear V1 microarray
165

166   To validate the transcriptomic data, 12 DEGs with varied ratios (between -1.9 and 2.9) were

167 tested by quantitative RT-PCR (qPCR; Table S1), on the two biological repeats used for transcriptomic

168 analyses. Considering the low number of DEGs found for apple in this study, we only tested two of

169 them in qPCR. As seen in Table 2 for pear, at 24 hpi, a majority of DEGs are up-regulated and at 72

hpi, a majority of DEG are down-regulated. qPCR was then performed essentially on DEGs with positive ratios at 24 hpi and negative ratios at 72 hpi (Table S1). The qPCR results confirmed the induced or repressed status of all tested DEGs.

Among the 1857 pear DEGs, the 1845 DEGs with *Arabidopsis* homologs (TAIR identifier) have been classified according to MapMan functional categories (Fig. 2). In order to highlight the enriched classes, the numbers of DEGs per category have been normalized to the numbers of *Arabidopsis* genes in each MapMan category and bootstraps have been done to provide a confidence estimate for the accuracy of the output (Fig. 2). We then more particularly explored the DEGs present in the following enriched classes: Hormone metabolism, Stress, Lipid metabolism, Signaling, Secondary metabolism, Cell wall, and depending on the defense pathways and responses identified, search for others related DEGs in enriched wider classes: Protein, RNA and Miscellaneous. The Table 3 gives the defense pathways and responses identified and the functional categories in which DEGs related to these pathways/responses have been found. The analyze of the 184 DEGs found (corresponding to 158 different functions) is expanded in the Discussion section (Metadata of the 184 DEGs in Table S2).

Table 3. Numbers of DEGs in the defense pathways and responses identified among the enriched functional categories analyzed in pear nonhost response to *V. inaequalis*

MapMan functional categories	Defense pathways and reponses identified												
	JA	SA/SAR	Calcium	ROS	Stomatal closure	HR	Cell Wall	Cuticle and waxes	Terpens and isoprenoids	Phenylpropanoids	Lignin	Coumarin and hydroxycinnamic acid	TOTAL
Protein	1					6						2	9
RNA	3					1		1		2			7
Miscellaneous		5				1		6			3	6	21
Hormone metabolism	10	2					1	1	3				17
Stress	4	11	2	1		3							21
Lipid Metabolism	5	1				8		7					21
Signalling		4	5	4	9	3							25
Secondary metabolism								4	9	5	11	2	31
Cell Wall							32						32
TOTAL	23	23	7	5	9	22	33	19	12	7	14	10	184

189   Among the 60 apple DEGs found, 17 have no *Arabidopsis* homolog, 12 more have

190   unpredicted function and for 11 more we could not find information (Metadata of the 60 DEGs in

191   Table S3). In view of our findings in pear / *V. inaequalis* nonhost interaction, 9 of the 20-remaining

192   apple DEGS could be relevant in apple / *V. pyrina* nonhost interaction and are further analyzed in the

193   Discussion section (Metadata of the 9 DEGs in Table S4).

Discussion

More frequent symptoms of nonhost resistance in pear versus apple at microscopic level

The apple 'Gala' remained completely symptomless after *V. pyrina* inoculations (Fig. 1E). This

is similar to the observation of Chevalier et al. [9] after inoculation of 'Gala' with another *V. pyrina*

strain. On the contrary, pear plants inoculated with *V. inaequalis* presented occasional pin points

symptoms (Fig. 1D) and chlorotic lesions (Fig. 1E). Chlorotic lesions had already been observed by

Chevalier et al. [9] after inoculation of the pear 'Pierre Corneille' with the *V. inaequalis* strain EUB04,

but pin points had never been reported in this nonhost interaction.

At the microscopic level, 14 days after inoculation, there was a clear reaction of the plant

cells in contact with the appressoria (accumulation of red autofluorescent compounds and

enlargement of these cells), which could indicate very small scale hypersensitive reactions (HR) (Fig. 1

H and I, I and K) in both plant species. No formation of subcuticular stroma and no conidiogenesis

were observed in the nonhost interactions, contrary to the host-resistance reactions [13]. These

observations are similar to the collapsed cells described by Chevalier et al. [9] in apple and pear

nonhost reactions, and to the rare HR-like reactions observed by Stehmann et al. [5] on apple

inoculated by *V. pyrina*.

Our results seems to indicate that the leaf surface morphology of apple and pear is equally

compatible with *V. pyrina* and *V. inaequalis* conidia germination, without specific inhibition at this

stage. Recognition probably occurs only at the appressorium site, leading to the cellular reactions

observed. These reactions were limited to a few cells without visible symptoms in apple / *V. pyrina*

interaction, but more extended in pear / *V. inaequalis* interaction and could occasionally produce

macroscopic symptoms.

Different patterns of global gene expression in nonhost resistance in pear versus apple

DEGs were analyzed by comparing transcript abundance in leaves between T0 and 24 hpi and

between T0 and 72 hpi, in the nonhost interactions 'Gala' / *V. pyrina* VP102 and 'Conference' / *V.*

inaequalis EUB05. This experimental design is open to criticism because it could include in results

DEGs responding to the inoculation method (spray of water), and DEGs which expression varies due

to leaves ageing. Water spray is effectively perceived as a mechanical stimulus by plants, the

Arabidopsis response being largely regulated by the JA pathway induction under the control of

MYC2/MYC3/MYC4 transcription factors [25]. But this response is transient and really fast as most of

genes differentially regulated peak within 30 minutes regain untreated transcriptional levels within 3

hours [25]. It seems therefore very unlikely that our analysis at 24 and 72 hpi includes genes

responding to water spray. Resistance due to leaves ageing i.e. ontogenic resistance has been

investigated in *Malus-Venturia* pathosystem at 72 and 96 hpi, and 5 genes have been identified

whose modulation could be linked to this resistance [21]. None of these apple genes, or homologs in

pear, were found differentially expressed at 72 hpi in our interactions (Table S5), which argue against

the presence of ageing responding genes in our results. In total, 60 DEGs in apple and 1857 DEGs in

pear were identified, which amounts to 0.19 % of all apple genes on the apple AryANE v2.0

microarray, and 4.23 % of all pear genes on the Pyrus v1.0 microarray (Table 2).

The very small number of DEGs (60) detected in the apple/ *V. pyrina* nonhost interaction at

24 or 72 hpi seems in agreement with the total absence of macroscopic symptoms observed during

this interaction, and the few small HR-like reactions detected at the microscopic level 14 days post

inoculation. Because these reactions involve only a few cells in the leaves, the changes in gene

expression are probably below the threshold of DEG detection applied in this experiment. It is also

possible that the mild response of apple to *V. pyrina* occurs later than 72 hpi, between 72 hpi and 6

days after inoculation. Indeed, Chevalier et al. [9] observed these rare HR from 6 days post

inoculation, with no more evolution until 14 days after inoculation. A later time (6 days post

inoculation) would have been necessary to conclude on this last hypothesis.

On the contrary to apple/ *V. pyrina* nonhost interaction, the number of DEGs (1857) detected

during the pear / *V. inaequalis* interaction is in the same order of magnitude as the number of DEGs

detected during pear host resistance to *V. pyrina* (see [13]). This could be consistent with the more

frequent observation in this interaction of microscopic HR-like reactions detected 14 days post

inoculation, and of occasional macroscopic symptoms of resistance (chlorotic lesions or pin points).

Hormone signaling pathways classically associated to resistance seems weakly involved in pear / *V.*

inaequalis interaction

We found pear DEGs that seems to indicate that the JA pathway was repressed. The JA

biosynthesis and metabolic conversions were reviewed by Wasternack et al. [26]. In our data, at 24

hpi, the first step of JA biosynthesis, corresponding to the conversion of linoleic acid in 12-oxo-

phytodienoic acid (OPDA), seems to be compromised: six out of seven lipoxygenases (LOX) (three

LOX1, two LOX2 and two LOX5) are repressed, the last one being induced (Fig. 3). OPDA produced in

the chloroplast is then transported to the peroxisome for subsequent conversion to JA via the action

of OPR3 (12-oxo-phytodienoic acid reductase) and β-oxidation enzymes (reviewed in [27] and in

[26]). In pear, three β-oxidation enzymes were found activated more or less rapidly: ACX4 (24 hpi),

MFP2 (72 hpi) and the thioesterase homolog to *At2g29590* (72 hpi), which could suggest that

constitutive OPDA stocks were turned into JA. But the early and long-lasting induction of *JMT* and

ST2A genes could be in favor of a rapid conversion of JA in inactive compounds, *JMT* induction

potentially being reinforced by *BBD1* repression (24 hpi). *BBD1* is actually known as a negative

regulator of *JMT* [28].

The behavior of some DEGs seemed indicate that the defense response depending on JA was

also repressed in pear (Fig. 3). The transcription activator MYC2 of JA-induced genes is known to be

repressed by its interaction with JAZ proteins (reviewed in [26]), and two *JAZ1* and one *JAZ3* coding

genes were found activated at 24 hpi in pear. UBP12 is known as a stabilizer of MYC2 [29]. In our

data, *UBP12* was found repressed at 72 hpi, which could reinforce the inactivation of MYC2. WRKY33

is known as an activator of the JA defense pathway [30] and WRK70 [31] or AS1 (or MYB91; [32]) as

inhibitors, and among JA-responsive proteins, the pathogenesis-related PR3, PR4 and PR12 act

downstream MYC2 activation [33]. In our data, accordingly with the repression of the activator

WRKY33 and the activation of the inhibitors WRK70 and AS1, some JA-responsive genes were also

found repressed, such as the chitinase coding genes *PR4* (also called *HEL*) and *ATEP3*. Furthermore,

no DEGs were found for PR3 and PR12 functions. To conclude, in the nonhost interaction between

pear and *V. inaequalis*, some JA could be produced, but is rapidly converted into inactive compounds

and the subsequent defense response seems to be repressed.

Pear DEGs were found that could indicate that the SA pathway was slightly engaged and

rapidly repressed (Fig. 3). *WRKY70* was induced at 72 hpi in our data. This transcription factor is

known as a negative regulator of SA biosynthesis but a positive regulator of SA-mediated defense

genes in *Arabidopsis* [34, 35, 36], among them *PR2*, *PR5* but not *PR1* [37]. *WRKY33* which is known as

a negative regulator of SA-responsive genes [38], was also repressed at 72 hpi in our data. PR2 and 5

are well-known anti-fungal proteins [39, 40, 41]. At 24 hpi a *PR2*, two *PR2-like*, a *PR5* and a *PR5-like*

coding genes were found induced in our work, another *PR2-like* and two others *PR5-like* being

repressed. The differential expression was maintained at 72 hpi for only two of the previously

activated ones. Furthermore no DEG was found for the PR1 function but three *PR1-like* genes were

found repressed: *ATPRB1* and genes homolog to *At5g57625* and *At4g33720*. *ATPRB1* was already

reported as repressed by SA treatment [42]. In our data, the WRKY70 transcription factor was

induced later than the induced PR genes so we could imagine that induced PR genes were activated

by another precocious regulation, such as an oxidative burst (see below), rather than by WRKY70.

Furthermore, WRKY70 induction seems not sufficient to enable a long lasting induction of these

defense genes.

Other pear DEGs seems to indicate that SA accumulation was transient. *CBP60a* [43], *ACA11*

[44], *EICBP.B* (or *CATMA1*; [45]), all three coding calcium-sensor proteins, are known as negative

regulators of SA accumulation and biosynthesis, as well as the light signaling factor *FAR1* [46] or the

SA glucosyltransferase *UGT74F1* which convert SA in inactive SA 2-O-beta-D-glucoside or the glucose

ester of SA [47]. On the contrary *EDS1*, *PAD4* (reviewed in [48]) and *MKS1* [49] are known as positive

regulators of SA accumulation. In our data, the repression of *CBP60a*, *ACA11* and *EICBP.B* genes could

sustain a SA biosynthesis and accumulation. In addition, *EDS1* activation could allow to consider a

positive feedback loop likely to potentiate SA action via EDS1 cytosolic homodimers, even though

PAD4 was repressed. But, as well as *WRKY70* induction, the repression of the MAPK *MKS1* and the

activation of the light signaling factor *FAR1* (2 times) or the SA glucosyltransferase *UGT74F1* could be

in favor of less free SA. Concerning SAR, *MES1* is known as required in healthy systemic tissues of

infected plants to release the active SA from methyl-SA, which serves as a long-distance signal for

SAR [48] and *ACBP6* may be involved in the generation of SAR inducing signal(s) [50]. In our data, SAR

seemed compromised given the repression of *ACBP6* at 24 hpi and *MES1* at 72 hpi. To conclude, in

the nonhost interaction between pear and *V. inaequalis*, the behavior of some DEGs led us to the

hypothesis that the SA pathway could be engaged but transiently and presumably reduced to the few

infection sites and not spread by SAR in healthy systemic tissues.

Calcium influx and reactive oxygen species (ROS) production seems to act as secondary messengers

and could lead to stomatal closure in pear / *V. inaequalis* interaction

Early responses of plants upon pathogen perception include calcium influx and ROS

production, which both act as secondary messengers ([51], reviewed in [10]). Three pear DEGs were

found that seems to indicate early increased cytosolic calcium level. The CSC (Calcium permeable

Stress-gated cation Channel) ERD4 (found two times) and the two glutamate receptors GLR3.4 and

GLR2.7, are known as calcium permeable channels [52, 53]. They were induced at 24 hpi in our data.

An increased cytosolic calcium level can lead to a pre-invasive defense response by stomatal closure

and promote the post-invasive defense response ROS accumulation [54].

Calcium influx has been reported to promote stomatal closure through the regulation of

potassium flux and the activation of anion channels in guard cells (reviewed in [10]). The stomata

closure is known to be induced via the inhibition of inward potassium currents which is achieved via

activation of calcium dependent protein kinases (CDPK) such as CPK13 and CPK8/CDPK19 [55, 56];

but also via activation of CBL1 of the CBL1-CIPK5 complex, which activates the GORK potassium

outward channel [57]. CPK13, CPK8/CDPK19 and CBL1 were all activated at 24 hpi in our data.

A NADPH oxidase *RBOHB* (respiratory burst oxidase homologs, RBOH) is early (i. e. 24 hpi)

and sustainably (through to 72 hpi) induced in the pear/*V. inaequalis* nonhost interaction, which

could suggest a rapid and maintained apoplastic ROS production. Indeed, the apoplastic ROS are

mainly produced by plasma membrane localized NADPH oxidases, cell wall peroxidases and amine

oxidases [58]. In addition, posttranslational regulation of RBOH is required for its activation and ROS

production. Calcium, phosphatidic acid, and direct interactors such as Rac1 GTPase and RACK1

(Receptor for Activated C-Kinase 1) have been reported to be positive regulators of RBOHs (reviewed

in [59]). For example, the Rac-like/ROP GTPase ARAC3 is known to interact with a RBOH to promote

ROS production [60]. In our data, RBOHB activity could also be supported by the presence of positive

regulators such as Rac-like/ROP GTPase. The three Rac-like/ROP GTPase *ARAC1*, *ARAC3* and the

homolog of *At4g03100* were induced at 24 hpi. CDPKs such as CPK1 are also known to activate

RBOHs in response to increased cytosolic calcium level [61]. But repression of *CPK1* in our data could

indicate that this way of activation did not function.

In response to abscisic acid (ABA) or microbe-associated molecular pattern (MAMP)

immunity, stomatal closure is known to be regulated by apoplastic ROS production (reviewed in [62])

and cysteine-rich receptor-like kinases (CRK) are also known to be elements between ROS production

and downstream signaling leading to stomatal closure, sometimes activated (CRK10), sometimes

inhibited (CRK2 and CRK29; [63]). Three DEGs coding for CRK were found in our data and the

repression of *CRK2* and *CRK29* (found two times) could be consistent with the stomata closure

previously hypothesized, but the repression of *CRK10* (found two times) was not. Beyond closure,

inhibition of stomatal development could be seen as an extreme defense. YODA (found two times)

and MPK6 (found two times) MAPKs belong to a pathway involved in the negative regulation of

stomata development [64]. These two genes were induced early in our data.

To conclude, in pear/*V. inaequalis* nonhost interaction, a calcium influx could lead to the

development of the stomatal closure pre-invasive defense, but could also promote a post-invasive

defense: apoplastic ROS accumulation. Apoplastic ROS, acting themselves as messengers, could come

to strengthen the stomatal closure (Fig. 4).

Transcription factors and sphingolipids could be implicated in maintaining HR under control in pear /

V. inaequalis interaction

ROS are known to mediate cellular signaling associated with defense-related gene

expression, HR i.e. the programmed cell death (PCD) at the site of infection during a pathogen attack,

and phytoalexin production [65]. *Arabidopsis thaliana* RCD1 regulator has been proposed to

positively regulate cell death in response to apoplastic ROS by protein-protein interactions with

transcription factors (reviewed in [66]) and WRKY70 and SGT1b were identified as cell death positive

regulators functioning downstream of RCD1 [66]. *RCD1* and *WRKY70* genes were found induced in

our data, at 24 hpi and 72 hpi respectively.

In *Arabidopsis*, the F-box protein CPR1, in association with the Skp1-Cullin-F-box (SCF)

ubiquitin ligase complex, targets for degradation NLR (nucleotide-binding domain and leucine-rich

repeats containing proteins) resistance protein such as SNC1, RPM1 or RPS2, to prevent

overaccumulation and autoimmunity (reviewed in [67]). A *Skp1-like* (*ASK19*; 72 hpi) gene and *CPR1*

(24 hpi) gene were found induced in our data. A gene coding for RPM1 function was also found

repressed at 24 hpi. These results could be in favor of the hypothesis that NLR receptors do not take

part in the HR development observed in the pear/*V. inaequalis* nonhost interaction (Fig. 1C, H, J). In

addition, the induction of an *AtSerpin1* gene homolog at 24 hpi (found two times) in our data could

be consistent with that hypothesis. Indeed, AtMC1 is a pro-death caspase-like protein required for

full HR mediated by intracellular NB-LRR immune receptor proteins such as RPP4 and RPM1 [68] and

AtSerpin1 is a protease inhibitor which block AtMC1 self-processing and inhibit AtMC1-mediated cell

death [69].

The differential expression of two others components of the proteasome pathway could be

in favor of the HR development: the induction of the *RIN3* ubiquitin E3 ligase (24 hpi) and the

repression of the *BRG3* ubiquitin E3 ligase (24 hpi). Indeed, RIN3 is known as positive regulator of

RPM1 dependent HR [70]. And BRG3 is known as a negative regulator of HR in plant/necrotrophic

pathogen interactions [71].

Sphingolipids are involved in the control of PCD, either as structural components of

membranes but also as initiators in the cell death regulatory pathway. According to Huby et al. [72],

free ceramides and long chain/sphingoid base components (LCBs) are able to trigger cell death, via

ROS production, whereas their phosphorylated counterparts, ceramide phosphates and long chain

base phosphate components (LCB-Ps) promote cell survival. The induction of PCD by LCB is based on

the activation of protein kinases, among them MPK6 [73]. As already mentioned, *MPK6* was found

early induced in our data and we found numerous DEGs in the nonhost interaction between pear and

V. inaequalis that seems to indicate the presence of free ceramides and LCB, which could possibly

participate to the HR development. Free LCB presence is suggested by the activations of *SBH1* (24

hpi), *SLD1* (24 hpi) and another sphingolipid Δ8 long-chain base desaturase homolog to *At2g46210*

(24 hpi; found two times), and their relative conversion in ceramides is suggested by the differential

expressions of the ceramide synthases *LOH2* (repressed at 24 hpi) and *LOH3/LAG13* (induced at 24

and 72 hpi). LCB non-conversion in phosphorylated counterparts could be shown by the *AtLCBK1*

repression (72 hpi) and free ceramides maintenance could be suggested by their non-conversion in

glycosyled ones given the repression of a glucosyl ceramide synthase homolog to *At2g19880* (24 hpi).

The differential expression of numerous known regulators of HR in our data seems again

consistent with the HR phenotype observed. The mechanosensor MSL10 and the calmodulin-

activated Ca^{2+} pump (autoinhibited Ca^{2+}-ATPase [ACA]) ACA11 were found differentially expressed: at

24 hpi *MSL10* was induced and *ACA11* was repressed. MSL10 is known as a positive regulator of cell

death [74] and ACA11 is known as a negative regulator of SA-dependent PCD [44]. Their modulation

could be linked with the hypothesized calcium influx discussed above [44, 75]. The participation of

the SA pathway in the development of the HR could also be supported by the repression of *EDR1* (at

72 hpi). Indeed, the MAPKKK EDR1 is known as a negative regulator of the SA-dependent HR

(reviewed in [76]).

Three other regulators of HR were found modulated in our data. The transcription factor *AS1*

(*MYB91*) was found induced at 24 hpi. It is known as a positive regulator of HR and implicated in the

JA pathway (reviewed in [31]). The transcription factor *WRKY40* was found repressed at 72 hpi. It is

known as a negative regulator of HR [77] and implicated in PTI [78]. Another negative regulator of HR

is the lipid-binding domains containing protein VAD1 [79]. It was found repressed at 72 hpi.

The behavior of another gene in our data, *UGT73B3,* could indicate that the developed HR

was contained due to intracellular ROS production and damages. The function *UGT73B3* was thus

activated (24 hpi). UGT73B3 is known as a restrictor of HR expansion via its action in detoxification of

ROS-reactive secondary metabolites (UGT73B3; [80]).

To conclude, in pear/*V. inaequalis* nonhost interaction, HR was spread out, potentially in link

with a calcium influx and ROS production via free sphingolipids accumulation and not via NLR

receptors. Furthermore, the behavior of eight regulators seems to indicate that the developed HR is

under control (Fig. 4).

Cell wall carbohydrates content and cuticle composition could be altered in pear / *V. inaequalis*

interaction

The first obstacle encountered by host as well as nonhost pathogens attempting to colonize

plant tissues is the plant cell wall, which is often covered with a cuticle. In nonhost resistance, in

which non adapted pathogens normally fail to penetrate nonhost plant cells, the cell wall is

considered an important factor and seen as a preinvasive penetration barrier, or as the onset place

of defensive signaling pathways [10, 54]. Plant cell wall alterations have been demonstrated to have

a significant impact on disease resistance and/or on abiotic stresses. These alterations can concern

the carbohydrates or the phenolic components, and results in impairing or overexpressing cell wall-

related genes (reviewed in [81] and [82]).

We found numerous genes related to the cell wall with a modified expression during nonhost

interaction between pear and *V. inaequalis*, among them about thirty related to the biosynthesis or

the modification of carbohydrates. These DEGs are presented in Table 4 except those related to the

lignin and other phenolic compounds, which will be discussed later. We saw in particular several

DEGs related to cellulose (8) and even more DEGs related to pectin (14) but no DEGs related to

callose.

Concerning these particular carbohydrate components, the model proposed by Bacete et al.

[81] is as follows. Firstly, alterations in cellulose biosynthesis from primary or secondary cell wall

trigger specific defensive responses, such as those mediated by the hormones JA, ET or abscisic acid

(ABA), activate biosynthesis of antimicrobial compounds, but also might attenuate pattern triggered

immunity (PTI) responses. Secondly, alterations of cell wall pectins, either in their overall content,

their degree of acetylation or methylation, activate specific defensive responses, such as those

regulated by JA or SA, and trigger PTI responses, probably mediated by damage-associated molecular

patterns like oligogalacturonides. Thus, even though our results do not completely support a role of

these DEGs, we suppose that the modified expression of cell wall related genes during nonhost

interaction between pear and *V. inaequalis* could be meaningful.

Table 4: Main DEGs related to cell wall carbohydrates synthesis/modification detected during

nonhost interaction pear/*V. inaequalis*.

	Gene	Action	Expression*
Primary cell wall			
Cellulose	CSLA2	synthesis	I
	PNT1	synthesis	I
	COBL2	deposition (GPI-anchored protein)	R
	AtGH9A4	catabolism	I
	XTR7	loosening	I
Hemi-cellulose (xyloglucan)	At5g15490	synthesis	I
Pectin	At3g42180	synthesis	I
	At4g01220	synthesis	I
	GHMP kinase	synthesis	I
	RHM1	synthesis	R
	PME At2g45220	methylesterification	I
	PME At2g46930	methylesterification	I
	PME At3g05910	methylesterification	R
	PME At1g02810	methylesterification	R
	PME44	methylesterification	R
	PG At3g16850	depolymerisation	I
	PG At3g59850	depolymerisation	I
	PG At4g13710	depolymerisation	I
	PG At3g62110	depolymerisation	R
	IDA	degradation	R
Arabinogalactan protein	AGP11	_	I
	AGP1	_	R
Secondary cell wall			
Cellulose	CESA09	synthesis	I
	CESA10	synthesis	I
	CSLG1	synthesis	I
Hemi-cellulose (xylan)	FRA8	synthesis	I
Undetermined			
Expansin	EXP15	loosening	I
	EXPB3	loosening	I
Hemi-cellulose	ATFUC1	modification	I
	XTH33	growth and assembling	R

 *I: induced, R: repressed

Concerning the cuticle layer, most cuticles are composed largely of cutin, an insoluble

polyester of primarily long-chain hydroxy fatty acids. This lipophilic cutin framework is associated

with hydrophobic compounds collectively referred to as waxes. The cuticle is also thought to contain

varying amounts of intermingled cell wall polysaccharides and sometimes also a fraction termed

cutan (reviewed in [83]). Cutin monomers are synthesized by the modification of plastid-derived 16C

and 18C fatty acids in the endoplasmic reticulum (ER), yielding variously oxygenated fatty acid–

glycerol esters referred to as monoacylglycerols, which polymerize upon arrival at the growing cuticle

(Fig. 5, reviewed in [83]).

C16 and C18 fatty acids are also important precursors of cuticular wax synthesis (Fig. 5).

Upon transport to the ER, the C16 and C18 fatty acids are extended to form very-long-chain fatty

acids (VCLFAs; C>20), and this extension is carried out by the fatty acid elongase (FAE) complex

located on the ER membrane. The very-long-chain FAs are then converted into the varied cuticular

waxes (primary alcohols, aldehydes, alkanes, secondary alcohols, ketones) by many ways (reviewed

in [84]).

Interestingly, we found three genes up-regulated 24 hpi belonging to the FAS (fatty acid

synthase) chloroplastic complex implicated in the production of the C16 precursor (Fig. 5): *ACCD*,

FabG and *MOD1* (found two times). ACCD encodes the carboxytransferase beta subunit of the

Acetyl-CoA carboxylase complex which catalyzes the first committed step in fatty acid synthesis: the

carboxylation of acetyl-CoA to produce malonyl-CoA. FabG and MOD1 are respectively a β-ketoacyl

ACP-reductase and an enoyl-ACP-reductase which catalyze respectively the conversion of

acetoacetyl-ACP into β-hydroxyacyl-ACP and the second reductive step from enoyl-ACP to butyryl-

ACP (reviewed in [84]).

In the ER, the four functions we found related to waxes biosynthesis in our data were

repressed at 24 hpi: *KCS4* (found two times), *CER1* and *CER3*, or 72 hpi: *ECR/CER10*. KCS4 and

ECR/CER10 belong to the FAE complex ([85]; [86]). The last two genes are implicated in aldehydes

(*CER1*) and alkanes (*CER1 and 3*) generation (reviewed in [84]). On the contrary, the eight genes we

found related to cutin biosynthesis were induced at 24 hpi except a gene homolog to *At5g14450*,

which was induced at 72 hpi. One of them is a glycerol-3-phosphate acyltransferase (GPAT) coding

gene: GPAT8, which catalyzes the transfer of a fatty acid from coenzyme A (CoA) to glycerol-3-

phosphate (Fig. 4; reviewed in [83]). GPAT8 function in cutin formation has been functionally

confirmed in association with GPAT4 [87]. The seven others genes code GDSL-lipases enzyme

(*At1g28600*, *At1g28660*, *At1g54790*, *At3g16370*, *At3g48460*, *AtCUS4*: *At4g28780*, *At5g14450*), some

of which have been shown to function as cutin synthase (Fig. 4; [88]; reviewed in [83]) and

polymerize monoacylglycerols.

We also found induced respectively at 24 and 72 hpi two genes involved in waxes and cutin

biosynthesis positive regulation: *MYB16* and *SHN1*. The SHN genes (*SHN1–SHN3*), a set of three

largely redundant APETALA 2 family transcription factors from *A. thaliana*, are regulators of floral

cutin and epidermal cell morphology. SHN1 is regulated by the MYB family transcription factor

MYB106, which, along with its paralog MYB16, controls many aspects of cuticle and epidermis

formation in *A. thaliana* (reviewed in [89] and [83]).

Cutin and cuticular waxes play an important role in plant-insect and plant-microbe

interactions. Numerous *Arabidopsis* mutants in cutin and waxes biosynthetic or transport genes, such

as Acyl-CoA binding proteins (ACBP), show varying degrees of cuticle impairment, alterations in cutin

and/or wax composition, and defects in SAR (reviewed in [84]). We found *ACBP6* repressed at 24 hpi.

That repression is not inconsistent with the previously described possible amplification of cutin

biosynthesis and polymerization, given that *acbp6* KO mutation is not associated with a defect in that

pathway [50]. That repression is also consistent with the SAR repression hypothesized above as the

acbp6 KO mutant show compromised SAR [50].

To conclude, our analysis of nonhost pear/V. *inaequalis* interaction could show an alteration

of the cuticle composition with more cutin and less waxes synthesis. The potential increase in cutin

polymerization could lead to a thickening of the cuticular layer to prevent fungus penetration via its

appressoria.

Secondary metabolism seems to lead to G unit lignin polymerization and simple coumarin or

hydroxycinnamic acid amine phytoalexins synthesis in pear / *V. inaequalis* interaction

As distinguished from primary metabolism, plant secondary metabolism refers to pathways

and small molecule products of metabolism that are non-essential for the survival of the organism.

But they are key components for plants to interact with the environment in the adaptation to both

biotic and abiotic stress conditions. Plant secondary metabolites are usually classified according to

their chemical structure. Several groups of large molecules, including phenolic acids and flavonoids,

terpenoids and steroids, and alkaloids have been implicated in the activation and reinforcement of

defense mechanisms in plants (reviewed in [90]).

Terpenoids and steroids, or isoprenoids, are components of both the primary and secondary

metabolisms in cells, and mono-, tri-, sesqui- and polyterpenes are considered as secondary

metabolites (reviewed in [91]). Our results on pear identified seven DEGs and five DEGs which could

belong to the chloroplastic methylerythritol posphate (MEP) and to the cytosolic mevalonic acid

(MVA) pathway of isoprenoids production respectively (Table 5), which results, among others

compounds, in tri- and sesquiterpenes secondary metabolites. The majority of these genes

contribute to produce primary metabolites according to Tetali [91]. Except *SMT2*, that we found

induced at 24 hpi, there is no report concerning a putative implication of others genes in plant biotic

resistance. SMT2 is involved in sterols production and *smt2* mutation was reported to compromise

bacterial resistance in *Nicotiana benthamiana* [92]. The hypothesis is that sterols regulate plant

innate immunity against bacterial host and nonhost infections by regulating nutrient efflux into the

apoplast. *V. inaequalis* is a hemibiotrophic pathogen which colonizes only the apoplast compartment

at the beginning of the interaction. Strong relative induction of *SMT2* in our data could indicate that a

similar mechanism of nutrient efflux regulation via sterols could take place to limit the fungus growth

in pear nonhost resistance against *V. inaequalis*.

Table 5: Main DEGs involved in biosynthetic pathways for terpenes and isoprenoids during pear/*V.*

***inaequalis* nonhost interaction.**

	Gene	Function	Expression*
Cytosolic MVA (mevalonic acid) pathway enzymes	HMGS	catalyze the second step of the pathway	R
	HMGR1	catalyze the third step of the pathway	R
	SMT2	sterols production	I
	FLDH	sesquiterpenes production	R
	SQE2	triterpenes production	I
Chloroplastic MEP (methylerythritol posphate) pathway enzymes	DXR	catalyzes the second step of the pathway	I
	GG reductase	chlorophylls production	R
	VTE4	tochopherols production	I
	KAO1	gibberellins production	R
	PDS2	plastoquinones production	I
	LYC	carotenoids production	I
	PGGT1	covalent attachment of a prenyl group to a protein	I

534 *I: induced, R: repressed
535
In our data, the other DEGs that were linked to secondary metabolism belong to the

phenylpropanoid pathway production (Fig. 6). Among them we found four genes which could belong

to the flavonoid production, all repressed, at 24 hpi (*DFR* and *DRM6*) or 72 hpi (*TT7* and *UGT71D1*).

DFR (dihydroflavonol reductase) is involved in flavan-3,4-ol production and TT7 (flavonoid 3'

hydroxylase) in dihydroquercetin production from dihydro-kaempferol, and UGT71D1 (glucosyl

transferase)   in   quercetin-glycoside   production   from   quercetin   (TAIR   database;

https://www.arabidopsis.org/index.jsp). DMR6 (flavone synthase) is involved in flavone production

from naringenin [93]. Thus flavonoid production does not seem to be favored, which is not consistent

with the induction of *MYB12* at 24 hpi, but consistent with *MYB4* induction at 72 hpi. MYB12 is

actually known as a positive regulator of flavonol biosynthesis in pear and apple fruits [94, 95]

whereas MYB4 is known as a negative regulator of this biosynthetic pathway [96].

Concerning the production of monolignols, precursors of lignin synthesis, some genes potentially related were found induced, others repressed. We found *CYP98A3* and *CAD9* (found two times) induced at 24 hpi and *HCT*, *CCR1* and a gene homolog to *At2g23910* (found two times, one time repressed at 24 hpi, one time repressed at 72 hpi, Fig. 6). *CYP98A3* encodes a C3H (coumarate 3-hydroxylase), *CAD9* encodes a CAD (cinnamyl alcohol dehydrogenase), HCT is an hydroxycinnamoyl-CoA shikimate/quinate hydroxycinnamoyl transferase, *CCR1* encodes a CCR (cinnamoyl-CoA reductase) and *At2g23910* encodes a CCR-related protein. (TAIR and KEGG databases (https://www.genome.jp/kegg/)).

Lignification is obtained by cross-linking reactions of the lignin monomers or by polymer–polymer coupling via radicals produced by oxidases such as peroxidases [97] and laccases [98]. However, while peroxidases are able to oxidize monolignols to produce H, G and S units of lignin, laccases only generate G units [97]. In our data, we found two laccases induced at 24 hpi: *LAC11* (found two times, one time induced at 24 and 72 hpi) and *LAC17* (found two times), and three peroxidases repressed at 24 hpi: *PRX17*, *PER47* and *PRX52* (also repressed at 72 hpi), which could be linked to lignin biosynthetic process (Fig. 6). According to Zhao et al. [98], LAC11 and LAC17, along with LAC4, play a critical role in lignification, and their results suggests that peroxidase and laccase do not serve redundant functions in lignification, at least in the vascular tissues of the stem and root of *Arabidopsis*. Participation in lignin formation has also been proved for PRX17 [99], PER47 [100] and PRX52 [101]. But there are currently no reports about a possible involvement of all these genes in lignification linked to biotic or abiotic stresses. Concerning nonhost resistance, two reports describe lignin accumulation/deposition involvement: one in apple fruit [102] and the other one in cowpea [103]. In the latest, authors showed that preferentially generated lignin units in this nonhost interaction are G units, just as it could be the case in our pear / *V. inaequalis* study. To summarize, it is tempting to think that modifications of expression observed for genes linked to lignin polymerization are relevant for the pear nonhost resistance against *V. inaequalis*, but further functional analysis should be conducted to conclude about the lignin status, such as histochemical

staining [104], content measure by absorbance [104] or Fourier-transform infrared (FTIR)

spectroscopy analysis [105].

The biosynthesis of two others types of phenylpropanoid compounds could appear to be

favored during pear nonhost resistance against *V. inaequalis*: simple coumarin on one hand and

hydroxycinnamic acid amides on the other hand. We found four *BGLU*-like genes induced: *BGLU47*

and *BGLC3* (at 24 hpi), *BGLU16* (at 72 hpi); *BGLU42* (at 24 and 72 hpi) (Fig. 6). These β-glucosidases

could be implied in simple coumarin path production from the cinnamic acid (KEGG database). Some

natural simple coumarins are known as antifungal compounds *in vitro* and have been developed as

fungicides [106]. Previous work on Hevea also reports the correlation between the resistance against

pathogenic fungi and the production of some coumarins, with antifungal activity *in vitro* [107]. We

also found induced at 24 hpi the genes *AACT1/ACT1*, *ATPAO5* and genes homologs to *At4g17830* and

At4g38220 (Fig. 6). AACT1/ACT1 catalyze the first specific step in branch pathway synthesizing

hydroxycinnamic acid amides from the p-Coumaroyl CoA or the feruloyl CoA and amines agmatine or

putrescine [108]. Hydroxycinnamic acid amides are produced in response to pathogenic infections

[108] and surface exported. Hydroxycinnamic acid amides are reported to participate in *Arabidopsis*

nonhost resistance against *Phytophthora infestans* via their inhibitory activity on spore germination

[109]. The three others genes belong to the arginine biosynthesis path (homologs to *At4g1783* and

At4g38220) and the arginine and proline metabolisms which produce the amines agmatine and

putrescine (*ATPAO5*) (KEGG database). Agmatine is directly produced from arginine thanks to an ADC

activity (arginine decarboxylase) and putrescine can be produced from spermidine thanks to a PAO

activity (polyamine oxidase). ATPAO5 catalyzes the conversion of spermine in spermidine. The

induction of these three last genes is therefore consistent with the hypothesis of amines production

in order to enable hydroxycinnamic acid amides synthesis. The induction of *C4H* at 24 hpi could also

favor hydroxycinnamic acid amides synthesis via p-Coumaroyl CoA biosynthesis promotion. C4H

(cinnamate 4-hydroxylase) catalyzes the production of p-Coumaric acid from Cinnamic acid and p-

Coumaric acid gives p-Coumaroyl CoA thanks to 4CL (4- coumarate-CoA ligase) (KEGG database).

Among the suite of defense components synthetized in nonhost as in host context, a

chemical barrier can be established via accumulation of a diverse array of secondary metabolites

rapidly produced upon pathogen infection, named phytoalexins, with toxic or inhibitory effects

(reviewed in [10]). Phytoalexins can be flavonoids, such as the pisatin of pea (in [110]) but also varied

phenylpropanoid compounds. In the nonhost interaction pear / *V. inaequalis*, the production of

flavonoid type phytoalexins does not seem to be favored, except possibly simple coumarin and

hydroxicinnamic acid amines.

Very limited transcriptomic modulation during apple / *V. pyrina* nonhost interaction

Only 60 DEGs were detected in the apple / *V. pyrina* nonhost interaction at 24 or 72 hpi, in

agreement with the total absence of macroscopic symptoms and few cells engaged in an HR-like

reaction observed at the microscopic level. Among these 60 DEGs, 36 have unpredicted function.

Among the 24 remaining DEGs, nine DEGS could be relevant in apple / *V. pyrina* nonhost interaction

in view of our findings in pear / *V. inaequalis* nonhost interaction. *ORG2* (*BHLH038*), a putative

integrator of various stress reactions [111] was induced at 24 hpi. Three genes were related to an

oxidative stress: *GASA10* was repressed at 24 hpi and *NRAMP3* and *AOR* were induced at 24 hpi.

GASA proteins have been suggested to regulate redox homeostasis via restricting the levels of OH. in

the cell wall [112]. The repression of this gene could be thus in favor of more OH. in the cell wall. The

oxidoreductase coding gene *AOR* is known in the chloroplast to contribute to the detoxification of

reactive carbonyls produced under oxidative stress [113]. *NRAMP* genes function as positive

regulators of ROS accumulation, especially during *Arabidopsis Erwinia chrisanthemi* resistance [114].

The induction (at 24 and 72 hpi) of another gene could suggest modifications at the cell wall level:

EXP8, an expansin coding gene involved in cell wall loosening (Tair database). We also found two

genes related to hormone pathways, one induced at 24 hpi: *WIN1* and the other one repressed at 72

hpi: *UBP12*. WIN1 is known as a negative regulator of SA pathway [115] and UBP12 as a positive

regulator of JA pathway via the stabilization of MYC2 [29]. In possible link with the JA pathway, we

also found *TPS21* induced at 24 hpi. TPS21 is involved in sesquiterpenes production and is promoted

by JA signal via MYC2 [116]. TPS21 is especially involved in the jasmonate-dependent defensive

emission of herbivore-induced volatiles in cranberries [117]. Finaly the last DEG we found potentially

relevant in apple / *V. pyrina* nonhost interaction could promote HR via ceramides accumulation.

ACD11 is repressed at 24 hpi in our data. In *acd11* mutants, the relatively abundant cell death

inducer phytoceramide rises acutely [118].

Because nonhost resistance of apple against *V. pyrina* has resulted in a very limited number

of cells engaged in an HR-like reaction, it has not been possible for us to exhaustively describe its

outcome at the transcriptomic level in the first three days of the interaction. Further insight with

later points of kinetic and more adapted techniques such as laser-assisted cell picking, prior to micro

arrays or RNA sequencing analysis (review in [119]) could provide more information in the future.

Comparison of pear resistances against the host pathogen *V. pyrina* and the nonhost pathogen *V.*

inaequalis

Perchepied et al. [13] performed a detailed transcriptomic analysis of the host resistance of

pear against *V. pyrina* strain VP102, deployed in a transgenic pear bearing the well-known apple *Rvi6*

resistance gene against *V. inaequalis*. Comparing this work to ours gives us the rare opportunity to

analyze similarities and differences between a host and a nonhost resistance in the same plant.

Concerning the recognition and early signaling steps of the interactions, many receptors and

co-receptors have been found induced in the host pear resistance, especially damage-associated

molecular patterns receptors such as RLK7, revealing that PTI and ETI must be engaged. We did not

find evidence of the mobilization of such receptors in the pear nonhost resistance. PTI and ETI

receptors are nonetheless reported as implicated in nonhost resistance (reviewed in [120] and [10]).

As we only analyzed post infection transcriptional modulations in the nonhost pear/*V. inaequalis*

interaction (at 24 and 72 hpi), one hypothesis to explain the lack of PTI and ETI receptors in our data

could be that these receptors were already present as preformed defenses and not particularly

induced by the infection onset. In pear nonhost interaction, the earliest likely signaling pathways we

were able to highlight are calcium influx and apoplastic ROS production. Calcium signaling seems to

be also implicated in pear host resistance, but less obviously than it seems in nonhost resistance.

Regarding the hormonal signaling pathways, the JA defense signaling pathway seems to be

repressed in pear nonhost resistance but was found quite activated in pear host resistance. The

JA/ethylene (ET) defense signaling pathway is known as an effective defense against necrotrophic

fungi in *Arabidopsis* [121]. Thus, it is not inconsistent to find the JA pathway potentially repressed in

the development of the pear nonhost resistance against the hemi-biotrophic pathogen *V. inaequalis*.

But it is very interesting to find this pathway rather induced in the development of the pear host

resistance against the other hemi-biotrophic pathogen *V. pyrina*. The SA signaling pathway is

commonly seen as the classical one triggered to resist biotrophic fungi in *Arabidopsis* [121], but it

seems only a little engaged in pear nonhost resistance , and is repressed in pear host resistance. If

this absence of SA implication is quite unexpected in pear host resistance against a hemi-biotrophic

fungus, it could be consistent with the report that the exact role of these key defense

phytohormones is unclear in nonhost resistance and remains to be established [54]. As shown by

Tsuda et al. [122], an explanation for the hormone pathways behavior in pear host resistance could

be that: as both the SA and JA/ET pathways positively contribute to immunity, a loss of signaling flow

through the SA pathway can be compensated by a rerouting signal through the JA/ET pathways. In

addition, independently of SA signaling, but in positive connection with JA signaling, SAR seems to be

engaged in distal tissues during pear host resistance. To conclude, in pear host as well as nonhost

resistances, classical resistance hormones SA and JA/ET, and the correlative PR gene defenses, seems

differently involved than in *Arabidopsis*.

The carbohydrate content of the cell-wall seems to be modified in response to the attacks by

the pathogens. Regarding cell-wall and cuticle, in pear host as well as nonhost resistances, the

numerous DEGs found could highlight important modifications. Similar modifications affected the

cellulose and mainly the pectin contents, but no callose production was observed. Regarding cuticle,

waxes production was induced in host resistance whereas it seems to be repressed in nonhost

resistance, in favor of cutin production / polymerization, which was also induced in host resistance.

To conclude, as a first obstacle encountered by host, as well as nonhost pathogens attempting to

colonize plant tissues, the plant cell wall and its cuticle seem to play a leading role in both pear host

and nonhost resistances.

Finally, the production of secondary metabolites and phenylpropanoids compounds in

particular, could be a major line of defense, in pear host as well as nonhost resistances, but with

divergences. If lignin and flavonoid productions are preponderant in pear host resistance against *V.*

pyrina, lignin implication in pear nonhost resistance seems less clear and flavonoids production

seems repressed. But the biosynthesis of two other types of phenylpropanoid-derived phytoalexins

could be favored during pear nonhost resistance: simple coumarin on one hand and hydroxycinnamic

acid amides on the other hand.

The comparative analysis between a host and a nonhost resistance in pear shows that, even

though specificities are observed, two major defense lines engaged seems shared: the cell wall and

its cuticle on one hand, the secondary metabolism with the phenylpropanoid pathway on the other

hand. Moreover, these defenses seem deployed independently of the SA signaling pathway, yet

widely recognized as the main defense hormone against biotrophic pathogens.

Conclusion

To our knowledge, our work is the first one published regarding a transcriptomic analysis of

post-infections events of a nonhost resistance to *Venturia sp.* in apple and pear. Here, our molecular

work on apple / *V. pyrina* nonhost resistance remains preliminary and in order to allow a deeper

deciphering, further analyses must be considered with the aid of tools adapted to this nonhost

resistance with very few cells engaged in an HR-like reaction, only visible at a microscopic level. In

pear, this deciphering allowed us to show that nonhost resistance against *V. inaequalis* involves

enough pathogen penetration in plant tissue to occasionally trigger visible HR, and develops post-

invasive defenses.

To summarize our findings on pear with a notion of cascading effect, we can propose the

following scenario (Fig. 4): once *V. inaequalis* presence is recognized by pear, a calcium cellular influx

could be induced and could lead to the possible development of a pre-invasive defense, the stomatal

closure, but could also promote a possible early post-invasive defense, an apoplastic ROS

accumulation. Apoplastic ROS, acting themselves as ubiquitous messengers, could come to reinforce

the stomatal closure but could also mediate cellular signaling possibly resulting in two post-invasive

defenses: HR development at infection sites, along with phytoalexin (simple coumarin and

hydroxicinnamic acid amines) production. The inferred alterations of the epidermis composition

(cellulose, pectin, lignin for the cell wall, and cutin for the cuticle), are presumed to strengthen this

physical barrier and could be seen as the development of another pre-invasive defense. Nonhost

resistance is defined as the resistance of an entire plant species against a specific parasite or

pathogen and is seen as the most durable resistance of plant. Thus, understanding the molecular

mechanisms underlying nonhost resistance can open up some interesting avenues to create

sustainable host resistances in the same plant species. Considering pear, in order to stop the

germination and entrance of hemibiotrophic host fungi such as *V. pyrina*, strengthening the cuticle

initial barrier via more cutin production and cross-link, or promoting the biosynthesis of phytoalexins

like hydroxycinnamic acid amines, appear as promising solutions.

Material and methods

Biological material

Apple plants from the cultivar 'Gala' and pear plants from the cultivar 'Conference' were

chosen because of their susceptibility to *V. inaequalis* and *V. pyrina*, respectively. The apple and pear

genotypes were multiplied *in vitro*, rooted and acclimatized in greenhouse as described previously

[123, 124].

For apple scab inoculation, the *V. inaequalis* monoconidial isolate used was EU-B05 from the

European collection of *V. inaequalis* of the European project Durable Apple Resistance in Europe

[125]. For pear scab inoculation, the monoconidial strain VP102 of *V. pyrina* was chosen for its

aggressiveness on 'Conference' [126].

Scab inoculation procedure

Greenhouse growth conditions and mode of inoculum preparation were as described in Parisi

and Lespinasse [127] for apple and Chevalier et al. [128] for pear. Briefly, the youngest leaf of actively

growing shoots was tagged and the plants inoculated with a conidial suspension (2×10^5 conidia ml^{-1})

of *Venturia pyrina* strain VP102 for apple and *Venturia inaequalis* strain EUB04 for pear. Symptoms

were recorded at 14, 21, 28, 35 and 42 days after inoculation on 20 actively growing shoots (one

shoot by plant) for each interaction. The type of symptoms was scored using the 6 class-scale of

Chevalier et al. [129].

Microscopic observations

Histological studies were made on samples stained with the fluorophore solophenylflavine

[130]. In brief, leaf discs were rinsed in ethanol 50° before staining in a water solution of

solophenylflavine 7GFE 500 (SIGMA-Aldrich, St Louis USA) 0.1% (v/v) for 10 min. The samples were

first rinsed in deionized water, then in 25% glycerol for 10 min. Finally, the leaf samples were

mounted on glass-slides in a few drops of 50% glycerol. They were examined with a wide-field

epifluorescence microscope BH2-RFC Olympus (Hamburg, D) equipped with the following filter

combination: excitation filter 395 nm and emission filter 504 nm.

750
Transcriptomics experiment

Leaf samples were immediately frozen in liquid nitrogen and kept at -80°C until analysis.

Sampling concerned the youngest expanded leaf of each plant labeled the day of the inoculation.

Each biological repeat is a pool of three leaves from three different plants and two biological repeats

(n=2) have been made by condition (genotype x treatment x time). Leaf samples taken just before

inoculation (T0) and at 24 and 72 hpi, were then used to perform transcriptomics analyses.

For RNA extraction, frozen leaves were ground to a fine powder in a ball mill (MM301,

Retsch, Hann, Germany). RNA was extracted with the kit NucleoSpin RNA Plant (Macherey Nagel,

Düren, Germany) according to the manufacturer's instructions but with a modification: 4% of PVP40

(4 g for 100 ml) were mixed with the initial lysis buffer RAP before use. Purity and concentration of

the samples were assayed with a Nanodrop spectrophotometer ND-1000 (ThermoFisher Scientific,

Waltham, MA, USA) and by visualization on agarose gel (1% (weight/volume) agarose, TAE 0.5x, 3%

(volume/volume)   Midori   green).   Intron-spanning   primers   (forward   primer:

CTCTTGGTGTCAGGCAAATG, reverse primer: TCAAGGTTGGTGGACCTCTC) designed on the *EF-1α* gene

(accession AJ223969 for apple and PCP017051 for pear, available at https://www.rosaceae.org/, with

the datasets on "*Pyrus communis* v1.0 draft genome") were used to check the absence of genomic

DNA contamination by PCR. The PCR reaction conditions were as follows: 95°C for 5 min, followed by

35 cycles at 95°C for 30 s, 60°C for 45 s, 72°C for 1 min, with a final extension at 72°C for 5 min. The

PCR products were separated on a 2% agarose gel.

Amplifications (aRNAs) were produced with MessageAmpII aRNA Kit (Ambion Invitrogen,

Waltham, MA, USA), from 300 ng total RNA. Then 5 µg of each aRNA were retrotranscribed and

labelled using a SuperScript II reverse transcriptase (Transcriptase inverse SuperScript™ II kit,

Invitrogen, Carlsbad, CA, USA) and fluorescent dyes: either cyanine-3 (Cy3) or cyanine-5 (Cy5)

(Interchim, Montluçon, France). Labeled samples (30 pmol each, one with Cy3, the other with Cy5)

were combined two by two, depending on the experimental design (i. e. for example, the

CF/EUB05/24 hpi sample and the CF/non inoculated sample are labeled with Cy3 and Cy5

respectively and pooled to co-hybridize the microarray) . For each comparison two biological

replicates were analyzed in dye-switch as described in Depuydt et al. [131]. Paired labeled samples

were then co-hybridized to Agilent microarray AryANE v2.0 (Agilent-070158_IRHS_AryANE-Venise,

GPL26767 at GEO: https://www.ncbi.nlm.nih.gov/geo/) for apple, or Pyrus v1.0 (Agilent-

078635_IRHS_Pyrus, GPL26768 at GEO) for pear, containing respectively 133584 (66792 sense and

66792 anti-sense probes) and 87812 (43906 sense and 43906 anti-sense probes) 60-mer

oligonucleotide probes. The hybridizations were performed as described in Celton, Gaillard et al.

[132] using a MS 200 microarray scanner (NimbleGen Roche, Madison, WI, USA).

For microarray analysis we designed two new chips. For apple we used a deduplicated

probeset from the AryANE v1.0 ([132]; 118740 probes with 59370 in sense and 59370 in anti-sense)

augmented by 14844 probes (7422 in sense and 7422 in anti-sense) designed on new gene

annotations from *Malus x domestica* GDDH13 v1.1 (https://iris.angers.inra.fr/gddh13 or

https://www.rosaceae.org/species/malus/malus_x_domestica/genome_GDDH13_v1.1).   These

probes target new coding genes with UTRs when available, manually curated micro-RNA precursors

and transposable elements. For transposable elements we used one consensus sequence for each

family and a randomly peaked number of elements proportional to their respective abundance in the

genome. The microarrays used in this study also have probes for coding genes of *V. inaequalis* but

they have not been taken into account.

For pear the design was done on the *Pyrus communis* Genome v1.0 Draft Assembly &

Annotation   available   on   GDR

(https://www.rosaceae.org/species/pyrus/pyrus_communis/genome_v1.0) web   site.   We   have

downloaded the reference genome and gene predictions fasta files and structural annotation gff file

the 21st of September 2015. Using home-made Biopython scripts we have extracted spliced CDS

sequences with 60 nucleotides before start and after stop codons to get UTR-like sequences likely to

be found on transcripts resulting in a fasta file containing 44491 sequences. These 60 nucleotides

increase the probability of finding specific probes on genes with high similarity. This file was sent to

the eArray Agilent probe design tool (https://earray.chem.agilent.com/earray/) to generate one

probe per gene prediction. Options used were: Probe Length: 60, Probe per Target: 1, Probe

Orientation: Sense, Design Options: Best Probe Methodology, Design with 3' Bias. The probeset was

then reverse-complemented to generate anti-sense probes and filtered to remove duplicated

probes. The final probeset contains 87812 unique probes targeting 1 (73612 probes) or more (14200

probes) potential transcript both in sense and anti-sense.

Normalization and statistical analyses performed to get normalized intensity values have

been done as in Celton, Gaillard et al. [132]. Briefly, data were normalized with the lowess method,

and differential expression analyses were performed using the lmFit function and the Bayes

moderated t-test with the LIMMA package in R software [133, 134]. The pipeline AnaDiff used for

these differential analyses is available at https://zenodo.org/record/6477918#.Yn5O3XVBzIE [135].

For each comparison and each probe, we retrieved a ratio of the logarithms of the fluorescence

intensities (one per compared sample: T0 versus 24 hpi or T0 versus 72 hpi in our case) and an

associated p-value. The applied p-value threshold to determine DEGs (differentially expressed genes)

was 0.01. Through blast analysis, a TAIR accession number (The *Arabidopsis* Information Resource;

https://www.arabidopsis.org/; [136]) has been linked to a majority of apple or pear

"probe/corresponding gene". Thanks to the Functional Classification SuperViewer tool [137], the

TAIR accessions have then been used to class DEGs in functional categories according to MapMan

software   (https://mapman.gabipd.org/homemapman.gabipd.org;   file

Ath_AGI_LOCUS_TAIR10_Aug2012.txt; [138]), and to highlight the enriched ones by calculating the

frequency of DEGs per category, normalized to the numbers of *Arabidopsis* genes in each MapMan

category, and bootstrapping the dataset to provide a confidence estimate for the accuracy of the

result.   The   detailed   analysis   of   DEGs   has   been   done   through   TAIR   and   KEGG

(https://www.genome.jp/kegg/) databases, and bibliography. Metadata for the 193 (184 for pear,

corresponding to 158 different functions and 9 for apple) DEGs discussed in this work are available in

Table S2 and S4 (Online only).

qPCR validation of transcriptomic data

In order to validate transcriptomic data, qPCR was performed on a selection of gene/sample

associations. First-strand cDNA was synthesized using total RNA (2.0 µg) in a volume of 30 µl of 5×

buffer, 0.5 µg of oligodT15 primer, 5 µl of dNTPs (2.5 mM each), and 150 units of MMLV RTase

(Promega, Madison, WI, USA). The mixture was incubated at 42°C for 75 min. qPCR was then

performed. Briefly, 2.5 µl of the appropriately diluted samples (1/16 dilution) were mixed with 5 µl of

PerfeCTa SYBR Green SuperMix for iQ kit (Quantabio, Beverly, MA, USA) and 0.2 or 0.6 µl of each

primer (10 µM) in a final volume of 10 µl. Primers were designed with Primer3Plus, their volumes

were according to their optimal concentration (determined for reaction efficiency near to 100%;

calculated as the slope of a standard dilution curve; [139]). Accessions, primer sequences and

optimal concentrations are indicated in Table S1. The reaction was performed on a CFX Connect Real-

Time System (BIO-RAD, Hercules, CA, USA) using the following program: 95°C, 5 min followed by 40

cycles comprising 95°C for 3 s, 60°C for 1 min. Melting curves were performed at the end of each run

to check the absence of primer-dimers and nonspecific amplification products. Expression levels

were calculated using the ΔΔCT method [140] and were corrected as recommended in

Vandesompele et al. [141], with three internal reference genes (GADPH, TUA and ACTIN 7 for apple,

GADPH, TUA and EF1α for pear) used for the calculation of a normalization factor. For each couple

DEG/sample (sample defining a plant, time, treatment and biological repeat combination), the ratio

was obtained by dividing the mean value of CT calculated from 3 technical repeats by the

normalization factor obtained for this sample.

Supplementary information

852   **Additional File 1: Table S1**, **S2, S3, S4 and S5**.

Abbreviations

ABA: abscisic acid; CDPK: calcium dependent protein kinase; CRK: cysteine-rich receptor-like kinase;

DEG: differentially expressed gene; DFR: dihydroflavonol 4-reductase; DGDG:

digalactosyldiacylglycerol; ET: ethylene; ER: endoplasmic reticulum; ETI: effector triggered immunity;

FAE: fatty acid elongase; GPAT: glycerol-3-phosphate acyltransferase; hpi: hours post inoculation; HR:

hypersensitive reaction; JA; jasmonic acid; LCB: long chain/sphingoid base component; LCB-Ps: long

chain base phosphate component; LOX: lipoxygenase; MAMP: microbe-associated molecular pattern;

OPDA: 12-oxo-phytodienoic acid; PCD: programed cell death; PTI: pattern triggered immunity; RBOH:

respiratory burst oxidase homolog; ROS: reactive oxygen species; SA: salicylic acid; SAR: systemic

acquired resistance

Declarations

Acknowledgements

The authors gratefully acknowledge the IRHS-ImHorPhen team of INRA Angers for technical

assistance in plant maintenance and the technical platforms ANAN and IMAC.

Authors contribution

EC, LP, and EV conceived the study. EC and EV supervised the study. ER and MB performed the

biological experiments. SG and SP performed the database work and assisted with the bioinformatics

analysis. EV wrote the original manuscript. EV and EC edited the manuscript. All authors have read

and agreed to the published version of the manuscript.

Funding

This project was funded by the Synthé-Poir-Pom project (Angers University) and by the TIFON project

(INRAE, department BAP).

Availability of data and materials

The datasets supporting the conclusion of this article are available in the Gene Expression Omnibus

(GEO) repository [https://www.ncbi.nlm.nih.gov/geo/] with GSE159179 and GSE159180 accession

numbers for apple and pear respectively [142, 143]. The pipeline AnaDiff used for differential

analyses is available at https://zenodo.org/record/6477918#.Yn5O3XVBzIE [135].

Ethics approval and consent to participate

Experimental research on plants in this work comply with relevant institutional, national, and

international guidelines and legislation.

Consent for publication

This section is not applicable.

Competing interests

The authors declare that they have no competing interests.

References

1. Potter D, Eriksson T, Evans RC, Oh S, Smedmark JEE, Morgan DR, Kerr M, Robertson KR, et al.

Phylogeny and classification of *Rosaceae*. Plant Syst Evol. 2007;266:5-43.

2. Campbell CS, Evans RC, Morgan DR, Dickinson TA, Arsenault MP. Phylogeny of subtribe *Pyrinae*

(formerly the *Maloideae*, *Rosaceae*): Limited resolution of a complex evolutionary history.

Plant Syst Evol. 2007;266: 119-145.

3. Wu J, Wang Z, Shi Z, Zhang S, Ming R, Zhu SL et al. The genome of pear (*Pyrus bretschneideri*

Redh.). Genome Res. 2013;23:396-408.

4. Velasco R, Zharkikh A, Affourtit J, Dhingra A, Cestaro A, Kalyanaraman A et al. The genome of the

domesticated apple (*Malus × domestica* Borkh.). Nature Genet. 2010;42:833–839.

5. Stehmann C, Pennycook S, Plummer K. Molecular identification of a sexual interloper: the pear

pathogen *Venturia pirina*, has sex on apple. Phytopathol. 2001;91:663-541.

6. Rossman A, Castlebury L, Aguirre-Hudson B, Berndt R, Edwards. (2647–2651) Proposals to

conserve the name *Venturia acerina* against *Cladosporium humile; Venturia borealis* against

Torulama culicola; Venturia carpophila against *Fusicladium amygdali* and *Cladosporium*

americanum; Sphaerella inaequalis (Venturia inaequalis) against *Spilocaea pomi, Fumago*

mali, Actinone macrataegi, Cladosporium dendriticum, Asteroma mali, and *Scolicotrichum*

venosum; and *Venturia pyrina* against *Helminthosporium pyrorum, Fusicladium virescens, F.*

fuscescens, Cladosporium polymorphum and *Passalora pomi (Ascomycota: Dothideomycetes)*.

Taxon. 2018;67:1209-1211.

7. Gonzalez-Dominguez E, Armengol J, Rossi V. Biology and epidemiology of *Venturia* species

affecting fruit crops: a review. Front Plant Sci. 2017;8: 1496.

8. Jiang S, Park P, Ishii H. Penetration behaviour of *Venturia nashicola* , associated with hydrogen

peroxide generation, in Asian and European pear leaves. J Phytopathol. 2014;162: 770-778.

9. Chevalier M, Bernard C, Tellier M, Audrain C, Durel CE. Host and non-host interactions of *Venturia*

inaequalis and *Venturia pirina* on *Pyrus communis* and *Malus x domestica*. Acta Hortic.

2004;663: 205-208.

10. Lee HA, Lee HY, Seo E, Lee J, Kim SB, Oh S, Choi E, Choi E, Lee SE, Choi D. Current understanding

of plant nonhost resistance. Mol Plant-Microbe Interact. 2017a;30:5-5.

11. Buron-Moles G, Wisniewski M, Viñas I, Teixidó N, Usall J, Droby S, Torres R. Characterizing the

proteome and oxi-proteome of apple in response to a host (Penicillium expansum) and a

non-host (Penicillium digitatum) pathogen. J Proteomics. 2015;114:136-151.

12. Vilanova L, Vall-Llaura N, Torres R, Usall J, Teixidó N, Larrigaudière C, Giné-Bordonaba J.

Penicillium expansum (compatible) and Penicillium digitatum (non-host) pathogen infection

differentially alter ethylene biosynthesis in apple fruit. Plant Physiol Biochem. 2017;120:132-

143.

13. Perchepied L, Chevreau E, Ravon E, Gaillard S, Pelletier S, Bahut M, Berthelot P, Cournol R,

Schouten HJ, Vergne E. Successful intergeneric transfer of a major apple scab resistance gene

(Rvi6) from apple to pear and precise comparison of the downstream molecular mechanisms

of this resistance in both species. BMC Genomics. 2021;22:843.

14. Yan Y, Zheng XF, Apaliya MT, Yang HJ, Zhang HY. Transcriptome characterization and expression

profile of defense-related genes in pear induced by *Meyerozyma guilliermondii*. Postharvest

Biol Technol. 2018;141:63-70.

15. Zhang Q, Zhao L, Li B, Gu X, Zhang X, Boateng NS, Zhang H. Molecular dissection of defense

response of pears induced by the biocontrol yeast, *Wicherhamomyces anomalus* using

transcriptomics and proteomics approaches. Biol Control. 2020;148:104305.

16. Wang H, lin J, Chang YH, Jiang CZ. Comparative transcriptomic analysis reveals that
ethylene/H2O2-mediated hypersensitive response and programmed cell death determine
the compatible interaction of sans pear and *Alternaria alternate.* Front Plant Sci.
2017b;8:196.

17. Xu M, Yang Q, Boateng NAS, Ahima J, Dou Y, Zhang H. Ultrastructure observation and
transcriptome analysis of *Penicillium expansum* invasion in postharvest pears. Postharvest
Biol Technol. 2020;165:111198.

18. Paris R, Cova V, Pagliarani G, Tartarini S, Konjanc M, Sansavini S. Expression profiling in HcrVf2-
transformed apple plants in response to Venturia inaequalis. Tree Genet Genome.
2009;5:81–91.

19. Cova V, Paris R, Toller C, Patocchi A, Belasco R, Komjanc M. Apple genes involved in the response
to Venturia inaequalis and salicylic acid treatment. Sci Hortic. 2017;226:157–72.

20. Paris R, Dondini L, Zannini G, Bastia D, Marasco E, Gualdi V, et al. dHPLC efficiency for semi-
automated cDNA-AFLP analyses and fragment collection in the apple scab-resistance gene
model. Planta. 2012;235:1065–80.

21. Gusberti M, Gessler C, Broggini GA. RNA-Seq analysis reveals candidate genes for ontogenic
resistance in Malus-Venturia pathosystem. PLoS One. 2013;8:e78457.

22. Sikorskaite-Gudziuniene S, Haimi P, Gelvonauskiene D, Stanys V. Nuclear proteome analysis of
apple scab cultivar 'Antonovka' accessions in response to apple scab (Venturia inaequalis).
Eur J Plant Pathol. 2017;148:771–84.

23. Masoodi KZ, Ahmed N, Mir MA, Bhat B, Shafi A, Mansoor S, Rasool RS, Yaseen M, Dar ZA, Mir JI,
Andrabi SM, Ganai NA. Comparative transcriptomics unravels new genes imparting scab
resistance in apple (Malus x domestica Borkh.). Funct Integr Genomics. 2022; Epub ahead of
print.

24. Mary-Huard T, Aubert J, Mansouri-Attia N, Sandra O, Daudin JJ. Statistical methodology for the
analysis of dye-switch microarray experiments. BMC Bioinformatics. 2008;9:98.

25. Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG,

Lama S, Singh KB, Ljung K, Ecker JR, Goossens A, Millar AH, Van Aken O. A

MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-

responsive gene expression and jasmonate levels. Proc Natl Acad Sci U S A. 2019;116:23345-

23356.

26. Wasternack C, Feussner I. The oxylipin pathways: biochemistry and function. Annu Rev Plant Biol.

2018;69:363-386.

27. Li C, Schilmiller AL, Liu G, Lee GI, Jayanty S, Sageman C, et al. Role of beta-oxidation in jasmonate

biosynthesis and systemic wound signaling in tomato. Plant Cell. 2005;17:971-986.

28. Seo JS, Koo YJ, Jung C, Yeu SY, Song JT, Kim JK, Choi Y, Lee JS, Do Choi Y. Identification of a novel

jasmonate-responsive element in the AtJMT promoter and its binding protein for AtJMT

repression. PLoS One. 2013;8:e55482.

29. Jeong JS, Jung C, Seo JS, Kim JK, Chua NH. The deubiquitinating enzymes UBP12 and UBP13

positively regulate MYC2 levels in jasmonate responses. Plant Cell. 2017;29:1406-1424.

30. Birkenbihl RP, Diezel C, Somssich IE. *Arabidopsis* WRKY33 is a key transcriptional regulator of

hormonal and metabolic responses toward *Botrytis cinerea* infection. Plant Physiol.

2012;159:266-85.

31. Kaurilind E, Xu E, Brosché M. A genetic framework for H2O2 induced cell death in *Arabidopsis*

thaliana. BMC Genomics. 2015;16:837.

32. Nurmberg PL, Knox KA, Yun BW, Morris PC, Shafiei R, Hudson A, Loake GJ. The developmental

selector AS1 is an evolutionarily conserved regulator of the plant immune response. Proc

Natl Acad Sci U.S.A. 2007;104:18795-187800.

33. Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA et al. Pathogenesis-related proteins and

peptides as promising tools for engineering plants with multiple stress tolerance. Microb Res.

2018; 212-213:29-37.

34. Li J, Brader G, Palva ET. The WRKY70 transcription factor: a node of convergence for jasmonate-
mediated and salicylate-mediated signals in plant defense. Plant Cell. 2004;16:319–331.

35. Li J, Brader G, Kariola T, Palva ET. WRKY70 modulates the selection of signaling pathways in plant
defense. Plant J. 2006;46:477–491.

36. Wang D, Amornsiripanitch N, Dong X. A genomic approach to identify regulatory nodes in the
transcriptional network of systemic acquired resistance in plants. PLoS Pathog. 2006;2:e123.

37. Li J, Zhong R, Palva ET. WRKY70 and its homolog WRKY54 negatively modulate the cell wall-
associated defenses to necrotrophic pathogens in *Arabidopsis*. PLoS One. 2017;12:e0183731.

38. Genot B, Lang J, Berriri S, Garmier M, Gilard F, Pateyron S, et al. Constitutively active *Arabidopsis*
MAP kinase 3 triggers defense responses involving salicylic acid and SUMM2 resistance
protein. Plant Physiol. 2017;174:1238-1249.

39. Hu X, Reddy AS. Cloning and expression of a PR5-like protein from *Arabidopsis*: inhibition of
fungal growth by bacterially expressed protein. Plant Mol Biol. 1997;34:949-59.

40. Mestre P, Arista G, Piron MC, Rustenholz C, Ritzenthaler C, Merdinoglu D, Chich JF. Identification
of a *Vitis vinifera* endo-β-1,3-glucanase with antimicrobial activity against *Plasmopara*
viticola. Mol Plant Pathol. 2017;18:708-719.

41. Zhang SB, Zhang WJ, Zhai HC, Lv YY, Cai JP, Jia F, Wang JS, Hu YS. Expression of a wheat β-1,3-
glucanase in Pichia pastoris and its inhibitory effect on fungi commonly associated with
wheat kernel. Protein Expres Purif. 2019;154:134-139.

42. Santamaria M, Thomson CJ, Read ND, Loake GJ; The promoter of a basic PR1-like gene, AtPRB1,
from *Arabidopsis* establishes an organ-specific expression pattern and responsiveness to
ethylene and methyl jasmonate. Plant Mol Biol. 2001;47:641-52.

43. Truman W, Sreekanta S, Lu Y, Bethke G, Tsuda K, Katagiri F, Glazebrook J. The CALMODULIN-
BINDING PROTEIN60 family includes both negative and positive regulators of plant immunity.
Plant Physiol. 2013;163:1741-1751.

44. Boursiac Y, Lee SM, Romanowsky S, Blank R, Sladek C, Chung WS, Harper JF. Disruption of the vacuolar calcium-ATPases in *Arabidopsis* results in the activation of a salicylic acid-dependent programmed cell death pathway. Plant Physiol. 2010;154:1158-71.

45. Huang J, Sun Y, Orduna AR, Jetter R, Li X. The Mediator kinase module serves as a positive regulator of salicylic acid accumulation and systemic acquired resistance. Plant J. 2019;98:842-852.

46. Wang W, Tang W, Ma T, Niu D, Jin JB, Wang H, Lin R. A pair of light signaling factors FHY3 and FAR1 regulates plant immunity by modulating chlorophyll biosynthesis. Journal of Integrative Plant Biol. 2015;58:91-103.

47. Song JT, Koo YJ, Seo HS, Kim MC, Choi YD, Kim JH. Overexpression of AtSGT1, an *Arabidopsis* salicylic acid glucosyltransferase, leads to increased susceptibility to *Pseudomonas syringae*. Phytochemistry. 2008;69:1128-34.

48. Vlot AC, Dempsey DA, Klessig DF. Salicylic acid, a multifaceted hormone to combat disease. Annu Rev Phytopathol. 2009;47:177-206.

49. Andreasson E, Jenkins T, Brodersen P, Thorgrimsen S, Petersen NH, Zhu S et al. The MAP kinase substrate MKS1 is a regulator of plant defense responses. EMBO J. 2005;24:2579-89.

50. Xia Y, Yu K, Gao QM, Wilson EV, Navarre D, Kachroo P, Kachroo A. Acyl CoA binding proteins are required for cuticle formation and plant responses to microbes. Front Plant Sci. 2012;3:224.

51. Boller T, Felix G. A renaissance of elicitors: perception of microbe-associated molecular patterns and danger signals by pattern recognition receptors. Annu Rev Plant Biol. 2009;60:379-406.

52. Vincill ED, Bieck AM, Spalding EP. Ca(2+) conduction by an amino acid-gated ion channel related to glutamate receptors. Plant Physiol. 2012;159:40-46.

53. Hou C, Tian W, Kleist T, He K, Garcia V, Bai F, Hao Y, Luan S, Li L. DUF221 proteins are a family of osmosensitive calcium-permeable cation channels conserved across eukaryotes. Cell Res. 2014;24:632-635.

54. Fonseca JP, Mysore KS. Genes involved in nonhost disease resistance as a key to engineer durable
1038   resistance in crops. Plant Sci. 2019;279:108-116.

1039   55. Ronzier E, Corratgé-Faillie C, Sanchez F, Prado K, Brière C, Leonhardt N, Thibaud JB, Xiong TC.
1040   CPK13, a noncanonical Ca2+-dependent protein kinase, specifically inhibits KAT2 and KAT1
1041   shaker K+ channels and reduces stomatal opening. Plant Physiol. 2014;166:314-26.

1042   56. Zou JJ, Li XD, Ratnasekera D, Wang C, Liu WX, Song LF, Zhang WZ, Wu WH. *Arabidopsis* CALCIUM-
1043   DEPENDENT PROTEIN KINASE8 and CATALASE3 function in abscisic acid-mediated signaling
1044   and H2O2 homeostasis in stomatal guard cells under drought stress. Plant Cell.
1045   2015;27:1445-1460.

1046   57. Förster S, Schmidt LK, Kopic E, Anschütz U, Huang S, Schlücking K, et al. Wounding-induced
1047   stomatal closure requires jasmonate-mediated activation of GORK K(+) channels by a Ca(2+)
1048   sensor-kinase CBL1-CIPK5 complex. Dev Cell. 2019;48:87-99.e6.

1049   58. Kadota Y, Shirasu K, Zipfel C. Regulation of the NADPH oxidase RBOHD during plant immunity.
1050   Plant Cell Physiol. 2015;56: 1472–1480.

1051   59. Adachi H, Yoshioka H. Kinase-mediated orchestration of NADPH oxidase in plant immunity. Brief
1052   Funct Genomics. 2015;14:253-259.

1053   60. Zhai L, Sun C, Feng Y, Li D, Chai X, Wang L, Sun Q, et al. AtROP6 is involved in reactive oxygen
1054   species signaling in response to iron-deficiency stress in *Arabidopsis thaliana*. FEBS Lett.
1055   2018;592:3446-3459.

1056   61. Gao X, Chen X, Lin W, Chen S, Lu D, Niu Y, Li L, Cheng C, McCormack M, Sheen J, Shan L, He P.
1057   Bifurcation of *Arabidopsis* NLR immune signaling via Ca^{2+}-dependent protein kinases. PLoS
1058   Pathog. 2013;9:e1003127.

1059   62. Qi J, Wang J, Gong Z, Zhou JM. Apoplastic ROS signaling in plant immunity. Curr Opin Plant Biol.
1060   2017;38:92-100.

63. Bourdais G, Burdiak P, Gauthier A, Nitsch L, Salojärvi J, Rayapuram C et al. CRK consortium large-

scale phenomics identifies primary and fine-tuning roles for CRKs in responses related to

oxidative stress. PLoS Genet. 2015;11:e1005373.

64. Sun T, Nitta Y, Zhang Q, Wu D, Tian H, Lee JS, Zhang Y. Antagonistic interactions between two

MAP kinase cascades in plant development and immune signaling. EMBO Rep.

2018;19:e45324.

65. O'Brien JA, Daudi A, Butt VS, Bolwell GP. Reactive oxygen species and their role in plant defence

and cell wall metabolism. Planta. 2012;236:765-779.

66. Brosché M, Blomster T, Salojärvi J, Cui F, Sipari N, Leppälä J et al. Transcriptomics and functional

genomics of ROS-induced cell death regulation by RADICAL-INDUCED CELL DEATH1. PLoS

Genet. 2014;10:e1004112.

67. Cheng YT, Li Y, Huang S, Huang Y, Dong X, Zhang Y, Li X. Stability of plant immune-receptor

resistance proteins is controlled by SKP1-CULLIN1-F-box (SCF)-mediated protein degradation.

Proc Natl Acad Sci U.S.A. 2011;108:14694-14699.

68. Coll NS, Vercammen D, Smidler A, Clover C, Van Breusegem F, Dangl JL, Epple P. *Arabidopsis* type

I metacaspases control cell death. Science. 2010;330:1393-1397.

69. Lema-Asqui S, Vercammen D, Serrano I, Valls M, Rivas S, Van Breusegem F, et al. AtSERPIN1 is an

inhibitor of the metacaspase AtMC1-mediated cell death and autocatalytic processing in

planta. New Phytol. 2018;218:1156-1166.

70. Kawasaki T, Nam J, Boyes DC, Holt BF, Hubert DA, Wiig A, Dangl JL. A duplicated pair of

Arabidopsis RING-finger E3 ligases contribute to the RPM1-and RPS2-mediated

hypersensitive response. Plant J. 2005;44: 258-270.

71. Luo H, Laluk K, Lai Z, Veronese P, Song F, Mengiste T. The *Arabidopsis* Botrytis Susceptible1

interactor defines a subclass of RING E3 ligases that regulate pathogen and stress responses.

Plant Physiol. 2010;154:1766-1782.

72. Huby E, Napier JA, Baillieul F, Michaelson LV, Dhondt-Cordelier S. Sphingolipids: towards an

integrated view of metabolism during the plant stress response. New Phytol. 2020;225:659-

670.

73. Saucedo-Garcia M, Guevara-Garcia A, Gonzalez-Solis A, Cruz-Garcia F, VazquezSantana S,

Markham JE, et al. MPK6, sphinganine and the LCB2a gene from serine palmitoyltransferase

are required in the signaling pathway that mediates cell death induced by long chain bases in

Arabidopsis. New Phytol. 2011;191:943–957.

74. Veley KM, Maksaev G, Frick EM, January E, Kloepper SC, Haswell ES. *Arabidopsis* MSL10 has a

regulated cell death signaling activity that is separable from its mechanosensitive ion channel

activity. Plant Cell. 2014;26:3115-3131.

75. Guerringue Y, Thomine S, Frachisse JM. Sensing and transducing forces in plants with MSL10 and

DEK1 mechanosensors. FEBS Lett. 2018;592:1968-1979.

76. Zhao C, Nie H, Shen Q, Zhang S, Lukowitz W, Tang D. EDR1 physically interacts with MKK4/MKK5

and negatively regulates a MAP kinase cascade to modulate plant innate immunity. PLoS

Genet. 2014;10:e1004389.

77. Lee MH, Jeon HS, Kim HG, Park OK. An *Arabidopsis* NAC transcription factor NAC4 promotes

pathogen-induced cell death under negative regulation by microRNA164. New Phytol.

2017b;214:343-360.

78. Najafi J, Brembu T, Vie AK, Viste R, Winge P, Somssich IE, Bones AM. PAMP-INDUCED SECRETED

PEPTIDE 3 modulates immunity in *Arabidopsis*. J Exp Bot. 2020;71:850-864.

79. Khafif M, Balagué C, Huard-Chauveau C, Roby D. An essential role for the VASt domain of the

Arabidopsis VAD1 protein in the regulation of defense and cell death in response to

pathogens. PLoS One. 2017;12:e0179782.

80. Simon C, Langlois-Meurinne M, Didierlaurent L, Chaouch S, Bellvert F, Massoud K, et al. The

secondary metabolism glycosyltransferases UGT73B3 and UGT73B5 are components of redox

status in resistance of *Arabidopsis* to *Pseudomonas syringae* pv. *tomato*. Plant Cell Environ.

2014;37:1114-1129.

81. Bacete L, Mélida H, Miedes E, Molina A. Plant cell wall-mediated immunity: cell wall changes

trigger disease resistance responses. Plant J. 2018;93:614-636.

82. Miedes E, Vanholme R, Boerjan W, Molina A. The role of the secondary cell wall in plant

resistance to pathogens. Front Plant Sci. 2014;5:358.

83. Fich EA, Segerson NA, Rose JK. The plant polyester cutin: biosynthesis, structure, and biological

roles. Ann Rev Plant Biol. 2016;67:207-233.

84. Lim GH, Singhal R, Kachroo A, Kachroo P. Fatty acid- and lipid-mediated signaling in plant defense.

Annu Rev Phytopathol. 2017;55:505-536.

85. Joubès J, Raffaele S, Bourdenx B, Garcia C, Laroche-Traineau J, Moreau P, Domergue F, Lessire R.

The VLCFA elongase gene family in *Arabidopsis thaliana*: phylogenetic analysis, 3D modelling

and expression profiling. Plant Mol Biol. 2008;67:547-66.

86. Lee SB and Suh MC. Advances in the understanding of cuticular waxes in *Arabidopsis thaliana* and

crop species. Plant Cell Rep. 2015;34:557-572.

87. Li Y, Beisson F, Koo AJ, Molina I, Pollard M, Ohlrogge J. Identification of acyltransferases required

for cutin biosynthesis and production of cutin with suberin-like monomers. Proc Natl Acad Sci

U.S.A. 2007;104:18339-18344.

88. Yeats TH, Huang W, Chatterjee S, Viart HM, Clausen MH, Stark RE, Rose JK. Tomato Cutin

Deficient 1 (CD1) and putative orthologs comprise an ancient family of cutin synthase-like

(CUS) proteins that are conserved among land plants. Plant J. 2014;77:667-675.

89. Cui F, Brosche M, Lehtonen MT, Amiryousefi A, Xu E, Punkkinen M, Valkonen JPT, Fujii H,

Overmyer K. Dissecting abscisic acid signaling pathways involved in cuticle formation. Mol

Plant. 2016;9: 926-938.

90. Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q**.** Response of plant secondary metabolites to

environmental factors. Molecules. 2018;23i:e762.

91. Tetali S. Terpenes and isoprenoids: a wealth of compounds for global use. Planta. 2019;249:1-8.

92. Wang K, Senthil-Kumar M, Ryu CM, Kang L, Mysore KS. Phytosterols play a key role in plant innate

immunity against bacterial pathogens by regulating nutrient efflux into the apoplast. Plant

Physiol. 2012;158:1789-1802.

93. Falcone Ferreyra ML, Emiliani J, Rodriguez EJ, Campos-Bermudez VA, Grotewold E, Casati P. The

identification of maize and *Arabidopsis* type I FLAVONE SYNTHASEs links flavones with

hormones and biotic interactions. Plant Physiol. 2015;169:1090-107.

94. Wang N, Xu H, Jiang S, Zhang Z, Lu N, Qiu H, Qu C, Wang Y, Wu S, Chen X. MYB12 and MYB22 play

essential roles in proanthocyanidin and flavonol synthesis in red-fleshed apple (*Malus*

sieversii f. *niedzwetzkyana*). Plant J. 2017a;90:276-292.

95. Zhai R, Zhao Y, Wu M, Yang J, Li X, Liu H, Wu T, Liang F, Yang C, Wang Z, Ma F, Xu L. The MYB

transcription factor PbMYB12b positively regulates flavonol biosynthesis in pear fruit. BMC

Plant Biol. 2019;19:85.

96. Wang XC, Wu J, Guan ML, Zhao CH, Geng P, Zhao Q. *Arabidopsis* MYB4 plays dual roles in

flavonoid biosynthesis. Plant J. 2020;101:637-652.

97. Fernández-Pérez F, Vivar T, Pomar F, Pedreño MA, Novo-Uzal E. Peroxidase 4 is involved in

syringyl lignin formation in *Arabidopsis thaliana*. J Plant Physiol. 2015a;175:86-94.

98. Zhao Q, Nakashima J, Chen F, Yin Y, Fu C, Yun J, Shao H, Wang X, Wang ZY, Dixon RA. Laccase is

necessary and nonredundant with peroxidase for lignin polymerization during vascular

development in *Arabidopsis*. Plant Cell. 2013;25:3976-3987.

99. Cosio C, Ranocha P, Francoz E, Burlat V, Zheng Y, Perry SE, Ripoll JJ, Yanofsky M, Dunand C. The

class III peroxidase PRX17 is a direct target of the MADS-box transcription factor AGAMOUS-

LIKE15 (AGL15) and participates in lignified tissue formation. New Phytol. 2017;213:250-263.

100. Tokunaga N, Kaneta T, Sato S, Sato Y. Analysis of expression profiles of three peroxidase genes

associated with lignification in *Arabidopsis thaliana*. Physiol Plantarum. 2009;136:237-249.

101. Fernández-Pérez F, Pomar F, Pedreño MA, Novo-Uzal E. The suppression of AtPrx52 affects

fibers but not xylem lignification in *Arabidopsis* by altering the proportion of syringyl units.

Physiol Plantarum. 2015b;154:395-406.

102. Vilanova L, Teixidó N, Torres R, Usall J, Viñas I. The infection capacity of *P. expansum* and *P.*

digitatum on apples and histochemical analysis of host response. International Journal of

Food Microbiol. 2012;157:360-367.

103. Fink W, Haug M, Deising H, Mendgen K. Early defence responses of cowpea *(Vigna sinensis* L.)

induced by non-pathogenic rust fungi. Planta. 1991;185:246-254.

104. Yu Y, Yu Y, Cui N, Ma L, Tao R, Ma Z, Meng X, Fan H. Lignin biosynthesis regulated by CsCSE1 is

required for Cucumis sativus defence to Podosphaera xanthii. Plant Physiol Biochem.

2022;186:88-98.

105. Ranade SS, Seipel G, Gorzsás A, García-Gil MR. Enhanced lignin synthesis and ecotypic variation

in defense-related gene expression in response to shade in Norway spruce. Plant Cell

Environ. 2022;45:2671-2681.

106. Song PP, Zhao J, Liu ZL, Duan YB, Hou YP, Zhao CQ, Wu M, Wei M, Wang NH, Lv Y, Han ZJ.

Evaluation of antifungal activities and structure-activity relationships of coumarin derivatives.

Pest Manag. Sci. 2017;73:94-101.

107. Giesemann A, Biehl B, Lieberei R. Identification of scopoletin as a phytoalexin of the rubber tree

Hevea brasiliensis. J Phytopathol. 1986;117:373–376.

108. Muroi A, Ishihara A, Tanaka C, Ishizuka A, Takabayashi J, Miyoshi H, Nishioka T. Accumulation of

hydroxycinnamic acid amides induced by pathogen infection and identification of agmatine

coumaroyltransferase in *Arabidopsis thaliana*. Planta. 2009;230:517-527.

109. Dobritzsch M, Lübken T, Eschen-Lippold L, Gorzolka K, Blum E, Matern A et al. MATE

transporter-dependent export of hydroxycinnamic acid amides. Plant Cell. 2016;28:583-96.

110. Celoy RM, VanEtten HD. (+)-Pisatin biosynthesis: from (-) enantiomeric intermediates via an

achiral 7,2'-dihydroxy-4',5'-methylenedioxyisoflav-3-ene. Phytochem. 2014;98:120-127.

111. Vorwieger A, Gryczka C, Czihal A, Douchkov D, Tiedemann J, Mock HP, et al. Iron assimilation

and transcription factor controlled synthesis of riboflavin in plants. Planta. 2007;226:147-

158.

112. Trapalis M, Li SF, Parish RW. The *Arabidopsis* GASA10 gene encodes a cell wall protein strongly

expressed in developing anthers and seeds. Plant Sci. 2017;260:71-79.

113. Yamauchi Y, Hasegawa A, Mizutani M, Sugimoto Y. Chloroplastic NADPH-dependent alkenal/one

oxidoreductase contributes to the detoxification of reactive carbonyls produced under

oxidative stress. FEBS Lett. 2012;586:1208-1213.

114. Segond D, Dellagi A, Lanquar V, Rigault M, Patrit O, Thomine S, Expert D. NRAMP genes function

in *Arabidopsis thaliana* resistance to *Erwinia chrysanthemi* infection. Plant J. 2009;58:195-

207.

115. Lee MW, Jelenska J, Greenberg JT. *Arabidopsis* proteins important for modulating defense

responses to *Pseudomonas syringae* that secrete HopW1-1. Plant J. 2008;54:452-445.

116. Hong GJ, Xue XY, Mao YB, Wang LJ, Chen XY. *Arabidopsis* MYC2 interacts with DELLA proteins in

regulating sesquiterpene synthase gene expression. Plant Cell. 2012;24:2635-2648.

117. Rodriguez-Saona CR, Polashock J, Malo EA. Jasmonate-mediated induced volatiles in the

American cranberry, *Vaccinium macrocarpon*: from gene expression to organismal

interactions. Front Plant Sci. 2013;4:115.

118. Simanshu DK, Zhai X, Munch D, Hofius D, Markham JE, Bielawski J, et al. *Arabidopsis* accelerated

cell death 11, ACD11, is a ceramide-1-phosphate transfer protein and intermediary regulator

of phytoceramide levels. Cell Rep. 2014;6:388-399.

119. Fink L, Kwapiszewska G, Wilhelm J, Bohle RM. Laser-microdissection for cell type- and

compartment-specific analyses on genomic and proteomic level. Exp Toxicol Pathol. 2006;57

Suppl 2:25-259.

120. Gill US, Lee S, Mysore KS. Host versus nonhost resistance: distinct wars with similar arsenals.

Phytopathol. 2015;105:580-587.

121. Pieterse CM, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SC. Hormonal modulation

of plant immunity. Annu Rev Cell Dev Biol. 2012;28:489-521.

122. Tsuda K, Sato M, Stoddard T, Glazebrook J, Katagiri F. Network properties of robust immunity in

plants. PLoS Genet. 2009;5(12):e1000772.

1218

123. Faize M, Malnoy M, Dupuis F, Chevalier M, Parisi L, Chevreau E. Chitinases of *Trichoderma*

atroviridae induce scab resistance and some metabolic changes in two cultivars of apple.

Phytopathol. 2003;93:1496-1504.

124. Leblay C, Chevreau E, Raboin LM. Adventitious shoot regeneration from in vitro leaves of several

pear cultivars (*Pyrus communis* L.). Plant Cell Tissue Organ Cult. 1991;25:99-105.

125. Lespinasse Y, Durel CE, Parisi L, Laurens F, Chevalier M, Pinet C. A European project: D.A.R.E.

Durable apple resistance in Europe. Acta Hortic. 2000;538:197–200.

126. Chevalier M, Tellier M, Lespinasse Y, Bruynincks M, Georgeault S. Behaviour studies of new

races of *Venturia pirina* isolated from 'Conference' cultivar on a range of pear cultivars. Acta

Hortic. 2008a;800:817-824.

127. Parisi L, Lespinasse Y. Pathogenicity of *Venturia inaequalis* strains of race 6 on apple clone

(*Malus sp.*). Plant Dis. 1996;80: 1179-1183.

128. Chevalier M, Tellier M, Lespinasse Y, Caffier V. How to optimize the *Venturia* pirina inoculation

on pear leaves in greenhouse conditions? Acta Hortic. 2008b;800: 913-920.

129. Chevalier M, Lespinasse Y, Renaudin S. A microscopic study of different classes of symptoms

coded by the *Vf* gene in apple resistance to scab (*Venturia inaequalis*). Plant Pathol. 1991;40:

249–256.

130. Hoch HC, Galvani CD, Szarowski DH, Turner JN. Two new fluorescent dyes applicable for

visualization of fungal cell walls. Mycologia. 2005;97: 580-588.

131. Depuydt S, Trenkamp S, Fernie AR, Elftieh S, Renou J-P, Vuylsteke M, Holster M, Vereecke D. An

integrated genomic approach to define niche establishment by *Rhodococcus fascians* . Plant

Physiol. 2009;149: 1366–1386.

132. Celton JM, Gaillard S, Bruneau M, Pelletier S, Aubourg S, Martin-Magniette ML, Navarro L,

Laurens F, Renou JP. Widespread anti-sense transcription in apple is correlated with siRNA

production and indicates a large potential for transcriptional and/or post-transcriptional

control. New Phytol. 2014;203:287-99.

133. Smyth G. "limma: linear models for microarray data", in Bioinformatics and Computational

Biology Solutions Using R and Bioconductor, eds Gentleman R., Carey V., Dudoit S., Irizarry R.,

Huber W. New York, NY: Springer, 397–420.

134. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, Smyth GK. limma powers differential

expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res.

2015;43:e47.135. Pelletier S. AnaDiff: A tool for differential analysis of microarrays and

RNAseq (v4.3). 2022. Zenodo. https://doi.org/10.5281/zenodo.6477918

135. Pelletier S. AnaDiff: A tool for differential analysis of microarrays and RNAseq (v4.3). 2022.

Zenodo. https://doi.org/10.5281/zenodo.6477918

136. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E. The *Arabidopsis*

information resource: Making and mining the "gold standard" annotated reference plant

genome. Genesis. 2015;53:474-85.

137. Provart, N., Zhu, T. A Browser-based Functional Classification SuperViewer for Arabidopsis

Genomics. Currents in Computational Molecular Biology. 2003;271-272.

138. Thimm O, Bläsing O, Gibon Y, Nagel A, Meyer S, Krüger P, Selbig J, Müller LA, Rhee SY, Stitt M.

MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic

pathways and other biological processes. Plant J. 2004;37:914-939.

139. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic

Acids Res. 2001;29, e45.

140. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative

PCR and the 2−ΔΔCT method. Methods. 2001;25:402-408.

141. Vandesompele, J, De Preter, K, Pattyn, F, Poppe, B, Van Roy, N, De Paepe, A et al. Accurate

normalization of real-time quantitative RT-PCR data by geometric averaging of multiple

internal control genes. Genome Biol. 2002;3(7):00341-003411.

142. Vergne E. Expression data for apple Rvi6 host resistance and nonhost resistance against scab.

2020. https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE159179

143. Vergne E. Expression data for 1) pear host resistance allowed by transgenic expression of apple

Rvi6   resistance   gene   and   2)   nonhost   resistance   against   scab.   2020.

https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE159180

1274

Figure legends

1276

Fig. 1: **Macro- and microscopic observations of nonhost interactions.**

V. pyrina VP102 strain / pear 'Conference' (A) and *V. inaequalis* EUB05 strain / apple 'Gala' (B) 21

days symptoms are shown as classical ones of susceptible host interactions, to compare to nonhost

ones. Binocular observation 21 days after *V. inaequalis* EUB05 strain inoculation on 'Conference' (C)

and (D) and *V. pyrina* VP102 strain inoculation on 'Gala' (E). Wide field fluorescence observations of:

'Conference' 3 days (F) and 14 days (H and J) after *V. inaequalis* EUB05 strain inoculation, 'Gala' 3

days (G) and 14 days (I and K) after *V. pyrina* VP102 strain inoculation. Ap: appressorium, C: conidia,

Gf: germination filament, Pp: pin point

1285

1286   **Fig. 2**: **Functional categories of DEGs at 24 or 72 hpi during pear response to *V. inaequalis*.**

1287   The 1845 DEGs with TAIR names have been classified according to MapMan functional categories. In

1288   order to highlight the enriched ones, the numbers of DEGs per category have been normalized to the

1289 numbers of *Arabidopsis* genes in each MapMan category and bootstraps have been done to provide

1290 a confidence estimate for the accuracy of the output.

1291

Fig. 3: DEGs involved in hormonal pathways during pear/*V. inaequalis* nonhost interaction.

A: DEGs involved in JA pathway; B: DEGs involved in SA pathway. Genes written in red are induced,

genes written in blue are repressed. ACA11: autoinhibited $Ca2^+$-ATPase, calmodulin-activated $Ca2^+$

pumps at the plasma membrane, endoplasmic reticulum, and vacuole. ACBP6: acyl-CoA-binding

protein. ACX4: acyl-CoA-oxidase1. AS1/MYB91: Asymmetric leaves 1 transcription factor, CAMTA1:

calmodulin-binding transcription activator, CBP60a: calmodulin-binding protein 60a, EDS1: enhanced

disease suceptibility 1. FAR1: FAR-red impaired response 1. G-box: cis-element in the promoter. JAZ:

jasmonate-zim domain protein, JMT: jasmonic acid carboxyl methyltransferase. LOX: lipoxygenase,

MES1: methylesterase 1. MFP2: multifunctional protein 2. MKS1: MAP kinase substrate 1. MYC2:

transcription factor. NINJA: novel interactor of JAZ. PAD4: phytoalexin deficient 4. UGT74F1:

glucosyltransferase. PR1-like (with ATPRB1), PR2, PR3, PR4 (HEL and ATEP3), PR5, PR12:

pathogenesis-related proteins. ST2A: sulfotransferase 2A. TPL: TOPLESS co-repressor. UBP12:

ubiquitin-specific protease 12. WRKY: transcription factor.

1305

Fig. 4: **Scenario of major events observed at three first days of pear/*V. inaequalis* nonhost**

interaction.

On the left side, events observed in a typical cell, on the right side, events observed in guard cells of a

stomata. A: apoplasm, AP: appressorium, C: cuticle, CBL1: calcineurin B-like protein 1, CDPK: Ca_2^+-

dependent protein kinases, CRK: cysteine-rich receptor-like kinase, CY: cytoplasm, CW: cell wall, HAA:

hydroxycinnamic acid amines, HR: hypersensitive response, JA: jasmonic acid, MB: plasma

membrane, LCB: Long Chain/sphingoid Base components, MPK6: Mitogen activated protein kinase 6,

MSL10: mechano-sensitive like 10, N: nucleus, PH: penetration hypha, PR: pathogenesis related

proteins, RBOHB: respiratory burst oxidase homolog B, ROS: reactive oxygen species, S: stomata, SA:

salycilic acid, SC: simple coumarins, SP: spore.

1316

Fig. 5: Main DEGs involved in cutin and wax biosynthesis during pear/V. inaequalis nonhost

interaction.

In green the chloroplast, in brown the endoplastic reticulum (ER) and in yellow the nucleus. Genes

written in red are induced, genes written in blue are repressed. FAS: Fatty Acid Synthase complex to

which belong ACCD (carboxytransferase beta subunit of the Acetyl-CoA carboxylase complex), FabG

(β-ketoacyl ACP-reductase) and MOD1 (enoyl-ACP-reductase) functions. FAE: fatty acid elongase

complex. KCS4 (3-ketoacyl-CoA synthase 4) and ECR/CER10 (trans-2-enoyl-CoA reductase) belong to

the FAE complex. CER1 (octadecanal decarbonylase) and CER3 are implicated in aldehydes (CER1)

and alkanes (CER1 and 3) generation in waxes biosynthesis. In cutin monomers synthesis, the ω-

hydroxylation of C16:0 and C18:1 is catalyzed by cytochrome P450 monooxygenase (CYP86A) and

LACS-encoded acyl-CoA synthetase may be required either to synthesize 16-hydroxy 16:0-CoA, a

substrate for ω-hydroxylase, or for membrane transfer of monomers. Finally, the mature

monoacylglycerol cutin monomers are generated by transfer of the acyl group from acyl-CoA to

glycerol-3-phosphate by glycerol-3-phosphate acyltransferase (GPAT) enzymes such as GPAT8. Some

GDSL-lipases enzyme (such as At1g28600, At1g28660, At1g54790, At3g16370, At3g48460, AtCUS4:

At4g28780, At5g14450) are then functioning as cutin synthase and polymerize cutin

monoacylglycerols. Transcription factors such as MYB16 and SHN1 are positive regulators of wax and

cutin biosynthesis. Adapted from Xia et al., 2009, [83] and [84].

1335

Fig. 6: Main DEGs involved in the phenylpropanoid pathway during Pear / *V. inaequalis* nonhost

interaction.

Genes framed in red are induced, genes frames in blue are repressed. Framed in black, the detail of

genes involved in flavonoids production and found in this interaction. Abbreviations: 4CL, 4-

coumarate-CoA ligase; AACT, anthocyanin 5-aromatic acyltransferase; ANR, anthocyanidin reductase;

ANS, anthocyanin synthase; BGLC or BGLU, β-glucosidases; C3H, coumarate 3-hydroxylase; C4H,

cinnamate 4-hydroxylase; CAD, cinnamyl alcohol dehydrogenase; CCoAOMT, caffeoyl-CoA O-

methyltransferase; CCR, cinnamoyl-CoA reductase; CHI, chalcone isomerase; CHS, chalcone synthase;

COMT, caffeic acid 3-O-methyltransferase; CPK, calcium-dependent protein kinase ; DFR,

dihydroflavonol reductase; DMR6, downy mildiou resistant 6; F3H, flavanone 3-hydroxylase; F3'H

flavonoid 3'-hydroxylase; FLS, flavonol synthase; FNS, flavone synthase; GGT1, gamma-glutamyl

transpeptidase 1; GT, glucosyl transferase; HCT, hydroxycinnamoyl-CoA shikimate/quinate

hydroxycinnamoyl transferase; LAC, laccase; LAR, leucoanthocyanidin reductase; OMT1, O-

methyltransferase 1; PAL, phenylalanine ammonia-lyase; PER or PRX, peroxidase; TT7, transparent

testa 7; UGFT, UDP-glucose flavonoid-3-O-glucosyltransferase; UGT71D1, UDP-glycosyltransferase

71D1.

Fig. 1

Number of DEG	Frequence of DEG normalize to the number of Arabidopsis genes in each class in MapMan	± Bootstrap standard deviation	P_value (>0,05 in bold)	Fonctional class
4	4.55	2.156	**8.385e-03**	polyamine metabolism (Input set freq.: 0; 0)
17	3.91	0.883	**8.663e-07**	TCA / org transformation (Input set freq.: 0; 0)
5	3.25	1.476	**0.013**	Biodegradation of Xenobiotics (Input set freq.: 0; 0)
7	2.65	1	**0.011**	tetrapyrrole synthesis (Input set freq.: 0; 0)
5	2.27	1.019	**0.045**	C1-metabolism (Input set freq.: 0; 0)
47	1.92	0.275	**7.666e-06**	secondary metabolism (Input set freq.: 0.02; 0.01)
8	1.84	0.698	**0.039**	glycolysis (Input set freq.: 0; 0)
137	1.83	0.154	**2.650e-12**	signalling (Input set freq.: 0.07; 0.04)
12	1.76	0.578	**0.021**	minor CHO metabolism (Input set freq.: 0; 0)
3	1.76	0.875	0.153	OPP (Input set freq.: 0; 0)
92	1.64	0.135	**1.012e-06**	transport (Input set freq.: 0.04; 0.03)
9	1.62	0.54	0.052	major CHO metabolism (Input set freq.: 0; 0)
38	1.61	0.247	**1.191e-03**	lipid metabolism (Input set freq.: 0.02; 0.01)
16	1.61	0.364	**0.019**	nucleotide metabolism (Input set freq.: 0; 0)
22	1.54	0.322	**0.013**	amino acid metabolism (Input set freq.: 0.01; 0)
45	1.51	0.221	**1.559e-03**	hormone metabolism (Input set freq.: 0.02; 0.01)
66	1.44	0.185	**6.346e-04**	cell (Input set freq.: 0.03; 0.02)
16	1.41	0.329	**0.041**	PS (Input set freq.: 0; 0)
2	1.4	0.78	0.253	N-metabolism (Input set freq.: 0; 0)
1	1.4	1.281	0.363	S-assimilation (Input set freq.: 0; 0)
1	1.4	1.017	0.363	gluconeogenesis / glyoxylate cycle (Input set freq.: 0; 0)
42	1.39	0.217	**6.642e-03**	cell wall (Input set freq.: 0.02; 0.01)
231	1.38	0.084	**7.613e-08**	RNA (Input set freq.: 0.12; 0.09)
93	1.37	0.136	**3.589e-04**	stress (Input set freq.: 0.05; 0.03)
116	1.32	0.114	**3.215e-04**	misc (Input set freq.: 0.06; 0.04)
1	1.3	0.802	0.369	fermentation (Input set freq.: 0; 0)
15	1.3	0.353	0.064	redox (Input set freq.: 0; 0)
45	1.04	0.144	0.060	development (Input set freq.: 0.02; 0.02)
256	0.96	0.053	**0.022**	protein (Input set freq.: 0.13; 0.14)
4	0.89	0.422	0.196	Co-factor and vitamine metabolism (Input set freq.: 0; 0)
6	0.72	0.28	0.113	mitochondrial electron transport / ATP synthesis (Input set freq.: 0; 0)
473	0.72	0.029	**6.015e-20**	not assigned (Input set freq.: 0.25; 0.35)
30	0.17	0.028	**1.079e-44**	DNA (Input set freq.: 0.01; 0.09)

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Lignin biosynthesis

LAC11, LAC17   PER17, PER47, PRX52

Lignin ← ← Monolignols

CAD

COMT

F5H

CCR   +at2g23910

CCoAOMT

C3H HCT

Shikimate pathway

Chorismate

Proanthocyanidins

MYB4
MYB12

Flavan-3-ols

4CL   C4H   PAL

ANR   LAR

ANS   DFR   F3H   CHI   CHS

F3'H TT7
GT UGT71D1

FNS DMR6

p-Coumaryol CoA   Phenylalanine

Anthocyanin

Coumarine

Quercetin,
Quercetin-glycoside

AACT1/ACT1

flavones

BGLU16, 42,
47, BGLC3

Flavonoid biosynthesis

Hydroxycinnamic acids

Hydoxycinnamic acid amines

FNS DMR6   flavones

Naringenin

Dihydrokaempferol → Kaempferol

TT7 F3'H

Dihydroquercetin → Quercetin → Quercetin-glycoside

GT UGT71D1

Fig. 6