# Title

**Phenotypic and transcriptomic analyses reveal major differences between apple and pear scab**

**nonhost resistance**

# Authors

Vergne E.[1*], Chevreau E.[1*], Ravon E.[1], Gaillard S.[1], Pelletier S.[1], Bahut M.[2], Perchepied L.[1]

# Affiliations

[1] Univ Angers, Institut Agro, INRAE, IRHS, SFR QUASAV, F-49000 Angers, France

[2] Univ Angers, SFR QUASAV, F-49000 Angers, France

*E. Vergne and E. Chevreau made equal contributions to this work.

# Corresponding author

E.Vergne

Emilie.vergne@inrae.fr

# Abstract

**Background.** Nonhost resistance is the outcome of most plant/pathogen interactions, but it has

rarely been described in Rosaceous fruit species. Apple (*Malus x domestica* Borkh.) is a nonhost for

*Venturia pyrina*, the scab species attacking European pear (*Pyrus communis* L.). Reciprocally, *P.*

*communis* is a nonhost for *Venturia inaequalis*, the scab species attacking apple. The major objective

of our study was to compare the scab nonhost resistance in apple and in European pear, at the

phenotypic and transcriptomic levels.

**Results.** Macro- and microscopic observations after reciprocal scab inoculations indicated that, after

a similar germination step, nonhost apple/*V. pyrina* interaction remained nearly symptomless,

whereas hypersensitive reactions were observed during nonhost pear/*V. inaequalis* interaction.

Comparative transcriptomic analyses of apple and pear nonhost interactions with *V. pyrina* and *V.*

*inaequalis*, respectively, revealed considerable differences. Very few differentially expressed genes

were detected during apple/*V. pyrina* interaction, which is consistent with a symptomless type I

nonhost resistance. On the contrary, numerous genes were differentially expressed during pear/*V.*

*inaequalis* interaction, as expected in a type II nonhost resistance involving visible hypersensitive

reaction. Pre-invasive defense, such as stomatal closure, was detected, as well as several post-

invasive defense mechanisms (apoplastic reactive oxygen species accumulation, phytoalexin

production and alterations of the epidermis composition). In addition, a comparative analysis

between pear scab host and nonhost interactions indicated that, although specificities were

observed, two major defense lines were shared in these resistances: cell wall and cuticle

modifications and phenylpropanoid pathway induction.

**Conclusion.** This first deciphering of the molecular mechanisms underlying a nonhost scab resistance

in pear offers new possibilities for the genetic engineering of sustainable scab resistance in this

species.

**Keywords:** apple, pear, nonhost resistance, transcriptomics

# Background

Apple (*Malus domestica* Borkh.) and European pear (*Pyrus communis* L.) are two closely related

species belonging to the *Rosaceae* family. Reclassification of the *Rosaceae* placed both *Pyrus* and

*Malus* genera in the subfamily *Spiraeoideae*, tribe *Pyreae* and subtribe *Pyrinae,* this subtribe

corresponding to the long-recognized subfamily *Maloideae* [1]. Efforts to resolve relationships within

this subtribe have frequently failed, and Campbell et al [2] concluded that the genera of this subtribe

*Pyreae* have not diverged greatly genetically. The recent sequencing of the pear genome [3] allowed

a precise comparison with the apple genome [4] and led to the estimation of a divergence time

between the two genera of ≈ 5.4 − 21.5 million years ago. Furthermore, apple and pear genomes

share similar chromosome number (n=17), structure and organization.

Scab disease, caused by *Venturia* spp., affects several rosaceous fruit tree species. These

hemibiotrophic pathogens can infect only a limited host-range during their parasitic stage, but they

can overwinter as saprophytes in the leaf litter of a larger range of plant species [5]. Scab disease is

caused by *V. inaequalis* on apple, by *V. pyrina* (formerly named *V. pirina* [6]) on European pear, and

by *V. nashicola* on Japanese (*P. pyrifolia* Nakai) and Chinese (*P. ussuriensis* Maxim) pears. Cross

inoculations of *Venturia* spp. on different rosaceous fruit trees indicates that these pathogens are

highly host specific, probably indicating a close co-evolution of these pathogens with their hosts [7].

A plant species unable to be successfully infected by all isolates of a pathogen species is considered

as a nonhost for this pathogen. Nonhost interactions of *Venturia* spp. on apple and pear have rarely

been described. Microscopic observations have been made on *P. communis / V. nashicola* [8] as well

as *M. domestica / V. pyrina* and *P. communis / V. inaequalis* [5, 9]. In all cases, conidia germinated

and produced appressoria and runner hypheae, but failed to establish a network of stroma. No

macroscopic symptoms were visible.

Because of its durability, nonhost resistance has attracted numerous studies over the last decade,

which have uncovered its multiple and complex defense components. The underlying mechanisms of

nonhost resistance comprise pre-invasion resistance with preformed or induced cell-wall defenses,

metabolic defense with phytoanticipin or phytoalexin accumulation, pattern-triggered immunity (PTI)

as well as elicitor-triggered immunity (ETI) and various signaling pathways [10]. To our knowledge,

the molecular bases of scab nonhost resistance of apple and pear have never been investigated. The

objectives of our study were 1) to precisely describe nonhost resistance symptoms in *M. domestica /*

*V.pyrina* and *P. communis /* V. *inaequalis* interactions 2) to analyze the underlying molecular

mechanisms of both nonhost interactions through a transcriptomic study 3) to compare the

mechanism of host [11] and nonhost scab resistance in apple and European pear.

## Results and discussion

Variable symptoms of nonhost resistance

Nonhost interactions were observed in a test performed on 'Gala' and 'Conference', all inoculated by a *V. pyrina* strain (VP102) and a *V. inaequalis strain* (VI EUB05). At the macroscopic level, a total absence of sporulation was observed on all nonhost interactions (Table 1). The apple 'Gala' remained completely symptomless after *V. pyrina* inoculations (Fig. 1C). This is similar to the observation of Chevalier et al [9] after inoculation of 'Gala' with another *V. pyrina* strain. On the contrary, pear plants inoculated with *V. inaequalis* presented occasional pin points symptoms (Fig. 1A) and chlorotic lesions (Fig. 1B). Chlorotic lesions had already been observed by Chevalier et al [9] after inoculation of the pear 'Pierre Corneille' with the *V. inaequalis* strain EUB04, but pin points had never been reported in this nonhost interaction. According to our observations, apple nonhost resistance could be classified as type I and pear as type II according to Mysore and Ryu [12] definition based on the absence/presence of visible HR reaction.

Table 1: Scab qualitative note of pear and apple lines inoculated with *V. pyrina* and *V. inaequalis.*

| | Percentage of plants in the different classes of symptoms, 42 days after inoculation | | | |
|---|---|---|---|---|
| Class of symptoms | *V. pyrina* strain VP102 | | *V. inaequalis* strain EUB05 | |
| | 'Conference' | 'Gala' | 'Conference' | 'Gala' |
| 0 | 0 | 100 | 90 | 0 |
| 1 | 0 | 0 | 5 | 0 |
| 2 | 0 | 0 | 5 | 0 |
| 3a | 0 | 0 | 0 | 0 |
| 3b | 0 | 0 | 0 | 0 |
| 4 | 100 | 0 | 0 | 100 |

Class 0: absence of symptoms
Class 1: hypersensitivity (pin points)
Class 2: resistance (chlorotic lesions, slight necrosis, crinkled aspect)
Class 3a: weak resistance (necrotic or chlorotic lesions with occasional very light sporulation)
Class 3b: weak susceptibility (clearly sporulating chlorotic or necrotic lesions
Class 4: susceptibility (sporulation only)

At the microscopic level, three days after inoculation, there was no clear difference between host

and nonhost interactions: the conidia of *V. inaequalis* and *V. pyrina* germinated equally on both hosts

forming one or two appressoria (Fig. 1 D and F). However, 14 days after inoculation, there was a clear

reaction of the plant cells in contact with the appressoria (accumulation of red autofluorescent

compounds and enlargement of these cells), which could indicate very small scale hypersensitive

reactions (HR) (Fig. 1 E and G) in both plant species. No formation of subcuticular stroma and no

conidiogenesis were observed in the nonhost interactions, contrary to the host-resistance reactions

[11]. These observations are similar to the collapsed cells described by Chevalier et al [9] in apple and

pear nonhost reactions, and to the rare HR-like reactions observed by Stehmann et al [5] on apple

inoculated by *V. pyrina*.

Our results indicate that the leaf surface morphology of apple and pear is equally compatible with *V.*

*pyrina* and *V. inaequalis* conidia germination, without specific inhibition at this stage. Recognition

probably occurs only at the appressorium site, leading to the cellular reactions observed. These

reactions were limited to a few cells without visible symptoms in apple / *V. pyrina* interaction, but

extended and produced macroscopic symptoms in pear / *V. inaequalis* interaction.

Different patterns of global gene expression in nonhost resistance in pear versus apple

Differentially expressed genes (DEGs) were analyzed by comparing transcript abundance in leaves

between T0 and 24 hours post inoculation (hpi) and between T0 and 72hpi, in the nonhost

interactions 'Gala' / *V. pyrina* VP102 and 'Conference' / *V. inaequalis* EUB05. This kinetic

experimental design is open to criticism because it does not exclude in results data corresponding on

one hand to genes responding not to the infection but to the inoculation method (spray of water),

and on the other hand to genes whose expression varies due to leaves ageing. Water spray is

effectively perceived as a mechanical stimulus by plants, the Arabidopsis response being largely

regulated by the JA pathway induction under the control of MYC2/MYC3/MYC4 transcription factors

[13]. But this response is transient and really fast as most of genes differently regulated peak within

30 minutes regain untreated transcriptional levels within 3 hours [13]. It seems therefore very

unlikely that our analysis at 24 and 72hpi includes genes responding to water spray. Resistance due

to leaves ageing i.e. ontogenic resistance has been investigated in *Malus-Venturia* pathosystem at 72

and 96hpi, and 5 genes have been identified whose modulation could be linked to this resistance

[14]. None of these apple genes, or homologs in pear, were found differentially expressed at 72hpi in

our interactions (Table S1), which argue against the presence of ageing responding genes in our

results. In total, 60 DEGs in apple and 1857 DEGs in pear were identified, which amounts to 0.19 % of

all apple genes on the apple AryANE v2.0 microarray, and 4.23 % of all pear genes on the Pyrus v1.0

microarray (Table 2).

**Table 2. Number of DEGs identified during apple and pear nonhost response to *V. pyrina* and *V.***

***inaequalis***

| | 'Gala' / VP102 | | 'Conference' / EUB05 | |
|---|---|---|---|---|
| | 24hpi | 72hpi | 24hpi | 72hpi |
| Total # of DEGs* | 49 | 11 | 1570 | 364 |
| DEGs in % of all genes on the microarray** | 0.16 | 0.03 | 3.58 | 0.83 |
| % upregulated DEGs | 67.3 | 36.4 | 74.5 | 25.5 |
| % of downregulated DEGs | 32.7 | 63.6 | 25.5 | 74.5 |
| | | | | |
| % of DEGs without TAIR name | 27.1 | 30.4 | 0.70 | 1.09 |

137    *: DEGs numbers were calculated using the p-adj values ≤ 0.01 as selection threshold
138    **: 31311 genes on the apple Ariane V2 microarray, 43906 genes on the pear V1 microarray
139

The very small number of DEGs detected in the apple nonhost interaction at 24 or 72hpi is in

agreement with the total absence of macroscopic symptoms observed during this interaction.

However, at the microscopic level, small HR-like reactions were detected in the apple / *V. pyrina*

interaction at 14 days post inoculation. Because theses reactions involve only a few cells in the

leaves, the changes in gene expression are probably below the threshold of DEG detection applied in

this experiment. It is also possible that the mild response of apple to *V. pyrina* occurs later than

72hpi, between 72hpi and 6 days after inoculation. Indeed, Chevalier et al [9] observed these rare HR

from 6 days post inoculation, with no more evolution until 14 days after inoculation. A longer time (6

days post inoculation) would have been necessary to conclude on this last hypothesis.

On the contrary, the number of DEGs detected during the pear / *V. inaequalis* interaction is in the

same order of magnitude as the number of DEGs detected during pear host resistance to *V. pyrina*

(see [11]). This is in agreement with the frequent observation of macroscopic symptoms of resistance

(chlorotic lesions or pin points) in this interaction. Among the 1857 pear DEGs, 80.2 % were only

detected at 24hpi and 15.4 % only at 72hpi, whereas 4.2 % were upregulated or down regulated

similarly at both time points of the kinetics. Among all the pear DEGs observed at 24 and 72hpi, the

proportion of up-regulated DEGs was higher (68.8 %) than the proportion of downregulated DEGs

(31.2 %). Using MapMan to map the DEGs TAIR names, we observed that the main functional

categories represented in this set of DEGs were similar to those observed during pear host resistance

to *V. pyrina* (see [11]): protein, RNA, signaling, transport and cell cycle (Fig. 2).

To basically validate the transcriptomic data, 12 DEGS with varied ratios (between -1.9 and 2.9) have

been tested by quantitative RT-PCR (qPCR; Table S2), on the two biological repeats used for

transcriptomic analyses. Considering the weak number of DEGs found for apple in this study, we only

tested two of them in qPCR. As seen in Table 2 for pear, at 24hpi, a majority of DEGs are up-regulated

and at 72hpi, a majority of DEG are down-regulated. qPCR was then performed essentially on DEGs

with positive ratios at 24phi and negative ratios at 72hpi (Table S2). The qPCR results confirmed the

induced or repressed status of all tested DEGs.

Weak involvement of hormone signaling pathways classically associated to resistance

Pear DEGs were found that indicate that the jasmonic acid (JA) pathway was repressed. The JA

biosynthesis and metabolic conversions were reviewed by Wasternack et al [15]. In our data, at

24hpi, the first step of JA biosynthesis, that is the conversion of linoleic acid in 12-oxo-phytodienoic

acid (OPDA), is compromised: six out of seven lipoxygenases (LOX) (three LOX1, two LOX2 and two

LOX5) are repressed, the last one being induced (Fig. 3). OPDA produced in the chloroplast is then

transported to the peroxisome for subsequent conversion to JA via the action of OPR3 (12-oxo-

phytodienoic acid reductase) and β-oxidation enzymes (reviewed in [16] and in [15]). In pear, three

β-oxidation enzymes were found activated more or less rapidly: ACX4 (24hpi), MFP2 (72hpi) and the

thioesterase homolog to *At2g29590* (72hpi), which suggests that constitutive OPDA stocks were

turned into JA. But the early and long-lasting induction of *JMT* and *ST2A* genes is in favor of a rapid

conversion of JA in inactive compounds, *JMT* induction being reinforced by *BBD1* repression (24hpi).

*BBD1* is actually known as a negative regulator of *JMT* [17].

The defense response depending on JA was also clearly repressed in pear (Fig. 3). The transcription

activator MYC2 of JA-induced genes is known to be repressed by its interaction with JAZ proteins

(reviewed in [15]), and two *JAZ1* and one *JAZ3* coding genes were found activated at 24hpi in pear.

UBP12 is known as a stabilizer of MYC2 [18]. In our data, *UBP12* was found repressed at 72hpi, which

reinforces the inactivation of MYC2. WRKY33 is known as an activator of the JA defense pathway [19]

and WRK70 [20] or AS1 (or MYB91; [21]) as inhibitors, and among JA-responsive proteins, the

pathogenesis-related PR3, PR4 and PR12 act downstream MYC2 activation [22]. In our data,

accordingly with the repression of the activator WRKY33 and the activation of the inhibitors WRK70

and AS1, some JA-responsive genes were also found repressed, such as the chitinase coding genes

*PR4* (also called *HEL*) and *ATEP3*. Furthermore, no DEGs were found for PR3 and PR12 functions. To

conclude, in the nonhost interaction between pear and *V. inaequalis*, some JA seems to be produced,

but rapidly converted in inactive compounds and the subsequent defense response is clearly

repressed.

Pear DEGs were found that seems to indicate that the salicylic acid (SA) pathway was slightly

engaged and rapidly repressed (Fig. 3). *WRKY70* was induced at 72hpi in our data. This transcription

factor is known as a negative regulator of SA biosynthesis but a positive regulator of SA-mediated

defense genes in Arabidopsis ([23]; [24]; [25]), among them *PR2*, *PR5* but not *PR1* [26]. WRKY33

which is known as a negative regulator of SA-responsive genes [27], was also repressed at 72hpi in

our data. PR2 and 5 are well-known anti-fungal proteins ([28]; [29]; [30]). At 24hpi a *PR2*, two *PR2-*

*like*, a *PR5* and a *PR5-like* coding genes were found induced in our work, another *PR2-like* and two

others *PR5-like* being repressed. The differential expression was maintained at 72hpi for only two of

the activated ones. Furthermore no DEG was found for the PR1 function but three *PR1-like* genes

were found repressed: *ATPRB1* and genes homolog to *At5g57625* and *At4g33720*. *ATPRB1* was

already reported as repressed by SA treatment [31]. In our data, the WRKY70 transcription factor was

induced later than the induced PR genes so we could imagine that induced PR genes were activated

by another precocious regulation, such as an oxidative burst (see below), rather than by WRKY70.

Furthermore, WRKY70 induction seems not sufficient to enable a long lasting induction of these

defense genes.

SA accumulation was also rather mixed. *CBP60a* [32], *ACA11* [33], *EICBP.B* (or *CATMA1*; [34]), all

three coding calcium-sensor proteins, are known as negative regulators of SA accumulation and

biosynthesis, as well as the light signaling factor *FAR1* [35] or the SA glucosyltransferase *UGT74F1*

which convert SA in inactive SA 2-O-beta-D-glucoside or the glucose ester of SA [36]. On the contrary

*EDS1*, *PAD4* (reviewed in [37]) and *MKS1* [38] are known as positive regulators of SA accumulation. In

our data, the repression of *CBP60a*, *ACA11* and *EICBP.B* genes sustained a SA biosynthesis and

accumulation. In addition, *EDS1* activation allowed to consider a positive feedback loop likely to

potentiate SA action via EDS1 cytosolic homodimers, even though *PAD4* was repressed. But, as well

as *WRKY70* induction, the repression of the MAPK *MKS1* and the activation of the light signaling

factor *FAR1* (2 times) or the SA glucosyltransferase *UGT74F1* were in favor of less free SA. Concerning

SAR, *MES1* is known as required in healthy systemic tissues of infected plants to release the active SA

from methyl-SA, which serves as a long-distance signal for systemic acquired resistance (SAR) [37]

and *ACBP6* may be involved in the generation of SAR inducing signal(s) [39]. In our data, SAR seemed

compromised given the repression of *ACBP6* at 24hpi and *MES1* at 72hpi. To conclude, in the nonhost

interaction between pear and *V. inaequalis*, the SA pathway could be engaged but transiently and

presumably reduced to the few infection sites and not spread by SAR in healthy systemic tissues.

Calcium influx and reactive oxygen species (ROS) production act as secondary messengers

and lead to stomatal closure

Early responses of plants upon pathogen perception include calcium influx and ROS production,

which both act as secondary messengers ([40], reviewed in [10]). Three pear DEGs were found that

indicate early increased cytosolic calcium level. The CSC (Calcium permeable Stress-gated cation

Channel) ERD4 (found two times) and the two glutamate receptors GLR3.4 and GLR2.7, are known as

calcium permeable channels ([41]; [42]). They were induced at 24hpi in our data. An increased

cytosolic calcium level can lead to a pre-invasive defense response by stomatal closure and promote

the post-invasive defense response ROS accumulation [43].

Calcium influx has been reported to promote stomatal closure through the regulation of potassium

flux and the activation of anion channels in guard cells (reviewed in [10]). The stomata closure is

known to be induced via the inhibition of inward potassium currents which is achieved via activation

of calcium dependent protein kinases (CDPK) such as CPK13 and CPK8/CDPK19 ([44]; [45]); but also

via activation of CBL1 of the CBL1-CIPK5 complex, which activates the GORK potassium outward

channel [46]. CPK13, CPK8/CDPK19 and CBL1 were all activated at 24 hpi in our data.

A NADPH oxidase *RBOHB* (respiratory burst oxidase homologs, RBOH) is early (i. e. 24hpi) and

sustainably (through to 72hpi) induced in the pear/*V. inaequalis* nonhost interaction suggesting a

rapid and maintained apoplastic ROS production. Indeed, the apoplastic ROS are mainly produced by

plasma membrane localized NADPH oxidases, cell wall peroxidases and amine oxidases [47]. In

addition, posttranslational regulation of RBOH is required for its activation and ROS production.

Calcium, phosphatidic acid, and direct interactors such as Rac1 GTPase and RACK1 (Receptor for

Activated C-Kinase 1) have been reported to be positive regulators of RBOHs (reviewed in [48]). For

example, the Rac-like/ROP GTPase ARAC3 is known to interact with a RBOH to promote ROS

production [49]. In our data, RBOHB activity was also supported by the presence of positive

regulators such as Rac-like/ROP GTPase. The three Rac-like/ROP GTPase *ARAC1*, *ARAC3* and the

homolog of *At4g03100* were induced at 24hpi. CDPKs such as CPK1 are also known to activate RBOHs

in response to increased cytosolic calcium level [50]. But repression of *CPK1* in our data seems to

indicate that this way of activation did not function.

In response to abscisic acid (ABA) or microbe-associated molecular pattern (MAMP) immunity,

stomatal closure is known to be regulated by apoplastic ROS production (reviewed in [51]) and

cysteine-rich receptor-like kinases (CRK) are also known to be elements between ROS production and

downstream signaling leading to stomatal closure, sometimes activated (CRK10), sometimes

inhibited (CRK2 and CRK29; [52]). Three DEGs coding for CRK were found in our data and the

repression of *CRK2* and *CRK29* (found two times) was consistent with the stomata closure previously

found, but the repression of *CRK10* (found two times) was not. Beyond closure, inhibition of stomatal

development could be seen as an extreme defense. YODA (found two times) and MPK6 (found two

times) MAPKs belong to a pathway involved in the negative regulation of stomata development [53].

These two genes were induced early in our data.

To conclude, in pear/*V. inaequalis* nonhost interaction, a calcium influx leads to the development of

the stomatal closure pre-invasive defense, but also promotes a post-invasive defense: apoplastic ROS

accumulation. Apoplastic ROS, acting themselves as messengers, come to strengthen the stomatal

closure (Fig. 4).

Transcription factors and sphingolipids maintain HR under control

ROS are known to mediate cellular signaling associated with defense-related gene expression,

hypersensitive response (HR) i. e. the programmed cell death (PCD) at the site of infection during a

pathogen attack, and phytoalexin production [54]. *Arabidopsis thaliana* RCD1 regulator has been

proposed to positively regulate cell death in response to apoplastic ROS by protein-protein

interactions with transcription factors (reviewed in [55]) and WRKY70 and SGT1b were identified as

cell death positive regulators functioning downstream of RCD1 [55]. *RCD1* and *WRKY70* genes were

found induced in our data, at 24hpi and 72hpi respectively.

In Arabidopsis, the F-box protein CPR1, in association with the Skp1-Cullin-F-box (SCF) ubiquitin ligase

complex, targets for degradation NLR (nucleotide-binding domain and leucine-rich repeats containing

proteins) resistance protein such as SNC1, RPM1 or RPS2, to prevent overaccumulation and

autoimmunity (reviewed in [56]). A *Skp1-like* (*ASK19*; 72hpi) gene and *CPR1* (24hpi) gene were found

induced in our data. A gene coding for RPM1 function was also found repressed at 24hpi. These

results are in favor of the hypothesis that NLR receptors do not take part in the HR development

observed in the pear/*V. inaequalis* nonhost interaction (Fig. 1A). In addition, the induction of an

*AtSerpin1* gene homolog at 24hpi (found two times) in our data is consistent with that hypothesis.

Indeed, AtMC1 is a pro-death caspase-like protein required for full HR mediated by intracellular NB-

LRR immune receptor proteins such as RPP4 and RPM1 [57] and AtSerpin1 is a protease inhibitor

which block AtMC1 self-processing and inhibit AtMC1-mediated cell death [58].

The differential expression of two others components of the proteasome pathway is in favor of an HR

development: the induction of the *RIN3* ubiquitin E3 ligase (24hpi) and the repression of the *BRG3*

ubiquitin E3 ligase (24hpi). Indeed, RIN3 is known as positive regulator of RPM1 dependent HR [59].

And BRG3 is known as a negative regulator of HR in plant/necrotrophic pathogen interactions [60].

Sphingolipids are involved in the control of PCD, either as structural components of membranes but

also as initiators in the cell death regulatory pathway. According to Huby et al [61], free ceramides

and long chain/sphingoid base components (LCBs) are able to trigger cell death, via ROS production,

whereas their phosphorylated counterparts, ceramide phosphates and long chain base phosphate

components (LCB-Ps) promote cell survival. The induction of PCD by LCB is based on the activation of

protein kinases, among them MPK6 [62]. As already mentioned, *MPK6* was found early induced in

our data and we found numerous DEGs in the nonhost interaction between pear and *V. inaequalis*

that indicate the presence of free ceramides and LCB, which possibly participate to the HR

development. Free LCB presence is demonstrated by the activations of *SBH1* (24hpi), *SLD1* (24hpi)

and another sphingolipid Δ8 long-chain base desaturase homolog to *At2g46210* (24hpi; found two

times), and their relative conversion in ceramides is demonstrated by the differential expressions of

the ceramide synthases *LOH2* (repressed at 24hpi) and *LOH3/LAG13* (induced at 24 and 72hpi). LCB

non-conversion in phosphorylated counterparts is shown by the *AtLCBK1* repression (72hpi) and free

ceramides maintenance is attested by their non-conversion in glycosyled ones given the repression

of a glucosyl ceramide synthase homolog to *At2g19880* (24hpi).

The differential expression of numerous known regulators of HR in our data is again consistent with

the HR phenotype observed. The mechanosensor MSL10 and the calmodulin-activated $Ca^{2+}$ pump

(autoinhibited $Ca^{2+}$-ATPase [ACA]) ACA11 were found engaged: at 24hpi *MSL10* was induced and

*ACA11* was repressed. MSL10 is known as a positive regulator of cell death [63] and ACA11 is known

as a negative regulator of SA-dependent programmed cell death [33]. Their modulation is linked with

the noticed calcium influx discussed above ([33]; [64]). The participation of the SA pathway in the

development of the hypersensitive response could also be supported by the repression of *EDR1* (at

72hpi). Indeed, the MAPKKK EDR1 is known as a negative regulator of the SA-dependent HR

(reviewed in [65]).

Three other regulators of HR were found modulated in our data. The transcription factor *AS1*

(*MYB91*) was found induced at 24hpi. It is known as a positive regulator of HR and implicated in JA

pathway (reviewed in [20]). The transcription factor *WRKY40* was found repressed at 72hpi. It is

known as a negative regulator of HR [66] and implicated in PTI [67]. Another negative regulator of HR

is the lipid-binding domains containing protein VAD1 [68]. It was found repressed at 72hpi.

The behavior of two others genes in our data seems to indicate that the developed HR was contained

due to intracellular ROS production and damages. The function *UGT73B3* and *CAT2* were thus

activated (24hpi). UGT73B3 and CAT2 are known as restrictors of HR expansion via their action in

ROS scavenging (CAT2; [69]) or in detoxification of ROS-reactive secondary metabolites (UGT73B3;

[70]).

To conclude, in pear/*V. inaequalis* nonhost interaction, HR was spread out, in link with the calcium

influx, but especially following apoplastic ROS production and ROS production via free sphingolipids

accumulation and not via NLR receptors. Furthermore, the behavior of not less than eight regulators

indicate that the developed HR is under control (Fig. 4).

Cell wall carbohydrates content and cuticle composition are altered

The first obstacle encountered by host as well as nonhost pathogens attempting to colonize plant

tissues is the plant cell wall, which is often covered with a cuticle. Preinvasive penetration barrier, as

a preformed physical barrier, or as the onset place of defensive signaling pathways, is considered an

important factor, especially in nonhost resistance in which non adapted pathogens normally fail to

penetrate nonhost plant cells when blocked by the cell wall ([10]; [43]). Plant cell wall alterations, of

the carbohydrates or the phenolic components, either by impairing or overexpressing cell wall-

related genes, have been demonstrated to have a significant impact on disease resistance and/or on

abiotic stresses (reviewed in [71] and [72]).

We found numerous genes related to the cell wall with a modified expression during nonhost

interaction between pear and *V. inaequalis*, among them about thirty related to the biosynthesis or

the modification of carbohydrates. These genes are presented in table 3, except those related to the

lignin and other phenolic compounds, which will be discussed later. We saw in particular several

genes related to cellulose (8) and even more genes related to pectin (14) but no genes related to

callose.

Concerning these particular carbohydrate components, the model proposed by Bacete et al [71] is as

follows. Firstly, alterations in cellulose biosynthesis from primary or secondary cell wall trigger

specific defensive responses, such as those mediated by the hormones JA, ET or abscisic acid (ABA),

activate biosynthesis of antimicrobial compounds, but also might attenuate pattern triggered

immunity (PTI) responses. Secondly, alterations of cell wall pectins, either in their overall content,

their degree of acetylation or methylation, activate specific defensive responses, such as those

regulated by JA or SA, and trigger PTI responses, probably mediated by damage-associated molecular

patterns like oligogalacturonides. Thus, even though our results do not completely support a role of

these genes, we think that the modified expression of cell wall related genes during nonhost

interaction between pear and *V. inaequalis* is meaningful.

**Table 3: Main DEGs related to cell wall carbohydrates synthesis/modification detected during non-**

**host interaction pear/*V. inaequalis*.**

| | Gene | Action | Expression* |
|---|---|---|---|
| | Primary cell wall | | |
| | CSLA2 | synthesis | I |
| | PNT1 | synthesis | I |
| Cellulose | COBL2 | deposition (GPI-anchored protein) | R |
| | AtGH9A4 | catabolism | I |
| | XTR7 | loosening | I |
| Hemi-cellulose (xyloglucan) | At5g15490 | synthesis | I |
| | At3g42180 | synthesis | I |
| | At4g01220 | synthesis | I |
| | GHMP kinase | synthesis | I |
| | RHM1 | synthesis | R |
| | PME At2g45220 | methylesterification | I |
| | PME At2g46930 | methylesterification | I |
| Pectin | PME At3g05910 | methylesterification | R |
| | PME At1g02810 | methylesterification | R |
| | PME44 | methylesterification | R |
| | PG At3g16850 | depolymerisation | I |
| | PG At3g59850 | depolymerisation | I |
| | PG At4g13710 | depolymerisation | I |
| | PG At3g62110 | depolymerisation | R |
| | IDA | degradation | R |
| Arabinogalactan protein | AGP11 | – | I |
| | AGP1 | – | R |
| | Secondary cell wall | | |
| | CESA09 | synthesis | I |
| Cellulose | CESA10 | synthesis | I |
| | CSLG1 | synthesis | I |
| Hemi-cellulose (xylan) | FRA8 | synthesis | I |

| | | Undetermined | | |
|---|---|---|---|---|
| Expansin | EXP15 | loosening | I | |
| | EXPB3 | loosening | I | |
| Hemi-cellulose | ATFUC1 | modification | I | |
| | XTH33 | growth and assembling | R | |

361 *I: induced, R: repressed
362

Concerning the cuticle layer, most cuticles are composed largely of cutin, an insoluble polyester of

primarily long-chain hydroxy fatty acids. This lipophilic cutin framework is associated with

hydrophobic compounds collectively referred to as waxes. The cuticle is also thought to contain

varying amounts of intermingled cell wall polysaccharides and sometimes also a fraction termed

cutan (reviewed in [73]). Cutin monomers are synthesized by the modification of plastid-derived 16C

and 18C fatty acids in the endoplasmic reticulum (ER), yielding variously oxygenated fatty acid–

glycerol esters referred to as monoacylglycerols, which polymerize upon arrival at the growing cuticle

(Fig. 5, reviewed in [73]).

C16 and C18 fatty acids are also important precursors of cuticular wax synthesis (Fig. 5). Upon

transport to the ER, the C16 and C18 fatty acids are extended to form very-long-chain fatty acids

(VCLFAs; C>20), and this extension is carried out by the fatty acid elongase (FAE) complex located on

the ER membrane. The very-long-chain FAs are then converted into the varied cuticular waxes

(primary alcohols, aldehydes, alkanes, secondary alcohols, ketones) by many ways (reviewed in [74]).

Interestingly, we found three genes upregulated 24hpi belonging to the FAS (fatty acid synthase)

chloroplastic complex implicated in the production of the C16 precursor (Fig. 5): ACCD, FabG and

MOD1 (found two times). ACCD encodes the carboxytransferase beta subunit of the Acetyl-CoA

carboxylase complex which catalyzes the first committed step in fatty acid synthesis: the

carboxylation of acetyl-CoA to produce malonyl-CoA. FabG and MOD1 are respectively a β-ketoacyl

ACP-reductase and an enoyl-ACP-reductase which catalyze respectively the conversion of

acetoacetyl-ACP into β-hydroxyacyl-ACP and the second reductive step from enoyl-ACP to butyryl-

ACP (reviewed in [74]).

In the ER, the four functions we found related to waxes biosynthesis in our data were repressed at

24hpi: *KCS4* (found two times), *CER1* and *CER3*, or 72hpi: *ECR/CER10*. KCS4 and ECR/CER10 belong to

the FAE complex ([75]; [76]). The last two genes are implicated in aldehydes (*CER1*) and alkanes

(*CER1 and 3*) generation (reviewed in [74]). On the contrary, the eight genes we found connected to

cutin biosynthesis were induced at 24hpi except a gene homolog to *At5g14450*, which was induced

at 72hpi. One of them is a glycerol-3-phosphate acyltransferase (GPAT) coding gene: GPAT8, which

catalyzes the transfer of a fatty acid from coenzyme A (CoA) to glycerol-3-phosphate (Fig. 4; reviewed

in [73]). GPAT8 function in cutin formation has been functionally confirmed in association with

GPAT4 [77]. The seven others genes code GDSL-lipases enzyme (*At1g28600*, *At1g28660*, *At1g54790*,

*At3g16370*, *At3g48460*, *AtCUS4: At4g28780*, *At5g14450*), some of which have been shown to

function as cutin synthase (Fig. 4; [78]; reviewed in [73]) and polymerize monoacylglycerols.

We also found induced respectively at 24 and 72hpi two genes involved in waxes and cutin

biosynthesis positive regulation: *MYB16* and *SHN1*. The SHN genes (*SHN1–SHN3*), a set of three

largely redundant APETALA 2 family transcription factors from *A. thaliana*, are regulators of floral

cutin and epidermal cell morphology. SHN1 is regulated by the MYB family transcription factor

MYB106, which, along with its paralog MYB16, controls many aspects of cuticle and epidermis

formation in *A. thaliana* (reviewed in [79] and [73]).

Cutin and cuticular waxes play an important role in plant-insect and plant-microbe interactions.

Numerous Arabidopsis mutants in cutin and waxes biosynthetic or transport genes, such as Acyl-CoA

binding proteins (ACBP), show varying degrees of cuticle impairment, alterations in cutin and/or wax

composition, and defects in SAR (reviewed in [74]). We found *ACBP6* repressed at 24hpi. That

repression is not inconsistent with the previously described amplification of cutin biosynthesis and

polymerization, given that *acbp6* KO mutation is not associated with a defect in that pathway [39].

That repression is also consistent with the SAR repression observed above as the *acbp6* KO mutant

show compromised SAR [39].

To conclude, our analysis of nonhost pear/V. *inaequalis* interaction identified an alteration of the

cuticle composition with more cutin and less waxes synthesis. The increase in cutin polymerization

could lead to a thickening of the cuticular layer to prevent fungus penetration via its appressoria.

Secondary metabolism leads to G unit lignin polymerization and simple coumarin or

hydrocinnamic acid amine phytoalexins synthesis

As distinguished from primary metabolism, plant secondary metabolism refers to pathways and small

molecule products of metabolism that are non-essential for the survival of the organism. But they are

key components for plants to interact with the environment in the adaptation to both biotic and

abiotic stress conditions. Plant secondary metabolites are usually classified according to their

chemical structure. Several groups of large molecules, including phenolic acids and flavonoids,

terpenoids and steroids, and alkaloids have been implicated in the activation and reinforcement of

defense mechanisms in plants (reviewed in [80]).

Terpenoids and steroids, or isoprenoids, are components of both the primary and secondary

metabolisms in cells, and mono-, tri-, sesqui- and polyterpenes are considered as secondary

metabolites (reviewed in [81]). Our results on pear identified seven DEGs and five DEGs belonging to

the chloroplastic methylerythritol posphate (MEP) and to the cytosolic mevalonic acid (MVA)

pathway of isoprenoids production respectively (Table 4), which results, among others compounds,

in tri- and sesquiterpenes secondary metabolites. The majority of these genes contribute to produce

primary metabolites according to Tetali [81]. Except *SMT2*, that we found induced at 24hpi, there is

no report concerning a putative implication of others genes in plant biotic resistance. SMT2 is

involved in sterols production and *smt2* mutation was reported to compromise bacterial resistance in

*Nicotiana benthamiana* [82]. The hypothesis is that sterols regulate plant innate immunity against

bacterial host and nonhost infections by regulating nutrient efflux into the apoplast. *V. inaequalis* is a

hemi biotrophic pathogen which colonizes only the apoplast compartment at the beginning of the

interaction. strong relative induction of *SMT2* in our data could indicate that a similar mechanism of

nutrient efflux regulation via sterols could take place to limit the fungus growth in pear nonhost

resistance against *V. inaequalis*.

**Table 4: Main DEGs involved in biosynthetic pathways for terpenes and isoprenoids during pear/*V.***

***inaequalis* non-host interaction.**

| | Gene | Function | Expression* |
|---|---|---|---|
| Cytosolic MVA (mevalonic acid) pathway enzymes | *HMGS* | catalyze the second step of the pathway | R |
| | *HMGR1* | catalyze the third step of the pathway | R |
| | *SMT2* | sterols production | I |
| | *FLDH* | sesquiterpenes production | R |
| | *SQE2* | triterpenes production | I |
| | *DXR* | catalyzes the second step of the pathway | I |
| Chloroplastic MEP (methylerythritol posphate) pathway enzymes | *GG reductase* | chlorophylls production | R |
| | *VTE4* | tochopherols production | I |
| | *KAO1* | gibberellins production | R |
| | *PDS2* | plastoquinones production | I |
| | *LYC* | carotenoids production | I |
| | *PGGT1* | covalent attachment of a prenyl group to a protein | I |

In our data, the other DEGs that were linked to secondary metabolism belong to the

phenylpropanoid pathway production (Fig. 6). Among them we found four genes belonging to the

flavonoid production, all repressed, at 24hpi (*DFR* and *DRM6*) or 72hpi (*TT7* and *UGT71D1*). DFR

(dihydroflavonol reductase) is involved in flavan-3,4-ol production and TT7 (flavonoid 3' hydroxylase)

in dihydroquercetin production from dihydro-kaempferol, and UGT71D1 (glucosyl transferase) in

quercetin-glycoside        production        from        quercetin        (TAIR        database;

https://www.arabidopsis.org/index.jsp). DMR6 (flavone synthase) is involved in flavone production

from naringenin [83]. Thus flavonoid production does not seem to be favored, which is not consistent

with the induction of *MYB12* at 24hpi, but consistent with *MYB4* induction at 72hpi. MYB12 is

actually known as a positive regulator of flavonol biosynthesis in pear and apple fruits ([84]; [85])

whereas MYB4 is known as a negative regulator of this biosynthetic pathway [86].

Concerning the production of monolignols, precursors of lignin synthesis, some genes were found

induced, others repressed. We found *CYP98A3* and *CAD9* (found two times) induced at 24hpi and

*HCT*, *CCR1* and a gene homolog to *At2g23910* (found two times, one time repressed at 72hpi)

repressed at 24hpi Fig. 6). *CYP98A3* encode a C3H (coumarate 3-hydroxylase), *CAD9* encode a CAD

(cinnamyl alcohol dehydrogenase), HCT is an hydroxycinnamoyl-CoA shikimate/quinate

hydroxycinnamoyl transferase, *CCR1* encode a CCR (cinnamoyl-CoA reductase) and *At2g23910*

encode a CCR-related protein. (TAIR and KEGG databases (https://www.genome.jp/kegg/)).

Lignification is obtained by cross-linking reactions of the lignin monomers or by polymer–polymer

coupling via radicals produced by oxidases such as peroxidases [87] and laccases [88]. However,

while peroxidases are able to oxidize monolignols to produce H, G and S units of lignin, laccases only

generate G units [87]. In our data, we found two laccases induced at 24hpi: *LAC11* (found two times,

one time induced at 24 and 72hpi) and *LAC17* (found two times), and three peroxidases repressed at

24hpi: *PRX17*, *PER47* and *PRX52* (also repressed at 72hpi), which can be linked to lignin biosynthetic

process (Fig. 6). According to Zhao et al. [88], LAC11 and LAC17, along with LAC4, play a critical role in

lignification, and their results suggests that peroxidase and laccase do not serve redundant functions

in lignification, at least in the vascular tissues of the stem and root of Arabidopsis. Participation in

lignin formation has also been proved for PRX17 [89], PER47 [90] and PRX52 [91]. But there are

currently no reports about a possible involvement of all these genes in lignification linked to biotic or

abiotic stresses. Concerning non-host resistance, two reports describe lignin

accumulation/deposition involvement: one in apple fruit [92] and the other one in cowpea [93]. In

the latest, authors showed that preferentially generated lignin units in this nonhost interaction are G

units, just as it seems to be the case in our pear / *V. inaequalis* study. To summarize, it is tempting to

think that modifications of expression observed for genes linked to lignin polymerization are relevant

for the pear nonhost resistance against *V. inaequalis*, but further functional analysis should be

conducted to conclude.

The biosynthesis of two others types of phenylpropanoid compounds appears to be favored during

pear nonhost resistance against *V. inaequalis*: simple coumarin on one hand and hydroxycinnamic

acid amides on the other hand. We found four *BGLU*-like genes induced: *BGLU47* and *BGLC3* (at

24hpi), *BGLU16* (at 72hpi); *BGLU42* (at 24 and 72hpi) (Fig. 6). These β-glucosidases could be implied

in simple coumarin path production from the cinnamic acid (KEGG database). Some natural simple

coumarins are known as antifungal compounds *in vitro* and have been developed as fungicides [94].

Previous work on Hevea also reports the correlation between the resistance against pathogenic fungi

and the production of some coumarins, with antifungal activity *in vitro* [95]. We also found induced

at 24hpi the genes *AACT1/ACT1*, *ATPAO5* and genes homologs to *At4g17830* and *At4g38220* (Fig. 6).

AACT1/ACT1 catalyze the first specific step in branch pathway synthesizing hydroxycinnamic acid

amides from the p-Coumaroyl CoA or the feruloyl CoA and amines agmatine or putrescine [96].

Hydroxycinnamic acid amides are produced in response to pathogenic infections [96] and surface

exported. Hydroxycinnamic acid amides are reported to participate in Arabidopsis nonhost resistance

against *Phytophthora infestans* via their inhibitory activity on spore germination [97]. The three

others genes belong to the arginine biosynthesis path (homologs to *At4g1783* and *At4g38220*) and

the arginine and proline metabolisms which produce the amines agmatine and putrescine (*ATPAO5*)

(KEGG database). Agmatine is directly produced from arginine thanks to an ADC activity (arginine

decarboxylase) and putrescine can be produced from spermidine thanks to a PAO activity (polyamine

oxidase). ATPAO5 catalyzes the conversion of spermine in spermidine. The induction of these three

last genes is therefore consistent with the hypothesis of amines production in order to enable

hydroxycinnamic acid amides synthesis. The induction of *C4H* at 24hpi could also favor

hydroxycinnamic acid amides synthesis via p-Coumaroyl CoA biosynthesis promotion. C4H

(cinnamate 4-hydroxylase) catalyzes the production of p-Coumaric acid from Cinnamic acid and p-

Coumaric acid gives p-Coumaroyl CoA thanks to 4CL (4- coumarate-CoA ligase) (KEGG database).

Among the suite of defense components synthetized in nonhost as in host context, a chemical barrier

can be established via accumulation of a diverse array of secondary metabolites rapidly produced

upon pathogen infection, named phytoalexins, with toxic or inhibitory effects (reviewed in [10]).

Phytoalexins can be flavonoids, such as the pisatin of pea (in [98]) but also varied phenylpropanoid

compounds. In the nonhost interaction pear / *V. inaequalis*, the production of flavonoid type

phytoalexins does not seem to be favored, except simple coumarin and hydroxicinnamic acid amines.

Very limited transcriptomic modulation during apple / *V. pyrina* nonhost interaction

Only 60 DEGs were detected in the apple / *V. pyrina* nonhost interaction at 24 or 72hpi, in agreement

with the total absence of macroscopic symptoms and few cells engaged in an HR-like reaction

observed at the microscopic level. Among these 60 DEGs, 36 have no known function. Among the 24

remaining DEGs, nine DEGS could be relevant in apple / *V. pyrina* nonhost interaction in view of our

findings in pear / *V. inaequalis* nonhost interaction. *ORG2* (*BHLH038*), a putative integrator of various

stress reactions [99] was induced at 24hpi. Three genes were related to an oxidative stress: *GASA10*

was repressed at 24hpi and *NRAMP3* and *AOR* were induced at 24hpi. GASA proteins have been

suggested to regulate redox homeostasis via restricting the levels of OH$^{\cdot}$ in the cell wall [100]. The

repression of this gene is thus in favor of more OH$^{\cdot}$ in the cell wall. The oxidoreductase coding gene

*AOR* is known in the chloroplast to contribute to the detoxification of reactive carbonyls produced

under oxidative stress [101]. *NRAMP* genes function as positive regulators of ROS accumulation,

especially during Arabidopsis *Erwinia chrisanthemi* resistance [102]. The induction (at 24 and 72hpi)

of another gene suggests modifications at the cell wall level: *EXP8*, an expansin coding gene involved

in cell wall loosening (Tair database). We also found two genes related to hormone pathways, one

induced at 24hpi: *WIN1* and the other one repressed at 72hpi: *UBP12*. WIN1 is known as a negative

regulator of SA pathway [103] and UBP12 as a positive regulator of JA pathway via the stabilization of

MYC2 [18]. In link with the JA pathway, we also found *TPS21* induced at 24hpi. TPS21 is involved in

sesquiterpenes production and is promoted by JA signal via MYC2 [104]. TPS21 is especially involved

in the jasmonate-dependent defensive emission of herbivore-induced volatiles in cranberries [105].

Finaly the last DEG we found relevant in apple / *V. pyrina* nonhost interaction could promote HR via

ceramides accumulation. *ACD11* is repressed at 24hpi in our data. In *acd11* mutants, the relatively

abundant cell death inducer phytoceramide rises acutely [106].

Because nonhost resistance of apple against *V. pyrina* is of a type I, with a very limited number of

cells engaged in an HR-like reaction, it has not been possible for us to exhaustively describe how this

interaction is expressed at the transcriptomic level in the first three days of the interaction. Further

insight with later points of cinetic and more adapted techniques such as laser-assisted cell picking,

prior to micro arrays or RNA sequencing analysis (review in [107]) could provide more information in

the future.

Comparison of pear resistances against the host pathogen *V. pyrina* and the nonhost

pathogen *V. inaequalis*

Perchepied et al [11] performed a detailed transcriptomic analysis of the host resistance of pear

against *V. pyrina* strain VP102, deployed in a transgenic pear bearing the well-known apple *Rvi6*

resistance gene against *V. inaequalis.* Comparing this work to ours gives us the rare opportunity to

analyze similarities and differences between a host and a nonhost resistance in the same plant. Only

four transcriptomic studies involving pear/pathogen interactions have been published so far. Yan et

al [108] reported the modulation of expression of 144 pear genes after fruit treatment by

*Meyerozyma guilliermondii,* an antagonistic yeast used for biocontrol of natural pear fruit decay.

Zhang et al [109] similarly reported the modulation of expression of 1076 pear genes after treatment

with *Wickerhamomyces anomalus,* another biocontrol agent. Using RNA-seq, Wang et al. [110]

reported a major role of ethylene signalization during the compatible interaction between *P. pyrifolia*

and *Alternaria alternata,* a necrotrophic pathogen. Finally, Xu et al. [111] applied RNA-seq to

characterize the genes of *Penicillium expansum* activated after infection of pear fruits. None of these

studies can be directly compared to our work on host and nonhost scab pear resistance.

Concerning the recognition and early signaling steps of the interactions, many receptors and co-

receptors have been found induced in the host pear resistance, especially damage-associated

molecular patterns receptors such as RLK7, revealing that PTI and ETI must be engaged. We did not

find evidence of the mobilization of such receptors in the pear nonhost resistance. PTI and ETI

receptors are nonetheless reported as implicated in nonhost resistance (reviewed in [112] and [10]).

As we only analyzed post infection transcriptional modulations in the nonhost pear/*V. inaequalis*

interaction (at 24 and 72hpi), one hypothesis to explain the lack of PTI and ETI receptors in our data

could be that these receptors were already present as preformed defenses and not particularly

induced by the infection onset. In pear nonhost interaction, the earliest signaling pathways we were

able to highlight are calcium influx and apoplastic ROS production. Calcium signaling seems to be also

implicated in pear host resistance, but less obviously than in nonhost resistance.

Regarding the hormonal signaling pathways, the JA defense signaling pathway was found repressed

in pear nonhost resistance but quite activated in pear host resistance. The JA/ethylene (ET) defense

signaling pathway is known as an effective defense against necrotrophic fungi in Arabidopsis [113].

Thus, it is not surprising to find the JA pathway repressed in the development of the pear nonhost

resistance against the hemi-biotrophic pathogen *V. inaequalis*. But it is very interesting to find this

pathway rather induced in the development of the pear host resistance against the other hemi-

biotrophic pathogen *V. pyrina*. The SA signaling pathway is commonly seen as the classical one

triggered to resist biotrophic fungi in Arabidopsis [113], but only a little engagement in pear nonhost

resistance has been observed, SA signaling being repressed in pear host resistance. If this absence of

SA implication is quite unexpected in pear host resistance against a hemi-biotrophic fungus, it is

consistent with the report that the exact role of these key defense phytohormones is unclear in

nonhost resistance and remains to be established [43]. As shown by Tsuda et al [114], an explanation

for the hormone pathways behavior in pear host resistance could be that: as both the SA and JA/ET

pathways positively contribute to immunity, a loss of signaling flow through the SA pathway can be

compensated by a rerouting signal through the JA/ET pathways. In addition, independently of SA

signaling, but in positive connection with JA signaling, SAR seems to be engaged in distal tissues

during pear host resistance. To conclude, in pear host as well as nonhost resistances, classical

resistance hormones SA and JA/ET, and the correlative PR gene defenses, seems differently involved

than in Arabidopsis.

The carbohydrate content of the cell-wall is modified in response to the attacks by the pathogens.

Regarding cell-wall and cuticle, in pear host as well as nonhost resistances, important modifications

were highlighted. Similar modifications affected the cellulose and mainly the pectin contents, but no

callose production was observed. Regarding cuticle, waxes production was induced in host resistance

whereas it was repressed in nonhost resistance, in favor of cutin production / polymerization, which

was also induced in host resistance. To conclude, as a first obstacle encountered by host, as well as

nonhost, pathogens attempting to colonize plant tissues, the plant cell wall and its cuticle seem to

play a leading role in both pear host and nonhost resistances.

Finally, the production of secondary metabolites and phenylpropanoids compounds in particular,

seems to be a major line of defense, in pear host as well as nonhost resistances, but with

divergences. If lignin and flavonoid productions are preponderant in pear host resistance against *V.*

*pyrina*, lignin implication in pear nonhost resistance is less clear and flavonoids production is

obviously repressed. But the biosynthesis of two other types of phenylpropanoid-derived

phytoalexins appears to be favored during pear nonhost resistance: simple coumarin on one hand

and hydroxycinnamic acid amides on the other hand.

The comparative analysis between a host and a nonhost resistance in pear shows that, even though

specificities are observed, the two major defense lines engaged are shared: the cell wall and its

cuticle on one hand, the secondary metabolism with the phenylpropanoid pathway on the other

hand. Moreover, these defenses seem deployed largely independently of the SA signaling pathway,

widely recognized as the main defense hormone against biotrophic pathogens.

**Conclusion**

As far as we know, our work is the first one published regarding a transcriptomic analysis of post-

infections events of a nonhost resistance to *Venturia sp.* in apple and pear. Velho and Stadik [115]

recently published a detailed description of the apple / *Colletotrichum higginsianum* nonhost

resistance, highlighting the accumulation of callose at the sites of penetration of the fungus. But no

data on gene expression was included. Here, our molecular work on apple / *V. pyrina* nonhost

resistance remains preliminary and in order to allow a deeper deciphering, further analyses must be

considered with the aid of tools adapted to this type I nonhost resistance with very few cells engaged

in an HR-like reaction, only visible at a microscopic level. In pear, this deciphering allowed us to show

that nonhost resistance against *V. inaequalis* is a type II one, which involves enough pathogen

penetration in plant tissue to trigger visible HR and develops post-invasive defenses.

To summarize our findings on pear with a notion of cascading effect, we can propose the following

scenario (Fig. 4): once *V. inaequalis* presence is recognized by pear, a calcium cellular influx is

induced and leads to the development of a pre-invasive defense, the stomatal closure, but also

promotes an early post-invasive defense, an apoplastic ROS accumulation. Apoplastic ROS, acting

themselves as ubiquitous messengers, come to reinforce the stomatal closure but also mediate

cellular signaling resulting in two post-invasive defenses: HR development at infection sites, along

with phytoalexin (simple coumarin and hydroxicinnamic acid amines) production. The observed

alterations of the epidermis composition (cellulose, pectin, lignin for the cell wall, and cutin for the

cuticle), are presumed to strengthen this physical barrier and can be seen as the development of

another pre-invasive defense. The calcium (action on pectin reviewed in [116]) and the ROS (action

on lignin, [117]; [118]; action on cuticle, [119]) have been linked to some type of epidermis

modifications and may participate in the proceeding of these defense in pear / *V. inaequalis* nonhost

interaction.

Nonhost resistance is defined as the resistance of an entire plant species against a specific parasite or

pathogen [120] and is seen as the most durable resistance of plant. Thus, understanding the

molecular mechanisms underlying nonhost resistance can open up some interesting avenues to

create sustainable host resistances in the same plant species. Considering pear, in order to stop the

germination and entrance of hemibiotrophic host fungi such as *V. pyrina*, strengthening the cuticle

initial barrier via more cutin production and cross-link, or promoting the biosynthesis of phytoalexins

like hydroxycinnamic acid amines, appear as promising solutions, relatively easy to engineer

regarding recent advances in biotechnology tools on this species ([121]; [122]; [123]).

**Material and methods**

Biological material

Apple plants from the cultivar 'Gala' and pear plants from the cultivar 'Conference' were chosen

because of their susceptibility to *V. inaequalis* and *V. pyrina*, respectively. The apple and pear

genotypes were multiplied in vitro, rooted and acclimatized in greenhouse as described previously

([124]; [125]).

For apple scab inoculation, the *V. inaequalis* monoconidial isolate used was EU-B05 from the

European collection of *V. inaequalis* of the European project Durable Apple Resistance in Europe

[126]. For pear scab inoculation, the monoconidial strain VP102 of *V. pyrina* was chosen for its

aggressiveness on 'Conference' [127].

Scab inoculation procedure

Greenhouse growth conditions and mode of inoculum preparation were as described in Parisi and

Lespinasse [128] for apple and Chevalier et al [129] for pear. Briefly, the youngest leaf of actively

growing shoots was tagged and the plants inoculated with a conidial suspension ($2 \times 10^5$ conidia ml$^{-1}$)

of *Venturia pyrina* strain VP102 for apple and *Venturia inaequalis* strain EUB04 for pear. Symptoms

were recorded at 14, 21, 28, 35 and 42 days after inoculation on 20 actively growing shoots (one

shoot by plant) for each interaction. The type of symptoms was scored using the 6 class-scale of

Chevalier et al [130].

Microscopic observations

Histological studies were made on samples stained with the fluorophore solophenylflavine [131]. In

brief, leaf discs were rinsed in ethanol 50° before staining in a water solution of solophenylflavine

7GFE 500 (SIGMA-Aldrich, St Louis USA) 0.1% (v/v) for 10 min. The samples were first rinsed in

deionized water, then in 25% glycerol for 10 min. Finally, the leaf samples were mounted on glass-

slides in a few drops of 50% glycerol. They were examined with a wide-field epifluorescence

microscope BH2-RFC Olympus (Hamburg, D) equipped with the following filter combination:

excitation filter 395 nm and emission filter 504 nm.

666
Transcriptomics experiment

Leaf samples were immediately frozen in liquid nitrogen and kept at -80°C until analysis. Sampling

concerned the youngest expanded leaf of each plant labeled the day of the inoculation. Each

biological repeat is a pool of three leaves from three different plants and two biological repeats (n=2)

have been made by condition (genotype x treatment x time). Leaf samples taken just before

inoculation (T0) and at 24 and 72hpi, were then used to perform transcriptomics analyses.

For RNA extraction, frozen leaves were ground to a fine powder in a ball mill (MM301, Retsch, Hann,

Germany). RNA was extracted with the kit NucleoSpin RNA Plant (Macherey Nagel, Düren, Germany)

according to the manufacturer's instructions but with a modification: 4% of PVP40 (4 g for 100 ml)

were mixed with the initial lysis buffer RAP before use. Purity and concentration of the samples were

assayed with a Nanodrop spectrophotometer ND-1000 (ThermoFisher Scientific, Waltham, MA, USA)

and by visualization on agarose gel (1% (weight/volume) agarose, TAE 0.5x, 3% (volume/volume)

Midori green). Intron-spanning primers (forward primer: CTCTTGGTGTCAGGCAAATG, reverse primer:

TCAAGGTTGGTGGACCTCTC) designed on the EF-1α gene (accession AJ223969 for apple and

PCP017051 for pear, available at https://www.rosaceae.org/, with the datasets on "Pyrus communis

v1.0 draft genome") were used to check the absence of genomic DNA contamination by PCR. The

PCR reaction conditions were as follows: 95°C for 5 min, followed by 35 cycles at 95°C for 30 s, 60°C

for 45 s, 72°C for 1 min, with a final extension at 72°C for 5 min. The PCR products were separated on

a 2% agarose gel.

Amplifications (aRNAs) were produced with MessageAmpII aRNA Kit (Ambion Invitrogen, Waltham,

MA, USA), from 300 ng total RNA. Then 5 µg of each aRNA were retrotranscribed and labelled using a

SuperScript II reverse transcriptase (Transcriptase inverse SuperScript™ II kit, Invitrogen, Carlsbad,

CA, USA) and fluorescent dyes: either cyanine-3 (Cy3) or cyanine-5 (Cy5) (Interchim, Montluçon,

France). Labeled samples (30 pmol each, one with Cy3, the other with Cy5) were combined two by

two, depending on the experimental design (i. e. for example, the CF/EUB05/24hpi sample and the

CF/non inoculated sample are labeled with Cy3 and Cy5 respectively and pooled to co-hybridize the

microarray) . For each comparison two biological replicates were analyzed in dye-switch as described

in Depuydt et al [132]. Paired labeled samples were then co-hybridized to Agilent microarray AryANE

v2.0 (Agilent-070158_IRHS_AryANE-Venise, GPL26767 at GEO: https://www.ncbi.nlm.nih.gov/geo/)

for apple, or Pyrus v1.0 (Agilent-078635_IRHS_Pyrus, GPL26768 at GEO) for pear, containing

respectively 133584 (66792 sense and 66792 anti-sense probes) and 87812 (43906 sense and 43906

anti-sense probes) 60-mer oligonucleotide probes. The hybridizations were performed as described

in Celton, Gaillard et al [133] using a MS 200 microarray scanner (NimbleGen Roche, Madison, WI,

USA).

For microarray analysis we designed two new chips. For apple we used a deduplicated probeset from

the AryANE v1.0 ([133]; 118740 probes with 59370 in sense and 59370 in anti-sense) augmented by

14844 probes (7422 in sense and 7422 in anti-sense) designed on new gene annotations from *Malus*

*domestica*          GDDH13          v1.1          (https://iris.angers.inra.fr/gddh13          or

https://www.rosaceae.org/species/malus/malus_x_domestica/genome_GDDH13_v1.1).          These

probes target new coding genes with UTRs when available, manually curated micro-RNA precursors

and transposable elements. For transposable elements we used one consensus sequence for each

family and a randomly peaked number of elements proportional to their respective abundance in the

genome. The microarrays used in this study also have probes for coding genes of *V. inaequalis* but

they have not been taken into account.

For pear the design was done on the *Pyrus communis* Genome v1.0 Draft Assembly & Annotation

available on GDR (https://www.rosaceae.org/species/pyrus/pyrus_communis/genome_v1.0) web

site. We have downloaded the reference genome and gene predictions fasta files and structural

annotation gff file the 21st of September 2015. Using home-made Biopython scripts we have

extracted spliced CDS sequences with 60 nucleotides before start and after stop codons to get UTR-

like sequences likely to be found on transcripts resulting in a fasta file containing 44491 sequences.

These 60 nucleotides increase the probability of finding specific probes on genes with high similarity.

This file was sent to the eArray Agilent probe design tool (https://earray.chem.agilent.com/earray/)

to generate one probe per gene prediction. Options used were: Probe Length: 60, Probe per Target:

1, Probe Orientation: Sense, Design Options: Best Probe Methodology, Design with 3' Bias. The

probeset was then reverse-complemented to generate anti-sense probes and filtered to remove

duplicated probes. The final probeset contains 87812 unique probes targeting 1 (73612 probes) or

more (14200 probes) potential transcript both in sense and anti-sense.

Normalization and statistical analyses performed to get normalized intensity values have been done

as in Celton, Gaillard et al [133]. For each comparison and each probe, we retrieved a ratio of the

logarithms of the fluorescence intensities (one per compared sample: T0 versus 24hpi or T0 versus

72hpi in our case) and an associated p-value. The applied p-value threshold to determine DEGs

(differentially expressed genes) was 0.01. Through blast analysis, a TAIR accession number (The

Arabidopsis Information Resource; https://www.arabidopsis.org/; [134]) has been linked to a

majority of apple or pear "probe/corresponding gene" and the couple "TAIR accession/ratio value"

has then been used to make a global analysis of functional categories observed in the Mapman

software (https://mapman.gabipd.org/homemapman.gabipd.org; [135]). The detailed analysis of

DEGs has been done through TAIR and KEGG (https://www.genome.jp/kegg/) databases, and

bibliography. Metadata for the 172 (162 for pear and 10 for apple) DEGs discussed in this work are

available in Table S3 and S4 (Online only).

qPCR validation of transcriptomic data

In order to validate transcriptomic data, qPCR was performed on a selection of gene/sample

associations. First-strand cDNA was synthesized using total RNA (2.0 μg) in a volume of 30 μl of 5×

buffer, 0.5 μg of oligodT15 primer, 5 μl of dNTPs (2.5 mM each), and 150 units of MMLV RTase

(Promega, Madison, WI, USA). The mixture was incubated at 42°C for 75 min. qPCR was then

performed. Briefly, 2.5 μl of the appropriately diluted samples (1/16 dilution) were mixed with 5 μl of

PerfeCTa SYBR Green SuperMix for iQ kit (Quantabio, Beverly, MA, USA) and 0.2 or 0.6 μl of each

primer (10 μM) in a final volume of 10 μl. Primers were designed with Primer3Plus, their volumes

were according to their optimal concentration (determined for reaction efficiency near to 100%;

calculated as the slope of a standard dilution curve; [136]). Accessions, primer sequences and

optimal concentrations are indicated in Table S2. The reaction was performed on a CFX Connect Real-

Time System (BIO-RAD, Hercules, CA, USA) using the following program: 95°C, 5 min followed by 40

cycles comprising 95°C for 3 s, 60°C for 1 min. Melting curves were performed at the end of each run

to check the absence of primer-dimers and nonspecific amplification products. Expression levels

were calculated using the ΔΔCT method [137] and were corrected as recommended in

Vandesompele et al [138], with three internal reference genes (GADPH, TUA and ACTIN 7 for apple,

GADPH, TUA and EF1α for pear) used for the calculation of a normalization factor. For each couple

DEG/sample (sample defining a plant, time, treatment and biological repeat combination), the ratio

was obtained by dividing the mean value of CT calculated from 3 technical repeats by the

normalization factor obtained for this sample.

**Supplementary information**

**Additional File 1: Table S1, S2, S3** and **S4**.

## Abbreviations

ABA: abscisic acid; CDPK: calcium dependent protein kinase; CRK: cysteine-rich receptor-like kinase; DEG: differentially expressed gene; DFR: dihydroflavonol 4-reductase; DGDG: digalactosyldiacylglycerol; ET: ethylene; ER: endoplasmic reticulum; ETI: effector triggered immunity; FAE: fatty acid elongase; GPAT: glycerol-3-phosphate acyltransferase; hpi: hours post inoculation; HR: hypersensitive reaction; JA; jasmonic acid; LCB: long chain/sphingoid base component; LCB-Ps: long chain base phosphate component; LOX: lipoxygenase; MAMP: microbe-associated molecular pattern; OPDA: 12-oxo-phytodienoic acid; PCD: programed cell death; PTI: pattern triggered immunity; RBOH: respiratory burst oxidase homolog; ROS: reactive oxygen species; SA: salicylic acid; SAR: systemic acquired resistance

## Declarations

Acknowledgements

The authors gratefully acknowledge the IRHS-ImHorPhen team of INRA Angers for technical assistance in plant maintenance and the technical platforms ANAN and IMAC.

Authors contribution

EC, LP, and EV conceived the study. EC and EV supervised the study. ER and MB performed the biological experiments. SG and SP performed the database work and assisted with the bioinformatics analysis. EV wrote the original manuscript. EV and EC edited the manuscript. All authors have read and agreed to the published version of the manuscript.

Funding

This project was funded by the Synthé-Poir-Pom project (Angers University) and by the TIFON project (INRAE, department BAP).

Availability of data and materials

The datasets supporting the conclusion of this article are available in the Gene Expression Omnibus

(GEO) repository [https://www.ncbi.nlm.nih.gov/geo/] with GSE159179 and GSE159180 accession

numbers for apple and pear respectively.

Ethics approval and consent to participate

Experimental research on plants in this work comply with relevant institutional, national, and

international guidelines and legislation.

Consent for publication

This section is not applicable.

Competing interests

The authors declare that they have no competing interests

**References**

1. Potter D, Eriksson T, Evans RC, Oh S, Smedmark JEE, Morgan DR, Kerr M, Robertson KR, et al.

Phylogeny and classification of *Rosaceae*. Plant Syst Evol. 2007;266:5-43.

2. Campbell CS, Evans RC, Morgan DR, Dickinson TA, Arsenault MP. Phylogeny of subtribe *Pyrinae*

(formerly the *Maloideae*, *Rosaceae*): Limited resolution of a complex evolutionary history.

Plant Syst Evol. 2007;266: 119-145.

3. Wu J, Wang Z, Shi Z, Zhang S, Ming R, Zhu SL et al. The genome of pear (*Pyrus bretschneideri*

Redh.). Genome Res. 2013;23:396-408.

4. Velasco R, Zharkikh A, Affourtit J, Dhingra A, Cestaro A, Kalyanaraman A et al. The genome of the

domesticated apple (*Malus × domestica* Borkh.). Nature Genet. 2010;42:833–839.

5. Stehmann C, Pennycook S, Plummer K. Molecular identification of a sexual interloper: the pear

pathogen *Venturia pirina*, has sex on apple. Phytopathol. 2001;91:663-541.

6. Rossman A, Castlebury L, Aguirre-Hudson B, Berndt R, Edwards. (2647–2651) Proposals to

conserve the name *Venturia acerina* against *Cladosporium humile; Venturia borealis* against

*Torulama culicola; Venturia carpophila* against *Fusicladium amygdali* and *Cladosporium*

*americanum; Sphaerella inaequalis (Venturia inaequalis)* against *Spilocaea pomi, Fumago*

*mali, Actinone macrataegi, Cladosporium dendriticum, Asteroma mali,* and *Scolicotrichum*

*venosum;* and *Venturia pyrina* against *Helminthosporium pyrorum, Fusicladium virescens, F.*

*fuscescens, Cladosporium polymorphum* and *Passalora pomi (Ascomycota: Dothideomycetes).*

Taxon. 2018;67:1209-1211.

7. Gonzalez-Dominguez E, Armengol J, Rossi V. Biology and epidemiology of *Venturia* species

affecting fruit crops: a review. Front Plant Sci. 2017;8: 1496.

8. Jiang S, Park P, Ishii H. Penetration behaviour of *Venturia nashicola* , associated with hydrogen

peroxide generation, in Asian and European pear leaves. J Phytopathol. 2014;162: 770-778.

9. Chevalier M, Bernard C, Tellier M, Audrain C, Durel CE. Host and non-host interactions of *Venturia*

*inaequalis* and *Venturia pirina* on *Pyrus communis* and *Malus x domestica.* Acta Hortic.

2004;663: 205-208.

10. Lee HA, Lee HY, Seo E, Lee J, Kim SB, Oh S, Choi E, Choi E, Lee SE, Choi D. Current understanding

of plant nonhost resistance. Mol Plant-Microbe Interact. 2017a;30:5-5.

11. Perchepied L, Chevreau E, Ravon E, Gaillard S, Pelletier S, Bahut M, Berthelot P, Cournol R,

Schouten HJ, Vergne E. Successful intergeneric transfer of a major apple scab resistance gene

(Rvi6) from apple to pear and precise comparison of the downstream molecular mechanisms

of this resistance in both species. BMC Genomics. 2021;22:843.

12. Mysore JS, Ryu CM. Nonhost resistance: how much do we know? Trends Plant Sci. 2004;9: 97-

104.

13. Van Moerkercke A, Duncan O, Zander M, Šimura J, Broda M, Vanden Bossche R, Lewsey MG,

Lama S, Singh KB, Ljung K, Ecker JR, Goossens A, Millar AH, Van Aken O. A

MYC2/MYC3/MYC4-dependent transcription factor network regulates water spray-

responsive gene expression and jasmonate levels. Proc Natl Acad Sci U S A. 2019;116:23345-

23356.

14. Gusberti M, Gessler C, Broggini GA. RNA-Seq analysis reveals candidate genes for ontogenic

resistance in Malus-Venturia pathosystem. PLoS One. 2013;8:e78457.

15. Wasternack C, Feussner I. The oxylipin pathways: biochemistry and function. Annu Rev Plant Biol.

2018;69:363-386.

16. Li C, Schilmiller AL, Liu G, Lee GI, Jayanty S, Sageman C, et al. Role of beta-oxidation in jasmonate

biosynthesis and systemic wound signaling in tomato. Plant Cell. 2005;17:971-986.

17. Seo JS, Koo YJ, Jung C, Yeu SY, Song JT, Kim JK, Choi Y, Lee JS, Do Choi Y. Identification of a novel

jasmonate-responsive element in the AtJMT promoter and its binding protein for AtJMT

repression. PLoS One. 2013;8:e55482.

18. Jeong JS, Jung C, Seo JS, Kim JK, Chua NH. The deubiquitinating enzymes UBP12 and UBP13

positively regulate MYC2 levels in jasmonate responses. Plant Cell. 2017;29:1406-1424.

19. Birkenbihl RP, Diezel C, Somssich IE. Arabidopsis WRKY33 is a key transcriptional regulator of

hormonal and metabolic responses toward *Botrytis cinerea* infection. Plant Physiol.

2012;159:266-85.

20. Kaurilind E, Xu E, Brosché M. A genetic framework for H2O2 induced cell death in *Arabidopsis*

*thaliana.* BMC Genomics. 2015;16:837.

21. Nurmberg PL, Knox KA, Yun BW, Morris PC, Shafiei R, Hudson A, Loake GJ. The developmental

selector AS1 is an evolutionarily conserved regulator of the plant immune response. Proc

Natl Acad Sci U.S.A. 2007;104:18795-187800.

22. Ali S, Ganai BA, Kamili AN, Bhat AA, Mir ZA, Bhat JA et al. Pathogenesis-related proteins and

peptides as promising tools for engineering plants with multiple stress tolerance. Microb Res.

2018; 212-213:29-37.

23. Li J, Brader G, Palva ET. The WRKY70 transcription factor: a node of convergence for jasmonate-

mediated and salicylate-mediated signals in plant defense. Plant Cell. 2004;16:319–331.

24. Li J, Brader G, Kariola T, Palva ET. WRKY70 modulates the selection of signaling pathways in plant

defense. Plant J. 2006;46:477–491.

225. Wang D, Amornsiripanitch N, Dong X. A genomic approach to identify regulatory nodes in the

transcriptional network of systemic acquired resistance in plants. PLoS Pathog. 2006;2:e123.

26. Li J, Zhong R, Palva ET. WRKY70 and its homolog WRKY54 negatively modulate the cell wall-

associated defenses to necrotrophic pathogens in Arabidopsis. PLoS One. 2017;12:e0183731.

27. Genot B, Lang J, Berriri S, Garmier M, Gilard F, Pateyron S, et al. Constitutively active Arabidopsis

MAP kinase 3 triggers defense responses involving salicylic acid and SUMM2 resistance

protein. Plant Physiol. 2017;174:1238-1249.

28. Hu X, Reddy AS. Cloning and expression of a PR5-like protein from Arabidopsis: inhibition of

fungal growth by bacterially expressed protein. Plant Mol Biol. 1997;34:949-59.

29. Mestre P, Arista G, Piron MC, Rustenholz C, Ritzenthaler C, Merdinoglu D, Chich JF. Identification

of a *Vitis vinifera* endo-β-1,3-glucanase with antimicrobial activity against *Plasmopara*

*viticola*. Mol Plant Pathol. 2017;18:708-719.

30. Zhang SB, Zhang WJ, Zhai HC, Lv YY, Cai JP, Jia F, Wang JS, Hu YS. Expression of a wheat β-1,3-

glucanase in Pichia pastoris and its inhibitory effect on fungi commonly associated with

wheat kernel. Protein Expres Purif. 2019;154:134-139.

31. Santamaria M, Thomson CJ, Read ND, Loake GJ; The promoter of a basic PR1-like gene, AtPRB1,

from Arabidopsis establishes an organ-specific expression pattern and responsiveness to

ethylene and methyl jasmonate. Plant Mol Biol. 2001;47:641-52.

32. Truman W, Sreekanta S, Lu Y, Bethke G, Tsuda K, Katagiri F, Glazebrook J. The CALMODULIN-

BINDING PROTEIN60 family includes both negative and positive regulators of plant immunity.

Plant Physiol. 2013;163:1741-1751.

33. Boursiac Y, Lee SM, Romanowsky S, Blank R, Sladek C, Chung WS, Harper JF. Disruption of the

vacuolar calcium-ATPases in Arabidopsis results in the activation of a salicylic acid-dependent

programmed cell death pathway. Plant Physiol. 2010;154:1158-71.

34. Huang J, Sun Y, Orduna AR, Jetter R, Li X. The Mediator kinase module serves as a positive

regulator of salicylic acid accumulation and systemic acquired resistance. Plant J. 2019;98:842-

852.

35. Wang W, Tang W, Ma T, Niu D, Jin JB, Wang H, Lin R. A pair of light signaling factors FHY3 and

FAR1 regulates plant immunity by modulating chlorophyll biosynthesis. Journal of Integrative

Plant Biol. 2015;58:91-103.

36. Song JT, Koo YJ, Seo HS, Kim MC, Choi YD, Kim JH. Overexpression of AtSGT1, an Arabidopsis

salicylic acid glucosyltransferase, leads to increased susceptibility to *Pseudomonas syringae*.

Phytochemistry. 2008;69:1128-34.

37. Vlot AC, Dempsey DA, Klessig DF. Salicylic acid, a multifaceted hormone to combat disease. Annu

Rev Phytopathol. 2009;47:177-206.

38. Andreasson E, Jenkins T, Brodersen P, Thorgrimsen S, Petersen NH, Zhu S et al. The MAP kinase

substrate MKS1 is a regulator of plant defense responses. EMBO J. 2005;24:2579-89.

39. Xia Y, Yu K, Gao QM, Wilson EV, Navarre D, Kachroo P, Kachroo A. Acyl CoA binding proteins are

required for cuticle formation and plant responses to microbes. Front Plant Sci. 2012;3:224.

40. Boller T, Felix G. A renaissance of elicitors: perception of microbe-associated molecular patterns

and danger signals by pattern recognition receptors. Annu Rev Plant Biol. 2009;60:379-406.

41. Vincill ED, Bieck AM, Spalding EP. Ca(2+) conduction by an amino acid-gated ion channel related

to glutamate receptors. Plant Physiol. 2012;159:40-46.

42. Hou C, Tian W, Kleist T, He K, Garcia V, Bai F, Hao Y, Luan S, Li L. DUF221 proteins are a family of

osmosensitive calcium-permeable cation channels conserved across eukaryotes. Cell Res.

2014;24:632-635.

43. Fonseca JP, Mysore KS. Genes involved in nonhost disease resistance as a key to engineer durable

resistance in crops. Plant Sci. 2019;279:108-116.

44. Ronzier E, Corratgé-Faillie C, Sanchez F, Prado K, Brière C, Leonhardt N, Thibaud JB, Xiong TC.

CPK13, a noncanonical Ca2+-dependent protein kinase, specifically inhibits KAT2 and KAT1

shaker K+ channels and reduces stomatal opening. Plant Physiol. 2014;166:314-26.

45. Zou JJ, Li XD, Ratnasekera D, Wang C, Liu WX, Song LF, Zhang WZ, Wu WH. Arabidopsis CALCIUM-

DEPENDENT PROTEIN KINASE8 and CATALASE3 function in abscisic acid-mediated signaling

and H2O2 homeostasis in stomatal guard cells under drought stress. Plant Cell.

2015;27:1445-1460.

46. Förster S, Schmidt LK, Kopic E, Anschütz U, Huang S, Schlücking K, et al. Wounding-induced

stomatal closure requires jasmonate-mediated activation of GORK K(+) channels by a Ca(2+)

sensor-kinase CBL1-CIPK5 complex. Dev Cell. 2019;48:87-99.e6.

47. Kadota Y, Shirasu K, Zipfel C. Regulation of the NADPH oxidase RBOHD during plant immunity.

Plant Cell Physiol. 2015;56: 1472–1480.

48. Adachi H, Yoshioka H. Kinase-mediated orchestration of NADPH oxidase in plant immunity. Brief

Funct Genomics. 2015;14:253-259.

49. Zhai L, Sun C, Feng Y, Li D, Chai X, Wang L, Sun Q, et al. AtROP6 is involved in reactive oxygen

species signaling in response to iron-deficiency stress in Arabidopsis thaliana. FEBS Lett.

2018;592:3446-3459.

50. Gao X, Chen X, Lin W, Chen S, Lu D, Niu Y, Li L, Cheng C, McCormack M, Sheen J, Shan L, He P.

Bifurcation of Arabidopsis NLR immune signaling via Ca²⁺-dependent protein kinases. PLoS

Pathog. 2013;9:e1003127.

51. Qi J, Wang J, Gong Z, Zhou JM. Apoplastic ROS signaling in plant immunity. Curr Opin Plant Biol.

2017;38:92-100.

52. Bourdais G, Burdiak P, Gauthier A, Nitsch L, Salojärvi J, Rayapuram C et al. CRK consortium large-

scale phenomics identifies primary and fine-tuning roles for CRKs in responses related to

oxidative stress. PLoS Genet. 2015;11:e1005373.

53. Sun T, Nitta Y, Zhang Q, Wu D, Tian H, Lee JS, Zhang Y. Antagonistic interactions between two

MAP  kinase  cascades  in  plant  development  and  immune  signaling.  EMBO  Rep.

2018;19:e45324.

54. O'Brien JA, Daudi A, Butt VS, Bolwell GP. Reactive oxygen species and their role in plant defence

and cell wall metabolism. Planta. 2012;236:765-779.

55. Brosché M, Blomster T, Salojärvi J, Cui F, Sipari N, Leppälä J et al. Transcriptomics and functional

genomics of ROS-induced cell death regulation by RADICAL-INDUCED CELL DEATH1. PLoS

Genet. 2014;10:e1004112.

56. Cheng YT, Li Y, Huang S, Huang Y, Dong X, Zhang Y, Li X. Stability of plant immune-receptor

resistance proteins is controlled by SKP1-CULLIN1-F-box (SCF)-mediated protein degradation.

Proc Natl Acad Sci U.S.A. 2011;108:14694-14699.

57. Coll NS, Vercammen D, Smidler A, Clover C, Van Breusegem F, Dangl JL, Epple P. Arabidopsis type

I metacaspases control cell death. Science. 2010;330:1393-1397.

58. Lema-Asqui S, Vercammen D, Serrano I, Valls M, Rivas S, Van Breusegem F, et al. AtSERPIN1 is an

inhibitor  of  the  metacaspase  AtMC1-mediated  cell  death  and  autocatalytic  processing  in

planta. New Phytol. 2018;218:1156-1166.

59. Kawasaki T, Nam J, Boyes DC, Holt BF, Hubert DA, Wiig A, Dangl JL. A duplicated pair of

Arabidopsis  RING-finger  E3  ligases  contribute  to  the  RPM1-and  RPS2-mediated

hypersensitive response. Plant J. 2005;44: 258-270.

60. Luo H, Laluk K, Lai Z, Veronese P, Song F, Mengiste T. The Arabidopsis Botrytis Susceptible1

interactor defines a subclass of RING E3 ligases that regulate pathogen and stress responses.

Plant Physiol. 2010;154:1766-1782.

61. Huby E, Napier JA, Baillieul F, Michaelson LV, Dhondt-Cordelier S. Sphingolipids: towards an

integrated view of metabolism during the plant stress response. New Phytol. 2020;225:659-

670.

62. Saucedo-Garcia M, Guevara-Garcia A, Gonzalez-Solis A, Cruz-Garcia F, VazquezSantana S,

Markham JE, et al. MPK6, sphinganine and the LCB2a gene from serine palmitoyltransferase

are required in the signaling pathway that mediates cell death induced by long chain bases in

Arabidopsis. New Phytol. 2011;191:943–957.

63. Veley KM, Maksaev G, Frick EM, January E, Kloepper SC, Haswell ES. Arabidopsis MSL10 has a

regulated cell death signaling activity that is separable from its mechanosensitive ion channel

activity. Plant Cell. 2014;26:3115-3131.

64. Guerringue Y, Thomine S, Frachisse JM. Sensing and transducing forces in plants with MSL10 and

DEK1 mechanosensors. FEBS Lett. 2018;592:1968-1979.

65. Zhao C, Nie H, Shen Q, Zhang S, Lukowitz W, Tang D. EDR1 physically interacts with MKK4/MKK5

and negatively regulates a MAP kinase cascade to modulate plant innate immunity. PLoS

Genet. 2014;10:e1004389.

66. Lee MH, Jeon HS, Kim HG, Park OK. An Arabidopsis NAC transcription factor NAC4 promotes

pathogen-induced cell death under negative regulation by microRNA164. New Phytol.

2017b;214:343-360.

67. Najafi J, Brembu T, Vie AK, Viste R, Winge P, Somssich IE, Bones AM. PAMP-INDUCED SECRETED

PEPTIDE 3 modulates immunity in Arabidopsis. J Exp Bot. 2020;71:850-864.

68. Khafif M, Balagué C, Huard-Chauveau C, Roby D. An essential role for the VASt domain of the

Arabidopsis VAD1 protein in the regulation of defense and cell death in response to

pathogens. PLoS One. 2017;12:e0179782.

69. Mittler R, Herr EH, Orvar BL, van Camp W, Willekens H, Inzé D, Ellis BE. Transgenic tobacco plants

with reduced capability to detoxify reactive oxygen intermediates are hyperresponsive to

pathogen infection. Proc Natl Acad Sci U.S.A. 1999;96:14165-70.

70. Simon C, Langlois-Meurinne M, Didierlaurent L, Chaouch S, Bellvert F, Massoud K, et al. The

secondary metabolism glycosyltransferases UGT73B3 and UGT73B5 are components of redox

status in resistance of Arabidopsis to *Pseudomonas syringae* pv. *tomato*. Plant Cell Environ.

2014;37:1114-1129.

71. Bacete L, Mélida H, Miedes E, Molina A. Plant cell wall-mediated immunity: cell wall changes

trigger disease resistance responses. Plant J. 2018;93:614-636.

72. Miedes E, Vanholme R, Boerjan W, Molina A. The role of the secondary cell wall in plant

resistance to pathogens. Front Plant Sci. 2014;5:358.

73. Fich EA, Segerson NA, Rose JK. The plant polyester cutin: biosynthesis, structure, and biological

roles. Ann Rev Plant Biol. 2016;67:207-233.

74. Lim GH, Singhal R, Kachroo A, Kachroo P. Fatty acid- and lipid-mediated signaling in plant defense.

Annu Rev Phytopathol. 2017;55:505-536.

75. Joubès J, Raffaele S, Bourdenx B, Garcia C, Laroche-Traineau J, Moreau P, Domergue F, Lessire R.

The VLCFA elongase gene family in *Arabidopsis thaliana*: phylogenetic analysis, 3D modelling

and expression profiling. Plant Mol Biol. 2008;67:547-66.

76. Lee SB and Suh MC. Advances in the understanding of cuticular waxes in *Arabidopsis thaliana* and

crop species. Plant Cell Rep. 2015;34:557-572.

77. Li Y, Beisson F, Koo AJ, Molina I, Pollard M, Ohlrogge J. Identification of acyltransferases required

for cutin biosynthesis and production of cutin with suberin-like monomers. Proc Natl Acad Sci

U.S.A. 2007;104:18339-18344.

78. Yeats TH, Huang W, Chatterjee S, Viart HM, Clausen MH, Stark RE, Rose JK. Tomato Cutin

Deficient 1 (CD1) and putative orthologs comprise an ancient family of cutin synthase-like

(CUS) proteins that are conserved among land plants. Plant J. 2014;77:667-675.

79. Cui F, Brosche M, Lehtonen MT, Amiryousefi A, Xu E, Punkkinen M, Valkonen JPT, Fujii H,

Overmyer K. Dissecting abscisic acid signaling pathways involved in cuticle formation. Mol

Plant. 2016;9: 926-938.

80. Yang L, Wen KS, Ruan X, Zhao YX, Wei F, Wang Q. Response of plant secondary metabolites to

environmental factors. Molecules. 2018;23i:e762.

81. Tetali S. Terpenes and isoprenoids: a wealth of compounds for global use. Planta. 2019;249:1-8.

82. Wang K, Senthil-Kumar M, Ryu CM, Kang L, Mysore KS. Phytosterols play a key role in plant innate
immunity against bacterial pathogens by regulating nutrient efflux into the apoplast. Plant
Physiol. 2012;158:1789-1802.

83. Falcone Ferreyra ML, Emiliani J, Rodriguez EJ, Campos-Bermudez VA, Grotewold E, Casati P. The
identification of maize and Arabidopsis type I FLAVONE SYNTHASEs links flavones with
hormones and biotic interactions. Plant Physiol. 2015;169:1090-107.

84. Wang N, Xu H, Jiang S, Zhang Z, Lu N, Qiu H, Qu C, Wang Y, Wu S, Chen X. MYB12 and MYB22 play
essential roles in proanthocyanidin and flavonol synthesis in red-fleshed apple (*Malus*
*sieversii* f. *niedzwetzkyana*). Plant J. 2017a;90:276-292.

85. Zhai R, Zhao Y, Wu M, Yang J, Li X, Liu H, Wu T, Liang F, Yang C, Wang Z, Ma F, Xu L. The MYB
transcription factor PbMYB12b positively regulates flavonol biosynthesis in pear fruit. BMC
Plant Biol. 2019;19:85.

86. Wang XC, Wu J, Guan ML, Zhao CH, Geng P, Zhao Q. Arabidopsis MYB4 plays dual roles in
flavonoid biosynthesis. Plant J. 2020;101:637-652.

87. Fernández-Pérez F, Vivar T, Pomar F, Pedreño MA, Novo-Uzal E. Peroxidase 4 is involved in
syringyl lignin formation in *Arabidopsis thaliana*. J Plant Physiol. 2015a;175:86-94.

88. Zhao Q, Nakashima J, Chen F, Yin Y, Fu C, Yun J, Shao H, Wang X, Wang ZY, Dixon RA. Laccase is
necessary and nonredundant with peroxidase for lignin polymerization during vascular
development in Arabidopsis. Plant Cell. 2013;25:3976-3987.

89. Cosio C, Ranocha P, Francoz E, Burlat V, Zheng Y, Perry SE, Ripoll JJ, Yanofsky M, Dunand C. The
class III peroxidase PRX17 is a direct target of the MADS-box transcription factor AGAMOUS-
LIKE15 (AGL15) and participates in lignified tissue formation. New Phytol. 2017;213:250-263.

90. Tokunaga N, Kaneta T, Sato S, Sato Y. Analysis of expression profiles of three peroxidase genes
associated with lignification in Arabidopsis thaliana. Physiol Plantarum. 2009;136:237-249.

91. Fernández-Pérez F, Pomar F, Pedreño MA, Novo-Uzal E. The suppression of AtPrx52 affects fibers

but not xylem lignification in Arabidopsis by altering the proportion of syringyl units. Physiol

Plantarum. 2015b;154:395-406.

92. Vilanova L, Teixidó N, Torres R, Usall J, Viñas I. The infection capacity of *P. expansum* and *P.*

*digitatum* on apples and histochemical analysis of host response. International Journal of

Food Microbiol. 2012;157:360-367.

93. Fink W, Haug M, Deising H, Mendgen K. Early defence responses of cowpea *(Vigna sinensis* L.)

induced by non-pathogenic rust fungi. Planta. 1991;185:246-254.

94. Song PP, Zhao J, Liu ZL, Duan YB, Hou YP, Zhao CQ, Wu M, Wei M, Wang NH, Lv Y, Han ZJ.

Evaluation of antifungal activities and structure-activity relationships of coumarin derivatives.

Pest Manag. Sci. 2017;73:94-101.

95. Giesemann A, Biehl B, Lieberei R. Identification of scopoletin as a phytoalexin of the rubber tree

*Hevea brasiliensis*. J Phytopathol. 1986;117:373–376.

96. Muroi A, Ishihara A, Tanaka C, Ishizuka A, Takabayashi J, Miyoshi H, Nishioka T. Accumulation of

hydroxycinnamic acid amides induced by pathogen infection and identification of agmatine

coumaroyltransferase in *Arabidopsis thaliana*. Planta. 2009;230:517-527.

97. Dobritzsch M, Lübken T, Eschen-Lippold L, Gorzolka K, Blum E, Matern A et al. MATE transporter-

dependent export of hydroxycinnamic acid amides. Plant Cell. 2016;28:583-96.

98. Celoy RM, VanEtten HD. (+)-Pisatin biosynthesis: from (-) enantiomeric intermediates via an

achiral 7,2'-dihydroxy-4',5'-methylenedioxyisoflav-3-ene. Phytochem. 2014;98:120-127.

99. Vorwieger A, Gryczka C, Czihal A, Douchkov D, Tiedemann J, Mock HP, et al. Iron assimilation and

transcription factor controlled synthesis of riboflavin in plants. Planta. 2007;226:147-158.

100. Trapalis M, Li SF, Parish RW. The Arabidopsis GASA10 gene encodes a cell wall protein strongly

expressed in developing anthers and seeds. Plant Sci. 2017;260:71-79.

101. Yamauchi Y, Hasegawa A, Mizutani M, Sugimoto Y. Chloroplastic NADPH-dependent alkenal/one

oxidoreductase contributes to the detoxification of reactive carbonyls produced under

oxidative stress. FEBS Lett. 2012;586:1208-1213.

102. Segond D, Dellagi A, Lanquar V, Rigault M, Patrit O, Thomine S, Expert D. NRAMP genes function

in *Arabidopsis thaliana* resistance to *Erwinia chrysanthemi* infection. Plant J. 2009;58:195-

207.

103. Lee MW, Jelenska J, Greenberg JT. Arabidopsis proteins important for modulating defense

responses to *Pseudomonas syringae* that secrete HopW1-1. Plant J. 2008;54:452-445.

104. Hong GJ, Xue XY, Mao YB, Wang LJ, Chen XY. Arabidopsis MYC2 interacts with DELLA proteins in

regulating sesquiterpene synthase gene expression. Plant Cell. 2012;24:2635-2648.

105. Rodriguez-Saona CR, Polashock J, Malo EA. Jasmonate-mediated induced volatiles in the

American cranberry, *Vaccinium macrocarpon*: from gene expression to organismal

interactions. Front Plant Sci. 2013;4:115.

106. Simanshu DK, Zhai X, Munch D, Hofius D, Markham JE, Bielawski J, et al. Arabidopsis accelerated

cell death 11, ACD11, is a ceramide-1-phosphate transfer protein and intermediary regulator

of phytoceramide levels. Cell Rep. 2014;6:388-399.

107. Fink L, Kwapiszewska G, Wilhelm J, Bohle RM. Laser-microdissection for cell type- and

compartment-specific analyses on genomic and proteomic level. Exp Toxicol Pathol. 2006;57

Suppl 2:25-259.

108. Yan Y, Zheng XF, Apaliya MT, Yang HJ, Zhang HY. Transcriptome characterization and expression

profile of defense-related genes in pear induced by *Meyerozyma guilliermondii*. Postharvest

Biol Technol. 2018;141:63-70.

109. Zhang Q, Zhao L, Li B, Gu X, Zhang X, Boateng NS, Zhang H. Molecular dissection of defense

response of pears induced by the biocontrol yeast, *Wicherhamomyces anomalus* using

transcriptomics and proteomics approaches. Biol Control. 2020;148:104305.

110. Wang H, lin J, Chang YH, Jiang CZ. Comparative transcriptomic analysis reveals that

ethylene/H2O2-mediated hypersensitive response and programmed cell death determine

the compatible interaction of sans pear and *Alternaria alternate.* Front Plant Sci.

2017b;8:196.

111. Xu M, Yang Q, Boateng NAS, Ahima J, Dou Y, Zhang H. Ultrastructure observation and

transcriptome analysis of *Penicillium expansum* invasion in postharvest pears. Postharvest

Biol Technol. 2020;165:111198.

112. Gill US, Lee S, Mysore KS. Host versus nonhost resistance: distinct wars with similar arsenals.

Phytopathol. 2015;105:580-587.

113. Pieterse CM, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SC. Hormonal modulation

of plant immunity. Annu Rev Cell Dev Biol. 2012;28:489-521.

114. Tsuda K, Sato M, Stoddard T, Glazebrook J, Katagiri F. Network properties of robust immunity in

plants. PLoS Genet. 2009;5(12):e1000772.

115. Velho AC, Stadnik MJ. Non-host resistance of arabidopsis and apple is associated with callose

accumulation and changes in preinfective structures of *Colletotrichum* species. Physiol Mol

Plant Pathol. 2020;110:101463.

116. Thor K. Calcium-nutrient and messenger. Front Plant Sci. 2019 ;10:440.

117. Das K and Roychoudhury A. Reactive oxygen species (ROS) and response of antioxidants as ROS-

scavengers during environmental stress in plants. Front Envir Sci. 2014;2:53.

118. Lehmann S, Serrano M, L'Haridon F, Tjamos SE, Metraux JP. Reactive oxygen species and plant

resistance to fungal pathogens. Phytochemistry. 2015;112:54-62.

119. Survila M, Davidsson PR, Pennanen V, Kariola T, Broberg M, Sipari N, Heino P, Palva ET.

Peroxidase-generated apoplastic ROS impair cuticle integrity and contribute to DAMP-elicited

defenses. Front Plant Sci. 2016; 7:1945.

120. Heath MC. Nonhost resistance and nonspecific plant defenses. Curr Opin Plant Biol. 2000;3:315-

319.

121. Charrier A, Vergne E, Dousset N, Richer A, Petiteau A, Chevreau E. Efficient targeted

mutagenesis in apple and first time edition of pear using the CRISPR/Cas9 system. Front Plant

Sci. 2019;10:40.

122. Chevreau E, Evans K, Montanari S Chagné D. Ch.19.9 Pyrus spp. Pear and Cydonia spp. Quince.

In: Litz RE, Pliego-Alfaro F, Hormaza JL, editors. Biotechnology of Fruit and Nut Crops, 2nd

edition. C.A.B. International, Wallingford, UK; 2020;p. 581-605.

123. Malabarba J, Chevreau E, Dousset N, Veillet F, Moizan J, Vergne E. New strategies to overcome

present CRISPR/Cas9 limitations in apple and pear: efficient dechimerization and base

editing. Int J Mol Sci. 2020;22:319.

124. Faize M, Malnoy M, Dupuis F, Chevalier M, Parisi L, Chevreau E. Chitinases of *Trichoderma*

*atroviridae* induce scab resistance and some metabolic changes in two cultivars of apple.

Phytopathol. 2003;93:1496-1504.

125. Leblay C, Chevreau E, Raboin LM. Adventitious shoot regeneration from in vitro leaves of several

pear cultivars (*Pyrus communis* L.). Plant Cell Tissue Organ Cult. 1991;25:99-105.

126. Lespinasse Y, Durel CE, Parisi L, Laurens F, Chevalier M, Pinet C. A European project: D.A.R.E.

Durable apple resistance in Europe. Acta Hortic. 2000;538:197–200.

127. Chevalier M, Tellier M, Lespinasse Y, Bruynincks M, Georgeault S. Behaviour studies of new

races of *Venturia pirina* isolated from 'Conference' cultivar on a range of pear cultivars. Acta

Hortic. 2008a;800:817-824.

128. Parisi L, Lespinasse Y. Pathogenicity of *Venturia inaequalis* strains of race 6 on apple clone

(*Malus sp.*). Plant Dis. 1996;80: 1179-1183.

129. Chevalier M, Tellier M, Lespinasse Y, Caffier V. How to optimize the *Venturia* pirina inoculation

on pear leaves in greenhouse conditions? Acta Hortic. 2008b;800: 913-920.

130. Chevalier M, Lespinasse Y, Renaudin S. A microscopic study of different classes of symptoms

coded by the *Vf* gene in apple resistance to scab (*Venturia inaequalis*). Plant Pathol. 1991;40:

249–256.

131. Hoch HC, Galvani CD, Szarowski DH, Turner JN. Two new fluorescent dyes applicable for

visualization of fungal cell walls. Mycologia. 2005;97: 580-588.

132. Depuydt S, Trenkamp S, Fernie AR, Elftieh S, Renou J-P, Vuylsteke M, Holster M, Vereecke D. An

integrated genomic approach to define niche establishment by *Rhodococcus fascians* . Plant

Physiol. 2009;149: 1366–1386.

133. Celton JM, Gaillard S, Bruneau M, Pelletier S, Aubourg S, Martin-Magniette ML, Navarro L,

Laurens F, Renou JP. Widespread anti-sense transcription in apple is correlated with siRNA

production and indicates a large potential for transcriptional and/or post-transcriptional

control. New Phytol. 2014;203:287-99.

134. Berardini TZ, Reiser L, Li D, Mezheritsky Y, Muller R, Strait E, Huala E. The Arabidopsis

information resource: Making and mining the "gold standard" annotated reference plant

genome. Genesis. 2015;53:474-85.

135. Thimm O, Bläsing O, Gibon Y, Nagel A, Meyer S, Krüger P, Selbig J, Müller LA, Rhee SY, Stitt M.

MAPMAN: a user-driven tool to display genomics data sets onto diagrams of metabolic

pathways and other biological processes. Plant J. 2004;37:914-939.

136. Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. Nucleic

Acids Res. 2001;29, e45.

137. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative

PCR and the 2−ΔΔCT method. Methods. 2001;25:402-408.

138. Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. et al. Accurate

normalization of real-time quantitative RT-PCR data by geometric averaging of multiple

internal control genes. Genome Biol. 2002;3(7):00341-003411.

1159

1160    **Figure legends**

1161

1162    **Fig. 1**: **Macro- and microscopic observations of nonhost interactions.**

1163 Binocular observation 21 days after *V. inaequalis* inoculation on 'Conference' (A) and (B) and *V.*

1164 *pyrina* inoculation on 'Gala' (C). Wide field fluorescence observations of: 'Conference' 3 days (D) and

1165 14 days (E) after *V. inaequalis* inoculation, 'Gala' 3 days (F) and 14 days (G) after *V. pyrina*

1166 inoculation. Ap: appressorium, C: conidia, Gf: germination filament, Pp: pin point

1167

1168 **Fig. 2**: **Functional categories of DEGs at 24 or 72hpi during pear response to *V. inaequalis*.**

1169 The number of up- or down-regulated DEGs is expressed as a percentage of the total number of

1170 genes present in the Pyrus v1.0 (87812 probes) microarray. DEGs are classified in functional

1171 categories according to MapMan 3.5.1R2 bins. Only bins with ≥ 6 DEGs are presented.

1172

**Fig. 3: DEGs involved in hormonal pathways during pear/*V. inaequalis* non-host interaction.**

A: DEGs involved in JA pathway; B: DEGs involved in SA pathway. Genes written in red are induced,

genes written in blue are repressed. ACA11: autoinhibited Ca2$^+$-ATPase, calmodulin-activated Ca2$^+$

pumps at the plasma membrane, endoplasmic reticulum, and vacuole. ACBP6: acyl-CoA-binding

protein. ACX4: acyl-CoA-oxidase1. AS1/MYB91: Asymmetric leaves 1 transcription factor, CAMTA1:

calmodulin-binding transcription activator, CBP60a: calmodulin-binding protein 60a, EDS1: enhanced

disease suceptibility 1. FAR1: FAR-red impaired response 1. G-box: cis-element in the promoter. JAZ:

jasmonate-zim domain protein, JMT: jasmonic acid carboxyl methyltransferase. LOX: lipoxygenase,

MES1: methylesterase 1. MFP2: multifunctional protein 2. MKS1: MAP kinase substrate 1. MYC2:

transcription factor. NINJA: novel interactor of JAZ. PAD4: phytoalexin deficient 4. UGT74F1:

glucosyltransferase. PR1-like (with ATPRB1), PR2, PR3, PR4 (HEL and ATEP3), PR5, PR12:

pathogenesis-related proteins. ST2A: sulfotransferase 2A. TPL: TOPLESS co-repressor. UBP12:

ubiquitin-specific protease 12. WRKY: transcription factor.

1186

**Fig. 4**: **Scenario of major events observed at three first days of pear/*V. inaequalis* non-host**

**interaction.**

On the left side, events observed in a typical cell, on the right side, events observed in guard cells of a

stomata. A: apoplasm, AP: appressorium, C: cuticle, CBL1: calcineurin B-like protein 1, CDPK: $Ca_2^+$-

dependent protein kinases, CRK: cysteine-rich receptor-like kinase, CY: cytoplasm, CW: cell wall, HAA:

hydroxycinnamic acid amines, HR: hypersensitive response, JA: jasmonic acid, MB: plasma

membrane, LCB: Long Chain/sphingoid Base components, MPK6: Mitogen activated protein kinase 6,

MSL10: mechano-sensitive like 10, N: nucleus, PH: penetration hypha, PR: pathogenesis related

proteins, RBOHB: respiratory burst oxidase homolog B, ROS: reactive oxygen species, S: stomata, SA:

salycilic acid, SC: simple coumarins, SP: spore.

1197

**Fig. 5: Main DEGs involved in cutin and wax biosynthesis during pear/V. inaequalis non-host**

**interaction.**

In green the chloroplast, in brown the endoplastic reticulum (ER) and in yellow the nucleus. Genes

written in red are induced, genes written in blue are repressed. FAS: Fatty Acid Synthase complex to

which belong ACCD (carboxytransferase beta subunit of the Acetyl-CoA carboxylase complex), FabG

(β-ketoacyl ACP-reductase) and MOD1 (enoyl-ACP-reductase) functions. FAE: fatty acid elongase

complex. KCS4 (3-ketoacyl-CoA synthase 4) and ECR/CER10 (trans-2-enoyl-CoA reductase) belong to

the FAE complex. CER1 (octadecanal decarbonylase) and CER3 are implicated in aldehydes (CER1)

and alkanes (CER1 and 3) generation in waxes biosynthesis. In cutin monomers synthesis, the ω-

hydroxylation of C16:0 and C18:1 is catalyzed by cytochrome P450 monooxygenase (CYP86A) and

LACS-encoded acyl-CoA synthetase may be required either to synthesize 16-hydroxy 16:0-CoA, a

substrate for ω-hydroxylase, or for membrane transfer of monomers. Finally, the mature

monoacylglycerol cutin monomers are generated by transfer of the acyl group from acyl-CoA to

glycerol-3-phosphate by glycerol-3-phosphate acyltransferase (GPAT) enzymes such as GPAT8. Some

GDSL-lipases enzyme (such as At1g28600, At1g28660, At1g54790, At3g16370, At3g48460, AtCUS4:

At4g28780, At5g14450) are then functioning as cutin synthase and polymerize cutin

monoacylglycerols. Transcription factors such as MYB16 and SHN1 are positive regulators of wax and

cutin biosynthesis. Adapted from Xia et al, 2009, [73] and [74].

1216

**Fig. 6: Main DEGs involved in the phenylpropanoid pathway during Pear / *V. inaequalis* non-host**

**interaction.**

Genes framed in red are induced, genes frames in blue are repressed. Framed in black, the detail of

genes involved in flavonoids production and found in this interaction. Abbreviations: 4CL, 4-

coumarate-CoA ligase; AACT, anthocyanin 5-aromatic acyltransferase; ANR, anthocyanidin reductase;

ANS, anthocyanin synthase; BGLC or BGLU, β-glucosidases; C3H, coumarate 3-hydroxylase; C4H,

cinnamate 4-hydroxylase; CAD, cinnamyl alcohol dehydrogenase; CCoAOMT, caffeoyl-CoA O-

methyltransferase; CCR, cinnamoyl-CoA reductase; CHI, chalcone isomerase; CHS, chalcone synthase;

COMT, caffeic acid 3-O-methyltransferase; CPK, calcium-dependent protein kinase ; DFR,

dihydroflavonol reductase; DMR6, downy mildiou resistant 6; F3H, flavanone 3-hydroxylase; F3'H

flavonoid 3'-hydroxylase; FLS, flavonol synthase; FNS, flavone synthase; GGT1, gamma-glutamyl

transpeptidase 1; GT, glucosyl transferase; HCT, hydroxycinnamoyl-CoA shikimate/quinate

hydroxycinnamoyl transferase; LAC, laccase; LAR, leucoanthocyanidin reductase; OMT1, O-

methyltransferase 1; PAL, phenylalanine ammonia-lyase; PER or PRX, peroxidase; TT7, transparent

testa 7; UGFT, UDP-glucose flavonoid-3-O-glucosyltransferase; UGT71D1, UDP-glycosyltransferase

71D1.

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

**Lignin biosynthesis**

LAC11, LAC17    PER17, PER47, PRX52

Lignin ← ← Monolignols

CAD

COMT

F5H

CCR    +at2g23910

Chorismate

CCoAOMT

**Shikimate pathway**

**Proanthocyanidins**

C3H HCT

MYB4
MYB12

**Flavan-3-ols**

ANR    LAR

ANS    DFR    F3H    CHI    CHS

4CL    C4H    PAL

F3'H TT7
GT UGT71D1

FNS DMR6

p-Coumaryol CoA

Phenylalanine

**Anthocyanin**

Quercetin,
Quercetin-glycoside

flavones

AACT1/ACT1

Coumarine

BGLU16, 42, 47, BGLC3

**Flavonoid biosynthesis**

**Hydroxycinnamic acids**

Hydoxycinnamic acid amines

Fig. 6