**A co-opted ISG15-USP18 binding mechanism normally reserved for deISGylation controls**

**type I IFN signalling**

Andri Vasou[1], Katie Nightingale[2], Vladimíra Cetkovská [1], Connor G.G. Bamford[3‡], Jelena

Andrejeva[1], Richard E. Randall[1], John McLauchlan[3], Michael P. Weekes[2], David J Hughes[1]*

[1]Biomedical Sciences Research Complex, School of Biology, University of St Andrews, St

Andrews, United Kingdom. [2]Cambridge Institute for Medical Research, Hills Road, Cambridge,

United Kingdom [3]MRC-University of Glasgow Centre for Virus Research, Glasgow, United

Kingdom.

‡ Present addresses: Wellcome-Wolfson Institute for Experimental Medicine, Queen's

University, Belfast UK.

* Corresponding author: djh25@st-andrews.ac.uk; Tel: +44 1334 467197

**Abstract**

Type I interferon (IFN) signalling induces the expression of several hundred IFN-stimulated

genes that provide an unfavourable environment for viral replication. To prevent an

overexuberant response and autoinflammatory disease, IFN signalling requires tight control.

One critical regulator is the ubiquitin-like protein ISG15, evidenced by autoinflammatory

disease in patients with inherited ISG15 deficiencies. Current models suggest that ISG15

stabilises USP18, a well-established negative regulator of IFN signalling. USP18 also functions

as an ISG15-specific peptidase, however its catalytic activity is dispensable for controlling IFN

signalling. Here, we show that the ISG15-dependent stabilisation of USP18 is necessary but

not sufficient for regulation of IFN signalling and that USP18 requires non-covalent

interactions with ISG15 to enhance its regulatory function. Intriguingly, this trait has been

acquired through co-option of a binding mechanism normally reserved for deISGylation,

identifying an unexpected new function for ISG15.

**Introduction**

The interferon (IFN) response plays a critical role in orchestrating protective antiviral immune responses to combat viral infections (1). Type I IFNs are widely expressed by all nucleated cells following viral infection (2) and signal through the IFN-α/β receptor (IFNAR), which consists of subunits IFNAR1 and IFNAR2 (3). Engagement of IFNAR triggers a phosphorylation cascade involving the reciprocal trans-phosphorylation of Janus kinase 1 (JAK1) and tyrosine kinase 2 (Tyk2) (4) and phosphorylation of the cytoplasmic tails of the receptor subunits creating a docking site for the recruitment and subsequent phosphorylation of signal transducer and activator of transcription 1 (STAT1) and STAT2 (5). Once activated, STAT1/2 heterodimers associate with IFN-regulatory factor 9 (IRF9) to form IFN-stimulated gene factor 3 (ISGF3), which binds to the IFN-stimulated response element (ISRE) within the promoters of interferon stimulated genes (ISGs) (1). Several ISGs have been shown to have specific antiviral activity and/or play a role in regulating the IFN response itself. Although this is a prompt and powerful defence against pathogens, a dysregulated type I IFN response can lead to autoinflammatory disease. Therefore, tight regulation of activating and inhibitory signals is of paramount importance for maintaining the protective host-defence nature of the response but limiting tissue damage.

ISG15, a ubiquitin-like (UbL) modifier, is synthesised from a precursor that is cleaved at the C-terminus to yield the mature 15-kDa protein with a C-terminal Leu-Arg-Leu-Arg-Gly-Gly (LRLRGG) tail (6,7). ISG15 exists as a free molecule but can also covalently bind to target proteins through the formation of an isopeptide bond between its terminal glycine and the lysine ε-amino group of the target protein (8), a process termed ISGylation (reviewed by (9,10)). ISGylation is reversible through the action of a deISGylase enzyme, the ubiquitin-specific protease 18 (USP18) (11). Independent to its isopeptidase activity on ISG15 (12,13), USP18 interacts with STAT2 to facilitate its recruitment to IFNAR2, where it can inhibit receptor dimerization by interfering with cytoplasmic interactions between IFNAR2 subunits (14–16).

Several reports now show that inherited ISG15-deficiency in humans causes type I interferonopathy and autoinflammatory disease (17–19). ISG15-deficient cells exhibited enhanced and prolonged ISG expression and a concomitant resistance to virus infection (18,20,21) a phenotype also associated with USP18 deficiency (13,22). Indeed, despite high levels of *USP18* transcription, USP18 protein levels in ISG15-deficient cells are very low

(18,20,21) and previous reports have shown that intracellular ISG15 is required for rescuing

USP18 from S-phase kinase associated protein 2 (SKP2)-mediated proteasomal degradation

(18,23).

Here, we show that, contrary to existing models, the ISG15-dependent stabilisation of

USP18 is necessary *but not sufficient* for regulation of type I IFN signalling. Furthermore,

independently of ISG15's ability to stabilise USP18, we show that ISG15 and USP18 must

interact for USP18 to negatively regulate type I IFN signalling. We find that the C-terminal di-

Gly moiety of ISG15 is critical for a stable ISG15-USP18 interaction and we unambiguously

demonstrate that abolishing the non-covalent interaction between ISG15 and USP18 results

in phenotypes associated with enhanced IFN-α signalling. Together our data illustrates that a

binding mechanism normally reserved for deISGylation has been co-opted to serve a crucial

role in the fine-tuning of early intracellular immune responses.

**Results**

**The C-terminal di-Gly motif of ISG15 is important for IFN-α signalling regulation.**

In work leading up to this report, we found that reconstituting expression of C-

terminal mutants of ISG15, where the terminal di-Gly motif was replaced with di-Ala, in ISG15-

deficient cells did not restore the regulation of IFN signalling. To dissect the role of the ISG15

C-terminus, we reconstituted expression of Myc-tagged ISG15, which retained its di-Gly motif

(ISG15.GG), or C-terminal mutants of ISG15, where the di-Gly motif was either replaced by di-

Ala (ISG15.AA) or deleted (ISG15.ΔGG), in our phenotypically validated A549-ISG15[-/-] cell line

(21). To better mimic physiological conditions, reconstituted ISG15 was placed under the

control of the native ISG15 promoter (pr15) (Fig. 1A). IFN-α stimulation induced expression

of Myc-ISG15 to levels similar to endogenous ISG15 in control A549s and, as expected,

ISG15.ΔGG and ISG15.AA did not ISGylate proteins (Fig. 1B). These experimentally tractable

cells are therefore valuable for deciphering the role of ISG15 or other ISGs during the innate

immune response.

We have previously shown that IFN-α treatment of A549-ISG15[-/-] cells results in

enhanced signalling characterised by elevated phospho-STAT1 levels and an augmented and

prolonged expression of ISGs (21). Here we determined the ability of C-terminal mutants of

ISG15 to regulates signalling. Cells were treated with IFN-α for 30 min, extensively washed,

and re-incubated in media without IFN-α. Cell lysates taken immediately after 30-min

treatment (0 min) and 30 min later (0.5 h) showed high levels of STAT1 phosphorylation in all

tested cell lines (Fig. 1C). Following 24-h treatment with IFN-α, there was evident expression

of the ISGs MxA and ISG15, Myc-ISG15 and enhanced expression of STAT1 (Fig. 1C).

Interestingly, although phospho-STAT1 levels had declined in A549 control and ISG15.GG-

expressing cells at 24 h after IFN-α treatment, levels were clearly higher in A549-ISG15[-/-] cells

and the cells expressing ISG15.AA and ISG15.ΔGG, signifying higher levels of IFN-α signalling

in these cells.

To confirm these observations at the level of ISG transcripts, cells were treated with

IFN-α for 24 h or left untreated. As expected, expression of *MxA* and *HERC5* mRNA was

significantly higher in A549-ISG15[-/-] cells compared to A549 control (2.6- and 3.3-fold

respectively, Fig. 1D). Consistent with the phosho-STAT1 levels, ISG mRNA levels were also

significantly higher in cells expressing the ISG15.AA and ISG15.ΔGG mutants. Intriguingly, this

demonstrated a gradient pattern of regulation where ISG15.GG-expressing cells regulated

similar to control A549 cells, cells expressing the ISG15.AA mutant were characterised by

higher ISG levels similar levels to A549-ISG15$^{-/-}$ cells, but cells expressing the ISG15.ΔGG

mutant consistently showed a trend towards an intermediate level of regulation. No

significant difference was observed between the ISG15.GG-expressing cells and the control,

suggesting that the level of ISG15 expression in this system is sufficient to regulate IFN

signalling similarly to unmodified control cells (Fig. 1D). Overall, these data show that the C-

terminal di-Gly motif is important for the negative regulation of type I IFN signalling.

**Quantitative analysis of global protein abundance shows enhanced IFN-α immune**

**responses demonstrates the importance of the ISG15 C-terminus for regulation.**

Next, we employed a multiplexed proteomic approach to obtain an unbiased, global

picture of the proteomic changes induced by IFN-α across our cell lines and to begin to tease

out the processes that could underpin interferonopathies. Cells were treated with IFN-α for

24, 48 or 72 h, and whole cell protein abundance measured using 16-plex TMT labelling and

MS3 mass spectrometry (Fig. 2A). In total, 7112 proteins were quantified (full data can be

visualised using the 'Plotter' in Supplementary File 1). As expected, the abundance of IFN-

stimulated proteins, such as ISG20 and IFIT1 was higher in the A549-ISG15$^{-/-}$ and ISG15.AA-

expressing cells compared to the A549 control and ISG15.GG-expressing cells (Fig. 2B). To

determine whether this trend was reflected more generally, ISGs were identified by (a)

comparison to the Interferome database, and (b) stringent criteria to identify proteins

upregulated by IFN-α compared to mock treatment in A549 control cells (>1.7-fold increase

in abundance, Fig. 2C). Ninety proteins identified by these criteria as interferon stimulated

were expressed at significantly higher levels by A549-ISG15$^{-/-}$ and ISG15.AA cells compared to

both A549 control and ISG15.GG cells (Fig. 2C). As expected, there was no statistically

significant difference in ISG expression between A549 control and ISG15.GG-expressing cells,

providing further evidence that genetic manipulation did not cause any inadvertent

phenotypic changes to the cells.

To determine whether specific pathways were enriched in the ISG15$^{-/-}$ and ISG15.AA

cells, we used DAVID analysis to examine proteins upregulated >1.7-fold compared to control

A549 cells. As anticipated, effector molecules involved in cellular immune responses, such as

innate immunity and antiviral defence, were enriched in all tested cell lines with several

overlapping factors in each cluster (Fig. 2D). Although fewer pathways were enriched in ISG15.GG cells suggesting tighter control than in cells expressing C-terminal mutants, antiviral defense, immunity and innate immunity pathways were identified suggesting that their control was not as tightly regulated compared to control A549 cells. This might be because, despite being inducible by IFN-α, lentivirally-delivered ISG15 transgenes are not in their authentic genomic loci and therefore expression is affected. Interestingly, components of IFN-γ mediated signalling and antigen processing and presentation pathways were selectively enriched in A549-ISG15$^{-/-}$ and ISG15.AA-expressing cells, for example transporter associated with antigen presentation 1 (TAP1) (Fig. 2D-E). One potential explanation is the selective enrichment in these cells of key regulatory factors induced by primary IFN-γ signalling (24,25), such as the interferon-regulatory factor 1 (IRF1) whose expression is induced by STAT1 homodimers typical of IFN-γ-induced ISGs with gamma-activated sequences (GAS) in their promoters and not STAT1-STAT2-IRF9 (ISGF3) complexes that characteristically induce genes with IFN-stimulated response elements (ISRE) activated by type I IFN signalling (although GAS genes can be activated upon type I IFN signalling as concentrations of STAT1-homodimers increase; Fig. 2E). It may be that enrichment with components of IFN-γ signalling derives through the enhanced levels of active STAT1 (Fig. 1C) and subsequent increased stoichiometry of STAT1 homodimer complexes in ISG15$^{-/-}$ and ISG15.AA cells, leading to gene expression through binding to GAS elements in the promoters of ISGs (26). Moreover, cell cycle components and several factors with nucleotide binding properties were also enriched in cells with a dysregulated IFN-α signalling response (Fig. 2D). In conclusion, using unbiased proteomic analyses, we have further confirmed that the C-terminal di-Gly domain of ISG15 is important for regulation of type I IFN responses. It is also possible that pathways associated with IFN-γ, which include genes associated with antiproliferative phenotypes and apoptosis, may underpin the pathogenesis of autoinflammatory diseases associated with loss of type I IFN signalling control.

**IFN-α-pretreatment leads to viral resistance in cell lines expressing the C-terminal mutants, ISG15.AA and ISG15.ΔGG.**

We and others have previously shown that pre-treatment of ISG15-deficient cells with IFN-α renders them resistant to viral infection (20,21). Because we have reported that viral resistance in ISG15-deficient cells is directly related to dysregulated type I IFN signalling (21),

this assay serves as an excellent model for investigating the biological implications of ISG15

loss-of-function and the regulatory role of the ISG15 di-Gly motif. Cell lines were primed with

IFN-α for 18 h, then infected with a recombinant parainfluenza virus type 5 (PIV5) expressing

mCherry (rPIV5-mCherry). Because the PIV5 V-protein targets STAT1 for proteasomal

degradation, rPIV5-mCherry can replicate in A549 control cells despite a primed IFN response,

albeit with reduced kinetics (27). By 48 h of infection in A549 control cells, there was a

corresponding increase in PIV5 NP expression, STAT1 was undetectable and MxA expression

was reduced (Fig. 3A). As anticipated, pretreatment with IFN-α rendered A549-ISG15[-/-] cells

resistant to PIV5 infection, whereas reconstituted expression of ISG15.GG reversed the

phenotype (Fig. 3A,B). Interestingly, like A549-ISG15[-/-] cells, NP expression was undetectable

at 24 h post-infection in both ISG15.AA and ISG15.ΔGG-expressing cells (Fig. 3A). Although

PIV5 infection recovered to some extent by 48 h, NP expression was still significantly reduced

by 90% and 45% at 48 h p.i. in cells expressing ISG15.AA and ISG15.ΔGG, respectively (Fig. 3B).

Intriguingly, this gradient in viral protein expression inversely correlated with the gradient

observed for ISG expression (Fig. 1D), linking the ability of ISG15 and ISG15 mutants to

regulate the magnitude of the antiviral response with their capacity to support infection.

Moreover, we used fluorescence microscopy to visualise mCherry expression and indicate

rPIV5-mCherry replication. Consistent with NP expression, no mCherry was detected in the

A549-ISG15[-/-] cells, with low or moderate levels in cells expressing the ISG15.AA and

ISG15.ΔGG mutants, respectively (Fig. 3C). Altogether, these experiments highlight the impact

on viral infection of the C-terminal di-Gly motif of ISG15, via its regulation of IFN signalling.

**202      The ISG15-dependent stabilisation of USP18 is necessary but not sufficient for the**

**203      regulation of IFN signalling.**

It has been established that ISG15 is crucial for sustaining the levels of USP18, a key

negative regulator of IFN signalling, by preventing its SKP2-mediated ubiquitination and

proteasomal degradation (18,20,23). Therefore, we reasoned that modifications to the ISG15

C-terminus might have affected its ability to stabilise USP18. To test this, cells were treated

with IFN-α for 24 and 48 h and cell lysates were subjected to immunoblot analysis.

Remarkably, although USP18 expression was dramatically reduced in A549-ISG15[-/-] cells, its

expression in ISG15.AA and ISG15.ΔGG expressing cells were comparable to levels observed

in ISG15.GG-expressing and control A549 cells (Fig. 4A). The proteomics analysis described in

Fig. 2 provided independent verification that ISG15 expression, independently of its ability to regulate signalling, was able to stabilise USP18 (Fig. 4B). Taken together, these experiments clearly show that, although the stabilization of USP18 is crucial, it is not sufficient for the regulation of type I IFN signalling.

**The C-terminal di-Gly motif of ISG15 is required for the ISG15-USP18 interaction.**

Because neither ISGylation (18,21) or ISG15-dependent stabilisation of USP18 are sufficient for the regulation of type I IFN signalling, the mechanism of ISG15 action with respect to the IFN pathway is likely to involve a non-covalent protein-protein interaction. The most likely candidate is USP18, given that during the deISGylation process, USP18 deconjugates ISG15 from target substrates in a process that requires engagement of USP18's catalytic triad with the C-terminal tail LRLRGG of ISG15 (28,29). We therefore assessed the ability of endogenous USP18 to interact with ISG15 and its C-terminal mutants in our reconstituted cell lines. Myc-tagged ISG15 was immunoprecipitated from ISG15-reconstituted cells following a 24-h IFN-α treatment. To verify that ISGylation was not necessary for the interaction between ISG15 and USP18, we knocked-out UBA7 (E1 enzyme essential for ISGylation) from our ISG15.GG-expressing cells by CRISPR/Cas9 genome editing (Fig. 5). Here we show that the ability of ISG15 mutants to bind USP18 mirrored the gradient pattern of ISG mRNA regulation (Fig. 1D) and corresponding effects on viral infection (Fig. 3). Reconstituted ISG15.GG efficiently bound USP18; however, the ISG15.AA mutant was unable to interact with USP18, whereas ISG15.ΔGG did interact but at reduced levels compared to ISG15.GG (Fig. 5). Notably, USP18 co-immunoprecipitated with ISG15.GG in UBA7$^{-/-}$ cells, confirming that the ISG15-USP18 interaction is not dependent on ISGylation (Fig. 5). Furthermore, we noted that all forms of ISG15 stabilised USP18 (see WCL samples, Fig. 5 lower panels), again confirming that ISG15 stabilised USP18 independently of their ability to regulate signalling and to add to this, stability is not dependent on their interaction. Collectively, these results indicate that the C-terminal di-Gly motif is important for ISG15-USP18 interaction and that non-covalent binding of ISG15 and USP18 is necessary to facilitate USP18's inhibition of type I IFN signalling.

**The ISG15-USP18 interaction is important for the tight regulation of IFN-α signalling.**

To independently assess the requirement of the ISG15-USP18 interaction for the
regulation of IFN signalling, we mutated the USP18 isoleucine residue at position 60
(USP18.I60N), which is known to abolish interactions with ISG15 (23). In addition, since
previous studies have shown that the mouse Usp18 (Ubp43) negatively regulates IFN
signalling in the absence of its isopeptidase activity (13), we additionally investigated the
catalytically inactive human mutant USP18.C64S. We first sought to evaluate the impact of
these point-mutations on ISG15-USP18 binding by coimmunoprecipitation (co-IP) (Fig. 6A).
Consistent with previous studies (23), our data demonstrated that the USP18.I60N mutant
was unable to interact with ISG15.GG, whereas the USP18.C64S mutant had stronger binding
affinity for ISG15 compared to wt USP18 (23) (Fig. 6A).

Next, we asked whether the USP18.I60N mutant could still be recruited to the IFNAR2
signalling complex despite its inability to interact with ISG15. Previous studies have shown
that STAT2 recruits USP18 to the IFNAR2 receptor, where it interferes with the cytosolic
interactions between the type I IFN receptor subunits (13,16,30). We co-expressed a V5-
tagged version of the IFNAR2 cytosolic domain (aa 265-515) and Myc-tagged STAT2 with Flag-
tagged wt USP18, USP18.I60N or USP18.C64S in HEK293T cells and performed co-IP assays
using anti-V5 antibody coupled to protein G Dynabeads (Fig. 6B). As expected, the IFNAR2
cytoplasmic domain interacted with STAT2 and wt USP18 (Fig. 6B). Intriguingly, both mutant
forms of USP18 co-immunoprecipitated with the IFNAR2 receptor subunit (Fig. 6B),
demonstrating that neither the I60N or C64S point mutations disrupted the recruitment of
USP18 to the receptor or the subsequent formation of the USP18-dependent type I IFN
receptor inhibitory complex.

In order to evaluate the functional consequences of these point mutants, we used
CRISPR/Cas9 genome editing to knock-out USP18 gene expression in A549 cells, as described
before (21), followed by lentiviral transduction to reconstitute inducible expression of Flag-
tagged wt or mutant forms of USP18 in the A549-USP18$^{-/-}$ cell line (Fig. 6C). To evaluate USP18
expression, A549 NC1-control cells, which express a negative control guide RNA (NC1) that is
nontargeting in humans, A549-USP18$^{-/-}$ and USP18-reconstituted derivatives were treated
with IFN-α for 48 h and cell lysates were subjected to immunoblot analysis (Fig. 6D).
Interestingly, the expression levels of the reconstituted forms of USP18 were higher
compared to NC1 control (Fig. 6D), perhaps because expression of reconstituted USP18 was
driven by the ISG15 promoter, which is strongly IFN-responsive (31), instead of its native

promoter. Consistent with previous observations (13), knockout of USP18 increased the levels

of ISG15 conjugates, whereas unexpectedly, reconstitution of wt USP18 resulted in a lower

level of ISGylation compared to the NC1 control cells. The expression of USP18.I60N mutant

was higher compared to the expression levels of wt USP18 and USP18.C64S; however, the

accumulation of ISGylated proteins was only marginally increased compared to the cell line

expressing wt USP18 (Fig. 6D). Interestingly, protein expression of ISGs, ISG15 and MxA,

appeared to be elevated in A549-USP18$^{-/-}$ and the USP18.I60N-expressing cells, signifying

higher levels of JAK/STAT signalling (Fig. 6D). As anticipated, expression of the catalytically

inactive mutant USP18.C64S resulted in higher levels of ISGylated proteins compared to the

cell line reconstituted with wt USP18 (Fig. 6D), and, similar to its mouse counterpart

(UBP43.C61S) (13), resulted in lower ISG15 protein expression compared to wt USP18.  These

observations indicate a dysregulation of the IFN response in an isopeptidase-independent

manner (Fig. 6D).

To further evaluate the importance of ISG15-USP18 binding in the regulation of IFN-α

signalling, as described in Fig. 1D, we measured the levels of *MxA* and *HERC5* gene expression

in these cells (Fig. 7A). Remarkably, *MxA* and *HERC5* expression was significantly higher in

USP168.I60N-expressing derivatives compared to NC1 control (averaging around 2.5-fold),

denoting that the lack of ISG15-USP18 interaction in these cells led to a dysregulated IFN

response, a phenotype similar to A549-USP18$^{-/-}$ cells (Fig. 7A). Notably, ISG expression levels

in cells expressing the USP18.C64S mutant were consistently lower than, but not significantly

different to, NC1 control cells.

We predicted that the elevated expression of ISGs in USP18.I60N-expressing cells

would engender resistance to infection. Using the assay described in Fig. 3, we observed that

IFN-α-pretreated USP18.I60N-expressing cells were largely resistant to infection, as similar to

A549-USP18$^{-/-}$ cells, expression of PIV5 NP was barely detectable (Fig. 7B). Quantitative

analysis of NP expression levels showed that viral replication was reduced more than 95% in

IFN-α-pretreated A549-USP18$^{-/-}$ and USP18.I60N-expressing cells, whereas the levels of viral

infection in cell lines expressing the wt and the catalytically inactive mutant USP18.C64S were

similar to NC1 control, confirming regulation in these cells (Fig. 7C). mCherry expression levels

further verified that IFN-α-pretreatment constrained virus replication in A549-USP18$^{-/-}$ and

USP18.I60N-expressing cells (Fig. 7D). Altogether, these experiments demonstrate that

disruption of the ISG15-USP18 interaction enhances IFN-mediated signalling, suggesting that

ISG15 plays a crucial role in the negative regulation of IFN signalling beyond its indirect

function as a USP18 stabiliser.

**The ISG15-USP18 interaction is important for the IFN-α-induced desensitization of IFN-α**

**signalling.**

Previous studies have shown that USP18 is crucial for the IFN-α-induced

desensitization of IFN-α signalling by disrupting the recruitment of IFNAR1 into the ternary

IFN-α-IFNAR1-IFNAR2 complex, decreasing the activation of signalling (15,16,30,32). To test

whether the ISG15-USP18 interaction is important for this USP18-dependent negative

regulation of IFN receptor plasticity, we established a desensitization assay based on previous

reports (15), where A549 NC1-control cells, A549-USP18$^{-/-}$ and USP18-reconstituted

derivatives were primed with IFN-α for 8 h or left untreated, washed extensively and then

maintained in medium without IFN for 16 h. During the prime-rest phase, USP18 is expressed

and prevents further signalling activation following additional stimulation. Following the 16-

h resting period, cells were stimulated with IFN-α for 30 min and cell lysates were subjected

to immunoblotting to assess phosho-STAT1 expression, which is indicative of early activation

of IFN signalling (Fig. 8A). As expected, priming with IFN-α decreased the responsiveness of

NC1-control cells to subsequent IFN stimulation as similar levels of phosho-STAT1 were

detected in primed and non-primed control cells. A similar phenotype was observed in A549-

USP18$^{-/-}$ cells reconstituted with wt USP18 or the catalytically inactive mutant USP18.C64S,

indicating that the sensitivity of these cells to IFN-α was downregulated following priming

with IFN-α (Fig. 8A). Consistent with previous reports (15), A549-USP18$^{-/-}$ cells were not

desensitized to IFN-α resulting to 3.8-fold increase in STAT1 phosphorylation when primed

with IFN-α compared to the non-primed control (Fig. 8B). Remarkably, cells expressing the

USP18.I60N mutant retained their responsiveness toward IFN-α similar to A549-USP18$^{-/-}$ cells

(Fig. 8A). Specifically, STAT1 phosphorylation in USP18.I60N-expressing cells showed 3.2-fold

increase following IFN-α priming compared to the non-primed control (Fig. 8B). Overall, our

data strongly suggest that the ISG15-USP18 interaction is important for the USP18-dependent

regulation ternary IFNAR complexes.

**Discussion**

The negative regulation of the type I IFN system is controlled at multiple levels by a

variety of mechanisms, involving sequestration of effector molecules and post-translational

modifications, such as ubiquitination or dephosphorylation (reviewed by Arimoto et al (33)).

Humans with ISG15-deficiency display abnormally strong type I IFN immunity highlighting the

emerging role of ISG15 as a central regulator of immunity (17,18,21,23). Current models

suggest that ISG15 antagonizes the SKP2-mediated ubiquitination and degradation of USP18,

promoting its functions (18,20,23). We extend these findings and show that the ISG15-

dependent stabilisation of USP18 is necessary *but not sufficient* to regulate IFN-I signalling

and that non-covalent binding of ISG15 and USP18 is also necessary to facilitate USP18's

inhibitory function (Fig. 9). Intriguingly, this is achieved through co-option of a binding

mechanism normally reserved for deISGylation and identifies an unanticipated new function

for the UbL ISG15.

It is well known that the conserved C-terminal di-Gly motif of ISG15 is essential for

ISGylation (34,35); therefore, the ISGylation-deficient mutants, ISG15.AA and ISG15.ΔGG,

have been extensively used for exploring the functional consequences of ISGylation. Here, we

have reported that the ISG15 C-terminal mutants display different propensities for binding

USP18 but, despite this, both mutants stabilize USP18 to levels comparable to wt ISG15 (Fig.

4). Although not having been previously assessed in the context of an IFN response, this is

consistent with previous reports showing that ISG15 can abrogate the USP18-SKP2 complex

and rescue USP18 from proteasomal degradation independent of its ability to bind USP18

(23). Importantly, our work strongly suggests that ISG15-USP18 binding is required for USP18-

dependent regulation of the type I IFN response as the level of binding to USP18 faithfully

reflected the level of IFN-α signalling regulation and the permissiveness of cells to viral

infection. We independently confirmed the importance of a ISG15-USP18 interaction as

expression of mutant USP18 unable to bind ISG15 (USP18-I60N) could not regulate type I IFN

signalling even though it could interact with the IFNAR2 signalling complex (Fig. 6). In line with

findings using mouse Usp18 (12,13), the protease activity of human USP18 remains

dispensable for signalling regulation in humans.

Previous studies have shown that USP18 desensitises type I IFN signalling (14–16).

Specifically, STAT2 recruits USP18 to the IFNAR2, where it interferes with the cytosolic

interactions between receptor subunits, impeding the recruitment of IFNAR1 into the ternary

complex (14,16). Remarkably, we demonstrate here that the non-covalent binding of ISG15

and USP18 is required for the USP18-dependent negative regulation of IFNAR dimerization

(Fig. 8). Consistent with previous reports (14), our data show that ISG15 appears dispensable

for the interaction of USP18 IFNAR2; therefore, we hypothesize that the ISG15's contribution

may be essential for further stabilizing the USP18-inhibitory complex, or for recruiting further

not yet identified interaction partners of USP18 or perhaps promoting a conformational

change in USP18 necessary for its inhibitory activity.

Structural studies have shown that the ISG15-bound USP18 adopts a different

structural confirmation, where the 'switching loop' in the thumb domain of USP18 acquires

an active conformation, enabling access of the LRLRGG C-terminal tail of ISG15 into the

catalytic cleft (28,29). Here, we showed that replacing the C-terminal di-Gly of ISG15 with di-

Ala completely abolishes ISG15-USP18 interaction (Fig. 5). Although this amino acid

substitution is considered subtle, it is possible that the two extra methyl groups present in

the di-Ala motif are not accessible to the tight catalytic cleft of USP18, abolishing the

interaction with the ISG15.AA mutant. This may also explain why the ISG15.ΔGG mutant,

which retains the LRLR peptide of the C-terminal tail, interacted to a greater extent with

USP18 compared to ISG15.AA. Moreover, in line with previous findings (23), we showed that

the USP18.I60N mutant was unable to bind ISG15 (Fig. 6A). Interestingly, Ile residue 60 does

not belong to the ISG15-binding boxes (28) but its proximity to the catalytic cysteine (C64)

may affect the conformational dynamics of the catalytic cleft, diminishing the interaction with

ISG15. Consistent with previous reports (23), we have observed that the catalytically inactive

mutant (USP18.C64S) has stronger binding affinity for ISG15, compared to wt USP18, and

there is a consistent pattern of stronger negative regulation (lower ISG expression, increased

desensitization to IFN-α) in cells expressing this mutant, supporting our observation that the

level of binding of ISG15 toward USP18 determines the level of IFN-α signalling regulation.

Hence, it is possible that the binding of ISG15 locks USP18 into a more stable structural

confirmation that may serve its regulatory functions at the level of IFNAR assembly. A caveat

to this model is that Isg15 is dispensable for Usp18-mediated regulation of type I IFN signalling

in mice (20).

That Isg15 does not stabilise Usp18 in mice (20), illustrates interesting interspecies

variation with regard to ISG15 function. Unlike the highly conserved ubiquitin (36), ISG15

possesses remarkable sequence variation between species, with sequence identity around

70% between human and mouse genes, suggesting that the different biochemical properties

of ISG15 between species may be key determinants of ISG15's species-specific functions

(37,38). Indeed, it has been shown that human ISG15 associates with higher affinity to USP18

compared to its mouse counterpart (20), which in agreement with our findings, suggests that

gain-of-function mutations in ISG15 and/or USP18 that facilitate stronger ISG15-USP18

interactions have been evolutionary selected in humans. Why this trait is apparent in humans

and not mice is of interest and may suggest that an additional, yet-to-be identified factor is

taking the place of ISG15 in mice, or that humans may require a different level of IFN-

regulatory control.

Intriguingly, our proteomics profiling and pathway analyses highlighted the

enrichment of components involved in IFN-γ signalling in A549-ISG15[-/-] and ISG15.AA-

expressing cells (Fig. 2). This may be due to the accumulation of STAT1 homodimers as a by-

product of enhanced type I immunity, leading to enrichment of GAS-containing genes

(24,25,39), such as IRF1 and MHC Class I proteins in these cells. IRF1 transcription factor itself

is involved in the regulation of genes implicated in antiproliferative (40) and antigen

processing pathways (41–43). These findings support our previous work that demonstrates

that ISG15 deficiency leads to translational regression following IFN-α treatment (21) and

further suggest that intervention strategies that target the ISG15-USP18 interaction may be

of therapeutic use in anticancer therapy.

In conclusion, we have demonstrated that intracellular ISG15 is essential to negatively

regulate IFN-α responses via its non-covalent interaction with USP18, thereby averting

autoinflammatory consequences of uncontrolled type I IFN signalling. This hitherto human-

specific trait has been acquired through co-option of a binding mechanism normally reserved

for the process of deISGylation. Further investigation is needed to decipher the biophysical

properties of the USP18-mediated inhibitory complex at the level of IFNAR assembly and

resolve how the binding of ISG15 to USP18 facilitates those interactions.

**Methods**

**Cells**

HEK293T (human embryonic kidney cell), A549 cells (human adenocarcinoma alveolar basal epithelial cells), and derivatives, were were grown as monolayers in Dulbecco's modified Eagles's medium (DMEM; Sigma) supplemented with 10% (v/v) foetal bovine serum (FBS, Biowest) and incubated in 5% (v/v) $CO_2$ at 37°C in a humidified incubator.A549-ISG15$^{-/-}$ cells (clone B8) were generated as previously described using CRISPR/Cas9n system; transfectants were enriched by treating cells with puromycin (1 μg/ml) for 2 days, then single-cell cloned and successful knockout cells were validated by immunoblot analysis [21]. Lentiviral technology was used to reconstitute expression of wt ISG15.GG (NCBI Ref Seq, NM_005101.3) or C-terminal mutants, ISG15.AA and ISG15.ΔGG in A549-ISG15$^{-/-}$ to generate the following derivative cell lines; A549-ISG15$^{-/-}$:prI5-GG, A549-ISG15$^{-/-}$:prI5-AA, A549-ISG15$^{-/-}$:prI5-ΔGG, respectively. Expression of wt and C-terminal mutant forms of ISG15 was driven by the native ISG15 promoter (pr15) (NCBI Ref seq, NG_033033.2) cloned using our lentiviral vector. An internal ribosome entry site (ISRE) downstream the ISG15 gene allows expression of puromycin resistance (pac) gene following induction of pr15. Hence, for puromycin selections, cells were primed with 1000 IU/ml IFN-α for 4 h prior treatment with puromycin (1 μg/ml) for 2 days. To generate the A549-ISG15$^{-/-}$:UBA7$^{-/-}$:pr15-GG cell line, A549-ISG15$^{-/-}$:pr15-GG cells were further modified to stably express *Streptococcus pyogenes* Cas9 and then transduced with UBA7 sgRNA-expressing lentiGuide-Puro (sgRNA sequences; sense: caccGCACACGGGTGACATCACTG; antisense: aaacCAGTGATGTCACCCGTGTGC) as described previously [21]. For generating A549-USP18$^{-/-}$ cell lines, A549 cells were modified to stably express *Streptococcus pyogenes* Cas9 and then transduced with USP18 sgRNA-expressing lentiGuide-Puro (sgRNA sequences; sense: caccgGGGGCCGCACTGCTTTCTGC; antisense: aaacGCAGAAAGCAGTGCGGCCCCc). Blasticidin/puromycin-resistant A549-USP18$^{-/-}$ cells were single-cell cloned in 96-well plates and successful knockout cells were validated by immunoblot analysis. Lentiviral technology using our pr15 lentiviral vector was used to reconstitute expression of wt USP18 (NCBI Ref seq NM_017414.4) or the point mutants USP18.I60N and USP18.C64S (generated by site-directed mutagenesis) in A549-USP18$^{-/-}$ to generate the A549-USP18$^{-/-}$:prI5-USP18.wt, A549-USP18$^{-/-}$:prI5-USP18.I60N and A549-USP18$^{-/-}$:prI5-USP18.C64S, respectively. All lentiviruses used in this study were generated in

HEK293T cells using a previously described self-inactivating lentiviral constitutive expression

system (44).

**Reverse transcription quantitative PCR**

To measure ISG expression levels, cells were treated with 1000 IU/ml IFN-α for 18 h and

intracellular total RNA was isolated with TRIZOL (ThermoFisher Scientific) and Direct-zol RNA

Miniprep Plus kits (Zymo Research), including removal of contaminating DNA following DNase

I treatment, following the manufacturer's instructions. A total of 500 ng of RNA was reverse

transcribed using LunaScript® reverse transcriptase (New England Biolabs) according to the

manufacturer's recommendations. Quantitative PCRs (qPCRs) were performed with PerfeCTa

SYBR green SuperMix (Quanta BioScience) using fast two-step cycling performed in a

Mx3005P real time PCR machine (Stratagene) and included an initial 2 min enzyme activation

step at 95°C, followed by 40 cycles of 5 s at 95°C and 20 s at 60°C. Melting curve analysis was

performed to verify amplicon specificity. For each assay, quantification of *β-ACTIN* mRNA was

used to normalize between samples and the average cycle threshold (CT) was determined

from three independent cDNA samples from independent cultures (technical replicates). Fold

changes were determined from three independent assays performed at different times

(biological replicates). Relative expression compared to non-treated control cells was

calculated using the ΔΔCT method. Primer sequences were: *MxA*

5'GCCTGCTGACATTGGGTATAA and 5'CCCTGAAATATGGGTGGTTCTC, *HERC5*

5'GACGAACTCTTGCACCGTCTC and 5'GCGTCCACAGTCATTTTCCAC, *β-ACTIN*

5'AGCGAGCATCCCCCAAAGTT and 5'AGGGCACGAAGGCTCATCATT.

**Viral infections and IFN treatment**

Recombinant mCherry-expressing parainfluenza virus type 5 (rPIV5-mCherry)

(provided by Professor Biao He, University of Georgia) (45) stocks were prepared by

inoculating Vero cells at a multiplicity of infection (MOI) of 0.001 with continual rocking at

37°C. Supernatants were harvested at 2 d p.i., clarified by centrifugation at 3,000 x g for 15

min, aliquoted and snap frozen. Titres were estimated by standard plaque assay on Vero cells

in 6-well plates.

For virus resistance assays, cell monolayers at 70-80% confluency were infected in 6-

well plates with virus diluted in 1 ml media supplemented with 2% (v/v) FBS to achieve an

MOI of 10. Virus adsorption was for 1 h with continual rocking at 37°C, after which 1 ml of

media supplemented with 2% (v/v) FBS was added to the viral inoculum and incubated in 5%

(v/v) $CO_2$ at 37°C until harvested. When cells were treated with IFN-α prior to infection (pre-

treated) this was done with 1000 IU/ml IFN-α2b (referred to as IFN-α from here on; IntronA,

Merck Sharp & Dohme Ltd). IFN-α remained on cells for the duration of experiments. Cells

were either processed for immunoblot analysis or observed with fluorescence microscopy

using EVOS M5000 Imaging System (pictures taken at 10X magnification).

**Immunoblotting**

Confluent monolayers in 6-well dishes were lysed with 250 µl 2 x Laemmli sample

buffer (4% w/v SDS, 20% v/v glycerol, 0.004% w/v bromophenol blue and 0.125 M Tris-HCl,

pH 6.8 with 10% v/v β-mercaptoethanol) for 10 min, incubated at 95°C for 10 min, sonicated

at 4°C with 3 cycles of 30 s on 30 s off in a Bioruptor Pico (Diagenode) and clarified by

centrifugation at 12,000 x g, 4°C for 10 min. SDS-PAGE in Tris-glycine-SDS running buffer and

immunoblotting followed standard techniques using the following antibodies: mouse

monoclonal anti-ISG15 F-9 (Santa Cruz Biotechnology Cat# sc166755), rabbit polyclonal anti-

ISG15 H-150 (Santa Cruz Biotechnology Cat# sc50366), rabbit polyclonal anti-MxA

(Proteintech Cat# 13750-1-AP), mouse monoclonal anti-UBA7 (anti-UBE1L B-7; Santa Cruz

Biotechnology Cat# sc-390097), rabbit anti-USP18 (Cell Signalling Technology Cat# 4813S),

mouse monoclonal anti-total STAT1 (N-terminus; BD Transduction Laboratories™ Cat#

610116), rabbit monoclonal anti-phosphorylated STAT1 (anti-phospho-STAT1 (Tyr701); Cell

Signalling Technology Cat# 9167), mouse monoclonal anti-PIV5 NP 125 (46), mouse anti-V5

tag Pk 336 (46), antibody mouse monoclonal anti-β-actin (Sigma Cat# A2066). For quantitative

immunoblots primary antibody-probed membranes were incubated with IRDye secondary

antibodies (LiCOR) and signals detected using an Odyssey CLx scanner. Data were processed

and analysed using Image Studio software (LiCOR).

**Immunoprecipitation**

Prior immunoprecipitation, HEK293T cells were transiently transfected to express the

proteins of interest using calcium-phosphate coprecipitation transfections. One day prior to

transfection, HEK293T cells were seeded in 6-well plates such that they are logarithmically

growing on the day of transfection (i.e., 50-60% confluent). A total of 200 µl calcium-

phosphate precipitate was prepared for each well by mixing each plasmid DNA (diluted in

total 90 µl in dH$_2$O) with 10 µl of 2.5M CaCl$_2$ solution.  DNA/CaCl$_2$ solutions were added

dropwise into 100 µl of 2× HEPES-buffered saline (HeBS) (50 mM HEPES, 0.28 M NaCl, 10 mM

KCl, 1.5 mM Na$_2$HPO$_4$, 12 mM D-glucose, pH 7.05) and incubated for 20 min at room

temperature (RT). Chloroquine diphosphate solution was added to cell culture media to 25

µM final concentration and calcium phosphate precipitate was added dropwise onto plate

and mixed gently. At 16 h after transfection, the calcium phosphate precipitate was removed,

and cells were incubated for further 24 h before subjected to co-immunoprecipitation (co-IP)

assays. The following vectors were used for co-IP assays: pLHCX-STAT2-Myc (kind gift from Dr

Michael Nevels, St Andrews University), pcDNA3.1-USP18.wt-3XFlag, pcDNA3.1-USP18.I60N-

3XFlag, pcDNA3.1-USP18.C64S-3XFlag, pcDNA3.1.ISG15.GG and pcDNA3.1-IFNAR2.CTD-V5,

which expresses the cytoplasmic C-terminal domain (CTD) of IFNAR2.

For co-IP assays, confluent monolayers of IFN-treated A549 derivatives grown in T150

cm$^2$ flasks or plasmid-transfected HEK293T cell cultures grown in 6-well plates were harvested

in phosphate-buffered saline (PBS), pelleted by centrifugation (300 × g, 5 min, 4°C) and

resuspended in 1 ml co-IP lysis buffer (50 mM Tris pH 7.5, 150 mM NaCl, 0.1% [v/v] Triton-X)

supplemented with 1x cOmplete$^{TM}$ protease inhibitor cocktail (Merck). The supernatant was

separated by centrifugation (12,000 x g, 10 min, 4°C) and incubated overnight at 4°C with

gentle rotation with 40 µl Pierce Anti-c-Myc Magnetic Beads (ThermoFisher Scientific Cat#

88842) or anti-Flag$^®$ M2 Magnetic Beads (Merck Cat# M8823) or anti-V5 tag antibody coupled

with Protein G Dynabeads$^{TM}$ (Invitrogen) by following manufacturer's instructions. Complexes

were washed three times with co-IP wash buffer (1X TBS; 50 mM Tris pH 7.5, 150 mM NaCl),

incubated in 50 µl 1X NuPage LDS sample buffer (ThermoFisher Scientific) for 20 min at RT

and then, further incubated for 10 min at 70°C. Beads were magnetically separated and 10%

(v/v) of β-mercaptoethanol was added in the eluted supernatant containing target antigens.

Immunoprecipitates were subjected to SDS-PAGE and immunoblotting.

**Desensitization assay.**

Cells grown to 70-80% confluency in 6-well plates were primed with 2000 IU/ml IFN-

α (equivalent to 20 ng/ml) for 8 h or left untreated. Following priming, cells were washed four

times with PBS and maintained in medium without IFN for 16 h (resting period) and then

stimulated with 2000 IU/ml IFN-α for 30 min. Cell lysates were subjected to SDS-PAGE and

immunoblotting.

**Tandem mass tags (TMT)-based quantitative proteomics**

Tandem mass tags (TMT)-based proteomics was used to quantify differences in

protein abundance following IFN-α treatment in A549 control cells, A549-ISG15$^{-/-}$ and the

A549-ISG15$^{-/-}$:prI5-AA and A549-ISG15$^{-/-}$:prI5-GG derivatives. Cells grown to 60-70%

confluency in T25 flasks were treated with 1000 IU/ml IFN-α for 24, 48 and 72 h or left

untreated for 72h. Whole cell lysate protein digestion was performed as described before

(47) for each time point. For lysis, cells were washed twice with PBS, and 250 ml lysis buffer

added (6 M Guanidine/50 mM HEPES pH 8.5). Cells were scraped in lysis buffer, vortexed

extensively and then sonicated. Cell debris was removed by centrifuging at 21,000 g for 10

min, twice. For the Dithiothreitol (DTT) was added to a final concentration of 5 mM and

samples were incubated for 20 min. Cysteines were alkylated with 14 mM iodoacetamide and

incubated 20 min at room temperature in the dark. Excess iodoacetamide was quenched with

DTT for 15 min. Samples were diluted with 200 mM HEPES pH 8.5 to 1.5 M Guanidine followed

by digestion at room temperature for 3 h with LysC protease at a 1:100 protease-to-protein

ratio. Samples were further diluted with 200 mM HEPES pH 8.5 to 0.5 M Guanidine. Trypsin

was then added at a 1:100 protease-to-protein ratio followed by overnight incubation at 37°C.

The reaction was quenched with 5% formic acid, then centrifuged at 21,000 g for 10 min to

remove undigested protein. Peptides were subjected to C18 solid-phase extraction (SPE, Sep-

Pak, Waters) and vacuum-centrifuged to near-dryness.

Samples were prepared for TMT labelling as previously described (47). Desalted

peptides were dissolved in 200 mM HEPES pH 8.5 and 25 mg of peptide labelled with TMT

reagent. TMT reagents (0.8 mg) were dissolved in 43 ml anhydrous aceto- nitrile and 3 ml

added to peptide at a final acetonitrile concentration of 30% (v/v). Following incubation at

room temperature for 1 h, the reaction was quenched with hydroxylamine to a final

concentration of 0.3% (v/v). Sample labelling was performed using 16-plex labelling reagent

(ThermoFisher Scientific$^{TM}$ CAT# A44520) and TMT-labelled samples were combined at a

1:1:1:1:1:1:1:1:1:1:1:1:1:1:1:1 ratio. Samples were vacuum-centrifuged to near dryness and

subjected to C18 SPE (Sep-Pak, Waters). An unfractionated single shot was analysed initially

to ensure similar peptide loading across each TMT channel, thus avoiding the need for

excessive electronic normalization. As all normalisation factors were >0.5 and <2, data for
each singleshot experiment was analysed with data for the corresponding fractions to
increase the overall number of peptides quantified.

TMT-labelled tryptic peptides were subjected to pH reversed-phase (HpRP)
fractionation using an Ultimate 3000 RSLC UHPLC system (Thermo Fisher Scientific) equipped
with a 2.1 mm internal diameter (ID) x 25 cm long, 1.7 mm particle Kinetix Evo C18 column
(Phenomenex). Mass spectrometry data was acquired using an Orbitrap Lumos and an
ultimate 3000 RSLC nano UHPLC equipped with a 300 mm ID x 5 mm Acclaim PepMap m-
Precolumn (Thermo Fisher Scientific) and a 75 mm ID x 50 cm 2.1 mm particle Acclaim
PepMap RSLC analytical column was used as described before (47).

**Data analysis of MS spectra**

For MS3-based TMT, as previously described (47), TMT tags on lysine residues and
peptide N termini (229.162932 Da) and carbamidomethylation of cysteine residues (57.02146
Da) were included as static modifications. Proteins were quantified by summing TMT reporter
ion counts across all matching peptide-spectral matches using 'MassPike', as described
previously (48). Briefly, a 0.003 Th window around the theoretical m/z of each reporter ion
(126, 127 n, 128 n) was scanned for ions, and the maximum intensity nearest to the the-
oretical m/z was used. An isolation specificity filter with a cutoff of 50% was employed to
minimise peptide co-isolation (48). Peptide-spectral matches with poor quality MS3 spectra
(more than 3 TMT channels missing and/or a combined S:N ratio of less than 100 across all
TMT reporter ions) or no MS3 spectra at all were excluded from quantitation. Peptides
meeting the stated criteria for reliable quantitation were then summed by parent protein, in
effect weighting the contributions of individual peptides to the total protein signal based on
their individual TMT reporter ion yields. Protein quantitation values were exported for further
analysis in Excel.

For protein quantitation, reverse and contaminant proteins were removed, then each
reporter ion channel was summed across all quantified proteins and normalised assuming
equal protein loading across all channels. Protein hits quantified by a single peptide were
removed from the dataset. The expression profile of each protein was observed after
comparing protein abundance to the condition (cell line/time point) with the highest MS

intensity score (set to 1) and normalised values were plotted against each time point for each

cell line (see Plotter in Supplementary File 1)

**Pathway Analysis**

625             To identify individual ISGs from our dataset, a list of 7112 gene symbols were searched

in 'Interferome v2.01' (http://interferome.its.monash.edu.au/interferome/home.jspx) (49).

The Interferome analysis was conducted on fold change values, which were calculated by

dividing the MS intensity score of each identified protein at a given time point post IFN

treatment (24, 48, 72 h) by the MS intensity score of the untreated control for each cell line.

A protein hit was considered to be an ISG if it was upregulated at least 1.7-fold in A549 control

cells following IFN-α treatment.

632             The Database for Annotation, Visualization and Integrated Discovery (DAVID) version

6.8 (https://david.ncifcrf.gov) was used to stringently identify cell line-specific enriched

pathways. An 'enrichment ratio' for each protein was obtained for each time point as follows

(MIS is MS intensity score):

$$Enrichment\ ratio = \left( \frac{Test\ cells\ MIS\ IFNa\ treated}{Test\ cells\ MIS\ untreated} \right) / \left( \frac{Control\ cells\ MIS\ IFNa\ treated}{Control\ cells\ MIS\ untreated} \right)$$

Protein hits upregulated at least 1.7-fold following enrichment were submitted using the

UniProt accession number and default medium classification stringency. Clusters were

considered significant if the Benjamini-Hochberg adjusted p-value was <0.05.

640             In the interest of simplicity, here we present the bioinformatic analyses only for the

641     48-h time point, which best represents the phenotypes observed in this experiment.

643     **Data availability**

644             The mass spectrometry proteomics data have been deposited to the

645     ProteomeXchange Consortium (http://www.proteomexchange.org) via the PRIDE (50)

646     partner repository with the dataset identifier (Accession # here).

648     **Figures**

649             Figures 1A, 2A, 6C and 9 were created with Biorender.com.

**Acknowledgments**

We are indebted to technical assistance provided by Dan Young. This work was

supported by grants from the Academy of Medical Sciences (SBF003/1028 to DJH), Wellcome

Trust Institutional Strategic Support Fund (to DJH), Wellcome Trust (101788/Z/13/Z to RER),

the UK Medical Research Council (MC_UU_12014/1 to JM and CGGB) and by the Wellcome

Trust via a Senior Clinical Research Fellowhship to MPW (108070/Z/15/Z).

**References**

1.  Ivashkiv LB, Donlin LT. Regulation of type I interferon responses. Nat Rev Immunol.
    2014;14(1):36–49.

2.  Isaacs A, Lindenmann J. Virus Interference. I. The Interferon. Proc R Soc London Ser B
    - Biol Sci. 1957;147(927):258–67.

3.  Kim SH, Cohen B, Novick D, Rubinstein M. Mammalian type I interferon receptors
    consists of two subunits: IFNaR1 and IFNaR2. Gene. 1997 Sep;196(1–2):279–86.

4.  Gauzzi MC, Velazquez L, McKendry R, Mogensen KE, Fellous M, Pellegrini S.
    Interferon-alpha-dependent activation of Tyk2 requires phosphorylation of positive
    regulatory tyrosines by another kinase. J Biol Chem. 1996 Aug 23;271(34):20494–500.

5.  Li X, Leung S, Kerr IM, Stark GR. Functional subdomains of STAT2 required for
    preassociation with the alpha interferon receptor and for signaling. Mol Cell Biol.
    1997 Apr;17(4):2048–56.

6.  Potter JL, Narasimhan J, Mende-Mueller L, Haas AL. Precursor processing of pro-
    ISG15/UCRP, an interferon-β-induced ubiquitin-like protein. J Biol Chem.
    1999;274(35):25061–8.

7.  Narasimhan J, Wang M, Fu Z, Klein JM, Haas AL, Kim JP. Crystal Structure of the
    Interferon-induced Ubiquitin-like. 2005;280(29):27356–65.

8.  Zou W, Papov V, Malakhova O, Kim K Il, Dao C, Li J, et al. ISG15 modification of
    ubiquitin E2 Ubc13 disrupts its ability to form thioester bond with ubiquitin. Biochem
    Biophys Res Commun. 2005 Oct 14;336(1):61–8.

9.  Zhang D, Zhang D-E. Interferon-Stimulated Gene 15 and the Protein ISGylation
    System. J Interf Cytokine Res. 2011;31(1):119–30.

10. Villarroya-beltri C, Guerra S, Sa F. ISGylation – a key to lock the cell gates for
    preventing the spread of threats. 2017;2961–9.

11. Malakhov MP, Malakhova OA, Il Kim K, Ritchie KJ, Zhang DE. UBP43 (USP18)
    specifically removes ISG15 from conjugated proteins. J Biol Chem.
    2002;277(12):9976–81.

12. Ketscher L, Hannß R, Moralese DJ, Basters A, Guerra S, Goldmann T, et al. Selective
    inactivation of USP18 isopeptidase activity in vivo enhances ISG15 conjugation and
    viral resistance. Proc Natl Acad Sci U S A. 2015;112(5):1577–82.

13. Malakhova OA, Kim K Il, Luo J-K, Zou W, Suresh Kumar KG, Fuchs SY, et al. UBP43 is a

novel regulator of interferon signaling independent of its ISG15 isopeptidase activity.

EMBO J. 2006;25:2358–67.

14.  Arimoto KI, Löchte S, Stoner SA, Burkart C, Zhang Y, Miyauchi S, et al. STAT2 is an

essential adaptor in USP18-mediated suppression of type i interferon signaling. Nat

Struct Mol Biol. 2017;24(3):279–89.

15.  François-Newton V, de Freitas Almeida GM, Payelle-Brogard B, Monneron D, Pichard-

Garcia L, Piehler J, et al. USP18-based negative feedback control is induced by type I

and type III interferons and specifically inactivates interferon α response. PLoS One.

2011;6(7).

16.  Wilmes S, Beutel O, Li Z, Francois-Newton V, Richter CP, Janning D, et al. Receptor

dimerization dynamics as a regulatory valve for plasticity of type I interferon

signaling. J Cell Biol. 2015;209(4):579–93.

17.  Bogunovic D, Byun M, Durfee LA, Abhyankar A, Sanal O, Mansouri D, et al.

Mycobacterial Disease and Impaired IFN-γ Immunity in Humans with Inherited ISG15

Deficiency. Science (80- ). 2012;337:1684–9.

18.  Zhang X, Bogunovic D, Payelle-Brogard B, Francois-Newton V, Speer SD, Yuan C, et al.

Human intracellular ISG15 prevents interferon-α/β over-amplification and auto-

inflammation. Nature. 2015;517(7532):89–93.

19.  Martin-Fernandez M, Bravo García-Morato M, Gruber C, Murias Loza S, Malik MNH,

Alsohime F, et al. Systemic Type I IFN Inflammation in Human ISG15 Deficiency Leads

to Necrotizing Skin Lesions. Cell Rep. 2020;31(6):107633.

20.  Speer SD, Li Z, Buta S, Payelle-Brogard B, Qian L, Vigant F, et al. ISG15 deficiency and

increased viral resistance in humans but not mice. Nat Commun. 2016;7(11496):DOI:

10.1038/ncomms11496.

21.  Holthaus D, Vasou A, Bamford CGG, Andrejeva J, Paulus C, Randall RE, et al. Direct

Antiviral Activity of IFN-Stimulated Genes Is Responsible for Resistance to

Paramyxoviruses in ISG15-Deficient Cells. J Immunol. 2020;205(1):261–71.

22.  Meuwissen MEC, Schot R, Buta S, Oudesluijs G, Tinschert S, Speer SD, et al. Human

USP18 deficiency underlies type 1 interferonopathy leading to severe pseudo-TORCH

syndrome. J Exp Med. 2016 Jun 27;213(7):1163–74.

23.  Vuillier F, Li Z, Commere PH, Dynesen LT, Pellegrini S. USP18 and ISG15 coordinately

impact on SKP2 and cell cycle progression. Sci Rep. 2019;9(1):1–11.

24.   Der SD, Zhou A, Williams BR, Silverman RH. Identification of genes differentially

regulated by interferon alpha, beta, or gamma using oligonucleotide arrays. Proc Natl

Acad Sci U S A. 1998 Dec 22;95(26):15623–8.

25.   Liu SY, Sanchez DJ, Aliyari R, Lu S, Cheng G. Systematic identification of type I and

type II interferon-induced antiviral factors. Proc Natl Acad Sci U S A.

2012;109(11):4239–44.

26.   Ivashkiv LB. IFNγ: signalling, epigenetics and roles in immunity, metabolism, disease

and cancer immunotherapy. Nat Rev Immunol. 2018;18(9):545–58.

27.   Didcock L, Young DF, Goodbourn S, Randall RE. The V protein of simian virus 5 inhibits

interferon signalling by targeting STAT1 for proteasome-mediated degradation. J

Virol. 1999 Dec;73(12):9928–33.

28.   Basters A, Geurink PP, Röcker A, Witting KF, Tadayon R, Hess S, et al. Structural basis

of the specificity of USP18 toward ISG15. Nat Publ Gr. 2017;24(3):270–8.

29.   Basters A, Knobeloch KP, Fritz G. How USP18 deals with ISG15-modified proteins:

structural basis for the specificity of the protease. FEBS J. 2018;285(6):1024–9.

30.   Arimoto K, Löchte S, Stoner SA, Burkart C, Zhang Y, Miyauchi S, et al. STAT2 is an

essential adaptor in USP18-mediated suppression of type I interferon signaling. Nat

Publ Gr. 2017;24(3):279–89.

31.   Farrell PJ, Broeze RJ, Lengyel P. Accumulation of an mRNA and protein in interferon-

treated Ehrlich ascites tumour cells. Nature. 1979 Jun;279(5713):523–5.

32.   Löchte S, Waichman S, Beutel O, You C, Piehler J. Live cell micropatterning reveals the

dynamics of signaling complexes at the plasma membrane. J Cell Biol.

2014;207(3):407–18.

33.   Arimoto K-I, Miyauchi S, Stoner SA, Fan J-B, Zhang D-E. Negative regulation of type I

IFN signaling. J Leukoc Biol. 2018 Jun;103(6):1099–116.

34.   Loeb KR, Haas AL. The interferon-inducible 15-kDa ubiquitin homolog conjugates to

intracellular proteins. J Biol Chem. 1992 Apr 15;267(11):7806–13.

35.   Lenschow DJ, Giannakopoulos N V, Gunn LJ, Johnston C, O'Guin AK, Schmidt RE, et al.

Identification of interferon-stimulated gene 15 as an antiviral molecule during Sindbis

virus infection in vivo. J Virol. 2005 Nov 15;79(22):13974–83.

36.   Catic A, Ploegh HL. Ubiquitin--conserved protein or selfish gene? Trends Biochem Sci.

2005 Nov 1;30(11):600–4.

37. Daczkowski CM, Dzimianski J V., Clasman JR, Goodwin O, Mesecar AD, Pegan SD.

Structural Insights into the Interaction of Coronavirus Papain-Like Proteases and

Interferon-Stimulated Gene Product 15 from Different Species. J Mol Biol.

2017;429(11):1661–83.

38. Dzimianski J V., Scholte FEM, Bergeron É, Pegan SD. ISG15: It's Complicated. J Mol

Biol. 2019 Oct 4;431(21):4203–16.

39. van den Elsen PJ. Expression regulation of major histocompatibility complex class I

and class II encoding genes. Front Immunol. 2011;2:48.

40. Rettino A, Clarke NM. Genome-wide Identification of IRF1 Binding Sites Reveals

Extensive Occupancy at Cell Death Associated Genes. J Carcinog Mutagen. 2013;(Spec

Iss Apoptosis).

41. Ma W, Lehner PJ, Cresswell P, Pober JS, Johnson DR. Interferon-γ rapidly increases

peptide transporter (TAP) subunit expression and peptide transport capacity in

endothelial cells. J Biol Chem. 1997;272(26):16585–90.

42. White LC, Wright KL, Felix NJ, Ruffner H, Reis LFL, Pine R, et al. Regulation of LMP2

and TAP1 genes by IRF-1 explains the paucity of CD8+ T cells in IRF-1(-/-) mice.

Immunity. 1996;5(4):365–76.

43. Chatterjee-Kishore M, LWright K, P-YTing J, RStark G. How Stat1 mediates constitutive

gene expression: a complex of unphosphorylated Stat1 and IRF1 supports

transcription of the LMP2 gene. EMBO J. 2000;19(15):4111–22.

44. Demaison C, Parsley K, Brouns G, Scherr M, Battmer K, Kinnon C, et al. High-Level

Transduction and Gene Expression in Hematopoietic Repopulating Cells Using a

Human Imunodeficiency Virus Type 1-Based Lentiviral Vector Containing an Internal

Spleen Focus Forming Virus Promoter. Hum Gene Ther. 2002 May 6;13(7):803–13.

45. He B, Paterson RG, Ward CD, Lamb RA. Recovery of infectious SV5 from cloned DNA

and expression of a foreign gene. Virology. 1997 Oct 27;237(2):249–60.

46. Randall RE, Young DF, Goswami KK, Russell WC. Isolation and characterization of

monoclonal antibodies to simian virus 5 and their use in revealing antigenic

differences between human, canine and simian isolates. J Gen Virol. 1987;68:2769–

80.

47. Nightingale K, Lin KM, Ravenhill BJ, Davies C, Nobre L, Fielding CA, et al. High-

Definition Analysis of Host Protein Stability during Human Cytomegalovirus Infection

Reveals Antiviral Factors and Viral Evasion Mechanisms. Cell Host Microbe.

2018;24(3):447-460.e11.

48.  McAlister GC, Nusinow DP, Jedrychowski MP, Wühr M, Huttlin EL, Erickson BK, et al.

MultiNotch MS3 enables accurate, sensitive, and multiplexed detection of differential

expression across cancer cell line proteomes. Anal Chem. 2014;86(14):7150–8.

49.  Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H, et al. INTERFEROME

v2.0: An updated database of annotated interferon-regulated genes. Nucleic Acids

Res. 2013;41(D1):1040–6.

50.  Vizcaíno JA, Csordas A, Del-Toro N, Dianes JA, Griss J, Lavidas I, et al. 2016 update of

the PRIDE database and its related tools. Nucleic Acids Res. 2016;44(D1):D447–56.

**Figure 1. Functional characterisation of A549-ISG15[-/-] cell lines reconstituted with the C-terminal ISG15 mutants, ISG15.AA and ISG15.ΔGG, or wild type ISG15, ISG15.GG. (A)** Schematic presentation of the lentiviral technology used to reconstitute ISG15 expression in A549-ISG15[-/-] using an inducible system where expression of ISG15 is under the control of its native promoter (pr15). The induction of pr15 is regulated by transcription factors physiologically upregulated by innate immune responses (e.g., IRF3 and ISGF3) or cell-cycle regulators (e.g. p53) **(B)** Immunoblot analysis of ISG15 expression induced by IFN-α treatment. A549 (CNTL), ISG15[-/-] and ISG15.AA-, ΔGG-, GG-expressing derivatives were treated with 1000 IU/ml IFN-α for 24 and 48 h or left untreated. Whole cell extracts were prepared and ISG15, MxA and β-actin proteins were analysed by immunoblot. **(C)** A549 (CNTL), 1SG15[-/-] and ISG15.AA-, ΔGG-, GG-expressing derivatives were treated with 1000 IU/ml IFN-α for 30 min, then extensively washed and media without IFN replaced. Cells were harvested at 0 and 30 min and 24 h after IFN-α removal and phopho-STAT1, total STAT1, MxA, ISG15 and β-actin were detected by immunoblot. Data in lanes 1 – 6 have previously been reported (21) **(D)** A549 (CNTL), ISG15[-/-] and ISG15.AA-, ΔGG-, GG-expressing derivatives were treated with 1000 IU/ml IFN-α for 24 h. Expression of ISGs was tested using reverse transcription quantitative PCR (RT-qPCR) with primers specific for *MxA* and *HERC5*. Relative expression was determined following SYBR Green quantitative PCR (qPCR) using ΔΔCt method. *β-Actin* expression was used to normalize between samples. Data are presented as a mean fold increase relative to IFN-α-treated A549 control cells (set to 1). Error bars represent the SD of the mean from three independent experiments performed on different occasions. Each experiment additionally included three technical replicates. Statistical significance was assessed using two-way ANOVA and Tukey multiple comparisons test; *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$, ****, $p < 0.0001$, n.s., no statistical significance.

**Figure 2. Quantitative temporal analysis of proteomes from IFN-α-treated ISG15[-/-] and reconstituted**

**cells (A)** Schematic presentation of the experimental workflow. **(B)** Abundance of representative

examples of well-known ISGs over time after IFN-α simulation, normalised to a maximum of 1 for each

protein. **(C)** Comparative quantitative analysis of ISG expression in A549 (CNTL), A549-ISG15[-/-] and

ISG15.AA- or ISG15.GG-expressing cells. ISGs were defined by (a) correspondence with the

Interferome database and (b) >1.7-fold increase in abundance in IFN-a stimulated compared to

unstimulated A549 control cells. Statistical significance was assessed using one-way ANOVA and Tukey

multiple comparisons test; ***, p < 0.001, ****, p < 0.0001, n.s., no statistical significance**. (D)**

Pathway analysis using DAVID software. Proteins >1.7-fold upregulated following IFN-α treatment in

each cell type compared to A549 control cells were searched using DAVID software using the default

medium classification stringency. Benjamini-Hochberg adjusted p-values are shown. **(E)** Expression

profiles of representative examples of well-characterised factors with roles in IFN-γ signalling and

antigen processing and presentation pathways. Protein abundance was calculated as described in (B).

**Figure 3. The ability of ISG15 C-terminus to regulate IFN-α signalling influences viral infection (A)**
A549 (CNTL), ISG15[-/-] and ISG15.AA-, ISG15.ΔGG- and ISG15.GG-expressing derivatives were treated
with 1000 IU/ml IFN-α for 18 h and then infected with rPIV5-cherry (MOI 10). Cells were harvested at
24 and 48 h p.i. and processed for immunoblot analysis using antibodies specific for PIV5 NP, STAT1,
MxA, ISG15 and β-actin. **(B)** Experiments described in (A) were performed independently three times
(infections were performed on three separate occasions), and NP and β-actin levels were quantified
using Image Studio software (LI-COR Biosciences). Signals were relative to those generated from IFN-
α-treated A549 cells infected for 48 h p.i. (set to 100%). Error bars = SD. Statistical significance was
assessed using two-way ANOVA and Tukey multiple comparisons test; ****, $p < 0.0001$, n.s., no
statistical significance. **(C)** Fluorescent imagining of mCherry expression, indicative of rPIV5-mCherry
infection, at the 48 hours post infection (h.p.i.) time point described in (A).

**Figure 4. The C-terminal di-Gly motif of ISG15 is not required for USP18 stabilisation**. **(A)** Immunoblot

analysis of USP18 expression in A549 (CNTL), ISG15$^{-/-}$ and ISG15.AA-, ΔGG-, GG-expressing derivatives

after treatment with 1000 IU/ml IFN-α for 24 h. Whole cell extracts were prepared and USP18, ISG15,

MxA and β-actin protein levels were analysed by immunoblot. **(B)** USP18 abundance in A549 (CNTL),

ISG15$^{-/-}$ and ISG15.AA and ISG15.GG-expressing derivative cells measured using quantitative tandem

mass tags (TMT)-based proteomic analysis (see Figure 2).

**Figure 5. The C-terminal di-Gly motif of ISG15 is crucial for the ISG15-USP18 interaction.**
Immunoprecipitation of Myc-tagged ISG15 expressed in ISG15.AA-, ΔGG-, GG-expressing cell lines
after treatment with 1000 IU/ml IFN-α for 24 h.  An ISG15$^{-/-}$ cell-line expressing a non-tagged form of
ISG15 was used as a negative control (left lane) and an ISG15.GG-expressing 1SG15$^{-/-}$.UBA7$^{-/-}$ cell line
was used as an ISGylation-deficient control (final lane). ISG15 was immunoprecipitated (IP) using anti-
c-Myc antibodies covalently coupled to magnetic beads. Immunoprecipitates (top) and whole cell
lysates (WCL; bottom) were subject to immunoblot analysis with antibodies to USP18, ISG15 and
UBA7. ISG15 expression in reconstituted cell lines was under the control of the ISG15 promoter (pr15)
and was therefore inducible by IFN.
893

**Figure 6. Functional characterisation of catalytically inactive and ISG15-binding mutants of USP18**

**(A)** Immunoprecipitation of Flag-tagged wt and mutant forms of USP18 in HEK293T cells co-

transfected with wt ISG15 (ISG15.GG) as indicated. Cells were lysed 48 h post-transfection and lysates

were immunoprecipitated with Flag-specific antibodies covalently coupled to magnetic beads.

Immunoprecipitates (top) and whole cell lysates (WCL; bottom) were subject to immunoblot with

antibodies to USP18 and ISG15. **(B)** Immunoprecipitation of V5-tagged IFNAR2 cytoplasmic domain in

HEK293T cells co-transfected with STAT2-Myc, USP18.wt-Flag, USP18.I60N-Flag or USP18.C64S-Flag

plasmids as indicated. Cells were lysed at 48 h post-transfection and lysates were immunoprecipitated

with anti-V5 antibody coupled to protein G dynabeads. Immunoprecipitates (V5-IP; top) and whole

cell lysates (WCL; bottom) were subject to immunoblot with antibodies to anti-V5 epitope tag,

STAT2and USP18. **(C)** Schematic presentation of the lentiviral technology used to reconstitute USP18

expression in A549-USP18$^{-/-}$ using an inducible system where USP18 expression is driven by the ISG15

promoter (pr15). **(D)** Immunoblot analysis of USP18 expression induced by IFN-α treatment. A549

(CNTL), USP18$^{-/-}$ and USP18.wt-,I60N-,C64S-expressing cell lines were treated with 1000 IU/ml IFN-α

for 48 h or left untreated. Whole cell extracts were prepared and USP18, ISG15, MxA and β-actin

protein levels were analysed by immunoblot.

**Figure 7. ISG expression is dysregulated in cells expressing USP18 mutant unable to bind ISG15. (A)**
A549 control cells expressing NC1 non targeting guide RNA (NC1-CNTL), A549-USP18$^{-/-}$ and USP18.wt,
I60N-, C64S-expressing derivatives (where USP18 expression was under the control of the ISG15
promoter (pr15) and therefore inducible by IFN) were treated with 1000 IU/ml IFN-α for 16 h.
Expression of ISGs was tested using reverse transcription quantitative PCR (RT-qPCR) with primers
specific for *MxA* and *HERC5*. Relative expression was determined following SYBR Green quantitative
PCR (qPCR) using ΔΔCt method. *β-Actin* expression was used to normalize between samples. Data
shown represent mean values from three independent experiments performed on different
occasions; error bars = SD. Statistical significance was assessed using two-way ANOVA and Tukey
multiple comparisons test; *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$, n.s., no statistical significance. **(B)**
A549 NC1-CNTL, A549-USP18$^{-/-}$ and USP18.wt-, I60N-, C64S-expressing derivatives were pre-treated
with 1000 IU/ml IFN-α for 16 h and then infected with rPIV5-mCherry (MOI 10). Cells were harvested
at 24 and 48 h.p.i. and processed for immunoblot analysis using antibodies specific for PIV5 NP, STAT1,
USP18 and β-actin. (**C**) Experiments described in (B) were performed independently three times
(infections were performed on three separate occasions), and NP and β-actin levels were quantified
using Image Studio software (LI-COR Biosciences). Signals were normalised to IFN-α-treated A549 cells
infected for 48 h p.i. (set to 100%). Data shown represent mean values from three independent
experiments; error bars = SD. Statistical significance was assessed using two-way ANOVA and Tukey
multiple comparisons test; ****, $p < 0.0001$, n.s., no statistical significance. **(D)** Fluorescent imagining

933    of mCherry expression, indicative of rPIV5-mCherry infection, at 48h p.i time point of experiment

934    described in (B).

**Figure 8. The ISG15-USP18 interaction is important for desensitization of IFN-α signalling. (A)** A549 control cells expressing NC1 non targeting guide RNA (NC1-CNTL), A549-USP18[-/-] and USP18.wt-, I60N, C64S-expressing cells were primed with 2000 IU/ml IFN-α (equivalent to 20 ng/ml) for 8 h or left untreated. Cells were washed, and re-incubated in medium without IFN for 16 h and then stimulated with 2000 IU/ml IFN-α for 30 min. Cell lysates were subject to immunoblot analysis with antibodies to anti-phospho-STAT1, USP18 and β-actin. **(B)** Experiments described in (A) were performed independently three times, and phospho-STAT1 and β-actin levels were quantified using Image Studio software (LI-COR Biosciences). Signals are presented as ratios of primed to non-primed control (a ratio of 1 is equivalent to no change). Data shown represent mean values from three independent experiments; error bars = SD. Statistical significance was assessed using one-way ANOVA and Tukey multiple comparisons test; *, $p < 0.05$, **, $p < 0.01$, ***, $p < 0.001$, ****, $p < 0.0001$, n.s., no statistical significance.

**Figure 9. Model.** IFN-α, expressed as a consequence of infection, binds to IFNAR and induces the formation of the IFNAR1-IFNAR2-IFN-α ternary complex. Signalling is transduced through the JAK/STAT pathway, culminating in the expression of several hundreds of ISGs. As ISGs, both USP18 and ISG15 are expressed and are both known to regulate the magnitude of type I IFN signalling. ISG15 promotes USP18 stability (1) by preventing SKP2-mediated ubiquitination and proteasomal degradation of USP18, making it available for STAT2-dependent recruitment to IFNAR2 and to inhibit ternary complex formation. How ISG15 protects USP18 from degradation is not known, but this does not require an interaction between ISG5 and USP18. Separately, through non-covalent interactions dependent on its C-terminal di-Gly, ISG15 facilitates USP18's inhibitory function (2). It is not currently known whether this interaction occurs at the receptor, or prior to USP18's recruitment to IFNAR2.