1    The catheterized bladder environment promotes Efg1- and Als1-dependent *Candida albicans*

2                                                     infection

4    Alyssa Ann La Bella[1], Marissa Jeme Andersen[1], Nicholas C. Gervais[2], Jonathan Jesus Molina[1],

5    Alex Molesan[1], Peter V. Stuckey[1], Lauren Wensing[2], Clarissa J. Nobile[3,4], Rebecca S. Shapiro[2],

6                         Felipe Hiram Santiago-Tirado[1]*, & Ana Lidia Flores-Mireles[1]*

8                 [1]Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA

9              [2]Department of Molecular and Cellular Biology, University of Guelph, Guelph, ON, Canada.

10               [3]Department of Molecular and Cell Biology, University of California, Merced, CA, USA

11                  [4]Health Sciences Research Institute, University of California, Merced, CA, USA

12
13               *Address correspondence to Ana Lidia Flores-Mireles (ALFM), afloresm@nd.edu,
14                     and Felipe Hiram Santiago-Tirado (FHST), fsantiago@nd.edu
15
16
17     Mailing address: 351 Galvin Life Science Center, Notre Dame, IN 46556, USA. Phone: (574)
18                         631-4964 (ALFM) and (574) 631-4096 (FHST).
19

22    **Short title:** Catheterized bladder environment induces *Candida* virulence

24    **Teaser:** This is the first study dissecting *C. albicans* pathogenesis during catheter-associated

25    urinary tract infections.

27    **Key words:** *C. albicans*; CAUTI; UTI; Efg1; Als1; filamentation; hyphae; biofilm; urine;

28    bladder; catheter; fungi; fibrinogen-binding protein.

**ABSTRACT**

Catheter-associated urinary tract infections (CAUTIs) account for 40% of all hospital-acquired infections. Given that 20-50% of all hospitalized patients receive a catheter, CAUTIs are one of the most common hospital-acquired infections and a significant medical complication as they result in increased morbidity, mortality, and an estimated annual cost of $340-370 million. *Candida spp.* – specifically *Candida albicans* – are a major causative agent of CAUTIs (17.8%), making it the second most common CAUTI uropathogen. Despite this frequent occurrence, the cellular and molecular details of *C. albicans* infection in the CAUTI microenvironment are poorly understood. Here, we characterize fungal virulence mechanisms and fungal biofilm formation during CAUTI for the first time. We found that the catheterized bladder environment triggers *Candida* virulence programs and robust biofilm formation through Efg1-dependent hyphal morphogenesis and Als1, an Efg1-downstream effector. Additionally, we show that the adhesin Als1 is necessary for *in vitro* and *in vivo C. albicans* biofilm formation dependent on the presence of fibrinogen (Fg), a coagulation factor released in the bladder due to the mechanical damage caused by urinary catheterization. Furthermore, in the presence of Fg, overexpression of *ALS1* in *C. albicans* led to enhanced colonization and dissemination, while deletion of *ALS1* reduced both outcomes during CAUTIs. Our study ultimately unveils the mechanism that contributes to fungal CAUTI, which may provide more effective targets for future therapies to prevent these infections.

## INTRODUCTION

*Candida albicans* is one of few fungal species that can cause serious infection in immunocompromised individuals and immunocompetent people with implanted medical devices (*1*). Normally, *C. albicans* is part of the healthy human microbiota, asymptomatically colonizing the gastrointestinal (GI) tract, reproductive tract, oral cavity, and skin of most humans (*1-3*). However, dysbiosis in the host environment, such as alterations in host immunity, stress, or resident microbiota, promotes *C. albicans* colonization and overgrowth, leading to biofilm formation (*1*).

*C. albicans* produces highly structured biofilms composed of multiple cell morphologies: yeast, pseudohyphae, and hyphae (*4, 5*). Hyphal formation, hyphae-specific gene expression, and biofilm formation are hallmarks of virulence, invasion, and immune evasion, resulting in tissue damage and invasive infection (*6*). *C. albicans* biofilm formation is initiated by the adherence of yeast cells to a surface, forming microcolonies, and is followed by proliferation of hyphae and pseudohyphae (*6*). Hyphal growth is triggered via numerous environmental signals including serum, neutral pH, and $CO_2$, and is controlled by master transcriptional regulators such as Enhanced filamentous growth protein 1 (Efg1) (*7*). Biofilm maturation occurs when the hyphal scaffold is encased in an extracellular polysaccharide matrix (*8*), and the final step involves dispersion of yeast cells to seed planktonic infections or to establish biofilms at new locations in the surrounding environment (*6*).

These robust and highly structured biofilms often develop on implanted devices, allowing *C. albicans* to colonize urinary and central venous catheters, pacemakers, mechanical heart valves, joint prostheses, contact lenses, and dentures (1-3). Biofilm formation on these devices predisposes patients to bloodstream infections and leads to invasive systemic infections of tissues

and organs (*1*). *Candida* biofilms on devices pose a serious threat since they are more resistant to

antifungal therapy and can evade protection from host defenses (*8*). Newer antifungal drug

classes, such as the echinocandins, that target the β-1,3 glucan component of the cell wall have

shown some efficacy against *C. albicans* biofilms (*2, 9*); however, echinocandin-resistant

*Candida spp.* have been recently emerging (*10*).

The 2016 National Healthcare Safety Network review found that *Candida spp.,*

specifically *C. albicans,* has become the second most prevalent uropathogen causing 17.8% of

catheter associated urinary tract infections (CAUTIs) (*11*). CAUTIs are the most common

nosocomial infection with more than 150 million individuals acquiring these infections per year

(*12*). The use of catheters and subsequent infections are not limited to hospital settings. In

nursing homes, 11.9% of residents use indwelling catheters (*13*), and ~50% of catheterized

residents will experience symptomatic CAUTIs (*14*). Despite *Candida spp*. prevalence in

CAUTIs, there are no studies available regarding the pathogenesis of *Candida spp*. during

CAUTI, and the roles of host and fungal factors remains unclear (*11, 15*).

Urinary catheterization causes dysbiosis of the bladder environment by compromising the

urothelium, inducing inflammation, interfering with normal micturition (voiding), and disrupting

host defenses in the bladder (*16-18*). The inflammation response caused by catheterization

exposes epithelial receptors and recruits host factors that can be recognized by the pathogen,

enabling microbial colonization and persistence within the urinary tract (*16-18*). Specifically,

host clotting factor 1, fibrinogen (Fg), is released into the bladder to heal damaged tissues and

prevent bleeding due to catheter-induced inflammation in both mice and humans (*19, 20*). Due to

the constant mechanical damage caused by the urinary catheter, Fg accumulates in the bladder

and deposits onto the catheter, where it is exploited by uropathogens (*19-22*).  In fact, in an *ex*

*vivo* study, we showed that a *C. albicans* CAUTI isolate colocalizes with Fg on urinary catheters

(*20*). This suggests that aspects of the catheterized bladder environment induce changes in *C.*

*albicans* virulence programming, prompting us to study potential fungal factors for CAUTI

pathogenesis.

Here, we characterized the growth, morphology, and biofilm formation in urine, as well

as the ability to cause CAUTIs, of five *C. albicans* clinical and laboratory strains. We found that

urine supported *C. albicans* growth, activated virulence by inducing hyphal morphology, and

enhanced Fg-dependent biofilm formation mediated by the transcriptional regulator Efg1.

Furthermore, we found that catheterized-induced inflammation, specifically Fg, is necessary for

fungal colonization of the urinary tract and further demonstrated that Efg1-dependent hyphal

formation is critical for CAUTI establishment. Importantly, we identified that the Efg1-induced

adhesin, Als1, is important for mediating Fg-fungal interactions *in vitro* and *in vivo* and its

expression is crucial for fungal CAUTIs. This suggests that the development of therapies that

target the Als1-mediated fungal-Fg interactions could potentially be useful in the treatment of *C.*

*albicans* CAUTIs.

**RESULTS**

*C. albicans* **survival and growth in urine promotes hyphal formation.**

*C. albicans* can infect a wide range of regions within the human body including the

oropharyngeal area, gastrointestinal tract, intra-abdominal area, skin, genitals, and urinary tract,

demonstrating its plasticity to survive and replicate in different host environments (*1, 2, 7, 12,*

*23, 24*). Since *C. albicans* is a prominent CAUTI pathogen, we compared growth and

morphology in nutrient-rich and restrictive urine conditions using three urinary clinical isolates

(Pt62, Pt65, and PCNL1 (*20, 25*)) and two laboratory strains (DAY286 and SC5314 (**Table S1**)).

Since hyphal formation is important for promoting disease, we used a hyphal defective double

deletion mutant strain in the SC5314 background, *efg1*Δ/Δ*cph1*Δ/Δ, to test whether hyphal

formation is important in the catheterized bladder environment (**Table S1**). Additionally,

microenvironment conditions induce *C. albicans* morphological changes that are associated with

virulence (*1, 5*). *C. albicans* can exhibit different morphologies (*6*), including yeast,

pseudohyphae, and hyphae (*5*). The hyphal morphology is associated with invasive growth and

increased fungal virulence (*7, 26*). Therefore, we also determined how the bladder environment

affects *C. albicans'* morphology.

For growth analysis, yeast extract peptone dextrose (YPD), a standard *C. albicans*

growth medium, and BHI, the isolation medium for the clinical strains, were considered rich

environments under both static and shaking conditions. Static growth was used to mimic the

bladder environment and shaking growth was used as a comparison with standard lab culture

conditions. For restrictive environments, we used human urine grown statically, supplemented

with 10% human serum, amino acids (AA), bovine serum albumin (BSA), or Fg to mimic

plasma protein extravasation in the catheterized bladder (*21*). Serum albumin and Fg were

chosen because they are two of the most abundant host proteins on catheters retrieved from

humans and mice, and it has been shown that other uropathogens use them as nutrient sources

(*19, 21, 27, 28*). Lastly, we used AA as a general nitrogen source. Samples were taken at 0, 24,

and 48 hours to assess growth by enumeration of colony forming units (CFUs) and morphology

by fluorescent microscopy. As expected, *C. albicans* strains grew in higher densities in rich

media with aeration while growth in static rich media was similar to any urine condition (**Fig. 1**

**and S1**). Growth in restrictive environments varied widely, but all strains were able to grow to

some extent. Human serum promoted growth of all strains by 24 hrs and of most strains by 48

hrs. A subsequent decline was observed by 48 hrs for Pt65 and SC5314, possibly because all

nutrients were consumed rapidly (**Fig. 1A-F**). BSA and AA moderately promoted growth in all

strains except for DAY286 where growth was inhibited when compared to growth in urine alone

(**Fig. 1A, D-F**). Fg enhanced growth of all strains compared to urine alone (**Fig. 1A-F**).

For morphology analysis, YPD and urine, with or without serum, were used as rich and

restrictive environments, respectively. The YPD conditions served as negative and positive

controls for filamentation with the absence or presence of serum, respectively (*29-31*). Cell

morphology analysis was done by quantifying the percentages of yeast, pseudohyphal, or hyphal

forms (**Fig. 1G-L, Table S2**). All strains showed predominantly yeast morphologies in YPD

medium, and YPD with serum induced pseudohyphal and hyphal morphologies in all strains,

except Pt65 and SC5314 *efg1*Δ/Δ*cph1*Δ/Δ. Notably, our analysis showed that urine conditions

promoted pseudohyphal and hyphal formation in all strains (**Fig.1G-K, Table S2**) except

SC5314 *efg1*Δ/Δ*cph1*Δ/Δ (**Fig. 1L, Table S2**). Pseudohyphal and hyphal morphologies were

further induced when urine was supplemented with human serum in all WT lab and clinical

strains. This suggests that the catheterized environment triggers *C. albicans* to transition from the

yeast to hyphal morphology.

**Fibrinogen enhances *C. albicans* biofilm formation.**

During candidiasis, *C. albicans* pseudohyphal and hyphal formation, observed on

catheters from CAUTI patients (*15, 25*) and rats (*32, 33*), induces expression of virulence genes

including adhesion factors (*4, 6, 34, 35*). Furthermore, our *ex vivo* study showed that the *C.*

*albicans* PCNL1 clinical isolate colocalized with Fg, a known binding and biofilm platform for

diverse uropathogens (*19, 20, 27, 36, 37*). Together, this suggests that Fg may be an important

factor promoting *C. albicans* CAUTI pathogenesis. Thus, we hypothesized that urine conditions

induce factors responsible for Fg-binding and biofilm formation.

Biofilm formation was compared in rich (YPD and BHI) and restrictive (human urine)

media as well as between BSA- and Fg-coated microplates (**Fig. 2A-F**). At 48 hrs,

immunostaining was performed to assess fungal biofilm biomass (*38*). We found that Fg

promoted biofilm formation of most strains under all conditions and was further enhanced by

human urine (**Fig. 2A-C**). Fg enhanced biofilm formation in DAY286 in YPD and urine but not

in BHI **(Fig. 2D).** SC5314 showed enhanced biofilm formation with Fg in all conditions **(Fig.**

**2E).** Importantly, SC5314 *efg1*Δ/Δ*cph1*Δ/Δ formed enhanced biofilms in rich media when

coated with Fg but was unable to do so in urine, regardless of the coated surface (**Fig. 2F**). This

suggests that not only is filamentation important for biofilm formation in urine environments, but

that biofilm formation mechanisms differ between urine and rich environments.

Further analysis of biofilm formation was performed on *C. albicans* biofilms grown in

urine on plates coated with either BSA, Fg, or fibrin meshes. Fibrin meshes were added because

in damaged tissue environments like the catheterized bladder, Fg is converted into fibrin fibers or

meshes to stop bleeding and allow tissue healing (*39*). Visual examination of biofilm formation

on BSA-, Fg- or fibrin-coated plates showed that Fg- and fibrin-coated plates supported robust

biofilm formation composed of yeast, pseudohyphae, and hyphae, while on BSA-coated plates,

sparse monolayers or small aggregates of cells were observed (**Fig. 2G-K, M-Q; Fig. S2**).

However, SC5314 *efg1*Δ/Δ*cph1*Δ/Δ showed a small yeast-exclusive monolayer of cells

regardless of BSA, Fg, or fibrin presence (**Fig. 2L and R; Fig. S2**). Together, this suggests

filamentation is important for Fg- and fibrin-dependent biofilm formation and that the regulation

of Fg-binding adhesins expressed during hyphal morphogenesis is controlled by Efg1, Cph1, or

their downstream effectors.

***C. albicans* hyphal formation and Fg interaction are critical for establishment of CAUTI.**

Based on these results and the importance of biofilm formation for infections such as

CAUTIs, we hypothesized that the *C. albicans* hyphal-deficient mutant strain would be unable to

colonize the bladder and catheter. To test this hypothesis, we compared the ability of the clinical

and laboratory strains to colonize the bladder both in the presence and absence of a catheter

using our established mouse model (*19, 21, 27, 28, 36, 37, 40-43*). Mouse bladders were

challenged with one of the strains, and 24 hours post infection (hpi), the mice were euthanized,

and fungal colonization was quantified via CFUs. Our results showed that catheterization

significantly promoted bladder colonization by all the clinical and lab strains compared to the

bladder colonization in non-catheterized mice (**Fig. 3**). Importantly, colonization by the hyphal-

deficient mutant strain, SC5314 *efg1*Δ/Δ*cph1*Δ/Δ, was significantly impaired (**Fig. 3E**), and in

the absence of a catheter, behaved and colonized to the same extent as the WT strain.  Since our

mouse model of CAUTI allows us to assess dissemination, we analyzed the fungal burden of the

kidneys, spleen, and heart after 24 hpi (**Fig. 3**). We found that urinary catheterization

significantly contributed to fungal spread of DAY286 to the kidneys and spleens (**Fig. 3D**) and

spread of SC5314 to the kidneys (**Fig. 3E**). Kidney colonization by Pt62 and Pt65 was 2-3 logs

higher than non-catheterized mice, trending to significance. Importantly, the hyphal-deficient

mutant strain, SC5314 *efg1*Δ/Δ*cph1*Δ/Δ, did not show differential dissemination between

catheterized and non-catheterized mice (**Fig. 3E**).

To further understand *C. albicans* morphogenesis, interaction with Fg, and spatial

colonization in the bladder during CAUTI, we performed histological analyses and

immunofluorescence (IF) microscopy of catheterized bladders at 24 hpi. Bladder colonization

was exceptionally robust and was visible in the H&E-stained whole bladders (blue arrow heads)

(**Fig. 4**). Consistently, our IF analysis showed the presence of *C. albicans* hyphal and

pseudohyphal morphologies in the lumen of the bladder in all clinical and laboratory strains (**Fig.**

**4A-E**), except the hyphal-deficient mutant strain, SC5314 *efg1Δ/Δcph1Δ/Δ* (**Fig. 4F**).

Furthermore, *C. albicans* cells in the catheterized bladder were found associated with Fg (**Fig. 4,**

see **Fig. S3-S8** for zoomed-in images). These data indicate that the changes induced by

catheterization promote fungal colonization of the urinary tract and further demonstrate that

hyphal morphogenesis, as well as Efg1 and Cph1 regulated pathways, are crucial for CAUTI

establishment.

*C. albicans* **interacts with Fg on the catheter surface** *in vivo* **during CAUTI.**

Based on our *in vitro* Fg-binding results (**Fig. 2**), we assessed *C. albicans*-Fg interactions

*in vivo* on the catheter during CAUTI. Catheters from mice infected with each strain were

retrieved 24 hpi, stained, and imaged. Except for the hyphal-deficient mutant strain, we found

that all strains formed a robust biofilm on the implanted catheter (**Fig. 4G-R** merge images; **Fig.**

**S9**, single channels), colonizing 59% to 79% of the surface of the catheter (**Table S3**). SC5314

WT showed 78.9 ± 14% catheter colonization, while SC5314 *efg1Δ/Δcph1Δ/Δ* showed 10.4 ±

7.5% catheter colonization, exhibiting a significant defect in colonization (**Fig. 4L-M, R; Table**

**S3**). Our IF analysis showed that *C. albicans* strains preferentially bound to deposited Fg on the

catheters (**Fig. 4H-M and Fig. S9**). Quantification of the staining found that 75-91% of the *C.*

*albicans* strains were colocalized with deposited Fg (**Fig. 4N-R and Fig. S9**). Although the

hyphal-deficient strain was only able to colonize 10% of the catheter (**Fig. 4M, Table S3**), its

cells were colocalizing with Fg (**Fig. 4R, Table S3**). This result further corroborates the

importance of hyphal morphogenesis for biofilm formation and Fg for catheter colonization

during CAUTI.

**Efg1 is critical for filamentation and for Fg-dependent biofilm formation in urine, and in**

**the development of CAUTI.**

We next explored the independent contributions of Efg1 and Cph1 to hyphal

morphogenesis, Fg-dependent biofilm formation, and colonization of the catheterized bladder

environment (**Fig. 5 and S10**). Using single deletion strains with their corresponding WT, we

found that the *cph1Δ/Δ* strain produced pseudohyphae and hyphae similar to WT while the

*efg1Δ/Δ* strain was predominantly in the yeast-form throughout all conditions (**Fig. S10).** Fg-

dependent biofilm formation was assessed and the *cph1Δ/Δ* strain exhibited a ~40% decrease in

biofilm formation when compared to the WT strains while the *efg1Δ/Δ* strain showed a ~95%

reduction, and the *EFG1* complemented strain restored biofilm formation to WT levels (**Fig.**

**5A**). Furthermore, analysis of fibrin interactions showed that the *cph1Δ/Δ* strain produced a

robust hyphal colonization surrounding the Fg fibers, differing from the *efg1Δ/Δ* strain, which

showed few yeast cells bound directly to the Fg mesh (**Fig. 5B and Fig. S2**). Analysis of

colonization in our mouse model showed that the *efg1Δ/Δ* strain significantly impaired bladder

and catheter colonization while the *EFG1* complemented and *cph1Δ/Δ* strains displayed

comparable colonization to the WT strain (**Fig. 5C**). Based on the role of Efg1 in CAUTI, *EFG1*

expression levels were analyzed in the 24 hpi catheterized and infected bladders with WT,

*efg1Δ/Δ*, and *EFG1* complemented strains. We found that *EFG1* is expressed during bladder

infection with the WT and *EFG1* complemented strains and that *EFG1* transcripts were

undetectable (ND) in the *efg1Δ/Δ* infected bladders as expected (**Fig. 5E**). Furthermore, analysis

of bladder tissue showed that the *cph1Δ/Δ* strain infected bladders had abundant hyphal burdens

similar to the WT strain while the *efg1Δ/Δ* strain infected bladders showed defective colonization

by sporadic yeast cells (**Fig. 5F, S11-S13**). These phenotypes exhibited by the *efg1Δ/Δ* strain

mimicked the phenotypes shown by the *efg1Δ/Δcph1Δ/Δ* strain, indicating that Efg1 is the major

transcriptional regulator responsible for the yeast to hyphal transition and for Fg-dependent

biofilm formation in the catheterized bladder environment.

Neutrophils are highly recruited into the catheterized bladder (*21, 28, 44*) and neutropenic

patients are more susceptible to *C. albicans* and bacterial dissemination originating from CAUTI

(*45-48*). We next analyzed neutrophil recruitment into the catheterized and infected bladder via

IF and flow cytometry (FC). Imaging of bladders showed that neutrophils were highly recruited

into the bladder, specifically in the areas with fungal colonization (**Fig. 4** and **Fig. S3-S8**). We

also observed *C. albicans* cells breaching the urothelium where they encounter a strong

neutrophil response (**Fig. 4C-E** and **Fig. S5-S7**). For example, PCNL1 was able to breach the

bladder lamina propria, inducing massive neutrophil recruitment to contain the infection (**Fig. 4C**

and **Fig. S5**). Pt62 and Pt65 strains were primarily found in the bladder lumen and their cells

were interacting with Fg and neutrophils (**Fig. 4A-B** and **Fig. S5-S6**). On the other hand, robust

fungal colonization and neutrophil recruitment was not observed in the *efg1Δ/Δcph1Δ/Δ* strain or

*efg1Δ/Δ* strain infected bladders (**Fig. 4D, 5F, S8,** and **S13**). Based on this data, using flow

cytometry, we quantified and compared bladder neutrophil recruitment in naïve and catheterized

bladders that were PBS-mock infected or infected with the WT, *efg1Δ/Δcph1Δ/Δ*, *efg1Δ/Δ*, and

*EFG1* complemented strains (**Fig. S14**). We observed that the catheterized bladder by itself

(PBS-mock infected) induced neutrophil recruitment; however, the infected bladder with the WT

strain further increased neutrophil recruitment (**Fig. 5E**), as we have observed via IF (**Fig. 5F**).

Furthermore, neutrophil recruitment in the bladder infected with the *efg1Δ/Δcph1Δ/Δ* and

*efg1Δ/Δ* strains showed levels comparable to the PBS-mock infected condition, while the *EFG1*

complemented strain was not significantly different from the WT strain (**Fig. 5D**). These data

demonstrate that in the catheterized bladder, infection with *C. albicans* leads to the recruitment

of neutrophils to the site of infection, and the virulent hyphal and pseudohyphal morphologies

are driven by Efg1.

**An Efg1-regulated adhesin is critical for biofilm formation and persistence during CAUTI.**

We have demonstrated that *C. albicans* interacts with Fg and its presence enhances

colonization in the catheterized bladder (**Fig. 2-3**), suggesting that urine conditions induce the

expression of Fg-binding adhesins. To identify adhesins important for attachment, colonization,

and persistence during CAUTI, we screened 12 adhesin mutant strains from the Shapiro library

(*49, 50*) (**Fig. 6A**) and 27 adhesin and biofilm-related mutants from the Noble library (*51*) (**Fig.**

**S15A**) for Fg-dependent biofilm formation under urine conditions. We found that of all adhesin

mutant strains, the *als1Δ/Δ* strain exhibited the most severe biofilm formation defect under urine

conditions (**Fig. 6A**), similar to the biofilm defects of the *efg1Δ/Δ* and *efg1Δ/Δcph1Δ/Δ* strains,

without compromising hyphal morphology (**Fig. 6A, 2F and Fig. S16**). Furthermore, the *ALS1*

complemented strain rescued biofilm formation to WT levels (**Fig.6B**; *ALS1 comp*). In addition,

we assessed biofilm formation by an *ALS1* overexpression strain (*ALS1* OE; obtained from the

Shapiro lab (*52*)). This strain not only rescued but enhanced biofilm formation to 8-fold higher

than the WT strain (**Fig. 6B**). Increased biomass was clearly visible in the *ALS1* OE strain and

due to insufficient antibody penetration, crystal violet was used to assess biomass (**Fig. S15B**).

Visualization of the WT, *als1*Δ/Δ, and *ALS1* OE strains interactions with fibrin showed that the

*als1*Δ/Δ strain displayed less colonization of the fibrin meshes while the *ALS1* OE strain

exhibited a more robust colonization than the WT strain (**Fig. 6C**).

We further compared the ability of the WT, *als1*Δ/Δ, *ALS1* complemented, and *ALS1* OE

strains to cause CAUTI. The *als1*Δ/Δ strain displayed a significant defect in bladder and catheter

colonization while the colonization of the *ALS1* complemented strain was similar to that of the

WT; in addition, the *ALS1* OE strain showed a significant increase in bladder, catheter, and

kidney colonization when compared to the WT strain (**Fig. 6D**). Als1 has been reported to be an

Efg1-downstream target (58). To confirm this, *ALS1* expression levels were analyzed in 24 hpi

catheterized and infected bladders with either WT, *efg1*Δ/Δ, and *EFG1* complemented strains.

We found that *ALS1* expression significantly decreased in bladders infected with the *efg1*Δ/Δ

strain compared with bladders infected with the WT and *EFG1* complemented strains (**Fig. 6E**).

Furthermore, imaging of the *als1*Δ/Δ strain infected bladders did not show fungal colonization

(**Fig. 6F**), whereas the *ALS1* OE strain infected bladders had abundant hyphal burdens similar to

the WT strain (**Fig. 6F; Fig. S17-S19**). Imaging of catheters found that the WT strain

colonization covered ~92.4% of the surface, the *als1*Δ/Δ strain covered ~10.73%, and the *ALS1*

OE strain covered ~87.5%. Of that colonization, the fungal population colocalizing with

deposited Fg was ~88.0% for the WT strain, ~93.0 for the *als1*Δ/Δ strain, and ~99.0% for the

*ALS1* OE strain (**Fig. 6G-H and Fig. S20**). The residual binding to Fg in the *als1*Δ/Δ strain was

similar to that of the *efg1*Δ/Δ*cph1*Δ/Δ strain, suggesting that this binding is mediated by other

adhesins. Together, these data show that Als1 is induced by Efg1 and that Als1 is the Fg-binding

protein critical for establishing colonization in the catheterized bladder and that modulation of

Als1 expression affects the outcome of infection.

**DISCUSSION**

*Candida spp*. CAUTIs are increasing, most commonly in prolonged urinary

catheterization and ICU settings (*53, 54*). Our results have shown that urinary catheterization

significantly promotes colonization of the bladder by all strains to $\geq 10^5$ CFUs/ml, meeting the

CDC's standards of UTI criteria (*55*), whereas in the absence of the catheter, colonization by all

strains was significantly lower at $\leq 10^3$ CFUs/ml (**Fig. 3**). It is possible that colonization by the

WT lab strain and the clinical strains is even higher than the CFU enumeration indicates since

hyphal cells are sticky multinucleated cells. Our model indicates that the prevalence of *C.*

*albicans* in the infected bladder is due to increased colonization resulting from changes in the

bladder environment caused by urinary catheterization.

In this study, we also found that Efg1-dependent hyphal morphogenesis and Als1-

mediated interactions with Fg and fibrin are critical for fungal CAUTI establishment (**Fig. 5-6**).

Importantly, Fg enhances *C. albicans* biofilm formation via interactions with the adhesin Als1.

Additionally, urine promotes Efg1 hyphal morphogenesis which is essential for biofilm

formation and infection during CAUTI. *C. albicans* hyphal formation induces strong neutrophil

recruitment in the bladder to control the fungal infection. Thus, the catheterized bladder creates

the ideal environment for *C. albicans* to colonize and persist in the host.

Host proteins have been shown to contribute to fungal biofilm formation (*33*). We

demonstrated that Fg is accumulated in the bladder and deposited on the catheters (*11, 20*) and

that the *C. albicans* PCNL1 clinical isolate colocalized with Fg deposited on urinary catheters

retrieved from a patient with preoperative negative urine culture (*20*). Moreover, a study from

the Andes group showed that Fg, as well as other host proteins, were associated with *C. albicans*

biofilms in urinary catheters retrieved from rats (*8*). Therefore, we focused on understanding the

role of Fg in *C. albicans* CAUTIs. We found that hyphal formation was critical for Fg-dependent

biofilm formation in urine conditions but not in YPD and BHI media, while the hyphal defective

mutant strain was not able to form Fg-dependent biofilms in urine, but could in rich media (**Fig.**

**1 and S1**). This seemingly contradictory result is not necessarily surprising since we have

observed that laboratory growth media does not fully recapitulate conditions found within the

host. For example, several factors critical for bacterial biofilm formation in CAUTI are

dispensable for *in vitro* biofilm formation when using conventional laboratory growth media (*19,*

*28*). This difference could be related to adhesins that are expressed in the yeast cells that may

contribute to biofilm growth in laboratory conditions. Thus, using conditions that closely mimic

the *in vivo* environment is important to identify physiologically relevant determinants for CAUTI

colonization and survival.

To further mimic the bladder environment, we incorporated the presence of host proteins,

BSA and Fg. These proteins are found in the catheterized bladder due to the inflammation

response caused by urinary catheter mechanical damage. Indubitably, we found that Fg enhances

fungal biofilm formation in urine conditions and during urinary catheterization, and that this is

mediated by the adhesin, Als1 (**Fig. 2, 6**). *C. albicans* expresses numerous adhesion proteins,

including the known Fg-binding protein Mp58 (Pra1 in SC5314); however, in urine conditions,

Als1 is the primary adhesin of Fg. We found that modulation of Als1 affected the outcome of

biofilm formation and infection, with the *C. albicans als1Δ/Δ* strain displaying defective Fg-

dependent biofilm formation and *in vivo* colonization. Conversely, the *ALS1* complemented

strain and the *ALS1* overexpression strain rescued Fg-dependent biofilm formation and promoted

CAUTI pathogenesis (**Fig. 6**). Previous structural studies showed that the N-terminal domain of

Als1 interacts with Fg (*56, 57*), binding to the Fg γ-chain via protein-protein interactions, similar

to Clf adhesins in *S. aureus* (*57, 58*). Als1 belongs to the agglutinin-like sequence (Als) protein

family of cell-surface glycoproteins, which are important for adhesion to host, bacteria, and

abiotic surfaces (*56*). Interestingly, Als3 and Als9-2 (an allelic variant of Als9) have been shown

to interact with Fg (*56, 57*); however, they are dispensable in Fg-dependent biofilm formation in

urine (**Fig. 6A**). This may suggest that these factors are not expressed under urine conditions.

Our results indicate that Als1 is a Efg1-downstream target in the catheterized bladder

environment. Studies have shown that Efg1 regulates *ALS3* expression under serum conditions

(*59*) and that Efg1 directly binds to the *ALS3* promoter during biofilm formation in Spider media

(*60*). Efg1 has both overlapping and distinct targets under different environmental conditions

(*59*). Therefore, future studies dissecting the Efg1 regulatory network in the catheterized bladder

will be crucial to understanding *C. albicans'* cellular signaling and to better manage the outcome

of infection.

It has been established that urinary catheterization causes mechanical damage to the

bladder, inducing epithelial wounding and triggering robust inflammation (*21, 61*). Additionally,

clinical studies have detected *C. albicans* hyphae and biofilms on indwelling urinary catheters

retrieved from patients with candiduria (*15, 62, 63*), suggesting fungal pathogenic activity during

urinary catheterization. Here, we demonstrated that urine induces hyphal morphogenesis *in vitro*

and *in vivo* (**Fig. 1 and Table S2**) and that Efg1-dependent hyphal programming is critical for

bladder and catheter colonization (**Fig. 5**). Recently, deletion of *EED1* in *C. albicans* was shown

to result in inhibition of hyphal formation but not virulence; this was attributed to the

overexuberant growth of the *eed1Δ/Δ* strain (*64*). Since we observed hyphal invasive

colonization in the catheterized bladder (**Fig. 4C-E; 5E**), in future experiments, we will explore

whether the *eed1Δ/Δ* strain is able to cause CAUTI, promote bladder tissue invasion,

colonization, and induce a strong neutrophil response.

Furthermore, strong neutrophil recruitment during hyphal colonization (for the WT,

*cph1Δ/Δ*, and *EFG1* complemented strains) in the catheterized bladder was observed whereas

colonization with hyphal deficient strains (*efg1Δ/Δcph1Δ/Δ* and *efg1Δ/Δ*) showed similar

neutrophil levels as the PBS-mock infected bladders (**Fig. 5**). This observation is important since

invasive fungi (in the hyphal form) evoke an active host immune and inflammatory response

while commensal fungi (in the yeast form) promote immune tolerance and "peaceful

coexistence" with the microbiota (*65, 66*). Our results strongly suggest that *C. albicans* has

pathogenic programming triggered by the catheterized bladder environment, resulting in fungal

CAUTI.

During CAUTI, neutrophils are heavily recruited to the bladder as a defense mechanism

against invading pathogens and their numbers increase with extended catheterization (*21, 44*). In

neutropenic mice, the bacterial burden increases 10-fold during CAUTI, suggesting that

neutrophils are controlling and containing the infection in the bladder (*21*). Similarly,

neutropenic patients developed candidemia from candiduria, suggesting that bladder recruited

neutrophils are critical to controlling fungal systemic dissemination (*45*). Neutrophils have

shown to be the major immune cells in the control of candidiasis (*67*) by phagocytizing and

killing yeast cells and short hyphae, while longer hyphae are killed by inducing neutrophil

extracellular traps (NETs), which release DNA, granule enzymes, and antimicrobial peptides

(*68*). Our future studies will be focused on understanding the immune cell strategies against

fungal CAUTIs and their role in containing the fungal infection in the bladder.

*C. albicans* occupies many niches in the human body, and morphological changes are

associated with the establishment of diseased states. This is important in the bladder, since it is

an open and dynamic system, where urine is constantly passing through. Therefore, in order to

establish a successful colonization, adhesion and biofilm formation on the urinary catheter is

essential (*19, 37*). Our results are consistent with that; *Candida* biofilms not only ensure

colonization but protect the growing cells from the hostile environment and potentiate the

establishment of the infection (*69*). This is the first study unveiling the *C. albicans* mechanism of

pathogenesis during CAUTI, finding that Efg1, and specifically its downstream target Als1, are

critical for establishing infection (**Fig. 7**). This study opens new avenues to understanding *C.*

*albicans* cellular programming in the catheterized bladder.  Importantly, the key interaction of

Als1-Fg, an initial step for colonization, could be exploited as a potential therapeutic avenue to

prevent *C. albicans* CAUTIs.

**MATERIALS AND METHODS**

**Ethics statement.** All animal care was consistent with the Guide for the Care and Use of

Laboratory Animals from the National Research Council. The University of Notre Dame

Institutional Animal Care and Use Committee approved all mouse infections and procedures as

part of protocol number 18-08-4792MD. For urine and blood collections, all donors signed an

informed consent form and protocols were approved by the Institutional Review Board of the

University of Notre Dame under study #19-04-5273 for urine and #18-08-4834 for blood.

**Urine Collection.** Human urine from at least two healthy female donors between the ages of 20 - 35 were collected and pooled. Donors did not have a history of kidney disease, diabetes, or recent antibiotic treatment. Urine was sterilized with a 0.22 μm filter (VWR 29186-212) and pH was normalized to 6.0-6.5. BSA (VWR 97061-48); supplemented urine was sterilized again using a 0.22 μm filter. When urine was supplemented with Fg (Enzyme Research Laboratories FIB 3), it was added directly to the sterilized urine and the urine was not sterilized after the addition of Fg.

**Fungal Culture Conditions.** All strains of *C. albicans* were cultured at 37 °C with aeration in 5 mL of YPD (10g/L Yeast Extract (VWR J850-500G), 20g/L Peptone (VWR J636-500G), 20 g/L Dextrose (VWR BDH9230-500G)) broth. For *in vivo* mouse experiments, *C. albicans* strains were grown static overnight in 10 mL of YPD.

**Growth Curves.** Growth curves were performed in glass test tubes (Thermo Fisher Scientific 14-961-29). Overnight cultures (all in stationary phase; measured using a UV/Vis Spectrophotometer) were normalized to ~$1\times10^6$ CFU/ml in 1xPBS (Sigma–Aldrich 1002786391). The culture was then diluted (1:1000) into human urine (supplemented with 1 mg/mL BSA, 1 mg/mL Fg, 50X amino acids, or 1 mg/mL human serum), BHI (incubated statically or shaking), or YPD (incubated statically or shaking) and were incubated in the test tube at 37ºC for 48 hours. At 0, 24, and 48 hours, samples of each condition were taken and analyzed by CFU counts.

**Morphology Assay and Analysis.**   All strains of *C. albicans* were grown in YPD with or

without serum and in human urine with or without serum. At 0, 24, and 48 hours, a sample of

each condition was taken, fixed with 10% formalin, and stained with 100 µg/mL of calcofluor.

Samples were viewed under a Zeiss inverted light microscope (Carl Zeiss, Thornwood, NY) with

the DAPI fluorescent channel. Random images were taken at 100x magnification and processed

with manual counting of yeast, pseudohyphae, and hyphae. Images (consisting of a 3 x 3 tiled

region, i.e. 9 fields of view) were randomly acquired and at least three images were analyzed per

condition. The total number of cells per phenotype were summed and divided by the total

number of cells to give the overall percentage of each cell type.

**Antibodies and dyes used in this study**.

**Primary antibodies**: Goat anti-fibrinogen (Sigma-Aldrich F8512), rabbit anti-*Candida*

(ThermoFisher Scientific PA1-27158), and rat anti-mouse Ly6G (BD Pharmingen 551459).

**Secondary antibodies**: Alexaflour 488-labeled donkey anti-goat (ThermoFisher Scientific SA5-

10086); Alexaflour 594-labeled donkey anti-rabbit (ThermoFisher Scientific SA5-10039);

Alexaflour 647-labeled donkey anti-rat (ThermoFisher Scientific SA5-10029); IRDye 800CW

donkey anti-goat; and IRDye 680LT donkey anti-rabbit. Alexaflour secondary antibodies were

purchased from Invitrogen Molecular Probes and IRDye conjugates secondary antibodies from

LI-COR Biosciences. **Dyes:** Calcofluor white (Fluorescent Brightener 28, Sigma F3543)

staining; Hoechst dye (Thermo Fisher Scientific 62249) staining; Hematoxylin and Eosin (H&E)

(vector Laboratories #H-3502).

**Biofilm formation assays.** Biofilm formation assays were performed in 96 well flat-bottomed plates (VWR 10861-562) coated with 100uL of BSA or Fg (150 µg/mL) incubated overnight at 4°C. The various strains were grown as described above and the inoculum normalized to ~1x10$^6$ CFUs/ml. Cultures were then diluted (1:1000) into YPD, BHI, or human urine. 100 µL of the inoculum were incubated in the wells of the 96 well plate at 37ºC for 48 hours while static.

Following the 48 hr incubation, the supernatant was removed from the plate and washed three times with 200 µL 1x PBS to remove unbound fungi. Plates were fixed with 10% neutralizing formalin (Leica 3800600) for 20 minutes and followed by three washes with PBS containing 0.05% Tween-20 (PBS-T). Blocking solution (PBS with 1.5% BSA and 0.1% sodium azide (Acros Organics 447811000)) was added to the plate for one hour at room temperature and then washed with PBS-T (3x). Biofilms were incubated with anti-*Candida* antibodies diluted into dilution buffer (PBS with 0.05% Tween (VWR M147-1L), 0.1% BSA) for two hours. Plates were washed three times with PBS-T and incubated for one hour with IRDye 680 LT donkey anti-rabbit secondary antibody solution at room temperature and washed with PBS-T (3x). As a final step, the biofilms were visualized by scanning the plates using the Odyssey Imaging System (LI-COR Biosciences) and the analyzed with Image Studio software to obtain the fluorescence intensities (LI-COR Version 5.2, Lincoln, NE).

**BSA or Fg-coated dishes and formation of Fibrin fibers/nets.** For these assays No. 0 cover glass glass-bottom 35 mm petri dish with a 14 mm microwell (MatTek P35G-0-14-C) were used. The dishes were coated with 150 µg/mL of BSA or Fg overnight at 4°C.  For fibrin fiber/nets formation, Fg and thrombin (Sigma-Aldrich T6884-250UN) were thawed at 37°C. 100 µl of 0.5 mg/ml Fg in PBS was added into the microwell glass-bottom and then 10 µl of 2 U/ml thrombin

was added to polymerize Fg into fibrin. Dishes were incubated at 37°C for 1 hour and kept

overnight at 4°C

**Visualization of biofilms and fungal-fibrin interactions.** The various strains were grown as

described above and the inoculum normalized to ~$1 \times 10^6$ CFUs/ml in PBS. These cultures were

then diluted (1:1000) into human urine, added to the BSA-, Fg-, or fibrin coated dishes and then

were incubated at 37ºC for 48 hours under static conditions. After incubation, dishes were then

washed three times with 1x PBS to remove unbound fungi, then dishes were fixed with 10%

neutralizing formalin solution for 20 minutes and washed with 1x PBS three times. Dishes were

blocked with blocking solution for an hour at room temperature as described above. Then BSA-

and Fg-coated dishes were incubated in primary antibody (rabbit anti-*Candida*) and for fibrin-

coated dishes were incubated with rabbit anti-*Candida* and goat anti-Fg antibodies. Incubation

with the primary antibodies was done for two hours followed by three washes with PBS-T. Then,

dishes were incubated for 1 hour with Alexaflour 594-labeled donkey anti-rabbit secondary

antibody for BSA- and Fg-coated dishes and Alexaflour 594-labeled donkey anti-rabbit and

Alexaflour 488-labeled donkey anti-goat antibodies for fibrin-coated dishes, followed by three

washes with PBS-T. BSA- and Fg- coated dishes were visualized with a Zeiss inverted light

microscope, and images were taken at different magnifications (10x, 20x, 40x and 100x). Zen

Pro and Fiji-ImageJ (*70*) softwares were used to analyze the images. For the fungal-fibrin

interaction, fibrin-coated dishes were visualized by Nikon A1-R/Multi-Photon Laser Scanning

Confocal Microscope and images were analyzed by IMARIS Image Analysis software and

ImageJ software (*70*).

**Crystal violet staining.** Following biofilm formation on Fg-coated microplate, the supernatant

was removed, and the plate was incubated in 200 µl of 0.5% crystal violet for 15 minutes.

Crystal violet stain was removed, and the plate was washed with water to remove remaining

stain. Plates were dried and then incubated with 200 µl of 33% acetic acid for 15 minutes. In a

new plate, 100 µl of the acetic acid solution was transferred, and absorbance values were

measured via a plate spectrophotometer at 595 nm.

*In vivo mouse* **model.** Mice used in this study were ~6-week-old female wild-type C57BL/6

mice purchased from Jackson Laboratory. Mice were subjected to transurethral implantation of a

silicone catheter and inoculated as previously described (*42*). Briefly, mice were anesthetized by

inhalation of isoflurane and implanted with a 6-mm-long silicone catheter (BrainTree Scientific

SIL 025). Mice were infected immediately following catheter implantation with 50 µl of

$\sim 1 \times 10^6$ CFUs/mL in PBS, of one of the fungal strains introduced into the bladder lumen by

transurethral inoculation. Mice were sacrificed at 24 hours post infection by cervical dislocation

after anesthesia inhalation and catheter, bladder, kidneys, spleen, and heart were aseptically

harvested. Organs were homogenized and catheters were cut in small pieces before sonication for

fungal CFU enumeration. A subset of catheters were fixed for imaging as described below and a

subset of bladders were fixed and processed for immunofluorescence and histology analysis as

described below.

**Catheter imaging and analysis.** Harvested catheters were fixed for imaging via standard IF

procedure as previously described (*41*). Briefly, catheters were fixed with formalin, blocked,

washed with 1x PBS, and incubated with the appropriate primary antibodies overnight. Catheters

were then incubated with secondary antibodies for 2 hrs at room temperature. Catheters were

washed with PBS-T and then a final wash with PBS. Catheters were visualized with the Odyssey

Imaging System and then analyzed using color pixel counter from Fiji-ImageJ software (*70*). The

number of pixels of each color was compared to the total number of pixels to identify percent

coverage of the catheter.

**IHC and H&E staining of mouse bladders.** Mouse bladders were fixed in 10% formalin

overnight, before being processed for sectioning and staining as previously described (*37*).

Briefly, bladder sections were deparaffinized, rehydrated, and rinsed with water. Antigen

retrieval was accomplished by boiling the samples in Na-citrate, washing in tap water, and then

incubating in 1x PBS three times. Sections were then blocked (1% BSA, 0.3% TritonX100

(Acros Organics 21568-2500) in 1x PBS) washed in 1x PBS, and incubated with appropriate

primary antibodies diluted in blocking buffer overnight at 4 °C. Next, sections were washed with

1x PBS, incubated with secondary antibodies for 2 hrs at room temperature, and washed once

more in 1x PBS prior to Hoechst dye staining. Secondary antibodies for immunohistochemistry

were Alexa 488 donkey anti-goat, Alexa 550 donkey anti-rabbit, and Alexa 650 donkey anti-rat.

Hematoxylin and Eosin (H&E) stain for light microscopy was done by the CORE facilities at the

University of Notre Dame (ND CORE). All imaging was done using a Zeiss inverted light

microscope and a Nikon A1-R/Multi-Photon Laser Scanning Confocal Microscope. Zen Pro,

ImageJ software, and IMARIS Image Analysis software were used to analyze the images.

**Measurement of *EFG1* and *ALS1* expression levels during *C. albicans* CAUTI.**  Total RNA

was extracted from infected and catheterized mouse bladders with the Qiagen RNeasy Kit (Cat.

No. 74004) and subsequently DNaseI treated. RNA was reverse transcribed into cDNA via

qScript cDNA synthesis kit (QuantaBio, Cat. No.101414) and incubated for 5 minutes at 25°C,

30 minutes at 42°C, 5 minutes at 85°C, and held at 4°C. qPCR was performed on QuantStudio 3

(Thermo Fisher Scientific) under the following conditions: $95°C^{5min}$ ( $95^{15sec}$, $60^{60sec}$)$_{35}$

$_{cycles}$ ($95^{15sec}$, $60^{60sec}$, $95^{15sec}$)$_{melt\ curve}$. Data were normalized using *ITS2* as an internal

housekeeping control gene (*71*) and WT as a sample calibrator and set the expression to 100%.

Data were analyzed via the ΔΔCT method. Primers are described in **Supplementary Table 4**.

**Neutrophil analysis.** Mouse bladders were harvested and minced into digestion buffer (34U/mL

Liberase and 100 µg/mL DNase1 in D-PBS). Bladder were incubated for 1 hours in a heat block

shaker at 37°C and 250rpm and every 15 min were vortexed in high speed for 30s. Digestion was

arrested by addition of D-PBS supplemented with 2% FBS and 0.2 µM EDTA and passed

through 40-µm cell strainers. Following a 20-minute incubation in Fc block (BD Biosciences),

cells were immunolabeled with the antibodies listed in **Supplementary Table 5**. Total cell

counts in the bladder were determined on a BD LSRFortessa X-20 system cytometer (using

DIVA software and analyzed by FlowJo software).

**Statistical analysis.** Data from at least 3 experiments were pooled for each assay. Two-tailed

Mann-Whitney *U* tests were performed with GraphPad Prism 5 software (GraphPad Software,

San Diego, CA) for all comparisons described in biofilm, CAUTI, and catheter coverage

experiments. Values represent means ± SEM derived from at least 3 independent experiments. *,

*P*<0.05; **, *P*<0.005; ***, *P*<0.0005; ns, difference not significant.

**REFERENCES**

1.     C. J. Nobile, A. D. Johnson, Candida albicans Biofilms and Human Disease. *Annu Rev Microbiol*
**69**, 71-92 (2015).
2.     C. Tsui, E. F. Kong, M. A. Jabra-Rizk, Pathogenesis of Candida albicans biofilm. *Pathog Dis* **74**,
ftw018 (2016).

3.   S. Duggan, I. Leonhardt, K. Hunniger, O. Kurzai, Host response to Candida albicans bloodstream
infection and sepsis. *Virulence* **6**, 316-326 (2015).
4.   F. L. Mayer, D. Wilson, B. Hube, Candida albicans pathogenicity mechanisms. *Virulence* **4**, 119-
128 (2013).
5.   D. S. Thompson, P. L. Carlisle, D. Kadosh, Coevolution of morphology and virulence in Candida
species. *Eukaryot Cell* **10**, 1173-1182 (2011).
6.   S. M. Noble, B. A. Gianetti, J. N. Witchley, Candida albicans cell-type switching and functional
plasticity in the mammalian host. *Nat Rev Microbiol* **15**, 96-108 (2017).
7.   P. E. Sudbery, Growth of Candida albicans hyphae. *Nat Rev Microbiol* **9**, 737-748 (2011).
8.   J. E. Nett, D. R. Andes, Contributions of the Biofilm Matrix to Candida Pathogenesis. *J Fungi
(Basel)* **6**,  (2020).
9.   S. P. Bachmann *et al.*, In vitro activity of caspofungin against Candida albicans biofilms.
*Antimicrob Agents Chemother* **46**, 3591-3596 (2002).
10.  K. E. Pristov, M. A. Ghannoum, Resistance of Candida to azoles and echinocandins worldwide.
*Clinical Microbiology and Infection* **25**, 792-798 (2019).
11.  A. Flores-Mireles, T. N. Hreha, D. A. Hunstad, Pathophysiology, Treatment, and Prevention of
Catheter-Associated Urinary Tract Infection. *Top Spinal Cord Inj Rehabil* **25**, 228-240 (2019).
12.  D. G. Maki, P. A. Tambyah, Engineering out the risk for infection with urinary catheters. *Emerg
Infect Dis* **7**, 342-347 (2001).
13.  M. A. Rogers *et al.*, Use of urinary collection devices in skilled nursing facilities in five states. *J
Am Geriatr Soc* **56**, 854-861 (2008).
14.  A. Montoya, L. Mody, Common infections in nursing homes: a review of current issues and
challenges. *Aging health* **7**, 889-899 (2011).
15.  J. F. Fisher, K. Kavanagh, J. D. Sobel, C. A. Kauffman, C. A. Newman, Candida urinary tract
infection: pathogenesis. *Clin Infect Dis* **52 Suppl 6**, S437-451 (2011).
16.  R. A. Garibaldi, J. P. Burke, M. L. Dickman, C. B. Smith, Factors predisposing to bacteriuria during
indwelling urethral catheterization. *The New England journal of medicine* **291**, 215-219 (1974).
17.  C. W. Norden, G. M. Green, E. H. Kass, Antibacterial mechanisms of the urinary bladder. *The
Journal of clinical investigation* **47**, 2689-2700 (1968).
18.  J. W. Warren, Catheter-associated urinary tract infections. *Infectious disease clinics of North
America* **11**, 609-622 (1997).
19.  A. L. Flores-Mireles, J. S. Pinkner, M. G. Caparon, S. J. Hultgren, EbpA vaccine antibodies block
binding of Enterococcus faecalis to fibrinogen to prevent catheter-associated bladder infection
in mice. *Sci Transl Med* **6**, 254ra127 (2014).
20.  A. L. Flores-Mireles *et al.*, Fibrinogen Release and Deposition on Urinary Catheters Placed during
Urological Procedures. *J Urol* **196**, 416-421 (2016).
21.  P. S. Guiton, T. J. Hannan, B. Ford, M. G. Caparon, S. J. Hultgren, Enterococcus faecalis
overcomes foreign body-mediated inflammation to establish urinary tract infections. *Infect
Immun* **81**, 329-339 (2013).
22.  M. J. Andersen *et al.*, Inhibiting host-protein deposition on urinary catheters reduces associated
urinary tract infections. *Elife* **11**,  (2022).
23.  R. Alves *et al.*, Adapting to survive: How Candida overcomes host-imposed constraints during
human colonization. *Plos Pathog* **16**,  (2020).
24.  A. L. Flores-Mireles, J. N. Walker, M. Caparon, S. J. Hultgren, Urinary tract infections:
epidemiology, mechanisms of infection and treatment options. *Nat Rev Microbiol* **13**, 269-284
(2015).
25.  J. N. Walker *et al.*, High-resolution imaging reveals microbial biofilms on patient urinary
catheters despite antibiotic administration. *World J Urol* **38**, 2237-2245 (2020).

26. R. Pukkila-Worley, A. Y. Peleg, E. Tampakakis, E. Mylonakis, Candida albicans Hyphal Formation
and Virulence Assessed Using a Caenorhabditis elegans Infection Model. *Eukaryotic Cell* **8**, 1750-
1758 (2009).
27. A. L. Flores-Mireles *et al.*, Antibody-Based Therapy for Enterococcal Catheter-Associated Urinary
Tract Infections. *Mbio* **7**, (2016).
28. P. S. Guiton, C. S. Hung, L. E. Hancock, M. G. Caparon, S. J. Hultgren, Enterococcal biofilm
formation and virulence in an optimized murine model of foreign body-associated urinary tract
infections. *Infect Immun* **78**, 4166-4175 (2010).
29. H. Bar-Yosef, N. Vivanco Gonzalez, S. Ben-Aroya, S. J. Kron, D. Kornitzer, Chemical inhibitors of
Candida albicans hyphal morphogenesis target endocytosis. *Sci Rep* **7**, 5692 (2017).
30. H. Watanabe, M. Azuma, K. Igarashi, H. Ooshima, Relationship between cell morphology and
intracellular potassium concentration in Candida albicans. *J Antibiot* **59**, 281-287 (2006).
31. Q. H. Feng, E. Summers, B. Guo, G. Fink, Ras signaling is required for serum-induced hyphal
differentiation in Candida albicans. *Journal of Bacteriology* **181**, 6339-6346 (1999).
32. J. E. Nett *et al.*, Rat indwelling urinary catheter model of Candida albicans biofilm infection.
*Infect Immun* **82**, 4931-4940 (2014).
33. J. E. Nett *et al.*, Host contributions to construction of three device-associated Candida albicans
biofilms. *Infect Immun* **83**, 4630-4638 (2015).
34. M. Schaller, C. Borelli, H. C. Korting, B. Hube, Hydrolytic enzymes as virulence factors of Candida
albicans. *Mycoses* **48**, 365-377 (2005).
35. P. L. Carlisle *et al.*, Expression levels of a filament-specific transcriptional regulator are sufficient
to determine Candida albicans morphology and virulence. *P Natl Acad Sci USA* **106**, 599-604
(2009).
36. J. R. Gaston *et al.*, Enterococcus faecalis Polymicrobial Interactions Facilitate Biofilm Formation,
Antibiotic Recalcitrance, and Persistent Colonization of the Catheterized Urinary Tract.
*Pathogens* **9**, (2020).
37. J. N. Walker *et al.*, Catheterization alters bladder ecology to potentiate Staphylococcus aureus
infection of the urinary tract. *P Natl Acad Sci USA* **114**, E8721-E8730 (2017).
38. C. Colomer-Winter, J. A. Lemos, A. L. Flores-Mireles, Biofilm Assays on Fibrinogen-coated
Silicone Catheters and 96-well Polystyrene Plates. *Bio Protoc* **9**, (2019).
39. J. W. Weisel, R. I. Litvinov, Fibrin Formation, Structure and Properties. *Subcell Biochem* **82**, 405-
456 (2017).
40. C. Colomer-Winter *et al.*, Manganese acquisition is essential for virulence of Enterococcus
faecalis. *Plos Pathog* **14**, e1007102 (2018).
41. C. Colomer-Winter, A. L. Flores-Mireles, S. Kundra, S. J. Hultgren, J. A. Lemos, (p)ppGpp and
CodY Promote Enterococcus faecalis Virulence in a Murine Model of Catheter-Associated
Urinary Tract Infection. *mSphere* **4**, (2019).
42. M. S. Conover, A. L. Flores-Mireles, M. E. Hibbing, K. Dodson, S. J. Hultgren, Establishment and
Characterization of UTI and CAUTI in a Mouse Model. *J Vis Exp*, e52892 (2015).
43. G. Di Venanzio *et al.*, Urinary tract colonization is enhanced by a plasmid that regulates
uropathogenic Acinetobacter baumannii chromosomal genes. *Nat Commun* **10**, (2019).
44. M. Rousseau *et al.*, Bladder catheterization increases susceptibility to infection that can be
prevented by prophylactic antibiotic treatment. *Jci Insight* **1**, (2016).
45. P. G. Pappas *et al.*, Clinical Practice Guideline for the Management of Candidiasis: 2016 Update
by the Infectious Diseases Society of America. *Clin Infect Dis* **62**, e1-50 (2016).
46. L. J. Conway, E. J. Carter, E. L. Larson, Risk Factors for Nosocomial Bacteremia Secondary to
Urinary Catheter-Associated Bacteriuria: A Systematic Review. *Urol Nurs* **35**, 191-203 (2015).

47.  M. T. Greene *et al.*, Predictors of hospital-acquired urinary tract-related bloodstream infection.
*Infect Control Hosp Epidemiol* **33**, 1001-1007 (2012).
48.  L. E. Nicolle, Catheter associated urinary tract infections. *Antimicrob Resist Infect Control* **3**, 23
(2014).
49.  S. Rosiana *et al.*, Comprehensive genetic analysis of adhesin proteins and their role in virulence
of Candida albicans. *Genetics* **217**,  (2021).
50.  R. S. Shapiro *et al.*, A CRISPR-Cas9-based gene drive platform for genetic interaction analysis in
Candida albicans. *Nat Microbiol* **3**, 73-82 (2018).
51.  S. M. Noble, S. French, L. A. Kohn, V. Chen, A. D. Johnson, Systematic screens of a Candida
albicans homozygous deletion library decouple morphogenetic switching and pathogenicity. *Nat*
*Genet* **42**, 590-598 (2010).
52.  N. C. Gervais *et al.*, Development and applications of a CRISPR activation system for facile
genetic overexpression in <em>Candida albicans</em>. *bioRxiv*, 2022.2008.2015.501889 (2022).
53.  M. S. Rishpana, J. S. Kabbin, Candiduria in Catheter Associated Urinary Tract Infection with
Special Reference to Biofilm Production. *J Clin Diagn Res* **9**, DC11-13 (2015).
54.  D. Keten *et al.*, Catheter-associated urinary tract infections in intensive care units at a university
hospital in Turkey. *Bosn J Basic Med Sci* **14**, 227-233 (2014).
55.  N. H. S. Network. (Center for Disease Control, 2021).
56.  L. L. Hoyer, E. Cota, Candida albicans Agglutinin-Like Sequence (Als) Family Vignettes: A Review
of Als Protein Structure and Function. *Front Microbiol* **7**, 280 (2016).
57.  P. S. Salgado *et al.*, Structural basis for the broad specificity to host-cell ligands by the
pathogenic fungus Candida albicans. *Proc Natl Acad Sci U S A* **108**, 15775-15779 (2011).
58.  J. Lin *et al.*, The peptide-binding cavity is essential for Als3-mediated adhesion of Candida
albicans to human cells. *J Biol Chem* **289**, 18401-18412 (2014).
59.  V. E. Glazier, EFG1, Everyone's Favorite Gene in Candida albicans: A Comprehensive Literature
Review. *Frontiers in Cellular and Infection Microbiology* **12**,  (2022).
60.  C. J. Nobile *et al.*, A recently evolved transcriptional network controls biofilm development in
Candida albicans. *Cell* **148**, 126-138 (2012).
61.  B. E. Glahn, O. Braendstrup, H. P. Olesen, Influence of drainage conditions on mucosal bladder
damage by indwelling catheters. II. Histological study. *Scand J Urol Nephrol* **22**, 93-99 (1988).
62.  E. M. Kojic, R. O. Darouiche, Candida infections of medical devices. *Clin Microbiol Rev* **17**, 255-
267 (2004).
63.  M. Azmy, N. Nawar, M. Mohiedden, L. Warille, Electron Microscopic Assay of Bacterial Biofilm
Formed on Indwelling Urethral Catheters. *Journal of the Egyptian Society of Parasitology* **46**,
475-484 (2016).
64.  C. Dunker *et al.*, Rapid proliferation due to better metabolic adaptation results in full virulence
of a filament-deficient Candida albicans strain. *Nat Commun* **12**, 3899 (2021).
65.  M. L. Wheeler, J. J. Limon, D. M. Underhill, Immunity to Commensal Fungi: Detente and Disease.
*Annu Rev Pathol* **12**, 359-385 (2017).
66.  I. D. Iliev, D. M. Underhill, Striking a balance: fungal commensalism versus pathogenesis. *Curr*
*Opin Microbiol* **16**, 366-373 (2013).
67.  J. V. Desai, M. S. Lionakis, The role of neutrophils in host defense against invasive fungal
infections. *Curr Clin Microbiol Rep* **5**, 181-189 (2018).
68.  S. Kenno, S. Perito, P. Mosci, A. Vecchiarelli, C. Monari, Autophagy and Reactive Oxygen Species
Are Involved in Neutrophil Extracellular Traps Release Induced by C. albicans Morphotypes.
*Front Microbiol* **7**, 879 (2016).
69.  M. Cavalheiro, M. C. Teixeira, Candida Biofilms: Threats, Challenges, and Promising Strategies.
*Front Med (Lausanne)* **5**, 28 (2018).

70.  J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**,
676-682 (2012).
71.  C. Y. Turenne, S. E. Sanche, D. J. Hoban, J. A. Karlowsky, A. M. Kabani, Rapid identification of
fungi by using the ITS2 genetic region and an automated fluorescent capillary electrophoresis
system. *J Clin Microbiol* **37**, 1846-1851 (1999).

**Acknowledgments:** We thank members of the A.L.F.M. and F.H.S.T. laboratories for their

helpful suggestions and insightful comments. Special thank you to Dr. Michael G. Caparon for

his comments. Special thank you to Dr. Sara Cole and the ND Integrated Imaging Facility for

tissue processing and support during imaging.   **Funding:** This work was supported by

institutional funds from the University of Notre Dame (to A.L.F.M. and F.H.S.T), National

Institutes of Health grant R01DK128805 (to A.L.F.M., and M.J.A.), Arthur J. Schmitt

Leadership Fellowship (A.A.L.), CIFAR Azrieli Global Scholar Award and Canadian Institutes

of Health Research (CIHR) Project Grant PJT162195 (to R.S.S.), and National Institutes of

Health grant R35GM124594 (to C.J.N.). **Author contributions** A.L.F.M., and F.H.S.T.

designed the experiments. A.A.L., M.J.A., A.M., J.J.M., and P.S. performed the studies. R.S.S.,

N.C.G., and L.W. designed and generated the adhesin mutant strain collection and the *ALS1* OE

strain. C.J.N.  generated the *efg1Δ/Δ, EFG1* complemented, *als1Δ/Δ*, and *ALS1* complemented

strains. A.A.L., A.L.F.M, and F.H.S.T. wrote the paper. All authors contributed to editing the

paper. **Competing interests:** C.J.N. is a cofounder of BioSynesis, Inc., a company developing

diagnostics and therapeutics for biofilm infections. All other authors declare no competing

financial interests.

 **Figures:**

 **Figure 1.** ***C. albicans* grows, survives, and transitions to the hyphal morphology when**

 **grown in urine.** **(A-F)** Growth curves of *C. albicans* strains Pt62 (A), Pt65 (B), PCNL1 (C),

 DAY230 (D), SC5314 (E), and SC5314 *efg1Δ/Δcph1Δ/Δ* (F), grown in human urine conditions

 alone, or urine conditions supplemented with 10% human serum (serum), fibrinogen (Fg),

 bovine serum albumin (BSA), or amino acids (AA). Fungal growth was determined by CFUs

 enumeration after 0, 24, and 48 hours. Except when indicated, all strains were grown under static

 conditions. Data presented shows the mean and standard error of the mean derived from three

 independent experiments with at least three technical replicates. **(G-M)** The morphology of *C.*

 *albicans* strains were evaluated after 0, 24, and 48 hours of growth in urine and YPD with or

 without 3% human serum. Images (consisting of a 3 x 3 tiled region, i.e. 9 fields of view) were

 randomly acquired and at least three images were analyzed per condition. The total number of

 cells per phenotype were summed and divided by the total number of cells to give the overall

 percentage of yeast, pseudohyphae, and hyphae (**M**). Differences between groups were tested for

 significance using the Mann-Whitney U test.  *, $P < 0.05$; **, $P < 0.005$; ***, $P < 0.0005$; and

 ****, $P < 0.0001$. P-value colors represent the comparison between the different populations:

 hyphae (red), pseudohyphae (orange), and yeast (black).

**Figure 2. Fibrinogen/fibrin interaction with *C. albicans* enhances biofilm formation.** (**A-F**) Immunostaining analysis of biofilm formation on BSA- or Fg- coated microplates by *C. albicans* strains when grown in YPD, BHI, or human urine. At 48 hrs, *C. albicans* biofilm formation was measured via fluorescence intensity using anti-*Candida* antibodies. Data presented shows the mean and standard error derived from three independent experiments with 24 technical replicates. Differences between groups were tested for significance using the Mann-Whitney U test. ****, P<0.0001; ns, not statistically different. (**G-L**) Microscopic visualization of 48 hrs *C. albicans* biofilms biomass on BSA- or Fg-coated glass bottom petri dishes grown in urine using anti-*Candida* antibodies. White squares represent the zoom-in area used for the higher magnification (x). (**M-R**) Microscopic visualization and 3D reconstruction of 48 hrs *C. albicans* biofilms on fibrin fibers/meshes grown in human urine using antibodies against Fg (anti-Fg; green) and *C. albicans* (anti-*Candida*; red). Scale bars: 50 µm.

**Figure 3. Urinary catheterization promotes fungal colonization and hyphal formation is required for CAUTI.** (**A-D**) Fungal burden of clinical and lab strains on the harvested organs and catheter. (**E**) Fungal burden comparison between SC5314 WT and SC5314 *efg1Δ/Δcph1Δ/Δ* strains. Values represent means ± SEM. The Mann-Whitney U test was used; *, P < 0.05 was considered statistically significant. **, P < 0.005; ns, values were not statistically significantly different. The horizontal bar represents the median value. The horizontal broken line represents the limit of detection of viable fungi. LOD; limit of detection. Animals that lost the catheter were not included in this work.

**Figure 4. Hyphal *C. albicans* cells invade the lumen of the catheterized bladder and interact**

**with fibrinogen on the bladders and catheters.** (**A-F**) Implanted and infected bladders and

catheters were recovered at 24 hpi. Bladders were subjected to analysis by H&E and IF staining.

For the IF analysis antibody staining was used to detect Fg (anti-Fg; green), *C. albicans* (anti–

*Candida*; red), and neutrophils (anti-Ly6G; white). Staining with DAPI (blue) delineated the

urothelium and cell nuclei (representative images). The white broken line separates the bladder

lumen (L) from the urothelium surface (U), the lamina propria (LP), and muscularis (M). H&E

stained bladder scale bars, 700 µm. White squares represent zoomed-in areas of higher

magnification (20x, 40x, 100x).    Blue arrow heads indicate *C. albicans* colonization. (**G-M**)

Implanted catheters were stained with antibodies to detect Fg (anti-Fg; green) and *C. albicans*

(anti-Candida; red). (**N-R**) Quantification of fungal colocalization with deposited Fg on the

catheter. The Mann-Whitney U test was used to analyze catheter colonization between SC5314

WT and hyphal mutant; *, P < 0.05; Values represent the means ± standard deviation derived

from co-localization of the catheter segments.  Non-implanted catheters were used as negative

controls (**G**).

**Figure 5. Efg1 is responsible for inducing hyphal morphogenesis and is critical for**

**establishment of CAUTI.** (**A**) Comparison of biofilm formation on Fg-coated microplates by

*C. albicans* strains when grown in urine for 48 hours. (**B**) Microscopic visualization and 3D

reconstruction of *C. albicans*-fibrin/meshes interaction grown in urine using antibodies against

Fg (anti-Fg; green) and *C. albicans* (anti-*Candida*; red). Scale bars: 50 µm for 40x. (**C**) Mice

were catheterized and challenged with $1x10^6$ CFUs of each strain, and burdens in organs or

catheters were quantitated as the number of CFUs recovered. (**D**) RNA was extracted from 24

hpi catheterized and infected bladders with either the WT, *efg1*Δ/Δ, and *EFG1* complemented

strains and *EFG1* expression analysis was assessed by qRT-PCR. (**E**) Neutrophil recruitment to

the bladder in response of catheterization and infection. (**F**) Mouse bladders were harvested

following infection and catheter implantation. Bladders were subjected to analysis by H&E and

IF staining (anti-Fg, green; anti-*Candida,* red; anti-Ly6G, white; DAPI, blue). Values represent

means ± SEM. The Mann-Whitney U test was used; P < 0.05 was considered statistically

significant. *, P < 0.05; **, P < 0.005; ***, P < 0.0005. The horizontal bar represents the median

value. The horizontal broken line represents the limit of detection of viable fungi. Animals that

lost the catheter were not included in this work. ND, Not detectable.

**Figure 6. Als1 is critical for Fg-dependent biofilm formation and establishment of CAUTI.**
Biofilm formation analysis on Fg-coated microplates by *C. albicans* adhesin mutant strains (**A**) or the *ALS1* OE strain (**B**) when grown in human urine. At 48 hrs, *C. albicans* biofilm formation was measured via fluorescence intensity using anti-*Candida* antibodies (**A**) or crystal violet (**B**). (**A-B**) Data presented shows the mean and standard error derived from three independent experiments with 24 technical replicates. (**C**) Microscopic visualization and 3D reconstruction of *C. albicans*-fibrin/meshes interaction grown in urine using antibodies against Fg (green) and *C. albicans* (red). Scale bars: 50 µm for 40x. (**D**) Fungal burden in organs or catheters at 24 hpi were quantitated as the number of CFUs recovered. € RNA was extracted from 24 hpi catheterized and infected bladders with either the WT, *efg1*Δ/Δ, and *EFG1* complemented strains and *ALS1* expression analysis was assessed by qRT-PCR. (**F**) H&E and IF analysis of catheterized and infected bladders. IF staining was performed to detect Fg (green), *C. albicans* (red), and neutrophils (white). DAPI (blue) was used to stain the cell nuclei and delineated the urothelium (representative images). L, represents; U, urothelium; LP, the lamina propria; and M, muscularis. Bladder scale bars, 700 µm. White squares represent zoomed-in areas of higher magnification (20x, 40x, and 100x). Blue arrow heads on H&E indicate *C. albicans* bladder colonization. (**G**) *C. albicans* and Fg colocalization in implanted catheters using antibodies against Fg (anti-Fg; green) and *C. albicans* (anti-*Candida*; red). (**H**) Quantification of fungal colocalization with deposited Fg on the catheter. For biofilm formation, fungal burden and colocalization analysis values represent means ± SEM. The Mann-Whitney U test was used; *, P < 0.05; **, P < 0.005; ***, P < 0.0005; and ****, P < 0.0001. The horizontal bar represents the median value. (**D**) The horizontal broken line represents the limit of detection of viable fungi. Animals that lost the catheter were not included in this work.

**Figure 7. *C. albicans* hyphal morphogenesis and Als1-Fg interactions are critical for**

**CAUTI.** Urinary catheterization results in mechanical damage to the bladder and leads to the

release of Fg, accumulating in the bladder and depositing on the urinary catheter. The Fg-coated

catheter serves as a scaffold for *C. albicans* attachment, leading to robust biofilm formation

consisting of yeast, pseudohyphae, and hyphae. *C. albicans* attachment to Fg can be attributed

primarily to the adhesin Als1. As a result of this adherence to Fg, *C. albicans* is able to persist in

the bladder and establish CAUTI.

Biofilms

Candida-fibrin interaction
40x

Als1

*Candida* Yeast

*Candida* Hyphae

Fibrinogen