SynergyFinder Plus: Toward Better Interpretation and Annotation of Drug Combination Screening Datasets

Shuyu Zheng, Wenyu Wang, Jehad Aldahdooh, Alina Malyutina, Tolou Shadbahr, Ziaurrehman Tanoli, Alberto Pessia, Jing Tang*

Research Program in Systems Oncology, Faculty of Medicine, University of Helsinki, Helsinki 00290, Finland

Running title: *Zheng S et al / SynergyFinderPlus for Drug Combination Data Analysis*

* Corresponding author.

E-mail: jing.tang@helsinki.fi (Tang J).

Word count: 3742

Reference count: 50

Figure count: 6

Supplementary figure count: 3

Supplementary table count: 1

Count of letters in the article title: 91

Count of letters in the running title: 47

Count of keywords: 5

Count of words in Abstract: 204

Abstract

Combinatorial therapies have been recently proposed to improve the efficacy of anticancer treatment. The SynergyFinder R package is a software used to analyze pre-clinical drug combination datasets. Here, we report the major updates to the SynergyFinder R package for improved interpretation and annotation of drug combination screening results. Unlike the existing implementations, the updated SynergyFinder R package includes five main innovations. (1) We extend the mathematical models to higher-order drug combination data analysis and implement dimension reduction techniques for visualizing the synergy landscape. (2) We provide a statistical analysis of drug combination synergy and sensitivity with confidence intervals and P values. (3) We incorporate a synergy barometer to harmonize multiple synergy scoring methods to provide a consensus metric for synergy. (4) We evaluate drug combination synergy and sensitivity to provide an unbiased interpretation of the clinical potential. (5) We enable fast annotation of drugs and cell lines, including their chemical and target information. These annotations will improve the interpretation of the mechanisms of action of drug combinations. To facilitate the use of the R package within the drug discovery community, we also provide a web server at www.synergyfinderplus.org as a user-friendly interface to enable a more flexible and versatile analysis of drug combination data.

Keywords:

SynergyFinder; Drug combinations; Synergy modeling; Drug discovery; Drug combination sensitivity analysis

Introduction

Many complex diseases including cancer and infectious diseases develop drug resistance [1,2]. To achieve more sustainable clinical efficacy, multi-targeted drug combinations have been proposed to tackle disease signaling pathways more systematically [3]. With the advances in high-throughput drug screening technologies, an increasing number of drug combinations are being tested in multiple disease models, such as cancer cell lines and patient-derived primary cultures. These preclinical drug combination experiments are conducted to identify the most synergistic and effective drug combinations that result in improved cellular responses, such as cell viability and toxicity. With the potential hits prioritized from screening, more functional studies are warranted to elucidate the mechanisms of action of the drug interactions, leading to the identification of drug combination response biomarkers that are critical for stratifying patients for more targeted therapies.

For the evaluation of the potential of a drug combination, mathematical and statistical models are needed to characterize the expectation if the drugs are not interactive; thereafter the boosting effects of the drug combination can be formally quantified. There are currently four major synergy models: highest single agent (HSA) [4], Loewe additivity (LOEWE) [5], Bliss independence (BLISS) [6], and zero interaction potency (ZIP) [7]. However, the overall effects of a drug combination may not lead to sufficient efficacy, despite a strong degree of interaction. Therefore, it is recommended to evaluate both synergy (*i.e.,* the degree of interactions) and sensitivity (*i.e.,* the overall treatment efficacy) simultaneously [8]. Previously developed tools that can analyze drug combination synergy include COMPUSYSN [9], BRAID [10], Combenfit [11], SynergyFinder [12], and Synergy [13]. However, these tools are designed mainly for analyzing two-drug combinations only. More recently, the SynergyFinder2 tool [14] has been developed, which extends to three-drug combinations. However, only the HSA model was appropriately constructed for its implementation. The key differences in the modeling are summarized in Table S1. Furthermore, the visualization of high-order drug combinations becomes non-trivial, as none of the existing visualization methods can deal with combinations of three or more drugs [15]. More importantly, there is a lack of implementation tools that can harmonize multiple synergy models to derive a consensus about the degree of interactions [16].

To address these limitations, we provide a major update of the SynergyFinder R package that enables novel functions, including (1) formal mathematical models to assess the synergy of high-order drug combinations and visualization of the degree of synergy using a dimension reduction

technique; (2) formal statistical methods to evaluate the significance of synergy; (3) implementation
of the synergy barometer to systematically compare the results from different synergy models; and
(4) implementation of a synergy-sensitivity (SS) plot to evaluate the potential of a drug combination
unbiasedly. (5) Finally, we develop data annotation tools to retrieve the pharmacological and
molecular profiles of the drugs and cells, facilitating the discovery of mechanisms of action of the
most synergistic and effective drug combinations. We provide a new website at
www.synergyfinderplus.org to allow users to upload their experimental results and run all the
analyses with the R package in the backend. All functions and source codes are freely accessible to
academic users (**Figure 1**).

Method

Synergy models for high-order drug combinations
Considering the response of a drug to be measured as a % inhibition y that ranges from 0 to 1, a
higher value indicates a better efficacy. For a combination that involves n drugs, the observed
combination response is denoted as y_c, whereas the observed monotherapy response of its constituent
drugs is $y_{i,i=1,...n}$. The expected combination response is determined by the assumption of non-
interaction. Currently there are four major synergy reference models [16,17]. For the HSA model, the
expected response is the highest monotherapy response, that is, $y_{HSA} = \max(y_1, ..., y_i, ..., y_n)$. For
the BLISS model, the expected response is derived from the probabilistic independence of the
monotherapy effects, that is, $y_{BLISS} = 1 - \prod_i(1 - y_i)$. For the LOEWE model, the expected response
satisfies $\sum_i(\frac{x_i}{f_i^{-1}(y_{LOEWE})}) = 1$, where x_i is the dose of each constituent drug in the combination, and
$f_i^{-1}(y_{LOEWE})$ is the inverse function of the dose-response curve. For the ZIP model, the expected
response satisfies $y_{ZIP} = 1 - \prod_i(1 - \hat{y}_i)$, where $\hat{y}_i$ is the predicted dose response of the monotherapy
by a monotonically increasing curve fitting model $\hat{y}_i = f_i(x_i)$. For example, a common choice for
modeling drug dose-response curves is the four-parameter log-logistic model $f(x) =$
$\frac{E_{min}+E_{max}(x/m)^\lambda}{1+(x/m)^\lambda}$, where $E_{min}, E_{max}, m,$ and λ are the minimal and maximal responses, IC50, and
slope of the dose-response curve, respectively.

Accordingly, the multi-drug synergy score for the observed combination response y_c can be
determined as follows:

$$S_{HSA} = y_c - \max(y_1, ..., y_i, ..., y_n). \tag{1}$$

$$S_{BLISS} = y_c - \left(1 - \prod_i (1 - y_i) \right) \tag{2}$$

$$= y_c - \left(\sum_i y_i - \sum_{i<j} y_i y_j + (-1)^{r+1} \times \sum_{i<j} \prod_j^r y_j + (-1)^{n+1} \times \prod_i y_i \right)$$

$$S_{LOEWE} = y_c - y_{LOEWE}, s.t. \sum_i \left(\frac{x_i}{f_i^{-1}(y_{LOEWE})} \right) = 1 \tag{3}$$

110 To determine the ZIP-based synergy score, y_c needs to be replaced with the predicted average
111 response $\hat{y}_c$ given by the curve fitting models f_i' to make it comparable to y_{ZIP}:

$$S_{ZIP} = \hat{y}_c - y_{ZIP} = \frac{1}{n} \sum_i f_i'(x_i) - \left(1 - \prod_i (1 - f_i(x_i)) \right), \tag{4}$$

$$\text{where } Emin(f_i') = f_{-i}'(x_{-i})$$

Namely, $f_i'(x)$ stands for the log-logistic model defined for the combination response at dose x of
drug i when the other drugs are present. Furthermore, $Emin$ of $f_i'(x)$ is determined by $f_{-i}'(x_{-i})$,
which is the fitted curve of the combination response while drug i is absent. Note that the ZIP model
captures the shift of potency for a drug combination in comparison to its monotherapy drugs;
therefore, the ZIP model compares the difference between fitted models for the drug combination
$f_i'(x_i)$ and for the monotherapy drug $f_i(x_i)$.

Visualization of higher-order synergy and sensitivity scores
The commonly utilized synergy or sensitivity landscape models are not directly applicable for the
visualization of higher-order drug combinations. It is non-trivial to determine the coordinates of the
multi-dimensional dose vectors in a two-dimensional space where similar dose vectors remain
proximal to each other. In this study we propose a dimension reduction technique that is based on
multidimensional scaling, similar to the recent application in transforming numerical data into images
[18]. For a drug combination in an n-dimensional dose space, with its coordinates of $X = (x_1, \ldots x_n)$,
we consider their rankings $R = (r_1, \ldots r_n)$. For example, a combination $X = (0.1, 10, 1000)$ nM in
three-dimensional dose space has a ranking vector R = (1, 3, 6), suggesting that 0.1 nM is the
minimum dose in the tested dose range of drug 1, 10 nM is the third smallest dose of drug 2, and
1000 nM is the sixth dosage of drug 3. Euclidean distance is a common choice for quantifying the
similarity between two dose ranking vectors. We utilize multi-dimensional scaling to minimize the
error of the pairwise distance in a two-dimensional space in which the synergy and sensitivity scores
can be visualized as a synergy landscape (**Figure 2**). Using the dose rankings as the input for multi-
dimensional scaling can ensure that the resulting two-dimensional coordinates are equally distanced
among the neighboring dose conditions, thus making the visualization easier to interpret.

Furthermore, for the case of two-drug combinations, the algorithm can converge to the actual dose

rankings, thus preserving the consistency across all orders of combinations. We also develop

functions to visualize the synergy and sensitivity scores for a specific dose condition in a grid of bar

plots, which can help identify the most synergistic and sensitive areas in the synergy landscape.

Statistical analysis

The statistical significance of synergy can be defined at a single dose or at the whole dose matrix

level. Consider a two-drug combination experiment as an example. We assume that the replicates of

drug responses are measured independently within each dose in the dose matrix (**Figure 3**). At each

dose level, we propose using a bootstrapping approach to determine the confidence intervals of the

synergy scores. To determine a bootstrap dose-response matrix, we sample the responses for each

dose combination from a normal distribution $N(\mu, \sigma)$, where μ and σ are the mean and standard

deviation of the replicates. The synergy scores for the HSA and BLISS models can be derived simply

by comparing the bootstrapped responses at the dose combination and at their corresponding

monotherapy doses (*i.e.*, using Equations (1) and (2)). However, for the LOEWE and ZIP models [7],

bootstrapped synergy scores will be derived using the curve fitting over the whole dose matrix (*i.e.,*

Equations (3) and (4)).

Suppose that B bootstrap samples are drawn from the replicates and suppose that $s_1, \ldots, s_B$ are the

estimates of the synergy scores determined from these bootstrap samples, with a mean of $\bar{s}$. We

determine the bootstrap standard error (SE) as follows:

$$SE = \sqrt{\frac{1}{B-1} \sum_{i=1}^{B} (s_i - \bar{s})^2} \tag{5}$$

The 95% confidence interval for the synergy score is approximately

$$[\bar{s} - 1.96SE, \bar{s} + 1.96SE]$$

At the whole dose matrix level, we provide an empirical P value to assess the significance of the

difference between the estimated average synergy score over the whole dose matrix and the expected

synergy score of zero under the null hypothesis of non-interaction.

Letting $s'_1, \ldots, s'_B$ are the estimates of the average synergy scores over the whole dose matrix from

the bootstrap samples, the P value is [19]:

$$p = \exp(-0.717z - 0.416z^2), \text{ where } z = abs(\bar{s}') / \sqrt{\frac{1}{B-1} \sum_{i=1}^{B} (s_i' - \bar{s}')^2}$$

For a scenario in which no replicates are available, the confidence interval cannot be determined at

the dose level. However, the *P* value of the average synergy score of the whole dose-response matrix

can still be derived by pooling the synergy scores together and then comparing them to zero. For both

scenarios, we report the overall *P* values of synergy and the confidence intervals at the dose-specific

level when replicates are available. We test the accuracy of the bootstrapping with the simulation

(File S1) and show that the *P* values for synergy scores determined from bootstrap are highly

correlated with the true *P* values (Figure S1).

Harmonization of multiple synergy models

As the SynergyFinder R package provides four synergy models (*i.e.*, HSA, BLISS, LOEWE, and

ZIP) that differ in their null hypotheses of non-interaction, the synergy scores do not necessarily

coincide with each other for a given dataset. To provide a more harmonized representation of the

synergy score results, we introduce a novel tool to enable a more systematic comparison of the

synergy scores. We leverage the implementation of the four models to derive the expected response

of non-interaction (*i.e.*, y_{HSA}, y_{BLISS}, y_{LOEWE}, and y_{ZIP}). Because all the expected responses share

the same unit as the observed response (*i.e.*, % inhibition), we may utilize a synergy barometer to

compare these values. The expected and observed responses for a given combination at a specific

dose are positioned on the same scale. With such a tool as the synergy barometer, one can evaluate

the degree of synergy of a specific model and easily understand the differences in the results. A strong

synergy should be concluded if the observed response goes beyond the expected responses of all the

four models.

Prioritization of drug combinations

In addition to synergy, the sensitivity of a drug combination is equally important [8]. Prioritization

of drug combinations based on the degree of synergy may only result in an excessive number of false

positives, as a clinical endpoint for approving a drug combination is commonly its efficacy rather

than the synergy. In cases where a strong synergy is obtained, the drug combination may not

necessarily reach sufficient levels of therapeutic efficacy. To capture the sensitivity of a drug

combination, we previously developed a combination sensitivity score (CSS) model that calculates

the relative inhibition of a drug combination based on the area under the log10 scaled dose-response

curves at the IC50 doses of the constituent drugs [8]. The CSS has the same unit as a drug response

(*i.e.*, % inhibition), which makes it directly comparable with the synergy scores defined in Equations

(1–4). Therefore, we propose a scatter plot of CSS versus synergy score for a set of drug

combinations. The proposed sensitivity-synergy (SS) plot significantly aids in the prioritization of drug combinations from a high-throughput experiment.

Annotation of drug combinations

Drugs and cell lines in a drug screening dataset are annotated by querying related databases. PubChem CID, Isomeric SMILES, and standard InChIKey for drugs are extracted from PubChem [20]. Furthermore, we provide the molecular formula, clinical stage, disease indication, cross references to multiple major chemical compound databases (*i.e.*, ChEMBL [21], ChEBI [22], DrugBank [23], BindingDB [24], PharmGKB [25], Guide to Pharmacology [26], Selleck, and ZINC [27]), and drug target profiles from MICHA [28]. Cell line annotation is obtained from Cellosaurus [29] (Version 37.0).

Results

Analysis of three-drug combinations

To test the algorithms designed for higher-order combinations, we utilize a recent anti-malaria study in which 16 triple artemisinin-based combination therapies have been tested against 15 parasite lines [30]. The study aimed to identify synergistic partner drugs that can be combined with artemisinin to overcome the emergence of malaria resistance. For each triple drug combination, a $10 \times 10 \times 12$ multi-dimensional dose matrix was constructed, resulting in n = 1200 viability values. The datasets are obtained from the NCATS data portal at https://tripod.nih.gov/matrix-client/.

We analyze the triple drug combinations in terms of their sensitivity and synergy and visualize them in surface plots using dimension reduction techniques. **Figure 4** shows a synergistic triple combination, pyronaridine tetraphosphate-piperaquine-darunavir ethanolate (PYR-PQP-DRV) compared to an antagonistic combination, pyronaridine tetraphosphate-piperaquine-lopinavir (PYR-PQP-LPV). Despite similar sensitivities (the average % inhibition is 78.87 for PYR-PQP-DRV and 74.97 for PYR-PQP-LPV), their synergy scores differ drastically (the average ZIP score is 12.02 for PYR-PQP-DRV and -13.39 for PYR-PQP-LPV), suggesting more complex behaviors of the drug interactions as the number of drugs increases. Because of the lack of computational tools for analyzing higher-order drug combinations, the authors of the anti-malaria studies provided only the results of pairwise drug combinations, whereas the actual interactions of the three drugs were unexplored. Therefore, our method may provide novel insights that leverage higher-order drug combination design in a more tailored manner.

Analysis of two-drug combinations with replicates

We obtain the ONEIL dataset from the DrugComb data portal [31]. The ONEIL data is a pan-cancer drug combination study in which 583 drug combinations were tested across 39 cell lines [32]. Four replicates were produced for each drug combination, making it a representative example dataset to analyze the significance of the synergy scores. In **Figure 5**, we show the selective drug combinations tested in the A2058 cell line (melanoma).

As shown in Figure 5A, the confidence intervals for the responses and synergy scores vary at different doses, suggesting that not all scores are equally significant. Furthermore, we observe a general trend that a higher response or synergy score tends to have a smaller confidence interval, consistent with the expectation of a typical drug screening experiment. To evaluate the synergy scores more systematically, we provide a synergy barometer for specific dose combinations. For example, for MK-1775 at 0.2 µM combined with niraparib at 2.75 µM, the drug combination response reaches 77.84% inhibition, as indicated by the pointer readout on the barometer (Figure 5B, left panel). Such a drug combination response is synergistic, as the expected responses of the HSA, LOEWE, BLISS, and ZIP models are much smaller, shown as the marks on the barometer. In contrast, paclitaxel at 0.0016 µM in combination with L-778123 at 2 µM shows a near-zero response (0.29% inhibition), which is much smaller than the expected response among all the synergy models (Figure 5B, right panel); therefore, it is considered strong antagonism. Using the synergy barometer, the actual drug combination response can be directly compared with the expectations of non-interaction among multiple synergy models.

Furthermore, we also show the CSS and ZIP synergy scores of all the drug combinations tested in the A2058 cell line in an SS plot (Figure 5C, Figure S3). CSS indicates the efficacy of a drug combination, whereas the synergy score indicates the degree of interaction. To prioritize potential drug combinations, it is necessary to identify those with higher CSS and higher synergy scores, that is, the top-right corner of the SS plot.

Annotation of mechanisms of action of drug combinations

Recent advances in machine learning and artificial intelligence studies have shown great potential in predictive modeling of drug synergies [33–35]. In these studies, the chemical structure information regarding drugs and the molecular profiles of cells are considered predictive features. However, for most of the drug combination datasets, annotations of drug and cell features are not provided. To

facilitate the interpretation of mechanisms of action of drug combinations, it would be beneficial to provide data integration tools for effective annotation and accurate matching to public databases such that more comprehensive features used for synergy predictions can be obtained. Efforts to annotate and harmonize existing drug screening data, such as DrugComb [31,36] and DrugCombDB [37], have significantly contributed to the development of data-driven pharmacological modeling [38–40]. For newly generated drug screening datasets, we develop the SynergyFinder Plus portal as a companion tool for retrieving publicly available information in a more automated fashion. With one click of the button, users can obtain cross-database identifiers of the drugs and cells, as well as additional information, such as the mechanism of action and disease indication (**Figure 6**).

We show how SynergyFinder Plus annotates a particular drug combination in the ONEIL [32] dataset (5-fluorouracil and vorinostat tested in the A2058 cell line, Figure 6A). After users upload the dataset, SynergyFinder Plus returns two annotation tables describing the cell and drug identities. For example, the cell annotation table includes synonyms, Cellosaurus accession, tissue origin, disease name, and disease NCI Thesaurus ID, whereas the drug annotation table includes InChIKey, isomeric SMILES, molecular formula, clinical trial phase, cross references, and disease indication. In addition, a third table listing primary and potent drug target information is provided. These annotations enable a more systematic exploration of drugs and cell lines tested in a high-throughput experiment (Figure 6B). With such a comprehensive collection of identifiers, we believe that the drug combination data can be better integrated with other related data sources for further downstream analysis.

Discussion

Synergistic and effective drug combinations have long been studied to improve disease management. Testing potential drug combinations using clinically representative cell cultures has become an important tool for detecting signs of drug sensitivity and resistance; in some cases, the results have already provided informative decision-making resources to guide patient stratification in clinical trials [41]. To facilitate the prioritization of drug combinations, informatics tools that enable formal assessment of the degree of synergy as well as sensitivity are highly needed. We provide a major update to the commonly used SynergyFinder R package, which models drug interactions for any higher-order combinations [15]. Higher-order drug combinations have been recently explored in lung cancer [42], colorectal cancer [43], and ovarian cancer [44]. Therefore, it is expected that promising multi-drug combinations will enter formal clinical trials for multiple diseases [45]. Furthermore, we develop a statistical analysis of drug combinations, which is also generally applicable for higher-

order combinations. Importantly, the mathematical models for higher-order combinations have been
consistent with those for two-drug combinations; therefore, their scores are readily comparable across
different orders. To visualize higher-order combinations, we develop a dimension reduction
technique to display the synergy and the sensitivity landscape on a two-dimensional plane.

In addition to providing multiple synergy models, we address the problem of harmonization. With
the increasing number of models that have been proposed to characterize drug synergy, there is an
increasing need to help users understand the differences between these models such that a user may
make an informed interpretation of the drug combination results. We propose the use of a synergy
barometer on which all the synergy models can be systematically compared with the observed drug
combination response. Despite only including the four major synergy models, HSA, BLISS, LOEWE,
and ZIP, we propose the incorporation of other synergy models that can readily have their synergy
scores put into the barometer for comparison with existing models. With the synergy barometer, we
hope that the drug discovery community may derive a consensus on the best reporting guideline for
drug combination data analysis, which has been long sought since the time of the Saariselkä
agreement [46] and thereafter [47].

We also provide the formal statistical analysis for replicates of drug combinations. The confidence
intervals and P values for all four synergy scores are comparable. Even for the cases in which no
replicates have been performed, the significance of synergy over all the dose conditions can be
provided, thus enabling a statistical evaluation of drug combinations in a primary screen that usually
does not provide replicates due to the high number of combinations. Once a drug combination hit
passes the initial screening based on the synergy barometer and the statistical significance, we
recommend a confirmation screen that involves more doses and replicates to formally evaluate its
synergy. If a user would like to study the mechanism of action of the drug combination, we also
provide integrative tools to retrieve the chemical structure information and other functional
annotations of the drugs, which will help in downstream analysis of the drug combinations. For
higher-order drug combinations, the interpretation of drug synergies is a challenging problem. We
hope that our tool provides the first evidence to pinpoint synergistic higher-order drug combinations.

Finally, we provide new features to harmonize the assessment of synergy and sensitivity. The
synergy models that have been developed in the SynergyFinder R package have the same scale as the
drug combination sensitivity method that was developed earlier, thus allowing a direct comparison in
an SS plot. We recommend that users consider the SS plot when reporting their drug combination

results, as sensitivity is the clinical endpoint for approving any drug or drug combinations. The drug combination sensitivity values also have the same scale as monotherapy drug responses in the unit of percentage inhibition; therefore, we can readily compare a drug combination with a single drug to evaluate their efficacy. With the harmonization of drug combination sensitivity and monotherapy sensitivity, we may provide an integrated data source for developing advanced machine learning approaches for predicting drug sensitivity, such as drug sensitivity datasets deposited in DrugComb [31]. To facilitate the FAIRification (findable, accessible, interoperable, and reusable characteristics) of drug combination analysis, we provide a web server at www.synergyfinderplus.org to allow users to upload the data and obtain the analysis results in the form of reports. In addition, we encourage users to deposit their experimental data in DrugComb at https://drugcomb.org/, currently one of the most comprehensive drug screening databases. With the algorithms and the source code made available through the SynergyFinder R package, we welcome the machine learning community to leverage the harmonized drug combination and monotherapy datasets for more robust and transferable predictions to further facilitate the drug combination discovery [34,48–50].

Availability

The www.synergyfinderplus.org web server is located at the CSC-IT Center for Science in Finland. All the source code for implementing the data analysis and visualization is available as the SynergyFinder R package at https://bioconductor.org/packages/release/bioc/html/synergyfinder.html.

Author CrediT statement

Shuyu Zheng: Conceptualization, Methodology, Software, Writing – Original draft preparation **Wenyu Wang**: Visualization, Writing – Original draft preparation **Jehad Aldahdooh**: Software **Alina Malyutina**: Methodology **Tolou Shadbahr**: Data Curation **Ziaurrehman Tanoli**: Resources **Alberto Pessia**: Methodology, Validation, Writing – Review & Editing **Jing Tang**: Supervision, Conceptualization, Methodology, Software, Writing – Original draft preparation

Acknowledgments

This work is supported by the European Research Council (ERC) starting grant DrugComb (Informatics approaches for the rational selection of personalized cancer drug combinations) [No. 716063], European Commission H2020 EOSC-life (Providing an open collaborative space for digital biology in Europe [No. 824087], and the Academy of Finland grant [No. 317680]. The Sigrid Jusélius Foundation grant. SZ and WW hold a salaried position funded by the University of Helsinki through

361   the Doctoral Program of Biomedicine (DPBM). WW also receives a personal grant from K. Albin

362   Johanssons stiftelse. AM holds a salaried position funded by the University of Helsinki through the

363   Doctoral Program of Integrative Life Science (ILS), and receives a personal grant from the Cancer

364   Foundation Finland.

Conflict of interests

None declared.

ORCID

0000-0003-0624-8077 (Shuyu Zheng)

0000-0003-2816-9740 (Wenyu Wang)

0000-0002-9947-4067 (Jehad Aldahdooh)

0000-0001-8220-5859 (Alina Malyutina)

0000-0001-5715-5933 (Tolou Shadbahr)

0000-0003-2435-9862 (Ziaurrehman Tanoli)

0000-0001-8607-9191 (Alberto Pessia)

0000-0001-7480-7710 (Jing Tang)

References

[1]   Vasan N, Baselga J, Hyman DM. A view on drug resistance in cancer. Nature 2019;575:299–309.

[2]   Nathan C. Resisting antimicrobial resistance. Nat Rev Microbiol 2020;18:259–60.

[3]   Boshuizen J, Peeper DS. Rational cancer treatment combinations: an urgent clinical need. Mol Cell 2020;78:1002–18.

[4]   Berenbaum MC. What is synergy? Pharmacol Rev 1989;41:93–141.

[5]   Loewe S. The problem of synergism and antagonism of combined drugs. Arzneimittelforschung 1953;3:285–90.

[6]   Bliss CI. The toxicity of poisons applied jointly. Ann App Biol 1939;26:585–615.

[7]   Yadav B, Wennerberg K, Aittokallio T, Tang J. Searching for drug synergy in complex dose–response landscapes using an interaction potency model. Comput Struct Biotechnol J 2015;13:504–13.

[8]   Malyutina A, Majumder MM, Wang W, Pessia A, Heckman CA, Tang J. Drug combination sensitivity scoring facilitates the discovery of synergistic and efficacious drug combinations in cancer. PLoS Computat Biol 2019;15:e1006752.

[9]   Chou T-C. Drug combination studies and their synergy quantification using the chou-talalay method. Cancer Res 2010;70:440–6.

[10]   Twarog NR, Stewart E, Hammill CV, Shelat AA. BRAID: A unifying paradigm for the analysis of combined drug action. Sci Rep 2016;6:25523.

[11] Di Veroli GY, Fornari C, Wang D, Mollard S, Bramhall JL, Richards FM, et al. Combenefit:
an interactive platform for the analysis and visualization of drug combinations. Bioinformatics
2016;32:2866–8.
[12] Ianevski A, He L, Aittokallio T, Tang J. SynergyFinder: a web application for analyzing drug
combination dose-response matrix data. Bioinformatics 2017;33:2413–5.
[13] Wooten DJ, Albert R. Synergy: a Python library for calculating, analyzing, and visualizing
drug combination synergy. Bioinformatics 2020;37:1473–4.
[14] Ianevski A, Giri AK, Aittokallio T. SynergyFinder 2.0: visual analytics of multi-drug
combination synergies. Nucleic Acids Res 2020;48:W488–93.
[15] Cokol-Cakmak M, Cetiner S, Erdem N, Bakan F, Cokol M. Guided screen for synergistic
three-drug combinations. PLoS One 2020;15:e0235929.
[16] Meyer CT, Wooten DJ, Lopez CF, Quaranta V. Charting the fragmented landscape of drug
synergy. Trends Pharmacol Sci 2020;41:266–80.
[17] Vlot AHC, Aniceto N, Menden MP, Ulrich-Merzenich G, Bender A. Applying synergy
metrics to combination screening data: agreements, disagreements and pitfalls. Drug Discov
Today 2019;24:2286–98.
[18] Bazgir O, Zhang R, Dhruba SR, Rahman R, Ghosh S, Pal R. Representation of features as
images with neighborhood dependencies for compatibility with convolutional neural networks.
Nat Commun 2020;11:4391.
[19] Altman DG, Bland JM. How to obtain the P value from a confidence interval. BMJ
2011;343:d2304.
[20] Kim S, Chen J, Cheng T, Gindulyte A, He J, He S, et al. PubChem in 2021: new data content
and improved web interfaces. Nucleic Acids Res 2021;49:D1388–95.
[21] Mendez D, Gaulton A, Bento AP, Chambers J, De Veij M, Félix E, et al. ChEMBL: towards
direct deposition of bioassay data. Nucleic Acids Res 2019;47:D930–40.
[22] Hastings J, Owen G, Dekker A, Ennis M, Kale N, Muthukrishnan V, et al. ChEBI in 2016:
Improved services and an expanding collection of metabolites. Nucleic Acids Res
2016;44:D1214–9.
[23] Wishart DS, Feunang YD, Guo AC, Lo EJ, Marcu A, Grant JR, et al. DrugBank 5.0: a major
update to the DrugBank database for 2018. Nucleic Acids Res 2018;46:D1074–82.
[24] Gilson MK, Liu T, Baitaluk M, Nicola G, Hwang L, Chong J. BindingDB in 2015: a public
database for medicinal chemistry, computational chemistry and systems pharmacology.
Nucleic Acids Res 2016;44:D1045–53.
[25] Whirl-Carrillo M, Huddart R, Gong L, Sangkuhl K, Thorn CF, Whaley R, et al. An evidence-
based framework for evaluating pharmacogenomics knowledge for personalized medicine.
Clin Pharmacol Ther 2021;110:563–72.
[26] Armstrong JF, Faccenda E, Harding SD, Pawson AJ, Southan C, Sharman JL, et al. The
IUPHAR/BPS Guide to PHARMACOLOGY in 2020: extending immunopharmacology
content and introducing the IUPHAR/MMV Guide to MALARIA PHARMACOLOGY.
Nucleic Acids Res 2020;48:D1006–21.
[27] Sterling T, Irwin JJ. ZINC 15 – ligand discovery for everyone. J Chem Inf Model
2015;55:2324–37.
[28] Tanoli Z, Aldahdooh J, Alam F, Wang Y, Seemab U, Fratelli M, et al. Minimal information
for chemosensitivity assays (MICHA): a next-generation pipeline to enable the FAIRification
of drug screening experiments. Brief Bioinform 2021;bbab350.
[29] Bairoch A. The Cellosaurus, a cell-line knowledge resource. J Biomol Tech 2018;29:25–38.
[30] Ansbro MR, Itkin Z, Chen L, Zahoranszky-Kohalmi G, Amaratunga C, Miotto O, et al.
Modulation of triple artemisinin-based combination therapy pharmacodynamics by
plasmodium falciparum genotype. ACS Pharmacol Transl Sci 2020;3:1144–57.

[31] Zagidullin B, Aldahdooh J, Zheng S, Wang W, Wang Y, Saad J, et al. DrugComb: an integrative cancer drug combination data portal. Nucleic Acids Res 2019;47:W43–51.

[32] O'Neil J, Benita Y, Feldman I, Chenard M, Roberts B, Liu Y, et al. An unbiased oncology compound screen to identify novel combination strategies. Mol Cancer Ther 2016;15:1155–62.

[33] Preuer K, Lewis RPI, Hochreiter S, Bender A, Bulusu KC, Klambauer G. DeepSynergy: predicting anti-cancer drug synergy with deep learning. Bioinformatics 2018;34:1538–46.

[34] Menden MP, Wang D, Mason MJ, Szalai B, Bulusu KC, Guan Y, et al. Community assessment to advance computational prediction of cancer drug combinations in a pharmacogenomic screen. Nat Commun 2019;10:2674.

[35] Li H, Li T, Quang D, Guan Y. Network propagation predicts drug synergy in cancers. Cancer Res 2018;78:5446–57.

[36] Zheng S, Aldahdooh J, Shadbahr T, Wang Y, Aldahdooh D, Bao J, et al. DrugComb update: a more comprehensive drug sensitivity data repository and analysis portal. Nucleic Acids Res 2021;49:W174–84.

[37] Liu H, Zhang W, Zou B, Wang J, Deng Y, Deng L. DrugCombDB: a comprehensive database of drug combinations toward the discovery of combinatorial therapy. Nucleic Acids Res 2020;48:D871–81.

[38] Douglass EF, Allaway RJ, Szalai B, Wang W, Tian T, Fernández-Torras A, et al. A community challenge for pancancer drug mechanism of action inference from perturbational profile data. BioRxiv 2020:2020.12.21.423514.

[39] Zhang B, Tang C, Yao Y, Chen X, Zhou C, Wei Z, et al. The tumor therapy landscape of synthetic lethality. Nat Commun 2021;12:1275.

[40] Shah K, Ahmed M, Kazi JU. The Aurora kinase/β-catenin axis contributes to dexamethasone resistance in leukemia. NPJ Precis Oncol 2021;5:13.

[41] Jones PA, Ohtani H, Chakravarthy A, De Carvalho DD. Epigenetic therapy in immune-oncology. Nat Rev Cancer 2019;19:151–61.

[42] Fernandes Neto JM, Nadal E, Bosdriesz E, Ooft SN, Farre L, McLean C, et al. Multiple low dose therapy as an effective strategy to treat EGFR inhibitor-resistant NSCLC tumours. Nat Commun 2020;11:3157.

[43] Ozkan-Dagliyan I, Diehl JN, George SD, Schaefer A, Papke B, Klotz-Noack K, et al. Low-dose vertical inhibition of the RAF-MEK-ERK cascade causes apoptotic death of *KRAS* mutant cancers. Cell Rep 2020;31:107764.

[44] Caumanns JJ, van Wijngaarden A, Kol A, Meersma GJ, Jalving M, Bernards R, et al. Low-dose triple drug combination targeting the PI3K/AKT/mTOR pathway and the MAPK pathway is an effective approach in ovarian clear cell carcinoma. Cancer Lett 2019;461:102–11.

[45] Van Cutsem E, Huijberts S, Grothey A, Yaeger R, Cuyle P-J, Elez E, et al. Binimetinib, encorafenib, and cetuximab triplet therapy for patients with *BRAF* V600E–mutant metastatic colorectal cancer: safety lead-in results from the phase III BEACON colorectal cancer study. J Clin Oncol 2019;37:1460–9.

[46] Greco W, Unkelbach H, Pöch G, Sühnel J, Kundi M, Bödeker W. Consensus on concepts and terminology for combined-action assessment: the Saariselkä agreement. Arch Complex Environ Stud 1992;4:65–9.

[47] Tang J, Wennerberg K, Aittokallio T. What is synergy? The Saariselkä agreement revisited. Front Pharmacol 2015;6:181.

[48] Wood K, Nishida S, Sontag ED, Cluzel P. Mechanism-independent method for predicting response to multidrug combinations in bacteria. Proc Natl Acad Sci U S A. 2012;109:12254–9.

[49] Adam G, Rampášek L, Safikhani Z, Smirnov P, Haibe-Kains B, Goldenberg A. Machine learning approaches to drug response prediction: challenges and recent progress. NPJ Precis Oncol 2020;4:19.

[50] Kim Y, Zheng S, Tang J, Jim Zheng W, Li Z, Jiang X. Anticancer drug synergy prediction in understudied tissues using transfer learning. J Am Med Inform Assoc 2021;28:42–51.

Figure legends

Figure 1 Summary of the SynergyFinder Plus workflow

We highlight the new features including: (1) the ability to analyze drug combinations of more than two drugs, (2) tailored statistical testing to account for uncertainty for data with replicates, (3) harmonized visualization of multiple synergy score metrics, (4) visualization of the relationship between sensitivity and synergy scores, and (5) a data annotation web server to query cell line and drug information.

Figure 2 Dimension reduction for the visualization of high-order drug combinations

In this example, the two-dimensional coordinates of each three-drug dose combination are determined by multidimensional scaling based on the similarity of their dose ranges, after which the synergy scores and sensitivity scores can be visualized on the map. Each dot in the map is annotated with its associated dose rankings for the three drugs ($r_1; r_2; r_3$). For example, a dot with a label 4;1;2 shows a combination involves 4^{th} dose of drug 1, 1^{st} dose of drug 2, and 2^{nd} dose of drug 3.

Figure 3 Statistical testing of synergy scores

Dose-specific confidence intervals and P values for the average synergy score can be derived using bootstrapping of the dose-response matrix.

Figure 4 Example of surface plots for higher-order drug combinations

A. The dose-response map for a synergistic drug combination (pyronaridine tetraphosphate-piperaquine-darunavir ethanolate, left panel) compared to an antagonistic drug combination (pyronaridine tetraphosphate-piperaquine-lopinavir, right panel). **B**. The ZIP synergy score map for the same drug combinations. For surface plots of other synergy scores, including HSA, BLISS, and LOEWE, see Figure S2.

Figure 5 Example of visualizations for two-drug combinations with replicates

A. An example of the confidence interval plots of a synergistic drug pair (MK-1775 and niraparib;

upper panel) and an antagonistic drug pair (paclitaxel and L-778123; lower panel). **B**. The synergy

barometer for a given dose combination (left panel: MK-1775 at 0.2 μM in combination with

niraparib at 2.75 μM; right panel: paclitaxel at 0.0016 μM in combination with L-778123 at 2 μM).

C. The synergy-sensitivity plot for drug pairs tested in the A2058 cell line.

Figure 6 Example of mechanisms of action annotation for drug combination

A. An example of a drug screening annotation table. The tested drug combination, 5-fluorouracil and

vorinostat, is further annotated with their chemical and target information. Cell line information for

the A2058 is also provided, including synonyms, tissue, and disease types. **B**. Summary of drug

clinical stage, drug target protein class, and cell line tissue origin for the ONEIL dataset.

Supplementary material

Figure S1 Accuracy of *P* values for synergy scores by bootstrapping

A. The expected monotherapy responses are defined by four-parameter log-logistic functions and the

expected combinational responses are defined by the non-interaction models. Based on the expected

values, the distribution of the test statistic under null hypothesis can be determined by simulations.

B. The scatter plots for comparing the true *P* values and bootstrapping *P* values for 200 samples.

Figure S2 Two-dimensional surface plots of synergy scores

A. HSA, **B.** BLISS, and **C.** LOEWE for a synergistic three-drug combination (pyronaridine

tetraphosphate-piperaquine-darunavir ethanolate, left panel) compared to an antagonistic drug

combination (pyronaridine tetraphosphate-piperaquine-lopinavir, right panel).

Figure S3 Synergy-sensitivity plots for HSA, BLISS, and LOEWE scores in relation to the

combination sensitivity score for the drug pairs tested in the A2058 cell line

Table S1 The key differences in the mathematical modeling between SynergyFinderPlus and

SynergyFinder2 for three-drug combinations

File S1 The simulation code for testing accuracy of *P* values for synergy scores by bootstrapping

561   **Figure 1**

 Figure 2

579 **Figure 3**

582   **Figure 4**

A

Dose response matrix

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Darunavir ethanolate

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Lopinavir

B

ZIP synergy score

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Darunavir ethanolate

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Lopinavir

Figure 5

589   **Figure 6**

590
591

592   **Figure S1**

A

595 **Figure S2**

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Darunavir ethanolate

Drug1: Pyronaridine tetraphosphate
Drug2: Piperaquine
Drug3: Lopinavir

HSA

BLISS

LOEWE

598 **Figure S3**

601 **Table S1 The key differences in the mathematical modeling between SynergyFinderPlus and**
602 **SynergyFinder2 for three-drug combinations**
603

	SynergyFinderPlus	SynergyFinder2
BLISS	$S_{BLISS} = E_{A,B,C} - (E_A + E_B + E_C - E_A E_B - E_A E_C - E_B E_C + E_A E_B E_C)$	$S_{BLISS} = E_{A,B,C} - (E_A + E_B + E_C - E_A E_B - E_A E_C - E_B E_C - E_A E_B E_C)$
LOEWE	$S_{LOEWE} = E_{A,B,C} - E_{LOEWE}$, $$s.t. \sum_{i \in \{A,B,C\}} \left(\frac{x_i}{f_i^{-1}(E_{LOEWE})} \right) = 1$$	$$S_{LOEWE} = \frac{a}{A} + \frac{b}{B} + \frac{c}{C}$$
ZIP	$S_{ZIP} = \hat{E}_{A,B,C} - \left(\hat{E}_A + \hat{E}_B + \hat{E}_C - \hat{E}_A \hat{E}_B - \hat{E}_A \hat{E}_C - \hat{E}_B \hat{E}_C + \hat{E}_A \hat{E}_B \hat{E}_C \right)$ $$\hat{E}_{A,B,C} = \frac{1}{3} \left(\frac{\hat{E}_{-A} + \left(\frac{x_A}{\hat{m}_A}\right)^{\hat{\lambda}_A}}{1 + \left(\frac{x_A}{\hat{m}_A}\right)^{\hat{\lambda}_A}} + \frac{\hat{E}_{-B} + \left(\frac{x_B}{\hat{m}_B}\right)^{\hat{\lambda}_B}}{1 + \left(\frac{x_B}{\hat{m}_B}\right)^{\hat{\lambda}_B}} + \frac{\hat{E}_{-C} + \left(\frac{x_C}{\hat{m}_C}\right)^{\hat{\lambda}_C}}{1 + \left(\frac{x_C}{\hat{m}_C}\right)^{\hat{\lambda}_C}} \right)$$	$S_{ZIP} = E_{A,B,C} - \left(\hat{E}_A + \hat{E}_B + \hat{E}_C - \hat{E}_A \hat{E}_B - \hat{E}_A \hat{E}_C - \hat{E}_B \hat{E}_C - \hat{E}_A \hat{E}_B \hat{E}_C \right)$

604