Transcriptomic changes highly similar to Alzheimer's disease are observed in a subpopulation of individuals during normal brain aging

Shouneng Peng[1,2,3], Lu Zeng[1,2,3], Jean-vianney Haure-mirande[1,2,3], Minghui Wang[1,2,3], Derek M. Huffman[4,5,6], Vahram Haroutunian[7,8], Michelle Erlich[9,10], Bin Zhang[1,2,3], Zhidong Tu[1,2,3*]

[1] Institute of Genomics and Multiscale Biology, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York City, NY 10029, USA

[2] Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York City, NY 10029, USA

[3] Icahn Institute for Data Science and Genomic Technology, Icahn School of Medicine at Mount Sinai, One Gustave L. Levy Place, New York, NY 10029, USA

[4] Departments of Medicine, Albert Einstein College of Medicine, 1300 Morris Park Avenue, New York, NY 10461, USA

[5] Institute for Aging Research, Albert Einstein College of Medicine, 1300 Morris Park Avenue, New York, NY 10461, USA

[6] Departments of Molecular Pharmacology, Medicine and Institute for Aging Research Albert Einstein College of Medicine, 1300 Morris Park Avenue, New York, NY 10461, USA

[7] Departments of Psychiatry and Neuroscience, Icahn School of Medicine at Mount Sinai, New York City, NY, USA

[8] Mental Illness Research, Education and Clinical Center (MIRECC), James J. Peters VA Medical Center, Bronx, NY, USA

[9] Department of Neurology, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[10] Departments of Pediatrics and Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, USA

[*]Corresponding author: Zhidong Tu

IMI 3-70F, 1425 Madison Ave

New York City, NY 10029

E-mail: zhidong.tu@mssm.edu

Office Phone: 212-659-8508

Abstract

Aging is a major risk factor for late-onset Alzheimer's disease (LOAD). How aging contributes to the development of LOAD remains elusive. In this study, we examined multiple large-scale transcriptomic data from both normal aging and LOAD brains to understand the molecular interconnection between aging and LOAD. We found that shared gene expression changes between aging and LOAD are mostly seen in the hippocampal and several cortical regions. In the hippocampus, the expression of phosphoprotein, alternative splicing and cytoskeleton genes are commonly changed in both aging and AD, while synapse, ion transport, and synaptic vesicle genes are commonly down-regulated. Aging-specific changes are associated with acetylation and methylation, while LOAD-specific changes are more related to glycoprotein (both up- and down-regulations), inflammatory response (up-regulation), myelin sheath and lipoprotein (down-regulation). We also found that normal aging brain transcriptomes from relatively young donors (45-70 years old) cluster into several subgroups and some subgroups showed gene expression changes highly similar to those seen in LOAD brains. Using brain transcriptomic data from another cohort of older individuals (> 70 years), we found that samples from cognitively normal older individuals clustered with the "healthy aging" subgroup while AD samples mainly clustered with the "AD similar" subgroups. This implies that individuals in the healthy aging subgroup will likely remain cognitively normal when they become older and vice versa. In summary, our results suggest that on the transcriptome level, aging and LOAD have strong interconnections in some brain regions in a subpopulation of cognitively normal aging individuals. This supports the theory that the initiation of LOAD occurs decades earlier than the manifestation of clinical phenotype and it may be essential to closely study the "normal brain aging" to identify the very early molecular events that may lead to LOAD development.

Keywords: Aging, LOAD, gene expression, brain transcriptome, brain aging subgroups, hippocampus, brain regions, Alzheimer's disease.

Introduction

Alzheimer's disease (AD) is the most common cause of dementia and about 6.2 million Americans live with the disease based on the Alzheimer's Association 2021 report (2021). Aging is the major risk factor for late-onset AD (LOAD), which represents over 95% of the AD cases that occurred at age 65 or older (Caselli et al., 2009). A study analyzing 1,246 subjects aged 30–95 years found that the risk of developing AD dramatically increases in APOE ε4 carriers who are 70 years or older (Jack et al., 2015). It has been well-recognized that normal brain aging and LOAD share multiple common features, e.g., aging brains often manifest certain degrees of cognitive impairment, memory loss, metabolic disturbances, bioenergetic deficits, and inflammation. Even though aging increases the risk of AD and the two processes share similarities in multiple aspects, a detailed brain-region-specific view of their interconnection at the molecular level is not fully available. It is also unclear which aging mechanisms are playing major contributions to AD development and why some individuals may age without major cognitive deficits while others develop AD (Koivisto et al., 1995; Herrup, 2010). To address these issues, we performed a global comparison of the transcriptomes from normal aging and LOAD brains across multiple regions in a hope to gain new insights into the molecular interconnection between aging and AD.

Despite the fact that many transcriptomic studies have been performed to investigate aging and LOAD independently, only a few studies have compared normal aging and LOAD transcriptomic data in a systematic way (Cribbs et al., 2012; Berchtold et al., 2013; Mastroeni et al., 2017; Lanke et al., 2018). For example, Berchtold et al. used microarray to profile 81 aging and AD brains and found that synapse-related genes showed progressive down-regulation in both aging and AD (Berchtold et al., 2013). Using the same dataset, Lanke et al. performed an integrated analysis on young, aging and AD brains by constructing co-expression network models, and found that modules associated with astrocytes, endothelial cells and microglial cells were up-regulated while modules associated with neurons, mitochondria and endoplasmic reticulum were down-regulated, and these modules significantly correlated with both AD and aging. All these studies greatly helped our

understanding of the interconnections between aging and AD; however, the previous studies

were limited in the brain regions examined and more importantly, they all treated aging and

AD samples as uniform groups, and did not consider the possible heterogeneity in either aging

or AD.

In this study, we compared the gene expression profiles of normal brain aging (age ≤ 70) with

LOAD (age ≥ 60) (Yang et al., 2015; Hodes and Buckholtz, 2016) to understand the similarity

and difference between aging and AD across multiple brain regions. We also considered brain

aging subgroups and compared the aging subgroups with LOAD.

Materials and Methods

Data collection and pre-processing

We compiled and processed multiple large-scale human brain aging and AD gene expression

datasets. We summarize and describe each dataset below and list all the data we studied in

Table 1.

GTEx brain data

GTEx brain gene expression data (v7 and v8) from 13 brain regions were downloaded from the

GTEx (Genotype-Tissue Expression) portal (Consortium, 2015) and NIH dbGaP database.

Donor ages ranged between 20-70 years and we removed samples from donors annotated with

any brain diseases from further analysis. We corrected gender, collection center, RIN (RNA

Integrity Number), PMI (postmortem interval), and top 3 genotype principal components (PCs)

to calculate gene expression associated with donors' chronological age.

UK brain data

Data from 134 post-mortem brain donors free of neurological conditions were obtained from

the UKBEC dataset (UK Brain Expression Consortium) (Millar et al., 2007; Trabzuni et al.,

2011). 10 brain regions were included, namely, cerebellum (CRBL, n = 131 samples), frontal

83 cortex (FCTX, n = 128), hippocampus (HIPP, n = 123), medulla (specifically inferior olivary

84 nucleus, MEDU, = 120), occipital cortex (specifically primary visual cortex, OCTX, n = 130),

85 putamen (PUTM, n= 130), substantia nigra (SNIG, n= 102), temporal cortex (TCTX, n = 120),

86 thalamus (THAL, n = 125) and intralobular white matter (WHMT, n = 132). To make the donor

87 age distribution comparable between GTEx and UK, we only considered samples from donors

88 whose age ≤ 70. The sample size for each brain region after filtering by age ≤ 70 is listed in

89 Table 1.

91 **Mount Sinai brain data**

92 364 human brains (238 females and 126 males) were accessed from the Mount Sinai/JJ Peters

93 VA Medical Center Brain Bank (MSBB–Mount Sinai NIH Neurobiobank) cohort. These

94 samples represented the full spectrum of cognitive and neuropathological disease severity in

95 the absence of discernable non-AD neuropathology. Donor ages for the samples ranged

96 61–108. For microarray profiling of 19 brain regions, a linear model was adopted to identify

97 genes differentially expressed among different disease stage groups using R package Limma

98 with default parameters and corrected for covariates including sex, postmortem interval (PMI),

99 pH, and race (Wang et al., 2016). For each brain regions, DEGs (Differential Expression

100 Genes) with FDR ≤ 0.05 in any of the 6 traits (CDR, Braak, CERAD, PLQ_Mn, NPrSum,

101 NTrSum in contrast of High vs. Low, Low vs. Normal or High vs. Normal) were combined as

102 the DEGs for that brain region (see Table S1)(Wang et al., 2016).

104 **Other brain data**

105 The AMP-AD knowledge portal (Hodes and Buckholtz, 2016) hosts datasets from three

106 studies: Mayo Clinic Brain Bank (Mayo), the Religious Orders Study and Memory and Aging

107 Project (ROSMAP) and Mount Sinai School of Medicine (MSSM). Mayo dataset included

108 temporal cortex (80 AD, 71 controls) and cerebellum (79 AD, 72 controls). ROSMAP dataset

109 contained samples from dorsolateral prefrontal cortex BA9 region (155 AD, 86 controls). The

110 MSSM cohort included samples from four brain regions: parahippocampal gyrus, inferior

111 frontal gyrus, superior temporal gyrus, and the frontal pole (n = 476, Table 1). DEGs were

112 downloaded from the portal and filtered for genes with FDR ≤ 0.05.

Jager AD gene signatures

De Jager et al. performed analyses on 478 ROSMAP dorsal lateral prefrontal cortex (DLPFC)

tissue samples (Mostafavi et al., 2018). 5 gene lists were considered containing genes whose

expression was associated with AD-related traits including clinical diagnosis of AD at the time

of death, cognitive decline, tau, amyloid, and pathologic diagnosis of AD.

Annse2018 and Rooij2019 AD gene expression signatures

Annse2018 profiled hippocampus CA1 gene expression in 10 males with age between 60 ~

81(5 AD and 5 controls) (Annese et al., 2018). DEGs with $|log2(Fold Change)| > 1$ and FDR $\leq$

0.05 were selected. Rooij2019 profiled gene expression of hippocampus samples from 18 AD

and 10 controls. DEGs with differential expression score $\geq$ 0.1 and FDR $\leq$ 0.05 were

considered (van Rooij et al., 2019).

Differential expression analysis and age-associated gene expression identification

Differential expression (DE) analysis in aging and AD was performed using the R package

edgeR and Limma (Robinson et al., 2010; Law et al., 2014) and we adjusted batch, RIN, sex,

and PMI in GTEx data and brain source, gender, batch effects and PMI in UK data. A linear

regression model was applied to identify gene expression changes associated with age and we

adjusted the same covariates as we did in the DE analysis (Yang et al., 2015; Zeng et al., 2020).

Subgroup identification using hierarchical clustering

We used hierarchical clustering method to identify subgroups in normal aging GTEx and UK

brain samples. We selected the top 5000 most variable genes which showed similar clustering

result with using all genes (the adjusted Rand index > 0.9) for the hierarchical clustering of

GTEx data. Ward.D2 method in the R hclust function was used (Murtagh and Legendre, 2014).

Deconvolution of GTEx bulk tissue gene expression data to infer cell type composition

The sorted single-cell RNAseq data (GSE125050, 4 cell types: NeuN+ neurons, GFAP+

astrocytes, CD31+ endothelial cells, and CD11b+ microglia/myeloid cells) in superior frontal

gyrus (SFG) from control samples with negligible amounts of amyloid and tau neuropathology

(56 cells, donor ages ranged 38-99 with mean 78.1 and standard deviation 15.5) was used as

reference (Srinivasan et al., 2020) to infer the cell-type composition of GTEx hippocampal

samples. According to a recently published deconvolution benchmark work (Avila Cobos et

al., 2020), we selected two top performing methods, namely the RLR (robust linear regression)

(Ripley et al., 2013) and MuSiC (Wang et al., 2019). We followed the recommended

processing procedures (no transformation and TMM normalization). For RLR method, we

used the "all marker selection strategy" (i.e., using all available markers). We used

non-parametric Kruskal-Wallis rank sum test (R function kruskal.test) for multiple group

comparison and Wilcox rank sum test for pairwise-comparison.

Comparison of PHG and GTEx hippocampal transcriptomes

We obtained gene expression data from 215 parahippocampal gyrus (PHG) samples which

were profiled at Mount Sinai (Wang et al., 2018). To compare the PHG and GTEx

hippocampal gene expression data, we first calculated $\log2(\text{tmp} + 1)$ for both datasets, merged

these datasets and then removed the batch effects using R ComBat package with AGE, PMI,

SEX, RIN as covariates. Negative values were assigned to 0 followed by sample-wide quantile

normalization. For the PHG data, we identified 22 normal (CDR = 0, braak score: bbscore $\leq$ 3,

CERAD = "NL") and 63 AD (CDR $\geq$ 1, bbscore $\geq$ 5, CERAD = "definite AD") samples. We

further selected the PHG samples with age > 70 and obtained 78 samples (19 normal, 59 AD).

Based on the GTEx top 5000 variance genes, we clustered PHG samples and found the normal

and AD samples were mixed to some degree. After removing the mixed samples, 51 samples

(14 normal and 37 AD samples) showed clear separation into two groups corresponding to

donors' AD status.

Functional enrichment analysis

We annotated the biological functions of each gene list using DVAID tool (Huang et al., 2009;

Sherman and Lempicki, 2009). We also performed pathway analysis using MetaCore

integrated software suite (https://portal.genego.com/; last accessed in November 2019) to

determine enriched biological processes. Signal transduction gene regulation network analysis

was based on SIGNOR 2.0 (Licata et al., 2020) from the SIGnaling Network Open Resource

(downloaded on Aug. 2020) using networkanalyst web tool (Zhou et al., 2019).

Results

1. A global comparison of aging and AD signatures across brain regions

1.1. Derive brain aging and AD signatures from multiple transcriptomic data

We collected multiple large-scale brain aging and AD gene expression datasets (Figure 1)

which are summarized in Tables 1 and S1. It is of note that to obtain "normal" aging gene

expression signatures, samples from donors annotated with any brain-related diseases were

removed in the GTEx data. We used a linear regression model to identify gene expression

associated with donors' chronological age and consider these genes as brain aging genes (we

call each list of brain aging genes as an aging signature for the corresponding brain region). We

derived aging signatures in 11 out of 13 brain regions from the GTEx data; two regions, i.e., the

brain spinal cord and substantia nigra showed no apparent age-associated genes (0 and 1 gene,

respectively) and were not considered for further analysis. Similarly, we derived aging

signatures from 8 out of the 10 brain regions profiled in UK data; substantia nigra and medulla

tissues showed no apparent aging signatures and were removed from further analysis.

Genes differentially expressed between AD and normal control samples were extracted from

previously published work (similarly we call them as AD signatures). These AD signatures are

MS AD sets from Mount Sinai Medical Center Brain Bank (MSBB) AD cohort (Wang et al.,

2016), and "Other" AD sets included AMP-AD knowledge portal data (Hodes and Buckholtz,

2016), Jager gene lists from the dorsal lateral prefrontal cortex (DLPFC) region (Mostafavi et

al., 2018), Annese2018 DEG list from hippocampal (HIPP) CA1 (denoted as Annese2018

HIPP)(Annese et al., 2018) and Rooij2019 DEG list in HIPP(van Rooij et al., 2019) (denoted

as Rooij2019 HIPP) (see Table 1 and Method).

1.2. A global comparison of aging and AD signatures across brain regions

We first identified genes that show similar gene expression regulation across multiple brain

regions in aging and AD datasets respectively (we call them global aging and AD signatures).

We found that 91 aging genes are consistently up or down-regulated in more than 4 out of 19

aging gene lists (11 GTEx and 8 UK brain regions). Among them, 27 are up-regulated and 64

are down-regulated (see Table S2) with age. Similarly, 86 AD signature genes show consistent

regulation in at least 9 out of 32 AD gene signature lists, among which 51 are consistently

up-regulated and 35 are consistently down-regulated in AD brains (see Table S2). We then

perform function annotation of these global aging and AD signatures.

1.2.1. The global aging and AD signatures show up-regulation of immune complement pathway genes and down-regulation of synaptic related signaling genes

We used the DAVID tool to annotate the functional enrichment and found the 91 aging genes

are enriched for synapse, nucleotide-binding and ion genes (see Table S3.1); while the 86 AD

signature genes are enriched for synapse, phosphoprotein and transport genes (see Table S3.4).

We further used MetaCore to annotate the function of consistently up- or down- regulated

aging and AD signature genes. The 27 up-regulated aging genes are enriched for regulation of

lipid metabolism, insulin regulation of glycogen metabolism (PHK gamma gene; FDR =

2.9E-2), immune/inflammation complement pathway (*C4B*, *C4*, *C4A*; FDR = 1.9E-06),

response to metal ion/transport categories (e.g., cellular response to copper ion; FDR =

1.7E-06), and dopamine metabolic process (*MAOB*, *MAO*; FDR= 3.8E-3) (see Table S3.2); the

64 down-regulated aging genes are enriched for *MAPK* (FDR = 1.2E-02) / *ASK1* (FDR =

2.4E-02) pathway, synaptic vesicle / calcium ion exocytosis categories, synaptic related

signaling (NMDA, glutamate) (see Table S3.3). The 51 up-regulated AD genes are involved in

immune complement pathway (GPCRs), TGF-beta receptor signaling (FDR = 4.6E-02),

cytoskeleton remodeling, and L-glutamate import/neurotransmitter transport (see Table S3.5);

35 down-regulated AD genes are involved in synapse and GABAergic neurotransmission,

glutamate secretion/glutamatergic pathways, vesicle fusion/recycling and transport,

cytoskeleton remodeling, cell adhesion and protein phosphorylation (see Table S3.6).

Therefore, both AD and aging global signature genes show up-regulation in immune

complement pathway and down-regulation in synapse (especially glutamate, synaptic vesicle

recycling) related pathways; while aging genes are more involved in metal ion, transport,

glycogen metabolism and AD signature genes are more involved in cytoskeleton remodeling,

cell adhesion and synaptic categories.

1.2.2. *SST* and *SVOP* are down-regulated and *FOXJ1*, *SLC44A1* are up-regulated in the

global aging and AD signatures

Two genes, *SST* and *SVOP* are consistently down-regulated in both global aging and AD gene

signatures, while *FOXJ1* and *SLC44A1* are consistently up-regulated.

SST (somatostatin) is a neuropeptide hormone that maintains permeability and integrity of the

blood-brain barrier (BBB) by regulating *LRP1* and *RAGE* expression. It abrogates Aβ-induced

JNK phosphorylation and expression of *MMP2* to maintain permeability and integrity of BBB

(Paik et al., 2019). Reduction of SST levels in the CSF and brain tissue is associated with

impaired cognitive function and memory loss (Solarski et al., 2018). *SVOP* (synaptic vesicle

2-related protein) can bind to adenine nucleotide (particularly NAD) via its C-terminal

extremity(Yao and Bajjalieh, 2009). Although the functions of *SVOP* remain obscure, the

evolutionary conservation and homology to transporters support it may play a role in molecular

trafficking in synaptic vesicles(Janz et al., 1998).

In mammalian cells, *FOXJ1* is a member of the Forkhead/winged helix (FOX) family of

transcription factors that is involved in ciliogenesis (Yu et al., 2008). It is shown to suppress

NFκB, a key regulator in the immune response (Lin et al., 2004). It is hypothesized that *FOXJ1*

may play a protective role involved in the pathophysiology of brain injury and may be required

for the differentiation of the cells (Jacquet et al., 2009). The solute carrier 44A1 (*SLC44A1*) is a

plasma membrane choline transporter. It is also a mitochondrial protein and acts as a choline

transmembrane transporter. Choline is essential for the synthesis of the neurotransmitter ACh

by cholinergic neurons for regulating neuronal activity. Choline deprivation in the central

nervous system reduces acetylcholine (ACh) release, memory retention, and spatial cognition

in the hippocampus (Canty and Zeisel, 1994; Nakamura et al., 2001; Zeisel, 2007).

2. Brain region-specific comparison of aging and AD signatures

In addition to comparing the global aging and AD signatures, we also compared aging and AD signatures in a brain region-specific manner.

2.1. Hypothalamus, hippocampus, and certain cortical regions show stronger age-related gene expression changes compared to other brain regions

As can be seen in Figure S1, the number of aging genes varies across brain regions. The regions showing a large number of aging genes in the GTEx data are hypothalamus (HYPO: 490 UP and 1160 DN genes), hippocampus (HIPP: 705 UP and 698 DN genes), and brain anterior cingulate cortex BA24 (BA24_AC: 320 UP and 888 DN genes). For the UK data, the brain regions showing a large number of aging genes are hippocampus (HIPP:432 UP and 324 DN genes) and temporal cortex (TCTX: 91 UP and 419 DN genes). It is of note that GTEx and UK do not cover the same brain regions, e.g., the hypothalamus was profiled by GTEx but not in the UK brain data. Among the 4 regions profiled by both GTEx and UK, (i.e., HIPP, CRBL, PUTM, FCTX), HIPP and CRBL show a higher number of aging genes and strong overlap between GTEx and UK aging signatures than the other two brain regions. Interestingly, the hippocampus aging signature significantly overlaps with aging signatures from several cortical regions, while the cerebellum aging signature has very little overlap with other regions. The cerebellum show the most distinguishable gene expression patterns from all other brain regions which was also reported by the Allen Brain Atlases study (Mahfouz et al., 2015).

In summary, brain aging signatures are highly region-specific. Hippocampus, hypothalamus, and cortex TCTX and BA24_AC are more affected by aging on the transcriptome level than other brain regions surveyed by the GTEx and UK brain data.

276
2.2. Common hippocampus aging genes between GTEx and UK data

In general, the aging signatures from GTEx and UK show large reproducibility in matched brain regions. Using hippocampus as an example, the 705 up-regulated GTEx aging genes significantly overlap with 431 up-regulated UK aging genes by 77 genes (FDR = 2.6E-31); and the 698 down-regulated GTEx aging genes overlap with 324 down-regulated UK aging genes

by 42 genes (FDR = 3.6E-12). The 119 commonly up- or down-regulated hippocampal aging

genes between GTEx and UK are enriched for phosphoprotein, alternative splicing,

acetylation, complement, and synapse (see Table S4.1). In addition to the immune /

inflammation categories found in previous global aging signatures, the 77 commonly

up-regulated aging genes are also enriched for *TGF-β* signal, transcription regulation (such as

mRNA transcription by RNA polymerase II with FDR of 2.2E-03, ATP-dependent chromatin

remodeling with FDR of 7.3E-03, blood vessel remodeling, membrane protein intracellular

domain proteolysis, protein transport, and metabolic process (see Table S4.2). Furthermore,

from signal transduction network analysis (see Table S4.4 and S4.5), the top signal network

contains 67 up-regulated genes and is connected with many metabolic-related pathways such

as endocrine resistance (FDR = 4.0E-15), *foxo* signaling pathway (FDR = 1.4E-13), thyroid

hormone signaling pathway (FDR = 8.7E-12), sphingolipid signaling pathway (FDR =

1.8E-10), neurotrophin signaling pathway (FDR = 1.8E-10), autophagy – animal (FDR =

3.6E-10), and insulin resistance (FDR = 1.5E-09).

The 42 commonly down-regulated aging genes between GTEx and UK are related to

insulin-like growth factor 1 (*IGF-1*) signaling, *ERK1/2* pathway (for caveolin-mediated

endocytosis or signal transduction), methylation, cell adhesion (e.g., synaptic contact), and

long-term potentiation pathways and previous reported energy-dependent pathways such as

synaptic categories (e.g. synapse vesicle), oxidative categories (see Table S4.3) based on

MetaCore analysis. *IGF-1* (insulin-like growth factor) is a growth factor and neurohormone

with some evidence suggesting its involvement in neurocognitive functions,

neuroinflammation, and amyloid-β clearance. Furthermore, from signal transduction network

analysis on the KEGG database (see Tables S4.6 and S4.7), the top signal network is involved

in circadian entrainment (FDR = 9.2E-07) and multiple synaptic-related pathways.

2.3. Brain aging signatures from the hypothalamus, the hippocampus, and BA24 anterior

cingulate cortex significantly overlap with AD signatures

After we evaluated the aging signatures between GTEx and UK data, we then compared the

aging signatures with AD signatures in a brain-region specific manner. We further collected

two hippocampal DEG lists (Rooij2019 and Annese2018, listed in "Other" AD sets) for the

following analysis. As can be seen in Figure 2, strong overlap between aging and AD

signatures are observed in hippocampus, hypothalamus, and several cortical regions. For

example, the GTEx hippocampus aging signature strongly overlaps with AD signatures

derived from either hippocampus or several cortex regions. For the 698 GTEx_HIPP_DN

aging genes, they overlap with Rooj2019_HIPP_DN by 288 genes (adjusted P-value =

2.78E-130); and the 705 GTEx_HIPP_UP genes overlap with Mayo_TCTX _UP (1578) by

150 genes (adjusted P-value = 6.62E-28). It is of note that as we divide aging and AD

signatures into up- and down-regulated genes, in most cases, the down-regulated aging genes

only significantly overlap with down-regulated AD genes, and vice versa, which supports that

these signatures represent real biological signals. The overlap pattern is highly brain region

specific. For example, GTEx_HIPP_UP (705) significantly overlap with AD signatures from

hippocampus, while it only overlaps with AD signatures from non-hippocampal regions such

as MS_BM44_IFG_UP (513) by 21 genes (adjusted P-value = 0.72). In both GTEx and UK

data, the aging signature in cerebellum shows relatively weak overlap with our AD signatures.

We also observed that the GTEx hippocampus and UK hippocampus aging signatures show

similar overlap pattern with AD signatures across different brain regions. Similarly, different

AD signatures such as Annesse2018 and Rooj2019 showed similar overlap pattern with aging

signatures across different brain regions, which further suggests that the aging and AD

signatures from different studies are biologically meaningful and contain real signals of aging

and AD.

2.4. Functional enrichment of aging specific, aging/AD common and AD specific genes in

hippocampus

Since aging and AD gene signatures show strong overlap in the hippocampus as seen in Figure

2, we focus on this brain region to investigate the functional enrichment of aging and AD

genes. To achieve better robustness, we annotated hippocampus aging genes derived from both

GTEx and UK with hippocampus AD genes derived from both Annese2018 and Rooij2019

(Figure 3). We divided genes in the aging and AD signatures into three categories: aging

specific signature genes (denoted as "ASGs"), aging-AD common signature genes (denoted as

"AADGs"), and AD specific signature genes (denoted as "ADSGs"). Phosphoprotein and

alternative splicing are enriched in ASGs, AADGs and ADSGs (also in both up- and

down-regulated genes); while glycosylation / glycoprotein, immune / inflammatory, stress

response, mitochondria are more enriched in ADSGs; acetylation, nucleus, and Ubl

conjugation are mostly enriched in ASGs (see Table S5.1). Down-regulated process of ASGs,

AADGs, and ADSGs are all associated with membrane/cytoskeleton categories.

Down-regulated AADGs are enriched for synapses (including cell junction,

cholinergic/GABAergic/dopaminergic/glutamatergic synapse, AMPA etc.) and ion/transport

(Figure 3 and Table S5); The comparison suggests that different from the normal aging

process, the AD-specific and aging/AD commonly affected genes are more enriched for

glycoprotein, inflammatory response, synapse and mitochondrial dysfunction, while the aging

process are more related to the nucleus, acetylation, coiled coil, which are less directly related

to neurodegenerative phenotype. The functional annotation of AADGs suggests that

dysregulation in transcription regulation, energy metabolism, membrane remodeling,

extracellular vesicles (EV) and synapse pathways have already initiated and developed to some

degrees in normal brain aging even though these individuals remain cognitively normal, while

other biological processes such as inflammation/immune response are further escalated in AD

patients.

3. Subgroup identification in normal aging brain hippocampus samples

3.1 Subgroups can be found in both GTEx and UK hippocampus datasets

Although we have shown that aging and AD share multiple gene expression changes in the

hippocampus, it is well known that not every old individual will develop AD. This suggests the

interconnection between normal aging and AD may occur at different intensity within the

aging population. To examine the potential heterogeneity of normal brain aging, we explored

subgrouping in both GTEx and UK hippocampus samples; and we observed that both GTEx

and UK samples can be divided into major subgroups. As shown in Figure 4, 56 GTEx

hippocampal samples with donors' age between 45 and 70 can form three major clusters

(named subgroups A, B and C). We noticed that samples from different age groups are

relatively evenly distributed across these subgroups, suggesting that subgrouping is not due to

difference in donor ages. Similar subgrouping is also observed in UK hippocampal data

(Figure 4B).

3.2 DEGs derived from comparing aging brain subgroups highly overlap with AD

signatures

We performed pair-wise comparisons to derive DEGs between subgroups in both GTEx and

UK datasets, respectively. We jointly considered these DEGs (FDR < 0.01) with 3

hippocampal AD signatures (MS, Rooij2019, and Annese2018). As shown in Figure 5, DEGs

derived from comparing GTEx subgroups B vs A (denoted as GTEx DEG BvsA) show strong

overlap with AD signatures from Rooij2019 and Annese2018. For the GTEx_DEG_BvsA

down-regulated genes (3018 genes), they overlap with Rooij2019 down-regulated genes (1622

genes) by 1274 genes while they only overlap with Rooij2019 up-regulated genes (1045 genes)

by 7 genes. This is highly significant, as Rooij2019 down-regulated genes (1622 genes)

overlap with another AD signature, Annese2018 down-regulated genes (1183 genes) by 614

genes, which is comparable to the overlap we see with aging subgroup DEGs. It is of note that

GTEx samples used in the subgroup analysis are from donors 45-70 years old without AD or

other types of brain diseases. Simply by performing subgrouping, the DEGs from comparing

these subgroups highly overlap with AD signatures from independent studies, which further

supports that gene expression changes in a subpopulation of the cognitively normal brains have

very strong interconnection with AD. Similarly, GTEx DEGs CvsA also strongly overlap with

AD signatures. Based on the pattern of overlap, we infer that GTEx subgroup A is more likely

to be a "healthy" aging subgroup while GTEx subgroups B and C are more similar to AD (here

denoted as "AD similar group"). Since GTEx DEG CvsB showed reverse overlap with AD

signatures from Annese2018 and Rooij2019, this indicates that subgroup B is more similarity

with AD samples compared to subgroup C. Therefore, the order of subgroups judged by how

similar they are with AD samples can be inferred as AD samples > subgroup B > subgroup C >

subgroup A, which is also consistent with the observation that GTEx DEG CvsA overlap with

GTEx DEG BvsA in the same gene regulation direction. Similarly, for the UK data, we also

observed very strong overlap between DEGs from comparing various subgroups and AD

signatures. For example, the UK_DEG_AvsC_DN (3144 genes) highly overlap with

Rooij2019_DN signature (1622 genes) by 762 genes. Similarly, we inferred the order of UK

subgroups judged by how similar they are with AD signatures as AD > UK subgroup A >> UK

subgroup B > UK subgroup C.

3.3 Function annotation of GTEx and UK subgroup DEGs with AD signatures

We next used the DAVID tool to annotate the functional enrichment for GTEx and UK

subgroup DEGs (Figure 6, Table S6). We focused on the DEGs (FDR < 0.01) derived from

comparing GTEx subgroup B (a subgroup that is more similar to AD) with the relatively

healthy aging subgroup A (named GTEx DEGs BvsA) and similarly for UK DEGs AvsC with

two AD signatures (Annese2018 and Rooij2019). Similar to previous annotations, we denote

the aging subgroup DEGs as ABGs and annotate the functional enrichment of ABSGs (aging

subgroup DEG specific genes), ADSGs, and ABADGs (common genes between aging

subgroup DEGs and AD signature genes). As shown in Table S6.1 and Figure 6, certain post

translational modifications (PTMs) (e.g., phosphoprotein, alternative splicing, Ubl

conjugation, acetylation), membrane and cytoskeleton categories are significantly enriched in

ABSGs, ADSGs, ABADGs and also enriched in both up- and down-regulated genes,

respectively. ADSGs (both up- and down-regulation) are enriched for glycoprotein, while

up-regulated ADSGs are more enriched for immunity, inflammatory response categories and

down-regulated ADSGs are enriched for calcium/ion categories. The up-regulated conserved

ABSGs show enrichment in cell cycle category and down-regulated conserved ABSGs show

enrichment in proteostasis such as proteasome, protein transport and protein binding. For

ABADGs, up-regulated genes are mainly enriched for transcription regulation and metabolism

categories (e.g. PI3K-Akt signaling pathway, MAPK signaling pathway, insulin resistance),

while down-regulated genes are mainly enriched for synapse, lipoprotein, and circadian

entrainment categories.

In summary, aging subgroup DEG functional analysis further suggests that remodeling of

PTMs (e.g., phosphoprotein and alternative splicing), proteostasis, cytoskeleton, and

metabolism are likely initiated in normal aging subgroups that show highly similar gene

expression changes with AD patients.

3.4 Differentially expressed genes between subgroups could be driven by difference in the

cellular compositions

The differentially expressed genes we observed between subgroups could be caused by either

the dysregulation of gene expressions or the change in the cellular compositions or both. Since

both GTEx and UK were bulk tissue gene expression data, the cellular composition

information was not available. To evaluate how likely the DEGs between the subgroups were

caused by different cellular compositions in the samples between subgroups, we used

computational tools to perform cell deconvolution. Based on a recently published

deconvolution benchmark work (Avila Cobos et al., 2020), we chose two top performing

methods, namely the RLR (robust linear regression)(Ripley et al., 2013) and MuSiC (Wang et

al., 2019) and followed the recommended data processing procedures (TMM normalization

and using all available markers) to perform cell deconvolution of the GTEx hippocampal data.

A recent sorted single-cell RNAseq dataset (Srinivasan et al., 2020) was chosen as the cell-type

specific gene expression reference data (see Methods). We obtained highly consistent results

from the RLR and MuSiC methods (the mean correlation between the estimated cell

proportions reported from RLR and MuSiC was 0.979). The neuron proportion is significantly

decreased in the AD similar subgroups; e.g., the MuSiC method reported that the mean and

standard deviation of the proportion of neurons in subgroup A is 0.498 ± 0.10, 0.221 ± 0.09 for

subgroup B, and 0.294 ± 0.11 for subgroup C; the difference between subgroups A and B is

significant with p-value = 1.3E-05, similarly the p-value for the difference between subgroups

A and C is 4.4E-05. The endothelial cell proportion is estimated to be significantly increased in

the AD similar subgroups; e.g., the MuSiC reported that the mean and standard deviation of

endothelial cell proportion in subgroup A is 0.407 ± 0.11, 0.695 ± 0.05 for subgroup B, and

0.566 ± 0.11 for subgroup C; the difference is significant between subgroups A and B with

p-value = 6.9E-06, and the p-value for the difference between subgroups A and C is 3.8E-04

(see Figure S2 A). Similar results were also observed from calculations using the RLR method

(see Figure S2 B).

3.5 GTEx "healthy aging" subgroup will likely remain cognitively normal in older ages as implied from joint analysis with another brain transcriptomic dataset

In both GTEx and UK datasets, we observed a subgroup that is quite different from AD and subgroups that are more similar to AD on the transcriptome level. We hypothesize that the subgroup less similar to AD is a "healthy aging" subgroup which will have a better chance of being cognitively normal if they could live into 70s or older. In contrast, the "AD similar" subgroups will likely have a higher chance of having cognitive deficits if these individual could live into 70s or older. To test this hypothesis, a longitudinal study that follows-up these subgroup individuals for several decades will be required. This is not feasible mainly because brain tissues are mostly available from postmortem donors which will not permit longitudinal studies. To evaluate the hypothesis, we have to rely on alternative approaches.

To indirectly test the hypothesis, we relied on another large brain transcriptomic dataset which profiled more than 200 brain tissues from the parahippocampal gyrus (PHG) region (Wang et al., 2018). Although PHG is a different brain region, it is next to the hippocampus and our analysis showed that the two transcriptomic datasets are comparable. From these PHG samples, we selected two groups of samples, i.e., control and AD samples. The normal control samples were chosen from donors with age > 70, CDR = 0, braak score (bbscore) $\leq$ 3, CERAD = 'NL' which has 19 samples. For this group, the donor ages ranged from 73 to 103, with mean and standard deviation of 83.3 $\pm$ 8.7 (yrs). The second group is the AD group, which we required donors to be > 70 yrs, CDR $\geq$ 1, bbscore $\geq$ 5, CERAD = 'definite AD'. 59 samples met these criteria, and the donor ages ranged from 71 to 104, with mean and standard deviation of 86.966 $\pm$ 8.4 (yrs). We merged these PHG data with the GTEx hippocampus RNA-seq data and corrected the batch effect using the R ComBat package (see method) to form a unified dataset. The joint processing introduced some changes to the GTEx gene expression. Therefore, we performed the hierarchical clustering on the 56 GTEx samples again and observed that the new subgrouping structure (subgroup B [n=11] > C [n=10] > A [n=35]) is highly similar to the previous clustering results (subgroup B' [n=9] > C' [n=11] > A' [n=36]). For example, the new

subgroup A (35 samples) overlaps with the previous subgroup A' (36 samples) by 32 samples.

We performed hierarchical clustering on the 78 PHG samples and found several normal and

AD samples were mixed together (see Figure 7A). To obtain a clear separation between control

and AD samples, we removed the mixed samples and obtained 14 normal (ages 73 ~ 103, 5

males and 9 females) and 37 AD samples (ages 73 ~ 104, 11 males and 26 females) which

formed two fully separated groups (see Figure 7B). We then assigned these PHG samples to the

GTEx subgroups by adding one PHG sample at a time to the 56 GTEx samples for hierarchical

clustering. We found that all the 14 control samples clustered with the GTEx "healthy aging"

subgroup A. For the 37 PHG AD samples, 5 (3 APOE ε3/ε4 and 2 APOE ε3/ε3) clustered with

GTEx subgroup B and 26 clustered with GTEx subgroup C (the two "AD similar" subgroups),

only 6 samples (all APOE ε3/ε3) clustered with the "healthy aging" subgroup A. This result

suggests that the gene expression of the "healthy aging" subgroup is mainly associated with

cognitively normal status in older individuals while the gene expression of the "AD similar"

subgroup is largely associated with AD in this older cohort.

Discussion

To better understand the interconnection between aging and LOAD, we collected gene

expression profiles from several large-scale transcriptomic datasets covering multiple brain

regions and systematically compared aging and LOAD gene expression signatures. Different

brain regions show varied levels of gene expression changes in aging and AD, respectively.

Among all the brain regions we studied, hippocampus is one of the top regions that show a very

strong interconnection between aging and AD on the transcriptomic level. We observed

common functional enrichment in aging and LOAD related to PTMs (especially for alternative

splicing and phosphoprotein), neurotransmission (especially glutamate), membrane,

cytoskeleton, and lipid metabolism. We also showed that gene expression changes in aging

brain has more association with acetylation while AD-specific gene expression changes are

more related to inflammatory response, glycoprotein, mitochondria and synapse. Importantly,

we demonstrated that cognitively normal brains are not homogeneous in their transcriptomes

and several major subgroups could be identified. By comparing gene expression among

different subgroups, we showed that gene expression changes in a subpopulation have strong

overlap with the AD signature, suggesting that although aging is a general risk factor for AD,

only a subset of individuals may experience gene expression changes to a level that

significantly facilitates the onset of the disease.

From our study, several biological processes shared between LOAD and normal aging could

point to the mechanisms of how aging and AD interconnect. First, PTMs related genes such as

alternative splicing and phosphoprotein are enriched in both up- and down-regulated ASGs,

AADGs/ABADGs and ADSGs (e.g., top function terms in Figures 3 and 6). PTMs regulation

plays critical role for synaptic plasticity at several levels (Kiltschewskij and Cairns, 2017) by

increasing proteome diversity through alternative splicing, or by enabling activity-dependent

regulation of mRNA localization, translation or degradation in the dendrite. It will be

interesting to investigate what events trigger the PTMs that are observed in aging and AD.

Second, the immune response strongly showed-up in AD signatures as can be seen in our

results and previous studies (Verbitsky et al., 2004; Reichwald et al., 2009; Bordner et al.,

2011; Cribbs et al., 2012). Since it was less significantly up-regulated in either ASGs/AADGs

or ABSGs/ABADGs, this supports that the escalated inflammation is a hallmark of AD,

relative to normal brain aging which is also known to be associated with low-grade

neuroinflammation. Whether the inflammatory response plays a key causative role in the early

stage of AD development or it is more of a reactive response to the upstream pathology

requires further investigation.

Since it is infeasible to directly test if individuals in the "healthy aging" subgroup A will

remain cognitively normal decades later, we relied on an alternative strategy in which we

compared transcriptomes between GTEx brain aging subgroups with a cohort of older

individuals who were either AD or cognitively normal. For the 51 PHG samples (14 control

and 37 AD samples), we observed that 31 out of 37 AD samples clustered with GTEx

subgroups B and C (AD similar subgroups); while all the 14 normal control samples clustered

with GTEx subgroup A, which presumably represents a healthy brain aging group. This result

suggests that the gene expression in subgroup A is strongly associated with normal cognitive

functions in both relatively young and older individuals, which supports that GTEx subgroup A

is a healthy aging subgroup and individuals in this subgroup will likely remain cognitively

normal when they get older. However, we did observe that several AD patients have their PHG

transcriptome clustered with GTEx subgroup A (6 out of 37 AD samples). This indicates that

an individual with a healthy aging gene expression pattern in the hippocampus or PHG can still

have AD. We think this could be explained by the heterogeneity of AD which is increasingly

known to have multiple subtypes. For example, at least four subtypes of AD (typical,

limbic-predominant, hippocampal-sparing, and minimal atrophy AD) have been reported

based on distribution of tau related pathology and regional brain atrophy (Ferreira et al., 2020).

It is possible that some subtypes of AD could have their hippocampal transcriptomes similar to

our healthy aging subgroup. For example, it has been reported that hippocampal-sparing AD

subtype has a lower frequency of APOE ε4 compared with typical and limbic-predominant AD

(Ferreira et al., 2020); interestingly, all the 6 AD samples clustered with the GTEx subgroup A

are from donors of APOE ε3/ε3. Since AD is not a single brain-region disease, to accurately

predict AD development based on gene expression or any type of brain region-specific data, we

believe that multiple regions should be examined and studied together.

The results from cell decomposition of the bulk gene expression data suggest that it is possible

that the differentially expressed gene expression between subtypes is at least partially due to

the changes in cell proportions across the subgroups. For example, the down-regulation of

synapse genes in the GTEx aging subgroup B might be explained by the decrease in the number

of neurons in the samples in this subgroup. However, the deconvolution methods we used all

assumed that the reference gene expressions do not change across conditions (in our case, the

aging subgroups), which may not necessarily be true. Therefore, although it is very likely that

cell compositions changed across subtypes, it is also possible that some gene expressions

changed in a cell-type specific manner across the aging subgroups. To fully resolve this issue,

single-cell profiling of samples from these aging subgroups will be highly useful.

Finally, we observed no significant difference in sex distribution among GTEx subgroups

(Fisher exact test, p-value = 0.81) and UK subgroups (p-value = 0.37). For example, for the 36

samples in GTEx subgroup A, 27 were males, and 9 were females; for the 9 samples in GTEx

subgroup B, 8 were males and 1 was female; and for GTEx subgroup C, 8 were males and 3

were females. Since there were only very limited number of female samples in the AD similar

subgroups, we did not have sufficient statistical power to investigate the sex-related difference

in these aging subgroups.

In summary, our study suggests that combined analysis of aging and LOAD can help us to

understand how aging may contribute to the development of LOAD. Since most genomic

studies on AD relied on AD samples from donors in 70s to 90s, these samples may not provide

the information for the molecular events occurred in the very early stage of the disease

development. As we demonstrated in this work, the brains from a subpopulation of cognitively

normal individuals in their 40s-70s already show gene expression changes similar to AD, this

supports that the initiation of LOAD could occur decades earlier than the manifestation of

clinical phenotypes and it may be critical to closely study cognitively normal individuals in

their 40s-60s to identify the triggering events in AD development.

Acknowledgements

This work is supported by NIH grants R01AG055501, R01AG067312 to Z.T. and D.M.H. and

R01AG057907 to B.Z. The content is solely the responsibility of the authors and does not

represent the official views of the National Institutes of Health. This work was also supported

in part through the computational resources and staff expertise provided by Scientific

Computing at the Icahn School of Medicine at Mount Sinai.

Author Contributions

ZT conceived the main concept of the work and BZ proposed the comparison with PHG data

for "validation". ZT and SP contributed to conceptualization, methodology, analysis, and

preparation of the manuscript. SP, LZ, JH, MW, DH, VH, ME, BZ helped to discuss and

improve the work. All authors have read and approved the manuscript for submission.

Competing interests

No competing interests declared.
607

Reference

(2021). 2021 Alzheimer's disease facts and figures. *Alzheimers Dement* 17(3)**,** 327-406. doi:
10.1002/alz.12328.
Annese, A., Manzari, C., Lionetti, C., Picardi, E., Horner, D.S., Chiara, M., et al. (2018).
Whole transcriptome profiling of Late-Onset Alzheimer's Disease patients provides
insights into the molecular changes involved in the disease. *Sci Rep* 8(1)**,** 4282. doi:
10.1038/s41598-018-22701-2.
Avila Cobos, F., Alquicira-Hernandez, J., Powell, J.E., Mestdagh, P., and De Preter, K. (2020).
Benchmarking of cell type deconvolution pipelines for transcriptomics data. *Nat*
Commun 11(1)**,** 5650. doi: 10.1038/s41467-020-19015-1.
Berchtold, N.C., Coleman, P.D., Cribbs, D.H., Rogers, J., Gillen, D.L., and Cotman, C.W.
(2013). Synaptic genes are extensively downregulated across multiple brain regions in
normal human aging and Alzheimer's disease. *Neurobiology of Aging* 34(6)**,**
1653-1661. doi: 10.1016/j.neurobiolaging.2012.11.024 PMID - 23273601.
Bordner, K.A., Kitchen, R.R., Carlyle, B., George, E.D., Mahajan, M.C., Mane, S.M., et al.
(2011). Parallel declines in cognition, motivation, and locomotion in aging mice:
association with immune gene upregulation in the medial prefrontal cortex. *Exp*
Gerontol 46(8)**,** 643-659. doi: 10.1016/j.exger.2011.03.003.
Canty, D.J., and Zeisel, S.H. (1994). Lecithin and choline in human health and disease. *Nutr*
Rev 52(10)**,** 327-339. doi: 10.1111/j.1753-4887.1994.tb01357.x.
Caselli, R.J., Dueck, A.C., Osborne, D., Sabbagh, M.N., Connor, D.J., Ahern, G.L., et al.
(2009). Longitudinal modeling of age-related memory decline and the APOE epsilon4
effect. *N Engl J Med* 361(3)**,** 255-263. doi: 10.1056/NEJMoa0809437.
Consortium, G.T. (2015). Human genomics. The Genotype-Tissue Expression (GTEx) pilot
analysis: multitissue gene regulation in humans. *Science* 348(6235)**,** 648-660. doi:
10.1126/science.1262110.
Cribbs, D.H., Berchtold, N.C., Perreau, V., Coleman, P.D., Rogers, J., Tenner, A.J., et al.
(2012). Extensive innate immune gene activation accompanies brain aging, increasing
vulnerability to cognitive decline and neurodegeneration: a microarray study. *J*
Neuroinflammation 9**,** 179. doi: 10.1186/1742-2094-9-179.
Ferreira, D., Nordberg, A., and Westman, E. (2020). Biological subtypes of Alzheimer disease:
A systematic review and meta-analysis. *Neurology* 94(10)**,** 436-448. doi:
10.1212/WNL.0000000000009058.

Herrup, K. (2010). Reimagining Alzheimer's disease--an age-based hypothesis. *J Neurosci*
30(50)**,** 16755-16762. doi: 10.1523/JNEUROSCI.4521-10.2010.

Hodes, R.J., and Buckholtz, N. (2016). Accelerating Medicines Partnership: Alzheimer's
Disease (AMP-AD) Knowledge Portal Aids Alzheimer's Drug Discovery through
Open Data Sharing. *Expert Opin Ther Targets* 20(4)**,** 389-391. doi:
10.1517/14728222.2016.1135132.

Huang, D.W., Sherman, B.T., and Lempicki, R.A. (2009). Bioinformatics enrichment tools:
paths toward the comprehensive functional analysis of large gene lists. *Nucleic acids*
research 37(1)**,** 1-13.

Jack, C.R., Jr., Wiste, H.J., Weigand, S.D., Knopman, D.S., Vemuri, P., Mielke, M.M., et al.
(2015). Age, Sex, and APOE epsilon4 Effects on Memory, Brain Structure, and
beta-Amyloid Across the Adult Life Span. *JAMA Neurol* 72(5)**,** 511-519. doi:
10.1001/jamaneurol.2014.4821.

Jacquet, B.V., Salinas-Mondragon, R., Liang, H., Therit, B., Buie, J.D., Dykstra, M., et al.
(2009). FoxJ1-dependent gene expression is required for differentiation of radial glia
into ependymal cells and a subset of astrocytes in the postnatal brain. *Development*
136(23)**,** 4021-4031.

Janz, R., Hofmann, K., and Sudhof, T.C. (1998). SVOP, an evolutionarily conserved synaptic
vesicle protein, suggests novel transport functions of synaptic vesicles. *J Neurosci*
18(22)**,** 9269-9281.

Kiltschewskij, D., and Cairns, M.J. (2017). Post-Transcriptional Mechanisms of Neuronal
Translational Control in Synaptic Plasticity. *Synaptic Plasticity***,** 15.

Koivisto, K., Reinikainen, K.J., Hanninen, T., Vanhanen, M., Helkala, E.L., Mykkanen, L., et
al. (1995). Prevalence of age-associated memory impairment in a randomly selected
population from eastern Finland. *Neurology* 45(4)**,** 741-747. doi:
10.1212/wnl.45.4.741.

Lanke, V., Moolamalla, S.T.R., Roy, D., and Vinod, P.K. (2018). Integrative Analysis of
Hippocampus Gene Expression Profiles Identifies Network Alterations in Aging and
Alzheimer's Disease. *Front Aging Neurosci* 10**,** 153. doi: 10.3389/fnagi.2018.00153.

Law, C.W., Chen, Y., Shi, W., and Smyth, G.K. (2014). voom: Precision weights unlock linear
model analysis tools for RNA-seq read counts. *Genome Biol* 15(2)**,** R29. doi:
10.1186/gb-2014-15-2-r29.

Licata, L., Lo Surdo, P., Iannuccelli, M., Palma, A., Micarelli, E., Perfetto, L., et al. (2020).
SIGNOR 2.0, the SIGnaling Network Open Resource 2.0: 2019 update. *Nucleic Acids*
Res 48(D1)**,** D504-D510. doi: 10.1093/nar/gkz949.

Lin, L., Spoor, M.S., Gerth, A.J., Brody, S.L., and Peng, S.L. (2004). Modulation of Th1
activation and inflammation by the NF-kappaB repressor Foxj1. *Science* 303(5660)**,**
1017-1020. doi: 10.1126/science.1093889.

Mahfouz, A., van de Giessen, M., van der Maaten, L., Huisman, S., Reinders, M., Hawrylycz,
M.J., et al. (2015). Visualizing the spatial gene expression organization in the brain
through non-linear similarity embeddings. *Methods* 73**,** 79-89.

Mastroeni, D., Khdour, O.M., Delvaux, E., Nolz, J., Olsen, G., Berchtold, N., et al. (2017).
Nuclear but not mitochondrial-encoded oxidative phosphorylation genes are altered in

aging, mild cognitive impairment, and Alzheimer's disease. *Alzheimers Dement* 13(5)**,**
510-519. doi: 10.1016/j.jalz.2016.09.003.

Millar, T., Walker, R., Arango, J.C., Ironside, J.W., Harrison, D.J., MacIntyre, D.J., et al.
(2007). Tissue and organ donation for research in forensic pathology: the MRC Sudden
Death Brain and Tissue Bank. *J Pathol* 213(4)**,** 369-375. doi: 10.1002/path.2247.

Mostafavi, S., Gaiteri, C., Sullivan, S.E., White, C.C., Tasaki, S., Xu, J., et al. (2018). A
molecular network of the aging human brain provides insights into the pathology and
cognitive decline of Alzheimer's disease. *Nat Neurosci* 21(6)**,** 811-819. doi:
10.1038/s41593-018-0154-9.

Murtagh, F., and Legendre, P. (2014). Ward's hierarchical agglomerative clustering method:
which algorithms implement Ward's criterion? *Journal of classification* 31(3)**,**
274-295.

Nakamura, S., Murayama, N., Noshita, T., Annoura, H., and Ohno, T. (2001). Progressive
brain dysfunction following intracerebroventricular infusion of beta(1-42)-amyloid
peptide. *Brain Res* 912(2)**,** 128-136. doi: 10.1016/s0006-8993(01)02704-4.

Paik, S., Somvanshi, R.K., and Kumar, U. (2019). Somatostatin Maintains Permeability and
Integrity of Blood-Brain Barrier in beta-Amyloid Induced Toxicity. *Mol Neurobiol*
56(1)**,** 292-306. doi: 10.1007/s12035-018-1045-5.

Reichwald, J., Danner, S., Wiederhold, K.H., and Staufenbiel, M. (2009). Expression of
complement system components during aging and amyloid deposition in APP
transgenic mice. *J Neuroinflammation* 6**,** 35. doi: 10.1186/1742-2094-6-35.

Ripley, B., Venables, B., Bates, D.M., Hornik, K., Gebhardt, A., Firth, D., et al. (2013).
Package 'mass'. *Cran r* 538**,** 113-120.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. *Bioinformatics* 26(1)**,**
139-140. doi: 10.1093/bioinformatics/btp616.

Sherman, B.T., and Lempicki, R.A. (2009). Systematic and integrative analysis of large gene
lists using DAVID bioinformatics resources. *Nature protocols* 4(1)**,** 44.

Solarski, M., Wang, H., Wille, H., and Schmitt-Ulms, G. (2018). Somatostatin in Alzheimer's
disease: A new Role for an Old Player. *Prion* 12(1)**,** 1-8. doi:
10.1080/19336896.2017.1405207.

Srinivasan, K., Friedman, B.A., Etxeberria, A., Huntley, M.A., van der Brug, M.P., Foreman,
O., et al. (2020). Alzheimer's Patient Microglia Exhibit Enhanced Aging and Unique
Transcriptional Activation. *Cell Rep* 31(13)**,** 107843. doi:
10.1016/j.celrep.2020.107843.

Trabzuni, D., Ryten, M., Walker, R., Smith, C., Imran, S., Ramasamy, A., et al. (2011). Quality
control parameters on a large dataset of regionally dissected human control brains for
whole genome expression studies. *J Neurochem* 119(2)**,** 275-282. doi:
10.1111/j.1471-4159.2011.07432.x.

van Rooij, J.G.J., Meeter, L.H.H., Melhem, S., Nijholt, D.A.T., Wong, T.H., Netherlands
Brain, B., et al. (2019). Hippocampal transcriptome profiling combined with
protein-protein interaction analysis elucidates Alzheimer's disease pathways and genes.
Neurobiol Aging 74**,** 225-233. doi: 10.1016/j.neurobiolaging.2018.10.023.

Verbitsky, M., Yonan, A.L., Malleret, G., Kandel, E.R., Gilliam, T.C., and Pavlidis, P. (2004).
Altered hippocampal transcript profile accompanies an age-related spatial memory
deficit in mice. *Learn Mem* 11(3)**,** 253-260. doi: 10.1101/lm.68204.
Wang, M., Beckmann, N.D., Roussos, P., Wang, E., Zhou, X., Wang, Q., et al. (2018). The
Mount Sinai cohort of large-scale genomic, transcriptomic and proteomic data in
Alzheimer's disease. *Sci Data* 5**,** 180185. doi: 10.1038/sdata.2018.185.
Wang, M., Roussos, P., McKenzie, A., Zhou, X., Kajiwara, Y., Brennand, K.J., et al. (2016).
Integrative network analysis of nineteen brain regions identifies molecular signatures
and networks underlying selective regional vulnerability to Alzheimer's disease.
Genome Med 8(1)**,** 104. doi: 10.1186/s13073-016-0355-3.
Wang, X., Park, J., Susztak, K., Zhang, N.R., and Li, M. (2019). Bulk tissue cell type
deconvolution with multi-subject single-cell expression reference. *Nat Commun* 10(1)**,**
380. doi: 10.1038/s41467-018-08023-x.
Yang, J., Huang, T., Petralia, F., Long, Q., Zhang, B., Argmann, C., et al. (2015). Synchronized
age-related gene expression changes across multiple tissues in human and the link to
complex diseases. *Sci Rep* 5**,** 15145. doi: 10.1038/srep15145.
Yao, J., and Bajjalieh, S.M. (2009). SVOP is a nucleotide binding protein. *PloS one* 4(4)**,**
e5315.
Yu, X., Ng, C.P., Habacher, H., and Roy, S. (2008). Foxj1 transcription factors are master
regulators of the motile ciliogenic program. *Nat Genet* 40(12)**,** 1445-1453. doi:
10.1038/ng.263.
Zeisel, S.H. (2007). Gene response elements, genetic polymorphisms and epigenetics influence
the human dietary requirement for choline. *IUBMB Life* 59(6)**,** 380-387. doi:
10.1080/15216540701468954.
Zeng, L., Yang, J., Peng, S., Zhu, J., Zhang, B., Suh, Y., et al. (2020). Transcriptome analysis
reveals the difference between "healthy" and "common" aging and their connection
with age-related diseases. *Aging Cell* 19(3)**,** e13121. doi: 10.1111/acel.13121.
Zhou, G., Soufan, O., Ewald, J., Hancock, R.E.W., Basu, N., and Xia, J. (2019).
NetworkAnalyst 3.0: a visual analytics platform for comprehensive gene expression
profiling and meta-analysis. *Nucleic Acids Res* 47(W1)**,** W234-W241. doi:
10.1093/nar/gkz240.

760
761
762
763

Figures Legends

Figure 1. Flowchart of the comparison between normal brain aging and AD. We collect gene expression profiles from a large number of normal aging brain and AD brain samples across multiple brain regions. We perform both a global comparison (Q1) and region-specific comparison (Q2) of aging and AD transcriptomes. We also consider the subgrouping in aging brain samples and compare the aging subgroups with AD (Q3).

Figure 2. Overlap between aging and AD signatures in different brain regions. AD signatures are plotted in rows and aging signatures are plotted in columns. We separate each signature into up- and down- regulated genes and the number of genes in each signature is listed after its ID. The number in the heatmap indicates how many genes are common in the corresponding aging and AD signatures while the color indicates the significance of the overlap.

Figure 3. Function annotation of hippocampus aging signatures (GTEx, UK) compared with AD signatures (Annese2018, Rooij2019). We consider the functional enrichment of aging specific genes (ASGs), conserved AD specific genes (ADSGs) between two AD signatures, and the overlap between aging and AD genes (AADGs) which represents genes shared by at least one aging list and one AD list.

We list the most representative function categories with FDR < 0.05. To reduce redundancy, only one representative functional category from each identified cluster of functions was selected.

Figure 4. Three major subgroups can be identified from either GTEx or UK hippocampus gene expression data. (**A).** Hierarchical clustering of GTEx hippocampus samples. 56 samples with donor age between 45 and 70 are plotted. Three subgroups are labeled in different colors in the dendrogram. The top color bar indicates that samples from different age groups are relatively evenly distributed into all the subgroups. (**B).** Hierarchical clustering of 70 UK hippocampus samples (45 ≤ age ≤ 75) also suggests these samples can be divided into three major subgroups.

Figure 5. Comparison of DEGs between GTEx subgroups and AD signatures from hippocampus. "GTEx DEG CvsB" represents DEGs derived from comparing subgroups C vs. B in GTEx hippocampus, similar naming format is used for the rest gene lists.

Figure 6. Function annotation of ABSGs, ABADGs, and ADSGs for a comparison of aging subgroup DEGs and AD signatures. The overlap of ABSGs between GTEx and UK is denoted as "Conserved ABSGs". The union of Rooij2019 and Annese2018 subtracting any ABGs is denoted as "ADSGs". We list the most representative function categories with FDR < 0.05. To reduce redundancy, only one representative functional category from each identified cluster of functions was selected.

Figure 7. Three major subgroups before and after mixed sample removal. (A). Hierarchical clustering of 78 PHG samples with age > 70. Three subgroups are labeled in different colors in the dendrogram. The color bar on the top indicates from the AD status of the sample: normal or AD. (**B).** Hierarchical clustering of 51 PHG samples with age > 70 after mixed samples being removed.

Tables

Table 1. List of brain transcriptomic datasets used for obtaining aging and AD gene signatures.

Datasets	Gene List ID	List Details (Tissue, traits)	# of genes (UP/DOWN)	# of protein coding genes	Sample Size
GTEx "Aging" sets (age ≤ 70)					
GTEx	G_AMY	Amygdala	50(19/31)	46(19/27)	46
GTEx	G _BA24_AC	Anterior_cingulate_cortex (BA24)	1460(473/987)	1208(320/888)	56
GTEx	G_CAU	Caudate (basal_ganglia)	77(50/27)	55(34/21)	75
GTEx	G_CRBH	Cerebellar_Hemisphere	322(138/184)	277(108/169)	73
GTEx	G_CRBL	Cerebellum	427(213/214)	340(162/178)	87
GTEx	G_CTX	Cortex	97(34/63)	78(30/48)	71
GTEx	G_ BA9 DLPFC	Frontal_Cortex (BA9)	15(8/7)	12(5/7)	61
GTEx	G_HIPP	Hippocampus	1804(950/854)	1403(705/698)	57
GTEx	G_HYPO	Hypothalamus	2140(822/1318)	1650(490/1160)	53
GTEx	G_NC	Nucleus_accumbens (basal_ganglia)	11(6/5)	9(5/4)	69
GTEx	G_PUTM	Putamen (basal_ganglia)	18(13/5)	17(13/4)	67
UK "Aging" sets (age ≤ 70)					
UK	UK_FCTX	Frontal cortex (FCTX)	170(50/120)	145(42/103)	94
UK	UK_TCTX	Temporal cortex (TCTX)	678(169/509)	510(91/419)	82
UK	UK_OCTX	Occipital cortex (OCTX)	54(30/24)	44(23/21)	95
UK	UK_WHMT	Intralobular white matter (WHMT)	464(267/197)	373(204/169)	93
UK	UK_CRBL	Cerebellum (CRBL)	186(101/85)	156(87/69)	92
UK	UK_PUTM	Putamen (PUTM)	65(30/35)	52(22/30)	95
UK	UK_THAL	Thalamus (THAL)	12(8/4)	9(5/4)	91
UK	UK_HIPP	Hippocampus (HIPP)	959(544/415)	755(431/324)	93
Other "AD" sets					
MSSM	MSSM_FP	BM10(MSSM_FP)	54(20/34)	47(19/28)	135
MSSM	MSSM_IFG	BM44(MSSM_IFG)	71(24/47)	61(24/37)	116
MSSM	MSSM_PHG	BM36(MSSM_PHG)	1577(936/641)	1391(868/523)	103
MSSM	MSSM_STG	BM22(MSSM_STG)	269(150/119)	242(142/100)	122
Mayo	Mayo_CRBL_Source *	Cerebellum	2632(1360/1272)	2282(1231/1051)	151
Mayo	Mayo_ CRBL	Cerebellum	1880(1008/872)	1600(901/699)	151
Mayo	Mayo_TCTX_Source *	Temporal cortex	3842(2034/1808)	3437(1933/1504)	151
Mayo	Mayo_TCTX	Temporal cortex	2951(1663/1288)	2628(1578/1050)	151
ROSMAP	ROSMAP_DLPFC	Dorsolateral prefrontal cortex (DLPFC)	162(89/73)	141(80/61)	241
Annese2018	Annese2018_HIPP	Hippocampus	2122(808/1314)	1925(742/1183)	10
Rooij2019	Rooij2019_HIPP	Hippocampus	2840(1109/1731)	2667(1045/1622)	28
Jager	Jager_Clinical_AD	Clinical_AD in DLPFC	855(466/389)	740(412/328)	478
Jager	Jager_Cognitive_decline	Cognitive_decline in DLPFC	3035(1481/1554)	2662(1281/1381)	478

Jager	Jager_Tau_tangles	Tau_tangles in DLPFC	238(155/83)	209(133/76)	478
Jager	Jager_B_amyloid	B_amyloid in DLPFC	2315(1158/1157)	2020(1060/960)	478
Jager	Jager_Patho_AD	Patho_AD in DLPFC	98(58/40)	85(51/34)	478

MS "AD" sets

MS	MS_BM10_FP	Frontal Pole	19(5/14)	17(5/12)	63
MS	MS_BM17_OCTX	Occipital Visual Cortex	345(325/21)	297(279/19)	53
MS	MS_BM20_ITG	Inferior Temporal Gyrus	397(397/0)	349(349/0)	58
MS	MS_BM21_MTG	Middle Temporal Gyrus	635(81/565)	571(66/516)	58
MS	MS_BM22_STG	Superior Temporal Gyrus	25(0/25)	24(0/24)	60
MS	MS_BM23_PCC	Posterior Cingulate Cortex	34(7/27)	29(7/22)	58
MS	MS_BM32_AC	Anterior Cingulate	239(33/207)	214(30/184)	59
MS	MS_BM36_PHG	Parahippocampal Gyrus	55(7/49)	48(6/42)	60
MS	MS_BM38_TP	Temporal Pole	44(28/16)	39(25/14)	58
MS	MS_BM4_PCG	Precentral Gyrus	131(8/123)	122(7/115)	49
MS	MS_BM44_IFG	Inferior Frontal Gyrus	874(589/566)	764(513/495)	53
MS	MS_BM46_DLPFC	Dorsolateral Prefrontal Cortex	58(11/47)	53(8/45)	57
MS	MS_BM7_SPL	Superior Parietal Lobule	346(240/110)	314(220/97)	50
MS	MS_BM8_SFG	Superior Prefrontal Gyrus	953(426/670)	869(386/617)	56
MS	MS_CD	Caudate Nucleus	31(30/1)	29(28/1)	52
MS	MS_HIPP	Hippocampus	107(25/82)	95(22/73)	55
MS	MS_PUTM	Putamen	94(52/42)	82(45/37)	52

✳ AMP-AD Mayo gene signature is based on SourceDiagnosis

Supporting Information

Supplementary Figures
Supplementary Tables

Normal aging brains

GTEx brain 13 regions
UK brain 10 regions

AD brains

AMP-AD 13 regions
MS Brain 19 regions
& Others

Q2: Brain region-specific comparison between aging and AD gene expression changes across brain regions

Aging gene regulation

Cross region comparison

AD gene regulation

Q3: Identify subgroups of aging and compare subgroup-DEGs with AD genes in hippocampus

Q1: Identify commonly regulated aging and AD genes across brain regions and how do they compare?

Common aging gene exp regulation across brain regions

Common AD gene exp regulation across brain regions

◯ Brain region specific gene expression changes

Aging subgroups

A   B   C

Aging subgroups

comparison

AD signature

A. Up hippocampus gene lists

Annese2018_742, GTEx_705, UK_431, Rooij2019_1045

Venn diagram values: 472, 245, 52, 54, 48, 401, 10, 8, 665, 5, 34, 79, 27, 25, 188

Conserved 188 ADSGs UP

Term	N	FDR$_{BH}$
Phosphoprotein	111	3.86E-05
Repressor	21	6.12E-05
inflammatory response	18	5.11E-04
Activator	20	6.35E-04
Cell membrane	53	6.35E-04
Glycoprotein	65	0.00386
Transcription regulation	40	0.00411
Stress response	7	0.00797
Polymorphism	132	0.01592
glycosylation site:N-linked (GlcNAc...)	63	0.03114
Cytoplasm	62	0.04739

288 AADGs UP ( )

Term	N	FDR$_{BH}$
Phosphoprotein	163	3.65E-06
Alternative splicing	192	1.89E-05
Staphylococcus aureus infection	9	7.82E-04
Coiled coil	70	0.001906
Actin-binding	15	0.002436
Cytoskeleton	33	0.007146
Polymorphism	199	0.007146
Cytoplasm	92	0.030856

771 ASGs UP ( )

Term	N	FDR$_{BH}$
Alternative splicing	500	6.90E-12
Phosphoprotein	401	1.59E-09
Coiled coil	181	1.00E-08
splice variant	384	8.13E-08
Acetylation	194	8.87E-08
Polymorphism	522	5.48E-06
Nucleus	261	7.68E-06
Disease mutation	134	0.001606
Metal-binding	178	0.003041
Endoplasmic reticulum	65	0.004370
Zinc-finger	97	0.005189
Protein transport	42	0.006566
Cytoskeleton	67	0.006720
Ciliopathy	15	0.006720
Cytoplasm	220	0.011919
Ubl conjugation	89	0.023305
Transcription regulation	115	0.030039
Repressor	38	0.030039

B. Down hippocampus gene lists

Annese2018_1183, GTEx_698, UK_324, Rooij2019_1622

Venn diagram values: 342, 184, 49, 15, 48, 509, 4, 8, 795, 7, 15, 157, 43, 108, 448

Conserved 448 ADSGs DN

Term	N	FDR$_{BH}$
Membrane	268	8.92E-22
Synapse	43	5.96E-17
myelin sheath	28	1.09E-13
Transport	98	1.22E-12
Synaptic vesicle cycle	19	7.76E-12
Phosphoprotein	238	8.88E-07
Ion transport	39	1.28E-06
mitochondrion	66	1.35E-06
Lipoprotein	44	1.09E-05
Alternative splicing	281	2.70E-05
Cytoplasmic vesicle	29	8.55E-05
Glycolysis	8	8.55E-05
Oxidative phosphorylation	16	3.34E-04
Metabolic pathways	60	3.34E-04
Glycoprotein	137	4.69E-04
Retrograde endocannabinoid signaling	13	8.44E-04
extracellular exosome	91	0.02873
Methylation	36	0.03057

439 AADGs DN ( )

Term	N	FDR$_{BH}$
Synapse	59	7.44E-32
plasma membrane	173	4.55E-16
Phosphoprotein	254	1.73E-12
Ion transport	43	4.15E-09
synaptic vesicle	15	6.77E-07
Kinase	41	1.41E-06
Alternative splicing	271	1.08E-04
cAMP signaling pathway	17	2.40E-04
Phosphatidylinositol signaling system	12	2.50E-04
Circadian entrainment	11	9.33E-04
Cytoskeleton	39	0.03103

541 ASGs DN ( )

Term	N	FDR$_{BH}$
Phosphoprotein	288	1.07E-07
Methylation	59	2.13E-06
protein binding	317	2.57E-06
Cytoplasm	182	4.08E-06
Kinase	47	4.08E-06
Ubl conjugation pathway	43	1.81E-05
Nucleotide-binding	82	4.40E-05
Transferase	79	4.53E-05
Neurogenesis	20	0.001289
Serine/threonine-protein kinase	26	0.001329
Acetylation	126	0.001329
Nucleus	179	0.001329
Endocytosis	21	0.004188

A. GTEx hippocampus 3 subgroups

B. UK hippocampus 3 subgroups

A. UP genes between aging and AD

B. DOWN (DN) genes between aging and AD

875 ABADGs UP ()

Term	N	FDR$_{BH}$
Phosphoprotein	537	4.96E-36
Activator	72	1.25E-10
Polymorphism	611	1.76E-10
protein binding	512	2.16E-10
Alternative splicing	550	3.01E-10
Repressor	64	1.58E-09
Cytoplasm	283	3.83E-08
Transcription	160	2.44E-07
Nucleus	298	2.99E-07
angiogenesis	37	3.87E-07
Ubl conjugation	120	1.44E-06
focal adhesion	47	1.88E-06
sequence variant	623	3.35E-06
Extracellular matrix	32	6.11E-06
Coiled coil	185	6.11E-06
Actin-binding	33	6.86E-06
PI3K-Akt signaling pathway	42	1.71E-05
Cell adhesion	45	3.90E-05
Cytoskeleton	81	1.37E-04
extracellular exosome	178	1.99E-04
Rho GTPase activation protein	18	2.24E-04
MAPK signaling pathway	30	9.90E-04
Insulin resistance	18	9.90E-04
membrane raft	25	0.00131

668 ADSGs UP ()

Term	N	FDR$_{BH}$
Glycoprotein	244	5.63E-16
Immunity	60	6.10E-16
Polymorphism	475	2.67E-10
Membrane	328	4.22E-10
Signal	207	1.05E-09
Receptor	102	1.20E-08
cell surface	52	2.23E-08
leukocyte migration	23	2.39E-07
Secreted	109	7.26E-07
sequence variant	479	9.80E-07
receptor activity	29	3.00E-06
Apoptosis	41	2.45E-05
Tuberculosis	26	3.54E-05
Osteoclast differentiation	22	3.54E-05
extracellular space	82	8.29E-05
Cell adhesion	36	1.74E-04
membrane raft	22	6.38E-04
Phagosome	21	7.32E-04

Conserved 420 ABSGs UP

Term	N	FDR$_{BH}$
Phosphoprotein	247	2.03E-12
Alternative splicing	271	6.80E-06
Cytoplasm	145	2.15E-05
splice variant	216	5.07E-05
Cytoskeleton	49	1.04E-04
Coiled coil	99	1.04E-04
Methylation	44	1.67E-04
Cell cycle	30	0.00282
Polymorphism	284	0.00282
Cell projection	32	0.00282
Actin-binding	17	0.00536
Microtubule	17	0.00630
Cell division	20	0.01103
Motor protein	11	0.01119
Nucleus	138	0.01162
Endocytosis	10	0.01814

1616 ABADGs DN ()

Term	N	FDR$_{BH}$
Synapse	156	2.62E-74
cell junction	159	5.76E-54
Membrane	877	1.75E-50
Transport	306	2.48E-31
Ion transport	144	1.67E-29
dendrite	100	3.08E-27
Ion channel	96	3.35E-25
axon	69	1.10E-19
Synaptic vesicle cycle	36	1.34E-18
Phosphoprotein	819	2.24E-18
Alternative splicing	995	8.21E-17
perikaryon	42	7.06E-16
myelin sheath	49	1.68E-14
Retrograde endocannabinoid signaling	39	1.55E-13
Oxidative phosphorylation	44	9.77E-13
neurotransmitter secretion	27	4.87E-12
Lipoprotein	126	1.03E-10
Synaptosome	23	2.09E-09
growth cone	35	2.55E-09
Oxytocin signaling pathway	41	2.67E-09
Circadian entrainment	31	6.13E-09
Long-term potentiation	25	1.17E-08

576 ADSGs ()

Term	N	FDR$_{BH}$
Calcium	62	9.35E-09
Ciliopathy	19	1.74E-06
Glycoprotein	182	7.06E-06
Disease mutation	115	1.30E-05
plasma membrane	76	3.07E-05
Dynein	9	1.42E-04
Cytoskeleton	7	5.39E-04
Signal	157	0.00109
Ion transport	37	0.00215
axoneme	12	0.00261
Polymorphism	380	0.00319
Membrane	253	0.00319
Secreted	83	0.00379
Ion channel	24	0.00519
extracellular region	72	0.00808
Alternative splicing	331	0.03113

Conserved 484 ABSGs DN

Term	N	FDR$_{BH}$
cytosol	154	4.40E-13
Phosphoprotein	264	1.62E-08
Acetylation	137	1.62E-08
Cytoplasm	174	3.65E-08
Protein transport	42	1.39E-07
Alternative splicing	305	1.10E-05
Nucleotide-binding	76	2.98E-05
mitochondrion	63	2.82E-04
GTP binding	27	0.00209
protein binding	268	0.00441
Transferase	65	0.00467
Transport	72	0.00640
Proteasome	8	0.00678
Nucleus	158	0.00764
Ubl conjugation pathway	32	0.00922

Venn diagram A: GTEx_2827, Annese2018_742, Rooij2019_1045, UK_3601 — 1906, 235, 345, 87, 90, 193, 51, 69, 38, 209, 112, 53, 60, 420, 2116

Venn diagram B: GTEx_3018, Annese2018_1183, Rooij2019_1622, UK_3144 — 1073, 337, 200, 38, 95, 91, 208, 19, 349, 414, 45, 303, 92, 484, 1761

A. PHG 3 subgroups before mixed sample removal

B. PHG 3 subgroups after mixed sample removal