Title: Male age and *Wolbachia* dynamics: Investigating how fast and why bacterial densities

and cytoplasmic incompatibility strengths vary

Short title: Male age and *Wolbachia* dynamics

Authors: J. Dylan Shropshire[1], Emily Hamant[1], and Brandon S. Cooper[1]

[1]Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA

*Correspondence to:

J. Dylan Shropshire, Missoula, MT, 59801, 423.930.6292, shropxp@gmail.com

ORCID iD: J. Dylan Shropshire (https://orcid.org/0000-0003-4221-2178), Emily Hamant

(https://orcid.org/0000-0001-5743-1731), Brandon S. Cooper (https://orcid.org/0000-0002-8269-

7731)

Author contributions:

JDS roles: Conceptualization, Data curation, Formal Analysis, Funding acquisition,

Investigation, Methodology, Supervision, Validation, Visualization, Writing - original draft, Writing

– review & editing. EH roles: Investigation, Validation, Writing – review & editing. BSC roles:

Project administration, Conceptualization, Funding acquisition, Supervision, Writing – original

draft, Writing – review & editing.

Keywords: aging, *Drosophila*, immunity, symbiosis, *w*Mel, *w*Ri

Abstract

Endosymbionts can influence host reproduction and fitness to favor their maternal

transmission. For example, endosymbiotic *Wolbachia* bacteria often cause cytoplasmic

incompatibility (CI) that kills uninfected embryos fertilized by *Wolbachia*-modified sperm.

Infected females can rescue CI, providing them a relative fitness advantage. *Wolbachia*-induced

CI strength varies widely and tends to decrease as host males age. Since strong CI drives

Wolbachia to high equilibrium frequencies, understanding how fast and why CI strength declines

with male age is crucial to explaining age-dependent CI's influence on *Wolbachia* prevalence.

Here, we investigate if *Wolbachia* densities and/or CI gene (*cif*) expression covary with CI-

strength variation and explore covariates of age-dependent *Wolbachia*-density variation in two

classic CI systems. *w*Ri CI strength decreases slowly with *Drosophila simulans* male age (6%/

day), but *w*Mel CI strength decreases very rapidly (19%/ day), yielding statistically insignificant

CI after only three days of *D. melanogaster* emergence. *Wolbachia* densities and *cif* expression

in testes decrease as *w*Ri-infected males age, but both surprisingly increase as *w*Mel-infected

males age, and CI strength declines. We then tested if phage lysis, Octomom copy number

(which impacts *w*Mel density), or host immune expression covary with age-

dependent *w*Mel densities—only host immune expression correlated with density. Together, our

results identify how fast CI strength declines with male age in two model systems and reveal

unique relationships between male age, *Wolbachia* densities, *cif* expression, and host immunity.

We discuss new hypotheses about the basis of age-dependent CI strength and its contributions

to *Wolbachia* prevalence.

Importance

   Wolbachia are the most common animal-associated endosymbionts due in large part to their manipulation of host reproduction. Many *Wolbachia* cause cytoplasmic incompatibility (CI) that kills uninfected host eggs. Infected eggs are protected from CI, favoring *Wolbachia* spread in natural systems and in transinfected mosquito populations where vector-control groups use strong CI to maintain pathogen-blocking *Wolbachia* at high frequencies for biocontrol of arboviruses. CI strength varies considerably in nature and declines as males age for unknown reasons. Here, we determine that CI strength weakens at different rates with age in two model symbioses. *Wolbachia* density and CI gene expression covary with *w*Ri-induced CI strength in *Drosophila simulans*, but neither explain rapidly declining *w*Mel-induced CI in aging *D. melanogaster* males. Patterns of host immune gene expression suggest a candidate mechanism behind age-dependent *w*Mel densities. These findings inform how age-dependent CI may contribute to *Wolbachia* prevalence in natural systems and potentially in transinfected systems.

Introduction

   Reproductive parasites manipulate host reproduction to facilitate their spread in host populations. These endosymbiotic microbes may kill or feminize males or induce parthenogenesis to bias sex ratios favoring females (1). More frequently, reproductive parasites cause cytoplasmic incompatibility (CI) that reduces embryonic viability when aposymbiotic females mate with symbiont-bearing males (**Fig. 1A**) (2). Females harboring a closely related symbiont are compatible with CI-causing symbiotic males of the same strain, providing symbiont-bearing females a relative advantage that encourages symbiont spread to high frequencies in host populations (3–8). Divergent *Cardinium* (9), *Rickettsiella* (10), *Mesenet* (11), and *Wolbachia* (12) endosymbionts cause CI. *Wolbachia* are the most common, infecting 40-65% of arthropod species (13, 14). *Wolbachia* cause CI in at least ten arthropod orders (2), and pervasive CI directly

contributes to *Wolbachia* spread and its status as one of the most common endosymbionts in

nature.

Within host populations, *Wolbachia* frequencies are governed by their effects on host

fitness (15–20), maternal transmission efficiency (21–23), and CI strength (% embryonic death)

(3, 5). CI strength varies from very weak to very strong and produces relatively low and high

infection frequencies, respectively. For example, *w*Yak in *Drosophila yakuba* causes weak CI

(~15%) and tends to occur at intermediate and often variable frequencies (~40-88%) in West

Africa (22, 24). Conversely, *w*Ri in *D. simulans* causes strong CI (~90%) and occurs at high and

stable frequencies (e.g., ~93% globally) (4, 25–27). In *D. melanogaster*, *w*Mel CI strength is

relatively weak (28–30), contributing to considerably differing infection frequencies on multiple

continents (31–35). In contrast, *w*Mel usually causes complete CI (no eggs hatch) in transinfected

Aedes aegypti mosquitoes (36–39). Vector-control groups use strong CI induced by *w*Mel and

other variants (e.g., *w*AlbB and *w*Pip) to either suppress mosquito populations through the release

of infected males (40–45) or to drive pathogen-blocking *Wolbachia* to high and stable frequencies

to inhibit pathogen spread (36, 46–49).

Despite CI's importance for explaining *Wolbachia* prevalence in natural systems and

reducing human disease transmission in transinfected mosquito systems, the mechanistic basis

of CI-strength variation remains unresolved. Two hypotheses are plausible. First, the bacterial-

density model predicts that CI is strong when bacterial density is high **(Fig. 1B)** (50). Indeed,

Wolbachia densities positively covary with CI strength across *Drosophila-Wolbachia* associations

(51, 52) and variable CI within strains (37, 38, 53–59). Second, the CI-gene-expression

hypothesis predicts higher CI-gene expression contributes to stronger CI **(Fig. 1B)** (60). In

Drosophila, two genes (*cifA* and *cifB*) associated with *Wolbachia*'s bacteriophage WO contribute

to CI when expressed in testes (60–64), and one gene (*cifA*) rescues CI when expressed in

ovaries (63–65). CI strength covaries with transgenic *cif* expression in *D. melanogaster* (60, 64),

and natural *cif* expression covaries with CI strength in *Habrobracon* ectoparasitoid wasps (66).

Bacterial density may explain CI strength via *cif* expression but may not perfectly align with CI

strength since *Wolbachia* variably express *cifs* across conditions that impact CI strength (60).

Thus, the bacterial-density and *cif*-expression hypotheses are not mutually exclusive. It remains

unknown if *cif* expression is responsible for CI-strength variation and if it covaries with *Wolbachia*

density in natural *Drosophila-Wolbachia* associations.

If symbiont density is a crucial factor governing CI strength, what governs the change in

density? There are several plausible drivers of *Wolbachia*-density variation. First, phage WO is a

temperate phage capable of cell lysis in some *Wolbachia* strains (66–70). Lytic phages form

particles that burst through the bacterial cell membrane, killing the bacterial host. The phage

density model proposes that as phage densities increase, *Wolbachia* densities decrease (**Fig.**

1B) (53). Temperature induced phage lysis covaries with lower *Wolbachia* densities and CI

strength in some parasitoid wasps (53, 66), though it is unknown if phage lysis influences

Wolbachia densities in any other systems. Second, *w*Mel *Wolbachia* have a unique ampliconic

gene region composed of eight genes termed "Octomom" (71–75). Octomom copy number varies

among *w*MelCS and *w*MelPop *Wolbachia* between host generations and positively covaries with

Wolbachia densities **(Fig. 1B)**, but effects of Octomom-dependent *Wolbachia* densities on CI

have not been investigated. Third, theory predicts that selection favors the evolution of host

suppressors (6), as observed for male killing (76–79). Indeed, CI strength varies considerably

across host backgrounds (24, 29, 39, 80–82), supporting a role for host genotype in CI-strength

variation. The genetic underpinnings and mechanistic consequences of host suppression remain

unknown, but two models have been proposed (2). The defensive model suggests that host CI

targets diverge to prevent interaction with *cif* products, and the offensive model suggests that host

products directly interfere with *Wolbachia* density or the proper expression of *cif* products (e.g.,

through immune regulation) **(Fig. 1B)**. Only a taxon-restricted gene of *Nasonia* wasps and host

transcriptional activity in *Drosophila* have been functionally determined to contribute to

Wolbachia-density variation (83, 84); thus, considerable work is necessary to uncover host

determinants of variation in *Wolbachia*-density. Since *Wolbachia* densities significantly contribute

to several phenotypes (54, 85), investigation of the causes of *Wolbachia*-density variation is sorely

needed.

CI strength within *Wolbachia*-host systems covaries with several factors, including

temperature (29, 37, 38, 53, 66), male mating rate (86, 87), male development time (88), rearing

density (88), nutrition (89), paternal grandmother age (30), and male age (3, 18, 27, 29, 86). Male

age does not always influence CI strength (90–92), but *w*Mel-infected *D. melanogaster* (29), *w*Ri-

infected *D. simulans* (3, 18, 27), and other *Wolbachia*-infected hosts tend to cause weaker CI as

males age (91, 93–95). CI seems to decline more slowly for *w*Ri (3, 18, 27) than for *w*Mel (3, 18,

27, 29), though the precise rates of CI-strength decline have not been estimated. While several

factors might contribute to age-dependent CI strength, the mechanistic underpinnings of this

phenotype remain unknown.

Here, we investigate rates of CI decline with male age and its mechanistic underpinnings

in two classic *Wolbachia* CI systems: *w*Ri and *w*Mel (25, 28, 32). These *Wolbachia* diverged up

to 6 million years ago and have unique *cif* repertoires (60, 63). We demonstrate that relative to

*w*Ri, *w*Mel-induced CI strength declines more than three times faster, disappearing in a matter of

days. We provide the first direct test of the *cif*-expression hypothesis in either system and the

highest resolution investigation of *Wolbachia*-density variation across age to date. Our results

suggest that *Wolbachia* density and *cif* expression in full-testes extracts cannot explain age-

dependent CI-strength relationships across *Wolbachia*-host associations and motivate future

work to investigate how host immunity could contribute to age-dependent *Wolbachia* densities.

We discuss how these data inform our understanding of the causes of CI-strength variation,

Wolbachia-density variation, and the consequences for *Wolbachia* prevalence in nature.

Figure 1. CI crossing relationships and potential causes of CI-strength variation. (A) CI
causes embryonic death, measured as reduced embryo hatch when infected males (filled symbol)
mate with uninfected females (unfilled symbol). All other crosses are compatible and have high
embryonic hatching. Importantly, infected females maternally transmit *Wolbachia* and can rescue
CI. (B) Schematic representation of factors that putatively impact *Wolbachia* densities, CI gene
expression, and CI strength.

Results

How much does CI strength vary with age?

CI manifests as embryonic lethality (**Fig. 1A**). As such, we measured CI strength as the

percent of embryos that hatch from a mating pair's clutch of offspring—high compatibility

corresponds with high hatching. Our experiments used males of different ages to test the impact

of male age on CI strength. Here, we defined age as days since eclosion where males paired

with females the day they eclosed were considered 0-days-old. For *w*Mel, we measured CI

strength daily across the first three days of male age (**Fig. 2A**) and separately every two days

across the first eight days of male age (**Fig. 2B**). This design enabled us to determine the rate

of CI decline and the ages where males no longer cause significant CI. Crossing uninfected *D.*

melanogaster females and males yielded high levels of compatibility (**Fig. 2A**; 95% confidence

interval of the mean = 74 - 93%). Young 0-day-old *w*Mel-infected males induced strong CI when

mated with uninfected females (95% interval = 9 - 27%). *w*Mel-infected females significantly

rescued CI caused by infected 0-day-old males (95% interval = 87 - 92%, P = 1.74E-12).

Crosses using older 1- (95% interval = 31 - 51%), 2- (95% interval = 53 - 73%), and 3-day-old

(95% interval = 69 - 83%) infected males trended toward progressively weaker CI (**Fig. 2A**).

Average *w*Mel CI strength decreased daily by 19.3%: 22.8% from 0- to 1-day-old males, 21.8%

from 1- to 2-day-old, and 13.4% from 2- to 3-day-old. Crosses between uninfected females and

3-day-old males (95% interval = 69 - 83%) did not cause significant CI, with egg hatch similar to

the compatible uninfected (95% interval = 74 - 93%; P = 0.35) and rescue (95% interval = 87 -

92%; P = 0.19) crosses. These data highlight the rapid decline of *w*Mel CI strength with *D.*

melanogaster male age.

In the experiment that includes older males (**Fig. 2B**), the uninfected cross also yielded

high compatibility (95% interval = 72 - 88%). 0-day-old infected males caused strong CI when

crossed with uninfected females (95% interval = 8 - 15%), and infected females significantly

rescued 0-day-old CI (95% interval = 83 - 91%; P = 2.51E-12). Compatibility increased as males

aged, where 2-day-old (95% interval = 59 - 73%) males caused significant CI and 4- (95%

interval = 66 - 83%), 6- (95% interval = 76 - 92%), and 8-day-old (95% interval = 77 - 91%)

infected males did not significantly inhibit egg hatch relative to the compatible uninfected cross

(P = 1 in all cases) (**Fig 2B**). Average *w*Mel CI strength decrease by approximately 19.3% each

day as *D. melanogaster* males aged, but this rate of decrease slowed each day, such that CI

was no longer statistically detectable once males were 3-days-old.

Next, we assessed age-dependent CI in *w*Ri-infected *D. simulans* (**Fig. 2C**). As

expected, uninfected *D. simulans* females and males were compatible (95% interval = 74 -

94%). Young 0-day-old *w*Ri-infected males caused strong CI when mated with uninfected

females (95% interval = 0 - 1%), and infected females significantly rescued 0-day-old CI (95%

interval = 59 - 84%; P = 1.83E-10). Older 4- (95% interval = 21 - 39%), 8- (95% interval = 54 -

64%), and 12-day-old (95% interval = 64 - 82%) infected males induced progressively weaker

CI as males aged. Average *w*Ri CI strength decreased by about 6.0% per day: 29.1% (7.3%/

day) from 0-day-old to 4-day-old males, 29.0% (7.3%/ day) from 4-day-old to 8-day-old, and

14.0% (3.5%/ day) from 8-day-old to 12-day-old. These data support a strong effect of *D.*

simulans male age on *w*Ri CI strength, but the daily decrease is more than three times slower

than what we observed for *w*Mel CI strength decline as *D. melanogaster* males age.

Figure 2. CI strength decreases as males age. (A) Hatch rate displaying CI strength with 0-,
1-, 2-, and 3-day-old *w*Mel-infected *D. melanogaster* males. (B) Hatch rate displaying CI
strength with 0-, 2-, 4-, 6-, and 8-day-old *w*Mel-infected *D. melanogaster* males. (C) Hatch rate
displaying CI strength with 0-, 4-, 8-, and 12-day-old *w*Ri-infected *D. simulans* males. Filled and
unfilled sex symbols represent infected and uninfected flies, respectively. Male age is displayed
to the right of the corresponding sex symbol. CI crosses are colored red, rescue crosses are

purple, and uninfected crosses are green. Boxplots represent median and interquartile ranges.
Letters to the right represent statistically significant differences based on α=0.05 calculated by
Dunn's test with correction for multiple comparisons between all groups —crosses that do not
share a letter are significantly different. *P*-values are reported in **Table S1**. These data
demonstrate that CI strength decreases with age in two *Wolbachia*-host associations and more
slowly in *w*Ri-infected *D. simulans*.

What causes CI strength to vary with age?

The bacterial-density and CI-gene-expression hypotheses are both proposed to explain

CI-strength variation. These hypotheses predict that *Wolbachia* density and/or *cif* expression

positively covary with CI strength. To elucidate the causes of declining CI strength with male

age, we tested both hypotheses in the context of rapidly declining *w*Mel CI strength and more

slowly declining *w*Ri CI strength.

Bacterial density differentially covaries with age between species.

We tested the bacterial density hypothesis by dissecting testes from siblings of flies used

in our CI assays above, extracting DNA, and measuring the relative abundance of a single-copy

Wolbachia gene (*ftsZ*) relative to a single-copy ultraconserved element (UCE) (96) of *Drosophila*

via qPCR. We selected a random infected sample from the 0-day-old age group as the

reference for all fold change analyses within each experiment. We report all qPCR data as fold

change relative to this control. Surprisingly, 0-day-old *D. melanogaster* testes had low *w*Mel

density (**Fig. 3A**; 95% interval = 0.53 - 1.01 fold change), and older 2- (95% interval = 0.92 -

1.11), 4- (95% interval = 0.96 - 1.72), 6- (95% interval = 1.17 - 1.49), and 8-day-old (95%

interval = 1.19 - 1.51) infected testes had progressively higher *w*Mel densities (**Fig. 3A**). *w*Mel

densities were significantly different among age groups according to a Kruskal-Wallis test (**Fig.**

3A; *P* = 1.1E-03). To test for a correlation between *w*Mel densities and CI strength, we

performed Pearson (r_p) and Spearman (r_s) correlations on the relationship between *w*Mel fold

change against median hatch rates from the associated age groups. *w*Mel densities were

significantly positively correlated with increasing compatibility (**Table S3**; $r_p = 0.75$, $P = 5.5E-06$;

$r_s = 0.77$, $P = 2.3E-06$). *w*Mel densities also covaried with age (**Fig. S1A**; $P = 0.02$) and

correlated with increasing compatibility (**Table S3**; $r_p = 0.64$, $P = 7.7E-04$; $r_s = 0.64$, $P = 7.4E-04$)

in the younger 0-, 1-, 2-, and 3-day-old *D. melanogaster* age group. This result was contrary to

our prediction that higher *w*Mel densities would be correlated with stronger CI and lower

compatibility.

Next, we tested the bacterial density model in *w*Ri-infected *D. simulans*. In contrast to

*w*Mel, *w*Ri-infected 0-day-old (95% interval = 0.82 - 1.36) *D. simulans* testes had the highest

*w*Ri densities and they consistently decreased in 4- (95% interval = 0.41 - 0.83), 8- (95%

interval 0.41 - 0.83), and 12-day-old (95% interval = 0.24 - 0.40) testes (**Fig. 3B**). *w*Ri densities

were significantly different among *D. simulans* age groups ($P = 3.9E-04$) and were significantly

negatively correlated with increasing compatibility (**Table S3**; $r_p = -0.84$, $P = 2.4E-07$; $r_s = -0.89$,

$P = 6.9E-09$).

In conclusion, these data fail to support the bacterial-density hypothesis for age-

dependent CI-strength variation in *w*Mel-infected *D. melanogaster* but support the hypothesis in

*w*Ri-infected *D. simulans*. Thus, *Wolbachia* densities from full-testis extracts cannot explain age-

dependent CI across *Wolbachia*-host associations, suggesting that other factors contribute to

these patterns.

Figure 3. Testing the bacterial density model for CI-strength variation. Fold change in
testes across male age for the relative expression of (A) wMel *ftsZ* to *D. melanogaster* UCE and
(B) wRi *ftsZ* to *D. simulans* UCE. Letters above data represent statistically significant
differences based on α=0.05 calculated by a Dunn's test with correction for multiple
comparisons between all groups—crosses that do not share a letter are significantly different.
Fold change was calculated as $2^{-\Delta\Delta cq}$. We selected a random infected sample from the youngest
0-day-old age group as the reference for all fold change analyses within each experiment. *P*-
values are reported in **Table S1**. These data demonstrate that *Wolbachia* density differentially
covaries with age between *Wolbachia*-host associations.

cif expression varies with age, but the direction differs between strains.

cif expression is hypothesized to control CI-strength variation within *Wolbachia*-host

associations (2, 60). *cif* loci are classified into five different phylogenetic clades called 'Types'

(60, 97–99). wMel has a single pair of Type I *cifs,* and wRi has two identical pairs closely related

to the wMel copy plus a divergent Type 2 pair (60). We investigated three questions regarding

cif expression. First, does *cif* expression change relative to the host as males age? We

expected *cif* expression per host cell would be the key determinant of CI-strength variation. To

test this, we used RT-qPCR to measure the transcript abundance of *cifA* and *cifB* and compared

their expression to β Spectrin (*βspec*), a *Drosophila* membrane protein with invariable

expression with age (see Materials and Methods for details). Second, does *cif* expression

decrease relative to *Wolbachia* as males age? Since *w*Mel densities increase with male age,

*w*Mel would need to express *cif$_{wMel[T1]}$* at lower levels in older males to allow *cif$_{wMel[T1]}$* to decrease

relative to the host. Finally, does *cifA* expression change relative to *cifB* as males age?

Evidence of differential localization of *cif* loci that covaries with age might indicate more complex

determinants of age-dependent CI based on the relative abundance of these products.

We started by investigating these question in *w*Mel-infected *D. melanogaster*. Contrary

to our first prediction, the relative expression of *cifA$_{wMel[T1]}$* to *D. melanogaster βspec* was lowest

in 0-day-old infected males (95% = 1.1 - 1.6) and consistently increased in 2- (95% interval =

1.5 - 3.2), 4- (95% interval = 1.9 - 2.3), 6- (95% interval = 2.1 - 2.8), and 8-day-old (95% interval

= 0.9 - 3.8) testes (**Fig. 4A**). Relative expression of *cifA$_{wMel[T1]}$* to *βspec* significantly varied

across male age (P = 8.4E-03) and was significantly positively correlated with increasing

compatibility (**Table S3**; r_p = 0.61, P = 6.4E-04; r_s = 0.59, P = 9.7E-04). Comparably, relative

expression of *cifB$_{wMel[T1]}$* to *βspec* significantly increased with male age (**Fig. S2A**; P = 7.3E-03).

Analysis of raw quantification cycle (C$_q$) variation with age supports increased *cifA$_{wMel[T1]}$* (**Fig.**

S2C; P = 3.1E-04) and *cifB$_{wMel[T1]}$* (**Fig. S2D**; P = 1.1E-03) expression; *βspec* C$_q$ does not vary

with age (**Fig. S2E**; P = 0.1) and *ftsZ* C$_q$ significantly decreases with age (**Fig. S2F**; P = 1.3E-

04). Thus, we report for the first time that *cif$_{wMel[T1]}$* expression relative to the host in full-testes

extracts is not sufficient to explain CI-strength variation, leading us to reject the hypothesis that

cif$_{wMel[T1]}$ expression in full-testes extracts can explain age-dependent *w*Mel CI strength.

Next, we investigated our second question: does *cif$_{wMel[T1]}$* expression vary relative to

Wolbachia as males age? Indeed, relative expression of *cifA$_{wMel[T1]}$* to *w*Mel *ftsZ* was highest in

0-day-old infected-*D. melanogaster* testes (95% interval = 0.9 - 1.1), and consistently decreased

in 2- (95% interval = 0.7 - 0.8), 4- (95% interval = 0.7 - 0.9), 6- (95% interval = 0.6 - 0.7), and 8-

day-old (95% interval = 0.4 - 0.9) testes (**Fig. 4B**). Relative expression of *cifA$_{wMel[T1]}$* to *w*Mel *ftsZ*

significantly varied with age (P = 2.9E-03) and was significantly correlated with increasing

compatibility (**Table S3**; r_p = -0.8, P = 4.0E-07; r_s = -0.7, P = 3.5E-05). Relative expression of

cifB_{wMel[T1]} to *w*Mel *ftsZ* did not significantly covary with age (**Fig. S2B**; P = 0.3), but was

significantly correlated with increasing compatibility (**Table S3**; r_p = -0.42, *P* = 3.7E-02; r_s = -

0.46, *P* = 2.2E-02). In summary, *cif_{wMel[T1]}* expression decreased relative to a *Wolbachia*

housekeeping gene with age, consistent with prior reports that *w*Mel expression of *cifA_{wMel[T1]}*

and *cifB_{wMel[T1]}* decrease as males age (60). However, since *cif_{wMel[T1]}* expression did not

decrease relative to the host with age, we conclude that the decrease in *cif_{wMel[T1]}* expression per

Wolbachia is insufficient to overcome the increase in *cif_{wMel[T1]}* expression caused by increased

*w*Mel density in full-testes extracts.

Finally, we tested if the relative expression of *cifA_{wMel[T1]}* to *cifB_{wMel[T1]}* varied with age.

Intriguingly, *cifA/B_{wMel[T1]}* relative expression did not significantly covary with age (**Fig. 4C**; *P* =

0.09), but was positively correlated with decreasing CI strength (**Table S3**; r_p = -0.61, *P* = 1.3E-

03; r_s = -0.46, *P* = 0.021). In summary, these data suggest that *cif_{wMel[T1]}* expression per *w*Mel

decreases as males age, that *cifA_{wMel[T1]}* expression decreases marginally faster than *cifB_{wMel[T1]}*,

and that overall *cif_{wMel[T1]}* expression increases relative to the host as males age and CI strength

decreases. This is the first report that CI strength is decoupled from *Wolbachia* densities and *cif*

expression in testes.

Next, we investigated the *cif*-expression hypotheses in *w*Ri. We predicted that *cif_{wRi[T1]}*

and/or *cif_{wRi[T2]}* expression would decrease relative to host expression. Since *w*Ri density

decreased with age, *cif* expression per *w*Ri would not need to change to accomplish this shift in

relative expression. As predicted, relative expression of *cifA_{wRi[T1]}* to *D. simulans βspec* was

highest in infected 0-day-old (95% interval = 0.7 - 1.7) testes, and declined in 4- (95% interval =

0.1 - 0.4), 8- (95% interval = 0.3 - 0.7), and 12-day-old (95% interval = 0.2 - 0.3) testes (**Fig.**

4D). Relative expression of *cifA_{wRi[T1]}* to *D. simulans βspec* significantly covaried with age (*P* =

1.2E-03) and was significantly correlated with decreasing CI strength (**Table S3**; r_p = -0.76; r_s = -

0.88). Similarly, relative expression of *cifB_{wRi[T1]}* (**Fig. S3A**; *P* = 2.3E-03), *cifA_{wRi[T2]}* (**Fig. S3C**; *P* =

1.9E-03), and *cifB_{wRi[T2]}* (**Fig. S3E**; *P* = 1.2E-03) to *D. simulans βspec* also decreased with age

and each were significantly correlated with decreasing CI strength (**Table S3**). These results

support the *cif*-expression hypothesis for age-dependent CI in *w*Ri.

As with *w*Mel-infected *D. melanogaster* testes, relative expression of *cifA$_{wRi[T1]}$* to *w*Ri

ftsZ significantly covaried with male age (**Fig. 4E**; P = 4.1E-02) and was significantly correlated

with decreasing CI strength (**Table S3**; r_p = -0.47, P = 0.032; r_s = -0.47, P = 0.033). However, 0-

(95% interval = 0.9 - 1.2), 4- (95% interval = 0.9 - 1.2), and 8-day-old (95% interval = 0.8 - 1.2)

testes had similar expression patterns, suggesting that expression in 12-day-old (95% interval =

0.5 - 0.9) testes drove this significant difference; though, a Dunn's test was unable to identify

significantly different pairs (**Fig. 4E**). Conversely, *cifB$_{wRi[T1]}$* (**Fig. S3B**; P = 0.6), *cifA$_{wRi[T2]}$* (**Fig.**

S3D; P = 0.2), and *cifB$_{wRi[T2]}$* (**Fig. S3F**; P = 0.2) expression relative to *w*Ri *ftsZ* did not vary with

age or decreasing CI strength (**Table S3**).

Finally, as with *w*Mel, we investigated the relationship between *cifA* and *cifB* expression

in *w*Ri across age and found similar results where *cifA$_{wRi[T1]}$* expression relative to *cifB$_{wRi[T1]}$*

expression did not significantly vary with male age (**Fig. 4F**; P = 0.2) but did significantly

correlate with increasing compatibility (**Table S3**; r_p = -0.44, P = 0.045; r_s = -0.46, P = 0.035).

Relative expression of *cifA$_{wRi[T1]}$* to *cifA$_{wRi[T2]}$* expression did not covary with age (**Fig. S3G**; P =

0.6) or increasing compatibility (**Table S3**; r_p = 0.01, P = 0.96; r_s = -0.05, P = 0.84). Analysis of

raw C_q values supported decreasing *cifA$_{wRi[T1]}$* (**Fig. S3H**; P = 1.0E-03), *cifB$_{wRi[T1]}$* (**Fig. S3I**; P =

8.1E-04), *cifA$_{wRi[T2]}$* (**Fig. S3J**; P = 1.8E-03), and *cifB$_{wRi[T2]}$* (**Fig. S3K**; P = 1.7E-03) expression

with male age; *D. simulans βspec* C_q did not vary with age (**Fig. S3L**; P = 0.6) and *w*Ri *ftsZ* C_q

significantly increased with age (**Fig. S3M**; P = 8.9E-04). In summary, *cif$_{wRi}$* expression

significantly decreased with age in *w*Ri testes, *cifA$_{wRi[T1]}$* expression decreased marginally faster

than *cifB$_{wRi[T1]}$* expression, and there was a small decrease in *cifA$_{wRi[T1]}$* expression relative to

*w*Ri but other *cif$_{wRi}$* loci do not follow similar trends.

In conclusion, we found that *w*Mel *cif* expression did not explain age-dependent CI-

strength variation. More specifically, *w*Mel's expression of *cif* genes decreased with age (60),

relative *w*Mel and *w*Ri *cifA*-to-*cifB* expression varied marginally with age, and *cif* expression

dynamics varied considerably across male age and differed between *w*Mel- and *w*Ri-infected

hosts.

Figure 4. Testing the *cif*-expression hypothesis for CI-strength variation. Fold change in
testes across male age for the relative expression of (A) *cifA*_{wMel[T1]} to *D. melanogaster βspec*,
(B) *cifA*_{wMel[T1]} to *w*Mel *ftsZ*, (C) *cifA*_{wMel[T1]} to *cifB*_{wMel[T1]}, (D) *cifA*_{wRi[T1]} to *D. simulans βspec*, (E)
*cifA*_{wRi[T1]} to *w*Ri *ftsZ*, and (F) *cifA*_{wRi[T1]} to *cifB*_{wRi[T1]}. Letters above data represent statistically
significant differences based on α=0.05 calculated by Dunn's test with correction for multiple
comparisons between all groups —crosses that do not share a letter are significantly different.
Fold change was calculated as $2^{-\Delta\Delta cq}$. We selected a random infected sample from the youngest
0-day-old age group as the reference for all fold change analyses within each experiment. *P*-
values are reported in **Table S1**. These data demonstrate that age-dependent *cif* expression is
variably related to host expression, *cif*_{wMel[T1]} expression decreases per *Wolbachia* with age, and
cifA/B relative expression only marginally decreases with age in both systems.

What causes Wolbachia density to vary with age?

We found that *Wolbachia* densities from full-testes extracts significantly increased with male age in *w*Mel-infected *D. melanogaster* and significantly decreased with male age in *w*Ri-infected *D. simulans*. The causes of age-dependent *Wolbachia*-density variation have not been explored. We tested three hypotheses. Namely, that phage lytic activity, Octomom copy number, or host immune expression may govern age-dependent *Wolbachia* densities.

Phage density does not covary with age-dependent Wolbachia density.

The phage density hypothesis predicts that *Wolbachia* density negatively covaries with phage lytic activity (53). Since phage lysis corresponds with increased phage copy number (53, 66), we tested the phage density model by measuring the relative abundance of phage to *Wolbachia ftsZ* using qPCR. *w*Mel and *w*Ri each harbor a unique set of phage haplotypes: *w*Mel has two phages (WOMelA and WOMelB), and *w*Ri has four (WORiA-C, WORiB is duplicated) (100). At least one of *w*Mel's phages is capable of particle production, but it is unknown if *w*Ri's phages yield viral particles (70). We monitored WOMelA and WOMelB of *w*Mel simultaneously using primers that target homologs present in a single copy in each phage. Conversely, we monitored WORiA, WORiB, and WORiC separately since shared homologs are too diverged to make suitable qPCR primers that match multiple phage haplotypes.

First, we evaluated the phage density model for *w*Mel. We predicted the relative abundance of WOMelA/B to decrease with *D. melanogaster* male age since *w*Mel density increases with age. However, there was no change in WOMelA/B abundance relative to *w*Mel *ftsZ* as males aged (**Fig. 5A**; $P = 0.3$), while WOMelA/B abundance relative to *D. melanogaster* UCE increased similar to *w*Mel density (**Fig. S4A**; $P = 3.0\text{E-}04$). Relative phage abundance was not significantly correlated with increasing compatibility (**Table S3**; $r_p = -0.065$, $P = 0.75$; $r_s =$

0.17, *P* = 0.39). Similarly, WOMelA/B significantly varied with age relative to UCE (**Fig. S4B**; *P*

= 0.049) but not *w*Mel *ftsZ* (**Fig. S4C**; *P* = 0.15) in the 0-, 1-, 2-, and 3-day-old age experiment.

Next, we predicted that WORi phage abundance would increase with decreasing *w*Ri

densities across *D. simulans* male age if governed by the phage density model. As with *w*Mel in

D. melanogaster, relative WORiB to *w*Ri *ftsZ* abundance did not significantly covary with male

age (**Fig. 5B**; *P* = 0.053) or correlate with increasing compatibility (**Table S3**; r_p = 0.032, *P* =

0.88; r_s = 0.12, *P* = 0.58). Relative WORiB to *D. simulans* UCE abundance increased with age,

similar to *w*Ri density (**Fig. S4D**; *P* = 4.4E-04). Comparably, WORiA (**Fig. S4E**; *P* = 0.3) and

WORiC (**Fig. S4F**; *P* = 0.4) abundance relative to *w*Ri did not vary with male age. These data

suggest that phage WO is unrelated to age-dependent *Wolbachia*-density variation in *w*Mel and

*w*Ri.

Octomom does not vary with age-dependent wMel density.

Only very closely related *w*Mel variants encode all eight Octomom genes (e.g., *w*Mel,

*w*MelCS, *w*MelPop). The relative abundance of Octomom to *Wolbachia* genes positively

covaries with *w*MelCS and *w*MelPop density (71–75), commonly changing between host

generations. A pair of repeat regions flank the Octomom genes and are hypothesized to be

involved in Octomom amplification. In *w*Mel, the 3' repeat region has a transposon insertion that

likely prevents Octomom amplification (71). As such, we predicted that Octomom copy number

would be invariable with age. We tested if Octomom copy number variation correlated with age-

dependent *w*Mel density variation using qPCR. Indeed, the relative abundance of an Octomom

gene (WD0509) to *w*Mel *ftsZ* did not covary with male age (**Fig. 5C**; *P* = 0.53) or correlate with

increasing compatibility (**Table S3**; r_s = -0.19, *P* = 0.36; r_s = 0.1, P = 0.61). Similar results were

observed in 0-, 1-, 2-, and 3-day-old *w*Mel-infected males **(Fig. S1B; Table S3)**. We conclude

that Octomom copy number is unrelated to the age-dependent increase in *w*Mel densities.

Relish expression is positively correlated with age-dependent wMel, but not wRi, densities.

Theory predicts that natural selection favors the evolution of host genes that suppress CI

(6). Manipulation of *Wolbachia* densities is one mechanism that may drive CI suppression (2).

Since the immune system is designed to control bacterial loads, we investigated the role of the

host immune system in *Wolbachia*-density variation across male age. The immune deficiency

(Imd) pathway is broadly involved in defense against gram-negative bacteria like *Wolbachia*

(101). Bacteria activate the Imd pathway by interacting with peptidoglycan (PG) recognition

proteins which start a signal cascade that results in the expression of the NF-κB transcription

factor Relish (*Rel*). Relish then activates antimicrobial peptide production. *Wolbachia* lack the

full suite of genes needed to synthesize PG (102–104) but can express the PG precursor lipid II

which is sufficient to activate the Imd pathway (104, 105).

We predicted that *D. melanogaster* Relish expression and *w*Mel density would be

correlated if the Imd pathway is involved in *w*Mel density regulation. Indeed, relative expression

of Relish to *βspec* significantly varied among age groups (P = 6.1E-4). However, relative

expression of Relish to *βspec* was lowest in 0-day-old (95% interval = 0.9 - 1.1) infected testes

and consistently increases in 2- (95% interval = 1.1 - 1.8), 4- (95% interval = 1.3 - 1.7), 6- (95%

interval = 1.9 - 2.3), and 8-day-old (95% interval = 1.5 - 3.9) testes (**Fig. 5D**). Relish expression

was significantly positively correlated with *w*Mel density within testes samples (**Fig. 5E**; r_p =

0.77, P = 2.5E-06; r_s = 0.87, P = 1.4E-06). In summary, *w*Mel density was strongly correlated

with increasing Relish expression, directly contrary to our prediction.

Conversely, relative expression of *D. simulans* Relish to *βspec* did not significantly

covary with age (**Fig. 5F**; P = 0.7), but remained positively correlated with the relative

expression of *w*Ri *ftsZ* to *βspec* within testes samples according to Pearson, but not Spearman,

analyses (**Fig. 5G**; r_p = 0.55, P = 0.012; r_s = 0.13, P = 0.59). In summary, Relish expression is

positively correlated with age-dependent *w*Mel densities in *D. melanogaster*, but less so in *w*Ri-

infected *D. simulans*, supporting a role for the Imd pathway in the regulation of at least *w*Mel

density variation. However, more work is necessary to determine if the correlation between age-

dependent immune expression and *Wolbachia* density in testes is causatively associated.

Figure 5. Testing the phage density, Octomom, and host immunity hypotheses for age-dependent *Wolbachia*-density variation. Fold change in testes across male age for the relative abundance or expression of (A) WOMelA/B to *w*Mel *ftsZ*, (B) WORiB to *w*Ri *ftsZ*, (C) Octomom gene WD0509 to *w*Mel *ftsZ*, (D) *D. melanogaster Rel* to *Bspec*, and (F) *D. simulans Rel* to *Bspec*. Correlation between the relative expression of *Rel* to *Bspec* and *ftsZ* to *Bspec* for (E) *w*Mel and (F) *w*Ri. Letters above data represent statistically significant differences based on α=0.05 calculated by Dunn's test with correction for multiple comparisons between all groups — crosses that do not share a letter are significantly different. (E, G) Pearson (top) and Spearman (bottom) correlations are reported. Linear regressions are plotted with 95% confidence intervals. Fold change was calculated as $2^{-\Delta\Delta cq}$. We selected a random infected sample from the youngest 0-day-old age group as the reference for all fold change analyses within each experiment. *P*-

values are reported in **Table S1**. These data demonstrate that age-dependent *Wolbachia*
densities are not controlled by phage WO lysis or Octomom copy number, but are correlated
with *Rel* expression in *D. melanogaster* and less so in *D. simulans*.

Discussion

Within *Wolbachia*-host systems, several factors influence CI strength (29, 30, 37, 38,

53, 66, 86–89), but male age can be particularly impactful (3, 18, 27, 29). Our results determine

how fast and investigate why CI strength declines as males age. First, we estimate that CI-

strength decreases rapidly for *w*Mel-infected *D. melanogaster* (19%/ day), becoming statistically

insignificant when males reach three days old. In contrast, *w*Ri causes intense CI that declines

more slowly (6%/ day), resulting in statistically significant CI through at least the first 12 days of

D. simulans male life. Second, *Wolbachia* densities and *cif* expression from full-testes extracts

increase in *w*Mel-infected *D. melanogaster* and decrease in *w*Ri-infected *D. simulans* as males

age and CI weakens. These results indicate that bacterial density and CI gene expression in

full-testes extracts cannot fully account for age-dependent CI strength across host-*Wolbachia*

associations. Third, while WO phage activity and Octomom copy number cannot explain

Wolbachia-density variation, *D. melanogaster* immune expression covaries with *w*Mel densities,

suggesting the host immune system may contribute to age-dependent *Wolbachia* density in *D.*

melanogaster, but much less so in *D. simulans*. Notably, transcript-based data (e.g., *cif* and

Relish) described here are subject to the caveat that mRNA levels may not correlate perfectly

with protein expression or activity. Future proteomics analyses will be needed to confirm these

trends hold at the protein level. We discuss how our discoveries inform the basis of age-

dependent CI-strength variation, how multiple mechanistic underpinnings may govern age-

dependent *Wolbachia* densities, and how age-dependent CI may contribute to *Wolbachia*

frequency variation observed in nature.

Wolbachia density and CI-gene expression in full-testes extracts do not fully explain age-

dependent CI-strength variation.

Despite support that CI strength is linked to *Wolbachia* density and *cif* expression across

and within systems (37, 38, 51–54, 60, 66), our observations add to a growing body of literature

suggesting *Wolbachia* densities in adult testes (30, 88) and, for the first time, *cif* expression, are

insufficient to explain CI-strength variation broadly. We discuss three hypotheses to explain the

disconnect between *Wolbachia* density and *cif* expression in full-testes extracts and CI strength

with male age. Note, however, that these results may also be explained by a decoupling of *cif*

transcription and protein translation, which will require future proteomics analyses to investigate.

First, the localization and density of *Wolbachia* and *cif* products within specific cells in

testes may more accurately predict CI strength. Indeed, the proportion of infected spermatocyte

cysts covaries with CI strength in natural and transinfected combinations of CI-inducing

Wolbachia and *D. melanogaster*, *D. simulans*, *D. yakuba*, *D. teissieri*, and *D. santomea* (51, 52).

Intriguingly, two *w*Ri-infected *D. simulans* strains whose *Wolbachia* cause variable CI did not

have different *Wolbachia* densities according to qPCR, but the number of infected sperm cysts

covaried with CI between strains (106). Thus, *Wolbachia* densities in full-testes extracts may not

reflect the cyst infection frequency, but it is unknown how generalizable this discrepancy is

across or within *Wolbachia*-host associations with variable CI strengths. It seems plausible that

while *w*Mel densities increase in the testes as males age, the proportion of infected

spermatocytes could decrease. Notably, since *w*Mel infections increase drastically as males

age, a considerable shift in localization and density dynamics would be necessary. Microscopy

assays will be required to test if *Wolbachia* and *cif* localization explains *w*Mel age-dependent CI-

strength variation.

Second, age-dependent CI may be governed by developmental constraints of CI-

susceptibility. For instance, the paternal grandmother age effect, where *Wolbachia*-infected

sons of older virgin females cause stronger CI than sons of younger females, covaries with

Wolbachia densities in embryos but not in adult males (30). Intriguingly, temperature-sensitive

CI-strength variation in *Cardinium*-infected *Encarsia* wasps is also decoupled from symbiont

densities, but CI strongly correlates with pupal development time (107, 108). *Cardinium* CI

effectors likely have more time to interact with host targets at critical stages of pupal

development when slowed by cool temperatures, despite lower *Cardinium* density (107, 108).

These studies suggest that sperm are modified in spermatogenesis before adult eclosion, and

that variation in symbiont densities during early development can contribute to CI-strength

variation. If modified sperm are primarily produced during pupal or larval development, younger

adult males would have a higher proportion of CI-modified sperm in their seminal vesicle than

older males since older males continue to produce sperm as adults. Intriguingly, remating

seems to weaken CI (86, 87), supporting this hypothesis. However, since CI strength decreases

faster in *D. melanogaster* than in *D. simulans*, this hypothesis predicts that adult *D. simulans*

sperm production is slower and/or CI modification occurs for an extended time. Functional work

is necessary to determine if CI modification is developmentally restricted.

Finally, age-dependent CI may be related to the availability of CI-effector targets with

male age and not the abundance of *cif* products. Indeed, the number of genes transcribed by *D.*

melanogaster increases from 7,000 in embryos to over 12,000 in adult males, and nearly a third

of genes are not expressed until the 3rd larval instar (109). As adult males age, the number of

transcribed genes continues to vary, though less so than during metamorphosis (109). These

data support the possibility that host targets of CI may vary in abundance as males age.

However, since transgenic *cif* expression can significantly enhance CI strength above wild-type

levels (60), there are circumstances when natural *cif* expression is not high enough to saturate

all targets. It is unknown if similar experimental approaches can strengthen age-dependent CI.

More work will be necessary to determine the host genes that modify CI and how those factors

vary in expression relative to CI strength.

Age-dependent bacterial density covaries with immune expression, not phage or

Octomom.

We report a strong relationship between male age and *Wolbachia* densities that differ

between systems: densities decrease in *w*Ri-infected *D. simulans* and increase in *w*Mel-infected

D. melanogaster. Reports of age-dependent variation in *Wolbachia* densities across age in

different tissues and sexes are common (51, 71, 90, 91, 110–112), but the basis of this variation

remains unexplored. We investigated the cause(s) of this variation for the first time. We

predicted that genes that covary with age-dependent densities might be causatively linked,

although additional experiments will be necessary to confirm this. First, we tested whether

phage or Octomom covary with age-dependent *Wolbachia* densities. Despite prior reports that

phage WO of *Nasonia* and *Habrobracon Wolbachia* can regulate temperature-dependent

Wolbachia densities (53, 66) and that Octomom copy number correlates with *w*MelCS and

wMelPop densities (72, 73), we found that neither covaries with age-dependent *Wolbachia*

densities in testes.

Next, we asked whether host immune gene expression correlates with age-dependent

Wolbachia densities. We report that Relish expression, which activates antimicrobial peptide

(AMP) production in the Imd pathway (101), strongly correlates with *w*Mel densities and is

highest when *w*Mel densities are high. This result was surprising since we predicted that

immune expression would hinder *Wolbachia* proliferation if it were correlated. Conversely,

Relish does not vary with *D. simulans* male age and is only very weakly correlated with *w*Ri

densities. It is plausible that the correlation between Relish expression and *w*Mel density

represents a spurious and non-causative association. Additionally, Relish transcription does not

necessarily equate to increased Relish activity and AMP production since endoproteolytic

cleavage is necessary to activate the Relish protein (113), but future analysis of AMP

expression will elucidate this. However, this correlation may represent a causative link between

age-dependent *w*Mel densities and immune expression. We propose two non-exclusive

hypotheses to explain this relationship.

First, *w*Mel rapidly proliferates as males age, elicit an immune response proportional to

their infection density, but evade the effects of immune activation. *Wolbachia* synthesize lipid II

(102–104), which is sufficient to activate the Imd pathway (104, 105), and increase AMP gene

expression when transinfection into novel host backgrounds (114–117), suggesting that

Wolbachia can trigger Imd activity. However, Relish and AMP expression do not vary with

Wolbachia infection state (118–124) or density (84, 121) in natural *Wolbachia*-host associations.

It has been proposed that *Wolbachia* evade the host immune system by residing in host-derived

membranes or bacteriocyte-like cells (125, 126). Thus, the correlation between Relish

expression and *w*Mel density may indicate that *w*Mel triggers the immune system but evades

the immune response, preventing its densities from decreasing. Notably, since this hypothesis

assumes that *w*Mel densities increase independent of Imd expression, it does not explain why

*w*Mel densities increase with age or why age-dependent *w*Mel and *w*Ri densities differ.

Second, age-dependent Imd expression increases independent of *Wolbachia* but

impacts *Wolbachia* densities. Aging in *D. melanogaster* is associated with increased expression

of AMPs, Relish, and other immune genes (127–133). Counterintuitively, age also covaries with

increased gut microbial loads and Imd activation in *D. melanogaster* (127–129, 134–136). Why

gut bacterial loads increase with *D. melanogaster* age and immune expression remains

unknown. However, age-dependent immune expression may damage the epithelium, lead to

dysbiosis through differential effects on gut microbial members, alter gut tissue renewal and

differentiation, and/or cause cellular inflammation (101, 137). In other words, the positive

correlation between Relish expression and wMel density may be caused by off-target effects of

immune expression on the cellular environment. To our knowledge, we report the first case

where endosymbiont densities increase with age-dependent immune expression, suggesting

that the cause(s) of age-dependent bacterial proliferation may apply to more than gut microbes.

Functional assays, such as Relish knockdowns, will be necessary to causatively link male age-

dependent *Wolbachia* densities and immune expression.

Age-dependent CI strength could contribute to *Wolbachia* frequency variation in nature.

We can consider our estimates of age-dependent CI strength in the context of an

idealized discrete-generation model of *Wolbachia* frequency dynamics first proposed by

Hoffmann et al. (1990). This model incorporates imperfect maternal transmission (μ), *Wolbachia*

effects on host fitness (F), and the proportion of embryos that hatch in a CI cross relative to

compatible crosses (H) (3). Across all experiments, CI strength ($s_h = 1 - H$) progressively

decreases as males age (**Table S2**): *w*Mel CI strength decreases quickly (Day 0 $s_h = 0.860$; Day

8 $s_h = -0.007$) and *w*Ri CI strength decreases relatively slowly (Day 0 $s_h = 0.991$; Day 8 $s_h =$

0.244). Small negative values of s_h indicate that the CI cross has a slightly higher egg hatch

than the compatible crosses.

*w*Ri occurs globally at high and relatively stable infection frequencies, consistent with

generally strong CI (4, 26), while *w*Mel varies in frequency on several continents. In eastern

Australia, *w*Mel frequencies range from ~90% in the tropical north to ~30% in the temperate

south (34). While the factors that maintain this cline are unresolved, mathematical modeling

suggests clinal differences in CI strength likely contribute (34). For example, CI must be

essentially nonexistent ($s_h \ll 0.05$) to explain relatively low *w*Mel frequencies observed in

temperate Australia, assuming little imperfect transmission ($\mu = 0.01 - 0.026$) (138). Conversely,

with $\mu = 0.026$ and similarly low-to-nonexistent CI ($s_h \leq 0.055$), large and positive *w*Mel effects

on host fitness ($F \sim 1.3$) are required to explain higher *w*Mel frequencies observed in the tropics.

Explaining higher tropical frequencies becomes easier with stronger CI ($s_h > 0.05$) or more

reliable *w*Mel maternal transmission ($\mu < 0.026$) (Kriesner et al. 2016).

So what is *w*Mel CI strength in nature? Field-collected males from near the middle of the

Australian cline to the northern tropics cause very weak ($s_h \sim 0.05$) to no CI (Hoffmann et al.

1998). These, and other data from the middle of the cline (29), led Kriesner et al. (2016) to

conjecture that the plausible range of s_h in subtropical/tropical Australian populations is $s_h = 0$ -

0.05, but < 0.1. In our study, only 6- ($s_h = -0.006$) and 8-day-old ($s_h = -0.007$) wMel-infected

males exhibited CI weaker than $s_h = 0.1$, suggesting that field-collected males causing little or

no CI (138) are older than four days. Interactions among male age, temperature, remating, and

other factors likely contribute to weaker CI in younger males (29, 37, 38, 53, 66, 86, 87). Future

analyses to disentangle the contributions of male age and other factors to CI-strength variation

are sorely needed. These estimates, along with estimates of *Wolbachia* transmission rate

variation across genetic and abiotic contexts (22), are ultimately required to better understand

Wolbachia frequency variation in host populations (7, 22, 24, 34, 139).

Conclusions.

Our results highlight that *Wolbachia* densities and *cif* expression from full-testes extracts

are insufficient to explain age-dependent CI strength. While age-dependent CI strength in wRi

aligns with the bacterial density and CI gene expression hypotheses without the need to

consider other factors, wMel CI strength cannot be explained by either of these hypotheses. We

propose that localization, development, and/or host genetic variation contribute to this

relationship. Moreover, wMel densities increase, and wRi decrease as their respective hosts

age. Neither phage WO nor Octomom explain age-dependent *Wolbachia* density, but variation

in these systems covaries with the expression of the immune gene Relish. This represents the

first report that the host immune system may contribute to variation in *Wolbachia* density in a

natural *Wolbachia*-host association. This work motivates an extensive analysis of *Wolbachia*

and *cif* expression in the context of localization and development and a thorough investigation of

the relationship between host genes and *Wolbachia* density and CI phenotypes. Finally,

Incorporating the age-dependency of CI into future modeling efforts may help improve our ability

to explain temporally and spatially variable *Wolbachia* infection frequencies, as incorporating

temperature effects on *w*Mel-like *Wolbachia* transmission has (22, 24, 140). Ultimately this will

help explain *Wolbachia's* status as the most prevalent endosymbiont in nature.

Materials and Methods

Fly lines

All fly lines used in this study are listed in **Table S4**. Uninfected flies were derived via

tetracycline treatment in prior studies (16, 60). Tetracycline cleared lines were used in

experiments over a year after treatment, mitigating the effects of antibiotics on mitochondria

(141). We regularly confirmed infection status by using PCR to amplify the *Wolbachia* surface

protein (*wsp*). An arthropod-specific 28S rDNA was amplified in a separate reaction and served

as a control for DNA quality and PCR inhibitors (24, 142). The y^1w^1 *D. melanogaster* line was

confirmed to be *w*Mel infected, as opposed to *w*MelCS, using IS5-WD1310 primers (143). DNA

was extracted for infection checks using a squish buffer protocol. Briefly, flies were

homogenized in 50 uL squish buffer per fly (100mL 1M Tris-HCL, 0.0372g EDTA, 0.1461g NaCl,

90 mL H_2O, 150uL Proteinase K), incubated at 65°C for 45m, incubated at 94°C for 4m,

centrifuged for 2m, and the supernatant was used immediately for PCR.

Fly care and maintenance

Flies were reared in vials with 10mL of food made of cornmeal (32.6%), dry corn syrup

(32%), malt extract (20.6%), inactive yeast (7.8%), soy flour (4.5%), and agar (2.6%). Fly stocks

were maintained at 23°C between experiments. Flies used for virgin collections were reared at

25°C, virgin flies were stored at 25°C, and experiments were performed at 25°C. Flies were

always kept on a 12:12 light:dark cycle. Flies were anesthetized using CO_2 for virgin collections

and dissections. During hatch-rate assays, flies were mouth aspirated between vials.

Hatch-rate assays

CI manifests as embryonic death. We measured CI as the percentage of embryos that fail to hatch into larva. Flies used in hatch rates were derived from vials where flies were given ~24hr to lay to control for rearing density (88). In the morning, virgin 6-8 day females were added individually to vials containing a small ice cream spoon filled with fly food. Spoon fly food was prepared as described above, but with blue food coloring added, 0.1g extra agar per 100mL of food, and fresh yeast smeared on top. After 4-5hr of acclimation, a single virgin male was added to each vial. The age of virgin males varied by experiment and cross. Paternal grandmother age was not controlled, but paternal grandmothers were non-virgin when setting up vials for fathers. Since *Wolbachia* densities associated with older paternal grandmothers are reduced upon mating (30), we do not expect variation in paternal grandmother *Wolbachia* densities across experiments or conditions. Vials with paired flies were incubated overnight at 25°C. Flies were then aspirated into new vials with a fresh spoon. Vials were incubated for another 24hr before flies were removed via aspirating. Embryos were counted on spoons immediately after flies were removed. After 48hr, the number of remaining unhatched eggs were counted. The percentage of embryos that hatched was then calculated.

Relative abundance assays

Siblings from hatch-rate assays were collected for DNA extractions. Virgin males were anesthetized and testes were dissected in chilled phosphate-buffered saline (PBS). Five pairs of testes were placed into a single 1.5mL Eppendorf tube and stored at -80°C until processing. All tissue was collected the day after the hatch-rate setup. Tissue was homogenized using a pestle, and the DNeasy Blood and Tissue kit (Qiagen) was used to extract and purify DNA.

689   qPCR was used to measure the relative abundance of host, *Wolbachia*, phage WO, and

690 Octomom DNA. Samples were tested in triplicate using Powerup SYBR Green Master Mix

691 (Applied Biosystems), which contains a ROX passive reference dye. Primers were designed

692 using Primer3 v2.3.7 in Geneious Prime (144). Host primers target an ultraconserved element

693 (UCE) *Mid1* identified previously (96). Phage genes were also identified from prior works (100).

694 Primers for *w*Mel's phages target both WOMelA (WD0288) and WOMelB (WD0634), while

695 those for *w*Ri are unique to a single phage haplotype. WORiA, WORiB, and WORiC were

696 measured with *w*Ri_012460, *w*Ri_005590/*w*Ri_010250, and *w*Ri_006880 primers, respectively.

697 Only *w*Mel has all eight Octomom genes (WD0507-WD0514) (71). We measured *w*Mel

698 Octomom copy number using primers targeting WD0509. Primer sequences and PCR

699 conditions are listed in **Table S5**. Fold difference was calculated as $2^{-\Delta\Delta Ct}$ for each comparison.

700 A random sample in the youngest age group was selected as the reference.

702 ***Gene expression assays***

703   Siblings from hatch-rate assays were collected for RNA extractions. Virgin males were

704 anesthetized, and testes were dissected in chilled RNase-free PBS. Fifteen pairs of testes were

705 placed into a single 2mL tube with 200uL of Trizol and four 3mm glass beads. Tissue was kept

706 on ice between dissections. Samples were then homogenized using a TissueLyser II (Qiagen)

707 at 25Hz for 2m, centrifuged, and stored at -80ºC until processing. All tissue was collected the

708 day after the hatch-rate setup.

709   Samples were thawed, 200uL of additional Trizol was added, and tissue was further

710 homogenized using a TissueLyser II at 25Hz for 2m. RNA was extracted using the Direct-Zol

711 RNA Miniprep kit (Zymo Research) following the manufacturer's recommendations, but with an

712 extra wash step. On-column DNase treatment was not performed. Instead, the 'rigorous'

713 treatment protocol from the DNA-free kit (Ambion) was used to degrade DNA in RNA samples.

714 Samples were confirmed DNA-free using PCR and gel electrophoresis for an arthropod-specific

28S rDNA (24, 142). The Qubit RNA HS Assay Kit (Invitrogen) was used to measure RNA

concentration. Samples within an experiment were diluted to the same concentration. RNA was

converted to cDNA using SuperScript IV VILO Master Mix (Invitrogen) with either 200ng or

500ng of total RNA per reaction depending on the experiment. qRT-PCR was performed using

1ng of cDNA per reaction using Powerup SYBR Green Master Mix (Applied Biosystems). All

samples were tested in triplicate.

Primers for expression included host reference, *Wolbachia* reference, *cif*, and host

immune genes. Primers to *Drosophila* genes for qRT-PCR were selected from FlyPrimerBank

(145). Since *Drosophila* expression patterns change with age (109), a host gene that is

invariable with male age was selected to act as a reference gene for relative expression

analyses. We selected an invariable gene using the *Drosophila* Gene Expression Tool (DGET)

to retrieve modENCODE gene expression data for ribosome and cytoskeletal genes (146).

DGET reports expression as Reads Per Kilobase of transcript, per million mapped reads

(RPKM), and included data for adult males 1, 5, and 30 days after eclosion. β-spec (1 Day = 81

RPKM, 5 Day = 80, 30 Day = 79) was selected because it is largely invariable across age. Our

results confirm invariable expression across male age (**Fig. S2E**; **Fig. S3L**). *D. melanogaster*

and *D. simulans* are identical across *βspec* primer binding sequences. All other primers were

designed using Primer3 in Geneious Prime (144) and are listed in **Table S5**. Fold difference

was calculated as $2^{-\Delta\Delta Ct}$ for each comparison. A random sample in the youngest age group was

selected as the reference.

Statistical analyses

All statistics were performed in R (147). Hatch-rate, relative-abundance, and expression

assays were analyzed using a Kruskal-Wallis followed by a Dunn's test with corrections for

multiple comparisons. Kruskal-Wallis and Dunn's *P*-values are reported in **Table S1**.

Correlations between hatch rate and expression or relative abundance measures were

performed using Pearson and Spearman correlations in GGPubR (148). Correlation statistics

are reported in **Table S3**. 95% confidence intervals were calculated using the classic MeanCI

function in DescTools (149). 95% BCa intervals were calculated using boot.ci in boot (150).

Samples with fewer than ten embryos laid were excluded from hatch-rate analyses. Samples

with a C_q standard deviation exceeding 0.4 between triplicate measures were excluded from

qPCR and qRT-PCR analyses. Figures were created using GGPlot2 (151), and figure

aesthetics were edited in Affinity Designer 1.8 (Serif Europe, Nottingham, UK).

Data availability

All data are made publicly available in the supplement of this manuscript.

Acknowledgments

We thank Michael Turelli for helpful feedback on our experimental design. We also thank

Will Conner for help identifying phage gene targets for qPCR, Mike Hague for support with BCa

estimates of *H*, Kelley Van Vaerenberghe and John Statz for review of earlier versions of the

manuscript, and Tim Wheeler for assisting in the laboratory. Seth Bordenstein provided

comments on a preprint. This work was supported by a National Institutes of Health R35

GM124701 to BSC and a National Science Foundation Postdoctoral Research Fellowship DBI-

2010210 to JDS. Any opinions, findings, conclusions, or recommendations expressed in this

material are those of the authors(s) and do not necessarily reflect the views of the National

Institutes of Health or the National Science Foundation.

References

1. Kaur R, Shropshire JD, Cross KL, Leigh B, Mansueto AJ, Stewart V, Bordenstein SR, Bordenstein SR. 2021. Living in the endosymbiotic world of *Wolbachia*: A centennial review. Cell Host & Microbe 29.

2. Shropshire JD, Leigh B, Bordenstein SR. 2020. Symbiont-mediated cytoplasmic incompatibility: what have we learned in 50 years? eLife 9:e61989.

3. Hoffmann A, Turelli M, Harshman L. 1990. Factors affecting the distribution of cytoplasmic incompatibility in *Drosophila simulans*. Genetics 126:933–948.

4. Kriesner P, Hoffmann AA, Lee SF, Turelli M, Weeks AR. 2013. Rapid sequential spread of two *Wolbachia* variants in *Drosophila simulans*. PLOS Pathogens 9:e1003607.

5. Hoffmann A, Turelli M. 1997., p. 42–80. *In* in Influential Passengers: Inherited Microorganisms and Arthropod Reproduction. Oxford University Press.

6. Turelli M. 1994. Evolution of incompatibility-inducing microbes and their hosts. Evolution 48:1500–1513.

7. Schuler H, Köppler K, Daxböck-Horvath S, Rasool B, Krumböck S, Schwarz D, Hoffmeister TS, Schlick-Steiner BC, Steiner FM, Telschow A, Stauffer C, Arthofer W, Riegler M. 2016. The hitchhiker's guide to Europe: the infection dynamics of an ongoing *Wolbachia* invasion and mitochondrial selective sweep in *Rhagoletis cerasi*. Molecular Ecology 25:1595–1609.

8. Narita S, Shimajiri Y, Nomura M. 2009. Strong cytoplasmic incompatibility and high vertical transmission rate can explain the high frequencies of *Wolbachia* infection in Japanese populations of *Colias erate* poliographus (Lepidoptera: Pieridae). Bull Entomol Res 99:385–391.

9. Hunter MS, Perlman SJ, Kelly SE. 2003. A bacterial symbiont in the Bacteroidetes induces cytoplasmic incompatibility in the parasitoid wasp *Encarsia pergandiella*. Proceedings of the Royal Society B: Biological Sciences 270:2185–2190.

10. Rosenwald LC, Sitvarin MI, White JA. 2020. Endosymbiotic Rickettsiella causes cytoplasmic incompatibility in a spider host. Proceedings of the Royal Society B: Biological Sciences 287:20201107.

11. Takano S, Gotoh Y, Hayashi T. 2021. "Candidatus *Mesenet longicola*": Novel Endosymbionts of *Brontispa longissima* that Induce Cytoplasmic Incompatibility. Microbial Ecology https://doi.org/10/gkbptr.

12. Yen JH, Barr AR. 1973. The etiological agent of cytoplasmic incompatibility in *Culex pipiens*. Journal of Invertebrate Pathology 22:242–250.

13. Zug R, Hammerstein P. 2012. Still a host of hosts for *Wolbachia*: analysis of recent data suggests that 40% of terrestrial arthropod species are infected. PLOS ONE 7:e38544.

14. Weinert LA, Araujo-Jnr EV, Ahmed MZ, Welch JJ. 2015. The incidence of bacterial endosymbionts in terrestrial arthropods. Proc R Soc B 282:20150249.

15. Brownlie JC, Cass BN, Riegler M, Witsenburg JJ, Iturbe-Ormaetxe I, McGraw EA, O'Neill SL. 2009. Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. PLoS Pathog 5:e1000368.

16. Hague MTJ, Caldwell CN, Cooper BS. 2020. Pervasive Effects of *Wolbachia* on Host Temperature Preference. mBio 11.

17. Teixeira L, Ferreira A, Ashburner M. 2008. The bacterial symbiont *Wolbachia* induces
resistance to RNA viral infections in *Drosophila melanogaster*. PLoS Biol 6:e2.
18. Weeks AR, Turelli M, Harcombe WR, Reynolds KT, Hoffmann AA. 2007. From parasite to
mutualist: Rapid evolution of *Wolbachia* in natural populations of *Drosophila*. PLOS
Biology 5:997–1005.
19. Hosokawa T, Koga R, Kikuchi Y, Meng X-Y, Fukatsu T. 2010. *Wolbachia* as a
bacteriocyte-associated nutritional mutualist. Proc Natl Acad Sci USA 107:769–774.
20. Moriyama M, Nikoh N, Hosokawa T, Fukatsu T. 2015. Riboflavin Provisioning Underlies
Wolbachia's Fitness Contribution to Its Insect Host. MBio 6:e01732-01715.
21. Carrington LB, Lipkowitz JR, Hoffmann AA, Turelli M. 2011. A re-examination of
Wolbachia-induced cytoplasmic incompatibility in California *Drosophila simulans*. PLOS
One 6:e22565.
22. Hague MTJ, Mavengere H, Matute DR, Cooper BS. 2020. Environmental and Genetic
Contributions to Imperfect *w*Mel-Like *Wolbachia* Transmission and Frequency Variation.
Genetics 215:1117–1132.
23. Serbus LR, Sullivan W. 2007. A cellular basis for *Wolbachia* recruitment to the host
germline. PLOS Pathogens 3:1930–1937.
24. Cooper BS, Ginsberg PS, Turelli M, Matute DR. 2017. *Wolbachia* in the *Drosophila
yakuba* complex: pervasive frequency variation and weak cytoplasmic incompatibility, but
no apparent effect on reproductive isolation. Genetics 205:333–351.
25. Hoffmann AA, Turelli M, Simmons GM. 1986. Unidirectional incompatibility between
populations of *Drosophila simulans*. Evolution 40:692–701.
26. Turelli M, Cooper BS, Richardson KM, Ginsberg PS, Peckenpaugh B, Antelope CX, Kim
KJ, May MR, Abrieux A, Wilson DA, Bronski MJ, Moore BR, Gao J-J, Eisen MB, Chiu JC,
Conner WR, Hoffmann AA. 2018. Rapid global spread of *w*Ri-like *Wolbachia* across
multiple *Drosophila*. Current Biology 28:963-971.e8.
27. Turelli M, Hoffmann AA. 1995. Cytoplasmic incompatibility in *Drosophila simulans*:
dynamics and parameter estimates from natural populations. Genetics 140:1319–1338.
28. Hoffmann AA. 1988. Partial cytoplasmic incompatibility between two Australian
populations of *Drosophila melanogaster*. Entomologia Experimentalis et Applicata 48:61–
67.
29. Reynolds KT, Hoffmann AA. 2002. Male age, host effects and the weak expression or
non-expression of cytoplasmic incompatibility in *Drosophila* strains infected by maternally
transmitted *Wolbachia*. Genetical Research 80:79–87.
30. Layton EM, On J, Perlmutter JI, Bordenstein SR, Shropshire JD. 2019. Paternal
grandmother age affects the strength of *Wolbachia*-induced cytoplasmic incompatibility in
Drosophila melanogaster. mBio 10.
31. Early AM, Clark AG. 2013. Monophyly of *Wolbachia pipientis* genomes within *Drosophila
melanogaster*: geographic structuring, titre variation and host effects across five
populations. Molecular Ecology 22:5765–5778.
32. Hoffmann AA, Clancy DJ, Merton E. 1994. Cytoplasmic incompatibility in Australian
populations of *Drosophila melanogaster*. Genetics 136:993–999.
33. Ilinsky YuYu, Zakharov IK. 2007. The endosymbiont *Wolbachia* in Eurasian populations
of *Drosophila melanogaster*. Russian Journal of Genetics 43:748.

34.   Kriesner P, Conner WR, Weeks AR, Turelli M, Hoffmann AA. 2016. Persistence of a
851   *Wolbachia* infection frequency cline in *Drosophila melanogaster* and the possible role of
852   reproductive dormancy. Evolution 70:979–997.
853   35.   Webster CL, Waldron FM, Robertson S, Crowson D, Ferrari G, Quintana JF, Brouqui J-M,
854   Bayne EH, Longdon B, Buck AH, Lazzaro BP, Akorli J, Haddrill PR, Obbard DJ. 2015.
855   The Discovery, Distribution, and Evolution of Viruses Associated with *Drosophila*
856   *melanogaster*. PLOS Biology 13:e1002210.
857   36.   Hoffmann AA, Montgomery BL, Popovici J, Iturbe-Ormaetxe I, Johnson PH, Muzzi F,
858   Greenfield M, Durkan M, Leong YS, Dong Y, Cook H, Axford J, Callahan AG, Kenny N,
859   Omodei C, McGraw EA, Ryan PA, Ritchie SA, Turelli M, O'Neill SL. 2011. Successful
860   establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission.
861   Nature 476:454–457.
862   37.   Ross PA, Axford JK, Yang Q, Staunton KM, Ritchie SA, Richardson KM, Hoffmann AA.
863   2020. Heatwaves cause fluctuations in *w*Mel *Wolbachia* densities and frequencies in
864   *Aedes aegypti*. PLOS Neglected Tropical Diseases 14:e0007958.
865   38.   Ross PA, Ritchie SA, Axford JK, Hoffmann AA. 2019. Loss of cytoplasmic incompatibility
866   in *Wolbachia*-infected *Aedes aegypti* under field conditions. PLOS Neglected Tropical
867   Diseases 13:e0007357.
868   39.   Walker T, Johnson PH, Moreira LA, Iturbe-Ormaetxe I, Frentiu FD, McMeniman CJ,
869   Leong YS, Dong Y, Axford J, Kriesner P, Lloyd AL, Ritchie SA, O'Neill SL, Hoffmann AA.
870   2011. The *w*Mel *Wolbachia* strain blocks dengue and invades caged *Aedes aegypti*
871   populations. Nature 476:450–453.
872   40.   Dobson SL, Fox Charles W., Jiggins Francis M. 2002. The effect of *Wolbachia*-induced
873   cytoplasmic incompatibility on host population size in natural and manipulated systems.
874   Proceedings of the Royal Society of London Series B: Biological Sciences 269:437–445.
875   41.   Lees RS, Gilles JR, Hendrichs J, Vreysen MJ, Bourtzis K. 2015. Back to the future: the
876   sterile insect technique against mosquito disease vectors. Current Opinion in Insect
877   Science 10:156–162.
878   42.   Nikolouli K, Colinet H, Renault D, Enriquez T, Mouton L, Gibert P, Sassu F, Cáceres C,
879   Stauffer C, Pereira R, Bourtzis K. 2018. Sterile insect technique and *Wolbachia* symbiosis
880   as potential tools for the control of the invasive species *Drosophila suzukii*. Journal of
881   Pest Science 91:489–503.
882   43.   O'Connor L, Plichart C, Sang AC, Brelsfoard CL, Bossin HC, Dobson SL. 2012. Open
883   release of male mosquitoes infected with a *Wolbachia* biopesticide: field performance and
884   infection containment. PLOS Neglected Tropical Diseases 6:e1797.
885   44.   Crawford JE, Clarke DW, Criswell V, Desnoyer M, Cornel D, Deegan B, Gong K, Hopkins
886   KC, Howell P, Hyde JS, Livni J, Behling C, Benza R, Chen W, Dobson KL, Eldershaw C,
887   Greeley D, Han Y, Hughes B, Kakani E, Karbowski J, Kitchell A, Lee E, Lin T, Liu J,
888   Lozano M, MacDonald W, Mains JW, Metlitz M, Mitchell SN, Moore D, Ohm JR, Parkes
889   K, Porshnikoff A, Robuck C, Sheridan M, Sobecki R, Smith P, Stevenson J, Sullivan J,
890   Wasson B, Weakley AM, Wilhelm M, Won J, Yasunaga A, Chan WC, Holeman J, Snoad
891   N, Upson L, Zha T, Dobson SL, Mulligan FS, Massaro P, White BJ. 2020. Efficient
892   production of male *Wolbachia*-infected *Aedes aegypti* mosquitoes enables large-scale
893   suppression of wild populations. Nature Biotechnology 38:482–492.

894   45.   Zheng X, Zhang D, Li Y, Yang C, Wu Y, Liang X, Liang Y, Pan X, Hu L, Sun Q, Wang X,
895   Wei Y, Zhu J, Qian W, Yan Z, Parker AG, Gilles JRL, Bourtzis K, Bouyer J, Tang M,
896   Zheng B, Yu J, Liu J, Zhuang J, Hu Z, Zhang M, Gong J-T, Hong X-Y, Zhang Z, Lin L, Liu
897   Q, Hu Z, Wu Z, Baton LA, Hoffmann AA, Xi Z. 2019. Incompatible and sterile insect
898   techniques combined eliminate mosquitoes. Nature https://doi.org/10/c8kg.

899   46.   O'Neill SL. 2018. The use of *Wolbachia* by the World Mosquito Program to interrupt
900   transmission of *Aedes aegypti* transmitted viruses. Advances in Experimental Medicine
901   and Biology 1062:355–360.

902   47.   O'Neill SL, Ryan PA, Turley AP, Wilson G, Retzki K, Iturbe-Ormaetxe I, Dong Y, Kenny
903   N, Paton CJ, Ritchie SA, Brown-Kenyon J, Stanford D, Wittmeier N, Anders KL, Simmons
904   CP. 2018. Scaled deployment of *Wolbachia* to protect the community from dengue and
905   other *Aedes* transmitted arboviruses. Gates Open Research 2:36.

906   48.   Utarini A, Indriani C, Ahmad RA, Tantowijoyo W, Arguni E, Ansari MR, Supriyati E,
907   Wardana DS, Meitika Y, Ernesia I, Nurhayati I, Prabowo E, Andari B, Green BR, Hodgson
908   L, Cutcher Z, Rancès E, Ryan PA, O'Neill SL, Dufault SM, Tanamas SK, Jewell NP,
909   Anders KL, Simmons CP, AWED Study Group. 2021. Efficacy of Wolbachia-Infected
910   Mosquito Deployments for the Control of Dengue. N Engl J Med 384:2177–2186.

911   49.   Gesto JSM, Pinto SB, Dias FBS, Peixoto J, Costa G, Kutcher S, Montgomery J, Green
912   BR, Anders KL, Ryan PA, Simmons CP, O'Neill SL, Moreira LA. 2021. Large-Scale
913   Deployment and Establishment of Wolbachia Into the Aedes aegypti Population in Rio de
914   Janeiro, Brazil. Frontiers in Microbiology 12:2113.

915   50.   Breeuwer JA, Werren JH. 1993. Cytoplasmic incompatibility and bacterial density in
916   *Nasonia vitripennis*. Genetics 135:565–574.

917   51.   Clark ME, Veneti Z, Bourtzis K, Karr TL. 2003. *Wolbachia* distribution and cytoplasmic
918   incompatibility during sperm development: the cyst as the basic cellular unit of CI
919   expression. Mechanisms of Development 120:185–198.

920   52.   Veneti Z, Clark ME, Zabalou S, Karr TL, Savakis C, Bourtzis K. 2003. Cytoplasmic
921   incompatibility and sperm cyst infection in different *Drosophila‐Wolbachia* associations.
922   Genetics 164:545–552.

923   53.   Bordenstein SR, Bordenstein SR. 2011. Temperature affects the tripartite interactions
924   between bacteriophage WO, *Wolbachia*, and cytoplasmic incompatibility. PLOS One
925   6:e29106.

926   54.   Foo IJ-H, Hoffmann AA, Ross PA. 2019. Cross-Generational Effects of Heat Stress on
927   Fitness and *Wolbachia* Density in *Aedes aegypti* Mosquitoes. Tropical Medicine and
928   Infectious Disease 4.

929   55.   Jia F-X, Yang M-S, Yang W-J, Wang J-J. 2009. Influence of continuous high temperature
930   conditions on *Wolbachia* infection frequency and the fitness of *Liposcelis tricolor*
931   (Psocoptera: Liposcelididae). Environmental Entomology 38:1365–1372.

932   56.   Binnington KC, Hoffmann AA. 1989. *Wolbachia*-like organisms and cytoplasmic
933   incompatibility in *Drosophila simulans*. Journal of Invertebrate Pathology 54:344–352.

934   57.   Bressac C, Rousset F. 1993. The Reproductive Incompatibility System in *Drosophila*
935   *simulans*: DAPI-Staining Analysis of the *Wolbachia* Symbionts in Sperm Cysts. Journal of
936   Invertebrate Pathology 61:226–230.

58. Clark ME, Veneti Z, Bourtzis K, Karr TL. 2002. The distribution and proliferation of the
intracellular bacteria *Wolbachia* during spermatogenesis in *Drosophila*. Mechanisms of
Development 111:3–15.

59. Karr TL, Yang W, Feder ME. 1998. Overcoming cytoplasmic incompatibility in *Drosophila*.
Proceedings of the Royal Society B: Biological Sciences 265:391–395.

60. LePage DP, Metcalf JA, Bordenstein SR, On J, Perlmutter JI, Shropshire JD, Layton EM,
Funkhouser-Jones LJ, Beckmann J, Bordenstein SR. 2017. Prophage WO genes
recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. Nature
543:243–247.

61. Beckmann J, Ronau JA, Hochstrasser M. 2017. A *Wolbachia* deubiquitylating enzyme
induces cytoplasmic incompatibility. Nature Microbiology 2:17007.

62. Chen H, Ronau JA, Beckmann J, Hochstrasser M. 2019. A *Wolbachia* nuclease and its
binding partner provide a distinct mechanism for cytoplasmic incompatibility. Proceedings
of the National Academy of Sciences 116:22314–22321.

63. Shropshire JD, Rosenberg R, Bordenstein SR. 2021. The impacts of cytoplasmic
incompatibility factor (*cifA* and *cifB*) genetic variation on phenotypes. Genetics 217:1–13.

64. Shropshire JD, Bordenstein SR. 2019. Two-by-one model of cytoplasmic incompatibility:
synthetic recapitulation by transgenic expression of *cifA* and *cifB* in *Drosophila*. PLOS
Genetics 15:e1008221.

65. Shropshire JD, On J, Layton EM, Zhou H, Bordenstein SR. 2018. One prophage WO
gene rescues cytoplasmic incompatibility in *Drosophila melanogaster*. Proceedings of the
National Academy of Sciences 115:4987–4991.

66. Nasehi SF, Fathipour Y, Asgari S, Mehrabadi M. 2021. Environmental Temperature, but
Not Male Age, Affects *Wolbachia* and Prophage WO Thereby Modulating Cytoplasmic
Incompatibility in the Parasitoid Wasp, *Habrobracon Hebetor*. Microbial Ecology
https://doi.org/10/gjxss6.

67. Bordenstein SR, Marshall ML, Fry AJ, Kim U, Wernegreen JJ. 2006. The tripartite
associations between bacteriophage, *Wolbachia*, and arthropods. PLOS Pathogens
2:384–393.

68. Masui S, Kamoda S, Sasaki T, Ishikawa H. 2000. Distribution and evolution of
bacteriophage WO in *Wolbachia*, the endosymbiont causing sexual alterations in
arthropods. Journal of Molecular Evolution 51:491–497.

69. Wright J, Sjostrand F, Portaro J, Barr A. 1978. Ultrastructure of Rickettsia-like
microorganism *Wolbachia pipientis* and associated virus-like bodies in mosquito *Culex*
pipiens. Journal of Ultrastructure Research 63:79–85.

70. Gavotte L, Vavre F, Henri H, Ravallec M, Stouthamer R, Boulétreau M. 2004. Diversity,
distribution and specificity of WO phage infection in Wolbachia of four insect species.
Insect Molecular Biology 13:147–153.

71. Chrostek E, Marialva MSP, Esteves SS, Weinert LA, Martinez J, Jiggins FM, Teixeira L.
2013. *Wolbachia* variants induce differential protection to viruses in *Drosophila*
melanogaster: a phenotypic and phylogenomic analysis. PLOS Genetics 9:e1003896.

72. Duarte EH, Carvalho A, Lopez-Madrigal S, Teixeira L. 2020. Regulation of *Wolbachia*
proliferation by the amplification and deletion of an addictive genomic island. bioRxiv
2020.09.08.288217.

73. Chrostek E, Teixeira L. 2015. Mutualism breakdown by amplification of *Wolbachia* genes.
PLOS Biology 13:e1002065.
74. Chrostek E, Teixeira L. 2018. Within host selection for faster replicating bacterial
symbionts. PLoS One 13:e0191530.
75. Woolfit M, Iturbe-Ormaetxe I, Brownlie JC, Walker T, Riegler M, Seleznev A, Popovici J,
Rancès E, Wee BA, Pavlides J, Sullivan MJ, Beatson SA, Lane A, Sidhu M, McMeniman
CJ, McGraw EA, O'Neill SL. 2013. Genomic evolution of the pathogenic Wolbachia strain,
wMelPop. Genome Biol Evol 5:2189–2204.
76. Hornett EA, Charlat S, Duplouy AMR, Davies N, Roderick GK, Wedell N, Hurst GDD.
2006. Evolution of Male-Killer Suppression in a Natural Population. PLOS Biology 4:e283.
77. Vanthournout B, Hendrickx F. 2016. Hidden suppression of sex ratio distortion suggests
Red queen dynamics between *Wolbachia* and its dwarf spider host. Journal of
Evolutionary Biology 29:1488–1494.
78. Reynolds LA, Hornett EA, Jiggins CD, Hurst GDD. 2019. Suppression of Wolbachia-
mediated male-killing in the butterfly Hypolimnas bolina involves a single genomic region.
PeerJ 7:e7677.
79. Hornett EA, Duplouy AMR, Davies N, Roderick GK, Wedell N, Hurst GDD, Charlat S.
2008. You can't keep a good parasite down: evolution of a male-killer suppressor
uncovers cytoplasmic incompatibility. Evolution 62:1258–1263.
80. Poinsot D, Bourtzis K, Markakis G, Savakis C, Mercot H. 1998. *Wolbachia* transfer from
Drosophila melanogaster into *D. simulans*: Host effect and cytoplasmic incompatibility
relationships. Genetics 150:227–237.
81. Veneti Z, Zabalou S, Papafotiou G, Paraskevopoulos C, Pattas S, Livadaras I, Markakis
G, Herren JK, Jaenike J, Bourtzis K. 2012. Loss of reproductive parasitism following
transfer of male-killing *Wolbachia* to *Drosophila melanogaster* and *Drosophila simulans*.
Heredity (Edinb) 109:306–312.
82. Zabalou S, Apostolaki A, Pattas S, Veneti Z, Paraskevopoulos C, Livadaras I, Markakis
G, Brissac T, Mercot H, Bourtzis K. 2008. Multiple rescue factors within a *Wolbachia*
strain. Genetics 178:2145–2160.
83. Funkhouser-Jones LJ, Opstal EJ van, Sharma A, Bordenstein SR. 2018. The Maternal
Effect Gene *Wds* Controls *Wolbachia* Titer in *Nasonia*. Current Biology 0.
84. Grobler Y, Yun CY, Kahler DJ, Bergman CM, Lee H, Oliver B, Lehmann R. 2018. Whole
genome screen reveals a novel relationship between Wolbachia levels and Drosophila
host translation. PLOS Pathogens 14:e1007445.
85. Lu P, Bian G, Pan X, Xi Z. 2012. *Wolbachia* induces density-dependent inhibition to
dengue virus in mosquito cells. PLOS Neglected Tropical Diseases 6:e1754.
86. Awrahman ZA, Champion de Crespigny F, Wedell N. 2014. The impact of *Wolbachia*,
male age and mating history on cytoplasmic incompatibility and sperm transfer in
Drosophila simulans. Journal of Evolutionary Biology 27:1–10.
87. De Crespigny FEC, Pitt TD, Wedell N. 2006. Increased male mating rate in *Drosophila* is
associated with *Wolbachia* infection. Journal of Evolutionary Biology 19:1964–1972.
88. Yamada R, Floate KD, Riegler M, O'Neill SL. 2007. Male development time influences the
strength of *Wolbachia*-induced cytoplasmic incompatibility expression in *Drosophila*
melanogaster. Genetics 177:801–808.

89. Clancy DJ, Hoffmann AA. 1998. Environmental effects on cytoplasmic incompatibility and
bacterial load in *Wolbachia*-infected *Drosophila simulans*. Entomologia Experimentalis et
Applicata 86:13–24.
90. Duron O, Fort P, Weill M. 2007. Influence of aging on cytoplasmic incompatibility, sperm
modification and *Wolbachia* density in *Culex pipiens* mosquitoes. Heredity 98:368–374.
91. Noda H, Koizumi Y, Zhang Q, Deng K. 2001. Infection density of *Wolbachia* and
incompatibility level in two planthopper species, *Laodelphax striatellus* and *Sogatella
furcifera*. Insect Biochem Mol Biol 31:727–737.
92. Kittayapong P, Mongkalangoon P, Baimai V, O'Neill SL. 2002. Host age effect and
expression of cytoplasmic incompatibility in field populations of Wolbachia-superinfected
Aedes albopictus. Heredity 88:270–274.
93. Calvitti M, Marini F, Desiderio A, Puggioli A, Moretti R. 2015. *Wolbachia* density and
cytoplasmic incompatibility in *Aedes albopictus*: concerns with using artificial *Wolbachia*
infection as a vector suppression tool. PLoS One 10:e0121813.
94. Jamnongluk W, Kittayapong P, Baisley KJ, O'Neill SL. 2000. Wolbachia Infection and
Expression of Cytoplasmic Incompatibility in Armigeres subalbatus (Diptera: Culicidae).
Journal of Medical Entomology 37:53–57.
95. Reynolds KT, Thomson LJ, Hoffmann AA. 2003. The effects of host age, host nuclear
background and temperature on phenotypic effects of the virulent *Wolbachia* strain
popcorn in *Drosophila melanogaster*. Genetics 164:1027–1034.
96. Kern AD, Barbash DA, Chang Mell J, Hupalo D, Jensen A. 2015. Highly constrained
intergenic *Drosophila* ultraconserved elements are candidate ncRNAs. Genome Biology
and Evolution 7:689–698.
97. Lindsey A, Rice DW, Bordenstein SR, Brooks AW, Bordenstein SR, Newton ILG. 2018.
Evolutionary genetics of cytoplasmic incompatibility genes *cifA* and *cifB* in prophage WO
of *Wolbachia*. Genome Biology and Evolution 10:434–451.
98. Bing X-L, Zhao D-S, Sun J-T, Zhang K-J, Hong X-Y. 2020. Genomic analysis of
Wolbachia from *Laodelphax striatellus* (Delphacidae, Hemiptera) reveals insights into its
"Jekyll and Hyde" mode of infection pattern. Genome Biology and Evolution
https://doi.org/10/gkbpnf.
99. Martinez J, Klasson L, Welch JJ, Jiggins FM. 2020. Life and death of selfish genes:
comparative genomics reveals the dynamic evolution of cytoplasmic incompatibility.
Molecular Biology and Evolution https://doi.org/10/gjfqg8.
100. Bordenstein SR, Bordenstein SR. 2016. Eukaryotic association module in phage WO
genomes from *Wolbachia*. Nature Communications 7:13155.
101. Myllymäki H, Valanne S, Rämet M. 2014. The *Drosophila* Imd Signaling Pathway. The
Journal of Immunology 192:3455–3462.
102. Hotopp JCD, Lin M, Madupu R, Crabtree J, Angiuoli SV, Eisen J, Seshadri R, Ren Q, Wu
M, Utterback TR, Smith S, Lewis M, Khouri H, Zhang C, Niu H, Lin Q, Ohashi N, Zhi N,
Nelson W, Brinkac LM, Dodson RJ, Rosovitz MJ, Sundaram J, Daugherty SC, Davidsen
T, Durkin AS, Gwinn M, Haft DH, Selengut JD, Sullivan SA, Zafar N, Zhou L, Benahmed
F, Forberger H, Halpin R, Mulligan S, Robinson J, White O, Rikihisa Y, Tettelin H. 2006.
Comparative Genomics of Emerging Human Ehrlichiosis Agents. PLOS Genetics 2:e21.

103.   Vollmer J, Schiefer A, Schneider T, Jülicher K, Johnston KL, Taylor MJ, Sahl H-G,
1069   Hoerauf A, Pfarr K. 2013. Requirement of lipid II biosynthesis for cell division in cell wall-
1070   less *Wolbachia*, endobacteria of arthropods and filarial nematodes. International journal
1071   of medical microbiology: IJMM 303:140–149.
1072   104.   Wilmes M, Meier K, Schiefer A, Josten M, Otten CF, Klöckner A, Henrichfreise B, Vollmer
1073   W, Hoerauf A, Pfarr K. 2017. AmiD Is a Novel Peptidoglycan Amidase in Wolbachia
1074   Endosymbionts of Drosophila melanogaster. Front Cell Infect Microbiol 0.
1075   105.   Stenbak CR, Ryu J-H, Leulier F, Pili-Floury S, Parquet C, Hervé M, Chaput C, Boneca IG,
1076   Lee W-J, Lemaitre B, Mengin-Lecreulx D. 2004. Peptidoglycan Molecular Requirements
1077   Allowing Detection by the Drosophila Immune Deficiency Pathway. The Journal of
1078   Immunology 173:7339–7348.
1079   106.   Clark ME, Karr TL. 2002. Distribution of *Wolbachia* within *Drosophila* reproductive tissue:
1080   implications for the expression of cytoplasmic incompatibility. Integrative and comparative
1081   biology 42:332–339.
1082   107.   Doremus MR, Stouthamer CM, Kelly SE, Schmitz-Esser S, Hunter MS. 2020. *Cardinium*
1083   Localization During Its Parasitoid Wasp Host's Development Provides Insights Into
1084   Cytoplasmic Incompatibility. Frontiers in Microbiology 11.
1085   108.   Doremus MR, Kelly SE, Hunter MS. 2019. Exposure to opposing temperature extremes
1086   causes comparable effects on *Cardinium* density but contrasting effects on *Cardinium*-
1087   induced cytoplasmic incompatibility. PLOS Pathogens 15:e1008022.
1088   109.   Graveley BR, Brooks AN, Carlson JW, Duff MO, Landolin JM, Yang L, Artieri CG, van
1089   Baren MJ, Boley N, Booth BW, Brown JB, Cherbas L, Davis CA, Dobin A, Li R, Lin W,
1090   Malone JH, Mattiuzzo NR, Miller D, Sturgill D, Tuch BB, Zaleski C, Zhang D, Blanchette
1091   M, Dudoit S, Eads B, Green RE, Hammonds A, Jiang L, Kapranov P, Langton L,
1092   Perrimon N, Sandler JE, Wan KH, Willingham A, Zhang Y, Zou Y, Andrews J, Bickel PJ,
1093   Brenner SE, Brent MR, Cherbas P, Gingeras TR, Hoskins RA, Kaufman TC, Oliver B,
1094   Celniker SE. 2011. The Developmental Transcriptome of *Drosophila melanogaster*.
1095   Nature 471:473–479.
1096   110.   Kaur R, Martinez J, Rota-Stabelli O, Jiggins FM, Miller WJ. 2020. Age, tissue, genotype
1097   and virus infection regulate *Wolbachia* levels in *Drosophila*. Molecular Ecology 29.
1098   111.   Tortosa P, Charlat S, Labbe P, Dehecq J-S, Barre H, Weill M. 2010. *Wolbachia* age-sex-
1099   specific density in *Aedes albopictus*: a host evolutionary response to cytoplasmic
1100   incompatibility? PLoS One 5:e9700.
1101   112.   Unckless RL, Boelio LM, Herren JK, Jaenike J. 2009. Wolbachia as populations within
1102   individual insects: causes and consequences of density variation in natural populations.
1103   Proc Biol Sci 276:2805–2811.
1104   113.   Stöven S, Ando I, Kadalayil L, Engström Y, Hultmark D. 2000. Activation of the
1105   Drosophila NF-κB factor Relish by rapid endoproteolytic cleavage. EMBO Rep 1:347–
1106   352.
1107   114.   Bian G, Xu Y, Lu P, Xie Y, Xi Z. 2010. The endosymbiotic bacterium *Wolbachia* induces
1108   resistance to dengue virus in *Aedes aegypti*. PLOS Pathogens 6:e1000833.
1109   115.   Kambris Z, Cook PE, Phuc HK, Sinkins SP. 2009. Immune activation by life-shortening
1110   *Wolbachia* and reduced filarial competence in mosquitoes. Science 326:134–136.

116.   Moreira LA, Iturbe-Ormaetxe I, Jeffery JA, Lu G, Pyke AT, Hedges LM, Rocha BC, Hall-
Mendelin S, Day A, Riegler M, Hugo LE, Johnson KN, Kay BH, McGraw EA, van den
Hurk AF, Ryan PA, O'Neill SL. 2009. A *Wolbachia* symbiont in *Aedes aegypti* limits
infection with dengue, Chikungunya, and Plasmodium. Cell 139:1268–1278.
117.   Xi Z, Gavotte L, Xie Y, Dobson SL. 2008. Genome-wide analysis of the interaction
between the endosymbiotic bacterium *Wolbachia* and its *Drosophila* host. BMC
Genomics 9:1.
118.   Wang L, Zhou C, He Z, Wang Z-G, Wang J-L, Wang Y-F. 2012. Wolbachia infection
decreased the resistance of Drosophila to lead. PLoS One 7:e32643.
119.   Lindsey ARI, Bhattacharya T, Hardy RW, Newton ILG. 2021. Wolbachia and Virus Alter
the Host Transcriptome at the Interface of Nucleotide Metabolism Pathways. mBio 12.
120.   Bourtzis K, Pettigrew MM, O'Neill SL. 2000. *Wolbachia* neither induces nor suppresses
transcripts encoding antimicrobial peptides. Insect molecular biology 9:635–639.
121.   Rancès E, Johnson TK, Popovici J, Iturbe-Ormaetxe I, Zakir T, Warr CG, O'Neill SL.
2013. The toll and Imd pathways are not required for *Wolbachia*-mediated dengue virus
interference. Journal of Virology 87:11945–11949.
122.   Wong ZS, Hedges LM, Brownlie JC, Johnson KN. 2011. *Wolbachia*-mediated
antibacterial protection and immune gene regulation in *Drosophila*. PLOS One 6:e25430.
123.   Zhang Y-K, Ding X-L, Rong X, Hong X-Y. 2015. How do hosts react to endosymbionts? A
new insight into the molecular mechanisms underlying the *Wolbachia*-host association.
Insect Molecular Biology 24:1–12.
124.   Simhadri RK, Fast EM, Guo R, Schultz MJ, Vaisman N, Ortiz L, Bybee J, Slatko BE,
Frydman HM. 2017. The Gut Commensal Microbiome of Drosophila melanogaster Is
Modified by the Endosymbiont Wolbachia. mSphere 2:e00287-17.
125.   Louis C, Nigro L. 1989. Ultrastructural evidence of *Wolbachia* rickettsiales in *Drosophila*
simulans and their relationships with unidirectional cross-incompatibility. Journal of
Invertebrate Pathology 54:39–44.
126.   Sacchi L, Genchi M, Clementi E, Negri I, Alma A, Ohler S, Sassera D, Bourtzis K, Bandi
C. 2010. Bacteriocyte-like cells harbour Wolbachia in the ovary of Drosophila
melanogaster (Insecta, Diptera) and Zyginidia pullula (Insecta, Hemiptera). Tissue Cell
42:328–333.
127.   Broderick NA, Buchon N, Lemaitre B. 2014. Microbiota-induced changes in *Drosophila*
melanogaster host gene expression and gut morphology. mBio 5:e01117-01114.
128.   Buchon N, Broderick NA, Chakrabarti S, Lemaitre B. 2009. Invasive and indigenous
microbiota impact intestinal stem cell activity through multiple pathways in *Drosophila*.
Genes & Development 23:2333–2344.
129.   Guo L, Karpac J, Tran SL, Jasper H. 2014. PGRP-SC2 promotes gut immune
homeostasis to limit commensal dysbiosis and extend lifespan. Cell 156:109–122.
130.   Ha E-M, Lee K-A, Park SH, Kim S-H, Nam H-J, Lee H-Y, Kang D, Lee W-J. 2009.
Regulation of DUOX by the Galphaq-phospholipase Cbeta-Ca2+ pathway in *Drosophila*
gut immunity. Developmental Cell 16:386–397.
131.   Landis GN, Abdueva D, Skvortsov D, Yang J, Rabin BE, Carrick J, Tavaré S, Tower J.
2004. Similar gene expression patterns characterize aging and oxidative stress in

Drosophila melanogaster. Proceedings of the National Academy of Sciences 101:7663–
7668.

132. Pletcher SD, Macdonald SJ, Marguerie R, Certa U, Stearns SC, Goldstein DB, Partridge
L. 2002. Genome-Wide Transcript Profiles in Aging and Calorically Restricted *Drosophila*
melanogaster. Current Biology 12:712–723.

133. Seroude L, Brummel T, Kapahi P, Benzer S. 2002. Spatio-temporal analysis of gene
expression during aging in *Drosophila melanogaster*. Aging Cell 1:47–56.

134. Clark RI, Salazar A, Yamada R, Fitz-Gibbon S, Morselli M, Alcaraz J, Rana A, Rera M,
Pellegrini M, Ja WW, Walker DW. 2015. Distinct Shifts in Microbiota Composition during
Drosophila Aging Impair Intestinal Function and Drive Mortality. Cell Reports 12:1656–
1667.

135. Mistry R, Kounatidis I, Ligoxygakis P. 2017. Interaction Between Familial Transmission
and a Constitutively Active Immune System Shapes Gut Microbiota in *Drosophila*
melanogaster. Genetics 206:889–904.

136. Ren C, Webster P, Finkel SE, Tower J. 2007. Increased Internal and External Bacterial
Load during *Drosophila* Aging without Life-Span Trade-Off. Cell Metabolism 6:144–152.

137. Buchon N, Silverman N, Cherry S. 2014. Immunity in *Drosophila melanogaster*--from
microbial recognition to whole-organism physiology. Nature Reviews Immunology
14:796–810.

138. Hoffmann AA, Hercus M, Dagher H. 1998. Population dynamics of the *Wolbachia*
infection causing cytoplasmic incompatibility in *Drosophila melanogaster*. Genetics
148:221–231.

139. Wheeler TB, Thompson V, Conner WR, Cooper BS. 2021. *Wolbachia* in the spittlebug
Prosapia ignipectus: Variable infection frequencies, but no apparent effect on host
reproductive isolation. bioRxiv 2021.02.25.432892.

140. Cooper BS, Vanderpool D, Conner WR, Matute DR, Turelli M. 2019. *Wolbachia*
acquisition by *Drosophila yakuba*-clade hosts and transfer of incompatibility loci between
distantly related *Wolbachia*. Genetics https://doi.org/10/ggh6t7.

141. Ballard JWO, Melvin RG. 2007. Tetracycline treatment influences mitochondrial
metabolism and mtDNA density two generations after treatment in *Drosophila*. Insect
Molecular Biology 16:799–802.

142. Meany MK, Conner WR, Richter SV, Bailey JA, Turelli M, Cooper BS. 2019. Loss of
cytoplasmic incompatibility and minimal fecundity effects explain relatively low *Wolbachia*
frequencies in *Drosophila mauritiana*. Evolution https://doi.org/10/gkbpks.

143. Riegler M, Sidhu M, Miller WJ, O'Neill SL. 2005. Evidence for a global *Wolbachia*
replacement in *Drosophila melanogaster*. Current Biology 15:1428–1433.

144. Kõressaar T, Lepamets M, Kaplinski L, Raime K, Andreson R, Remm M. 2018.
Primer3_masker: integrating masking of template sequence with primer design software.
Bioinformatics 34:1937–1938.

145. Hu Y, Sopko R, Foos M, Kelley C, Flockhart I, Ammeux N, Wang X, Perkins L, Perrimon
N, Mohr SE. 2013. FlyPrimerBank: an online database for *Drosophila melanogaster* gene
expression analysis and knockdown evaluation of RNAi reagents. G3 3:1607–1616.

146. Hu Y, Comjean A, Perrimon N, Mohr S. 2017. The *Drosophila* Gene Expression Tool
(DGET) for expression analyses. BMC Bioinformatics 18:98.

147.   R Core Team. 2020. R: A Language and Environment for Statistical Computing. R
1199   Foundation for Statistical Computing, Vienna, Austria.
1200   148.   Kassambara A. 2020. ggpubr: "ggplot2" Based Publication Ready Plots.
1201   149.   Signorell A. 2021. Tools for Descriptive Statistics [R package DescTools version 0.99.41].
1202   Comprehensive R Archive Network (CRAN).
1203   150.   Canty A, Ripley BD. 2021. boot: Bootstrap R (S-Plus) Functions.
1204   151.   Wickham H. 2016. ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New
1205   York.

1206

Supporting information

Figure S1. Testing the bacterial density and Octomom copy number hypotheses for CI strength variation in young *w*Mel-infected *D. melanogaster*. Fold change across male age for (A) *w*Mel *ftsZ* relative to *D. melanogaster* UCE and (B) Octomom gene WD0509 to *w*Mel *ftsZ*. Letters above data represent statistically significant differences based on α=0.05 calculated by Dunn's test with correction for multiple comparisons between all groups —crosses that do not share a letter are significantly different. Fold change was calculated as $2^{-\Delta\Delta cq}$. *P*-values are reported in Table S1.

Figure S2. Testing the *cif*-expression hypothesis for *w*Mel CI strength variation. Fold change across male age for the relative expression of (A) $cifB_{wMel[T1]}$ to *D. melanogaster βspec* and (B) $cifB_{wMel[T1]}$ to *w*Mel *ftsZ*. Raw C_q values for (C) $cifA_{wMel[T1]}$, (D) $cifB_{wMel[T1]}$, (E) *D. melanogaster βspec*, and (D) *w*Mel *ftsZ*. Letters above data represent statistically significant differences based on α=0.05 calculated by Dunn's test with correction for multiple comparisons between all groups —crosses that do not share a letter are significantly different. Fold change was calculated as $2^{-\Delta\Delta cq}$. *P*-values are reported in Table S1.

Figure S3. Testing the *cif*-expression hypothesis for *w*Ri CI strength variation. Fold change across male age for the relative expression of (A) $cifB_{wRi[T1]}$ to *D. simulans βspec*, (B) $cifB_{wRi[T1]}$ to *w*Ri *ftsZ*, (C) $cifA_{wRi[T2]}$ to *D. simulans βspec*, (D) $cifA_{wRi[T2]}$ to *w*Ri *ftsZ*, (E) $cifB_{wRi[T2]}$ to *D. simulans βspec*, (F) $cifB_{wRi[T2]}$ to *w*Ri *ftsZ*, and (G) $cifA_{wRi[T1]}$ to $cifA_{wRi[T2]}$. Raw C_q values for (H) $cifA_{wRi[T1]}$, (I) $cifB_{wRi[T1]}$, (J) $cifA_{wRi[T2]}$, (K) $cifB_{wRi[T2]}$, (L) *D. simulans βspec*, and (M) *w*Ri *ftsZ*. Letters above data represent statistically significant differences based on α=0.05 calculated by Dunn's test with correction for multiple comparisons between all groups —crosses that do not share a letter are significantly different. Fold change was calculated as $2^{-\Delta\Delta cq}$. *P*-values are reported in Table S1.

Figure S4. Testing the phage density model for *Wolbachia*-density variation. Fold change across male age for the relative abundance of (A) WOMelA/B to *D. melanogaster* UCE in the 0-8 day old male age experiment, (B) WOMelA/B to *D. melanogaster* UCE in the 0-3 day old male age experiment, (C) WOMelA/B to *w*Mel *ftsZ* in the 0-3 day old male age experiment, (D) WORiB to *D. simulans* UCE, (E) WORiA to *w*Ri *ftsZ*, and (F) WORiC to *w*Ri *ftsZ*. Letters above data represent statistically significant differences based on α=0.05 calculated by Dunn's test with correction for multiple comparisons between all groups —crosses that do not share a letter are significantly different. Fold change was calculated as $2^{-\Delta\Delta cq}$. *P*-values are reported in Table S1.

Table S1. *P*-values associated with all statistical comparisons.

Table S2. Parameter estimates of H and Sh. Hatch-rate data used for estimates are derived from experiments in Figure 2.

Table S3. Correlations between relative abundance or relative expression and hatch rate. Correlations are organized by the relevant hypothesis: bacterial density, phage density, host immunity, Octomom, and *cif* expression. Factors positively correlated with hatch rate are negatively correlated with CI strength, and vice versa. Pearson and Spearman *P*-values are bold if < 0.05, underlined if < 0.01, and double-underlined if < 0.001. *w*Mel "old cohort" refers to the experiment presented in Fig. 2B where 0-, 2-, 4-, 6-, and 8-day-old males are tested for CI. *w*Mel "young cohort" refers to the experiment in Fig. 2A where 0-, 1-, 2-, and 3-day-old males

1256 are tested for CI. *w*Ri data comes from siblings of males used in Fig. 2C. Median hatch rates
1257 were used for correlations and were taken from relevant experiments in Fig. 2.
1258
1259 **Table S4.** ***Drosophila*** **lines used in this study.**
1260
1261 **Table S5. Primers used in this study.**
1262
1263 **Supporting data file 1. All hatch-rate and qPCR data generated in this study.**