Sequence features of retrotransposons allow for epigenetic variability

Kevin R. Costello[1,2], Amy Leung[1], Candi Trac[1], Michael Lee[1,2], Mudasar Basam[1], J. Andrew

Pospisilik[3] and Dustin E. Schones[1,2]*

[1]Department of Diabetes Complications and Metabolism, Beckman Research Institute

[2]Irell and Manella Graduate School of Biological Sciences, City of Hope, Duarte, CA 91010

[3]Van Andel Research Institute, Grand Rapids, MI, 49503, USA.

Abstract

Transposable elements (TEs) are mobile genetic elements that make up a large fraction of

mammalian genomes. While select TEs have been co-opted in host genomes to have function,

the majority of these elements are epigenetically silenced by DNA methylation in somatic cells.

However, some TEs in mice, including the Intracisternal A-particle (IAP) subfamily of

retrotransposons, have been shown to display interindividual variation in DNA methylation.

Recent work has revealed that IAP sequence differences and strain-specific KRAB zinc finger

proteins (KZFPs) may influence the methylation state of these IAPs. However, the mechanisms

underlying the establishment and maintenance of interindividual variability in DNA methylation

still remain unclear. Here we report that sequence content and genomic context influence the

likelihood that IAPs become variably methylated. IAPs that differ from consensus IAP

sequences have altered KZFP recruitment that can lead to decreased KAP1 recruitment when

in proximity of constitutively expressed genes. These variably methylated loci have a high CpG

density, similar to CpG islands, and can be bound by ZF-CxxC proteins, providing a potential

mechanism to maintain this permissive chromatin environment and protect from DNA

methylation. These observations indicate that variably methylated IAPs escape silencing

through both attenuation of KZFP binding and recognition by ZF-CxxC proteins to maintain a

hypomethylated state.

Introduction

Sequences derived from transposable elements (TEs) make up a large fraction of mammalian genomes (Bourque et al., 2018). TEs are silenced by both histone modifications and DNA methylation to prevent these elements from having deleterious impact on the host genome. Certain TEs, however, can escape silencing and behave as sites of interindividual epigenetic variability, or (Bakshi et al., 2016; Dolinoy et al., 2007a; Gunasekara et al., 2019; Hernando-Herraez et al., 2015; Oey et al., 2015; Reiss et al., 2010). These loci have been reported to be sensitive to environmental stimuli such as diet and exposure to toxins (Dolinoy et al., 2007b; Morgan et al., 1999; Rosenfeld et al., 2013; Waterland et al., 2007; Waterland and Jirtle, 2003), though this field remains an area of active research (Bertozzi et al., 2021). The well-most studied metastable epiallele is present in the Agouti viable yellow mouse, where genetically identical mice display phenotypic diversity that manifests as a range of coat colors and susceptibility to obesity (Bernal et al., 2011; Rakyan et al., 2002). This variable coat color has been linked to the methylation state of a novel insertion of an Intracisternal A-particle (IAP) retrotransposon (Dolinoy et al., 2010). Variable methylation of this IAP contributes to variable transcription of the *agouti* gene across all tissues and variation in coat color, hyperphagia and hyperinsulinemia (Bernal et al., 2011; Rakyan et al., 2002). These observations in the Agouti viable mouse model highlights the impact of variable methylation of TEs on phenotypic outputs. Thus, elucidating the mechanisms underlying the establishment and maintenance of these variably methylated TEs is important to further our understanding of non-Mendelian phenotypic variability and epigenetic plasticity (Ecker et al., 2018, 2017; Tejedor and Fraga, 2017).

Recent work profiling interindividual epigenetic variability in mice identified roughly 50 loci that are variably methylated in a manner similar to the IAP that drives the agouti mouse phenotype (Adams et al., 2012; Bertozzi et al., 2020; Elmer et al., 2020; Kazachenka et al., 2018). (Adams et al., 2012; Kazachenka et al., 2018). The majority of these variably methylated loci are IAP

retrotransposons (VM-IAPs) (Bertozzi et al., 2020; Faulk et al., 2013; Kazachenka et al., 2018).

IAPs are silenced in a sequence-specific manner by KRAB-domain containing zinc finger

proteins (KZFPs) (Coluccio et al., 2018). KZFPs recruit KAP1, also known as TRIM28, which, in

turn, recruits repressive protein complexes to deposit H3K9me3 and DNA methylation at these

loci (Bulut-Karslioglu et al., 2014; Ecco et al., 2017, 2016; Schultz et al., 2002; Turelli et al.,

2014). While this has been thought to happen primarily in embryonic stem cells, recent work has

demonstrated that KZFPs and KAP1 regulated TEs in somatic cells as well (Ecco et al., 2016).

KZFPs have evolved in clusters through imperfect gene duplication and spontaneous mutations

that can allow for the recognition of novel TEs (Imbeault et al., 2017; Kauzlaric et al., 2017).

Over time, TEs can mutate and escape from KZFP targeted silencing (Imbeault et al., 2017;

Kauzlaric et al., 2017; Wolf et al., 2020). It has been proposed that these KZFPs are engaged in

an evolutionary arms race with TEs to maintain their silencing (Jacobs et al., 2014).

IAPs have an unusually high CpG content that is more similar to CpG islands (CGIs) than to the

rest of the mouse genome (Elmer et al., 2020; Kazachenka et al., 2018). CGIs are generally

hypomethylated and can recruit ZF-CxxC proteins, which help maintain a permissive chromatin

environment (Clouaire et al., 2012; Gu et al., 2018; Mikkelsen et al., 2007; Thomson et al.,

2010; van de Lagemaat et al., 2018). Notable CxxC domain containing proteins include CFP1,

which is a subunit of the SET1 complex that deposit H3K4me3, and TET1/TET3, which are

responsible for the active removal of methylation from CpG dinucleotides to maintain a

hypomethylated state (Gu et al., 2018; Tahiliani et al., 2009; Williams et al., 2011). Mammalian

genomes in general have an underrepresentation of CpG dinucleotides as a result of

deamination at methylated cytosines over evolutionary time (Duncan and Miller, 1980; Lewis et

al., 2016; Shen et al., 1994). The "richness" of CpG dinucleotides at IAPs is potentially due to

the fact these elements are more recent additions to the genome that have yet to undergo

'evolutionary' deamination at methylated CpG dinucleotides. It has been previously shown that

recently evolved CpG dense TEs have the potential to be hypomethylated when there is no

targeted suppression of these elements (Jacobs et al., 2014; Long et al., 2016; Ward et al.,

2013).

Based on this evidence, we postulated that variably methylated CpG dense IAPs can have

sequence variants that allow for partial escape from KZFP-mediated silencing (Bertozzi et al.,

2020; Faulk et al., 2013; Kazachenka et al., 2018) and subsequently be bound by ZF-CxxC

proteins, thereby preventing stable repression. Focusing on the IAPLTR1 subfamily (identified

as IAPLTR1_Mm in RepBase,, we performed multiple sequence alignment and hierarchical

clustering of these elements and confirmed that the VM-IAPs have a unique sequence that is

distinct from other IAPLTRs. These IAPLTRs have altered KZFP binding and diminished KAP1

recruitment. Relative to silenced IAPs - that are found in gene poor regions or proximal to tissue

specific transcripts – we find that VM-IAPs are over-represented proximal to constitutively

expressed genes. Importantly, we find that CpG dense TEs (regardless of subfamily) have

increased recruitment of ZF-CxxC proteins such as CFP1 and TET1 in the absence of KZFP-

mediated silencing.

Results

IAP sequence influences KZFP recruitment and establishment of variable methylation

Recent work has identified novel VM loci, which are largely IAPs of the IAPLTR1 and IAPLTR2

subfamilies, across tissues in C57BL/6J mice (Bertozzi et al., 2020; Elmer et al., 2020;

Kazachenka et al., 2018). IAPLTR1 and IAPLTR2 are among the most evolutionarily recent

IAPs (Qin et al., 2010) As KZFPs are responsible for silencing IAP elements in a sequence-

specific manner (Coluccio et al., 2018), we examined if the previously identified VM-IAPs have

shared sequence variants that could allow for escape from KZFP-mediated silencing at both the

LTR and internal element. While many older IAPs have lost their internal elements over time,

many IAPLTR1 and IAPLTR2, which among the most evolutionarily recent IAPLTRs, flank an

IAPEz-int (Qin et al., 2010; Stoye, 2001). We performed multiple sequence alignment and

hierarchical clustering of all LTRs belonging to the IAPLTR1 and IAPLTR2 subfamilies that were

greater than 300 bps long using the ETE3 pipeline (Huerta-Cepas et al., 2016). This analysis

treated all LTRs independently, regardless of whether the LTR was a solo LTR or part of an

ERV. We also performed multiple sequence alignment and hierarchical clustering the first

150bps of IAPEz-ints flanked by either IAPLTR1 or IAPLTR2 elements, which has previously

been show to recruit the KZFP Gm14419 (Wolf et al., 2020). This profiling resulted in the

characterization of four sequence 'clades' for IAPLTR1s (Figure 1A, Figure 1 - figure

supplement 1,2 and supplement file 1), two sequence 'clades' for the IAPLTR2s (Figure 1 -

figure supplement 1 and supplemental file 1), and three sequence clades for IAPEz-ints. All

IAPLTRs flanking an IAPEz-int belonged to the same sequence clade.

To determine whether these clades have altered KZFP recruitment, we used published KZFP

ChIP-seq profiles (Wolf et al., 2020) to survey the occupancy of KZFPs at each identified clade.

We aligned all ChIP-seq libraries to the mm10 genome with multimapping, but allowed no

SNPs, as we were specifically interested in identifying which sequences the KZFPs could bind

(see Methods for details). We identified that clade 1 IAPLTR1 elements have evidence of

binding by ZFP989 and Gm21082, while clade 2 IAPLTR1 elements have evidence of binding

by ZFP429 (Figure 1A and figure supplement 3). However, clade 3 and clade 4 IAPLTR1

elements do not show evidence of binding by these KZFPs and have decreased KAP1 binding

(Figure 1A,B and figure supplement 3). Similar analysis with IAPLTR2s demonstrated that clade

1 IAPLTR2s are bound by ZFP429, while clade 2 IAPLTR2 elements are largely bound by

ZFP989 (Figure 1 - figure supplement 4). Both clades have evidence of KAP1 binding (Figure 1

– figure supplement 4). When profiling KZFP and KAP1 occupation at the IAPEz-int variants, we

observed that clade γ elements displayed the weakest Gm14419 binding and the lowest KAP1

signal compared to clades α and β (Figure 1A,B). Additionally, we observed that IAPLTR2s can

flank either clade α or β IAPEz-ints, (Figure 1 – figure supplement 5), while IAPLTR1s can flank

any IAPEz-int, including IAPEz-int clade γ elements that have decreased KAP1 binding (Figure

1C). These results suggest that both LTR and internal sequence variants can result in altered

KZFP recruitment and KAP1-mediated silencing.

Finally, we were interested in profiling whether the elements that have decreased KZFP

mediated silencing were significantly more likely to be variably methylated. We observed that

VM-IAPLTR1s were significantly enriched at IAPs with IAPLTR1 clade 3 elements flanking

IAPEz-int clade γ elements (Figure 1C), and VM-IAPLTR2 elements were significantly enriched

at solo IAPLTR2 clade 1 elements (p-value < 1e-16 as determined using a Fisher exact test)

(Figure 1 - figure supplement 5). Consistent with previous work (Bertozzi et al., 2020; Faulk et

al., 2013; Kazachenka et al., 2018), these results suggest that sequence is a contributing factor

in the establishment of VM-IAPs. Overall, these results demonstrate that the IAPs that are most

prone to be variably methylated are those that are escaping KZFP mediated silencing..

VM-IAP are proximal to constitutively expressed transcripts and enhancer elements

While sequence has been shown to be a factor in the establishment of variable methylation

(Bertozzi et al., 2020; Faulk et al., 2013; Kazachenka et al., 2018), not all IAPs with sequences

identical to VM-IAPs are variably methylated (Figure 1A), as has been reported previously

(Kazachenka et al., 2018). Previous studies have identified that CTCF enrichment, a protein

involved in regulating chromatin architecture, has been bound at these VM-IAPs when

hypomethylated(Elmer et al., 2020; Kazachenka et al., 2018). However, as CTCF binding motifs

appear to be present at both VM-IAPs and silenced IAPs (Elmer et al., 2020) we examined

whether genomic context had an impact on the establishment of variable methylation at the VM-

IAPs. We used proximity of the IAP to constitutively expressed genes (Li et al., 2017) and

ENCODE annotated enhancer elements from any cell type (ENCODE Project Consortium et al.,

2020) as our proxy for a euchromatic environment (see Methods for details). We found that 83%

of the VM-IAPLTR1s were within 50kb of a constitutively expressed gene, while the remaining

17% of VM-IAPLTR1s were less than 1kb from an annotated enhancer element. In contrast,

only 12% of the non-VM IAPs were proximal to a constitutively expressed gene and 7% were

proximal to an enhancer (Figure 1D). Clade 3 VM-IAPLTR1s were more likely to be proximal to

a constitutively expressed gene than other clade 3 IAPLTR1s regardless of the FPKM or

distance profiled (Figure 1 – figure supplement 6). This indicates that while sequence allows

these elements to have the potential to become variably methylated, VM-IAPs are present in a

permissive chromatin environment to escape being silenced, as heterochromatin spreading

from neighboring loci may drive silencing of these elements.

Conservation of IAPs between mouse strains

IAP elements are a potential source of novelty between mouse strains and VM-IAPs in

particular have been shown to be highly polymorphic (Gagnier et al., 2019; Kazachenka et al.,

2018; Nellåker et al., 2012; Rebollo et al., 2020). In order to investigate the relationship between

IAP sequence, potential for variable methylation and conservation across strains, we leveraged

structural variant differences between mouse strains identified by the Sanger mouse genome

project (Keane et al., 2011). We found that clade 3-4 IAPLTR1s, which have the lowest KAP1

binding of the IAPLTR1 clades and IAPEz-int clade γ elements are more polymorphic across

mouse strains than the other clades (Figure 1E). Additionally, we found that IAPLTR2 clade 1

elements, which are more prone to be variably methylated, are more likely to be polymorphic

across mouse strains (Figure 1 – figure supplement 7). Thus, IAPs that have diminished KZFP

recruitment are more likely to be polymorphic between mouse strains (Figure 1- figure

supplement 9).

IAPs that have loss of KZFP binding are bound by the ZF-CxxC containing proteins TET1 and

CFP1.

It has previously been reported that IAPLTR1s and IAPLTR2s have a significantly higher CpG

density than the genomic average (Elmer et al., 2020; Kazachenka et al., 2018). Given the

prevalence of variable methylation at these elements, we wanted to assess the relationship

between CpG content and variable methylation directly. For each TE subfamily, we profiled the

CpG density of all LTRs (see Methods for details) and the percentage of elements that were

variably methylated using the previously identified list of VM-TEs in C57BL/6J mice

(Kazachenka et al., 2018). We observed that TE subfamilies with a higher average CpG density

had a larger percentage of elements that are variably methylated, with IAPLTRs being the most

CpG dense and having the largest percentage of variably methylated elements (Figure 2A). The

CpG content of VM-IAPLTR1s and VM-IAPLTR2s is similar to non-VM-IAPLTR1s and VM-

IAPLTR2s, indicating that high CpG content alone is not sufficient to induce variable methylation

(Figure 2B).

As CpG dense loci are capable of being recognized and bound by ZF-CxxC proteins

(Blackledge et al., 2010; Thomson et al., 2010), we examined the ability of ZF-CxxC proteins to

bind to IAPLTR1s. Utilizing a publicly available ChIP-seq dataset for the ZF-CxxC-domain

containing protein TET1 in mouse embryonic stem cells (mESCs) (Gu et al., 2018) we identified

increased TET1 binding at clade3 IAPLTR1s elements, which contains the most variably

methylated elements (Figure 2C and Figure 2 – figure supplement 1). We furthermore profiled

CpG methylation of VM-IAPLTRs in *Tet1* knockout (KO) and *Tet1/Dnmt3a* double knockout

(DKO) mESCs (Gu et al., 2018) and found that VM-IAPs displayed a significant increase in CpG

methylation in the *Tet1* KO cells (Figure 2D). This increase in methylation is reversed when both

Dnmt3a and *Tet1* are knocked out in mESCs (Figure 2D). We also profiled the recruitment of

another ZF-CxxC protein, CFP1, using existing CFP1 ChIP-seq data from C57Bl/6 mice and

observed CFP1 binding appears to be specific to VM-IAPLTRs (Figure 2 -figure supplement 2).

As CFP1 is a subunit of the SET1 complex responsible for depositing H3K4me3, we examined

publicly available H3K4me3 ChIP-seq data from B cells (Adams et al., 2012) to determine if

H3K4me3 is enriched at VM-IAPs. Consistent with previous observations (Bertozzi et al., 2020),

we observed that VM-IAPs were more likely to ben enriched for H3K4me3 compared to non-

variable IAPs (Figure 2 - figure supplement 3). *De novo* motif discovery applied to VM-IAP

sequences furthermore revealed known recognition sequences for ZF-CxxC proteins such as

CFP1 (Figure 2 - figure supplement 4). ZF-CxxC protein binding has been shown to protect the

underlying DNA from methylation (Blackledge et al., 2010; Thomson et al., 2010). From these

results we propose that recruitment of ZF-CxxC proteins, which have been shown to be involved

in maintaining a euchromatic state at CpG islands, may be involved in the establishment of

variable methylation at VM-IAPs in mice.

To determine if evolutionarily recent CpG-dense TEs are also more prone to be variably

methylated in humans, we used a previously identified list of loci that display interindividual

epigenetic variation in humans (Gunasekara et al., 2019). We determined the observed

frequency at which variably methylated loci were found to be present on a TEs and compared

this to an expected distribution if the variable methylation was only due to chance. Expected

distribution was determined by selecting random loci in the genome the same size of the

variably methylated loci. We observed that evolutionarily recent TEs were overrepresented in

the variably methylated dataset (Figure 3A). TE subfamily ages were obtained from DFAM

(Hubley et al., 2016). We also found the TE subfamilies with a higher CpG density are more

likely to be variably methylated (Figure 3B and Figure 3 – figure supplement 1), with the highly

CpG dense transposable element subfamily LTR12C having both one of the highest CpG

densities and one of highest percentage of elements that are variably methylated. LTR12Cs are

recently evolved TEs that have previously been shown to have latent regulatory potential (Ward

et al., 2013) and can act as cryptic promoter elements in cancer cells (Babaian and Mager,

2016). These results demonstrate a correlation between CpG density and variable methylation

in both mice and humans.

CpG dense TEs are hypomethylated and recruit ZF-CxxC proteins in the absence of KZFP-

mediated silencing

To further investigate the relationship between KZFP/KAP1 silencing, DNA methylation, and ZF-

CxxC binding to CpG dense TEs, we profiled DNA methylation and ZF-CxxC binding in Tc1

mouse liver. This "transchromosomic" mouse carries the majority of human chromosome 21

(Hsa21), which possesses human specific transposable elements without the corresponding

KZFPs for silencing (O'Doherty et al., 2005) and has previously been used to identify the latent

regulatory potential of primate specific TEs (Jacobs et al., 2014; Long et al., 2016; Ward et al.,

2013). We performed MinION nanopore sequencing to globally profile DNA methylation of

human chromosome 21 in Tc1 mouse liver and compared this with existing DNA methylation

data from human liver cells (Li et al., 2016)(Figure 4A,B). Consistent with previous reports

(Jacobs et al., 2014; Long et al., 2016; Ward et al., 2013), we found that TE-derived CGIs were

hypomethylated in the absence of KZFP mediated silencing (Figure 4B). In contrast, TE-derived

CGIs that can be targeted by KZFPs shared between mice and humans were largely methylated

(Figure 4 - figure supplement 1). These silenced TE-derived CGIs are largely Alu elements that

are derived from the same ancestral sequence as mouse B2 SINE elements and can be

silenced by the KZFP ZNF182, which is present in both mice and humans (Imbeault et al.,

2017). Additionally, other primate specific TEs that are not CpG rich are methylated in both the

Tc1 and human livers (Figured 4B), highlighting the importance of CpG density in escaping CpG

methylation at TEs.

To profile ZF-CxxC occupancy at these hypomethylated TEs, we performed CUT&RUN for

CFP1, a ZF-CxxC protein expressed in somatic tissue, in Tc1 mouse livers and compared this

with existing CFP1 ChIP-seq data from human erythroblasts (van de Lagemaat et al., 2018). We

observed that CpG dense TEs that are not targeted by KZFPs displayed CFP1 recruitment,

while the methylated CpG poor TEs have no CFP1 signal (Figure 4C). This CFP1 binding was

also unique to the Tc1 mice as there was no CFP1 binding at these loci in humans (Figure 4C),

while non-TE CGIs in both the mouse and human genomes have CFP1 binding (Figure 4 -

figure supplement 2). Together this work demonstrates that in the absence of targeted KZFP

mediated silencing, TEs with high CpG content can be hypomethylated and bound by ZF-CxxC

proteins.

Trim28 haploinsufficiency activates evolutionarily recent TEs

As TRIM28/KAP1 is a key protein involved in the establishment of heterochromatin at TEs, we

profiled *Trim28* haploinsufficient mice to determine if decreased abundance of TRIM28 allows

for CpG dense TEs to escape silencing. *Trim28* haploinsufficiency has previously been shown

to impact the Agouti viable yellow mouse coat color (Daxinger et al., 2016) and drive a bimodal

obesity phenotype in FVB/NJ mice (Dalgaard et al., 2016). We performed both H3K4me3 ChIP-

seq and CFP1 CUT&RUN in the livers of Trim28 haploinsufficient (*Trim28*$^{+/D9}$) mice and their

wild type littermates (Methods; Figure 5A). We identified 69 loci that had novel H3K4me3 signal

in the *Trim28*$^{+/D9}$ mice compared to control (Methods; Figure 6 - figure supplement 1). The

majority of these loci with novel H3K4me3 recruitment were evolutionarily recent TEs (Figure 6B

and Figure 6 - supplement 2). These loci also had an increase in CFP1 signal in *Trim28*$^{+/D9}$ mice

relative to control mice (Figure 6C and Figure 6 - figure supplement 1). Additionally, we profiled

global changes in expression of TE subfamilies in the *Trim28*$^{+/D9}$ mice compared to wild type

mice using our previously generated RNA-seq datasets (Dalgaard et al., 2016). We identified

that TEs which have a significant increase in gene expression in the *Trim28*$^{+/D9}$ mice compared

to wild type mice have significantly higher CpG density than expected when compared to non-

responsive TE subfamilies (Figure 6D). Together this work demonstrates that loss of KAP1

mediated silencing allows for some evolutionary recent TEs to be active and recruit ZF-CxxC

proteins.

Discussion

This work provides the foundation for a deeper understanding of how mammalian genomes are

shaped during evolution. It has been proposed that genomes are locked in an "evolutionary

arms race" between endogenous retrovirus and KZFPs to maintain proper silencing of the

genome. It The variable methylation observed at IAPs could be the result of the IAPs winning

this "battle". We observed that VM-IAPs have unique sequence variants that allow them to have

diminished KZFP recruitment. This provides an explanation to previous works that identified

sequence biases for variably methylated IAPs in mice (Bertozzi et al., 2020; Faulk et al., 2013;

Kazachenka et al., 2018). We also found that IAPs with the same sequence variants as the VM-

IAPs are more likely to be polymorphic between individual mouse strains. Interestingly, we find

that the IAPLTRs flanking the "master" IAP that has been proposed to be responsible for an

expansion of IAPs in C3H/HeJ mice (Rebollo et al., 2020) has the same sequence as the clade

3 IAPLTR1s (Figure 6 - figure supplement 1). Together this implies that these VM-IAPs may be

more active in certain mouse strains and may be propagating due to a lack of repression.

Our work also highlights the potential importance of CpG density in the silencing of the

endogenous retroviruses. We observed that VM-IAPs have recruitment of ZF-CxxC proteins. In

the absence of targeted silencing, CpG-dense TEs have the potential to become

hypomethylated and recruit ZF-CxxC proteins, while CpG poor TEs remain methylated. As TEs

can lose CpG density over time, the observation that the CpG rich TEs have the potential for

reactivation supports the idea that deamination of methylated loci could be one means for

driving permanent silencing of TEs over time (Long et al., 2016).

Based on our results, we propose a model that VM-IAP loci are established as a result of partial

KZFP mediated silencing at CpG dense TEs, which can then be protected from silencing

through recruitment of ZF-CxxC containing proteins such as TET1 (Figure 6). This potentially

results in a stochastically silenced locus, which is then inherited by all tissues during

development. Together our work has significant implications in furthering our understanding of

how TEs can alter both the genomes and epigenomes of mammals.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Strain, strain background (*Mus musculus*)	*Trim29$^{+/D9}$*	(Blewitt et al., 2005)	(RRID:MGI:3 821610)	Haploinsufficient for *Trim28*
Strain, strain background (*Mus musculus*)	B6129S-Tc(HSA21) 1TybEmcf/J	The Jackson Laboratory	Stock No: **010801 (**JAX) (RRID:IMSR_ JAX:010801)	2Mb of a freely segregating human fragment of Chr21
Antibody	Rabbit Polyclonal anti-CFP1 antibody	Millipore	ABE211 (RRID:AB_10 806210)	CUT&RUN (1:50 dilution)
Antibody	Rabbit polyclonal anti-H3K4me3 Antibody	Abcam	ab8580 (RRID:AB_30 6649)	ChIP-seq (2µg antibody per 25µg chromatin)

Multiple sequence alignment and hierarchical clustering

IAPLTR1_Mm, IAPLTR2_Mm, and IAPEz-int elements were defined by RepeatMasker

(RRID:SCR_012954) (Smit, AFA, Hubley, R & Green, P, 2010) in the mm10 genome. Only

IAPLTRs greater than 300bps in length were profiled. For IAPEz-ints, only IAPEz-int elements

flanked by either IAPLTR1 or IAPLTR2 elements were selected for profiling, and only the first

150bps of the 5' end of the IAPEz-int were aligned and clustered. All elements were processed

using the ETE3 pipeline (Huerta-Cepas et al., 2016) using MAFFT (RRID:SCR_011811) with

default settings (Katoh and Standley, 2013), and bases that were found in less than 10% of the

samples were trimmed from the multiple sequence alignment using trimAL (RRID:SCR_017334)

(Capella-Gutiérrez et al., 2009). The multiple sequence alignments were then hierarchically

clustered using PhyML (RRID:SCR_014629) (Guindon et al., 2010). Sequence clades were

empirically selected.

Alignment of existing ChIP-seq data

Reads were trimmed using Trimgalore version 0.5.0 (RRID:SCR_011847), which utilizes

cutadapt (RRID:SCR_011841) (Martin, 2011), and were aligned to either the mm10 genome for

mouse data, hg38 genome for human data, or a custom assembly that included both the mm10

genome and human chromosome 21 for the Tc1 mice. Reads were aligned using bowtie1

version 1.2.3 (RRID:SCR_005476) retaining only reads that could be mapped to unambiguously

to a single locus using the -m 1 option (Langmead et al., 2009). Aligned reads were sorted using

samtools version 1.10 (RRID:SCR_002105) (Li et al., 2009) and filtered to remove duplicate

reads using the MarkDuplicates function of Picardtools version 2.21.1 (RRID:SCR_006525).

Regions of enrichment were called using the callpeaks function of MACS2 version 2.2.5

(RRID:SCR_01329) (Zhang et al., 2008) with a q-val threshold of 1e-3.

KZFP and KAP1 data in Figure 1 was aligned using different parameters to allow for potential

multimapping as the reads were too short to provide confident unique mapping at repetitive

elements. These reads were aligned to the mm10 genome using bowtie2 version 2.3.5.1

(RRID:SCR_016368) with the --end-to-end --very-sensitive options (Langmead and Salzberg,

2012). Aligned reads were sorted using samtools version 1.10 (RRID:SCR_002105) (Li et al.,

2009) and filtered for duplicate reads using the MarkDuplicates function of Picardtools version

2.21.1 (RRID:SCR_006525). Regions of enrichment were called using MACS2 version 2.2.5

callpeaks (RRID:SCR_01329) (Zhang et al., 2008) with a q-val threshold of 1e-3.

For all datasets, Bedgraph files were generated using bedtools version 2.29.0 genomecov

(RRID:SCR_006646) with the -bg -ibam options (Quinlan and Hall, 2010). BigWigs

(RRID:SCR_007708) were generated using the UCSCtools bedGraphToBigWig (Karolchik et

al., 2004). Heatmaps of ChIP-seq signal across the IAP multiple sequence alignments were

generated using a custom script to profile the read coverage at each base and were visualized

using pheatmap (RRID:SCR_016418). All other heatmaps and aggregate plots of loci that

extend were generated using deeptools (RRID:SCR_016366) (Ramírez et al., 2016).

Genomic context analysis

We used proximity to a constitutively expressed gene or annotated enhancer element as a

proxy for profiling a euchromatic environment. Constitutive expressed genes were identified as

genes that have a fragments per kilobase of transcript per million mapped reads (FPKM) greater

than 2 in more than 90% of the samples previously profiled in (Li et al., 2017), while enhancers

were all ENCODE annotated enhancer elements (RRID:SCR_006793) (ENCODE Project

Consortium et al., 2020). While FPKM of >2 was used, observed trends appeared regardless of

FPKM threshold set (Figure 1 – figure supplement 6).

374 Polymorphism analysis between mouse strains

Coordinates of the structural variants between mouse strains were obtained from the Sanger

mouse genome project (RRID:SCR_006239) (Keane et al., 2011). Deletions between mouse

strains were extracted for each of the profiled mouse strains. Bedtools version 2.29.0 intersect

(RRID:SCR_006646) (Quinlan and Hall, 2010) with the -f 1 option was used to determine if an

IAP was fully deleted in each mouse strain relative to C57BL/6J. Phylogeny of the mouse

strains was obtained from previous work (Nellåker et al., 2012). For branches where multiple

mouse strains are similarly diverged from C57BL6/J, the IAP was considered to be present if the

IAP existed in any of profiled mice at that branch.

384 Profiling of CpG score

CpG score was calculated as previously described (Gardiner-Garden and Frommer, 1987).

Briefly, we calculated CpG score as Obs/Exp CpG = Number of CpG * N / (Number of C *

Number of G). When profiling the CpG density of across the TE using sliding 200 bp tiles with

10 bp steps between windows and selected the most CpG dense portion of the TE to remove

and impact of TE size. Window size was chosen to match the minimum size of a CpG island

(Gardiner-Garden and Frommer, 1987). Differences in CpG density across an element was

most pronounced on L1 elements, which can contain a CpG island in their 5' UTR but have low

CpG density across the rest of the element. TEs with low copy numbers, such as IAPLTR4,

were removed from this analysis.

395 MinION sequencing and alignment

Livers from B6129S-Tc(HSA21)1TybEmcf/J mice (RRID:IMSR_JAX:010801) were purchased

from JAX. Purified DNA from Tc1 mouse liver was prepared using the SQK-LSK109 ligation

sequencing kit and run on R9 flow cells purchased from Nanopore. 1ug of DNA was end

repaired with NEBNext FFPE DNA repair and Ultra II End-prep purchased from New England

Biolabs. The repaired DNA was cleaned with KAPA Pure Beads. Adapters were ligated using

NEBNext quick T4 ligase purchased from New England Biolabs. Prepared libraries were mixed

with Sequencing Buffer from the SQK-LSK109 kit and approximately 200ng was loaded onto the

MinION flowcell. Reads were aligned using bwa mem with the options -x ont2d -t 100

(RRID:SCR_010910) (Li, 2013) to a custom assembly that included all of the mm10 genome

and human chromosome 21, and then sorted using samtools version 1.10 (RRID:SCR_002105)

(Li et al., 2009). Reads were processed using minimap2 (RRID:SCR_018550) with the options -

a -x map-ont (Li, 2018). Methylation state of CpGs was called using nanopolish

(RRID:SCR_016157) with the options call-methylation -t 100 (Loman et al., 2015). Only loci with

greater than 5x coverage were considered in the analysis. CpG islands were obtained from the

UCSC table browser (Gardiner-Garden and Frommer, 1987; Karolchik et al., 2004). Methylation

percentage was averaged across CpG islands.

CFP1 CUT&RUN sequencing and alignment

CUT&RUN was performed as previously described (Skene and Henikoff, 2017) on livers from

B6129S-Tc(HSA21)1TybEmcf/J mice (RRID:IMSR_JAX:010801) purchased from JAX and

Trim28$^{+/D9}$ livers using the CFP1 antibody (RRID:AB_10806210) (ABE211, Millipore). Briefly,

unfixed permeabilized cells are incubated with the CFP1 antibody fused to A-Micrococcal

Nuclease. Fragmented DNA was isolated and sequenced on an Illumina HiSeq 2500 System.

We assessed standard QC measures on the FASTQ file using FASTQC (RRID:SCR_014583)

and adapters were trimmed using Trimgalore version 0.5.0 (RRID:SCR_011847). Trimmed

reads were aligned to a custom assembly that included both the mm10 genome and human

chromosome 21, or only the mm10 genome using bowtie1 version 1.2.3 (RRID:SCR_005476)

using the -m 1 option to only retain reads that could be mapped to unambiguously to a single

locus (Langmead et al., 2009). Aligned reads were sorted using samtools version 1.10

(RRID:SCR_002105) (Li et al., 2009) and filtered for duplicate reads using the MarkDuplicates

function of Picardtools version 2.21.1 (RRID:SCR_006525). Regions of CFP1 enrichment were

called using the call peaks function of MACS2 version 2.2.5 (RRID:SCR_01329) (Zhang et al.,

2008) with a *q*-val threshold of 1e-3. Only peaks found in two animals were retained to remove

biological noise. Additionally, due to small fragment size, loci that could not be mapped

unambiguously by 90 bp fragments were removed from consideration using the deadzones tool

from RSEG (RRID:SCR_007695) (Song and Smith, 2011).

Animals

The generation of *Trim28*[+/D9] mice (RRID:MGI:3821610) has been described elsewhere (Blewitt

et al., 2005). Animals were kept on a 12 hr light/dark cycle with free access to food and water

and housed in accordance with international guidelines. Livers were isolated at 2 weeks of age

and immediately snap-frozen for further processing.

Tc1 mouse livers (RRID:IMSR_JAX:010801), freshly harvested and snap frozen, were

purchased from JAX Laboratories (Stock No: 010801).

ChIP-seq analysis of H3K4me3 ChIP-seq

Chromatin immunoprecipitation was performed with an H3K4me3 antibody (RRID:AB_306649)

(ab8580, Abcam) as previously described (Leung et al., 2013). Isolated DNA was sequenced on

an Illumina HiSeq 2500 System. We assessed standard QC measures on the FASTQ file using

FASTQC (RRID:SCR_014583) and adapters were trimmed using Trimgalore version 0.5.0

(RRID:SCR_011847). Reads were aligned to the mm10 genome using bowtie1 version 1.2.3

(RRID:SCR_005476) using the -m 1 option to only retain reads that could be mapped to

unambiguously to a single locus (Langmead et al., 2009). Aligned reads were sorted using

samtools version 1.10 (RRID:SCR_002105) (Li et al., 2009) and filtered for duplicate reads

using the MarkDuplicates function of Picardtools version 2.21.1 (RRID:SCR_006525). Regions

of enrichment were called using the call peaks function of MACS2 version 2.2.5

(RRID:SCR_01329) (Zhang et al., 2008) with a q-val threshold of 1e-3. Loci that displayed

enrichment of H3K4me3 in at least two animals were retained to remove biological noise. The

loci with increased H3K4me3 as shown in Fig 6*B-C* were identified by finding peaks present in

Trim28$^{+/D9}$ mice that were absent from WT mice and also had 3x more H3K4me3 signal in

Trim28$^{+/D9}$ mice compared to WT. Heatmaps and aggregate plots were generated using

deeptools (RRID:SCR_016366) (Ramírez et al., 2016).

Differential expression analysis of repetitive element subfamilies.

RNA-seq reads were downloaded from ENA:PRJEB11740, and adapters were trimmed using

Trimgalore version 0.5.0 (RRID:SCR_011847). Reads were then aligned and processed using

the RepEnrich2 pipeline RRID:SCR_021733 as previously described (Criscione et al., 2014).

This package sums the number of reads mapping to each repetitive element subfamily.

Differentially expressed subfamilies between *Trim28$^{+/D9}$* and wild type mice were identified using

DEseq2 (RRID:SCR_015687) (Love et al., 2014) with a p-val threshold of 0.05.

Acknowledgements

We thank Vanessa Wegert for valuable technical support. We would also like the thank the and

members of the Schones lab for helpful comments and suggestions. This work was supported

by the National Institutes of Health, grants R01DK112041, R01CA220693 (D.E.S.). The

research reported in this publication included work performed in the Integrative Genomics and

Analytical Cytometry Cores supported by the National Cancer Institute of the National Institutes

of Health under award number P30CA033572.

Availability of data and materials

All datasets generated in this study have been submitted to the NCBI Gene Expression

Omnibus (GEO; https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE176176) under

accession no. GSE176176. Reviewer access token gtmroekqrnshheb

Public data used:

Mouse KZFP and KAP1 ChIP-seq from C57Bl/6J mice GEO: GSE115291. Mouse TET1 ChIP-

seq and mouse WGBS for WT, Tet1KO and Dnmt3a/Tet1 DKO from J1 mouse ESCs

GEO:GSE100957. Mouse H3K4me3 ChIP-seq from C57BL/6J B-cells and T-cells GEO:

GSM2480410. CFP1 ChIP-seq from C57BL/6J Brain tissue GEO:GSE18578. Human

erythrocyte CFP1 ChIP-seq GEO:GSM3132538. WGBS from human liver GEO:GSM1716957.

Trim28 haploinsufficient and wild type mouse RNA-seq from an FVB background was obtained

from ENA:PRJEB11740. List of VM-loci in humans were obtained from published work

(Gunasekara et al., 2019). List of VM-loci in mice were obtained from published work (Elmer et

al., 2020).

Figure Legends

Figure 1: **Sequence and chromatin context influence the establishment of a VM-IAPLTRs.**

A) All IAPLTR1 elements larger than 300 bps and the first 150bps of IAPEz-ints flanked by

IAPLTR1 or IAPLTRs were clustered by sequence using PhyML with default settings. Major

sequence variants for IAPLTR1s were separated into separate clades: clade1 (blue), clade 2

(Green), clade 3 (red) and clade 4 (orange). Major sequence variants for IAPEz-ints were

separated into separate clades: clade α (dark blue), clade β (light blue), and clade γ (yellow).

KZFP and KAP1 ChIP-seq signal was mapped across the consensus IAPLTR1 sequence as

determined by MAFFT multiple sequence alignment. Gaps in the multiple sequence alignment

for the IAPLTR1 sequence are displayed as grey. Heatmaps for KZFP and KAP1 ChIP-seq

signals centered on the 5' end of the IAPEz-int elements are show as well B) Average KAP1

ChIP-seq signal across all IAPLTR1 and IAPEz-int clades. Each data point refers to the average

KAP1 ChIP-seq signal at an individual IAPLTR1 element. Mean and standard deviation for each

clade is shown as well. C) Distribution of the IAPLTR1 elements clades (outer circle) and

IAPEz-int clades (inner circle) for all IAP elements and VM-IAP elements. D) Percent of VM-

IAPLTR1s clade 3 elements and non VM-IAPLTR1 clade 3 elements that are within 50kb of a

constitutively expressed gene or 1kb enhancer element, as a proxy for constitutive euchromatin

environment. E) Conservation of IAPLTR1 and IAPEz-int variants across mouse strains.

Presence or absence of a IAP was determined using structural variants identified from the

Sanger mouse genome project. Mouse KZFP and KAP1 ChIP-seq date are from GEO:

GSE115291. VM-loci coordinates were obtained from (Elmer et al., 2020).

Figure 2: **Divergent VM-IAPs elements have high CpG density and recruitment of ZF-CxxC**

proteins. A) Percent of ERV LTRs elements that are variably methylated for a given TE

subfamily and average maximum CpG score of the TE subfamily. The size of each dot is

determined by the number of VM-loci for each subfamily, with the largest dots indicating the

greatest number of variably methylated elements. Average CpG score was determined by

identifying the most CpG dense 200bp window of each LTR and calculating the average CpG

score for the whole subfamily. B) Average CpG density of silenced and variably methylated

IAPLTRs, as well as the average CpG score of a randomly selected background the same size

as the IAPLTRs. Each dot refers to an individual IAPLTR element in the mm10 genome. Mean

and standard deviation for each group is shown. C) Aggregate plots of TET1 ChIP-seq signal

across all identified IAPLTR1 clades. D) Average methylation percentage of non-variable and

VM-IAPLTR elements in wild type, Tet1 knockout (KO), and Dnmt3a/Tet1 double knockout cells

(DKO). Each dot refers to the average methylation of an individual IAPLTR element. Only CpGs

with >5x coverage were retained to calculate methylation. Significance determined using a

Wilcoxon rank sum test. *** indicates a p.value <0.0001. Bars for mean and standard deviation

is shown as well for figure 2B and 2D. TET1 ChIP-seq from GEO:GSE100957. Mouse *Tet1* and

Tet1/Dnmt3a DKO WGBS from GEO:GSE134396

Figure 3: **Variably methylated loci in humans are enriched for evolutionarily recent CpG-**

dense TEs. A) Observed over expected distribution of VM-TE elements in humans stratified by

the evolutionary age of the TE. Expected distribution was determined using a random sampling

of the hg38 genome the same size as the VM-loci. Evolutionary age for each TE subfamily was

obtained from DFAM. B) Scatterplot showing the percent of LTR elements which display VM for

a given TE subfamily and average CpG score of the TE subfamily. The size of each dot is

determined by the number of VM-loci for each subfamily. Average CpG score was determined

by identifying the most CpG dense 200bp window of each ERV and calculating the average

CpG score for the whole subfamily. The coordinates of VM-loci for humans were obtained from

(Gunasekara et al., 2019).

Figure 4: **CpG dense TEs are hypomethylated and recruit ZF-CxxC proteins in the**

absence of KZFP mediated silencing. A) UCSC genome browser screenshot of an LTR12C

element which is hypomethylated in Tc1 mice and shows novel CFP1 recruitment in Tc1 mice.

B) CpG methylation of TE CpG islands (CGIs) and other TEs for human chromosome 21 in both

Tc1 mouse and human genomes. Each data point refers to an individual CpG island that had

greater than 5x coverage. Mean and standard deviation for DNA methylation at is shown as

well. C) CFP1 signal in Tc1 mice and humans at the human TE CGIs and other TEs (from

Figure 4B) on human chromosome 21. Human CFP1 ChIP-seq from GEO:GSM3132538.

WGBS from human liver GEO:GSM1716957

Figure 5: **Trim28 haploinsufficiency leads to activation of evolutionarily recent and CpG**

dense TEs. A) Genome screenshot of an IAPLTR2 element with novel H3K4me3 enrichment in

Trim28 haploinsufficient mice. B) Breakdown of loci with novel H3K4me3 in Trim28

haploinsufficient mice. Age of each TE was determined by DFAM. C) Aggregate plots of

H3K4me3 and CFP11 signal across all loci that have novel H3K4me3 signal in Trim28

haploinsufficient mice D) CpG Score of TE subfamilies with a global increase expression in

Trim28 haploinsufficient mice and a random selection of non-responsive TEs subfamilies.

Expression levels for each TE subfamily was determined using RepEnrich, and Deseq2 was

used to determine TE subfamilies with a significant increase in expression in the Trim28

haploinsufficient mice. Bar is placed at mean and error bars cover one standard deviation. Each

data point refers to an individual TE subfamily. *P*-values for CpG density difference was

calculated using a Wilcoxon rank sum test. *Trim28* haploinsufficient and wild type mouse RNA-

seq was obtained from ENA:PRJEB11740.

Figure 6: **Model for variable methylated transposable elements.** Loci with high CpG density

and loss of KZFP binding have the potential to recruit ZF-CxxC proteins to protect these TEs

from being silenced. However, elements with high CpG density but strong KZFP recruitment will

remain methylated.

Supplemental Legends

Figure 1 – figure supplement 1: Multiple sequence alignment of IAPLTR1s, IAPLTR2, and the

first 150bps of IAPEz-int flanked by IAPLTR1/2 aligned by MAFFT with default settings(Katoh

and Standley, 2013). Bases present in less than 10% of IAPLTR elements were trimmed from

the alignment. Sequences were clustered using PhyML with default settings. Each base is

visualized by a different color: thymine is orange, adenine is dark grey, cytosine is yellow, and

guanine is light grey. Bases with no alignment are white.

Figure 1 – figure supplement 2: Consensus sequence of the IAPLTR1 clades identified in Figure

1 – Supplemental 1. Bases present in less than 10% of IAPLTR elements were trimmed from

the alignment. CpG dinucleotides are highlighted in light grey.

Figure 1 – figure supplement 3: Gm21082 and KAP1 ChIP-seq signal mapped across all

IAPLTR1 elements. Consensus sequence of the IAPLTR1 clades at the region of increased

ZFP989/Gm21082 binding. Arrows indicate locations of indels between clades. Previously

identified Gm21082 binding motif from (Wolf et al., 2020).

Figure 1 – figure supplement 4: ZFP429, ZFP989, and KAP1 ChIP-seq signal mapped across

all IAPLTR2 elements.

Figure 1 – figure supplement 5: Association of IAPLTR2 and IAPEz-int clades for all IAPLTR2s

and VM-IAPLTR2.

Figure 1 – figure supplement 6: The percent of VM-IAPLTR1s clade 3 elements and non VM-

IAPLTR1 clade 3 elements proximal to a constitutively expressed genes. Each point refers to

the percentage of IAPLTR1 proximal to a constitutively expressed gene at a given value.

Distances were sampled every 2500bps.

Figure 1 – figure supplement 7: Conservation of IAPLTR1 and IAPEz-int variants across mouse

strains. Presence or absence of a IAP was determined using structural variants identified from

the Sanger mouse genome project. If the entire IAP was found to be missing from the other

mouse assembly, the IAP was determined to be not conserved.

Figure 1 – figure supplement 8: Model of IAPLTR and IAPEz-int interactions the height of each

IAPEz-int refers to the percentage of IAP elements.

Figure 2 – figure supplement 1: TET1 ChIP-seq signal mapped across all IAPLTR1 elements.

Figure 2 – figure supplement 2: A) Aggregate plots of Cfp1 signal at variably methylated and

other IAPLTRs. Heatmaps of Cfp1 signal at B) VM-IAPLTRs and C) all other IAPLTR1 and

IAPLTR2 elements (2819 elements). Reads were uniquely aligned using bowtie1. Data from

GEO:GSE18578

Figure 2 – figure supplement 3: A) Observed over expected H3K4me3 signal at variably

methylated and non-variably methylated IAPLTRs. Expected H3K4me3 signal determined using

average H3K4me3 signal at 50 randomly selected loci repeated 1000 times. B) H3K4me4 ChIP-

seq signal at all VM-IAPLTRs, silenced IAPLTRs and a random sampling of 50 IAPLTRs that

were not variably methylated. C) Aggregate plots of H3K4me3 ChIP-seq signal at the three

groups identified in Figure 2 – supplemental 1C.

Figure 2 – figure supplement 4: MEME identified motifs present in VM-IAPLTRs that contain

CFP1 binding sites. *P*-values obtained from MEME.

Figure 3 – figure supplement 1: Scatter plot showing the percent of SINEs and LINEs that

display variable methylation for a given TE subfamily and average CpG score of the TE

subfamily in humans.

Figure 4 – figure supplement 1: CpG methylation of all TEs on human chromosome 21. TEs

were separated by CpG density and their capability of being silenced by a KZFP element

shared between mouse and humans. Largely, these silenced repeat CGIs were found at Alu

elements.

Figure 4 – figure supplement 2: CFP1 signal at all non-repeat derived CpG Islands on human

chromosome 21 in Tc1 mice and humans.

Figure 5 – figure supplement 1: Heatmap and aggregate plots of H3K4me3 and CFP1 signal at

loci with novel H3K4me3 signal in Trim28 D9/+ mice.

Figure 5 – figure supplement 2: Evolutionary age of all TEs in the mm10 genome. Evolutionary

age was obtained from Dfam (Hubley et al., 2016).

Figure 6 – figure supplement 1: Comparison of the IAPLTR1 associated with the "master" IAP

element identified in the C3H mice compared to the consensus sequence for each clade. Only a

portion of the IAPLTR1 containing identifiable sequence variants for each clade is shown to

improve visibility.

References

Adams D, Altucci L, Antonarakis SE, Ballesteros J, Beck S, Bird A, Bock C, Boehm B, Campo
E, Caricasole A, Dahl F, Dermitzakis ET, Enver T, Esteller M, Estivill X, Ferguson-Smith A,
Fitzgibbon J, Flicek P, Giehl C, Graf T, Grosveld F, Guigo R, Gut I, Helin K, Jarvius J, Küppers
R, Lehrach H, Lengauer T, Lernmark Å, Leslie D, Loeffler M, Macintyre E, Mai A, Martens JHA,
Minucci S, Ouwehand WH, Pelicci PG, Pendeville H, Porse B, Rakyan V, Reik W, Schrappe M,
Schübeler D, Seifert M, Siebert R, Simmons D, Soranzo N, Spicuglia S, Stratton M,
Stunnenberg HG, Tanay A, Torrents D, Valencia A, Vellenga E, Vingron M, Walter J, Willcocks
S. 2012. BLUEPRINT to decode the epigenetic signature written in blood. *Nat Biotechnol*
30:224–226.

Babaian A, Mager DL. 2016. Endogenous retroviral promoter exaptation in human cancer. *Mob*
DNA **7**:24.

Bakshi A, Herke SW, Batzer MA, Kim J. 2016. DNA methylation variation of human-specific Alu
repeats. *Epigenetics* **11**:163–173.

Bernal AJ, Murphy SK, Jirtle RL. 2011. Chapter 15 - Mouse Models of Epigenetic Inheritance In:
Tollefsbol T, editor. Handbook of Epigenetics. San Diego: Academic Press. pp. 233–249.

Bertozzi TM, Becker JL, Blake GET, Bansal A, Nguyen DK, Fernandez-Twinn DS, Ozanne SE,
Bartolomei MS, Simmons RA, Watson ED, Ferguson-Smith AC. 2021. Variably methylated
retrotransposons are refractory to a range of environmental perturbations. *Nat Genet* **53**:1233–
1242.

Bertozzi TM, Elmer JL, Macfarlan TS, Ferguson-Smith AC. 2020. KRAB zinc finger protein
diversification drives mammalian interindividual methylation variability. *Proc Natl Acad Sci U S A*
117:31290–31300.

Blackledge NP, Zhou JC, Tolstorukov MY, Farcas AM, Park PJ, Klose RJ. 2010. CpG islands
recruit a histone H3 lysine 36 demethylase. *Mol Cell* **38**:179–190.

Blewitt ME, Vickaryous NK, Hemley SJ, Ashe A, Bruxner TJ, Preis JI, Arkell R, Whitelaw E.
2005. An N-ethyl-N-nitrosourea screen for genes involved in variegation in the mouse. *Proc Natl*
Acad Sci U S A **102**:7629–7634.

Bulut-Karslioglu A, De La Rosa-Velazquez IA, Ramirez F, Barenboim M, Onishi-Seebacher M,
Arand J, Galan C, Winter GE, Engist B, Gerle B, O'Sullivan RJ, Martens JH, Walter J, Manke T,
Lachner M, Jenuwein T. 2014. Suv39h-dependent H3K9me3 marks intact retrotransposons and
silences LINE elements in mouse embryonic stem cells. *Mol Cell* **55**:277–290.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. 2009. trimAl: a tool for automated
alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* **25**:1972–1973.

Clouaire T, Webb S, Skene P, Illingworth R, Kerr A, Andrews R, Lee J-H, Skalnik D, Bird A.
2012. Cfp1 integrates both CpG content and gene activity for accurate H3K4me3 deposition in
embryonic stem cells. *Genes Dev* **26**:1714–1728.

Coluccio A, Ecco G, Duc J, Offner S, Turelli P, Trono D. 2018. Individual retrotransposon
integrants are differentially controlled by KZFP/KAP1-dependent histone methylation, DNA
methylation and TET-mediated hydroxymethylation in naive embryonic stem cells. *Epigenetics*
Chromatin **11**:7.

Criscione SW, Zhang Y, Thompson W, Sedivy JM, Neretti N. 2014. Transcriptional landscape of
repetitive elements in normal and cancer human cells. *BMC Genomics* **15**:583.

Dalgaard K, Landgraf K, Heyne S, Lempradl A, Longinotto J, Gossens K, Ruf M, Orthofer M,
Strogantsev R, Selvaraj M, Lu TT-H, Casas E, Teperino R, Surani MA, Zvetkova I, Rimmington
D, Tung YCL, Lam B, Larder R, Yeo GSH, O'Rahilly S, Vavouri T, Whitelaw E, Penninger JM,
Jenuwein T, Cheung C-L, Ferguson-Smith AC, Coll AP, Körner A, Pospisilik JA. 2016. Trim28
Haploinsufficiency Triggers Bi-stable Epigenetic Obesity. *Cell* **164**:353–364.

Daxinger L, Oey H, Isbel L, Whitelaw NC, Youngson NA, Spurling A, Vonk KKD, Whitelaw E.
2016. Hypomethylation of ERVs in the sperm of mice haploinsufficient for the histone
methyltransferase Setdb1 correlates with a paternal effect on phenotype. *Sci Rep* **6**:25004.

Dolinoy DC, Das R, Weidman JR, Jirtle RL. 2007a. Metastable Epialleles, Imprinting, and the
Fetal Origins of Adult Diseases. *Pediatr Res* **61**:30R.

Dolinoy DC, Huang D, Jirtle RL. 2007b. Maternal nutrient supplementation counteracts
bisphenol A-induced DNA hypomethylation in early development. *Proc Natl Acad Sci U S A*
104:13056–13061.

Dolinoy DC, Weinhouse C, Jones TR, Rozek LS, Jirtle RL. 2010. Variable histone modifications
at the A(vy) metastable epiallele. *Epigenetics* **5**:637–644.

Duncan BK, Miller JH. 1980. Mutagenic deamination of cytosine residues in DNA. *Nature*
287:560–561.

Ecco G, Cassano M, Kauzlaric A, Duc J, Coluccio A, Offner S, Imbeault M, Rowe HM, Turelli P,
Trono D. 2016. Transposable Elements and Their KRAB-ZFP Controllers Regulate Gene
Expression in Adult Tissues. *Dev Cell* **36**:611–623.

Ecco G, Imbeault M, Trono D. 2017. KRAB zinc finger proteins. *Development* **144**:2719–2729.

Ecker S, Chen L, Pancaldi V, Bagger FO, Fernández JM, Carrillo de Santa Pau E, Juan D,
Mann AL, Watt S, Casale FP, Sidiropoulos N, Rapin N, Merkel A, BLUEPRINT Consortium,
Stunnenberg HG, Stegle O, Frontini M, Downes K, Pastinen T, Kuijpers TW, Rico D, Valencia
A, Beck S, Soranzo N, Paul DS. 2017. Genome-wide analysis of differential transcriptional and
epigenetic variability across human immune cell types. *Genome Biol* **18**:18.

Ecker S, Pancaldi V, Valencia A, Beck S, Paul DS. 2018. Epigenetic and Transcriptional
Variability Shape Phenotypic Plasticity. *Bioessays* **40**. doi:10.1002/bies.201700148

Elmer JL, Hay AD, Kessler NJ, Bertozzi TM, Ainscough E, Ferguson-Smith AC. 2020. Genomic
properties of variably methylated retrotransposons in mouse. *bioRxiv*.
doi:10.1101/2020.10.21.349217

ENCODE Project Consortium, Moore JE, Purcaro MJ, Pratt HE, Epstein CB, Shoresh N, Adrian

J, Kawli T, Davis CA, Dobin A, Kaul R, Halow J, Van Nostrand EL, Freese P, Gorkin DU, Shen
Y, He Y, Mackiewicz M, Pauli-Behn F, Williams BA, Mortazavi A, Keller CA, Zhang X-O,
Elhajjajy SI, Huey J, Dickel DE, Snetkova V, Wei X, Wang X, Rivera-Mulia JC, Rozowsky J,
Zhang Jing, Chhetri SB, Zhang Jialing, Victorsen A, White KP, Visel A, Yeo GW, Burge CB,
Lécuyer E, Gilbert DM, Dekker J, Rinn J, Mendenhall EM, Ecker JR, Kellis M, Klein RJ, Noble
WS, Kundaje A, Guigó R, Farnham PJ, Cherry JM, Myers RM, Ren B, Graveley BR, Gerstein
MB, Pennacchio LA, Snyder MP, Bernstein BE, Wold B, Hardison RC, Gingeras TR,
Stamatoyannopoulos JA, Weng Z. 2020. Expanded encyclopaedias of DNA elements in the
human and mouse genomes. *Nature* **583**:699–710.

Faulk C, Barks A, Dolinoy DC. 2013. Phylogenetic and DNA methylation analysis reveal novel
regions of variable methylation in the mouse IAP class of transposons. *BMC Genomics* **14**:48.

Gagnier L, Belancio VP, Mager DL. 2019. Mouse germ line mutations due to retrotransposon
insertions. *Mob DNA* **10**:15.

Gardiner-Garden M, Frommer M. 1987. CpG islands in vertebrate genomes. *J Mol Biol*
196:261–282.

Gu T, Lin X, Cullen SM, Luo M, Jeong M, Estecio M, Shen J, Hardikar S, Sun D, Su J, Rux D,
Guzman A, Lee M, Qi LS, Chen J-J, Kyba M, Huang Y, Chen T, Li W, Goodell MA. 2018.
DNMT3A and TET1 cooperate to regulate promoter epigenetic landscapes in mouse embryonic
stem cells. *Genome Biol* **19**:88.

Guindon S, Dufayard J-F, Lefort V, Anisimova M, Hordijk W, Gascuel O. 2010. New algorithms
and methods to estimate maximum-likelihood phylogenies: assessing the performance of
PhyML 3.0. *Syst Biol* **59**:307–321.

Gunasekara CJ, Scott CA, Laritsky E, Baker MS, MacKay H, Duryea JD, Kessler NJ, Hellenthal
G, Wood AC, Hodges KR, Gandhi M, Hair AB, Silver MJ, Moore SE, Prentice AM, Li Y, Chen R,
Coarfa C, Waterland RA. 2019. A genomic atlas of systemic interindividual epigenetic variation
in humans. *Genome Biol* **20**:105.

Hernando-Herraez I, Heyn H, Fernandez-Callejo M, Vidal E, Fernandez-Bellon H, Prado-
Martinez J, Sharp AJ, Esteller M, Marques-Bonet T. 2015. The interplay between DNA
methylation and sequence divergence in recent human evolution. *Nucleic Acids Res* **43**:8204–
8214.

Hubley R, Finn RD, Clements J, Eddy SR, Jones TA, Bao W, Smit AFA, Wheeler TJ. 2016. The
Dfam database of repetitive DNA families. *Nucleic Acids Res* **44**:D81-9.

Huerta-Cepas J, Serra F, Bork P. 2016. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. *Mol Biol Evol* **33**:1635–1638.

Imbeault M, Helleboid PY, Trono D. 2017. KRAB zinc-finger proteins contribute to the evolution
of gene regulatory networks. *Nature* **543**:550–554.

Jacobs FM, Greenberg D, Nguyen N, Haeussler M, Ewing AD, Katzman S, Paten B, Salama
SR, Haussler D. 2014. An evolutionary arms race between KRAB zinc-finger genes ZNF91/93
and SVA/L1 retrotransposons. *Nature* **516**:242–245.

Karolchik D, Hinrichs AS, Furey TS, Roskin KM, Sugnet CW, Haussler D, Kent WJ. 2004. The
UCSC Table Browser data retrieval tool. *Nucleic Acids Res* **32**:D493-6.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. *Mol Biol Evol* **30**:772–780.

Kauzlaric A, Ecco G, Cassano M, Duc J, Imbeault M, Trono D. 2017. The mouse genome
displays highly dynamic populations of KRAB-zinc finger protein genes and related genetic
units. *PLoS One* **12**:e0173746.

Kazachenka A, Bertozzi TM, Sjoberg-Herrera MK, Walker N, Gardner J, Gunning R, Pahita E,
Adams S, Adams D, Ferguson-Smith AC. 2018. Identification, Characterization, and Heritability
of Murine Metastable Epialleles: Implications for Non-genetic Inheritance. *Cell* **175**:1259-
1271.e13.

Keane TM, Goodstadt L, Danecek P, White MA, Wong K, Yalcin B, Heger A, Agam A, Slater G,
Goodson M, Furlotte NA, Eskin E, Nelläker C, Whitley H, Cleak J, Janowitz D, Hernandez-
Pliego P, Edwards A, Belgard TG, Oliver PL, McIntyre RE, Bhomra A, Nicod J, Gan X, Yuan W,
van der Weyden L, Steward CA, Balasubramaniam S, Stalker J, Mott R, Durbin R, Jackson IJ,
Czechanski A, Assunção JAG, Donahue LR, Reinholdt LG, Payseur BA, Ponting CP, Birney E,
Flint J, Adams DJ. 2011. Mouse genomic variation and its effect on phenotypes and gene
regulation. *Nature* **477**:289–294.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. *Nat Methods*
9:357–359.

Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome. *Genome Biol* **10**:R25.

Leung A, Trac C, Jin W, Lanting L, Akbany A, Sætrom P, Schones DE, Natarajan R. 2013.
Novel long noncoding RNAs are regulated by angiotensin II in vascular smooth muscle cells.
Circ Res **113**:266–278.

Lewis CA Jr, Crayle J, Zhou S, Swanstrom R, Wolfenden R. 2016. Cytosine deamination and
the precipitous decline of spontaneous mutation during Earth's history. *Proc Natl Acad Sci U S*
A **113**:8194–8199.

Li B, Qing T, Zhu J, Wen Z, Yu Y, Fukumura R, Zheng Y, Gondo Y, Shi L. 2017. A
Comprehensive Mouse Transcriptomic BodyMap across 17 Tissues by RNA-seq. *Sci Rep*
7:4200.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**:3094–
3100.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv [q-bioGN].

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map format
and SAMtools. *Bioinformatics* **25**:2078–2079.

Li X, Liu Y, Salz T, Hansen KD, Feinberg A. 2016. Whole-genome analysis of the methylome
and hydroxymethylome in normal and malignant lung and liver. *Genome Res* **26**:1730–1741.

Loman NJ, Quick J, Simpson JT. 2015. A complete bacterial genome assembled de novo using
only nanopore sequencing data. *Nat Methods* **12**:733–735.

Long HK, King HW, Patient RK, Odom DT, Klose RJ. 2016. Protection of CpG islands from DNA
methylation is DNA-encoded and evolutionarily conserved. *Nucleic Acids Res* **44**:6693–6706.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**:550.

Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing reads.
EMBnet.journal **17**:10–12.

Mikkelsen TS, Ku M, Jaffe DB, Issac B, Lieberman E, Giannoukos G, Alvarez P, Brockman W,
Kim T-K, Koche RP, Lee W, Mendenhall E, O'Donovan A, Presser A, Russ C, Xie X, Meissner
A, Wernig M, Jaenisch R, Nusbaum C, Lander ES, Bernstein BE. 2007. Genome-wide maps of
chromatin state in pluripotent and lineage-committed cells. *Nature* **448**:553–560.

Morgan HD, Sutherland HG, Martin DI, Whitelaw E. 1999. Epigenetic inheritance at the agouti
locus in the mouse. *Nat Genet* **23**:314–318.

Nellåker C, Keane TM, Yalcin B, Wong K, Agam A, Belgard TG, Flint J, Adams DJ, Frankel WN,
Ponting CP. 2012. The genomic landscape shaped by selection on transposable elements
across 18 mouse strains. *Genome Biol* **13**:R45.

O'Doherty A, Ruf S, Mulligan C, Hildreth V, Errington ML, Cooke S, Sesay A, Modino S, Vanes
L, Hernandez D, Linehan JM, Sharpe PT, Brandner S, Bliss TVP, Henderson DJ, Nizetic D,
Tybulewicz VLJ, Fisher EMC. 2005. An aneuploid mouse strain carrying human chromosome
21 with Down syndrome phenotypes. *Science* **309**:2033–2037.

Oey H, Isbel L, Hickey P, Ebaid B, Whitelaw E. 2015. Genetic and epigenetic variation among
inbred mouse littermates: identification of inter-individual differentially methylated regions.
Epigenetics Chromatin **8**:54.

Qin C, Wang Z, Shang J, Bekkari K, Liu R, Pacchione S, McNulty KA, Ng A, Barnum JE, Storer
RD. 2010. Intracisternal A particle genes: Distribution in the mouse genome, active subtypes,
and potential roles as species-specific mediators of susceptibility to cancer. *Mol Carcinog*
49:54–67.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* **26**:841–842.

Rakyan VK, Blewitt ME, Druker R, Preis JI, Whitelaw E. 2002. Metastable epialleles in
mammals. *Trends Genet* **18**:348–351.

Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dündar F, Manke
T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic*
Acids Res **44**:W160-5.

Rebollo R, Galvão-Ferrarini M, Gagnier L, Zhang Y, Ferraj A, Beck CR, Lorincz MC, Mager DL.
2020. Inter-Strain Epigenomic Profiling Reveals a Candidate IAP Master Copy in C3H Mice.
Viruses **12**. doi:10.3390/v12070783

Reiss D, Zhang Y, Rouhi A, Reuter M, Mager DL. 2010. Variable DNA methylation of
transposable elements: the case study of mouse Early Transposons. *Epigenetics* **5**:68–79.

Rosenfeld CS, Sieli PT, Warzak DA, Ellersieck MR, Pennington KA, Roberts RM. 2013.
Maternal exposure to bisphenol A and genistein has minimal effect on A(vy)/a offspring coat
color but favors birth of agouti over nonagouti mice. *Proc Natl Acad Sci U S A* **110**:537–542.

Schultz DC, Ayyanathan K, Negorev D, Maul GG, Rauscher FJ. 2002. SETDB1: a novel KAP-1-
associated histone H3, lysine 9-specific methyltransferase that contributes to HP1-mediated
silencing of euchromatic genes by KRAB zinc-finger proteins. *Genes Dev* **16**:919–932.

Shen JC, Rideout WM 3rd, Jones PA. 1994. The rate of hydrolytic deamination of 5-
methylcytosine in double-stranded DNA. *Nucleic Acids Res* **22**:972–976.

Skene PJ, Henikoff S. 2017. An efficient targeted nuclease strategy for high-resolution mapping
of DNA binding sites. *Elife* **6**. doi:10.7554/eLife.21856

Smit, AFA, Hubley, R & Green, P. 2010. Repeat-Masker Open-3.0.

Song Q, Smith AD. 2011. Identifying dispersed epigenomic domains from ChIP-Seq data.
Bioinformatics **27**:870–871.

Stoye JP. 2001. Endogenous retroviruses: still active after all these years? *Curr Biol* **11**:R914-6.

Tahiliani M, Koh KP, Shen Y, Pastor WA, Bandukwala H, Brudno Y, Agarwal S, Iyer LM, Liu
DR, Aravind L, Rao A. 2009. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in
mammalian DNA by MLL partner TET1. *Science* **324**:930–935.

Tejedor JR, Fraga MF. 2017. Interindividual epigenetic variability: Sound or noise? *Bioessays*
39. doi:10.1002/bies.201700055

Thomson JP, Skene PJ, Selfridge J, Clouaire T, Guy J, Webb S, Kerr AR, Deaton A, Andrews
R, James KD, Turner DJ, Illingworth R, Bird A. 2010. CpG islands influence chromatin structure
via the CpG-binding protein Cfp1. *Nature* **464**:1082–1086.

Turelli P, Castro-Diaz N, Marzetta F, Kapopoulou A, Raclot C, Duc J, Tieng V, Quenneville S,
Trono D. 2014. Interplay of TRIM28 and DNA methylation in controlling human endogenous
retroelements. *Genome Res* **24**:1260–1270.

van de Lagemaat LN, Flenley M, Lynch MD, Garrick D, Tomlinson SR, Kranc KR, Vernimmen
D. 2018. CpG binding protein (CFP1) occupies open chromatin regions of active genes,
including enhancers and non-CpG islands. *Epigenetics Chromatin* **11**:59.

Ward MC, Wilson MD, Barbosa-Morais NL, Schmidt D, Stark R, Pan Q, Schwalie PC, Menon S,
Lukk M, Watt S, Thybert D, Kutter C, Kirschner K, Flicek P, Blencowe BJ, Odom DT. 2013.
Latent Regulatory Potential of Human-Specific Repetitive Elements. *Mol Cell* **49**:262–272.

Waterland RA, Jirtle RL. 2003. Transposable Elements: Targets for Early Nutritional Effects on
Epigenetic Gene Regulation. *Mol Cell Biol* **23**:5293–5300.

Waterland RA, Travisano M, Tahiliani KG. 2007. Diet-induced hypermethylation at agouti viable
yellow is not inherited transgenerationally through the female. *FASEB J* **21**:3380–3385.

Williams K, Christensen J, Pedersen MT, Johansen JV, Cloos PAC, Rappsilber J, Helin K.
2011. TET1 and hydroxymethylcytosine in transcription and DNA methylation fidelity. *Nature*
473:343–348.

Wolf G, de Iaco A, Sun M-A, Bruno M, Tinkham M, Hoang D, Mitra A, Ralls S, Trono D,
Macfarlan TS. 2020. KRAB-zinc finger protein gene expansion in response to active
retrotransposons in the murine lineage. *Elife* **9**. doi:10.7554/eLife.56337

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,
Brown M, Li W, Liu XS. 2008. Model-based Analysis of ChIP-Seq (MACS). *Genome Biol*
9:R137.

A

B

C

All IAPLTR1
with IAPEz-ints

VM-IAPLTR1
with IAPEz-ints

102   56   133

4
2

IAPLTR1 ■ Clade 1 ■ Clade 2 ■ Clade 3 ■ Clade 4
IAPEz-int ■ Clade α ■ Clade β ■ Clade γ □ Solo LTR

D

VM clade3
IAPLTR1

other clade3
IAPLTR1

■ Neither
■ Proximal to
an enhancer
■ Proximal to a
consitutively
expressed gene

Percent of IAPs

E

C57Bl6/NJ

NZO

DBA/AJ/NOD/AKR/
BALB/C3H/CBA

Lp/129

IAPLTR1
▼ Clade 1
▲ Clade 2
■ Clade 3
● Clade 4

IAPEz-int
— Clade α
— Clade β
— Clade γ

Percent Conserved

A Mouse ERV LTRs

IAPLTR1
IAPLTR2

VM %

# of LTRs
●100
●75
●50
●25

Ave CpG Score

B

CpG Score

Background   non VM-IAPLTR   VM-IAPLTR

C

TET1 ChIP

— Clade1
— Clade2
— Clade3
— Clade4

-2   center   2Kb

D

WT

Tet1 KO

Dnmt3a/
Tet1 DKO

Methylation

***
***

A

chr12:90564525-90565363

200bp ⊢————⊣

WT
H3K4me3

Trim28$^{+/D9}$
H3K4me3

IAPLTR2   Lx4A

B

Loci with novel
H3K4me3 in *Trim28*$^{+/D9}$

Percentage

Transposon

Unique
Muridae
Rodentia
Mus

C

            Wild Type   *Trim28*$^{+/D9}$

H3K4me3
ChIP-seq

CFP1
Cut&Run

-4 Peak   4kb   -4 Peak   4kb
    center   center

D

Trim28$^{+/D9}$ vs WT mice

p.val=0.0002

CpG Score

Increased   No change
expression