Dynamic genome plasticity during unisexual reproduction in the human fungal pathogen *Cryptococcus deneoformans*

Ci Fu, Aaliyah Davy, Simeon Holmes, Sheng Sun, Vikas Yadav, Asiya Gusa, Marco A. Coelho, and Joseph Heitman

Department of Molecular Genetics and Microbiology, Duke University Medical Center, Durham, NC, 27710, USA

Corresponding author:

Joseph Heitman, MD, PhD

Department of Molecular Genetics and Microbiology

Duke University Medical Center

322 CARL Building Box 3546

Durham, NC 27710, USA

Tel: 919-684-2824

Fax: 919-684-2790

Email: heitm001@duke.edu

Key words: ploidy duplication, cell cycle, cyclin, sexual reproduction, segmental aneuploidy

Abstract

Genome copy number variation occurs during each mitotic and meiotic cycle and it is crucial for organisms to maintain their natural ploidy. Defects in ploidy transitions can lead to chromosome instability, which is a hallmark of cancer. Ploidy in the haploid human fungal pathogen *Cryptococcus neoformans* is exquisitely orchestrated and ranges from haploid to polyploid during sexual development and under various environmental and host conditions. However, the mechanisms controlling these ploidy transitions are largely unknown. During *C. deneoformans* (formerly *C. neoformans* var. *neoformans*, serotype D) unisexual reproduction, ploidy increases prior to the onset of meiosis, can be independent from cell-cell fusion and nuclear fusion, and likely occurs through an endoreplication pathway. To elucidate the molecular mechanisms underlying this ploidy transition, we identified twenty cell cycle-regulating genes encoding cyclins, cyclin-dependent kinases (CDK), and CDK regulators. We characterized four cyclin genes and two CDK regulator genes that were differentially expressed during unisexual reproduction and contributed to diploidization. To detect ploidy transition events, we generated a ploidy reporter, called *NURAT*, which can detect copy number increases via double selection for nourseothricin-resistant, uracil-prototrophic cells. Utilizing this ploidy reporter, we showed that ploidy transition from haploid to diploid can be detected during the early phases of unisexual reproduction. Interestingly, selection for the *NURAT* reporter revealed several instances of segmental aneuploidy of multiple chromosomes, which conferred azole resistance in some isolates. These findings provide further

52 evidence of ploidy plasticity in fungi with significant biological and public health

53 implications.

Author Summary

Ploidy is an intrinsic fundamental feature of all eukaryotic organisms, and ploidy variation and maintenance are critical to the organism survival and evolution. Fungi exhibit exquisite plasticity in ploidy variation in adaptation to various environmental stresses. For example, the haploid opportunistic human fungal pathogen *C. deneoformans* can generate diploid blastospores during unisexual reproduction and also forms polyploid titan cells during host infection, however, the mechanisms underlying these ploidy transitions are largely unknown. In this study, we elucidated the genetic regulatory circuitry governing ploidy duplication during *C. deneoformans* unisexual reproduction through the identification and characterization of cell cycle regulators that are differentially expressed during unisexual reproduction. We showed that four cyclin and two cyclin-dependent kinase regulator genes function in concert to orchestrate ploidy transition during unisexual reproduction. To trace and track ploidy transition events, we also generated a ploidy reporter and revealed the formation of segmental aneuploidy in addition to diploidization, illustrating the diverse mechanisms of genome plasticity in *C. deneoformans*.

Introduction

Ploidy refers to the total number of chromosomal sets in a cell. Variations in ploidy are prevalent among both prokaryotic and eukaryotic organisms and have a profound effect on cellular phenotypes. Polyploidization has been suggested to provide adaptive advantages to environmental stresses through increases in gene copy number [1, 2]. Cells can achieve polyploidization through either genome doubling within a single species, called auto-polyploidization, or via hybridization of genomes from different species, termed allo-polyploidization [2]. Upon polyploidization, cells experience the immediate impacts of having twice the genome content, which can include changes in cell size, genome stability, and gene expression. Despite these often drastic and deleterious changes, cells regularly tolerate ploidy transitions during mitotic and meiotic cell cycles, in which the entire genome undergoes duplication and reduction [1].

In the fungal kingdom, ploidy variation among natural isolates of a single species is a common phenomenon [3]. For example, the baker's yeast *Saccharomyces cerevisiae*, which is an evolutionary product of ancient allo-polyploidization between two different ancestral species, has natural isolates with ploidy ranging from haploid to tetraploid [3-6]. *Candida albicans*, which was once thought to be an obligate diploid human fungal pathogen, has been shown to form haploid, triploid, and tetraploid cells [7, 8]. Nondiploid *C. albicans* cells have increased genomic instability and often return to a diploid or near-diploid state through auto-diploidization of the haploid genome or concerted chromosome loss of tetraploid cells, as *C. albicans* lacks a complete meiotic chromosomal

reduction cycle [8, 9]. In the syncytial hyphae of the filamentous fungus *Ashbya*

gossypii, nuclear ploidy ranges from haploid to higher than tetraploid within the

same hyphal compartment, and the degree of ploidy variation increases with

hyphal aging and decreases upon exposure to cellular stress [10]. The

prevalence of polyploidy in fungi illustrates how these genomic changes can

provide efficient strategies for fungal cells to rapidly adapt to their environment

[11].

Ploidy in the opportunistic human fungal pathogen *Cryptococcus* exhibits

exquisite plasticity during sexual reproduction and under host infection conditions

[12-14]. Cryptococcal infection can cause fatal cryptococcal meningitis in

immunocompromised patients. The mortality rate of cryptococcal meningitis is as

high as 70% for patients receiving treatment in resource-limited countries due to

a lack of cost-effective therapeutics, and mortality is 100% in those left untreated

[15]. *Cryptococcus* species have a bipolar mating system and undergo bisexual

reproduction, while *C. deneoformans* can also undergo unisexual reproduction in

the absence of a mating partner of the opposite mating type [16, 17]. During

bisexual reproduction, haploid *MAT*α and *MAT***a** cells undergo cell-cell fusion to

achieve genome doubling, while during unisexual reproduction, haploid cells

achieve genome doubling either via whole-genome duplication or through cell-

cell fusion events between cells of the same mating type [18]. In the natural

environment, *Cryptococcus* is largely present as haploid yeast cells, but diploid

cells of a single mating type (mainly αAAα) have also been documented,

demonstrating that the presence of unisexual reproduction in nature can

generate ploidy variation [19]. Besides ploidy transitions during sexual

reproduction, *Cryptococcus* can also form polyploid giant cells, termed titan cells,

during host infection [12, 20]. The ploidy of titan cells can reach up to 64 or more

copies of the genome, which is accompanied with morphological changes,

including increased cell size up to 100 μm in diameter (compared to standard

haploid cells that are 5 to 9 μm in diameter) and a thickened cell wall with a

dense cross-linked capsule [21, 22]. The formation of titan cells in host lung

tissue has been shown to enable fungal evasion of phagocytosis by host immune

cells and enhance fungal virulence [23, 24]. Polyploid titan cells can further

produce haploid and aneuploid progeny with enhanced tolerance to stressors

within the host environment, and meiotic genes have also been shown to be

activated in this niche [25, 26]. To utilize this ploidy plasticity, *Cryptococcus* has

evolved an elegant ploidy transition machinery that can be activated in response

to mating cues, environmental stresses, and host conditions.

The environmental stimuli that trigger diploid and polyploid cell formation

in *C. deneoformans* during unisexual reproduction and host infection have been

characterized and include cell density and quorum sensing molecules, nutrient

starvation, and serum [26-30]. However, the molecular mechanisms underlying

these ploidy transitions are less clear. In other eukaryotic organisms, increases in

ploidy are achieved primarily through endoreplication, during which, cells

undergo multiple rounds of S phase without entering mitosis and cytokinesis [31,

32]. This abnormal cell cycle is regulated by the same group of cyclins and

cyclin-dependent kinases that govern the progression of the mitotic cell cycle [31,

32]. For example, in flies and mammals, oscillation of cyclin E and cyclin-

dependent kinase 2 activity is required for endocycles of S phase [31]. In

Schizosaccharomyces pombe, mutants lacking the P34^{cdc2}P56^{cdc13} mitotic B

cyclin complex undergo multiple rounds of S phase and generate polyploid

progeny [33]. In *S. cerevisiae*, cell cycle progression is regulated by activation of

the cyclin-dependent kinase Cdc28 through binding of G1/S/G2/M-phase specific

cyclins [34]. Periodical oscillation of B-type cyclin *CLB6* in *clb1-5Δ* cells can drive

S. cerevisiae cells to re-enter S phase without undergoing mitosis and results in

polyploid cell formation [35]. In *C. neoformans*, it was recently shown that

reduced cyclin *CLN1* expression in cells arrested in G2 phase can lead to titan

cell formation [36]. Thus, it is likely that concerted regulation of these cell cycle

regulators in *C. deneoformans* contributes to diploidization during unisexual

reproduction.

In this study, we sought to identify cell cycle regulators that govern ploidy

transitions during unisexual reproduction in *C. deneoformans*. Because cell cycle

progression in *S. cerevisiae* is governed by transcript levels of cyclins, we initially

identified 20 putative cell cycle-regulating genes and examined their transcription

levels during unisexual reproduction [34]. Among them, six genes were

differentially expressed during unisexual reproduction compared to mitotic yeast

growth. Further examination of the ploidy of blastospores, the diploid products of

wild-type unisexual reproduction, confirmed that these genes are required for

ploidy transitions during unisexual reproduction. We also developed a *NURAT*

ploidy reporter to detect ploidy transition events and were able to detect both

diploidization as well as aneuploid and segmental aneuploid formation events

during both mitotic growth and unisexual reproduction, all of which underlie ploidy

plasticity in *Cryptococcus* species.

Results

Identification of cell cycle regulators involved in unisexual reproduction

In fungi, ploidy duplication is a prerequisite for meiosis during sexual reproduction and is largely achieved through gamete fusion. However, cell fusion and nuclear fusion are dispensable during unisexual reproduction in *C. deneoformans* and it has been proposed that an endoreplication pathway drives the haploid to diploid transition [18, 37]. To elucidate the endoreplication pathway for unisexual reproduction, we sought to identify cell cycle regulators that are critical for this ploidy transition. Because cyclin abundance and turnover regulate cyclin-dependent kinase (CDK) activities and drive cell cycle progression [34], we searched for cyclins in the *C. deneoformans* JEC21 genome on FungiDB (www.fungidb.org) [38] and identified 51 candidate genes (Table S1). Based on the annotated protein function for each gene, 20 genes were selected with predicted functions in the following three categories: cyclin (9), cyclin-dependent kinase (6), and CDK regulator (5) (Table S1).

Transcriptional profiling during unisexual reproduction revealed that genes involved in the pheromone response pathway, meiosis, and spore production were activated between 24 and 48 hours upon mating induction [27, 39]. We hypothesized that cell cycle genes important for ploidy duplication might be differentially expressed during unisexual reproduction. To determine this, we compared the expression levels of these putative cell cycle genes in wild-type *C. deneoformans* XL280α cells after incubation for 36 hours on mating-inducing

conditions (V8 agar medium) to yeast cell growth conditions (nutrient-rich YPD

medium) by qRT-PCR with *KAR5* serving as a negative control that has been

previously shown to be expressed at a comparable level under these two

conditions [18]. We found four cyclin genes and two CDK regulator genes were

significantly differentially expressed: *PCL2*, *CLB3*, and *CKS2* were

downregulated, while *PCL6*, *PCL9*, and *CKS1* were upregulated (Figure 1A).

Interestingly, none of the predicted cyclin-dependent kinase genes were

differentially expressed (Figure S1A).

In agreement with our findings, all six genes were previously shown to be

differentially expressed after growth on V8 medium for 12, 24, and 48 hours

compared to growth on YPD for 12 hours (Figure S1B). *PCL6*, *PCL9*, and *CKS1*

expression levels peaked at either the 24- or the 48-hour time point on V8

medium (Figure S1B). *CLB3* and *CKS2* were down-regulated on V8 medium,

while *PCL2* had an initial upregulation on V8 at the 12-hour time point and then

was down-regulated at later time points (Figure S1B) [39]. Three of the four

cyclin genes, *PCL2*, *PCL6*, and *PCL9*, are Pho85 cyclins with predicted functions

in regulating the cyclin-dependent kinase Pho85 in *S. cerevisiae* [40]. *CKS1* and

CKS2 are predicted to encode regulatory subunits for Cdc28, the master CDK for

cell cycle progression in *S. cerevisiae* (Table S1) [41].

To determine whether the differentially expressed cyclin and CDK

regulator genes are required for unisexual reproduction, we generated two

independent deletion mutants for each gene except for *CLB3*, for which we

generated a galactose-inducible allele under the control of the *GAL7* promoter

due to technical difficulty in deleting *CLB3* in our studies (Figure S2A). Deletion

of *PCL2, PCL6*, *CKS1*, or suppressed expression of *CLB3* caused a mild

reduction in hyphae production during unisexual reproduction, whereas deletion

of *PCL9* or *CKS2* did not cause any defect in hyphae formation (Figure 1B).

Deletion of *CKS1* or suppressed expression of *CLB3* also caused a defect in

sporulation leading to the production of bald basidia (basidia lacking spores),

strikingly different from the typical wild-type basidia with four chains of spores

produced (Figure 1B). These results suggest *CKS1* and *CLB3* are required for

the mitotic cycles during spore genesis, and they may also directly contribute to

cell cycle progression during the meiotic cycle. The differential expression

patterns and the observed morphological defects for the *pcl2Δ*, *pcl6Δ*, *cks1Δ*, and

P_{GAL7}-*CLB3* strains suggest that these cell cycle regulatory genes play critical

roles during *C. deneoformans* unisexual reproduction. Interestingly, deletion of

these cell cycle regulating genes had a smaller impact on bisexual reproduction,

especially that deletion of *CKS1* did not block basidium spore chain production

(Figure S2B), further corroborating the hypothesis that the expression of these

cell cycle regulating genes is coordinated during unisexual reproduction.

CKS1 and CLB3 promote G2/M phase progression

To examine if these cell cycle regulating genes also function during yeast

growth, we stained yeast cells with DAPI, which stains nuclei, and calcofluor

white (CFW), which stains chitin in the cell wall, to observe yeast cell morphology

for these mutant strains. Deletion of *CKS1* and suppressed expression of *CLB3*

induced pseudohyphal growth whereas other deletion mutants or the expression

of P_{GAL7}-*CLB3* in the presence of galactose all produced yeast cells with normal

morphologies (Figure S3), suggesting disruption in cell cycle progression can

trigger pseudohyphal formation in *C. deneoformans*, similar to previous findings

in *C. albicans* [42, 43].

To determine if these genes are involved in cell cycle progression during

yeast growth, we arrested cells from overnight cultures in liquid YPD medium at

the G1/S phase with hydroxyurea (arrest was subsequently released by

removing the reagent), and at G2/M phase with nocodazole [44-47]. Deletion of

CKS1 and suppressed expression of *CLB3* failed to respond to cell cycle arrest

reagents and cells were arrested at G2/M phase in the overnight culture even

before the treatment (Figure S4 and Table S4), providing strong evidence that

CKS1 and *CLB3* promote G2/M phase progression during yeast growth. This cell

cycle arrest may also contribute to the observed pseudohyphal growth in *cks1*Δ

mutants and the P_{GAL7}-*CLB3* strain in the presence of glucose (Figure S3).

Cell cycle regulators contribute to ploidy duplication during unisexual

reproduction

Cell cycle arrest at the G2 phase in large cell populations triggered by high

temperature or nocodazole has been shown to promote hyphal growth in *C.*

deneoformans [48], which is a hallmark of unisexual reproduction, illustrating a

potential intrinsic association between ploidy transition and unisexual

reproduction. To further characterize this association, we examined the ploidy

distribution in populations of wild-type and mutant cells grown overnight on YPD

and V8 agar media. Interestingly, although some mutants showed hyphal growth

defects during unisexual reproduction (Figure 1B), all samples exhibited similar

population distributions on mating-inducing medium compared to nutrient-rich

YPD medium (Figure S5). These results suggest that cell cycle arrest at G2

phase alone is not sufficient to promote unisexual reproduction, and that the

ploidy transition required for unisexual meiosis likely occurs in a small portion of

the cell population. Suppression of *CLB3* expression on V8 medium led to cell

cycle arrest at the G2 phase and caused defects in both hyphal growth and

sporulation (Figures 1B and 1C), further suggesting that cell cycle arrest at G2

phase is not sufficient to drive unisexual reproduction.

To understand how these cell cycle-regulating genes govern ploidy

duplication, we examined the ploidy of blastospores, which are cells produced by

mitotic budding directly from and along hyphae during unisexual reproduction. In

the wild type, all blastospores tested from ten different budding sites of different

hyphae were diploid except for two isolates that were aneuploid and originated

from the same budding site (Figure 2 and Table S2). These findings are in

agreement with previous studies and provide evidence that diploidization occurs

during or prior to blastospore formation [16, 18]. Compared to the wild type,

pcl2Δ, *pcl6Δ*, *cks1Δ*, and *cks2Δ* mutant strains produced blastospores with lower

germination rates and suppressed expression of *CLB3* caused a severe defect in

blastospore germination (Table S2). Ploidy determination for these germinated

blastospores showed that all of the gene deletions (except *PCL2*) as well as

suppressed expression of *CLB3* caused a defect in diploid blastospore formation

(Figure 2 and Table S2). More than 50% of the *pcl6Δ* blastospores produced

mixed populations of haploid and diploid cells, while about 80% of the *pcl9Δ* and

50% of the *cks2Δ* blastospores were haploid (Figure 2 and Table S2). Deletion of

CKS1 and suppressed expression of *CLB3* led to the production of only haploid

blastospores (Figure 2 and Table S2). However, due to a defect in blastospore

germination present in these mutants (Table S2), we could not determine the

ploidy status for these ungerminated blastospores. Collectively, our data suggest

that these cell cycle regulating genes contribute to ploidy duplication during

unisexual reproduction.

To further decipher the mixed haploid and diploid cell populations

observed among the *pcl6Δ* blastospores and determine if *PCL6* is required for

diploid maintenance, we streaked cells derived from diploid wild-type, diploid

pcl6Δ, and mixed haploid and diploid *pcl6Δ* blastospores for single colonies and

subsequently determined their ploidy (Table S3). Interestingly, all single colonies

derived from diploid wild type and *pcl6Δ* blastospores were diploid, while all

single colonies except one derived from mixed haploid-diploid *pcl6Δ* blastospores

were either haploid or diploid (Table S3). The observation of stable mitotically

passaged diploid cells suggests that *PCL6* is not required for diploid

maintenance during mitotic growth.

Detecting ploidy transitions during unisexual reproduction with a ploidy sensor

Because the 1N and 2N population distributions largely remained the same during unisexual reproduction and vegetative growth conditions in bulk culture (Figure S5), we hypothesized that the ploidy duplication required for meiosis during unisexual reproduction might only be occurring in the sub-population of cells that are committed to unisexual hyphal growth. To track this hypothesized diploidization/endoreplication event, we engineered a genetic construct called *NURAT*, similar to the *UAU1* cassette developed in *Candida albicans* [49], which allows for the detection of copy number increases in the genomic regions harboring this construct, which could be due to either aneuploidy formation or whole-genome duplication (Figure 3A) . The *NURAT* construct encodes a functional *URA5* gene flanked by truncated 5' and 3' *NAT* cassette sequences that share 530 bp of the *NAT* coding sequence (CDS), which allows homologous recombination to yield a functional allele of the *NAT* cassette and thus conferring nourseothricin resistance (Figure 3A). We integrated the *NURAT* ploidy reporter into a previously identified safe haven locus on Chromosome (Chr) 1 in *MAT***a** and *MAT*α strains in which the native *URA5* gene had been replaced by the hygromycin resistance *HYG* cassette (Figure S6) [50]. This ploidy reporter allows selection of nourseothricin resistant (NAT^R) and uracil-prototrophic (Ura$^+$) progeny; however, it depends on a copy number increase prior to homologous recombination in one of the two *NURAT* cassettes. If homologous recombination occurred between the truncated *NAT* CDSs in

haploid cells prior to diploidization, the nourseothricin-sensitive (NAT^S) and

uracil-prototrophic (Ura$^+$) haploid cell would become nourseothricin resistant

(NAT^R) and prevent selection of the second copy of *NURAT* cassette due to the

loss of the Ura$^+$ marker. Similarly, if homologous recombination occurred in both

copies of the *NURAT* construct in diploidized cells, both *NURAT* cassettes

become active *NAT* markers, which prevents the selection of NAT^R Ura$^+$ diploid

cells (Figure 3A). Normal homologous recombination functioning during either

mitotic or meiotic growth is a prerequisite for the ploidy sensor to detect copy

number variance.

To test the robustness of the *NURAT* reporter in detecting diploid cells

versus haploid cells, we generated two diploid *NURAT*/*NURAT* strains through

blastospore dissection of the haploid *NURAT* strain (Figure S6) and performed

fluctuation assays (Figure S7). The haploid and diploid *NURAT* strains share the

same genomic sequences and only differ in ploidy. In overnight liquid cultures,

haploid and diploid *NURAT* strains exhibited similar *NURAT* recombination

frequencies despite diploid *NURAT* strains having two copies of the *NURAT*

construct (Figure S8A). Among the NAT^R colonies, about 1% of the haploid

*MAT***a** cells and 0.04% of the *MAT*α cells were Ura$^+$, whereas 73% and 79% of

the diploid cells were Ura$^+$, suggesting homologous recombination occurred in

only one copy of the *NURAT* construct in most diploid cells (Figure S8A).

Interestingly, haploid and diploid NAT^R, Ura$^+$ colonies maintained their ploidy

(Figure S8A). Our findings suggest that in the overnight cultures of haploid

strains, very few cells undergo diploidization. Nevertheless, using the *NURAT*

construct, we were still able to detect possible aneuploidy of chromosome 1 (on

which the *NURAT* construct is located) that occurred at a low frequency.

To examine whether the *NURAT* reporter could be used to detect ploidy

transition events during unisexual reproduction, we incubated haploid and diploid

MATα NURAT strains on both YPD and V8 agar media for 36 and 60 hours

(Figure S7). After 36 hours of incubation on the mating-inducing V8 medium, only

haploid *NURAT* cells displayed a significant increase in *NURAT* recombination

compared to cells incubated on YPD (Figure S8B). Interestingly, both haploid

and diploid cells showed a significant increase in *NURAT* recombination after

incubation for 60 hours on V8 medium (Figure 3B), illustrating a possible

elevated rate in homologous recombination under mating-inducing conditions.

Among the recombined NAT^R colonies, haploid cells had a much lower

percentage of Ura^+ colonies compared to diploid cells (about 1% in haploid and

50-70% in diploid) (Figure3B). Additionally, activation of unisexual reproduction

after longer incubation on V8 medium did not lead to increased numbers of NAT^R,

Ura^+ cells in the diploid populations, but a significant increase of NAT^R, Ura^+ cells

was observed in the haploid populations (Figures 3B and S8). FACS analyses of

individual colonies showed that there were more diploid colonies than haploid

cells after longer incubation on V8 (Figures 3B and S8), suggesting that

diploidization occurs during unisexual reproduction, and the *NURAT* construct

can indeed detect ploidy transition events. However, the low frequency of

detected diploidization events in this assay also suggests that ploidy duplication

during early time points of unisexual reproduction is occurring in a sub-population

of cells whereas other cells within the mating patch undergo mitotic growth as

haploid isolates. It is important to note that the sensitivity of this ploidy sensor to

detect ploidy changes is limited by the frequency of mitotic recombination of the

NURAT reporter.

Segmental aneuploidy occurs during both mitosis and meiosis

To understand the nature of the NAT^R, Ura$^+$ colonies, which harbor both

NAT and *NURAT* alleles (referred to subsequently as *NURAT*/*NAT* strains), we

performed whole-genome Illumina sequencing of five *NURAT*/*NAT* colonies

derived from mitotic passages of both the *MAT***a** and *MAT*α *NURAT* strains

(Figure S6). Chromosome alterations were inferred from changes in coverage of

reads aligned to a newly obtained, chromosome-level, reference genome

assembly of *C. deneoformans* XL280α, generated *de novo* by Oxford Nanopore

sequencing (see methods for details). Instead of observing the expected

aneuploidy of Chr 1 where the *NURAT* construct has been inserted (the safe

haven locus is located near the end of the chromosome arm), segmental

aneuploidy and about one extra copy of the region harboring the *NURAT*

construct was present in all of the *NURAT*/*NAT* colonies (Figures 4A, 4B, and

S9). Besides the segmentally duplicated region on Chr 1 that was selected for,

other chromosomal abnormalities were also detected, including segmental

duplications of Chr 2 in the diploid *MAT***a** *ura5Δ NURAT*/*NAT*-4 isolate, Chr 6 in

the *MAT*α *ura5Δ NURAT*/*NAT*-1 isolate, Chr 10 in *MAT***a** *ura5Δ NURAT*/*NAT*-2

isolate, and Chr 13 in the *MAT***a** *ura5Δ NURAT*/*NAT*-1 and -2 isolates (Figure 4A).

Interestingly the segmentally duplicated region on Chr 13 in *MAT***a** *NURAT*/*NAT*

colonies 1 and 2 spanned the centromere, which could potentially give rise to a

dicentric chromosome. Loss of chromosomal segments (inferred as regions with

lower read coverage) were only detected in the context of the diploid *MAT***a**

ura5Δ NURAT/*NAT*-4 isolate (on Chrs 2 and 4).

Segmentally duplicated regions on Chr 1 in all progeny also harbored a

drug efflux pump gene, *AFR1*, which has been shown to confer fluconazole

resistance in *Cryptococcus* (Figure 4B) [51]. Interestingly, three out of five strains

tested were found to be resistant to fluconazole, suggesting additional epistatic

interactions with the *AFR1* gene or unidentified mutations mitigated the

fluconazole resistance phenotype conferred by *AFR1* gene copy number

increase (Figure 4C). We also found that two of the strains were hypersensitive

to 37°C, a phenotype that has been associated with aneuploidy in *C. neoformans*

(Figure 4C) [14].

Segmental aneuploidies are formed via multiple mechanisms

To elucidate the mechanism(s) giving rise to segmental aneuploidy, we

analyzed the genomic regions flanking the segmental duplications detected in

different *NURAT*/*NAT* progeny. Specifically, we assessed if read-pairs aligned to

those regions in the XL280α reference genome had unexpected separation

distances, anomalous orientations, or if the forward and reverse reads of a pair

aligned to different chromosomes, all indicative of structural changes. This

analysis revealed distinct modes of segmental aneuploid formation among

different *NURAT*/*NAT* progeny (Figure S9). Sequencing reads bridging

segmentally duplicated regions on different chromosomes were detected in three

isolates suggesting potential fusion events may have occurred between

segmentally duplicated regions: Chrs 1 and 13 (a1-a2) and Chrs 1 and 6 (b1-b2)

in the *MAT**a** ura5Δ NURAT*/*NAT*-1 strain, Chrs 1 and 13 (a1-a2 and c1-c2) and

Chrs 1 and 10 (b1-b2) in the *MAT**a** ura5Δ NURAT*/*NAT*-2 strain, and Chrs 1 and

2 (a1-a2) and Chrs 2 and 4 (c1-c2) in the *MAT**a** ura5Δ NURAT*/*NAT*-4 strain

(Figure S9). In all three isolates, one segmentally duplicated region

encompasses the centromere (Chr 13 in the *MAT**a** ura5Δ NURAT*/*NAT*-1 and -2,

and Chr 1 in *MAT**a** ura5Δ NURAT*/*NAT*-4) allowing the opportunity for

neochromosome formation through fusion of segmentally duplicated regions

originating from different chromosomes. Segmental aneuploidy formation via

tandem duplications were detected for Chr 1 in the *MAT**a** ura5Δ NURAT*/*NAT*-3

strain and for Chrs 1 and 6 in the *MAT*α *ura5Δ NURAT*/*NAT*-1 strain, in which

large and small inversion events were detected, suggesting that complex

chromosomal rearrangements are also associated with segmental aneuploidy

formation. Transposable element movements have been shown to be highly

mutagenic, especially under host infection or temperature stress [52].

Interestingly, T1 and T3/CNIRT4 transposon movements were detected in

sequences flanking some of the *NURAT*/*NAT* progeny, suggesting that

transposable elements may also contribute to formation of some of the

segmental aneuploidies (Figure S9).

The segmentally duplicated regions were further analyzed through

separation of chromosomes via CHEF gel electrophoresis followed by chromo-

blotting with probes specific to these regions. These methods revealed that

various forms of karyotypic changes were present in the *NURAT*/*NAT* progeny

(Figure S10). The *NAT* probe for the *NURAT* construct in the *MAT***a** *ura5Δ*

NURAT/*NAT*-1 strain and both the *NAT* and *URA5* probes in the *MAT***a** *ura5Δ*

NURAT/*NAT*-2 strain hybridized to a smaller chromosome than expected (Figure

S10A, green arrows). Additionally, the probe specific to the duplicated region on

Chr 13 in the *MAT***a** *ura5Δ NURAT*/*NAT*-1 strain hybridized to two smaller

chromosomes, supporting the hypothesis that segmentally duplicated regions

can form neochromosomes (Figure S10C, green arrows). Hybridization of the

NAT and *URA5* probes to a smaller chromosome in the *MAT***a** *ura5Δ*

NURAT/*NAT*-3 and the *MAT*α *ura5Δ NURAT*/*NAT*-1 strains suggested the

segmentally duplicated events did not likely occur directly on Chr 1 (Figure S10A,

red arrows). Conversely, hybridization of a probe specific to the segmentally

duplicated region on Chr 6 in the *MAT*α *ura5Δ NURAT*/*NAT*-1 strain supported

the tandem duplication of the region within Chr 6, as detected by whole-genome

sequencing (Figure S9 and S10B, green arrow). Overall, the different types of

segmental aneuploidy formation illustrate a significant level of genomic and

ploidy plasticity in *C. deneoformans*.

Cell cycle regulators contribute to segmental aneuploidy formation

To investigate how the identified cell cycle-regulating genes impact ploidy

transitions during unisexual reproduction, we generated mutant strains containing

both the *NURAT* construct and the *ura5* gene deletion through meiotic crosses

and performed fluctuation assays (Figure S7). Compared to the wild type, genetic

deletion of *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* all significantly reduced

NURAT recombination frequencies. *pcl6Δ* and *cks1Δ* mutants exhibited the most

severe defects, suggesting modulation of ploidy transitions by these cell cycle-

regulating genes could influence the regulation of homologous recombination

during unisexual reproduction (Figure 5A). Interestingly, suppressed expression

of *CLB3* significantly increased *NURAT* recombination whereas upregulated

expression of *CLB3* (by supplementing galactose in V8 medium) significantly

reduced *NURAT* recombination, indicating that *CLB3* plays an opposite role in

contributing to homologous recombination frequencies during unisexual

reproduction. Addition of glucose or galactose in V8 medium also increased

homologous recombination frequencies in the wild type, which is likely due to

robust vegetative growth in the presence of excess nutrients.

Despite the significantly reduced *NURAT* recombination frequencies

observed in the deletion mutants, they produced significantly different

frequencies of NAT^R, Ura$^+$ colonies than the wild type except for *pcl9Δ* mutants.

pcl2Δ, *pcl6Δ*, and *cks2Δ* mutants produced zero or only one NAT^R, Ura$^+$ colony

(Figure 5A), suggesting that these three genes all function in driving

diploidization and segmental aneuploidy formation during unisexual reproduction.

In contrast, *cks1Δ* mutants, which had a severe defect in *NURAT* recombination,

produced significantly more NAT^R, Ura$^+$ colonies than the wild type (56% in

cks1Δ-1 and 32% in *cks1Δ-2* compared to 0.9% in wild type) (Figure 5A),

indicating that *CKS1* plays an inhibitory role in diploidization and segmental

aneuploidy formation. Interestingly, both suppressed and upregulated expression

of *CLB3* increased the frequency of NAT^R, Ura$^+$ colonies relative to the wild type

(Figure 5B), suggesting that *CLB3* expression levels modulate diploidization and

segmental aneuploid formation.

492   To evaluate the impact of these cell cycle genes on ploidy transitions, the

average ploidy was calculated for all *NURAT*/*NAT* colonies of wild type and

mutants (Figure 5). For *pcl9Δ* mutants that produced a comparable number of

NAT^R, Ura$^+$ colonies to wild type, deletion of *PCL9* significantly increased the

average ploidy level when outliers with ploidies above 1.6 were removed from the

dataset (Figure S11A). Both suppression and upregulation of *CLB3* expression

significantly decreased the average ploidy of the *NAT*/*NURAT* colonies when the

same outliers were removed (Figure S11B). Taken together, our findings suggest

that cell cycle-regulating genes act in concert to control ploidy transitions during

unisexual reproduction.

Discussion

Genome size changes occur during both mitotic and meiotic cycles and disruption of these processes leads to dire consequences on cellular viability and fertility. During mitosis, the whole genome duplicates during S phase and is governed by cell cycle regulators, while in fungi, during meiosis, ploidy duplication is accomplished via cell-cell fusion and nuclear fusion between mating partners [1, 32]. Besides these two fundamental cellular processes, ploidy transitions also occur during unisexual reproduction and titan cell formation in the human fungal pathogen *C. deneoformans*, in which ploidy increases through putative endoreplication pathways [13, 18, 21, 36]. To examine the mechanism(s) underlying the ploidy transition from haploid to diploid during unisexual reproduction, we identified and characterized six cell cycle-regulating genes that contribute to diploidization.

Among the identified twenty putative cell cycle-regulating genes, four cyclin genes and two CDK regulator genes were differentially expressed in *C. deneoformans* during unisexual reproduction compared to mitotic growth, whereas none of the cyclin-dependent kinase genes exhibited expression differences. This finding is in agreement with previous studies in *S. cerevisiae* that found fluctuations in transcript levels of cyclin genes, but not CDK genes, drive cell cycle progression [34]. Three of the four cyclin genes, *PCL2*, *PCL6*, and *PCL9*, are Pho85 cyclins (Pcls), which in *S. cerevisiae* interact with the cyclin-dependent kinase Pho85 [53]. Pcl2 and Pcl9 belong to the Pcl1,2-like subfamily and are expressed during G1 phase activating Pho85, which promotes

the G1/S phase transition [40, 53]. Pcl6 belongs to the Pho80 subfamily and

functions with Pho85 in activating the serine/threonine protein phosphatase Glc7,

which modulates kinetochore-microtubule interactions during M phase [53]. In *C. deneoformans*, expression of *PCL6* and *PCL9* was highly induced during

unisexual reproduction, and their deletion caused a defect in diploid blastospore

formation, while in contrast, *PCL2* was down-regulated during unisexual

reproduction and its deletion did not cause a defect in diploid blastospore

formation, suggesting concerted regulation of these cyclins are critical in driving

diploidization during unisexual reproduction, likely via modulation of Pho85

activity. The fourth identified gene, *CLB3*, belongs to B-type cyclins and its

expression was reduced during unisexual reproduction. However, repression of

CLB3 during unisexual reproduction blocked diploid blastospore formation. In *S. cerevisiae*, *CLB3* is a nonessential cyclin gene expressed during G2/M phase,

activating Cdc28 and promoting G2/M phase transition [34]. In *C. deneoformans*,

CLB3 was also not essential, and Pengjie Hu and colleagues have successfully

deleted *CLB3* utilizing a CRISPR-mediated transformation technique [54, 55].

Our failure in deleting this gene was likely due to the combinatory effect of low

biolistic transformation efficiency and a severe defect in cytokinesis of the mutant.

Deletion of *CLB3* caused defects in stress resistance, melanin production, and

capsule formation, and the deletion strain was avirulent in a mouse infection

model [54, 56]. Interestingly, different from the deletion mutants of the other five

cell cycle regulating genes, deletion of *CLB3* blocked basidium spore chain

production during bisexual reproduction [54], suggesting that Clb3 plays a pivot

548 role in sporogenesis during both modes of sexual reproduction in *C.*

549 *deneoformans.* In this study, suppression of *CLB3* expression led to cell cycle

550 arrest at G2/M phase, which is consistent with the phenotype observed for the

551 deletion mutant, suggesting that Clb3 contributes to G2/M phase progression, a

552 conserved role for Clb3 in *S. cerevisiae* [34, 54].

553   The two differentially expressed CDK regulator genes *CKS1* and *CKS2*

are both homologs of *S. cerevisiae CKS1*, which is required for G1/S and G2/M

phase transitions [41]. In *S. cerevisiae*, Cks1 functions as a phosphor-adaptor

protein for the CDK inhibitor Sic1 and the G1/S phase Cln-Cdc28 complex, which

facilitates phosphorylation and destruction of Sic1 at the onset of S phase [57].

Similar to *C. deneoformans*, humans have two homologs of the *S. cerevisiae*

CKS1 gene, and both can complement a null mutation of *CKS1* in *S. cerevisiae*

[58]. Interestingly, the *C. deneoformans CKS1* and *CKS2* genes were oppositely

regulated during unisexual reproduction. Deletion of *CKS1*, which is upregulated

during unisexual reproduction, caused a more severe phenotype during

unisexual spore production and diploid blastospore formation compared to

deletion of *CKS2*, which is downregulated during unisexual reproduction. These

results suggest a functional divergence between *CKS1* and *CKS2* in *C.*

deneoformans.

567   Among the six characterized cyclin and CDK regulator genes, *PCL9*,

CLB3, and *CKS2* were previously shown to be periodically expressed during the

cell cycle in *C. neoformans* with a peak at G2/M phase for *PCL9* and a peak at

G1/S phase for *CLB3* and *CKS2* [59]. Although *PCL2* and *CKS1* were not

periodically expressed during the cell cycle, expression of both genes peaked at

G1/M phase during the first cell cycle in synchronized cells in *C. neoformans*,

supporting a role for these cell cycle regulating genes in cell cycle progression

[59]. Four of these cell cycle-regulating genes (*CLB3*, *PCL2*, *CKS1*, and *CKS2*)

were highly expressed in the G1/S phase under rich growth conditions in *C.*

neoformans; however, deletion of *CKS1* and repression of *CLB3* in *C.*

deneoformans caused a prominent phenotype in G2/M phase arrest. More

interestingly, *CKS1* and *CLB3* expression were differentially regulated during

unisexual reproduction, suggesting a rewired transcriptional coordination of

different cell cycle-regulating genes may be is required for genome duplication

and diploid genome maintenance before the onset of meiosis during unisexual

reproduction. However, mechanisms underlying this process remain to be

elucidated.

Different from unisexual reproduction, ploidy increases during titan cell

formation was recently shown to be modulated by *CLN1* expression in *C.*

neoformans [36]. Interestingly, the orthologous *CLN1* gene (CNM00990) in *C.*

deneoformans was not differentially expressed during unisexual (Figure 1A).

Under host environment, unbudded *C. neoformans* cells were arrested in G2 with

a 2C genome and reduced *CLN1* expression allowed 2C cells to re-enter the

G1/S phase without mitosis to form polyploid titan cells [36] These distinct cyclin

expression regulations during different cellular and developmental processes

highlight the exquisite genome and ploidy adaptability of *Cryptococcus* species in

response to various host and environmental stress cues.

To further detect ploidy transition events during unisexual reproduction,

we generated the *NURAT* ploidy reporter, using which, we were able to detect

diploid cells under both mating-suppressive (YPD) and mating-inducing (V8)

culture conditions by selecting NAT^R, Ura$^+$ colonies. Although more diploid cells

were detected on V8 compared to YPD, the overall frequency of diploid cells was

low, suggesting that diploidization may occur in only a small number of cells that

are primed for unisexual reproduction, diploidization events could be transient, or

the relative sensitivity of detection with the *NURAT* reporter is limited by the

frequency of recombination.

Among the selected NAT^R, Ura$^+$ colonies from both YPD and V8 culture

condition, many remained haploid. Whole-genome sequencing of these haploid

colonies showed that these colonies had segmental aneuploidies of the region on

Chr 1 where the *NURAT* cassette was inserted. Besides the selected region on

Chr 1, other chromosomes which were not under selection also exhibited

segmental aneuploidy formation. Segmental aneuploidy represents a form of

chromosome instability, a hallmark of tumorigenesis, which occurs via breakage-

fusion-bridge cycles of duplication and multiple amplifications of certain

chromosome regions [60]. In the human fungal pathogen *C. albicans*, segmental

aneuploidy has been observed in azole-resistant isolates derived clinically and

experimentally, with multiple amplified regions containing genes, such as *TAC1*

and *ERG11* that contribute to azole tolerance [61-63]. In *C. albicans*, segmental

aneuploidy is observed exclusively in regions flanked by long inverted repeat

sequences, which function in repairing the breakage-fusion-bridge cycles during

the formation of segmental aneuploidy [63, 64]. Long-repeat sequences are

distributed throughout the *C. albicans* genome, suggesting a strong potential for

segmental aneuploidy formation [63]. Interestingly, in the absence of the drug,

the azole-resistant, segmentally aneuploid isolates can quickly lose the multiple

amplified copies and return to euploid chromosomal karyotypes and azole

susceptibility, suggesting a selective pressure may be required for the

maintenance of segmental aneuploidy [64].

Different from the segmental aneuploidy formation in *C. albicans*, none of

the characterized segmental aneuploid regions observed in this study are flanked

by inverted long-repeat sequences. Instead, many segmental aneuploid regions

are flanked by T1 or T3/CNIRT4 DNA transposons. In *C. neoformans*,

movements of both DNA transposons have been observed at elevated rates *in*

vivo and at host temperature and contribute to development of resistance against

multiple antifungal drugs [52]. It is likely that transposons play a role in segmental

aneuploidy formation and contribute to chromosome instability. Based on

detection of discordant read-pairs at the borders of the segmental aneuploid

regions, there appears to be two likely modes of segmental aneuploidy formation.

In the *MAT***a** *ura5Δ NURAT/NAT*-1 and -2 strains, forward and reverse reads of a

subset of the aligned read-pairs were found on different chromosomes, and one

of the regions always contained a centromere, suggesting segmental aneuploidy

formation may occur in conjunction with chromosomal fusion and neo-

chromosome formation. This hypothesis was confirmed by binding of a

chromoblot probe for the *NAT* cassette to two distinct chromosomes. On the

other hand, in the *MAT**a** ura5Δ NURAT*/*NAT*-3 and *MAT*α *ura5Δ NURAT*/*NAT*-1

strains, segmental aneuploidy formed on the same chromosome and involved

tandem duplications and inversions, likely through breakage-fusion-bridge cycles

similar to what has been observed in *C. albicans*. However, chromoblot analysis

for the segmental aneuploid regions only confirmed the size increase in Chr 6 for

the *MAT*α *ura5Δ NURAT*/*NAT*-1 strain. Surprisingly, chromoblot analysis for both

the *URA5* gene and *NAT* cassette detected the presence of a smaller

chromosome in the *MAT**a** ura5Δ NURAT*/*NAT*-3 and the *MAT*α *ura5Δ*

NURAT/*NAT*-1 strains, suggesting additional segmental aneuploidy formation

mechanisms occur and remain to be elucidated.

Recent experimental selection experiments have shown that breakage or

deletion of centromeres in *Cryptococcus* species can induce chromosomal

translocation and chromosomal fusion events, which may lead to reproductive

isolation, underlying the involvement of karyotypic variation in speciation during

evolution [65, 66]. However, detection of chromosomal instability events is

extremely difficult as naturally occurring karyotypic changes are rare events

derived from defective mitosis and that abnormal karyotypes often result in

cellular fitness costs [67, 68]. Here we showed that the *NURAT* cassette has

tremendous potential for isolating cells with altered karyotypes, which may prove

to be a valuable tool for elucidating mechanisms underlying chromosome

instability.

Utilizing the *NURAT* cassette, we characterized the involvement of cell

cycle-regulating genes in diploidization during unisexual reproduction. However,

because diploidization could be detected at a low frequency during the early

stages of unisexual reproduction, double selection of the *NURAT* reporter yielded

mostly aneuploid isolates. Deletion of cyclin and CDK regulator genes and

altered expression of *CLB3* all impacted *NURAT* recombination, suggesting

perturbation of cell cycle progression suppresses homologous recombination.

Because the *NURAT* reporter requires homologous recombination to detect

ploidy transition events, defects in this function will likely limit the use of *NURAT*

in probing ploidy transition. Despite this limitation, deletion of *PCL9* or *CKS1* did

not block double selection for the *NURAT* and the recombined *NAT* cassettes.

Interestingly, deletion of *PCL2*, *PCL6*, and *CKS2* almost completely prohibited

double selection for the ploidy reporter, while these three genes had a minimal

impact on diploid blastospore formation, suggesting diploidization and aneuploidy

formation may require distinct cell cycle regulatory circuitries.

Detection of the segmental aneuploidy events in this study illuminate yet

another example of the diverse mechanisms of genome plasticity in *C.*

neoformans. Unlike diploidization during unisexual reproduction and

polyploidization during titan cell formation in the host environment, segmental

aneuploidy is more likely to be the result of a rare mitotic error than to be

regulated by defined genetic pathways. Despite the rarity, three out of the five

characterized segmental aneuploid isolates exhibited resistance to the antifungal

azole drug fluconazole compared to their progenitor strain, which was likely due

to an increase in the copy number of the azole efflux pump gene *AFR1*,

suggesting that, under selection, these rare events could provide fitness benefits

for these cells to adapt to environmental stresses [51]. However, the prevalence

and biological significance of segmental aneuploidy in different environmental

and clinical isolates or strains with different ploidy status requires further

experimental exploration.

Materials and methods

Strains, media, and growth conditions

Strains and plasmids used in this study are listed in Table S5. All strains were generated in the congenic *MAT***a** and *MAT*α XL280 strain backgrounds [69]. Strains were frozen at -80°C in glycerol and maintained on Yeast Extract Peptone Dextrose or Glucose (YPD or YPG) agar medium for routine use. Strains harboring dominant selectable markers were grow on YPD or YPG medium supplemented with 100 μg/mL nourseothricin (NAT), 200 μg/mL G418 (NEO), or 200 μg/mL hygromycin (HYG). Synthetic dextrose or galactose medium without uracil (SD-URA or SG-URA) was used to select uracil prototrophic progeny. Unisexual and bisexual mating assays were induced on either 5% V8 juice agar medium (pH = 7) or Murashige and Skoog (MS) medium minus sucrose (Sigma-Aldrich) in the dark at room temperature for the designated time period.

Identification of putative cell cycle genes

To identify genes involved in cell cycle control, the key word cyclin was used to search against the *C. deneoformans* JEC21 genome on FungiDB (www.fungidb.org) [38]. BLASTP searches were performed for candidate cell cycle regulating genes against the *S. cerevisiae* genome database (www.yeastgenome.org), and then reciprocal BLASTP searches of top candidate

genes in *S. cerevisiae* were conducted against the *C. deneoformans* JEC21

genome database to provide putative gene names and predicted protein

functions (Table S1).

Expression levels of the putative cell cycle genes during unisexual

reproduction

The wild-type XL280α strain was grown overnight in YPD liquid medium

and adjusted to OD_{600}=2 in sterile H_2O. Then 10 µl of cells were spot inoculated

on V8 (mating-inducing condition) and YPD (non-mating condition) agar media

and incubated for 36 hours. RNA extraction and qRT-PCR were performed as

previously described [18]. Gene expression levels were normalized using the

endogenous reference gene *GPD1* and determined by using the comparative

ΔΔCt method. Expression fold change on V8 versus YPD agar media for each

putative cell cycle-regulating gene was compared to *KAR5*, which has been

previously shown to be expressed in the XL280α strain at a comparable level on

V8 and YPD agar [18]. Primers used for qRT-PCR are listed in Table S6.

Deletion of putative cell cycle genes and conditional expression of *CLB3*

The primers used in this section are listed in Table S6. Coding sequences

(CDS) for six differentially expressed putative cell cycle genes *PCL2*, *PCL6*,

PCL9, *CLB3*, *CKS1*, and *CKS2* were replaced by the dominant selectable marker

NEO cassette through homologous recombination as previously described [70].

In brief, for each gene, a deletion construct consisting of 1 kb upstream and 1 kb downstream sequences flanking the CDS and the *NEO* cassette was generated by overlap PCR, and then the deletion construct was introduced into the wild type XL280α strain via biolistic transformation. Stable transformants were selected on YPD medium supplemented with G418 (200 mg/l) and gene replacements were confirmed by PCR. Two independent deletion mutants were generated for *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* deletion mutants.

Biolistic transformation using the deletion construct for *CLB3* failed to generate a deletion mutant. To study *CLB3*, a conditional expression allele of *CLB3* under a galactose inducible promoter was generated by replacing a 300-bp region upstream of the *CLB3* start codon with a *NEO* cassette followed by a 1034-bp promoter sequence for the *GAL7* (CNM00600) (P_{GAL7}) gene using the TRACE method [55, 71]. To generate the regulated expression construct, the *NEO* cassette, the 1034-bp P_{GAL7} sequence, the 1117-bp upstream and 1019 bp downstream sequences of the 300-bp region were cloned into the pXL1 plasmid using the Gibson cloning method resulting in plasmid pSH5 [72]. Then the regulated expression construct was PCR amplified from the plasmid pSH5 using primer pair JOHE45301/JOHE46452. For the sgRNA expression construct, the U6 promoter and the sgRNA scaffold that share 20 bp of overlapping sequence targeting the 300 bp region were amplified from XL280α genomic DNA and the plasmid pYF515 respectively, and then the intact sgRNA expression construct was generated by overlap PCR [55, 73]. The *CAS9* expression construct was amplified from the plasmid pXL1-CAS9-HYG [55]. The regulated expression

construct, the sgRNA expression construct, and the *CAS9* expression construct

were transformed into wild-type XL280α cells through electroporation using a

BIO-RAD Gene Pulser. Stable transformants were selected on YPG medium

supplemented with G418 and the correct integration of the *GAL7* promoter was

confirmed in the transformant CF1715 by PCR.

To validate that the *GAL7* promoter could regulate *CLB3* expression, wild-

type XL280α and CF1715 (*NEO-P$_{GAL7}$-CLB3*) strains were grown overnight in

liquid YPD medium and adjusted to OD_{600}=2 in sterile H_2O. Then 10 µl of cells

were spot-inoculated on YPD, YPG, V8, V8 + 2% glucose, and V8+ 2%

galactose agar media and incubated for 36 hours. RNA extraction and qRT-PCR

were performed as previously described to determine the expression level of

CLB3 [18].

Microscopy

To test whether the putative cell cycle regulating genes contribute to

sexual reproduction, wild type XL280α, two independent deletion mutants for

PCL2, *PCL6*, *PCL9*, *CKS1*, and *CKS2*, and the conditional expression strain for

CLB3 were spot-inoculated on MS agar medium and incubated for up to three

weeks for unisexual reproduction; and *MAT**a*** and *MAT*α cells of the wild type

XL280 and deletion mutants for *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* were

equally mixed and spot-inoculated on MS agar medium and incubated up to two

weeks for bisexual reproduction. Hyphal growth on the edge of mating patches,

basidia, and spore chains were captured at specified time points using a Nikon

Eclipse E400 microscope equipped with a Nikon DXM1200F camera.

To observe yeast cell morphology, wild-type XL280α, two independent

deletion mutants for *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2*, and the conditional

expression strain for *CLB3* were grown overnight in liquid YPD or YPG medium.

Yeast cells were then fixed in 3.7% formaldehyde, membrane permeabilized in 1%

Triton X100, and strained with 3 µg/ml Hoechst 33342 (Thermo Fisher) and 1

µg/ml calcofluor white (CFW) (Sigma). Stained yeast cells were imaged using a

ZEISS Imager widefield fluorescence microscope and images were processed

using the software FIJI.

Basidiospore and blastospore dissection

Dissections of basidiospore and blastospores were performed using a

fiber optic needle spore dissecting system as previously described [18, 74]. To

obtain meiotic progeny, mating patches were inoculated on MS medium and

incubated in the dark at room temperature for 1-2 weeks to allow basidiospore

chain formation. Basidiospores were transferred onto YPD agar medium (YPG

was used when strains expressing *CLB3* under a galactose-inducible promoter

were involved), and individual basidiospores were separated. To dissect

blastospores, mating patches were prepared similarly but incubated for 3-4

weeks or longer until hyphae grew further away from yeast cells on the agar

surface, then the agar block containing the entire mating patch was excised and

transferred to a YPD or YPG agar medium plate with an equivalent size of agar

removed from the mating patch agar block, and nascent blastospores produced

along the growing hyphae were separated onto fresh YPD or YPG medium.

Flow cytometry

To determine ploidy, actively growing cells on solid agar medium were

collected, fixed in ethanol, stained with propidium iodide, and analyzed by

Fluorescence Activated Cell Sorting (FACS) using a BD FACSCanto II analyzer

as previously described [18, 75]. XL280α and MN142.6 (XL280α/α

$ura5\Delta$::*NAT*/$ura5\Delta$::*NEO*) were used as haploid and diploid controls, respectively.

All FACS data were analyzed in FlowJo.

To determine whether the putative cell cycle regulating genes contributed

to cell cycle progression, wild-type and deletion mutant cells were treated with

hydroxyurea or nocodazole to arrest cells at G1/S and G2/M, respectively [44,

45]. For G1/S arrest, cells were grown in YPD liquid medium overnight, washed

in H_2O, readjusted to OD_{600} = 0.2 in 2 ml fresh YPD liquid medium, regrown at

30°C for 3 hours to reach exponential growth, and then hydroxyurea was added

to the growing culture at a final concentration of 90 mM. Cells were then grown

for an additional 3 hours to arrest cells at G1/S phase. Half of the volume of

arrested cells was collected and fixed in 70% ethanol, and the other half was

fixed after cell cycle release from G1/S arrest by growing in fresh YPD liquid

medium for 90 minutes. For G2/M arrest, cells were prepared in the same

manner and grown in the presence of nocodazole at a final concentration of 100

nM for 5 hours, and arrested cells were fixed in 70% ethanol. Fixed cells were

then stained with propidium iodide and analyzed by FACS following the method

described above. For the conditional expression strain of *CLB3*, the experiment

was repeated in both YPD and YPG liquid media.

To analyze population ploidy dynamics during mating, the wild-type and

deletion mutants were grown on YPD and V8 agar media for 24 hours and cell

ploidy was determined by FACS. The *CLB3* conditional expression strain and

wild type were grown on YPD, YPG, V8, V8 + 2% glucose, and V8 + 2%

galactose agar media for 24 hours.

Generation of the ploidy sensor *NURAT*

The ploidy sensor plasmid pNURAT was generated using the Gibson

cloning method [72]. First, the truncated 5' and 3' *NAT* cassette sequences

sharing 530 bp of the *NAT* CDS were PCR amplified from the plasmid pAI3 using

primer pairs JOHE40975/JOHE41548 and JOHE41547/JOHE40976, the *URA5*

expression cassette was amplified from XL280α genomic DNA using the primer

pair JOHE41549/JOHE41550, and the plasmid backbone was amplified from

plasmid pAI3 using the primer pair JOHE41352/JOHE41353. These PCR

products share 20 bp overlapping sequences and were assembled together to

generate pNURAT where the *URA5* expression cassette was inserted between

the truncated 5' and 3' *NAT* cassette sequences. To insert the ploidy sensor into

the genome, the safe haven locus was identified on Chr 1 in *C. deneoformans*

using the identified safe haven locus in *C. neoformans*, and the plasmid pCF3

(*SH-NEO)* targeting the *C. deneoformans* safe haven locus was generated by

swapping the *C. neoformans* sequence and the *NAT* cassette in pSDMA25 with

the *C. deneoformans* sequences and the *NEO* cassette [50]. The *NURAT*

construct was then PCR amplified from pNURAT and inserted into pCF3 to yield

pCF7 (*SH-NURAT-NEO*) using the Gibson method. pCF7 was linearized with

PacI and introduced into XL280α via biolistic transformation. Insertion of *NURAT-*

NEO at the safe haven locus was verified in the resulting transformant CF1300

by junction PCRs and southern blot.

To generate the deletion construct for the endogenous *URA5* gene, the

HYG cassette was PCR amplified from the plasmid pAG32 and inserted between

5' and 3' sequences flanking the *URA5* CDS using overlap PCR. The deletion

construct was then introduced into XL280**a** via biolistic transformation, and

replacement of the *URA5* CDS by the *HYG* cassette in the resulting transformant

CF1321 was verified by junction and spanning PCRs.

To generate strains carrying both *SH-NURAT-NEO* and *ura5Δ::HYG*,

CF1300 (XL280α *SH-NURAT-NEO*) was crossed with CF1321 (XL280**a**

ura5Δ::HYG), and basidiospores were dissected following methods described

above. Progeny were streaked on YPD+NAT, YPD+NEO, YPD+HYG, and SD-

URA media to check viability phenotypes on each medium. NAT-sensitive and

NEO- and HYG-resistant progeny that could grow on SD-URA medium were

PCR genotyped for the mating-type locus, deletion of the *URA5* gene, and the

presence of the *NURAT-NEO* construct at the safe haven locus. Two progeny

CF1348 (XL280**a** *ura5Δ::HYG SH-NURAT-NEO*) and CF1349 (XL280α

ura5Δ::HYG SH-NURAT-NEO) were verified and selected for further analyses.

To generate diploid strains carrying the ploidy sensor, blastospores were

dissected from CF1349 and two diploid progeny (CF1610 and CF1611 XL280

α/α *ura5Δ::HYG/ura5Δ::HYG SH-NURAT-NEO/SH-NURAT-NEO*) were obtained.

To introduce the ploidy sensor into the deletion mutant strains lacking the

putative cell cycle-regulating genes, *MAT*α *pcl2Δ::NEO*, *MAT*α *pcl6Δ::NEO*,

*MAT*α *pcl9Δ::NEO*, *MAT*α *cks1Δ::NEO*, and *MAT*α *cks2Δ::NEO* were first

crossed with XL280**a** to obtain deletion mutants of each gene in the *MAT***a**

background, and then *MAT***a** *pcl2Δ::NEO* (CF1510), *MAT***a** *pcl6Δ::NEO* (CF1534),

*MAT***a** *pcl9Δ::NEO* (CF1798), *MAT***a** *cks1Δ::NEO* (CF1526), and *MAT***a**

cks2Δ::NEO (CF1516) were crossed with XL280α *ura5Δ::HYG SH-NURAT-NEO*

(CF1349). Basidiospores were dissected from each cross, and NAT-sensitive

and NEO- and HYG-resistant progeny that could grow on SD-URA medium were

PCR genotyped for the mating-type locus, deletion of the putative cell cycle gene,

deletion of the *URA5* gene, and the presence of the *NURAT-NEO* construct at

the safe haven locus. For each cell cycle gene, two *MAT*α progeny with the

desired genotype were obtained except for *PCL6*, where one *MAT*α and one

*MAT***a** progeny were obtained. For *CLB3*, the conditional expression strain

XL280α P_{GAL7}-*CLB3-NEO* (CF1715) was crossed with XL280**a** *ura5Δ::HYG SH-*

NURAT-NEO (CF1348), and basidiospores were dissected on YPG agar medium.

NAT-sensitive, NEO- and HYG-resistant progeny that could grow on SG-URA

medium were genotyped for the mating-type locus, the presence of the

conditional expression construct for *CLB3*, deletion of the *URA5* gene, and the

presence of the *NURAT-NEO* construct at the safe haven locus. One progeny

with the desired genotype was obtained (CF1835).

Detection of ploidy transition events using the ploidy sensor *NURAT*

To test whether the ploidy sensor *NURAT* could detect ploidy transition

events, fluctuation assays were performed using haploid and diploid wild-type

strains carrying the ploidy sensor. Overnight cultures for CF1348, CF1349,

CF1610, and CF1611 were washed once and adjusted to $OD_{600} = 5$ in sterile

H_2O, and 100 µl of cells were spot inoculated on YPD or V8 pH = 7.0 agar

medium and incubated in the dark at room temperature for 36 or 60 hours. After

incubation, cells were collected, suspended in 300 µl sterile H_2O, and serially

diluted by 10-fold seven times. 200 µl of cells from each of the last two serial

dilutions were plated on YPD agar medium to estimate the number of colony

forming units (CFUs), and 200 µl of the undiluted and the 10-fold diluted cell

suspensions were plated on YPD agar medium supplemented with NAT to select

for progeny with the recombined *NURAT* construct. *NURAT* recombination

events per million CFU was used to determine the recombination frequency.

Nourseothricin-resistant (NAT[R]) colonies were then replica plated onto SD-URA

medium to select NAT[R] colonies that were uracil prototrophic (Ura[+]). The

percentages of Ura[+] colonies among NAT[R] colonies were calculated to determine

the double selection efficiency of the *NURAT* and *NAT* genetic constructs. For

each experiment, up to eight colonies were tested for ploidy by FACS analyses.

For each strain, five biological replicates were performed for each condition.

To study whether the identified putative cell cycle genes impact ploidy

transitions during early mating, fluctuation assays were performed for *PCL2*

(CF1779 and CF1780), *PCL6* (CF1773 and CF1774), *PCL9* (CF1806 and

CF1807), *CKS1* (CF1784 and CF1787), and *CKS2* (CF1770 and 1772) genetic

deletion mutants following the method above by incubating cells on V8 pH = 7.0

agar medium for 60 hours. For *CLB3* expression, cells of the wild type (CF1349)

and the conditional expression strain for *CLB3* (CF1835) were incubated on V8,

V8 2% glucose, and V8 2% galactose agar media for 60 hours.

DNA preparation, Nanopore sequencing and assembly of *C. deneoformans*

XL280

The DNA for Nanopore sequencing of the XL280α genome was isolated

as described previously [66]. The DNA was enriched for high molecular weight,

and purified DNA was tested for its quality using NanoDrop. The samples were

sequenced on the MinION system using an R9.4.1 Flow-Cell and the SQK-

LSK109 library preparation kit. Nanopore sequencing was performed at the

default voltage for 48 hours as per the MinION sequencing protocol provided by

the manufacturer. MinION sequencing protocol and setup was controlled using

the MinKNOW software. Base-calling was performed with Guppy v4.2.2 using the

parameters: config dna_r9.4.1_450bps_fast.cf --gscore_filtering, and the sequence

reads obtained were used for genome assembly.

Canu v2.0 [76] was used to assemble the genome of XL280α using reads

that were longer than 10 kb (-minReadLength=10000), which yielded an

estimated genome size of 19.4 Mb. The genome assembly was checked for

integrity by mapping the Canu-corrected reads back to the genome assembly

using minimap2 v2.14 and duplicated small contigs were discarded. These steps

resulted in the generation of a chromosome-level genome assembly consisting of

14 nuclear contigs plus the mitochondrial genome. The genome assembly was

then error-corrected via one round of Nanopolish v0.13.2 (using nanopore reads;

https://github.com/jts/nanopolish) and five rounds of Pilon v1.23 polishing (using

Illumina reads; https://github.com/broadinstitute/pilon) [77]. After the polishing,

telomere sequences were identified in each chromosome, and any additional

sequences flanking the telomere ends were trimmed after validation by Nanopore

read-mapping. The chromosomes were numbered based on their synteny with

the JEC21 genome [78]. Repetitive DNA content, including transposable

elements, was analyzed with RepeatMaster version open-4.0.7 (using RepBase-

20170127 and Dfam Consensus-20170127). Centromeres were predicted by

detection of centromere-associated LTR elements previously reported in *C.*

neoformans (Tcn1 to Tcn6) [79], and further refined by mapping onto the XL280

assembly using the position of each of the centromere flanking genes previously

identified in *C. deneoformans* [80], using BLAST analyses. Both nanopore and

Illumina data for the XL280 genome have been deposited at the NCBI under the

accession number PRJNA720102.

Illumina genome sequencing and read coverage assessment

To understand the nature of double selection of the *NURAT* construct and

NAT marker in non-diploidization events, whole-genome Illumina sequencing

was performed for parental strains CF1300, CF1321, CF1348 and CF1349, and

five NAT^R, Ura^+ progeny (CF1354, CF1355, CF1356, CF1357, and CF1358)

obtained through mitotic passaging on YPD agar medium. Genomic DNA was

extracted following method as previously described [81]. Short-read library

preparation and genome sequencing were conducted at the University of North

Carolina at Chapel Hill's Next Generation Sequencing Facility. Paired 151-base

reads were generated in an Illumina Hiseq2500 system.

To detected aneuploidy events (including segmental or whole-

chromosome aneuploidy), Illumina paired-end reads were filtered with the default

parameters of Trimmomatic v0.36 [82], and subsequently mapped to the *C.*

deneoformans XL280 reference genome using the BWA-MEM short-read aligner

(v0.7.17-r1188) with default settings [83]. Picard tools, integrated in the Genome

Analysis Toolkit (GATK) v4.0.1.2 [84], was used to sort the resulting files by

coordinate, to fix read groups (modules: SORT_ORDER=coordinate;

'AddOrReplaceReadGroups') and to mark duplicates. Aneuploidy events were

inferred from read counts calculated in 1-kb non-overlapping windows across the

genome using the module "count_dna" from the Alfred package (v0.1.7)

(https://github.com/tobiasrausch/alfred). These counts were subjected to median

normalization and log2 transformation and the data was converted into a tiled

data file (.tdf) using "igvtools toTDF" and plotted as a heatmap in IGV viewer

v2.8.0 [85]. Structural events including inversions, duplications, and

translocations/fusions were inferred based on the manual inspection of

discordant read pairs, with LL/RR reads implying inversions and RL reads

implying duplications with respect to the reference. These sets of reads are

represented in IGV with different color codes after grouping and color alignments

by insert size and pair orientation.

Stress response phenotype

To test whether segmental aneuploidy conferred phenotypic variance to

heat stress and the antifungal drug fluconazole, haploid and diploid wild-type

strains and the five mitotically passaged double selection progeny (CF1354,

CF1355, CF1356, CF1357, and Cf1358) were cultured overnight at 30°C in liquid

YPD medium. The cells were subsequently washed once with water, adjusted to

OD600 = 0.8, 10-fold serially diluted, and spot inoculated on YPD and YPD agar

medium supplemented with 8 µg/ml fluconazole. YPD plates were incubated at

30°C and 37°C, and the fluconazole plates were incubated at 30°C for 48 to 72

hours [86].

Pulsed-field gel electrophoresis (PFGE) and chromoblot analysis

PFGE and chromoblot analyses were performed as previously described [87]. CHEF gels were run using 1% agarose gel in 0.5X TBE at 14°C for 96 hours with a ramped switching time from 260 seconds to 560 seconds at a voltage of 3V/cm. To separate smaller chromosomes, CHEF gels were run for 40 hours with a ramped switching time from 50 to 76 seconds at a voltage of 5 V/cm. For chromoblot analyses, probes were designed to hybridize to *URA5*, the *NAT* cassette, and the segmental aneuploid regions on Chrs 2, 6, and 13. Primers used to PCR amplify the probes were listed in Table S6.

Statistical analyses

Graph preparation and statistical analyses were performed using the Graphpad Prism 8 program. Student's t-test was performed for each pairwise comparison. p values lower than 0.05 were considered statistically significant (* indicates $0.01 < p \leq 0.05$, ** indicates $0.001 < p \leq 0.01$, *** indicates $0.0001 < p \leq 0.001$, and **** indicates $p \leq 0.0001$).

Acknowledgements

We thank Shelby Priest for critical reading of the manuscript and thank the

helpful suggestions and discussions from members of the Heitman lab.

Reference

1.   Otto SP. The evolutionary consequences of polyploidy. Cell. 2007;131(3):452-62. Epub 2007/11/06. doi: 10.1016/j.cell.2007.10.022. PubMed PMID: 17981114.

2.   Van de Peer Y, Mizrachi E, Marchal K. The evolutionary significance of polyploidy. Nat Rev Genet. 2017;18(7):411-24. Epub 2017/05/16. doi: 10.1038/nrg.2017.26. PubMed PMID: 28502977.

3.   Todd RT, Forche A, Selmecki A. Ploidy variation in fungi: Polyploidy, aneuploidy, and genome evolution. Microbiol Spectr. 2017;5(4):doi: 10.1128/microbiolspec.FUNK-0051-2016. Epub 2017/07/29. doi: 10.1128/microbiolspec.FUNK-0051-2016. PubMed PMID: 28752816; PubMed Central PMCID: PMCPMC5656283.

4.   Marcet-Houben M, Gabaldon T. Beyond the whole-genome duplication: Phylogenetic evidence for an ancient interspecies hybridization in the baker's yeast lineage. PLoS Biol. 2015;13(8):e1002220. Epub 2015/08/08. doi: 10.1371/journal.pbio.1002220. PubMed PMID: 26252497; PubMed Central PMCID: PMCPMC4529251.

5.   Zhu YO, Sherlock G, Petrov DA. Whole genome analysis of 132 clinical *Saccharomyces cerevisiae* strains reveals extensive ploidy variation. G3 (Bethesda). 2016;6(8):2421-34. Epub 2016/06/19. doi: 10.1534/g3.116.029397. PubMed PMID: 27317778; PubMed Central PMCID: PMCPMC4978896.

6.   Strope PK, Skelly DA, Kozmin SG, Mahadevan G, Stone EA, Magwene PM, et al. The 100-genomes strains, an *S. cerevisiae* resource that illuminates its natural phenotypic and genotypic variation and emergence as an opportunistic pathogen. Genome Res. 2015;25(5):762-74. Epub 2015/04/05. doi: 10.1101/gr.185538.114. PubMed PMID: 25840857; PubMed Central PMCID: PMCPMC4417123.

7.   Selmecki A, Forche A, Berman J. Genomic plasticity of the human fungal pathogen *Candida albicans*. Eukaryot Cell. 2010;9(7):991-1008. Epub 2010/05/25. doi: 10.1128/EC.00060-10. PubMed PMID: 20495058; PubMed Central PMCID: PMCPMC2901674.

8.   Hickman MA, Zeng G, Forche A, Hirakawa MP, Abbey D, Harrison BD, et al. The 'obligate diploid' *Candida albicans* forms mating-competent haploids. Nature. 2013;494(7435):55-9. Epub 2013/02/01. doi: 10.1038/nature11865. PubMed PMID: 23364695; PubMed Central PMCID: PMCPMC3583542.

9.   Bennett RJ, Johnson AD. Completion of a parasexual cycle in *Candida albicans* by induced chromosome loss in tetraploid strains. EMBO J. 2003;22(10):2505-15. Epub 2003/05/14. doi: 10.1093/emboj/cdg235. PubMed PMID: 12743044; PubMed Central PMCID: PMCPMC155993.

10.   Anderson CA, Roberts S, Zhang H, Kelly CM, Kendall A, Lee C, et al. Ploidy variation in multinucleate cells changes under stress. Mol Biol Cell. 2015;26(6):1129-40. Epub 2015/01/30. doi: 10.1091/mbc.E14-09-1375. PubMed PMID: 25631818; PubMed Central PMCID: PMCPMC4357512.

11.   Berman J. Ploidy plasticity: a rapid and reversible strategy for adaptation to stress. FEMS Yeast Res. 2016;16(3). Epub 2016/03/08. doi: 10.1093/femsyr/fow020. PubMed PMID: 26945893.

12.   Zaragoza O, Garcia-Rodas R, Nosanchuk JD, Cuenca-Estrella M, Rodriguez-Tudela JL, Casadevall A. Fungal cell gigantism during mammalian infection. PLoS Pathog.

2010;6(6):e1000945. Epub 2010/06/30. doi: 10.1371/journal.ppat.1000945. PubMed PMID:
20585557; PubMed Central PMCID: PMCPMC2887474.
13.   Okagaki LH, Wang Y, Ballou ER, O'Meara TR, Bahn YS, Alspaugh JA, et al. Cryptococcal
titan cell formation is regulated by G-protein signaling in response to multiple stimuli. Eukaryot
Cell. 2011;10(10):1306-16. Epub 2011/08/09. doi: 10.1128/EC.05179-11. PubMed PMID:
21821718; PubMed Central PMCID: PMCPMC3187071.
14.   Ni M, Feretzaki M, Li W, Floyd-Averette A, Mieczkowski P, Dietrich FS, et al. Unisexual
and heterosexual meiotic reproduction generate aneuploidy and phenotypic diversity *de novo* in
the yeast *Cryptococcus neoformans*. PLoS Biol. 2013;11(9):e1001653. doi:
10.1371/journal.pbio.1001653. PubMed PMID: 24058295; PubMed Central PMCID:
PMC3769227.
15.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, et al. Global
burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect
Dis. 2017;17(8):873-81. Epub 2017/05/10. doi: 10.1016/S1473-3099(17)30243-8. PubMed PMID:
28483415.
16.   Lin X, Hull CM, Heitman J. Sexual reproduction between partners of the same mating
type in *Cryptococcus neoformans*. Nature. 2005;434(7036):1017-21. Epub 2005/04/23. doi:
10.1038/nature03448. PubMed PMID: 15846346.
17.   Ene IV, Bennett RJ. The cryptic sexual strategies of human fungal pathogens. Nat Rev
Microbiol. 2014;12(4):239-51. doi: 10.1038/nrmicro3236. PubMed PMID: 24625892.
18.   Fu C, Heitman J. *PRM1* and *KAR5* function in cell-cell fusion and karyogamy to drive
distinct bisexual and unisexual cycles in the *Cryptococcus* pathogenic species complex. PLoS
Genet. 2017;13(11):e1007113. Epub 2017/11/28. doi: 10.1371/journal.pgen.1007113. PubMed
PMID: 29176784; PubMed Central PMCID: PMCPMC5720818.
19.   Lin X, Patel S, Litvintseva AP, Floyd A, Mitchell TG, Heitman J. Diploids in the
Cryptococcus neoformans serotype A population homozygous for the α mating type originate via
unisexual mating. PLoS Pathog. 2009;5(1):e1000283. doi: 10.1371/journal.ppat.1000283.
PubMed PMID: 19180236; PubMed Central PMCID: PMC2629120.
20.   Okagaki LH, Strain AK, Nielsen JN, Charlier C, Baltes NJ, Chretien F, et al. Cryptococcal
cell morphology affects host cell interactions and pathogenicity. PLoS Pathog.
2010;6(6):e1000953. doi: 10.1371/journal.ppat.1000953. PubMed PMID: 20585559; PubMed
Central PMCID: PMCPMC2887476.
21.   Zaragoza O, Nielsen K. Titan cells in *Cryptococcus neoformans*: cells with a giant impact.
Curr Opin Microbiol. 2013;16(4):409-13. doi: 10.1016/j.mib.2013.03.006. PubMed PMID:
23588027; PubMed Central PMCID: PMC3723695.
22.   Mukaremera L, Lee KK, Wagener J, Wiesner DL, Gow NAR, Nielsen K. Titan cell
production in *Cryptococcus neoformans* reshapes the cell wall and capsule composition during
infection. Cell Surf. 2018;1:15-24. Epub 2018/08/21. doi: 10.1016/j.tcsw.2017.12.001. PubMed
PMID: 30123851; PubMed Central PMCID: PMCPMC6095662.
23.   Crabtree JN, Okagaki LH, Wiesner DL, Strain AK, Nielsen JN, Nielsen K. Titan cell
production enhances the virulence of *Cryptococcus neoformans*. Infect Immun.
2012;80(11):3776-85. Epub 2012/08/15. doi: 10.1128/IAI.00507-12. PubMed PMID: 22890995;
PubMed Central PMCID: PMCPMC3486048.
24.   Okagaki LH, Nielsen K. Titan cells confer protection from phagocytosis in *Cryptococcus
neoformans* infections. Eukaryot Cell. 2012;11(6):820-6. Epub 2012/05/01. doi:
10.1128/EC.00121-12. PubMed PMID: 22544904; PubMed Central PMCID: PMCPMC3370461.
25.   Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, et al. Polyploid titan
cells produce haploid and aneuploid progeny to promote stress adaptation. mBio.

2015;6(5):e01340-15. Epub 2015/10/16. doi: 10.1128/mBio.01340-15. PubMed PMID: 26463162;
PubMed Central PMCID: PMCPMC4620463.

26. Zhao Y, Wang Y, Upadhyay S, Xue C, Lin X. Activation of meiotic genes mediates ploidy
reduction during cryptococcal infection. Curr Biol. 2020;30(8):1387-96 e5. Epub 2020/02/29. doi:
10.1016/j.cub.2020.01.081. PubMed PMID: 32109388.

27. Tian X, He G-J, Hu P, Chen L, Tao C, Cui Y-L, et al. *Cryptococcus neoformans* sexual
reproduction is controlled by a quorum sensing peptide. Nat Microbiol. 2018;3(6):698-707. doi:
10.1038/s41564-018-0160-4.

28. Hommel B, Mukaremera L, Cordero RJB, Coelho C, Desjardins CA, Sturny-Leclere A, et al.
Titan cells formation in *Cryptococcus neoformans* is finely tuned by environmental conditions
and modulated by positive and negative genetic regulators. PLoS Pathog. 2018;14(5):e1006982.
Epub 2018/05/19. doi: 10.1371/journal.ppat.1006982. PubMed PMID: 29775480.

29. Trevijano-Contador N, de Oliveira HC, Garcia-Rodas R, Rossi SA, Llorente I, Zaballos A, et
al. *Cryptococcus neoformans* can form titan-like cells *in vitro* in response to multiple signals.
PLoS Pathog. 2018;14(5):e1007007. Epub 2018/05/19. doi: 10.1371/journal.ppat.1007007.
PubMed PMID: 29775477.

30. Dambuza IM, Drake T, Chapuis A, Zhou X, Correia J, Taylor-Smith L, et al. The
Cryptococcus neoformans Titan cell is an inducible and regulated morphotype underlying
pathogenesis. PLoS Pathog. 2018;14(5):e1006978. Epub 2018/05/19. doi:
10.1371/journal.ppat.1006978. PubMed PMID: 29775474.

31. Zielke N, Edgar BA, DePamphilis ML. Endoreplication. Cold Spring Harb Perspect Biol.
2013;5(1):a012948. Epub 2013/01/04. doi: 10.1101/cshperspect.a012948. PubMed PMID:
23284048; PubMed Central PMCID: PMCPMC3579398.

32. Fox DT, Duronio RJ. Endoreplication and polyploidy: insights into development and
disease. Development. 2013;140(1):3-12. doi: 10.1242/dev.080531.

33. Hayles J, Fisher D, Woollard A, Nurse P. Temporal order of S phase and mitosis in fission
yeast is determined by the state of the p34cdc2-mitotic B cyclin complex. Cell. 1994;78(5):813-
22. Epub 1994/09/09. doi: 10.1016/s0092-8674(94)90542-8. PubMed PMID: 8087848.

34. Lew DJ, Weinert T, Pringle JR. Cell cycle control in *Saccharomyces cerevisiae*. In: Pringle
JR, Broach JR, Jones EW, editors. The Molecular and Cellular Biology of the Yeast *Saccharomyces*:
Cell Cycle and Cell biology. Cold Spring Harbor, NY: Cold Spring Harbor Press; 1997. p. 607-95.

35. Simmons Kovacs LA, Mayhew MB, Orlando DA, Jin Y, Li Q, Huang C, et al. Cyclin-
dependent kinases are regulators and effectors of oscillations driven by a transcription factor
network. Mol Cell. 2012;45(5):669-79. Epub 2012/02/07. doi: 10.1016/j.molcel.2011.12.033.
PubMed PMID: 22306294; PubMed Central PMCID: PMCPMC3578314.

36. Altamirano S, Li Z, Fu MS, Ding M, Fulton SR, Yoder JM, et al. The cyclin Cln1 controls
polyploid titan cell formation following a stress-induced G2 arrest in *Cryptococcus*. mBio.
2021;12(5):e02509-21. doi: doi.org/10.1128/mBio.02509-21.

37. Fu C, Coelho MA, David-Palma M, Priest SJ, Heitman J. Genetic and genomic evolution of
sexual reproduction: echoes from LECA to the fungal kingdom. Curr Opin Genet Dev. 2019;58-
59:70-5. Epub 2019/09/02. doi: 10.1016/j.gde.2019.07.008. PubMed PMID: 31473482; PubMed
Central PMCID: PMCPMC6889014.

38. Stajich JE, Harris T, Brunk BP, Brestelli J, Fischer S, Harb OS, et al. FungiDB: an integrated
functional genomics database for fungi. Nucleic Acids Res. 2012;40(Database issue):D675-81. doi:
10.1093/nar/gkr918. PubMed PMID: 22064857; PubMed Central PMCID: PMCPMC3245123.

39. Liu L, He GJ, Chen L, Zheng J, Chen Y, Shen L, et al. Genetic basis for coordination of
meiosis and sexual structure maturation in *Cryptococcus neoformans*. eLife. 2018;7:e38683.

Epub 2018/10/04. doi: 10.7554/eLife.38683. PubMed PMID: 30281018; PubMed Central PMCID:
PMCPMC6235564.
40.   Measday V, Moore L, Retnakaran R, Lee J, Donoviel M, Neiman AM, et al. A family of
cyclin-like proteins that interact with the Pho85 cyclin-dependent kinase. Mol Cell Biol.
1997;17(3):1212-23. Epub 1997/03/01. doi: 10.1128/mcb.17.3.1212. PubMed PMID: 9032248;
PubMed Central PMCID: PMCPMC231846.
41.   Tang Y, Reed SI. The Cdk-associated protein Cks1 functions both in G1 and G2 in
Saccharomyces cerevisiae. Genes Dev. 1993;7(5):822-32. Epub 1993/05/01. doi:
10.1101/gad.7.5.822. PubMed PMID: 8491379.
42.   Berman J. Morphogenesis and cell cycle progression in *Candida albicans*. Curr Opin
Microbiol. 2006;9(6):595-601. Epub 2006/10/24. doi: 10.1016/j.mib.2006.10.007. PubMed PMID:
17055773; PubMed Central PMCID: PMCPMC3552184.
43.   Senn H, Shapiro RS, Cowen LE. Cdc28 provides a molecular link between Hsp90,
morphogenesis, and cell cycle progression in *Candida albicans*. Mol Biol Cell. 2012;23(2):268-83.
Epub 2011/11/18. doi: 10.1091/mbc.E11-08-0729. PubMed PMID: 22090345; PubMed Central
PMCID: PMCPMC3258172.
44.   Kozubowski L, Yadav V, Chatterjee G, Sridhar S, Yamaguchi M, Kawamoto S, et al.
Ordered kinetochore assembly in the human-pathogenic basidiomycetous yeast *Cryptococcus
neoformans*. mBio. 2013;4(5):e00614-13. Epub 2013/10/03. doi: 10.1128/mBio.00614-13.
PubMed PMID: 24085781; PubMed Central PMCID: PMCPMC3791896.
45.   Janbon G, Maeng S, Yang DH, Ko YJ, Jung KW, Moyrand F, et al. Characterizing the role
of RNA silencing components in *Cryptococcus neoformans*. Fungal Genet Biol. 2010;47(12):1070-
80. Epub 2010/11/12. doi: 10.1016/j.fgb.2010.10.005. PubMed PMID: 21067947; PubMed
Central PMCID: PMCPMC3021383.
46.   Slater ML. Effect of reversible inhibition of deoxyribonucleic acid synthesis on the yeast
cell cycle. J Bacteriol. 1973;113(1):263-70. Epub 1973/01/01. doi: 10.1128/jb.113.1.263-
270.1973. PubMed PMID: 4120066; PubMed Central PMCID: PMCPMC251626.
47.   Jacobs CW, Adams AE, Szaniszlo PJ, Pringle JR. Functions of microtubules in the
Saccharomyces cerevisiae cell cycle. J Cell Biol. 1988;107(4):1409-26. Epub 1988/10/01. doi:
10.1083/jcb.107.4.1409. PubMed PMID: 3049620; PubMed Central PMCID: PMCPMC2115239.
48.   Fu J, Morris IR, Wickes BL. The production of monokaryotic hyphae by *Cryptococcus
neoformans* can be induced by high temperature arrest of the cell cycle and is independent of
same-sex mating. PLoS Pathog. 2013;9(5):e1003335. doi: 10.1371/journal.ppat.1003335.
PubMed PMID: 23658522; PubMed Central PMCID: PMC3642078.
49.   Enloe B, Diamond A, Mitchell AP. A single-transformation gene function test in diploid
Candida albicans. J Bacteriol. 2000;182(20):5730-6. Epub 2000/09/27. doi:
10.1128/jb.182.20.5730-5736.2000. PubMed PMID: 11004171; PubMed Central PMCID:
PMCPMC94694.
50.   Arras SD, Chitty JL, Blake KL, Schulz BL, Fraser JA. A genomic safe haven for mutant
complementation in *Cryptococcus neoformans*. PLoS ONE. 2015;10(4):e0122916. Epub
2015/04/10. doi: 10.1371/journal.pone.0122916. PubMed PMID: 25856300; PubMed Central
PMCID: PMCPMC4391909.
51.   Sionov E, Lee H, Chang YC, Kwon-Chung KJ. *Cryptococcus neoformans* overcomes stress
of azole drugs by formation of disomy in specific multiple chromosomes. PLoS Pathog.
2010;6(4):e1000848. doi: 10.1371/journal.ppat.1000848.
52.   Gusa A, Williams JD, Cho JE, Averette AF, Sun S, Shouse EM, et al. Transposon
mobilization in the human fungal pathogen *Cryptococcus* is mutagenic during infection and
promotes drug resistance *in vitro*. Proc Natl Acad Sci U S A. 2020;117(18):9973-80. Epub

2020/04/19. doi: 10.1073/pnas.2001451117. PubMed PMID: 32303657; PubMed Central PMCID:
PMCPMC7211991.
53.   Huang D, Friesen H, Andrews B. Pho85, a multifunctional cyclin-dependent protein
kinase in budding yeast. Mol Microbiol. 2007;66(2):303-14. Epub 2007/09/14. doi:
10.1111/j.1365-2958.2007.05914.x. PubMed PMID: 17850263.
54.   Hu P, Liu L, Ke W, Tian X, Wang L. A cyclin protein governs the infectious and sexual life
cycles of *Cryptococcus neoformans*. Sci China Life Sci. 2020:doi: 10.1007/s11427-020-1697-3.
Epub 2020/11/10. doi: 10.1007/s11427-020-1697-3. PubMed PMID: 33165808.
55.   Fan Y, Lin X. Multiple applications of a transient CRISPR-Cas9 coupled with
electroporation (TRACE) Ssystem in the *Cryptococcus neoformans* species complex. Genetics.
2018;208(4):1357-72. Epub 2018/02/16. doi: 10.1534/genetics.117.300656. PubMed PMID:
29444806; PubMed Central PMCID: PMCPMC5887135.
56.   Fitch I, Dahmann C, Surana U, Amon A, Nasmyth K, Goetsch L, et al. Characterization of
four B-type cyclin genes of the budding yeast *Saccharomyces cerevisiae*. Mol Biol Cell.
1992;3(7):805-18. Epub 1992/07/01. doi: 10.1091/mbc.3.7.805. PubMed PMID: 1387566;
PubMed Central PMCID: PMCPMC275636.
57.   Kõivomägi M, Valk E, Venta R, Iofik A, Lepiku M, Balog ER, et al. Cascades of multisite
phosphorylation control Sic1 destruction at the onset of S phase. Nature. 2011;480(7375):128-
31. Epub 2011/10/14. doi: 10.1038/nature10560. PubMed PMID: 21993622; PubMed Central
PMCID: PMCPMC3228899.
58.   Richardson HE, Stueland CS, Thomas J, Russell P, Reed SI. Human cDNAs encoding
homologs of the small p34Cdc28/Cdc2-associated protein of *Saccharomyces cerevisiae* and
Schizosaccharomyces pombe. Genes Dev. 1990;4(8):1332-44. Epub 1990/08/01. doi:
10.1101/gad.4.8.1332. PubMed PMID: 2227411.
59.   Kelliher CM, Leman AR, Sierra CS, Haase SB. Investigating conservation of the cell-cycle-
regulated transcriptional program in the fungal pathogen, *Cryptococcus neoformans*. PLoS Genet.
2016;12(12):e1006453. Epub 2016/12/06. doi: 10.1371/journal.pgen.1006453. PubMed PMID:
27918582; PubMed Central PMCID: PMCPMC5137879.
60.   Vanneste E, Voet T, Le Caignec C, Ampe M, Konings P, Melotte C, et al. Chromosome
instability is common in human cleavage-stage embryos. Nat Med. 2009;15(5):577-83. Epub
2009/04/28. doi: 10.1038/nm.1924. PubMed PMID: 19396175.
61.   Selmecki A, Forche A, Berman J. Aneuploidy and isochromosome formation in drug-
resistant *Candida albicans*. Science. 2006;313(5785):367-70. Epub 2006/07/22. doi:
10.1126/science.1128242. PubMed PMID: 16857942; PubMed Central PMCID:
PMCPMC1717021.
62.   Selmecki A, Gerami-Nejad M, Paulson C, Forche A, Berman J. An isochromosome confers
drug resistance *in vivo* by amplification of two genes, *ERG11* and *TAC1*. Mol Microbiol.
2008;68(3):624-41. Epub 2008/03/28. doi: 10.1111/j.1365-2958.2008.06176.x. PubMed PMID:
18363649.
63.   Todd RT, Wikoff TD, Forche A, Selmecki A. Genome plasticity in Candida albicans is
driven by long repeat sequences. eLife. 2019;8:e45954. Epub 2019/06/08. doi:
10.7554/eLife.45954. PubMed PMID: 31172944; PubMed Central PMCID: PMCPMC6591007.
64.   Todd RT, Selmecki A. Expandable and reversible copy number amplification drives rapid
adaptation to antifungal drugs. eLife. 2020;9:e58349. Epub 2020/07/21. doi:
10.7554/eLife.58349. PubMed PMID: 32687060; PubMed Central PMCID: PMCPMC7371428.
65.   Schotanus K, Heitman J. Centromere deletion in *Cryptococcus deuterogattii* leads to
neocentromere formation and chromosome fusions. eLife. 2020;9:e56026. Epub 2020/04/21.
doi: 10.7554/eLife.56026. PubMed PMID: 32310085; PubMed Central PMCID: PMCPMC7188483.

66.   Yadav V, Sun S, Coelho MA, Heitman J. Centromere scission drives chromosome
shuffling and reproductive isolation. Proc Natl Acad Sci U S A. 2020;117(14):7917-28. Epub
2020/03/21. doi: 10.1073/pnas.1918659117. PubMed PMID: 32193338; PubMed Central PMCID:
PMCPMC7149388.
67.   Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, et al. Effects of
aneuploidy on cellular physiology and cell division in haploid yeast. Science.
2007;317(5840):916-24. doi: 10.1126/science.1142210.
68.   Williams BR, Prabhu VR, Hunter KE, Glazier CM, Whittaker CA, Housman DE, et al.
Aneuploidy affects proliferation and spontaneous immortalization in mammalian cells. Science.
2008;322(5902):703-9. Epub 2008/11/01. doi: 10.1126/science.1160058. PubMed PMID:
18974345; PubMed Central PMCID: PMCPMC2701511.
69.   Zhai B, Zhu P, Foyle D, Upadhyay S, Idnurm A, Lin X. Congenic strains of the filamentous
form of *Cryptococcus neoformans* for studies of fungal morphogenesis and virulence. Infect
Immun. 2013;81(7):2626-37. doi: 10.1128/IAI.00259-13. PubMed PMID: 23670559; PubMed
Central PMCID: PMCPMC3697605.
70.   Davidson RC, Cruz MC, Sia RA, Allen B, Alspaugh JA, Heitman J. Gene disruption by
biolistic transformation in serotype D strains of *Cryptococcus neoformans*. Fungal Genet Biol.
2000;29(1):38-48. doi: 10.1006/fgbi.1999.1180. PubMed PMID: 10779398.
71.   Ruff JA, Lodge JK, Baker LG. Three galactose inducible promoters for use in *C.*
neoformans var. *grubii*. Fungal Genet Biol. 2009;46(1):9-16. Epub 2008/10/28. doi:
10.1016/j.fgb.2008.10.003. PubMed PMID: 18952189; PubMed Central PMCID:
PMCPMC2654232.
72.   Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. Enzymatic
assembly of DNA molecules up to several hundred kilobases. Nat Methods. 2009;6(5):343-5. doi:
10.1038/nmeth.1318. PubMed PMID: 19363495.
73.   Fang Y, Cui L, Gu B, Arredondo F, Tyler BM. Efficient genome editing in the oomycete
Phytophthora sojae using CRISPR/Cas9. Curr Protoc Microbiol. 2017;44:21A 1 1-A 1 6. Epub
2017/02/07. doi: 10.1002/cpmc.25. PubMed PMID: 28166383.
74.   Idnurm A. A tetrad analysis of the basidiomycete fungus *Cryptococcus neoformans*.
Genetics. 2010;185:153-63. Epub 2010/02/17. doi: genetics.109.113027 [pii]

10.1534/genetics.109.113027. PubMed PMID: 20157004.
75.   Tanaka R, Taguchi H, Takeo K, Miyaji M, Nishimura K. Determination of ploidy in
Cryptococcus neoformans by flow cytometry. J Med Vet Mycol. 1996;34(5):299-301. PubMed
PMID: 8912162.
76.   Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH, Phillippy AM. Canu: scalable and
accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Res.
2017;27(5):722-36. Epub 2017/03/17. doi: 10.1101/gr.215087.116. PubMed PMID: 28298431;
PubMed Central PMCID: PMCPMC5411767.
77.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an
integrated tool for comprehensive microbial variant detection and genome assembly
improvement. PLoS ONE. 2014;9(11):e112963. Epub 2014/11/20. doi:
10.1371/journal.pone.0112963. PubMed PMID: 25409509; PubMed Central PMCID:
PMCPMC4237348.
78.   Loftus BJ, Fung E, Roncaglia P, Rowley D, Amedeo P, Bruno D, et al. The genome of the
basidiomycetous yeast and human pathogen *Cryptococcus neoformans*. Science.
2005;307(5713):1321-4. Epub 2005/01/18. doi: 10.1126/science.1103773. PubMed PMID:
15653466; PubMed Central PMCID: PMCPMC3520129.

79.   Janbon G, Ormerod KL, Paulet D, Byrnes EJ, 3rd, Yadav V, Chatterjee G, et al. Analysis of
the genome and transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex RNA
expression and microevolution leading to virulence attenuation. PLoS Genet.
2014;10(4):e1004261. doi: 10.1371/journal.pgen.1004261. PubMed PMID: 24743168; PubMed
Central PMCID: PMCPMC3990503.
80.   Yadav V, Sun S, Billmyre RB, Thimmappa BC, Shea T, Lintner R, et al. RNAi is a critical
determinant of centromere evolution in closely related fungi. Proc Natl Acad Sci U S A.
2018;115(12):3108-13. Epub 2018/03/07. doi: 10.1073/pnas.1713725115. PubMed PMID:
29507212; PubMed Central PMCID: PMCPMC5866544.
81.   Pitkin JW, Panaccione DG, Walton JD. A putative cyclic peptide efflux pump encoded by
the *TOXA* gene of the plant-pathogenic fungus *Cochliobolus carbonum*. Microbiology.
1996;142:1557-65. doi: Doi 10.1099/13500872-142-6-1557. PubMed PMID:
WOS:A1996UR35800028.
82.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30(15):2114-20. Epub 2014/04/04. doi:
10.1093/bioinformatics/btu170. PubMed PMID: 24695404; PubMed Central PMCID:
PMCPMC4103590.
83.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform.
Bioinformatics. 2009;25(14):1754-60. Epub 2009/05/20. doi: 10.1093/bioinformatics/btp324.
PubMed PMID: 19451168; PubMed Central PMCID: PMCPMC2705234.
84.   DePristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, et al. A framework for
variation discovery and genotyping using next-generation DNA sequencing data. Nat Genet.
2011;43(5):491-8. Epub 2011/04/12. doi: 10.1038/ng.806. PubMed PMID: 21478889; PubMed
Central PMCID: PMCPMC3083463.
85.   Robinson JT, Thorvaldsdottir H, Winckler W, Guttman M, Lander ES, Getz G, et al.
Integrative genomics viewer. Nat Biotechnol. 2011;29(1):24-6. Epub 2011/01/12. doi:
10.1038/nbt.1754. PubMed PMID: 21221095; PubMed Central PMCID: PMCPMC3346182.
86.   Fu C, Donadio N, Cardenas ME, Heitman J. Dissecting the roles of the calcineurin
pathway in unisexual reproduction, stress responses, and virulence in *Cryptococcus*
deneoformans. Genetics. 2018;208(2):639-53. Epub 2017/12/14. doi:
10.1534/genetics.117.300422. PubMed PMID: 29233811; PubMed Central PMCID:
PMCPMC5788528.
87.   Findley K, Sun S, Fraser JA, Hsueh YP, Averette AF, Li WJ, et al. Discovery of a modified
tetrapolar sexual cycle in *Cryptococcus amylolentus* and the evolution of *MAT* in the
Cryptococcus species complex. PLoS Genet. 2012;8(2):e1002528. doi: ARTN e1002528

10.1371/journal.pgen.1002528. PubMed PMID: WOS:000300725500044.

1341

1342

1343 **Figure legends**

1344

Figure 1. Identification of cell cycle regulating genes involved in unisexual

reproduction. (A) Differential expression patterns of cell cycle regulating genes

in wild type XL280α cells upon incubation for 36 hours on mating-inducing V8

agar medium versus nutrient rich YPD agar medium were examined by qRT-

PCR. The error bars represent the standard deviation of the mean for three

biological replicates. Red and green colors indicate genes that are significantly

down- and up- regulated during unisexual reproduction compared to the control

gene *KAR5*, respectively. * indicates $0.01 < p \leq 0.05$ and ** indicates $0.001 < p \leq$

0.01. (B) Wild type XL280α, conditional expression mutant of *CLB3*, and deletion

mutants of individual cell cycle regulating genes were grown on MS medium to

assess unisexual hyphal growth and spore formation. Hyphal growth on the edge

of each colony was imaged after 7 days and the scale bar represents 500 µm.

Spore formation was imaged after three weeks and the scale bar represents 10

µm. (C) Wild type and the conditional expression strain for *CLB3* were grown on

YPD, YPG, V8, V8 glucose, and V8 galactose for 24 hours. Ploidy for the cell

populations were determined by FACS.

1361

Figure 2. Cell cycle regulating genes contribute to blastospore

diploidization during unisexual reproduction. Ploidy of single colonies derived

from microscopically dissected blastospores were determined by FACS.

Schematic diagram showing basidiospores and blastospores and representative

gating strategy for single cells were provided at top left. Representative FACS

results for haploid (red), diploid (blue), aneuploid (purple), or mixed

haploid/diploid (yellow) were overlay-plotted with half offset.

1369

Figure 3. Ploidy sensor reports ploidy transition events during both mitotic

and meiotic growth. (A) Schematic diagram shows the mechanism that the

ploidy sensor construct *NURAT* detects ploidy transition depends on homologous

recombination in one *NURAT* allele after the ploidy transition events. (B) A

haploid and two diploid *ura5Δ* strains harboring *NURAT* constructs were

incubated on YPD and V8 agar medium for 60 hours. The number of *NURAT*

recombination events per million CFU was plotted to compare the recombination

frequencies. NAT^R colonies were replica-plated onto SD-URA agar medium to

obtain cells that contain both *NURAT* and *NAT* constructs. The percentile of

NURAT presence among NAT^R colonies were plotted in the bar graph. Mean

values of five independent experiments were plotted for the *NURAT*

recombination frequencies and the Ura$^+$ percentiles among NAT^R colonies; error

bars represent standard deviations. * indicates $0.01 < p \leq 0.05$ and ** indicates

$0.001 < p \leq 0.01$. Individual *NAT*/*NURAT* colonies were tested for ploidy by

FACS and plotted in the violin plot.

1385

Figure 4. Ploidy sensor detects auto-diploidization and segmental

aneuploid formation. (A) A *MATα NURAT*, *MATa ura5Δ*, *MATa ura5Δ NURAT*,

MATα ura5Δ NURAT, and four *MATa ura5Δ NAT/NURAT* and one *MATα ura5Δ*

NAT/NURAT colonies derived through mitotic passaging were subjected to

Illumina whole-genome sequencing. Read depth across all 14 chromosomes was

plotted for each strain. Ploidy was determined by FACS and is listed at the end of

each sequencing result. Centromeres and mating-type loci are indicated by grey

and green bars, respectively. (B) Read depth for Chr 1 and regions containing

the *AFR1* gene, which encodes a drug efflux pump, and the safe haven locus,

where NURAT is inserted. Both loci are present in duplicated chromosomal

segments among all *NAT/NURAT* progeny. Transposable elements were

highlighted and labeled in green below the Chr 1 read depth plot. (C) Human

host temperature tolerance and fluconazole resistance phenotypes were

examined for these five *NAT/NURAT* strains with haploid and diploid wild-type

controls. Cells were 10-fold serial-diluted and spotted on YPD and YPD

supplemented with 8 μg/ml fluconazole, and then incubated at either 30°C or

37°C for two days.

1403

Figure 5. Cell cycle regulating genes contribute to segmental aneuploid

formation. (A) Deletion of *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* reduced

NURAT recombination frequency; deletion of *CKS1* increased the percentile of

NURAT presence among NAT[R] colonies; deletion of *PCL2*, *PCL6*, and *CKS2*

blocked segmental aneuploidy formation. Student's t-test with Bonferroni

correction for 10 repeated tests was performed to compare each mutant with the

wild type. *p* value lower than 0.005 (*) was considered statistically significant. (B)

Suppressed expression of *CLB3* increased *NURAT* recombination frequency and

the percentile of *NURAT* presence among NAT^R colonies, while increased

expression of *CLB3* reduced *NURAT* recombination frequencies but increased

the percentile of *NURAT* presence among NAT^R colonies. Neither down

regulation or up regulation of *CLB3* blocked segmental aneuploid formation.

Mean values of five independent experiments were plotted for the *NURAT*

recombination frequencies and the Ura$^+$ percentiles among NAT^R colonies; error

bars represent standard deviations. Student's t-test with Bonferroni correction for

9 repeated tests was performed for each pairwise comparison. *p* value lower

than 0.0056 (*) was considered statistically significant.

1421

Supporting information

S1 Fig. Expression of putative cyclin dependent kinases and expression profiles of the putative cell cycle regulators. (A) Differential expression patterns of genes encoding putative cyclin dependent kinases in wild-type XL280α cells incubated for 36 hours on mating-inducing V8 agar medium versus nutrient-rich YPD agar medium were examined by qRT-PCR. (B) Relative expression levels for the six differentially expressed cell cycle regulators were extrapolated from a time-course transcriptional profiling study of the wild-type strain XL280α during unisexual reproduction [39]. Expression levels on YPD medium after incubation for 12h and V8 agar medium after incubation for 12h, 24h, and 48h were plotted for these putative cell cycle genes. A black dashed line was drawn to indicate the time point assayed for these genes in this study.

S2 Fig. Expression of *CLB3* under the galactose-inducible promoter P_{GAL7} and bisexual mating phenotypes of the cell cycle regulating gene deletion mutants. (A) *CLB3* was expressed under the control of galactose-inducible promoter P_{GAL7}. Compared to the expression level of the wild type on YPD agar medium, P_{GAL7}-*CLB3* was upregulated 5.2- and 5.8-fold on YPG and V8 galactose agar media and downregulated 38.8-, 15.4-, 10.9-, and 9.9-fold on YPD, MS, V8, and V8 glucose agar media, respectively. The error bars represent the standard deviation of the mean for three biological replicates. (B) *MAT**a*** and *MAT*α cells of wild type XL280 and deletion mutants for *PCL2*, *PCL6*, *PCL9*,

CKS1, and *CKS2* were equally mixed and inoculated on MS medium to assess bisexual hyphal growth and spore formation. Hyphal growth on the edge of each colony was imaged after three days and the scale bar represents 200 μm. Spore formation was imaged after eleven days and the scale bar represents 50 μm.

S3 Fig. Deletion of *CKS1* and repressed expression of *CLB3* result in pseudo-hyphal growth. Wild type, deletion mutants of *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* were grown in liquid YPD overnight, and the conditional expression strain for *CLB3* was grown in liquid YPD and YPG medium overnight. Cells were stained with Calcofluor white and DAPI. The scale bar represents 10 μm

S4 Fig. Deletion of *CKS1* and reduced expression of *CLB3* arrest cells at G2 phase. Overnight culture in YPD for the wild type and deletion mutant strains of *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2*, and overnight culture in YPD for the conditional expression strain for *CLB3* were arrested by hydroxyurea and nocodazole to assess whether these genes regulate cell cycle progression. Cells were arrested in G1 by hydroxyurea and released to S/G2 after removal of hydroxyurea. Nocodazole arrested cells at S/G2 phase. Ploidy for the cell populations were determined by FACS.

S5 Fig. Population ploidy distribution is similar between mating-inducing and -suppressing conditions. Wild type and deletion mutants of *PCL2*, *PCL6*, *PCL9*, *CKS1*, and *CKS2* were grown on YPD and V8 agar media for 24 hours.

S6 Fig. Schematic diagram for the generation of strains carrying the *NURAT* ploidy sensor. A *MATα NURAT* strain (CF1300 XL280α *SH-NURAT-NEO*) was crossed with a *MAT**a** ura5Δ* strain (CF1321 XL280**a** *ura5Δ::HYG*) to generate *MAT**a*** and *MATα ura5Δ NURAT* strains (CF1348 and CF1349). The *MAT**a** ura5Δ NURAT/NAT*-1, -2, -3, and -4 strains (CF1354, CF1355, CF1356, and CF1347) and the *MATα ura5Δ NURAT/NAT*-1 strain (CF1358) were generated through mitotic passages of CF1348 and CF1349, respectively. All above nine strains were subjected to Illumina whole-genome sequencing. Diploid *MATα ura5Δ NURAT/NURAT* strains (CF1610 and CF1611 α/α *ura5Δ/ura5Δ NURAT/NURAT*) were generated by dissecting blastospores from CF1349. Ploidy of all strains were confirmed by FACS.

S7 Fig. Schematic diagram for the ploidy transition detection assays using the *NURAT* ploidy sensor. Overnight cultures of strains carrying the *NURAT* construct were washed and inoculated on V8 or YPD medium for the designated time period. Cells were then plated on YPD medium supplemented with nourseothricin to select for cells with a recombined, functional *NAT* construct. Colonies derived from these cells were replica plated onto SD-URA medium to

screen for NAT^R cells that retained an intact *NURAT* construct. NAT^R, Ura$^+$ colonies were then tested for ploidy by FACS.

S8 Fig. Ploidy sensor reports ploidy transition events during both mitotic and meiotic growth. Frequencies of *NURAT* recombination in haploid strains *MAT**a** NURAT* (CF1348, only overnight culture was tested) and *MATα NURAT* (CF1349), and diploid strains *MAT α NURAT/NURAT*-1 and *MATα NURAT/NURAT*-2 (CF1610 and CF1611) grown (A) as overnight cultures in liquid YPD medium and (B) on V8 or YPD agar medium for 36 hours (scatter dot plots). NAT^R colonies were replica-plated onto SD-URA medium to obtain NAT^R, Ura$^+$ colonies (bar graphs), and ploidy for these colonies was assessed by FACS (violin plots). Mean values of five independent experiments were plotted for the *NURAT* recombination frequencies and the Ura$^+$ percentiles among NAT^R colonies; error bars represent standard deviations. Student's T-test was performed for each pairwise comparison. ** indicates $0.001 < p \leq 0.01$.

S9 Fig. Flanking sequences of segmented regions show distinct modes of segmental aneuploid formation. For each *NURAT/NAT* progeny, sequencing reads at the borders of segmentally duplicated regions were analyzed. Blue, red, and green bubbles indicate forward and reverse reads that were aligned to two different chromosomal positions. Sequence alignments of these reads were shown in the panels on the right of the chromosome diagrams. Chimeric reads

aligning to two different chromosomal positions were highlighted in connected boxes. Sequencing reads aligned to segmentally duplicated regions from three chromosomes were identified in the *MAT***a** *ura5Δ NURAT*/*NAT*-1, -2, and -4 strains, suggesting fusion of these regions. T1 and T3/CNIRT4 transposable element movements were detected flanking some of the regions in the *MAT***a** *ura5Δ NURAT*/*NAT*-1, -2, and -3 strains and the *MAT*α *ura5Δ NURAT*/*NAT*-1 strain. In the *MAT*α *ura5Δ NURAT*/*NAT*-1 strain, tandem duplication and inversion events were detected in the segmentally duplicated regions.

S10 Fig. Karyotypic changes are associated with segmental aneuploid formation. CHEF gel electrophoresis separation of chromosomes was performed under different conditions to separate larger or smaller chromosomes. Karyotypic changes (highlighted in green and red arrows) were observed for strains with segmental aneuploidy (*MAT***a** *ura5Δ NAT*/*NURAT*-1, -2, -3, -4 and *MAT*α *ura5Δ NAT*/*NURAT*-1) compared with wild type and parental strains. Chromoblot analyses with probes recognizing (A) *URA5* and *NAT*, and segmental aneuploid portions of (B) Chrs 2 and 6 and (C) Chr 13 confirmed the karyotypic changes. Strains are highlighted in red when the probed sequences are within segmental aneuploid regions.

S11 Fig. *PCL9* and *CLB3* contribute to aneuploidy formation. Individual colonies with ploidies above 1.6 were identified as outliers and removed from

Figure 5 and data were replotted for (A) *pcl9Δ* and *cks1Δ* mutants and (B) P_{UGE2}-*CLB3*. Student's t-tests with Bonferroni correction for 4 and 9 repeated tests were performed for each pairwise comparison for panel A and panel B, respectively. *p* value lower than 0.0125 (A) or 0.0056 (B) was considered statistically significant (*).

A

Putative cyclin genes

Relative expression under mating inducing versus mating suppressing conditions

KAR5, PCL2 CNA01720, CNA04230, SSN8 CNA04250, PCL9 CNE04400, CLB3 CNG01990, PCL1 CNJ00160, CLB2 CNK01480, PCL6 CNM00990, CLB4 CNI01430, CCL1

Putative CDK regulators

CNE00250, PHO81 CNE02070, ALY2 CNE04100, Putative CKR CNF01190, CKS2 CNF02160, CKS1

B

WT, P_{GAL7}-CLB3 (off), pcl2Δ-1, pcl2Δ-2, pcl6Δ-1, pcl6Δ-2

pcl9Δ-1, pcl9Δ-2, cks1Δ-1, cks1Δ-2, cks2Δ-1, cks2Δ-2

WT, P_{GAL7}-CLB3 (off), pcl2Δ-1, pcl2Δ-2, pcl6Δ-1, pcl6Δ-2, pcl9Δ-1, pcl9Δ-2, cks1Δ-1, cks1Δ-2, cks2Δ-1, cks2Δ-2

C

WT, P_{GAL7}-CLB3

Count

1N, 2N, 4N

YPD, YPG, V8, V8 + Glucose, V8 + Galactose