SARS-CoV-2 Infects Syncytiotrophoblast and Activates Inflammatory Responses in the Placenta

Lissenya B. Argueta[1,*], Lauretta A. Lacko[2,*], Yaron Bram[3,*], Takuya Tada[4], Lucia Carrau[5], Tuo Zhang[6], Skyler Uhl[5], Brienne C. Lubor[1], Vasuretha Chandar[3], Cristianel Gil[2], Wei Zhang[6], Brittany Dodson[7], Jeroen Bastiaans[1], Malavika Prabhu[7], Christine M. Salvatore[9], Yawei J. Yang [7,8], Rebecca N. Baergen[8], Benjamin R. tenOever[5], Nathaniel R. Landau[4], Shuibing Chen[2,#], Robert E. Schwartz[3,#], Heidi Stuhlmann[1,10,#]

Department of Cell and Developmental Biology, Weill Cornell Medicine, 1300 York Avenue, New York 10065, NY, USA[1]
Department of Surgery, Weill Cornell Medicine, New York, NY, USA[2]
Division of Gastroenterology and Hepatology, Department of Medicine, Weill Cornell Medicine, New York, NY, USA[3]
Department of Microbiology, NYU Grossman School of Medicine, New York, NY, USA[4]
Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY, USA[5]
Genomics Resources Facility, Weill Cornell Medicine, New York, NY, USA[6]
Department of Obstetrics and Gynecology, Weill Cornell Medicine, New York, NY, USA[7]
Department of Pathology and Laboratory Medicine, Weill Cornell Medicine, New York, NY, USA[8]
Department of Pediatrics, Division of Pediatric Infectious Diseases, Weill Cornell Medicine, New York, NY, USA[9]
Department of Pediatrics, Weill Cornell Medicine, New York, NY, USA[10]

*These authors contributed equally

#Corresponding Authors:

Heidi Stuhlmann PhD (lead contact) hes2011@med.cornell.edu

Robert Schwartz MD-PhD res2025@med.cornell.edu

Shuibing Chen PhD shc2034@med.cornell.edu

Abstract

SARS-CoV-2 infection during pregnancy leads to an increased risk of adverse pregnancy outcomes. Although the placenta itself can be a target of virus infection, most neonates are virus free and are born healthy or recover quickly. Here, we investigated the impact of SARS-CoV-2 infection on the placenta from a cohort of women who were infected late during pregnancy and had tested nasal swab positive for SARS-CoV-2 by qRT-PCR at delivery. SARS-CoV-2 genomic and subgenomic RNA was detected in 23 out of 54 placentas. Two placentas with high virus content were obtained from mothers who presented with severe COVID-19 and whose pregnancies resulted in adverse outcomes for the fetuses, including intrauterine fetal demise and a preterm delivered baby still in newborn intensive care. Examination of the placental samples with high virus content showed efficient SARS-CoV-2 infection, using RNA in situ hybridization to detect genomic and replicating viral RNA, and immunohistochemistry to detect SARS-CoV-2 nucleocapsid protein. Infection was restricted to syncytiotrophoblast cells that envelope the fetal chorionic villi and are in direct contact with maternal blood. The infected placentas displayed massive infiltration of maternal immune cells including macrophages into intervillous spaces, potentially contributing to inflammation of the tissue. *Ex vivo* infection of placental cultures with SARS-CoV-2 or with SARS-CoV-2 spike (S) protein pseudotyped lentivirus targeted mostly syncytiotrophoblast and in rare events endothelial cells. Infection was reduced by using blocking antibodies against ACE2 and against Neuropilin 1, suggesting that SARS-CoV-2 may utilize alternative receptors for entry into placental cells.

Introduction

The global pandemic resulting from the novel coronavirus, Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has already taken a devastating toll, with over 175 million total cases and more than 3.8 million deaths worldwide. SARS-CoV-2 which causes Coronavirus Disease 2019 (COVID-19) has significant clinical variability. In severe cases SARS-CoV-2 causes a respiratory illness, whose defining features are an imbalanced inflammatory host response, reduced innate antiviral defenses and an inflammatory "cytokine storm", endothelial damage, coagulopathies and thrombosis in several tissues from infected patients (Blanco-Melo et al. 2020).

To date, our understanding of how SARS-CoV-2 infection impacts pregnancy, including the health of COVID-19 positive mothers and their babies, remains incomplete. Pregnant women with symptomatic COVID-19 infections are more likely to be admitted to the intensive care unit (ICU), and have statistically higher maternal death rates when compared to non-pregnant infected women (Zambrano et al. 2020). While preterm deliveries occur more often in women with suspected or confirmed SARS-CoV-2 infection, no increase in stillbirth or early neonatal death was found (Mullins et al. 2021). Prospective and retrospective studies showed that pregnant women infected with SARS-CoV-2 are at increased risk of adverse events, including higher rates of cesarean section and increased post-partum complications (Woodworth et al. 2020; Prabhu et al. 2020; Marín Gabriel et al. 2020). While vertical transmission of SARS-CoV-2 from mother to fetus has been reported in a few cases (Hecht et al. 2020a; Vivanti et al. 2020; Taglauer et al. 2020; Facchetti et al. 2020; Woodworth et al. 2020; Hecht et al. 2020b; Alamar et al. 2020), most studies did not detect viral transmission (Penfield et al.

2020; Baergen and Heller 2020; Prabhu et al. 2020; Salvatore et al. 2020; Edlow et al.

2020; Schwartz 2020; Della Gatta et al. 2020; Kimberlin and Stagno 2020).

Several studies have detected SARS-CoV-2 infection of the placentas from women

who tested positive for the virus at, or prior to, delivery. In some cases, the placenta

displayed signs of inflammation. These placentas were found to have increased

vascular malperfusion indicative of thrombi in fetal vessels (Baergen and Heller 2020;

Vivanti et al. 2020; Prabhu et al. 2020; Shanes et al. 2020) and infiltration of maternal

immune cells (Hosier et al. 2020; Facchetti et al. 2020; Debelenko et al. 2021; Garrido-

Pontnou et al. 2021; Lu-Culligan et al. 2021; Morotti et al. 2021; Schwartz et al. 2021).

Whether inflammation results from virus infection of the mother or direct infection of the

placenta remains unresolved, as this may depend on the gestational age of the fetus.

Virus infection is known to impair placental function. Virus-associated inflammation

during pregnancy can result in chronic cardiovascular disease, diabetes and obesity

later in life (Burton, Fowden, and Thornburg 2016). Little is known however about the

effect of SARS-CoV-2 on placental function.

SARS-CoV-2 utilizes ACE2 (Angiotensin-converting enzyme 2) as the primary

receptor (Hoffmann et al. 2020), and Neuropilin-1 (NRP1) as a coreceptor (Cantuti-

Castelvetri et al. 2020; Daly et al. 2020), in concert with the two proteinases TMPRSS2

(Transmembrane protease serine 2) (Hoffmann et al. 2020) and CTSL (Ou et al. 2020)

and the pro-protein convertase furin (Shang et al. 2020), amongst others (Wei et al.

2021; Daniloski et al. 2021; Wang et al. 2021; Schneider et al. 2021) for cell entry. All of

the viral entry receptors are expressed at significant levels in first and second trimester

placentas. However, at term they are expressed at lower levels at the maternal-fetal

interface, including the placenta and the chorioamniotic membranes (Pique-Regi et al.

2020; Li et al. 2020; Singh, Bansal, and Feschotte 2020; Taglauer et al. 2020; Lu-

Culligan et al. 2021; Baston-Buest et al. 2011). Whether alternative entry mechanisms

are exploited by SARS-CoV-2 in the placenta is not known.

In this study we were interested to understand the impact of SARS-CoV-2 infection

late in pregnancy on placental function. Using a cohort of 54 women who tested positive

for SARS-CoV-2 at the time of delivery, we report on placental infections, placental

pathologies and *in vivo* inflammatory responses to infection. Furthermore, we present *in*

situ studies of infected placentas as well as *ex vivo* placental explant cultures that

investigate susceptibility of placental cells to SARS-CoV-2 infection.

Results

Clinical presentations of SARS-CoV-2 positive mothers, fetal outcomes and placental pathologies

All placental samples in the present study were provided by the Department of Pathology and Laboratory Medicine at Weill Cornell Medicine. A cohort of 54 women who were identified as positive for SARS-CoV-2 by RT-PCR from nasopharyngeal swabs at the time of admission for delivery at NY Presbyterian Hospital-Weill Cornell was included (P1–P54). As controls, a cohort of 5 women who tested negative for SARS-CoV-2 (C1-C5), and a cohort of 5 SARS-CoV-2 negative women who presented various placental inflammatory pathologies (I1-I5) were also included in the study (Table 1).

The pregnant women ranged in age from 16 to 51 years, with a majority in their 20's and 30's. Two pregnancies resulted in intrauterine fetal demise (IUFD) (P2, P6), and one fetus, delivered preterm at 25 weeks of gestation, was admitted to the neonatal intensive care unit (NICU) where the infant has remained for 4 months (P1). All neonates were tested by nasopharyngeal swabs for SARS-CoV-2 at 24 hours, and none were positive. Among the placentas delivered from mothers who tested positive for SARS-CoV-2, 31% (17 cases) presented fetal vascular malperfusion (FVM), 19% (10 cases) displayed maternal vascular malperfusion (MVM), and 7% (4 cases) overlapped for both placental pathologies. None of the healthy control placentas from SARS-CoV-2 negative mothers displayed FVM or MVM (Table 1).

RNA was isolated from all placental samples and subjected qRT-PCR to determine the presence of genomic and replicating SARS-CoV-2 RNA. 22 out of 54 placentas from

SARS-CoV-2 positive mothers showed presence of viral RNA (42%), 2 of those were

highly positive (4%), 10 positive (19%) and 10 were borderline positive (19%) (Table 1

and Figure 1A). Presence of SARS-CoV-2 in the placenta did not correlate with

observed FVM: Of the 22 positive placentas, 10 displayed FVM while 12 were without

FVM (Table 1).

Strikingly, the three pregnancies from SARS-CoV-2 positive mothers that resulted in

IUFD or admission of the neonate to the NICU delivered placentas that were highly

positive (P1 and P2) or positive (P6) for SARS-CoV-2 (Table 1, grey shaded rows).

Placental syncytiotrophoblast are the primary target for SARS-CoV-2 infection of

pregnant females at term

To determine whether the placenta itself was infected by SARS-CoV-2, qRT-PCR

using primers against SARS-CoV-2-N was run on RNA isolated from FFPE patient

placenta slides. This provided us with 5 distinct cohorts for this study depicted in Table

1: High Positive (P1, P2; ddCT value > 9), Positive (P3-P12; ddCT > 4.5), and

Borderline Positive (P13-P22). We also ran qRT-PCR on RNA from patient placenta

samples from SARS-CoV-2 negative mothers (C1-C5) as well as SARS-CoV-2 negative

mothers that had unrelated inflammatory pathologies (I1-I5) (Figure 1A and Table 1). To

confirm presence of viral RNA, the presence of a distinct amplicon on PCR melt curve

and on gels run on RNA samples distinguished between the positive and negative

samples from all of the placenta samples obtained from COVID-positive mothers (data

not shown).

To identify the cells in the placental chorionic villi that were infected by SARS-CoV-2,

adjacent placental sample sections (10 microns apart) from the different cohorts were

stained by hematoxylin and eosin (H&E), or for the presence of replicating viral RNA, for

the presence of SARS-CoV-2 nucleocapsid protein (SARS-CoV-2-N), and for the

presence CD163$^+$ Hofbauer cells (HBC) and macrophages using

immunohistochemistry. SARS-CoV-2 RNA was detected by *in situ* hybridization in the

high positive samples, but not in negative control samples (Figure 1B). Presence of

SARS-CoV-2 RNA was restricted to the Keratin-7 (KRT7)-positive syncytial trophoblast

layer which anatomically encapsulate the chorionic villi structures (Figure 1B). Similarly,

expression of the SARS-CoV-2 N protein was detected in adjacent sections within the

same villi. Localization of the N protein was restricted to the syncytiotrophoblast layers

in the high positive placentas. Interestingly, the syncytial trophoblast layer of the high

positive sample had significantly fewer nuclei and appeared damaged as indicated by

the hematoxylin counterstaining. Importantly, at low magnification, massive infiltration of

maternal immune cells, including CD163+ macrophages detected in the high positive

samples, but not in the controls (Figure 1B and not shown). This result is consistent with

the pathology report for the sample P2 of chronic histiocytic intervillositis (CHI) (Table

1). Intravilllous HBC and intervillous maternal macrophages did not show infection with

SARS-CoV-2, evidenced by the absence of SARS-CoV-2 RNA and N protein. In

summary, these results indicate that syncytiotrophoblast are the primary targets for

SARS-CoV-2 infection in the placenta, and that the massive maternal immune cell

migration occurred in response to the SARS-CoV-2 infection either in the mother or in

the placenta.

Placental explant and cell cluster cultures are permissive to pseudo-entry virus

and infection can be blocked by anti-ACE2 and anti-NRP1 antibodies

To determine the SARS-CoV-2 tropism and infection of term placentas, we used

fresh placental isolates from SARS-CoV-2 negative mothers obtained immediately post-

delivery. After removal of the fetal chorionic plate and maternal decidua, samples

containing terminal, intermediate and stem chorionic villi were used for the preparation

of placental villi explant cultures (Figure 2C). In addition, placental cell clusters were

prepared by enzymatic digestion of the chorionic villi, followed by filtration that allows

passage of small cell clusters. Placental cultures were infected with a dual

nanoluciferase/green fluorescent protein (GFP) reporter lentiviral vector pseudotyped

with SARS-CoV-2 spike (S) (Tada et al. 2020), and luciferase activity was quantified 72

hpi. Lentiviral reporter viruses pseudotyped with vesicular stomatitis virus G protein

(VSV-G) were used as a positive control for infection (Figure 2). A comparison of the

infectivity showed that both pseudotyped viruses infected the placental cultures at

similar levels. Explant cultures consistently showed an approximately 5-fold lower

infection as compared to clusters or single cells, likely due to the reduced surface

accessibility. Infectivity was significantly reduced by adding the human

immunodeficiency virus (HIV) reverse transcriptase (RT) inhibitor nevirapine (NVP),

indicating that the luciferase activity was primarily due to viral entry and not carry-over

from residual viral particles in the cultures (Figure 2A). The major SARS-CoV-2 entry

factors, ACE2 and TMPRSS2 are expressed in the placenta, albeit their expression is

significantly decreased in the third trimester (Pique-Regi et al. 2020; Singh, Bansal, and

Feschotte 2020; Ouyang et al. 2021). Furthermore, NRP1 has been identified as a

novel host factor for SARS-CoV-2 (Cantuti-Castelvetri et al. 2020; Daly et al. 2020) and

is expressed on syncytiotrophoblast (Arad et al. 2017; Baston-Buest et al. 2011). To

determine if ACE2 and/or NRP1 facilitate infection in the placenta, we pre-treated

placental cell clusters with anti-ACE2 or anti-NRP1 blocking antibody prior to infection.

Both antibodies reduced infectivity of SARS-CoV-2 S protein pseudotyped lentivirus in

placental cell clusters by about 50%, while anti-ACE2 blocking antibody did not reduce

infectivity of VSV-G pseudotyped lentivirus. Pre-treatment with both antibodies did not

result in further reduction of infectivity, suggesting the possibility of alternative

receptor(s) (Figure 2B).

To determine which cells are targeted by the pseudotyped virus, placental explant

cultures were infected for 72 hours with lentivirus pseudotyped by SARS-CoV-2 spike

(S) protein, and live GFP could be visualized in the infected explant cultures, with a

more robust signal observed in the pseudotyped VSV-G infected cultures (Figure 2C).

Explant cultures were then processed for immunofluorescence staining and analyzed

for co-localization of the GFP reporter with KRT-7/Cytokeratin (trophoblast marker) and

CD31 (endothelial marker). GFP was detected in small patches of syncytiotrophoblast

cells located on the outer perimeter of the chorionic villi, but not in endothelial cells

(Figure 2D). Similar results were obtained after infection with VSV-G pseudotyped

lentivirus, with more robust infection visualized by live fluorescence microscopy on the

infected explant cultures, whereas no GFP signal was found in mock-infected explant

cultures (Figure 2D).

Primary placental cell clusters are permissive to SARS-CoV-2

To further determine the intrinsic susceptibility of placental cells to SARS-CoV-2,

primary placental cell clusters were infected *ex vivo*. Placentas were isolated from

healthy term deliveries as described above, digested into cell clusters of approximately

50-100 cells and plated on Matrigel-coated plates. Cell clusters were infected with live

SARS-CoV-2 virus (Isolate USA-WA1/2020, multiplicity of infection, MOI=1). Cells were

collected 24 hpi and virus load analyzed by qRT-PCR and immunofluorescence

staining.

qRT-PCR analysis using primers targeting subgenomic N transcripts demonstrated

robust SARS-CoV-2 viral replication in primary human placental cell clusters at 24 hpi

(Figure 3A). To determine which cells of primary placental cell clusters were susceptible

to SARS-CoV-2 infection, infected cell clusters were immunostained for SARS-CoV-2

nucleocapsid protein (SARS-N) and cell type specific markers for trophoblast cells

(KRT7) and endothelial cells (CD31). Three-dimensional reconstruction of confocal

imaging confirmed the presence of SARS-CoV-2-N protein in KRT7+

syncytiotrophoblast (Figure 3B). Co-localization of SARS-CoV-2-N protein was found in

multiple clusters of KRT7+ syncytiotrophoblast cells (Figure 3C). In addition to the

positive staining for SARS-CoV-2-N in syncytiotrophoblast, there were rare CD31+

endothelial cells that also stained positively for SARS-CoV-2-N protein (Figure 3C).

Discussion

The placenta is a vital organ that provides the gestational interface between mother

and fetus. Compromised maternal health and environmental insults, such as viral

infections, can result in placental dysfunction and lead to pregnancy complications with

increased morbidity and mortality for the mother and fetus (Rossant and Cross 2001;

Maltepe, Bakardjiev, and Fisher 2010; John and Hemberger 2012). Impairment of

placental function can also developmentally program the fetus for chronic disease later

in life, including cardiovascular disease, diabetes and obesity (Burton, Fowden, and

Thornburg 2016).

The aim of our study was to investigate the impact of late pregnancy SARS-CoV-2

infection on maternal and fetal health and proper placental function. Within a cohort of

54 placental samples from women who tested positive for SARS-CoV-2 at delivery, 22

were positive for genomic and replicating viral RNA. The observed percentage of

positive samples was higher compared to other studies (Hecht et al. 2020b; Facchetti et

al. 2020; Debelenko et al. 2021; Lu-Culligan et al. 2021) and likely reflects the fact that

New York City was at the epicenter for COVID-16 in March-May of 2020. Furthermore,

many of the placental samples were obtained from deliveries where the mothers or

infants presented with clinical pathologies which may bias sample collection.

Quantification of virus content in the placental samples revealed that the 2 cases with

high SARS-CoV-2 presence in the placenta were from pregnancies with adverse fetal

outcome, including fetal demise and preterm delivery. In contrast, only 1 out of 10

pregnancies with medium viral content resulted in IUFD, and this may have been

triggered by poorly controlled maternal T2D. Of the remaining pregnancies with medium

or low virus content in the placenta, all babies tested negative for SARS-CoV-2 and

were healthy at discharge. It will be important to follow up on the health of these infants

to investigate possible long-term effects of SARS-CoV-2. None of the pregnancies of

SARS-CoV-2 negative healthy controls (n=5) or inflammation controls (n=5) resulted in

fetal demise.

Upon examining sections from placentas with high virus content, we detected SARS-

CoV-2 RNA and protein in a large fraction of syncytiotrophoblast, the single cell layer

enveloping the fetal chorionic villi situated at the interphase to maternal blood. No virus

was detected in fetal macrophages (Hofbauer cells), other cell types inside the villi,

including stromal and endothelial cells, or outside the villi. Several recent reports also

provided evidence for SARS-CoV-2 infection restricted to syncytiotrophoblast of

placentas from SARS-CoV-2 positive mothers (Alamar et al. 2020; Mulvey et al. 2020;

Hecht et al. 2020b; Penfield et al. 2020; Hosier et al. 2020; Vivanti et al. 2020; Taglauer

et al. 2020; Facchetti et al. 2020). However, two reports on a preterm placenta and a

placenta from a newborn with vertically SARS-CoV-2 noted also presence of SARS-

CoV-2 in other cell types, including Hofbauer cells and stromal cells inside the villi, and

maternal macrophages and epithelial cells at the maternal-fetal interface (Verma et al.,

2021, Fachetti et al, 2020). It is possible that SARS-CoV-2 infection in these cases

occurred at earlier gestational stages and allowed for additional viral spread beyond the

syncytiotrophoblast layer.

Placentas in our study with high virus presence also displayed massive infiltration of

maternal immune cells, including macrophages into the intervillous space. However, we

did not detect SARS-CoV-2 in the infiltrating immune cells. Several recent studies

reported similar findings; interestingly they were most prominently found in placentas

from live-borne and stillborn neonates that had tested positive for SARS-CoV-2. These

finding included intervillous infiltration by inflammatory immune cells, chronic histiocytic

intravillositis with trophoblast necrosis, and increased fibrin deposition (Debelenko et al.

2021; Facchetti et al. 2020; Garrido-Pontnou et al. 2021; Lu-Culligan et al. 2021;

Schwartz et al. 2021; Morotti et al. 2021; Verma et al. 2021). Furthermore,

transcriptome data presented in one of these studies showed localized inflammatory

responses to systemic SARS-CoV-2 infection in the placenta, even if SARS-CoV-2 virus

was not detected (Lu-Culligan et al. 2021).

Considering the low numbers of placental infections by SARS-CoV-2 so far seen

clinically, we decided to complement our *in vivo* studies by using *ex vivo* placental

explant and cell cluster culture models to study virus entry. We showed infection with

SARS-CoV-2 virus or SARS-CoV-2 spike S pseudotyped lentivirus targeted

predominantly syncytiotrophoblast and, in rare instances, endothelial cells. Term

placentas express very low levels of the SARS-CoV-2 receptor ACE2 and the co-factor

TMPRSS2 (Pique-Regi et al. 2020; Singh, Bansal, and Feschotte 2020; Ouyang et al.

2021). Recently, Lu-Culligan et al. (Lu-Culligan et al. 2021) reported on increased levels

of placental ACE2 expression in COVID-19 positive mothers; whereas a second

publication reported on a decrease of ACE2 expression and dysregulation of the renin-

angiotensin system (Verma et al. 2021). We asked here if alternative entry factors might

be used by SARS-CoV-2 to infect placental cells. One likely candidate is the

transmembrane protein NRP1, which has recently been identified as a host factor that

facilitates SARS-CoV-2 cell entry and infectivity (Cantuti-Castelvetri et al. 2020; Daly et

al. 2020). NRP1 was originally identified as a co-receptor for Vascular endothelial

growth factor (VEGF) on endothelial cells but is expressed also at the maternal-fetal

interface in decidual cells and syncytiotrophoblast, and is thought to play important roles

during pregnancy and in the immune system (Arad et al. 2017; Baston-Buest et al.

2011). We show that infection by SARS-CoV-2 S pseudotyped lentivirus can be partially

inhibited by using blocking anti-ACE2 or anti-NRP1 antibodies, whereas infection by

VSV-G pseudotyped lentivirus is not blocked, suggesting that SARS-CoV-2 may use

NRP1 as an alternative entry factor, and suggesting the existence of additional entry

factors in the placenta.

The present study focused on late cohort infections from mothers who tested

positive at the time of delivery. It will be important to study the impact of early cohort

infections in mothers who are serologically positive at delivery but negative for viral

RNA, as infection during the first and second trimester may affect placental

development and morphogenesis and result in different placental pathologies and

clinical outcomes for mother and fetus.

Materials and Methods

Cell Lines

Vero E6 (African green monkey [Chlorocebus aethiops] kidney) were obtained from ATCC (https://www.atcc.org/). Vero E6 and A549 (adenocarcinomic human alveolar basal epithelial cell line)-ACE2 cells were cultured in Dulbecco's Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 100 U/mL penicillin and 100 µg/mL streptomycin, and maintained at 37°C with 5% CO_2.

SARS-CoV-2 propagation and infection

SARS-CoV-2 isolate USA-WA1/2020 (NR-52281) was provided by the Center for Disease Control and Prevention (CDC) and obtained through BEI Resources, NIAID, NIH. SARS-CoV-2 was propagated in Vero E6 cells in DMEM supplemented with 2% FBS, 4.5 g/L D-glucose, 4 mM L-glutamine, 10 mM Non-essential amino acids, 1 mM sodium pyruvate and 10 mM HEPES using a passage-2 stock of virus. Three days after infection, supernatant containing propagated virus was filtered through an Amicon Ultra 15 (100 kDa) centrifugal filter (Millipore Sigma) at ~4000 rpm for 20 minutes. Flow-through was discarded and virus was resuspended in DMEM supplemented as described above. Infectious titers of SARS-CoV-2 were determined by plaque assay in Vero E6 cells in Minimum Essential Media supplemented with 2% FBS, 4 mM L-glutamine, 0.2% BSA, 10 mM HEPES and 0.12% $NaHCO_3$ and 0.7% agar. All MOI values were based on titer determined from plaque assays on Vero E6 cells. All work involving live SARS-CoV-2 was performed in the CDC/USDA-approved biosafety level-3

(BSL-3) facility of the Icahn School of Medicine at Mount Sinai in accordance with

institutional biosafety requirements.

Placental Samples

Placental tissues from SARS-CoV-2 positive women and controls were obtained at

delivery at Weill Cornell-NY Presbyterian by the Department of Pathology and

Laboratory Medicine at Weill Cornell Medicine. All women admitted for delivery were

tested by nasal swabs for acute SARS-CoV-2 infection by qRT-PCR, and serologically

for previous infection at Weill Cornell Medicine Department of Pathology and Laboratory

Medicine. Infants were tested for SARS-CoV-2 at birth and 1 week of age by nasal

swabs and RT-PCR. Placental samples were fixed for 48 hours in formalin and then

processed and embedded into formalin fixed paraffin embedded (FFPE) blocks by the

pathology department. FFPE placenta samples from 5 healthy women who tested

negative for SARS-CoV-2 served as controls. An additional 5 FFPE placental samples

with inflammation pathologies, obtained from SARS-CoV-2 negative patients, were also

included in the study. Unstained sections and H&E sections of the FFPE blocks were

performed at the Weill Cornell Clinical & Translational Science Center (CTSC) core

facility. Additional H&E staining was performed by the Weill Cornell Histology core

facility.

SARS-CoV-2 Detection in RNA from FFPE Placental Sections by qRT-PCR

Total RNA samples were prepared from FFPE placental tissue sections, followed by

DNaseI treatment using manufacturer's instructions (Qiagen RNeasy FFPE kit Cat#

73604). To quantify viral replication, as measured by the expression of nucleocapsid

sub genomic viral RNA along with the housekeeping gene GAPDH, two-step RT-qPCR

was performed using LunaScript® RT SuperMix Kit (E3010L) for cDNA synthesis and

Luna® Universal qPCR Master Mix (NEB #M3003) for RT-qPCR. Quantitative real-time

PCR reactions were performed on CFX384 Touch Real-Time PCR Detection System

(BioRad). The sequences of primers/probes are provided below.

SARS-CoV-2-N

Forward 5' CTCTTGTAGATCTGTTCTCTAAACGAAC 3'

Reverse 5' GGTCCACCAAACGTAATGCG 3'

GAPDH

Forward 5' CATCACCATCTTCCAGGAGCGAGAT 3'

Reverse 5' GAGGCATTGCTGATGATCTTGAGGC 3'

qRT-PCR graphs were generated using GraphPad Prism software.

RNA *In Situ* Hybridization to Detect SARS-CoV2 RNA on FFPE Placental Sections

Probe design. Probes were designed with a 20-25 nucleotides homology to SARS-

CoV-2 genomic RNA and were assessed by NCBI BLAST to exclude off target binding

to other cellular transcripts. IDT OligoAnalyzer (Integrated DNA Technologies) was used

to identify probe pairs with similar thermodynamic properties, melting temperature 45-

60ºC, GC content of 40-55%, and low self-complementary. The 3' end of each one of

the probes used for proximity ligation signal amplification is designed with a partially

complementary sequence to the 61bp long backbone and partially to the 21bp insert as

described previously (Yang et al. 2020).

Tissue viral RNA staining pretreatment. Sections of FFPE placental samples were deparaffinized using 100% xylenes, 5 min at room temperature, repeated twice. Slides were rinsed in 100% ethanol, 1 min at room temperature, twice and air dried. Endogenous peroxidase activity was quenched by treating the samples with 0.3% hydrogen peroxide, 10 min at room temperature followed by washing with DEPC treated water. Samples were incubated 15 min at 95-100 °C in antigen retrieval solution (ACDBio, Newark, CA, USA) rinsed in DEPC treated water and dehydrated in 100% ethanol, 3 min at room temperature and air dried. Tissue sections were permeabilized 30 min at 40°C using RNAscope protease plus solution (ACDBio, Newark, CA, USA) and rinsed in DEPC treated water.

SARS-CoV-2 RNA detection by probes proximity ligation. Hybridization was performed overnight at 40°C in a buffer based on DEPC-treated water containing 2× SSC, 20% formamide (Thermo Fischer Scientific, Waltham, MA, USA), 2.5 % (vol/vol) polyvinylsulfonic acid, 20 mM ribonucleoside vanadyl complex (New England Biolabs, Ipswich, MA, USA), 40 U/ml RNasin (Promega, Madison, WI, USA), 0.1% (vol/vol) Tween 20 (Sigma Aldrich), 100 µg/ml salmon sperm DNA (Thermo Fisher Scientific, Waltham, MA, USA), 100 µg/ml yeast RNA (Thermo Fisher Scientific, Waltham, MA, USA). DNA probes dissolved in DEPC-treated water were added at a final concentration of 100nM (Integrated DNA Technologies, Coralville, IA, ISA). Samples were washed briefly and incubated in 2× SSC, 20% formamide, 40 U/ml RNasin at 40 °C and then washed four times (5 min each) in PBS, 0.1% (vol/vol) Tween 20, and 4 U/ml RNasin (Promega, Madison, WI, USA). Slides were then incubated with 100 nM insert/backbone oligonucleotides in PBS, 1× SSC, 0.1% (vol/vol) Tween 20, 100 µg/ml

salmon sperm DNA (Thermo Fisher Scientific, Waltham, MA, USA), 100 µg/ml yeast

RNA (Thermo Fisher Scientific, Waltham, MA, USA), 40 U/ml RNasin at 37 °C. After

four washes, tissues were incubated at 37°C with 0.1 U/µl T4 DNA ligase (New England

Biolabs, Ipswich, MA, USA) in 50mM Tris-HCl, 10mM $MgCl_2$, 1mM ATP, 1mM DTT,

250µg/ml BSA, 0.05% Tween 20, 40 U/ml RNasin, followed by incubation with 0.1 U/µl

phi29 DNA polymerase in 50 mM Tris–HCl, 10 mM $MgCl_2$, 10 mM $(NH_4)_2SO_4$, 250µM

dNTPs, 1mM DTT, 0.05% Tween 20, 40 U/ml RNasin pH 7.5 at 30 °C. Slides were

washed and endogenous biotin was blocked using Avidin/Biotin blocking kit (Vector

laboratories, Burlingame, CA, USA) according to the manufacture instructions. Rolling

cycle amplicons were identified using a biotin labeled DNA probe at a concentration of 5

nM at 37 °C in PBS, 1× SSC, 0.1% Tween 20, 100 µg/ml salmon sperm DNA, 100

µg/ml yeast RNA, 1 hr. After washing, samples were incubated with 1:100 diluted

streptavidin-HRP (Thermo Fisher Scientific, Waltham, MA, USA) in PBS, 60 min at

room temperature followed by washing. Labeling was accomplished using EnzMet kit

(Nanoprobes, Yaphank, NY, USA) according to manufacture instructions. Slides were

further labeled with rabbit anti-cytokeratin 1:250 (Dako Z0622), overnight 4°C. After

washing, samples were incubated with 1:1000 with anti-rabbit alkaline phosphatase

antibody (1:1000, Jackson immunoresearch, Baltimore, PA, USA) and stained using

Fast Red substrate kit according the manufacture instructions (Abcam, Cambridge, MA,

USA). Hematoxylin was used for counterstaining (Vector laboratories, Burlingame, CA,

USA), and samples were mounted in Permount (Fischer Scientific, Waltham, MA, USA).

Proximity Ligation Probes	Sequence
SARS Cov2-1	TGA GTT GGA CGT GTG TTT TCA AAA AAA AAA ACT CAG TCG TGA

	CAC TCT T
SARS Cov2-2	AGC ACG TCG CGA ACC TGT AAA AAA AAA AGA CGC TAA TAT CGT GAC C
SARS Cov2-3	AAT GCA CTC AAG AGG GTA GCA AAA AAA AAA ACT CAG TCG TGA CAC TCT T
SARS Cov2-4	GCT TTA CCA GCA CGT GCT AGA AAA AAA AAA AGA CGC TAA TAT CGT GAC C
SARS Cov2-5	TCC AAA GGC AAT AGT GCG ACA AAA AAA AAA ACT CAG TCG TGA CAC TCT T
SARS Cov2-6	ATG GCA ACC AAC ATA AGA GAA AAA AAA AAA AGA CGC TAA TAT CGT GAC C
SARS Cov2-7	CCA GTT GAA ACT ACA AAT GGA AAA AAA AAA CTC AGT CGT GAC ACT CTT
SARS Cov2-8	ACA ACA CCT AGC TCT CTG AAG AAA AAA AAA AGA CGC TAA TAT CGT GAC C
SARS Cov2-9	GAA ACA CAC AAC AGC ATC GTA AAA AAA AAA CTC AGT CGT GAC ACT CTT
SARS Cov2-10	CAC TAG ACC TTG AGA TGC ATA AAA AAA AAA GAC GCT AAT ATC GTG ACC
SARS Cov2-11	GTC TTT CAG TAC AGG TGT TAA AAA AAA AAA CTC AGT CGT GAC ACT CTT
SARS Cov2-12	TGA GCG TTT CTG CTG CAA AAA AAA AAA AAA GAC GCT AAT ATC GTG ACC
Insert	/5Phos/ACGACTGAGTTTGGTCACGAT
Backbone	/5Phos/ATTAGCGTCCAGTGAATGCGAGTCCGTCTAGGAGAGTAGTACA GCAGCCGTCAAGAGTGTC
Detection	/5BiosG/ACGACTGAGTTTGGTCACGAT

Placental Explants and Cluster Cultures

Fresh de-identified placentas from SARS-CoV-2-negative mothers were collected

within 30 min to 2 hours post-delivery from Labor & Delivery at WCM/NYP. Collection of

placentas was performed under an approved IRB exempt protocol (#20-07022453, Weill

Cornell Medicine.) Tissue samples were dissected by removing the fetal chorionic plate

and any remaining maternal decidual tissue. Primary explant cultures (1cm x 1cm x

2cm) containing terminal, intermediate and stem chorionic villi were further dissected,

washed in ice cold 1X PBS to remove maternal blood, and plated into 48-well plastic

dishes in DMEM/F12 medium supplemented with 10% FBS and 100 U/mL penicillin,

100 µg/mL streptomycin and 0.25 µg/mL amphotericin B, as previously described

(Massimiani et al. 2019).

In addition, placental cell clusters were prepared from fresh chorionic villi tissue

samples by mincing with scissors and 10 blade scalpels. The minced tissue was

digested using 0.2 mg/mL collagenase/ 0.8U/mL dispase (Roche) and recombinant

DNAse I (Sigma) in MACS buffer (PBS/2mM EDTA/ 0.5% bovine serum albumin (BSA))

at 42^{o}C with agitation by pipetting with a 5 ml stripette. The digested tissue was filtered

through 100 µ filters (Corning 352360), and red blood cells (RBC) were removed using

RBC Lysis Buffer (Biolegend 420301). Clusters were washed once in MACS buffer,

examined for viability with Trypan Blue (GIBCO) and plated onto Matrigel-coated 96-

well dishes and µ-slide 8-well chamber slides (ibidi GmbH, Germany) at confluent

density in DMEM/F12 supplemented with 10% FBS and penicillin/streptomycin/

fungizone, and were incubated at 37°C with 5% CO_2 for 24 hours prior to infection with

pseudovirus to allow for attachment,

For infection with SARS-CoV-2, Sections of fresh chorionic villi (2g) were minced

with sterile scalpels, digested in Accutase (Innovative Cell Technologies) for 7 min or

isolated using a human umbilical cord dissociation kit (Millitenyi Biotec 130-105-737),

and filtrated through a 100 µm cell strainer (Falcon) to obtain cell clusters of ~50-100

cells. Red blood cells were lysed using RBC Lysis Buffer (Biolegend), washed with

PBS-0.5% BSA, and resuspended in medium (DMEM-10%FBS-1% Pen-Strep-

Glutamax). Cell viability was determined with Trypan blue (Gibco). Cell clusters were

plated on Matrigel (Corning, hESC-qualified)-coated plates at 4x10^5/well in 24-well

plates or 3x10^4/well in glass-like polymer bottom 96-well plates (CellVis).

Infection of Ex Vivo Placental Cultures

Infection of Explants and Placental clusters with Pseudovirus. Lentiviruses encoding dual Nanoluciferase/GFP reporter lentivirus and pseudotyped by SARS-CoV-2 spike (S) protein (D614G) or VSV-G were prepared as previously described (Tada et al. 2020). The viruses were concentrated 10-fold by ultracentrifugation and titers were quantified by reverse transcriptase assay. Placental explant cultures and cell clusters were infected with 10 µl SARS-CoV-2 S or VSV-G pseudotyped lentivirus (Tada et al. 2020). To determine whether NRP1 or ACE2 facilitates the infection of placental cells, placental cell clusters were pretreated for 30 min with anti-NRP1 mAb (R&D Systems, AF3870) or anti-ACE2 mAb (Agilent, AG-20A-0032-C50). Infected placental clusters were lysed 72 hours post-infection. Luciferase activity was measured using a Promega Nano-Glo Assay Kit and read on an Envision microplate luminometer (Perkin Elmer).

Infection of Placental Clusters with Live SARS-CoV-2. Placental cell clusters were infected with live SARS-CoV-2 (isolate USA-WA1/2020 (NR-52281) at an MOI of 0.1 and 1 or mock-infected at day-1 in culture as recently described (Yang et al. 2020). At the indicated hpi, cells were washed three times with PBS. For RNA analysis cells were lysed in TRIzol (Invitrogen). For immunofluorescence staining cells were fixed in 4% formaldehyde for 60 min at room temperature. All work involving live SARS-CoV-2 was performed in the CDC/USDA-approved BSL-3 facility of the Icahn School of Medicine at Mount Sinai in accordance with institutional biosafety requirements.

qRT-PCR for Viral Load of SARS-CoV-2 Infected Placental Clusters. Total RNA was extracted using Trizol (Thermo Fisher Scientific, Waltham, MA, USA) followed by

ezDNAse treatment (Thermo Fisher Scientific, Waltham, MA, USA) per manufacturer's

instructions. To quantify viral replication, measured by the accumulation of subgenomic

N transcripts, one-step quantitative real-time PCR was performed using the SuperScript

III Platinum SYBR Green One-Step qRT–PCR Kit (Invitrogen) with primers specific for

TRS (listed above) and beta-actin (ACTB) as an internal reference (listed below), as

previously described (Yang et al. 2020). Reactions were performed on a QuantStudio 6

Flex Real Time PCR Instrument (Applied Biosystems). The delta-delta-cycle threshold

($\Delta\Delta$CT) was determined relative to ACTB and mock-infected samples. Graphs were

generated using GraphPad Prism software.

Primer Name	Sequence (5'-3')
ACTB-Forward	CGTCACCAACTGGGACGACA
ACTB-Reverse	CTTCTCGCGGTTGGCCTTGG

Immunostaining of FFPE Placental Sections and Infected Placental Cell Clusters

IHC for SARS-CoV2 and Hofbauer Cells on FFPE Slides. Immunohistochemistry

(IHC) was performed on FFPE slides using ImmPRESS Reagent kit (Vector

laboratories, Burlingame, CA, USA). FFPE slides were dewaxed in a hybrid oven for 45

minutes at 55^{o}C and then rehydrated using xylenes followed by a standard ethanol

gradient. Antigen retrieval was performed using sodium citrate buffer, pH 6.1 in a

steamer for 35 minutes. Slides were blocked using 2.5% horse serum (Vector

laboratories) for 1 hour at room temperature and then incubated overnight at 4°C in a

humid chamber with primary antibodies (SARS-CoV-2-N,GeneTex GTX635679, at

1:100; CD163,Novus Biologicals NBP2-48846, 1:250) diluted in 1% BSA/0.1% Triton-X

PBS (PBST). Slides were treated with 3% hydrogen peroxide at room temperature

(Sigma H1009), washed 3 times with 0.1% PBST and then incubated for 1 hour at room

temperature with ImmPRESS anti-rabbit peroxidase conjugated antibody (Vector

Laboratories, Burlingame, CA, USA). Slides were again washed 3 times with 0.1%

PBST with final wash in PBS prior to developing using freshly prepared DAB substrate

(Vector Labs SK-4100). Slides were rinsed with water and counterstained with

Hematoxylin (RICCA Chemical Company, Arlington, TX, USA) to mark the nuclei.

Stained slides were dehydrated using an increasing ethanol gradient, treated with

xylenes, and then mounted with Permount solution (Thermo Fisher Scientific, Waltham,

MA, USA). Brightfield images were acquired using a Zeiss microscope (Carl Zeiss,

Germany).

IF Staining for Pseudovirus Infected Placental Explants/Clusters. For

immunofluorescence (IF) analysis, SARS-CoV-2 GFP-pseudotyped virus infected

explant cultures were drop-fixed overnight in 4% paraformaldehyde (PFA) in PBS

containing $Ca2^+/Mg^{2+}$ at 4^oC on a rocker 72 hours post-infection. The fixed explants

were then dehydrated with 30% sucrose in PBS overnight at 4^oC on a rocker. Explants

were embedded in optimal cutting temperature compound (OCT) on dry ice. Frozen

blocks were sectioned on a cryomicrotome at 10 micron thickness. Explant culture

sections were blocked for 1 hour in 10% donkey serum (Jackson ImmunoResearch

labs, Westgrove, PA) in 0.1% PBST. Primary antibodies (rabbit anti-cytokeratin 1:1000

(Dako Z0622), sheep anti-human CD31 1:500 (BD AF806) and chicken anti-GFP

1:1000 (Abcam ab13970)) were diluted in 10% donkey serum-0.1% PBST and

incubated overnight at 4^oC followed by incubation with secondary antibodies

(AlexaFlour647-donkey anti-rabbit, AlexaFlour594-donkey anti-sheep, and

AlexaFlour488-donkey anti-chicken, Jackson ImmunoResearch labs, Westgrove, PA)).

The clusters were then stained using 4',6-diamidino-2-phenylindole (DAPI).

Slides were mounted with coverslips using ProLong Gold Antifade Mountant with DAPI

(Thermo Fisher Scientific, Waltham, MA, USA). Fluorescence microscopy was

performed using a Zeiss fluorescent microscope and image analysis was done using

ImageJ software.

Immunofluorescence Staining for SARS-CoV-2 of Infected Placental Clusters.

PFA-fixed cells were blocked in 5% normal donkey serum in PBS-0.05% Triton X-0.01%

Saponin (PBS-TSP). Primary antibodies (SARS-CoV2-N, Genetex GTX635679, 1:200;

KRT7, Agilent Dako M701829-2, 1:400; PECAM1, R&D AF806, 1:1000) were incubated

overnight at 4degC in block, followed by incubation in secondary antibodies

(AlexaFluor488-donkey-anti-mouse, AlexaFluor594-donkey-anti-rabbit, AlexaFluor647-

donkey-anti-sheep, ThermoFisher, 1:500) in PBS-TSP, and counterstaining with DAPI

(Thermo Fisher Scientific, Waltham, MA, USA). Images were acquired using a Zeiss

LSM 800 Confocal microscope and processed using Imaris software (Bitplane).

Acknowledgements

We thank the patients, their families, and healthcare workers fighting the COVID-19

pandemic. This work was supported by a Weill Cornell Medicine COVID-19 Research

Grant (H.S., R.E.S., R.N.B. Baergen), the NCI (R01CA234614) and NIAID

(2R01AI107301) and NIDDK (R01DK121072) to Department of Medicine, Weill Cornell

Medicine (R.E.S.), NIDDK (R01DK119667, R01DK119667-02S1) to S.C. R.E.S. and

S.C. are supported as an Irma Hirschl Trust Research Award Scholar. LBA was

supported in part by NYSTEM Training grant. L.A.L. is supported by an F32 post-

doctoral fellowship from the National Institute of Health (1F32HD096810-01A1) and

Weill Cornell Medicine Research Assistance for Primary Parents Award. N.R.L was

supported by grants from the NIH (DA046100, AI122390 and AI120898). T.T. was

supported by the Vilcek/Goldfarb Fellowship Endowment Fund. We would also like to

acknowledge Michael D. Glendenning for his technical assistance with the IHC and the

WCM Histology Core.

Competing Interests

R.E.S. is on the scientific advisory board of Miromatrix Inc and is a consultant and

speaker for Alnylam Inc.

References

Alamar, I., M. H. Abu-Arja, T. Heyman, D. J. Roberts, N. Desai, P. Narula, and B. Dygulska. 2020. 'A Possible Case of Vertical Transmission of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in a Newborn With Positive Placental In Situ Hybridization of SARS-CoV-2 RNA', *J Pediatric Infect Dis Soc*, 9: 636-39.

Arad, A., S. Nammouz, Y. Nov, G. Ohel, J. Bejar, and Z. Vadasz. 2017. 'The Expression of Neuropilin-1 in Human Placentas From Normal and Preeclamptic Pregnancies', *Int J Gynecol Pathol*, 36: 42-49.

Baergen, R. N., and D. S. Heller. 2020. 'Placental Pathology in Covid-19 Positive Mothers: Preliminary Findings', *Pediatr Dev Pathol*, 23: 177-80.

Baston-Buest, D. M., A. C. Porn, A. Schanz, J. S. Kruessel, W. Janni, and A. P. Hess. 2011. 'Expression of the vascular endothelial growth factor receptor neuropilin-1 at the human embryo-maternal interface', *Eur J Obstet Gynecol Reprod Biol*, 154: 151-6.

Blanco-Melo, D., B. E. Nilsson-Payant, W. C. Liu, S. Uhl, D. Hoagland, R. Møller, T. X. Jordan, K. Oishi, M. Panis, D. Sachs, T. T. Wang, R. E. Schwartz, J. K. Lim, R. A. Albrecht, and B. R. tenOever. 2020. 'Imbalanced Host Response to SARS-CoV-2 Drives Development of COVID-19', *Cell*, 181: 1036-45.e9.

Burton, G. J., A. L. Fowden, and K. L. Thornburg. 2016. 'Placental Origins of Chronic Disease', *Physiol Rev*, 96: 1509-65.

Cantuti-Castelvetri, L., R. Ojha, L. D. Pedro, M. Djannatian, J. Franz, S. Kuivanen, F. van der Meer, K. Kallio, T. Kaya, M. Anastasina, T. Smura, L. Levanov, L. Szirovicza, A. Tobi, H. Kallio-Kokko, P. Österlund, M. Joensuu, F. A. Meunier, S. J. Butcher, M. S. Winkler, B. Mollenhauer, A. Helenius, O. Gokce, T. Teesalu, J. Hepojoki, O. Vapalahti, C. Stadelmann, G. Balistreri, and M. Simons. 2020. 'Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity', *Science*, 370: 856-60.

Daly, J. L., B. Simonetti, K. Klein, K. E. Chen, M. K. Williamson, C. Antón-Plágaro, D. K. Shoemark, L. Simón-Gracia, M. Bauer, R. Hollandi, U. F. Greber, P. Horvath, R. B. Sessions, A. Helenius, J. A. Hiscox, T. Teesalu, D. A. Matthews, A. D. Davidson, B. M. Collins, P. J. Cullen, and Y. Yamauchi. 2020. 'Neuropilin-1 is a host factor for SARS-CoV-2 infection', *Science*, 370: 861-65.

Daniloski, Z., T. X. Jordan, H. H. Wessels, D. A. Hoagland, S. Kasela, M. Legut, S. Maniatis, E. P. Mimitou, L. Lu, E. Geller, O. Danziger, B. R. Rosenberg, H. Phatnani, P. Smibert, T. Lappalainen, B. R. tenOever, and N. E. Sanjana. 2021. 'Identification of Required Host Factors for SARS-CoV-2 Infection in Human Cells', *Cell*, 184: 92-105.e16.

Debelenko, L., I. Katsyv, A. M. Chong, L. Peruyero, M. Szabolcs, and A. C. Uhlemann. 2021.
'Trophoblast damage with acute and chronic intervillositis: disruption of the placental
barrier by severe acute respiratory syndrome coronavirus 2', *Hum Pathol*, 109: 69-79.
Della Gatta, A. N., R. Rizzo, G. Pilu, and G. Simonazzi. 2020. 'Coronavirus disease 2019 during
pregnancy: a systematic review of reported cases', *Am J Obstet Gynecol*, 223: 36-41.
Edlow, A. G., J. Z. Li, A. Y. Collier, C. Atyeo, K. E. James, A. A. Boatin, K. J. Gray, E. A. Bordt,
L. L. Shook, L. M. Yonker, A. Fasano, K. Diouf, N. Croul, S. Devane, L. J. Yockey, R.
Lima, J. Shui, J. D. Matute, P. H. Lerou, B. O. Akinwunmi, A. Schmidt, J. Feldman, B. M.
Hauser, T. M. Caradonna, D. De la Flor, P. D'Avino, J. Regan, H. Corry, K. Coxen, J.
Fajnzylber, D. Pepin, M. S. Seaman, D. H. Barouch, B. D. Walker, X. G. Yu, A. J.
Kaimal, D. J. Roberts, and G. Alter. 2020. 'Assessment of Maternal and Neonatal SARS-
CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in
Pregnancies During the COVID-19 Pandemic', *JAMA Netw Open*, 3: e2030455.
Facchetti, F., M. Bugatti, E. Drera, C. Tripodo, E. Sartori, V. Cancila, M. Papaccio, R. Castellani,
S. Casola, M. B. Boniotti, P. Cavadini, and A. Lavazza. 2020. 'SARS-CoV2 vertical
transmission with adverse effects on the newborn revealed through integrated
immunohistochemical, electron microscopy and molecular analyses of Placenta',
EBioMedicine, 59: 102951.
Garrido-Pontnou, M., A. Navarro, J. Camacho, F. Crispi, M. Alguacil-Guillén, A. Moreno-Baró, J.
Hernandez-Losa, M. Sesé, Y. Cajal S. Ramón, I. Garcia Ruíz, B. Serrano, P. Garcia-
Aguilar, A. Suy, J. C. Ferreres, and A. Nadal. 2021. 'Diffuse trophoblast damage is the
hallmark of SARS-CoV-2-associated fetal demise', *Mod Pathol*: 1-6.
Hecht, J. L., B. Quade, V. Deshpande, M. Mino-Kenudson, D. T. Ting, N. Desai, B. Dygulska, T.
Heyman, C. Salafia, D. Shen, S. V. Bates, and D. J. Roberts. 2020a. 'SARS-CoV-2 can
infect the placenta and is not associated with specific placental histopathology: a series
of 19 placentas from COVID-19-positive mothers', *Mod Pathol*: 1-12.
Hoffmann, M., H. Kleine-Weber, S. Schroeder, N. Krüger, T. Herrler, S. Erichsen, T. S.
Schiergens, G. Herrler, N. H. Wu, A. Nitsche, M. A. Müller, C. Drosten, and S.
Pöhlmann. 2020. 'SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is
Blocked by a Clinically Proven Protease Inhibitor', *Cell*, 181: 271-80.e8.
Hosier, H., S. F. Farhadian, R. A. Morotti, U. Deshmukh, A. Lu-Culligan, K. H. Campbell, Y.
Yasumoto, C. B. Vogels, A. Casanovas-Massana, P. Vijayakumar, B. Geng, C. D. Odio,
J. Fournier, A. F. Brito, J. R. Fauver, F. Liu, T. Alpert, R. Tal, K. Szigeti-Buck, S.
Perincheri, C. Larsen, A. M. Gariepy, G. Aguilar, K. L. Fardelmann, M. Harigopal, H. S.

Taylor, C. M. Pettker, A. L. Wyllie, C. D. Cruz, A. M. Ring, N. D. Grubaugh, A. I. Ko, T. L. Horvath, A. Iwasaki, U. M. Reddy, and H. S. Lipkind. 2020. 'SARS-CoV-2 infection of the placenta', *J Clin Invest*, 130: 4947-53.

John, R., and M. Hemberger. 2012. 'A placenta for life', *Reprod Biomed Online*, 25: 5-11.

Kimberlin, D. W., and S. Stagno. 2020. 'Can SARS-CoV-2 Infection Be Acquired In Utero?: More Definitive Evidence Is Needed', *JAMA*.

Li, M., L. Chen, J. Zhang, C. Xiong, and X. Li. 2020. 'The SARS-CoV-2 receptor ACE2 expression of maternal-fetal interface and fetal organs by single-cell transcriptome study', *PLoS One*, 15: e0230295.

Lu-Culligan, A., A. R. Chavan, P. Vijayakumar, L. Irshaid, E. M. Courchaine, K. M. Milano, Z. Tang, S. D. Pope, E. Song, C. B. F. Vogels, W. J. Lu-Culligan, K. H. Campbell, A. Casanovas-Massana, S. Bermejo, J. M. Toothaker, H. J. Lee, F. Liu, W. Schulz, J. Fournier, M. C. Muenker, A. J. Moore, L. Konnikova, K. M. Neugebauer, A. Ring, N. D. Grubaugh, A. I. Ko, R. Morotti, S. Guller, H. J. Kliman, A. Iwasaki, and S. F. Farhadian. 2021. 'Maternal respiratory SARS-CoV-2 infection in pregnancy is associated with a robust inflammatory response at the maternal-fetal interface', *Med (N Y)*, 2: 591-610.e10.

Maltepe, E., A. I. Bakardjiev, and S. J. Fisher. 2010. 'The placenta: transcriptional, epigenetic, and physiological integration during development', *J Clin Invest*, 120: 1016-25.

Marín Gabriel, M. A., M. Reyne Vergeli, S. Caserío Carbonero, L. Sole, T. Carrizosa Molina, I. Rivero Calle, I. Cuadrado Pérez, B. Álvarez Fernández, A. Forti Buratti, and A. Fernández-Cañadas Morillo. 2020. 'Maternal, Perinatal and Neonatal Outcomes With COVID-19: A Multicenter Study of 242 Pregnancies and Their 248 Infant Newborns During Their First Month of Life', *Pediatr Infect Dis J*, 39: e393-e97.

Massimiani, M., L. A. Lacko, C. S. Burke Swanson, S. Salvi, L. B. Argueta, S. Moresi, S. Ferrazzani, S. E. Gelber, R. N. Baergen, N. Toschi, L. Campagnolo, and H. Stuhlmann. 2019. 'Increased circulating levels of Epidermal Growth Factor-like Domain 7 in pregnant women affected by preeclampsia', *Transl Res*, 207: 19-29.

Morotti, D., M. Cadamuro, E. Rigoli, A. Sonzogni, A. Gianatti, C. Parolin, L. Patanè, and D. A. Schwartz. 2021. 'Molecular Pathology Analysis of SARS-CoV-2 in Syncytiotrophoblast and Hofbauer Cells in Placenta from a Pregnant Woman and Fetus with COVID-19', *Pathogens*, 10.

Mullins, E., M. L. Hudak, J. Banerjee, T. Getzlaff, J. Townson, K. Barnette, R. Playle, A. Perry, T. Bourne, and C. C. Lees. 2021. 'Pregnancy and neonatal outcomes of COVID-19:

681 coreporting of common outcomes from PAN-COVID and AAP-SONPM registries',
682 *Ultrasound Obstet Gynecol*, 57: 573-81.

683 Mulvey, J. J., C. M. Magro, L. X. Ma, G. J. Nuovo, and R. N. Baergen. 2020. 'Analysis of
684 complement deposition and viral RNA in placentas of COVID-19 patients', *Ann Diagn*
685 *Pathol*, 46: 151530.

686 Ou, X., Y. Liu, X. Lei, P. Li, D. Mi, L. Ren, L. Guo, R. Guo, T. Chen, J. Hu, Z. Xiang, Z. Mu, X.
687 Chen, J. Chen, K. Hu, Q. Jin, J. Wang, and Z. Qian. 2020. 'Characterization of spike
688 glycoprotein of SARS-CoV-2 on virus entry and its immune cross-reactivity with SARS-
689 CoV', *Nat Commun*, 11: 1620.

690 Ouyang, Y., T. Bagalkot, W. Fitzgerald, E. Sadovsky, T. Chu, A. Martínez-Marchal, M. Brieño-
691 Enríquez, E. J. Su, L. Margolis, A. Sorkin, and Y. Sadovsky. 2021. 'Term Human
692 Placental Trophoblasts Express SARS-CoV-2 Entry Factors ACE2, TMPRSS2, and
693 Furin', *mSphere*, 6.

694 Penfield, C. A., S. G. Brubaker, M. A. Limaye, J. Lighter, A. J. Ratner, K. M. Thomas, J. A.
695 Meyer, and A. S. Roman. 2020. 'Detection of severe acute respiratory syndrome
696 coronavirus 2 in placental and fetal membrane samples', *Am J Obstet Gynecol MFM*, 2:
697 100133.

698 Pique-Regi, R., R. Romero, A. L. Tarca, F. Luca, Y. Xu, A. Alazizi, Y. Leng, C. D. Hsu, and N.
699 Gomez-Lopez. 2020. 'Does the human placenta express the canonical cell entry
700 mediators for SARS-CoV-2?', *Elife*, 9.

701 Prabhu, M., K. Cagino, K. C. Matthews, R. L. Friedlander, S. M. Glynn, J. M. Kubiak, Y. J. Yang,
702 Z. Zhao, R. N. Baergen, J. I. DiPace, A. S. Razavi, D. W. Skupski, J. R. Snyder, H. K.
703 Singh, R. B. Kalish, C. M. Oxford, and L. E. Riley. 2020. 'Pregnancy and postpartum
704 outcomes in a universally tested population for SARS-CoV-2 in New York City: a
705 prospective cohort study', *BJOG*, 127: 1548-56.

706 Rossant, J., and J. C. Cross. 2001. 'Placental development: lessons from mouse mutants', *Nat*
707 *Rev Genet*, 2: 538-48.

708 Salvatore, C. M., J. Y. Han, K. P. Acker, P. Tiwari, J. Jin, M. Brandler, C. Cangemi, L. Gordon,
709 A. Parow, J. DiPace, and P. DeLaMora. 2020. 'Neonatal management and outcomes
710 during the COVID-19 pandemic: an observation cohort study', *Lancet Child Adolesc*
711 *Health*, 4: 721-27.

712 Schneider, W. M., J. M. Luna, H. H. Hoffmann, F. J. Sánchez-Rivera, A. A. Leal, A. W.
713 Ashbrook, J. Le Pen, I. Ricardo-Lax, E. Michailidis, A. Peace, A. F. Stenzel, S. W. Lowe,

M. R. MacDonald, C. M. Rice, and J. T. Poirier. 2021. 'Genome-Scale Identification of

SARS-CoV-2 and Pan-coronavirus Host Factor Networks', *Cell*, 184: 120-32.e14.

Schwartz, D. A. 2020. 'An Analysis of 38 Pregnant Women with COVID-19, Their Newborn

Infants, and Maternal-Fetal Transmission of SARS-CoV-2: Maternal Coronavirus

Infections and Pregnancy Outcomes', *Arch Pathol Lab Med*.

Schwartz, D. A., M. Baldewijns, A. Benachi, M. Bugatti, R. R. J. Collins, D. De Luca, F.

Facchetti, R. L. Linn, L. Marcelis, D. Morotti, R. Morotti, W. T. Parks, L. Patanè, S.

Prevot, B. Pulinx, V. Rajaram, D. Strybol, K. Thomas, and A. J. Vivanti. 2021. 'Chronic

Histiocytic Intervillositis With Trophoblast Necrosis Is a Risk Factor Associated With

Placental Infection From Coronavirus Disease 2019 (COVID-19) and Intrauterine

Maternal-Fetal Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2)

Transmission in Live-Born and Stillborn Infants', *Arch Pathol Lab Med*, 145: 517-28.

Shanes, E. D., L. B. Mithal, S. Otero, H. A. Azad, E. S. Miller, and J. A. Goldstein. 2020.

'Placental Pathology in COVID-19', *Am J Clin Pathol*, 154: 23-32.

Shang, J., Y. Wan, C. Luo, G. Ye, Q. Geng, A. Auerbach, and F. Li. 2020. 'Cell entry

mechanisms of SARS-CoV-2', *Proc Natl Acad Sci U S A*.

Singh, M., V. Bansal, and C. Feschotte. 2020. 'A Single-Cell RNA Expression Map of Human

Coronavirus Entry Factors', *Cell Rep*, 32: 108175.

Tada, T., C. Fan, J. S. Chen, R. Kaur, K. A. Stapleford, H. Gristick, B. M. Dcosta, C. B. Wilen,

C. M. Nimigean, and N. R. Landau. 2020. 'An ACE2 Microbody Containing a Single

Immunoglobulin Fc Domain Is a Potent Inhibitor of SARS-CoV-2', *Cell Rep*, 33: 108528.

Taglauer, E., Y. Benarroch, K. Rop, E. Barnett, V. Sabharwal, C. Yarrington, and E. M.

Wachman. 2020. 'Consistent localization of SARS-CoV-2 spike glycoprotein and ACE2

over TMPRSS2 predominance in placental villi of 15 COVID-19 positive maternal-fetal

dyads', *Placenta*, 100: 69-74.

Verma, S., C. S. Joshi, R. B. Silverstein, M. He, E. B. Carter, and I. U. Mysorekar. 2021. 'SARS-

CoV-2 colonization of maternal and fetal cells of the human placenta promotes alteration

of local renin-angiotensin system', *Med (N Y)*, 2: 575-90.e5.

Vivanti, A. J., C. Vauloup-Fellous, S. Prevot, V. Zupan, C. Suffee, J. Do Cao, A. Benachi, and D.

De Luca. 2020. 'Transplacental transmission of SARS-CoV-2 infection', *Nat Commun*,

11: 3572.

Wang, R., C. R. Simoneau, J. Kulsuptrakul, M. Bouhaddou, K. A. Travisano, J. M. Hayashi, J.

Carlson-Stevermer, J. R. Zengel, C. M. Richards, P. Fozouni, J. Oki, L. Rodriguez, B.

Joehnk, K. Walcott, K. Holden, A. Sil, J. E. Carette, N. J. Krogan, M. Ott, and A. S.

Puschnik. 2021. 'Genetic Screens Identify Host Factors for SARS-CoV-2 and Common
Cold Coronaviruses', *Cell*, 184: 106-19.e14.

Wei, J., M. M. Alfajaro, P. C. DeWeirdt, R. E. Hanna, W. J. Lu-Culligan, W. L. Cai, M. S. Strine,
S. M. Zhang, V. R. Graziano, C. O. Schmitz, J. S. Chen, M. C. Mankowski, R. B. Filler,
N. G. Ravindra, V. Gasque, F. J. de Miguel, A. Patil, H. Chen, K. Y. Oguntuyo, L.
Abriola, Y. V. Surovtseva, R. C. Orchard, B. Lee, B. D. Lindenbach, K. Politi, D. van Dijk,
C. Kadoch, M. D. Simon, Q. Yan, J. G. Doench, and C. B. Wilen. 2021. 'Genome-wide
CRISPR Screens Reveal Host Factors Critical for SARS-CoV-2 Infection', *Cell*, 184: 76-
91.e13.

Woodworth, K. R., E. O. Olsen, V. Neelam, E. L. Lewis, R. R. Galang, T. Oduyebo, K. Aveni, M.
M. Yazdy, E. Harvey, N. D. Longcore, J. Barton, C. Fussman, S. Siebman, M. Lush, P.
H. Patrick, U. A. Halai, M. Valencia-Prado, L. Orkis, S. Sowunmi, L. Schlosser, S.
Khuwaja, J. S. Read, A. J. Hall, D. Meaney-Delman, S. R. Ellington, S. M. Gilboa, and V.
T. Tong. 2020. 'Birth and Infant Outcomes Following Laboratory-Confirmed SARS-CoV-2
Infection in Pregnancy - SET-NET, 16 Jurisdictions, March 29-October 14, 2020',
MMWR Morb Mortal Wkly Rep, 69: 1635-40.

Yang, L., Y. Han, B. E. Nilsson-Payant, V. Gupta, P. Wang, X. Duan, X. Tang, J. Zhu, Z. Zhao,
F. Jaffré, T. Zhang, T. W. Kim, O. Harschnitz, D. Redmond, S. Houghton, C. Liu, A. Naji,
G. Ciceri, S. Guttikonda, Y. Bram, D. T. Nguyen, M. Cioffi, V. Chandar, D. A. Hoagland,
Y. Huang, J. Xiang, H. Wang, D. Lyden, A. Borczuk, H. J. Chen, L. Studer, F. C. Pan, D.
D. Ho, B. R. tenOever, T. Evans, R. E. Schwartz, and S. Chen. 2020. 'A Human
Pluripotent Stem Cell-based Platform to Study SARS-CoV-2 Tropism and Model Virus
Infection in Human Cells and Organoids', *Cell Stem Cell*, 27: 125-36.e7.

Zambrano, L. D., S. Ellington, P. Strid, R. R. Galang, T. Oduyebo, V. T. Tong, K. R. Woodworth,
J. F. Nahabedian, 3rd, E. Azziz-Baumgartner, S. M. Gilboa, and D. Meaney-Delman.
2020. 'Update: Characteristics of Symptomatic Women of Reproductive Age with
Laboratory-Confirmed SARS-CoV-2 Infection by Pregnancy Status - United States,
January 22-October 3, 2020', *MMWR Morb Mortal Wkly Rep*, 69: 1641-47.

Figure legends

<u>Figure 1.</u> **SARS-CoV-2 virus is present in placentas from infected mothers and results in**

inflammatory responses. (A) Graph showing ΔΔCT values of RNA samples isolated from

FFPE patient placenta slides from the different cohorts. A student's t-test comparing the 3

positive cohorts (High Positive, Positive, Borderline Positive) to the negative control cohort

resulted in statistically significant higher viral load in the High Positive and Positive cohorts (***

= p-value < 0.001). (B) Brightfield microscopy images of a representative COVID High positive

patient (P3) and a representative negative control patient sample (C1). Slides were stained by

H&E, *in situ* PLAYR for SARS-CoV2-RNA counterstained for syncytial trophoblast marker

cytokeratin (KRT7, red), and by immunohistochemistry for SARS-CoV2-N protein (brown) as

well as for CD163+ Hofbauer cells (HBC). Scale bars = 100µm.

<u>Figure 2.</u> **Placental explant and cell clusters can be infected by SARS-CoV-2 S**

protein pseudotyped lentivirus and infection can be blocked by anti-ACE2 and anti-NRP1

antibodies. (A) Graphs showing relative luminescence units (RLU) from infected explant

cultures 72 hpi with the addition of reverse transcriptase inhibitor, Nevirapine (NVP). (B) Graphs

showing RLU from infected isolated primary placental clusters 72 hpi with the addition of

blocking antibodies against ACE2, NRP1. Statistical analysis was performed using one-way

Anova (** = p-value < 0.005, *** = p-value < 0.001). (C) Brightfield and live fluorescence

microscopy images of cultured placental explants Mock (left column), or 72hpi with either Lenti-

SARS-CoV2-S Pseudovirus (center column) or Lenti-VSV-G (right column). (D) Fluorescence

microscopy on mock (top row) and Lenti-SARS-CoV2-S infected (center row) or Lenti-VSV-G

infected (bottom row) explant sections stained for the GFP reporter (green) syncytial trophoblast

marker, cytokeratin (KRT7, grey), endothelial marker CD31 (red) and DAPI nuclear stain (blue).

Scale bars = 500µm.

<u>Figure 3</u>. **Primary human placenta cells can be infected with SARS-CoV-2** *ex*

vivo. (A) qRT-PCR analysis of relative viral N subgenomic RNA expression in primary placental

cell clusters infected with SARS-CoV-2 *ex vivo* (MOI=1) for 24 hours and normalized to ACTB

levels. (mean+/- SD; n=12 from 4 repeated experiments; ****$p<0.0001$) . (B) Three-dimensional

reconstruction of confocal imaging of primary placental cell clusters infected with SARS-CoV-

2 *ex vivo* (MOI=1) at 24hpi, stained for trophoblast marker KRT7 (green), SARS-N (red),

endothelial marker CD31 (grey), and DAPI (blue). Scale bar = 30µm. (C) Confocal imaging of

primary placental cell clusters infected with MOCK (top rows) or SARS-CoV-2 (MOI=1, bottom

rows) *ex vivo* at 24hpi, stained for trophoblast marker KRT7 (green), SARS-N (red), endothelial

marker CD31 (grey), and DAPI (blue). Arrows indicate presence of SARS-N nucleocapsid

protein in trophoblast and endothelial cells. Scale bar = 20µm.

Table 1. Clinical presentations of SARS-CoV-2 positive mothers, fetal outcomes and placental

pathologies.

Overview of 65 Patients included in study. 55 COVID-positive, 10 COVID-negative.

Abbreviations: FVM: Fetal Vascular Malperfusion, MVM: Maternal Vascular Malperfusion, DFM:

Decreased Fetal Movement, MCI: Massive Chronic Intervillositis, MFI: Maternal Floor Infarction,

CHI: Chronic Histiocytic Intervillositis, IUFD: Intra-Uterine Fetal Demise, T2D: Type 2 Diabetes,

Mec: Meconium, IVT: Intervillous Thrombi, VUE: Villositis of Unknown Etiology/Chronic

Villositis, ACA: Acute Chorioamniotis, IAI: Intra-Amniotic Infection/Chorioamnioitis, HTN:

Hypertension, IUGR: Intra-Uterine Growth Restriction, GDM: Gestational Diabetes Mellitus,

PPROM: Preterm Premature Rupture of Membranes, PTL: Preterm Labor, PAPP-A: Pregnancy-

associated Plasma Protein A, UCTD: Undifferentiated Connective Tissue Disorder, HCV:

Hepatitis C Virus, ITP: Immune Thrombocytopenic Purpura. IUGR: intrauterine growth

restriction. ICP: intrahepatic cholestasis of pregnancy. Gray Shaded Rows = Fetal Demise/NICU

admission.

Table 1

	Case	Mat Age	Gest Age	Mother COVID +/-	Patient History	Birth weight	Apgar 1 min	Apgar 5 min	FVM	MVM	Other pathology	Placenta Viral RNA
High Positive	P1	35	25	+	Symptomatic COVID-19, DFM, Delivered due to nonreassuring fetal status	650	1	8	-	-	MCI	+
	P2	34	30	+	Symptomatic COVID-19, DFM, IUFD	1389	0	0	-	-	MFI, CHI	+
Positive	P3	29	40	+	Symptomatic COVID-19	3400	9	9	+	-	-	+
	P4	40	36	+	T2D (poorly controlled), Placenta previa	2680	9	9	-	-	-	+
	P5	16	32	+	Symptomatic COVID-19, PTL	1740	9	8	-	+		+
	P6	26	37	+	Asymptomatic COVID-19, T2D (poorly controlled), DFM, IUFD	3200	0	0	-	-	Villous Dysmaturity	+
	P7	31	39	+	Symptomatic COVID-19	3140	9	9	-	-	Mec, IVT	+
	P8	30	38	+	Symptomatic COVID-19	3910	9	9	+	-	VUE, Mec	+
	P9	41	39	+	Asymptomatic COVID-19	3770	9	9	-	-	VUE	+
	P10	28	39	+	Asymptomatic COVID-19	3300	9	9	-	-	Mec	+
	P11	34	37	+	ICP	2900	9	9	-	-	-	+
	P12	31	40	+	Asymptomatic COVID-19	3340	9	9	-	-	Mec	+
Borderline (+)	P13	30	38	+	Symptomatic COVID-19, Intrapartum chorioamnionitis	3360	9	9	+	-	Mec, Furcate cord	+
	P14	26	38	+	Asymptomatic COVID-19	3050	9	9	+	-	-	+
	P15	19	38	+	Symptomatic COVID-19	2390	9	9	+	+	ACA	+
	P16	28	40	+	Asymptomatic COVID-19	3820	9	9	+	-	Mec	+
	P17	28	41	+	Symptomatic COVID-19	4020	8	9	+	-	VUE, IDA, Mec	+
	P18	41	40	+	Symptomatic COVID-19	4115	9	9	+	-	Mec	+
	P19	32	38	+	Asymptomatic COVID-19	3160	9	9	-	-	Twisted Cord	+
	P20	26	39	+	Symptomatic COVID-19	3720	9	9	+	-	Hofbauer hyperplasia	+
	P21	20	38	+	Symptomatic COVID-19	3685	6	9	+	-	IAI, Mec	+
	P22	25	39	+	Asymptomatic COVID-19	3000	9	9	-	-	Villitis	+
Negative Samples	P23	40	39	+	Asymptomatic COVID-19	3720	9	9	+	-	-	-
	P24	40	37	+	Symptomatic COVID-19, PEC	2060	8	9	-	+	Mec	-
	P25	38	39	+	Symptomatic COVID-19, ITP, Protein S deficiency, Gestational HTN	3890	9	9	+	-	Funisitis	-
	P26	26	40	+	Chronic HTN, Symptomatic COVID-19	3799	9	9	+	-	-	-
	P27	37	39	+	Asymptomatic COVID-19. Autoimmune gastritis	2415	9	9	-	+	-	-
	P28	40	33	+	Symptomatic COVID-19, IUGR, PEC, Delivered for nonreassuring fetal status	1690	8	8	-	+	Mec	-
	P29	33	35	+	Symptomatic COVID-19, Dichorionic twins, PEC	2280 (A) / 2810 (B)	8 (A) / 8 (B)	9 (A) / 9 (B)	+	+	VUE	-
	P30	23	39	+	Asymptomatic COVID-19	3580	8	9	-	-	Villitis	-
	P31	25	38	+	Asymptomatic COVID-19	3920	9	9	-	-	Mec	-
	P32	34	39	+	Symptomatic COVID-19	3360	9	9	-	-	Mec	-
	P33	40	37	+	Asymptomatic COVID-19	3400	8	9	-	-	-	-
	P34	37	41	+	Symptomatic COVID-19	3900	9	9	-	-	IAI, Mec	-
	P35	39	37	+	Asymptomatic COVID-19	2650	9	9	-	-	Villous dysmaturity	-
	P36	34	39	+	Asymptomatic COVID-19	2510	9	9	-	+	-	-
	P37	33	23	+	COVID-19 remote from delivery, fetal anencephaly, intrapartum fetal demise	370	0	0	-	-	-	-
	P38	30	39	+	Asymptomatic COVID-19	3910	9	9	+	-	Villitis	-
	P39	31	40	+	Symptomatic COVID-19 remote from delivery	3200	8	9	-	-	-	-
	P40	30	39	+	Symptomatic COVID-19 remote from delivery	3650	9	9	+	-	Chorionic cysts, IAI, Mec	-
	P41	27	41	+	Symptomatic COVID-19, Low PAPP-A, Gestational HTN	3630	8	9	-	-	Focal chorangiosis	-
	P42	23	37	+	Symptomatic COVID-19 remote from delivery	2510	9	9	-	-	IVT	-
	P43	31	36	+	ICP, Asymptomatic COVID-19	3290	9	9	-	-	IVT, Mec	-
	P44	29	37	+	Symptomatic COVID-19, low PAPP-A	2350	8	9	-	+	-	-
	P45	29	36	+	Genetic carrier for hearing loss	2510	9	9	-	-	-	-
	P46	40	38	+	Symptomatic COVID-19 remote from delivery	2820	9	9	-	+	-	-
	P47	32	40	+	Symptomatic COVID-19 remote from delivery	3360	9	9	-	+	VUE	-
	P48	51	37	+	Symptomatic COVID-19 remote from delivery, T2D	3080	9	10	-	-	VUE	-
	P49	41	38	+	Symptomatic COVID-19 remote from delivery, Asthma	2990	8	9	+	+	-	-
	P50	38	39	+	Symptomatic COVID-19 remote from delivery	3010	9	9	-	-	VUE, IVT	-
	P51	38	39	+	Symptomatic COVID-19 remote from delivery	3840	9	9	-	-	-	-
	P52	33	38	+	Symptomatic COVID-19, Long QT syndrome, remote from delivery	3005	9	9	-	-	-	-
	P53	38	36	+	Symptomatic COVID-19 remote from delivery, Dichorionic twins, PTL	2680 (A) / 2740 (B)	9 (A) / 9 (B)	9 (A) / 9 (B)	-	-	-	-
	P54	35	39	+	Symptomatic COVID-19 remote from delivery	2870	9	9	+	+	Villitis	-
Negative Controls	C1	32	40	-	Subglottic stenosis	NA	9	9	-	-	Mec	-
	C2	29	39	-	Low PAPP-A, UCTD, celiac disease	3470	9	9	-	-	Mec	-
	C3	39	34	-	PPROM	2320	9	9	-	-	IVT	-
	C4	31	40	-	COVID-19 in first trimester	3392	9	9	-	-	IVT	-
	C5	36	39	-	Gestational HTN, GDM	3277	9	9	-	-	ACA	-
Inflammatory Samples	I1	32	38	-	Intrapartum chorioamnionitis	3145	9	9	-	-	ACA, Acute funisitis, Mec	-
	I2	28	40	-	No medical history	3100	9	9	-	-	VUE, ACA	-
	I3	33	38	-	Opioid use disorder, HCV, Placental abruption	2664	9	9	+	-	VUE, Acute funisitis	-
	I4	42	38	-	Gestational HTN, GDM	2891	9	9	-	-	ACA, Acute funisitis	-
	I5	34	39	-	ITP	3447	7	9	-	+	ACA, Acute funisitis, Mec	-

Table 1. Clinical presentations of SARS-CoV-2 positive mothers, fetal outcomes and placental pathologies. DFM: Decreased Fetal Movement, MCI: Massive Chronic Intervillositis, MFI: Maternal Floor Infarction, CHI: Chronic Histiocytic Intervillositis, IUFD: Intra-Uterine Fetal Demise, T2D: Type 2 Diabetes, Mec: Meconium, IVT: Intervillous Thrombi, VUE: Villositis of Unknown Etiology/ Chronic Villositis, ICP: Intrahepatic Cholestasis of Pregnancy, GBS: Group B Streptococcus+ , TOLAC: Trial of Labor After Cesarean, ACA: Acute Chorioamnionitis, BMI: Body Mass Index, TAB: Therapeutic Abortion, IDA: Iron Deficiency Anemia, IAI: Intra-Amniotic Infection, PPH: Post-Partum Hemorrhage, HTN: Hypertension, PEC: Preeclampsia SF: Severe Features, PCS: Pelvic Congestion Syndrome, NI: Class I No signs or symptoms, Di/Di: Dichorionic/Diamniotic, D&C: Dilation & Curettage, GDM: Gestational Diabetes Mellitus, PROM: Premature Rupture of Membranes, PTL: Pret-term Labor, PAPP-A: Pregnancy-associated Plasma Protein A, UCTD: Undifferentiated Connective Tissue Disorder, PIH: Pregnancy-Induced/Gestational Hypertension, HCV: Hepatitis C Virus+, ITP: Immune Thrombocytic Purpura

Figure 1

A

ΔΔCT Values

- High Positive
- Positive
- Borderline Positive
- Negative Controls
- Inflammatory Negative
- Positive Control (Lung)

B

	H & E	SARS-CoV-2 RNA/KRT7	SARS-CoV2-N	CD163+ HBC
High Positive (P2)				
Negative Control (C1)				

Figure 2