Chromosome-level *de novo* genome assembly of *Telopea speciosissima* (New South Wales waratah) using long-reads, linked-reads and Hi-C

Running title: A reference genome for waratah (Proteaceae)

Stephanie H Chen[1,2], Maurizio Rossetto[2,3], Marlien van der Merwe[2], Patricia Lu-Irving[2], Jia-Yee S Yap[2,3], Hervé Sauquet[4,5], Greg Bourke[6], Timothy G Amos[1], Jason G Bragg[2,5,✉], Richard J Edwards[1,✉]

[1]School of Biotechnology and Biomolecular Sciences, UNSW Sydney, High St, Kensington, NSW 2052, Australia

stephanie.h.chen@unsw.edu.au, t.amos@garvan.org.au, richard.edwards@unsw.edu.au

[2]Research Centre for Ecosystem Resilience, Australian Institute of Botanical Science, The Royal Botanic Garden Sydney, Mrs Macquaries Rd, Sydney, NSW 2000, Australia

maurizio.rossetto@botanicgardens.nsw.gov.au, marlien.vandermerwe@botanicgardens.nsw.gov.au, patricia.lu-irving@botanicgardens.nsw.gov.au, samantha.yap@botanicgardens.nsw.gov.au, jason.bragg@botanicgardens.nsw.gov.au

[3]Queensland Alliance of Agriculture and Food Innovation, University of Queensland, St Lucia 4072, Australia

[4]National Herbarium of New South Wales, Royal Botanic Gardens and Domain Trust, Mrs Macquaries Rd, Sydney, NSW 2000, Australia

herve.sauquet@botanicgardens.nsw.gov.au

[5]School of Biological, Earth and Environmental Sciences, UNSW Sydney, High St, Kensington, NSW 2052, Australia

[6]Blue Mountains Botanic Garden, Bells Line of Road, Mount Tomah, NSW 2758, Australia

greg.bourke@botanicgardens.nsw.gov.au

✉ Corresponding authors

ORCID iD

28   SHC 0000-0001-8844-6864

MR 0000-0002-4878-9114

30   MVDM 0000-0003-1307-5143

PL-I 0000-0003-1116-9402

32   JSY 0000-0002-9141-6006

HS 0000-0001-8305-3236

34   TGA 0000-0002-5829-6655

JGB 0000-0002-7621-7295

36   RJE 0000-0002-3645-5539

38   ABSTRACT

40   *Telopea speciosissima,* the New South Wales waratah, is an Australian endemic woody shrub

in the family Proteaceae. Waratahs have great potential as a model clade to better

42   understand processes of speciation, introgression and adaptation, and are significant from a

horticultural perspective. Here, we report the first chromosome-level genome for *T.*

44   *speciosissima*. Combining Oxford Nanopore long-reads, 10x Genomics Chromium linked-

reads and Hi-C data, the assembly spans 823 Mb (scaffold N50 of 69.0 Mb) with 97.8 % of

46   Embryophyta BUSCOs complete. We present a new method in Diploidocus

(https://github.com/slimsuite/diploidocus) for classifying, curating and QC-filtering scaffolds,

48   which combines read depths, k-mer frequencies and BUSCO predictions. We also present a

new tool, DepthSizer (https://github.com/slimsuite/depthsizer), for genome size estimation

50   from the read depth of single copy orthologues and estimate the genome size to be

approximately 900 Mb. The largest 11 scaffolds contained 94.1 % of the assembly,

52   conforming to the expected number of chromosomes ($2n$ = 22). Genome annotation

predicted 40,158 protein-coding genes, 351 rRNAs and 728 tRNAs. We investigated

54   *CYCLOIDEA* (*CYC*) genes, which have a role in determination of floral symmetry, and confirm

the presence of two copies in the genome. Read depth analysis of 180 'Duplicated' BUSCO

56   genes suggest almost all are real duplications, increasing confidence in protein family

analysis using annotated protein-coding genes, and highlighting a possible need to revise the

58   BUSCO set for this lineage. The chromosome-level *T. speciosissima* reference genome

(Tspe_v1) provides an important new genomic resource of Proteaceae to support the

60   conservation of flora in Australia and further afield.

62   **Keywords**: Telopea, waratah, genome assembly, reference genome, long-read sequencing,

Hi-C

INTRODUCTION

Telopea R.Br. is an eastern Australian genus of five species of large, long-lived shrubs in the

68   flowering plant family Proteaceae. The New South Wales waratah, *Telopea speciosissima*

(Sm.) R.Br., is a striking and iconic member of the Australian flora, characterised by large,

70   terminal inflorescences of red flowers (Figure 1). It has been the state floral emblem of New

South Wales since 1962 and was one of the first Australian plant species collected for

72 cultivation in Europe (Nixon, 1987). The species is endemic to the state of New South Wales,

occurring on sandstone ridges in the Sydney region. Previous studies have investigated

74 variation among *Telopea* populations by phenetic analysis of morphology (Crisp & Weston,

1993) and evolutionary relationships using cladistics (Weston & Crisp, 1994). Population

76 structure and patterns of divergence and introgression between *T. speciosissima* populations

have been characterised using several loci (Rossetto et al., 2011). Further, microsatellite

78 data and modelling suggest a history of allopatric speciation followed by secondary contact

and hybridization among *Telopea* species (Rossetto et al., 2012). These studies point to the

80 great potential of *Telopea* as a model clade for understanding processes of divergence,

environmental adaptation and speciation. Our understanding of these processes can be

82 greatly enhanced by a genome-wide perspective, enabled by a reference genome (Ellegren

et al., 2012; Hoban et al., 2016; Lewin et al., 2018; Radwan & Babik, 2012; Seehausen et al.,

84 2014).

86 Genome sequencing efforts have traditionally focused on model species, crops and their

wild relatives, resulting in a highly uneven species distribution of reference genomes across

88 the plant tree of life (Royal Botanic Gardens, Kew, 2017). Despite Proteaceae occurring

across several continents and encompassing 81 genera and ca. 1700 species (Mast et al.,

90 2008; Weston, 2006), the only publicly available reference genome in the family is a widely-

grown cultivar of the most economically important crop in the family, *Macadamia*

92 *integrifolia* (macadamia nut) HAES 74 (Nock et al., 2016, 2020). Waratahs are significant to

the horticultural and cut flower industries, with blooms cultivated for the domestic and

94 international markets (Offord et al., 1987; Worrall & Gollnow, 2013). A reference genome

will accelerate efforts in breeding for traits such as resistance to pests and diseases (e.g.

96 *Phytophthora* and *Cylindrocapon destructans* infection; Summerell, 1997; Summerell et al.,

1990) as well as desirable floral characteristics (Offord, 2003, 2006).

Technological advances in sequencing and decreasing costs will facilitate the generation of

100 more flowering plant reference genomes, including within the Proteaceae family, and

advance research into links between the evolution of genomes and traits that exhibit

102 exceptional diversity, such as floral morphology (Soltis & Soltis, 2014; Zheng et al., 2021).

CYCLOIDEA (*CYC*) genes belong to the TPC transcription factor gene family, and are known to

104 have an essential role in determining floral symmetry and inflorescence architecture in many

angiosperm lineages (Busch & Zachgo, 2009; Fambrini & Pugliesi, 2017; Horn et al., 2015;

106 Luo et al., 1996); studies have characterised recurrent duplications of members of the *CYC2*

clade, especially in eudicots (Howarth & Donoghue, 2006), including Fabales (Citerne et al.,

108 2003; Feng et al., 2006), Asterales (Chapman et al., 2008), and Lamiales (Yang et al., 2015;

Zhong & Kellogg, 2015). In Proteaceae, a single duplication of *CYC*-like genes occurred prior

110 to diversification and two genes, *ProtCYC1* and *ProtCYC2*, have been characterised (Citerne

et al., 2017). In particular, *Grevillea juniperina* has been studied in detail (Damerval et al.,

112 2019) and the existence of both *ProtCYC1* and *ProtCYC2* in *Telopea mongaensis* has been

supported by phylogenetic analysis (Citerne et al., 2017). However, *CYC* copy number has

114 not been established in *T. speciosissima.*

116 Here, we provide a high quality chromosome-level *de novo* assembly of the *Telopea*

speciosissima genome, using Oxford Nanopore long-reads, 10x Genomics Chromium linked-

118 reads and Hi-C proximity ligation scaffolding, which will serve as an important platform for

evolutionary genomics and the conservation of the Australian flora. We present an analysis

120 of *CYC* genes in the genome to contribute to the understanding of floral evolution in the

Proteaceae family.

MATERIALS AND METHODS

Sampling and DNA extraction

126 Young leaves (approx. 8 g) were sampled from the reference genome individual (NCBI

BioSample SAMN18238110) where it grows naturally along the Tomah Spur Fire Trail (-

128 33.53° S, 150.42° E) on land belonging to the Blue Mountains Botanic Garden, Mount Tomah

in New South Wales, Australia. Leaves were immediately frozen in liquid nitrogen and stored

130 at -80° C prior to extraction.

132 High-molecular-weight (HMW) genomic DNA (gDNA) was obtained using a sorbitol pre-wash

step prior to a CTAB extraction adapted from Inglis et al. (2018). The gDNA was then purified

134 with AMPure XP beads (Beckman Coulter, Brea, CA, USA) using a protocol based on

Schalamun et al. (2019) – details available on protocols.io (Lu-Irving & Rutherford, 2021).

136 The quality of the DNA was assessed using Qubit, NanoDrop and TapeStation 2200 System

(Agilent, Santa Clara, CA, USA).

ONT PromethION sequencing

140   We performed an in-house sequencing test on the MinION (MinION, RRID:SCR_017985)

using a FLO-MINSP6 (R9.4.1) flow cell with a library prepared with the ligation kit (SQK-

142   LSK109). The remaining purified genomic DNA was sent to the Australian Genome Research

Facility (AGRF) where size selection was performed to remove small DNA fragments using

144   the BluePippin High Pass Plus Cassette on the BluePippin (Sage Science, Beverly, MA, USA).

Briefly, 10 µg of DNA was split into 4 aliquots (2.5 µg) and diluted to 60 µL in TE buffer. Then,

146   20 µL of RT equilibrated loading buffer was added to each aliquot and mixed by pipetting.

Samples were loaded on the cassette by removing 80 µL of buffer from each well and adding

148   80 µL of sample or external marker. The cassette was run with the 15 kb High Pass Plus

Marker U1 cassette definition. Size selected fractions (approximately 80 µL) were collected

150   from the elution module following a 30 min electrophoresis run. The library was prepared

with the ligation sequencing kit (SQK-LSK109). The sequencing was performed using

152   MinKNOW v.19.12.2 (MinION) and v12.12.8 (PromethION) and MinKNOW Core v3.6.7 (in-

house test), v3.6.8 (AGRF MinION) and v3.6.7 (AGRF PromethION). A pilot run was first

154   performed on the MinION using the FLO-MIN106 (R9.4.1) flow cell followed by two FLO-

PRO002 flow cells (R9.4) on the PromethION (PromethION, RRID:SCR_017987)

Basecalling was performed after sequencing with GPU-enabled Guppy v3.4.4 using the high-

158   accuracy flip-flop models, resulting in 54x coverage. The output from all ONT basecalling was

pooled for adapter removal using Porechop (Porechop, RRID:SCR_016967) v.0.2.4 (Wick et

160   al., 2017) and quality filtering (removal of reads less than 500 bp in length and Q lower than

7) with NanoFilt (NanoFilt, RRID:SCR_016966) v2.6.0 (De Coster et al., 2018) followed by

162   assessment using FastQC (FastQC, RRID:SCR_014583) v0.11.8 (Andrews, 2010).

164 **10x Genomics Chromium sequencing**

High-molecular-weight gDNA was sent to AGRF for 10x Genomics Chromium sequencing.

166 Size selection was performed to remove DNA fragments <40 kb using the BluePippin 0.75 %

Agarose Gel Cassette, Dye Free on the BluePippin (Sage Science, Beverly, MA, USA). Briefly, 5

168 µg of DNA was diluted to 30 µL in TE buffer and 10 µL of RT equilibrated loading buffer was

added to each aliquot and mixed by pipetting. Samples were loaded on the cassette by

170 removing 40 µL of buffer from each well and adding 40 µL of sample or external marker. The

cassette was run with the 0.75 % DF Marker U1 high-pass 30-40 kb v3 cassette definition.

172 Size selected fractions (approximately 40 µL) were collected following the 30 min

electrophoresis run. The library was prepared using the Chromium Genome Library Kit & Gel

174 Bead Kit and sequenced (2 x 150 bp paired-end) on the NovaSeq 6000 (Illumina NovaSeq

6000 Sequencing System, RRID:SCR_016387) with NovaSeq 6000 SP Reagent Kit (300 cycles)

176 and NovaSeq XP 2-Lane Kit for individual lane loading.

178 **Hi-C sequencing**

Hi-C library preparation and sequencing was conducted at the Ramaciotti Centre for

180 Genomics at the University of New South Wales (UNSW Sydney) using the Phase Genomics

Plant kit v3.0. The library was assessed using Qubit and the Agilent 2200 TapeStation system

182 (Agilent Technologies, Mulgrave, VIC, Australia). A pilot run on an Illumina iSeq 100 with 2 x

150 bp paired end sequencing run was performed for QC using hic_qc v1.0 (Phase Genomics,

184 2019) with i1 300 cycle chemistry. This was followed by sequencing on the Illumina NextSeq

500 (Illumina NextSeq 500, RRID:SCR_014983) with 2 x 150 bp paired-end high output run

186 and NextSeq High Output 300 cycle kit v2.5 chemistry.

188 **Genome assembly and validation**

Our assembly workflow consisted of assembling a draft long-read assembly, hybrid polishing

190 of the assembly with long- and short-reads, and scaffolding the assembly into chromosomes

using Hi-C data (Figure 2). Computational steps were carried out on the UNSW Sydney

192 cluster Katana.

194 The first stage of our assembly approach involved comparing three long-read assemblers

using the ONT data as input: NECAT v0.01 (Chen et al., 2021), Flye (Flye, RRID:SCR_017016)

196 v2.6 (Kolmogorov et al., 2019) and Canu (Canu, RRID:SCR_015880) v1.9 (Koren et al., 2017).

The genome size parameter used for the assemblers was 1,134 Mb, as previously reported

198 for *Telopea truncata* (Jordan et al., 2015). We later refined genome size estimates for *T.*

speciosissima (see 'DepthSizer: genome size estimation using single-copy orthologue

200 sequencing depths' section below). We chose the draft long-read assembly for use in

downstream assembly steps based on contiguity (N50), BUSCO completeness and assembly

202 size in relation to the DepthSizer estimated genome size. As a comparison to the long-read

assemblies, the 10x data were assembled with Supernova (Supernova assembler,

204 RRID:SCR_016756) v2.1.1 (Weisenfeld et al., 2017) with 332 Mb reads subsampled by

Supernova (54x raw coverage, as recommended by Supernova documentation) as input. We

206 generated pseudohaploid output (pseudohap2 output '1').

Assembly completeness and accuracy

Completeness was initially evaluated by BUSCO (BUSCO, RRID:SCR_015008) v3.0.2b (Simão et al., 2015), implementing BLAST+ v2.2.31, Hmmer v3.2.1 and EMBOSS v6.6.0 with the embryophyta_odb9 dataset (n = 1,440). To investigate the robustness of BUSCO completeness statistics, assemblies were also evaluated with BUSCO v5.0.0 (Manni et al., 2021), implementing BLAST+ v2.11.0 (Altschul et al., 1990), SEPP v4.3.10 (Mirarab et al., 2011) and Hmmer (Hmmer, RRID:SCR_005305) v3.3 (Eddy, 2011), against the embryophyta_odb10 dataset (n = 1,614). BUSCO results were calculated with both Augustus (Augustus, RRID:SCR_008417) v3.3.2 (Stanke & Morgenstern, 2005) and MetaEuk v732bcc4b91a08e69950ce0e25976f47c3bb6b89d (Levy Karin et al., 2020) as the gene predictor.

BUSCO results were collated using BUSCOMP (BUSCO Compilation and Comparison Tool; RRID:SCR_021233) v0.11.0 (Stuart et al., 2021) to better evaluate the gains and losses in completeness between different assembly stages, and compare different BUSCO versions. Assembly quality (QV) was also estimated using k-mer analysis of trimmed and filtered 10x linked-read data by Merqury v1.0 with k = 20 (Rhie et al., 2020). First, 30 bp from the 5' end of read 1 and 10 bp from the 5' end of read 2 were trimmed using BBmap (BBmap, RRID:SCR_016965) v38.51 (Bushnell, 2014). In addition, reads were trimmed to Q20, then those shorter than 100 bp were discarded.

Genome size estimation and ploidy

230   *Telopea speciosissima* has been reported as a diploid (2*n* = 22) (Darlington & Wylie, 1956;

Ramsay, 1963). We confirmed the individual's diploid status using Smudgeplot v0.2.1

232   (Ranallo-Benavidez et al., 2019). The 1C-value of *T. truncata* (Tasmanian waratah) has been

estimated at 1.16 pg (1.13 Gb) using flow cytometry (Jordan et al., 2015). We used the 10x

234   data to estimate the genome size using Supernova v2.1.1 and GenomeScope

(GenomeScope, RRID:SCR_017014) v1.0 (Vurture et al., 2017).

We sought to refine the genome size estimate of *T. speciosissima* using the ONT data and

238   draft genome assemblies, implementing a new tool, DepthSizer

(https://github.com/slimsuite/depthsizer, RRID:SCR_021232, **Box 1**). ONT reads were

240   mapped onto each draft genome using Minimap2 (Minimap2, RRID:SCR_018550) v2.17 (Li,

2018) (--secondary=no -ax map-ont). The single-copy read depth for each assembly was then

242   calculated as the modal read depth across single copy complete BUSCO genes, which should

be reasonably robust to poor-quality and/or repeat regions within these genes (Edwards et

244   al., 2021).

246   **DepthSizer benchmarking**

DepthSizer was benchmarking using five PacBio reference genomes, plus the high-quality

248   genome assembly and PacBio long reads for the German Shepherd Dog (Field et al., 2020;

Table S1). Accuracy was calculated as the estimated genome size, divided by the

250   documented genome size. Additional benchmarking of the robustness of DepthSizer

predictions was performed using ONT and PacBio sequence data for three high-quality dog

252   genomes: Basenji (Edwards et al., 2021), Dingo (Yadav et al., 2020), and German Shepherd

Dog (Field et al., 2020). Raw reads from each technology were analysed independently using

254   both the breed-specific reference genome, and the CanFam 3.1 dog reference

(GCA_000002285.2; Lindblad-Toh et al., 2005). For all benchmarking, reads were mapped

256   with Minimap2 (Minimap2, RRID:SCR_018550) v2.17 (Li, 2018). Summary violin plots were

generated with ggstatsplot (Patil, 2021) in R.

Box 1. DepthSizer: genome size estimation using single-copy orthologue sequencing depths

GitHub: https://github.com/slimsuite/depthsizer

Genome size prediction is a fundamental task in genome assembly. DepthSizer is a tool for estimating genome size using single-copy long-read sequencing depth profiles.

By definition, sequencing depth (X) is the volume of sequencing divided by the genome size. Given a known volume of sequencing, it is therefore possible to estimate the genome size by estimating the achieved sequencing depth. DepthSizer works on the principle that the modal read depth across single copy BUSCO genes provides a good estimate of the true depth of coverage. This assumes that genuine single copy depth regions will tend towards the same, true, single copy read depth. In contrast, assembly errors or collapsed repeats within those genes, or incorrectly-assigned single copy genes, will give inconsistent read depth deviations from the true single copy depth. The exception is regions of the genome only found on one haplotig – half-depth alternative haplotypes for regions also found in the main assembly –

---

such as heterogametic sex chromosomes (Edwards et al., 2021), but these are unlikely to outnumber genes present in single copy on both homologous chromosomes. As a consequence, the dominant (i.e. modal) depth across these regions should represent single copy (2n) sequencing depth. First, the distribution of read depth for all single copy genes is generated using Samtools (Samtools, RRID:SCR_002105) v0.11 (Li et al., 2009) mpileup, and the modal peak calculated using a smoothed 'density' function of R (R Project for Statistical Computing, RRID:SCR_001905) v3.5.3 (R Core Team, 2019) to allow non-integer estimation (see DepthSizer documentation for details). Genome size, G, was then estimated from the modal peak single-copy depth, X_{sc}, and the total volume of sequencing data, T, using the formula: $G = T / X_{SC}$.

DepthSizer has six different genome size adjustment modes that modify T using different core assumptions (see documentation for details):

- None: no adjustment. Assumes zero contamination and perfect read mapping.

- IndelRatio: adjusts total sequencing volume for mismatch between read data being mapped and assembly coverage. Assumes no contamination in raw reads.

- CovBases: sets T as the total number of sequencing read bases covering the assembly. (Assembly Length x Mean depth)

- MapBases: sets T as the total number bases from sequencing reads mapped on to the genome. Assumes perfect mapping and all unmapped reads are contamination.

- MapAdjust: adjusts total sequencing volume by the ratio of mapped reads to mapped bases to account for depth losses during mapping. Assumes no contamination in raw reads.

- MapRatio: adjusts the MapBases by the IndelRatio sequencing:mapping bias.

It is expected that the true genome size should fall between IndelRatio (upper) and MapRatio (lower). CovBases should provide an absolute lower bound for genome size. If there is a very large difference between CovBases and MapBases, this could indicate a problem with the reads and/or assembly (e.g. some kind of incompatibility) and will result in a very inaccurate MapAdjust. If there is a very big difference between MapBases and None, this could indicate a very incomplete assembly, or a lot of contamination. In these cases, it is advisable to establish which before deciding which prediction size to use.

Benchmarking on PacBio data from six model organisms demonstrates robust genome size estimates, with a tendency to slightly overestimate genome size as expected (Figure 3, Table S1 and Table S2). Additional benchmarking on three high-quality canid genomes further revealed robustness to both assembly used (breed-specific genome versus CanFam v3.1) and sequencing technology (PacBio vs ONT), although PacBio data appears to over-estimate genome size more than ONT data (Figure S1).

Figure 3. Model organism benchmarking of DepthSizer with six core methods using Minimap2 read mapping and Mpileup depth calculations with a) accuracy for each species and b) estimated genome size vs published genome size.

Assembly tidying and contamination screening

The draft genome was screened and filtered to remove contamination, low-quality contigs

and putative haplotigs, using more rigorous refinement of the approach taken for the

Canfam_GSD (German Shepherd) and CanFam_Bas (Basenji) dog reference genomes

(Edwards et al., 2021; Field et al., 2020), implemented in Diploidocus v0.9.6

(https://github.com/slimsuite/diploidocus, RRID:SCR_021231, **Box 2**).

BUSCO Complete genes were used to estimate a single-copy read depth of 54X. This was

used to set low-, mid- and high-depth thresholds for Purge Haplotigs (Purge_haplotigs,

RRID:SCR_017616) v20190612 (Roach et al., 2018) (implementing Perl v5.28.0, BEDTools

(BEDTools, RRID:SCR_006646) v2.27.1 (Quinlan & Hall, 2010), R v3.5.3 (R Core Team, 2019),

and SAMTools v1.9 (Li et al., 2009) of 13X, 40X and 108X. For the draft genome, convergence

was reached after three cycles with 148 core sequences and 62 repeat sequences retained

(see Table S6 for summary of cycles and Table S7 for full output).

Box 2. Automated genome assembly tidying with Diploidocus

GitHub: https://github.com/slimsuite/diploidocus

Diploidocus is a tool that assists with tidying and curating genome assemblies. The tool

combines read depth, KAT k-mer frequencies, Purge Haplotigs depth bins, Purge Haplotigs

best sequence hits, BUSCO gene predictions, telomere prediction and vector contamination

---

into a single seven-part (PURITY|DEPTH|HOM|TOP|MEDK|BUSCO+EXTRA) classification (Table S4). Diploidocus then performs a hierarchical rating of scaffolds, based on their classifications and compiled data (Table S5 and Figure 4). Based on these ratings, sequences are divided into sets:

1. Core. Predominantly diploid scaffolds and unique haploid scaffolds with insufficient evidence for removal.

2. Repeats. Unique haploid scaffolds with insufficient evidence for removal but dominated by repetitive sequences. High coverage scaffolds representing putative collapsed repeats.

3. Quarantine. Messy repetitive sequences and strong candidates for alternative haplotigs.

4. Junk. Low coverage, short and/or high-contaminated sequences.

If any sequences are marked as 'Quarantine' or 'Junk', sequences in the 'Core' and 'Repeat' sets are retained and used as input for another round of classification and filtering.

First, the assembly is screened against the NCBI UniVec database (ftp://ftp.ncbi.nlm.nih.gov/pub/UniVec/, downloaded 05/08/2019) to identify and remove contaminants. Hits are first scored using rules derived from NCBI Vecscreen (https://www.ncbi.nlm.nih.gov/tools/vecscreen/) and regions marked as 'Terminal' (within 25 bp of a sequence end), 'Proximal' (within 25 bases of another match) or 'Internal' (>25 bp from sequence end or vecsreen match). Then, any segment of fewer than 50 bases between

two vector matches or between a match and a sequence end are marked as 'Suspect'. In our experience, default Vecscreen parameters appear prone to excessive false positives in large genomes (data not shown), and so Diploidocus features two additional contaminant identification filters. First, the 'Expected False Discovery Rate' (eFDR) is calculated for each contaminant sequence. This is simply the BLAST+ Expect value for that hit, divided by the total number of hits at that Expect value threshold. Any hits with an eFDR value exceeding the default threshold of 1.0 were filtered from the vecscreen results. Short matches in long-read assemblies are unlikely to be real contamination and a second filter was applied, restricting contaminant screening to a minimum hit length of 50 bp. Finally, the percentage coverage per scaffold is calculated from the filtered hits. This is performed first for each contaminant individually, before being collapsed into total non-redundant contamination coverage per query. Diploidocus then removes any scaffolds with at least 50 % contamination, trims off any vector hits within 1 kb of the scaffold end, and masks any remaining vector contamination of at least 900 bp. This masking replaces every other base with an N to avoid an assembly gap being inserted: masked regions should be manually fragmented if required. Diploidocus can also report the number of mapped long reads that completely span regions flagged as contamination.

After contamination screening, a sorted BAM file of ONT reads mapped to the filtered assembly is generated using Minimap2 v2.17 (−ax map-ont --secondary = no) (Li, 2018). Purge Haplotigs coverage bins were adjusted to incorporate zero-coverage bases, excluding assembly gaps (defined as 10+ Ns). Counts of Complete, Duplicate and Fragmented BUSCO

genes were also generated for each sequence. General read depth statistics for each

sequence were calculated with BBMap v38.51 pileup.sh (Bushnell, 2014). The sect function

of KAT (KAT, RRID:SCR_016741) v2.4.2 (Mapleson et al., 2017) was used to calculate k-mer

frequencies for the 10x linked reads (first 16 bp trimmed from read 1), and the assembly

itself. Telomeres were predicted using a method adapted from

https://github.com/JanaSperschneider/FindTelomeres, searching each sequence for 5'

occurrences of a forward telomere regular expression sequence, C{2,4}T{1,2}A{1,3}, and 3'

occurrences of a reverse regular expression, T{1,3}A{1,2}G{2,4}. Telomeres were marked if at

least 50 % of the terminal 50 bp matches the appropriate sequence.

Figure 4. Diploidocus scaffold rating process based on a six-part classification. Asterisks

indicate any class value is accepted. Phases are executed in order. Consequently, rules for

later phases appear less restrictive than the full set of criteria required to receive that rating.

276 **Assembly polishing and gap-filling**

The assembly was first long-read polished with Racon (Racon, RRID:SCR_017642) v1.4.5

278 (Vaser et al., 2017) using the parameters -m 8 -x -6 -g -8 -w 500 and medaka v1.0.2 (Oxford

Nanopore Technologies Ltd., 2018) using the r941_prom_high_g303 model. Then, the 10x

280 reads were incorporated by short-read polishing using Pilon (Pilon, RRID:SCR_014731) v1.23

(Walker et al., 2014) with reads mapped using Minimap2 v2.12 (Li, 2018) and correcting for

282 indels only; we found correcting for indels only resulted in a higher BUSCO score than

correcting for indels and SNPs following the steps described in this section. We scaffolded

284 using SSPACE-LongRead v1.1 (Boetzer & Pirovano, 2014) with -k 1 followed by gap-filling

using gapFinisher v20190917 (Kammonen et al., 2019) with default parameters. After

286 another round of long-read polishing with Racon v1.4.5 (Vaser et al., 2017) and medaka

v1.0.2 (Oxford Nanopore Technologies Ltd., 2018), we moved forward with a second round

288 of tidying in Diploidocus v0.9.6 (default mode).

290 **Hi-C scaffolding**

Hi-C data were aligned to the draft genome assembly using the Juicer (Juicer,

292 RRID:SCR_017226) pipeline v1.6 (Durand et al., 2016) then scaffolds were ordered and

orientated using the 3D *de novo* assembly pipeline (3D de novo assembly, RRID:SCR_017227)

294 v180922 (Dudchenko et al., 2017). The contact map was visualised using Juicebox Assembly

Tools v1.11.08 and errors over 3 review rounds were corrected manually to resolve 11

296 chromosomes (Dudchenko et al., 2018). The resulting assembly was tidied again using

Diploidocus v0.10.6 (default mode).

Final polishing and assembly clean-up

300 A further round of long-read polishing with Racon v1.4.5 (Vaser et al., 2017) and medaka

v1.0.2 (Oxford Nanopore Technologies Ltd., 2018) was performed as described above. We

302 then short-read polished using Pilon v1.23 (Walker et al., 2014). Two Pilon strategies were

applied: (1) indel-only correction; (2) indel and SNP correction. We retained the indel and

304 SNP corrected assembly as it resulted in a marginally higher BUSCO score compared to indel

only correction (1311 vs 1310 complete BUSCOs); there was no change to contig nor scaffold

306 numbers. A final hybrid polish was performed using Hypo v1.0.3 (Kundu et al., 2019). The

assembly was concluded with a final tidy with Diploidocus v0.14.1 (default mode). All gaps in

308 the assembly were then standardised to 100 bp.

310 Genome-wide heterozygosity was estimated using trimmed 10x reads with GenomeScope

(Vurture et al., 2017) from the k-mer 20 histogram computed using Jellyfish (Jellyfish,

312 RRID:SCR_005491) v2.2.10 (Marçais & Kingsford, 2011).

314 **Genome annotation**

The genome was annotated using the homology-based gene prediction program GeMoMa

316 (GeMoMa, RRID:SCR_017646) v1.7.1 (Keilwagen et al., 2019) with four reference genomes

downloaded from NCBI: *Macadamia integrifolia* (SCU_Mint_v3, GCA_013358625.1),

318 *Nelumbo nucifera* (Chinese Lotus 1.1, GCA_000365185.2), *Arabidopsis thaliana* (TAIR10.1,

GCA_000001735.2) and *Rosa chinensis* (RchiOBHm-V2, GCA_002994745.2). The annotation

files for *M. integrifolia* were downloaded from the Southern Cross University data repository

([doi.org/10.25918/5e320fd1e5f06](doi.org/10.25918/5e320fd1e5f06)). *Macadamia* (Nock et al., 2020) and *Nelumbo* (Ming et

al., 2013) genomes were chosen as they are related to *Telopea* i.e. in the order Proteales.

The other two high-quality genomes represented the core eudicots and included the model

flowering plant *Arabidopsis* (Lamesch et al., 2012) and *Rosa* (Hibrand Saint-Oyant et al.,

2018) where the publication focused on genetic regulators of ornamental traits which is of

interest for *Telopea*. Annotation completeness was assessed using BUSCO v3.0.2b and v5.0.0

in proteome mode.

Ribosomal RNA (rRNA) genes were predicted with Barrnap (Barrnap, RRID:SCR_015995) v0.9

(Seemann, 2018) and transfer RNAs (tRNAs) were predicted with tRNAscan-SE (tRNAscan-SE,

RRID:SCR_010835) v2.05 (Lowe & Chan, 2016), implementing Infernal (Infernal,

RRID:SCR_011809) v1.1.2 (Nawrocki & Eddy, 2013). A set of 2,419 tRNAs was initially

predicted and filtered to 760 using the recommended protocol for eukaryotes. Then, 22

tRNAs with mismatched isotype and 10 with unexpected anticodon were removed to form

the high-confidence set.

The genome has also been annotated by the NCBI Eukaryotic Genome Annotation Pipeline

using RNAseq data from other Proteaceae (RefSeq accession GCF_018873765.1).

Genome-wide copy number analysis

Estimated single-copy (2n) sequencing depth was calculated for different regions of the

342 genome using the same smoothed density profile as employed by DepthSizer (Box 1) and

comparing this to the BUSCO-derived single-copy (2n) sequencing depth of DepthSizer. This

344 analysis was performed on: (1) BUSCO v5 (MetaEuk) single-copy 'Complete' genes; (2)

BUSCO v5 'Duplicated' genes; (3) All NCBI gene annotations; (4) Each final assembly scaffold;

346 (5) 100 kb non-overlapping windows across the genome. For convenience, this method has

been made available as DepthKopy (https://github.com/slimsuite/depthkopy).

Repeat annotation

350 Following the approach from the *Macadamia integrifolia* genome paper (Nock et al., 2020),

we identified and quantified repeats in the Telopea genome as well as the other four species

352 used in the GeMoMa annotation for comparison. A custom repeat library was generated

with RepeatModeler (RepeatModeler, RRID:SCR_015027) v2.0.1 (-engine ncbi) and the

354 genome was masked with RepeatMasker (RepeatMasker, RRID:SCR_012954) v4.1.0 (Tarailo-

Graovac & Chen, 2009), both with default parameters. The annotation table was generated

356 using the buildSummary.pl RepeatMasker script.

358 **Orthologous clusters and synteny analyses**

Synteny between the *Telopea* (Tspe_v1) and *Macadamia* (SCU_Mint_v3) genomes was

360 explored with satsuma2 version untagged-2c08e401140c1ed03e0f with parameters -l 3000 -

do_refine 1 -min_matches 40 -cutoff 2 -min_seed_length 48 and visualised with the

362 ChromosomePaint function (Grabherr et al., 2010) and MizBee v1.0 (Meyer et al., 2009). The

protein sequences of Tspe_v1 and the four species used in the GeMoMa annotation were

364 clustered into orthologous groups and tests for gene ontology (GO) enrichment were

conducted for waratah-specific clusters using OrthoVenn2 (Xu et al., 2019). Intersection of

366 clusters was visualised using the R package UpSetR (Conway et al., 2017).

368 ***CYCLOIDEA* transcription factor gene family analysis**

Complete and partial protein sequences for *CYCLOIDEA* transcription factors were

370 downloaded from NCBI using identifiers listed in Table S3 of Citerne et al., 2017. GABLAM

v2.30.5 (Davey et al., 2006) was used to identify all homologous proteins (BLAST+ v2.11.0,

372 blastp e-value <1e-4) in the waratah GeMoMa annotation, which was annotated with

protein descriptions from closest Swissprot hits using SAAGA v0.7.6 (Stuart et al., 2021).

374 Each *Telopea speciosissima* homologue was then used as query sequence for HAQESAC

v1.14.0 (Edwards et al., 2007) to generate a high-quality multiple sequence alignment and

376 inferred phlyogenetic tree of close homologues (limited to a maximum of 100 closest hits). A

search database was constructed from all angiosperm proteins in Uniprot (taxid 3398), the

378 three reference proteomes used for GeMoMa annotation (*Macadamia integrifolia*, *Nelumbo

nucifera* and *Rosa chinensis*), and all angiosperm reference proteomes from Quest For

380 Orthologues (March 2021 release; (Forslund et al., 2018). To this were added the original

CYC sequences and full GeMoMa annotation of *T. speciosissima*. BLAST+ searches and

382 HAQESAC runs were controlled by MultiHAQ v1.5.0 (Jones et al., 2011). To generate a

comprehensive but non-redundant tree of *CYC* genes, all homologues meeting initial

384 HAQESAC screening criteria (min 40 % global identity and 60 % global coverage to query, <50

% gaps relative to nearest homologue) were combined into a single non-redundant dataset

386 of *CYCLOIDEA* homologues and their homologues. A candidate *Telopea CYCLOIDEA*-like 1

gene (TSPEV1G03060) was identified based on SAAGA annotation and HAQESAC

388 homologues. This was used as a query for a second, manually curated HAQESAC run against

the full non-redundant protein dataset, screening out any proteins with an unknown species

390 designation (including sequence assigned the 9MAGSP species code). Multiple sequence

alignments were performed with Clustal Omega (Clustal Omega, RRID:SCR_001591) v1.2.4

392 (Sievers et al., 2011). The final tree was generated with IQ-TREE (IQ-TREE, RRID:SCR_017254)

v2.0.4 (Nguyen et al., 2015) with 1,000 bootstraps.

RESULTS AND DISCUSSION

High-quality chromosome-level Tspe_v1 reference genome

398 The ONT, 10x and Hi-C sequencing yielded a total of 48.3, 123.4 and 25.0 Gb of sequence,

respectively (Table 1). At the initial long-read assembly stage, NECAT resulted in the most

400 contiguous assembly, at 365 contigs and the highest BUSCO completeness at 81.2 %. This

was followed by Flye at 2,484 contigs and 81.0 % complete, then Canu at 3,983 contigs at

402 78.4 % complete. The BUSCO completeness of the 10x pseudohaploid assembly was higher

than each of the long-read assemblies at 91.8 %. However, the 10x assembly had much

404 lower contiguity at 43,951 contigs, as expected (Table S3). Whilst Supernova had a higher

BUSCO completeness (91.9 % versus 81.2 %), NECAT was orders of magnitude better in

406 terms of contiguity (10.7 Mb N50 on 365 contigs vs 874 kb N50 on 27,610 scaffolds).

Furthermore, BUSCOMP analysis revealed that the NECAT assembly contained more

408 complete BUSCO genes when base accuracy is not considered (Figure 5; Supplementary Files

– BUSCOMP full report). Guided by these metrics, NECAT was selected as the core assembly

410   for additional processing. We confirmed the individual's diploid status with Smudgeplot

     (Figure S2a).

     Rounds of polishing and tidying improved the contiguity and quality of the genome as the

414   genome progressed through the assembly workflow (Table S3). The first round of polishing

     markedly improved the BUSCO score – long-read polishing increased complete BUSCOs from

416   1,532 (v0.2) to 1,590 (v0.3) and short-read polishing further increased this to 1,602 (v0.4).

     The assembly was scaffolded by SSPACE-LongRead from 209 contigs into 138 scaffolds,

418   however, no gaps were filled by gapFinisher. After further long-read polishing, a run of

     Diploidocus (v0.7) retained 128 scaffolds out of 138, which consisted of 87 core, 41 repeat,

420   10 quarantine and 0 junk scaffolds. Following incorporation of Hi-C data, the assembly was

     in 2,357 scaffolds, and the N50 increased substantially from 16.5 Mb to 68.9 Mb.

422   Surprisingly, the contig number increased considerably from 148 to 3,537, suggesting that

     the Hi-C data and NECAT assembly were frequently in conflict. The resulting assembly was

424   tidied with Diploidocus and 1643 scaffolds (824,534,974 bp) were retained out of 2,357

     (833,952,765 bp; 1,347 core, 296 repeat, 548 quarantine and 166 junk scaffolds). The

426   removal of many sequences by Diploidocus, and the less contiguous initial assemblies from

     widely-used long-read assemblers Canu and Flye (Table S3), suggest that the NECAT

428   assembly contained erroneously joined sequences, and these were corrected by Hi-C.

     However, it is also possible that limitations of the Hi-C library contributed to the high degree

430   of fragmentation. The assembly contiguity improved to 1,399 scaffolds and 1,595 contigs

     following a further round of long-read polishing (Table S3). Following hybrid polishing with

432   Hypo (v0.9), the number of scaffolds remained as 1,399 and the BUSCO score improved

slightly. Notably, Hypo polishing improved the Merqury QV score from 29.8 to 33.9. A final

434   iteration of Diploidocus Tidy removed 72 putative haplotigs and 38 low quality 'junk'

scaffolds, keeping 1,084 core and 250 repetitive scaffolds.

The conclusion of the assembly workflow produced an 823.3 Mb haploid genome assembly

438   (Tspe_v1) on 1,289 scaffolds, with an N50 of 69.0 Mb and L50 of 6 (Table 2). The Hi-C data

facilitated scaffolding into 11 chromosomes (Figure 6), conforming to previous cytological

440   studies (Darlington & Wylie, 1956), and the anchored proportion of Tspe_v1 spanned 94.2 %

of the final assembly; the chromosomes were numbered by descending length (Table S8) as

442   this is the first instance *Telopea* chromosomes have been studied in detail.

444   From a core set of 1,614 single-copy orthologues from the Embryophyta lineage, 97.8 %

were complete in the assembly (86.7 % as single-copy, 11.2 % as duplicates), 1.7 % were

446   fragmented and only 0.5 % were not found, suggesting that the assembly includes most of

the waratah gene space. Interestingly, BUSCO scores vary by many percentages between

448   different BUSCO versions and gene predictors. BUSCO v5.0.0 with MetaEuk as the gene

predictor consistently produced the highest scores (Table S3). BUSCO v3.0.2b with Augustus

450   benchmarked the assembly against 1,440 single-copy orthologues only found 91.3 %

complete in the assembly (81.5 % as single-copy, 9.7 % as duplicates), with 2.9 % fragmented

452   and 5.8 % missing. BUSCO v5.0.0 with Augustus as the gene predictor reported higher scores

than v3.0.2b but lower than when MetaEuk was used as the gene predictor (Table S3). We

454   recovered a maximal non-redundant set of 1,549 complete single copy BUSCOs across the

set of assemblies. BUSCOMP analysis revealed that only one gene out of 1614 was not found

456   by BUSCO v5 MetaEuk in any version of the assembly (Figure 5; Supplementary File –

BUSCOMP full report). The Tspe_v1 assembly completeness is favourable in comparison to

458   the *Macadamia integrifolia* (SCU_Mint_v3) assembly (Nock et al., 2020), which also

combined long-read and Illumina sequences (BUSCO v5 MetaEuk 96.7 % vs 81.9 % complete,

460   respectively, in the anchored portion of the assembly). The Merqury QV score of the

assembly was 34.03, indicating a base-level accuracy of >99.99 % (Figure S3). Genome-wide

462   heterozygosity was estimated to be 0.756 % (Figure S2b).

464   **The *Telopea speciosissima* genome is approximately 900 Mb**

The 1C-value of *T. truncata* (Tasmanian waratah) has been estimated at 1.16 pg (1.13 Gb)

466   using flow cytometry (Jordan et al., 2015). Supernova v2.1.1 predicted a genome size of 953

Mb from the assembly of the 10x linked-reads whilst GenomeScope predicted a smaller

468   genome of 794 Mb from the same data (Figure S2b). DepthSizer analysis of the six different

versions of the genome assembly (four raw assemblies, Tspe_v1, and Tspe_v1

470   chromosomes) estimated the genome size of *T. speciosissima* to fall within a range from 850

Mb to 950 Mb (Table S9), and shows good robustness to both assembly version and BUSCO

472   dataset used (Figure 7). This falls between the Supernova and GenomeScope estimates. We

report an estimated genome size of approximately 900 Mb, considering the mean of

474   estimates of the six adjustment methods using the BUSCO v5 MetaEuk data, based on the

highest quality Tspe_v1 assemblies.

    The majority of Tspe_v1 is at single-copy (2n) read depth

478 Read depth copy number analysis reveals that the majority of the assembly is at the

expected $2n$ depth (Figure 8). Single-copy 'Complete' BUSCO genes strongly cluster around

480 CN = 1, further supporting the robustness of the method underpinning DepthSizer. Notably,

the 180 'Duplicated' BUSCO genes are also predominantly at single-copy depth, with a

482 similar copy number distribution to the BUSCOs classified as single-copy and complete. This

indicates that the vast majority are likely to be real duplications found in *T. speciosissima*,

484 with only a few representing potential sequencing errors (Table S10). This was supported by

HAQESAC phylogenetic analysis of all 180 genes (Supplementary File –

486 Tspe_v1.buscodup_HAQESAC.zip). Copy number analysis of all 14,882 NCBI annotated genes

shows a similar clustering around a median copy number of 1. However, the mean copy

488 number is surprisingly high at 2.36. Further inspection of the data revealed that this is being

driven by a reasonably small number of very high copy number genes, derived from highly

490 collapsed repeat regions (Table S11). This is further supported by the elevated mean copy

number for both whole scaffolds and 100 kb windows. This is consistent with the

492 identification by Diploidocus of 250 repetitive scaffolds, and a final assembly of approx. 91.5

% of the predicted genome size. Consistent with other Hi-C scaffolded assemblies (e.g. Rhie

494 et al., 2021), it is likely that Tspe_v1 still contains some misassemblies that will need to be

corrected with additional curation in future.

Repetitive elements and gene prediction

498 The *Telopea* genome is highly repetitive, with repeats accounting for 62.3 % of the total

sequence length and has a similar repeat content to *Macadamia*, previously reported as

500 55.1 % (Nock et al., 2020) and found to be 58.5 % in our analyses (Table S12). Class I

transposable elements (TEs) or retrotransposons were the most pervasive classified repeat

502   class (20.3 % of the genome) and were dominated by long terminal repeat (LTR)

retrotransposons (18.1 %). Class II TEs (DNA transposons) only accounted for 0.03 % of the

504   genome. A high percentage of repeats remained unclassified (40.6 %) and the genome will

serve as a resource for future studies into repetitive elements in *Telopea* and related

506   species.

508   Genome annotation predicted 40,126 protein-coding genes and 46,842 mRNAs in the *T.*

speciosissima assembly, which fits the expectation for plant genomes (Sterck et al., 2007). Of

510   these genes, 38,427 appeared in the 11 chromosomes (Table S8). Of 1,440 Embryophyta

orthologous proteins, 94.0 % were complete in the annotation (79.3 % as single-copy, 14.7 %

512   as duplicates), 3.4 % were fragmented and 2.6 % were missing. Additionally, 351 rRNA genes

and a set of 728 high-confidence transfer RNAs (tRNAs) were predicted. The NCBI

514   Annotation Release 100 had a higher completeness, as expected, than the GeMoMa

annotation; of 1,614 Embryophyta genes, 98.3 % were complete in the annotation (54.2 % as

516   single-copy, 44.1 % as duplicated), 1.1 % were fragmented and 0.6 % were missing. When

comparing the assembly completeness with proteome completeness using BUSCO v3.0.2b,

518   the proteome completeness at 94.0 % (79.3 % as single-copy and 14.7 % as duplicated) was

unexpectedly higher than the genome completeness at 91.3 % (81.5 % as single-copy and 9.7

520   % as duplicated). However, this issue was resolved with a later version of BUSCO (v5.0.0).

The improvements in BUSCO likely meant that genes could be better discerned in the

522   genome assembly, where they are more difficult to identify, compared to a proteome.

524   An inverse pattern in the incidence of genes and repeats was observed across all

chromosomes, with repeat content generally peaking towards the centre of each

526   chromosome (Figure 9), suggesting predominantly metacentric and submetacentric

chromosomes. This pattern may represent enriched repeat content and reduced coding

528   content in pericentromeric regions, although further study is required to identify the

centromeres (Jiang et al., 2003; Oliveira & Torres, 2018; Simon et al., 2015).

BUSCO completeness statistics must be matched by version and gene predictor

532   One surprising observation from our BUSCO analysis was a jump in completeness of over 6 %

when moving from BUSCO v3 Augustus predictions to BUSCO v5 MetaEuk predictions (Figure

534   5 and Table S3). This is explained in part by the change to the lineage database used.

However, completeness scores for BUSCO v5 Augustus are only about 3 % higher. This is

536   particularly pronounced for the raw assemblies, where Augustus scores can be over 10 %

lower than MetaEuk scores. Great care must be taken in naïve comparison of published

538   BUSCO scores, even if using the same version of BUSCO. MetaEuk scores seem to be both

higher and more stable. However, nucleotide sequences for Complete BUSCO genes are

540   currently only output from Augustus mode. We have therefore updated BUSCOMP to

extract the missing sequences from MetaEuk runs so that they can be used with

542   downstream tools such as BUSCOMP that require these sequences.

544   **Orthologous clusters and synteny between *Telopea* and *Macadamia***

The five species formed 24,140 clusters: 23,031 orthologous clusters (containing at least 2

546   species) and 1,109 single-copy gene clusters. There were 9,463 orthologous families

common to all of the species. The three members of the order Proteales (*T. speciosissima*,

548   *M. integrifolia* and *N. nucifera*) shared 456 families (Figure 10 and Figure S4). Tests for GO

enrichment of 912 waratah-specific clusters identified 12 significant terms (Table S13). The

550   most enriched GO terms were DNA recombination (GO:0006310, P = 1.8 x 10^{-27}),

retrotransposon nucleocapsid (GO:0000943, P = 3.5 x 10^{-12}) and DNA integration

552   (GO:0015074, P = 4.1 x 10^{-11}).

554   The *Macadamia* genome (2*n* = 28) has six more chromosomes than the *Telopea* genome (2*n*

= 22), but the two species have similar estimated genome sizes – 896 Mb (Nock et al., 2020)

556   compared to 874 Mb. It is thought that the ancestral Proteaceae had a chromosome number

of *x* = 7 (Carta et al., 2020; L. A. S. Johnson & Briggs, 1963, 1975; Murat et al., 2017),

558   although the occurrence of paleo-polyploidy in family has been debated (Stace et al., 1998).

Overall, synteny analyses reveal an abundance of interchromosomal rearrangements

560   between the *Telopea* and *Macadamia* genomes (Figure 11), reflecting the long time since

their divergence (73-83 Ma; Sauquet et al., 2009). However, a number of regions exhibit

562   substantial collinearity, for example, *Telopea* chromosome 09 and *Macadamia* chromosome

11 (Figure S5).

CYC gene copy number and the genetic control of floral symmetry

566   In total, 210 predicted waratah sequences (longest isoform per gene) were identified as

homologous to the 49 Citerne et al. *CYC* protein sequences. Of these, 198 generated

568   multiple sequence alignments and phylogenetic trees. These combined to form a non-

redundant dataset of 12,238 proteins. HAQESAC reduced this to a high-quality alignment of

570   46 homologous proteins, including two waratah proteins, TSPEV1G03060 – *CYC1* and

        TSPEV1G20406 – *CYC2.* Consistent with previous work (Citerne et al., 2017), these two

572   proteins belonged to two distinct clades (Figure 12). While the exact role of the two

        paralogues in determining floral symmetry in Proteaceae would require a study of gene

574   expression and remains incompletely understood in the species examined so far (Citerne et

        al., 2017; Damerval et al., 2019), this is the first study to quantify the total number of

576   *CYCLOIDEA* paralogues in Proteaceae based on a complete genome sequence. Our results

        hence lend further support to the pattern of a single gene duplication in the stem lineage of

578   Proteaceae that had so far emerged from Sanger and transcriptome sequencing.

580   **A molecular resource for biodiversity genomics**

        The *T. speciosissima* reference genome will enable genome-scale research into Proteaceae

582   evolution, at a wide range of scales. At shallower evolutionary scales, the *Telopea* genus

        contains five species that exhibit genetic variation consistent with a history of divergence

584   and introgression, likely driven by climatic change (Rossetto et al., 2011, 2012). Recent

        studies highlight the power of genome-scale approaches for inferring demographic change

586   and mechanistic forces that have influenced such clades, often making use of heterogenetity

        in patterns of variation across whole genomes (Choi et al., 2021; Soltis & Soltis, 2021). We

588   expect the waratah genome to similarly facilitate studies that provide new insights about

        historical gene flow and selection, in changing environments.

CONCLUSIONS

We present a high-quality annotated chromosome-level reference genome of *Telopea*

594 *speciosissima* assembled from Oxford Nanopore long-reads, 10x Genomics Chromium

linked-reads and Hi-C (823 Mb in length, N50 of 69.9 Mb and BUSCO completeness of 97.8

596 %): the first for a waratah, and only the second publicly available Proteaceae reference

genome. We envisage these data will be a platform to underpin evolutionary genomics, gene

598 discovery, breeding and the conservation of Proteaceae and the Australian flora.

600 ACKNOWLEDGEMENTS

602 We thank Stuart Allan for providing access to the sequenced plant and assistance with

sample collection at Blue Mountains Botanic Garden and Carolyn Connelly for facilitating

604 access to lab materials at the Royal Botanic Garden Sydney. We acknowledge Chris Jackson

for advice on repeat annotation. We thank the members of UNSW Research Technology

606 Services, particularly Duncan Smith, for help with software installation on the high-

performance computing cluster Katana. We acknowledge Mabel Lum for assistance with the

608 Bioplatforms Australia data portal. ONT and 10x sequencing were conducted at the

Australian Genome Research Facility (AGRF). Hi-C library prep and sequencing was

610 conducted at the Ramaciotti Centre for Genomics at the University of New South Wales.

612 FUNDING

614   We would like to acknowledge the contribution of the Genomics for Australian Plants

Framework Initiative consortium

616   (https://www.genomicsforaustralianplants.com/consortium/) in the generation of data used

in this publication. The Initiative is supported by funding from Bioplatforms Australia

618   (enabled by NCRIS), the Ian Potter Foundation, Royal Botanic Gardens Foundation (Victoria),

Royal Botanic Gardens Victoria, the Royal Botanic Gardens and Domain Trust, the Council of

620   Heads of Australasian Herbaria, CSIRO, Centre for Australian National Biodiversity Research

and the Department of Biodiversity, Conservation and Attractions, Western Australia. SHC

622   was supported through an Australian Government Research Training Program Scholarship.

RJE was funded by the Australian Research Council (LP160100610 and LP18010072).

AUTHOR CONTRIBUTIONS

JGB coordinated the project. MR, MvdM, PL-I, HS, GB, JGB and RJE designed the study and

628   funded the project. GB provided the samples. PL-I and J-YSY performed optimised DNA

extraction protocols and performed extractions. SHC performed the genome assembly,

630   scaffolding and annotation. RJE conceptualised and developed Diploidocus and DepthSizer.

TGA and RJE performed the DepthSizer benchmarking analysis. RJE performed the copy

632   number analysis and *CYC* phylogenetics. SHC, RJE and JGB wrote the manuscript. All authors

edited and approved the final manuscript.

REFERENCES

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search

638   tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data*.

640   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Boetzer, M., & Pirovano, W. (2014). SSPACE-LongRead: Scaffolding bacterial draft genomes using

642   long read sequence information. *BMC Bioinformatics*, *15*(1), 211.

https://doi.org/10.1186/1471-2105-15-211

644   Busch, A., & Zachgo, S. (2009). Flower symmetry evolution: Towards understanding the abominable

mystery of angiosperm radiation. *BioEssays*, *31*(11), 1181–1190.

646   https://doi.org/10.1002/bies.200900081

Bushnell, B. (2014). *BBMap: A fast, accurate, splice-aware aligner*.

648   https://sourceforge.net/projects/bbmap/

Carta, A., Bedini, G., & Peruzzi, L. (2020). A deep dive into the ancestral chromosome number and

650   genome size of flowering plants. *New Phytologist*, *228*(3), 1097–1106.

https://doi.org/10.1111/nph.16668

652   Chapman, M. A., Leebens-Mack, J. H., & Burke, J. M. (2008). Positive selection and expression

divergence following gene duplication in the sunflower *CYCLOIDEA* gene family. *Molecular*

654   *Biology and Evolution*, *25*(7), 1260–1273.

Chen, Y., Nie, F., Xie, S.-Q., Zheng, Y.-F., Dai, Q., Bray, T., Wang, Y.-X., Xing, J.-F., Huang, Z.-J., Wang,

656   D.-P., He, L.-J., Luo, F., Wang, J.-X., Liu, Y.-Z., & Xiao, C.-L. (2021). Efficient assembly of

nanopore reads via highly accurate and intact error correction. *Nature Communications*,

658   *12*(1), 60. https://doi.org/10.1038/s41467-020-20236-7

Choi, J. Y., Dai, X., Alam, O., Peng, J. Z., Rughani, P., Hickey, S., Harrington, E., Juul, S., Ayroles, J. F.,

660   Purugganan, M. D., & Stacy, E. A. (2021). Ancestral polymorphisms shape the adaptive

radiation of Metrosideros across the Hawaiian Islands. *Proceedings of the National Academy*

662   *of Sciences*, *118*(37). https://doi.org/10.1073/pnas.2023801118

Citerne, H. L., Luo, D., Pennington, R. T., Coen, E., & Cronk, Q. C. B. (2003). A Phylogenomic

664   investigation of *CYCLOIDEA*-like TCP genes in the Leguminosae. *Plant Physiology*, *131*(3),

1042–1053. https://doi.org/10.1104/pp.102.016311

666 Citerne, H. L., Reyes, E., Le Guilloux, M., Delannoy, E., Simonnet, F., Sauquet, H., Weston, P. H.,

Nadot, S., & Damerval, C. (2017). Characterization of *CYCLOIDEA*-like genes in Proteaceae, a

668   basal eudicot family with multiple shifts in floral symmetry. *Annals of Botany*, *119*(3), 367–

378. https://doi.org/10.1093/aob/mcw219

670 Conway, J. R., Lex, A., & Gehlenborg, N. (2017). UpSetR: An R package for the visualization of

intersecting sets and their properties. *Bioinformatics*, *33*(18), 2938–2940.

672   https://doi.org/10.1093/bioinformatics/btx364

Crisp, M. D., & Weston, P. H. (1993). Geographic and ontogenetic variation in morphology of

674   Australian waratahs (*Telopea*: Proteaceae). *Systematic Biology*, *42*(1), 49–76. JSTOR.

https://doi.org/10.2307/2992556

676 Damerval, C., Citerne, H., Conde e Silva, N., Deveaux, Y., Delannoy, E., Joets, J., Simonnet, F., Staedler,

Y., Schönenberger, J., Yansouni, J., Le Guilloux, M., Sauquet, H., & Nadot, S. (2019).

678   Unraveling the developmental and genetic mechanisms underpinning floral architecture in

Proteaceae. *Frontiers in Plant Science*, *10*, 18. https://doi.org/10.3389/fpls.2019.00018

680 Darlington, C. D., & Wylie, A. P. (1956). *Chromosome atlas of flowering plants*. George Allen and

Unwin Ltd.

682 Davey, N. E., Shields, D. C., & Edwards, R. J. (2006). SLiMDisc: Short, linear motif discovery, correcting

for common evolutionary descent. *Nucleic Acids Research*, *34*(12), 3546–3554.

684   https://doi.org/10.1093/nar/gkl486

De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., & Van Broeckhoven, C. (2018). NanoPack:

686   Visualizing and processing long-read sequencing data. *Bioinformatics*, *34*(15), 2666–2669.

https://doi.org/10.1093/bioinformatics/bty149

688   Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim, M. S.,

Machol, I., Lander, E. S., Aiden, A. P., & Aiden, E. L. (2017). De novo assembly of the *Aedes*

690   *aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science*, *356*(6333), 92–95.

https://doi.org/10.1126/science.aal3327

692   Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R., Pham, M.,

Hilaire, B. G. S., Yao, W., Stamenova, E., Hoeger, M., Nyquist, S. K., Korchina, V., Pletch, K.,

694   Flanagan, J. P., Tomaszewicz, A., McAloose, D., Estrada, C. P., Novak, B. J., … Aiden, E. L.

(2018). The Juicebox Assembly Tools module facilitates de novo assembly of mammalian

696   genomes with chromosome-length scaffolds for under $1000. *BioRxiv*, 254797.

https://doi.org/10.1101/254797

698   Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., & Aiden, E. L.

(2016). Juicer provides a one-click system for analyzing loop-resolution Hi-C experiments. *Cell*

700   *Systems*, *3*(1), 95–98. https://doi.org/10.1016/j.cels.2016.07.002

Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLOS Computational Biology*, *7*(10), e1002195.

702   https://doi.org/10.1371/journal.pcbi.1002195

Edwards, R. J., Field, M. A., Ferguson, J. M., Dudchenko, O., Keilwagen, J., Rosen, B. D., Johnson, G. S.,

704   Rice, E. S., Hillier, L. D., Hammond, J. M., Towarnicki, S. G., Omer, A., Khan, R., Skvortsova, K.,

Bogdanovic, O., Zammit, R. A., Aiden, E. L., Warren, W. C., & Ballard, J. W. O. (2021).

706   Chromosome-length genome assembly and structural variations of the primal Basenji dog

(*Canis lupus familiaris*) genome. *BMC Genomics*, *22*(1), 188. https://doi.org/10.1186/s12864-

708   021-07493-6

Edwards, R. J., Moran, N., Devocelle, M., Kiernan, A., Meade, G., Signac, W., Foy, M., Park, S. D. E.,

710   Dunne, E., Kenny, D., & Shields, D. C. (2007). Bioinformatic discovery of novel bioactive

peptides. *Nature Chemical Biology*, *3*(2), 108–112. https://doi.org/10.1038/nchembio854

712   Ellegren, H., Smeds, L., Burri, R., Olason, P. I., Backström, N., Kawakami, T., Künstner, A., Mäkinen, H.,

Nadachowska-Brzyska, K., Qvarnström, A., Uebbing, S., & Wolf, J. B. W. (2012). The genomic

714   landscape of species divergence in *Ficedula* flycatchers. *Nature*, *491*(7426), 756–760.

https://doi.org/10.1038/nature11584

716   Fambrini, M., & Pugliesi, C. (2017). *CYCLOIDEA* 2 clade genes: Key players in the control of floral

symmetry, inflorescence architecture, and reproductive organ development. *Plant Molecular*

718   *Biology Reporter*, *35*(1), 20–36. https://doi.org/10.1007/s11105-016-1005-z

Feng, X., Zhao, Z., Tian, Z., Xu, S., Luo, Y., Cai, Z., Wang, Y., Yang, J., Wang, Z., Weng, L., Chen, J.,

720   Zheng, L., Guo, X., Luo, J., Sato, S., Tabata, S., Ma, W., Cao, X., Hu, X., … Luo, D. (2006).

Control of petal shape and floral zygomorphy in *Lotus japonicus*. *Proceedings of the National*

722   *Academy of Sciences*, *103*(13), 4970–4975. https://doi.org/10.1073/pnas.0600681103

Field, M. A., Rosen, B. D., Dudchenko, O., Chan, E. K. F., Minoche, A. E., Edwards, R. J., Barton, K.,

724   Lyons, R. J., Tuipulotu, D. E., Hayes, V. M., D. Omer, A., Colaric, Z., Keilwagen, J., Skvortsova,

K., Bogdanovic, O., Smith, M. A., Aiden, E. L., Smith, T. P. L., Zammit, R. A., & Ballard, J. W. O.

726   (2020). Canfam_GSD: De novo chromosome-length genome assembly of the German

Shepherd Dog (*Canis lupus familiaris*) using a combination of long reads, optical mapping,

728   and Hi-C. *GigaScience*, *9*(giaa027). https://doi.org/10.1093/gigascience/giaa027

Forslund, K., Pereira, C., Capella-Gutierrez, S., da Silva, A. S., Altenhoff, A., Huerta-Cepas, J., Muffato,

730   M., Patricio, M., Vandepoele, K., Ebersberger, I., Blake, J., Fernández Breis, J. T., Quest for

Orthologs Consortium, Boeckmann, B., Gabaldón, T., Sonnhammer, E., Dessimoz, C., Lewis,

732   S., & Quest for Orthologs Consortium. (2018). Gearing up to handle the mosaic nature of life

in the quest for orthologs. *Bioinformatics (Oxford, England)*, *34*(2), 323–329.

734   https://doi.org/10.1093/bioinformatics/btx542

Grabherr, M. G., Russell, P., Meyer, M., Mauceli, E., Alföldi, J., Di Palma, F., & Lindblad-Toh, K. (2010).

736   Genome-wide synteny through highly sensitive sequence alignment: Satsuma.

Bioinformatics, *26*(9), 1145–1151. https://doi.org/10.1093/bioinformatics/btq102

738   Gu, Z., Gu, L., Eils, R., Schlesner, M., & Brors, B. (2014). Circlize implements and enhances circular

visualization in R. *Bioinformatics*, *30*(19), 2811–2812.

740   https://doi.org/10.1093/bioinformatics/btu393

Hibrand Saint-Oyant, L., Ruttink, T., Hamama, L., Kirov, I., Lakhwani, D., Zhou, N. N., Bourke, P. M.,

742   Daccord, N., Leus, L., Schulz, D., Van de Geest, H., Hesselink, T., Van Laere, K., Debray, K.,

Balzergue, S., Thouroude, T., Chastellier, A., Jeauffre, J., Voisine, L., … Foucher, F. (2018). A

744   high-quality genome sequence of *Rosa chinensis* to elucidate ornamental traits. *Nature*

Plants, *4*(7), 473–484. https://doi.org/10.1038/s41477-018-0166-1

746   Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M. L., Reed, L.

K., Storfer, A., & Whitlock, M. C. (2016). Finding the genomic basis of local adaptation:

748   Pitfalls, practical solutions, and future directions. *The American Naturalist*, *188*(4), 379–397.

https://doi.org/10.1086/688018

750   Horn, S., Pabón-Mora, N., Theuß, V. S., Busch, A., & Zachgo, S. (2015). Analysis of the CYC/TB1 class

of TCP transcription factors in basal angiosperms and magnoliids. *The Plant Journal*, *81*(4),

752   559–571. https://doi.org/10.1111/tpj.12750

Howarth, D. G., & Donoghue, M. J. (2006). Phylogenetic analysis of the "ECE" (CYC/TB1) clade reveals

754   duplications predating the core eudicots. *Proceedings of the National Academy of Sciences*,

103(24), 9101–9106.

756   Inglis, P. W., Pappas, M. de C. R., Resende, L. V., & Grattapaglia, D. (2018). Fast and inexpensive

protocols for consistent extraction of high quality DNA and RNA from challenging plant and

758   fungal samples for high-throughput SNP genotyping and sequencing applications. *PLOS ONE*,

13(10), e0206085. https://doi.org/10.1371/journal.pone.0206085

760   Jiang, J., Birchler, J. A., Parrott, W. A., & Kelly Dawe, R. (2003). A molecular view of plant

centromeres. *Trends in Plant Science*, *8*(12), 570–575.

762   https://doi.org/10.1016/j.tplants.2003.10.011

Johnson, L. A. S., & Briggs, B. G. (1963). Evolution in the Proteaceae. *Australian Journal of Botany*,

764   *11*(1), 21–61.

Johnson, L. A. S., & Briggs, B. G. (1975). On the Proteaceae—The evolution and classification of a

766   southern family. *Botanical Journal of the Linnean Society*, *70*(2), 83–182.

https://doi.org/10.1111/j.1095-8339.1975.tb01644.x

768   Jones, B. M., Edwards, R. J., Skipp, P. J., O'Connor, C. D., & Iglesias-Rodriguez, M. D. (2011). Shotgun

proteomic analysis of *Emiliania huxleyi*, a marine phytoplankton species of major

770   biogeochemical importance. *Marine Biotechnology (New York, N.Y.)*, *13*(3), 496–504.

https://doi.org/10.1007/s10126-010-9320-0

772   Jordan, G. J., Carpenter, R. J., Koutoulis, A., Price, A., & Brodribb, T. J. (2015). Environmental

adaptation in stomatal size independent of the effects of genome size. *New Phytologist*,

774   *205*(2), 608–617. https://doi.org/10.1111/nph.13076

Kammonen, J. I., Smolander, O.-P., Paulin, L., Pereira, P. A. B., Laine, P., Koskinen, P., Jernvall, J., &

776   Auvinen, P. (2019). GapFinisher: A reliable gap filling pipeline for SSPACE-LongRead

scaffolder output. *PLOS ONE*, *14*(9), e0216885.

778   https://doi.org/10.1371/journal.pone.0216885

Keilwagen, J., Hartung, F., & Grau, J. (2019). GeMoMa: Homology-based gene prediction utilizing

780   intron position conservation and RNA-seq data. *Methods in Molecular Biology (Clifton, N.J.)*,

1962, 161–177. https://doi.org/10.1007/978-1-4939-9173-0_9

782   Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone reads using

        repeat graphs. *Nature Biotechnology*, *37*(5), 540–546. https://doi.org/10.1038/s41587-019-

784   0072-8

        Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017). Canu:

786   Scalable and accurate long-read assembly via adaptive k-mer weighting and repeat

        separation. *Genome Research*, gr.215087.116. https://doi.org/10.1101/gr.215087.116

788   Kundu, R., Casey, J., & Sung, W.-K. (2019). *HyPo: Super fast and accurate polisher for long read

        genome assemblies*. https://doi.org/10.1101/2019.12.19.882506

790   Lamesch, P., Berardini, T. Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R., Dreher, K.,

        Alexander, D. L., Garcia-Hernandez, M., Karthikeyan, A. S., Lee, C. H., Nelson, W. D., Ploetz, L.,

792   Singh, S., Wensel, A., & Huala, E. (2012). The Arabidopsis Information Resource (TAIR):

        Improved gene annotation and new tools. *Nucleic Acids Research*, *40*(D1), D1202–D1210.

794   https://doi.org/10.1093/nar/gkr1090

        Levy Karin, E., Mirdita, M., & Söding, J. (2020). MetaEuk—Sensitive, high-throughput gene discovery,

796   and annotation for large-scale eukaryotic metagenomics. *Microbiome*, *8*(1), 48.

        https://doi.org/10.1186/s40168-020-00808-x

798   Lewin, H. A., Robinson, G. E., Kress, W. J., Baker, W. J., Coddington, J., Crandall, K. A., Durbin, R.,

        Edwards, S. V., Forest, F., Gilbert, M. T. P., Goldstein, M. M., Grigoriev, I. V., Hackett, K. J.,

800   Haussler, D., Jarvis, E. D., Johnson, W. E., Patrinos, A., Richards, S., Castilla-Rubio, J. C., …

        Zhang, G. (2018). Earth BioGenome Project: Sequencing life for the future of life. *Proceedings

802   of the National Academy of Sciences*, *115*(17), 4325–4333.

        https://doi.org/10.1073/pnas.1720115115

804   Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18), 3094–

        3100. https://doi.org/10.1093/bioinformatics/bty191

806   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R.,

       & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map

808   format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

       https://doi.org/10.1093/bioinformatics/btp352

810   Lindblad-Toh, K., Wade, C. M., Mikkelsen, T. S., Karlsson, E. K., Jaffe, D. B., Kamal, M., Clamp, M.,

       Chang, J. L., Kulbokas, E. J., & Zody, M. C. (2005). Genome sequence, comparative analysis

812   and haplotype structure of the domestic dog. *Nature*, *438*(7069), 803–819.

       Lowe, T. M., & Chan, P. P. (2016). tRNAscan-SE On-line: Integrating search and context for analysis of

814   transfer RNA genes. *Nucleic Acids Research*, *44*(W1), W54-57.

       https://doi.org/10.1093/nar/gkw413

816   Lu-Irving, P., & Rutherford, S. (2021). *High molecular weight DNA extraction from leaf tissue*.

       dx.doi.org/10.17504/protocols.io.bu9ynz7w

818   Luo, D., Carpenter, R., Vincent, C., Copsey, L., & Coen, E. (1996). Origin of floral asymmetry in

       Antirrhinum. *Nature*, *383*(6603), 794–799.

820   Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO update: Novel

       and streamlined workflows along with broader and deeper phylogenetic coverage for scoring

822   of eukaryotic, prokaryotic, and viral genomes. *Molecular Biology and Evolution*, *38*(10),

       4647–4654. https://doi.org/10.1093/molbev/msab199

824   Mapleson, D., Garcia Accinelli, G., Kettleborough, G., Wright, J., & Clavijo, B. J. (2017). KAT: A K-mer

       analysis toolkit to quality control NGS datasets and genome assemblies. *Bioinformatics*,

826   *33*(4), 574–576. https://doi.org/10.1093/bioinformatics/btw663

       Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting of

828   occurrences of k-mers. *Bioinformatics*, *27*(6), 764–770.

       https://doi.org/10.1093/bioinformatics/btr011

830   Mast, A. R., Willis, C. L., Jones, E. H., Downs, K. M., & Weston, P. H. (2008). A smaller *Macadamia*

from a more vagile tribe: Inference of phylogenetic relationships, divergence times, and

832   diaspore evolution in *Macadamia* and relatives (tribe Macadamieae; Proteaceae). *American*

Journal of Botany, *95*(7), 843–870.

834   Meyer, M., Munzner, T., & Pfister, H. (2009). MizBee: A multiscale synteny browser. *IEEE*

Transactions on Visualization and Computer Graphics, *15*(6), 897–904.

836   https://doi.org/10.1109/TVCG.2009.167

Ming, R., VanBuren, R., Liu, Y., Yang, M., Han, Y., Li, L.-T., Zhang, Q., Kim, M.-J., Schatz, M. C.,

838   Campbell, M., Li, J., Bowers, J. E., Tang, H., Lyons, E., Ferguson, A. A., Narzisi, G., Nelson, D.

R., Blaby-Haas, C. E., Gschwend, A. R., … Shen-Miller, J. (2013). Genome of the long-living

840   sacred lotus (*Nelumbo nucifera* Gaertn.). *Genome Biology*, *14*(5), 1–11.

https://doi.org/10.1186/gb-2013-14-5-r41

842   Mirarab, S., Nguyen, N., & Warnow, T. (2011). SEPP: SATé-Enabled Phylogenetic Placement. In

Biocomputing 2012 (pp. 247–258). World Scientific.

844   https://doi.org/10.1142/9789814366496_0024

Murat, F., Armero, A., Pont, C., Klopp, C., & Salse, J. (2017). Reconstructing the genome of the most

846   recent common ancestor of flowering plants. *Nature Genetics*, *49*(4), 490–496.

https://doi.org/10.1038/ng.3813

848   Nawrocki, E. P., & Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics, *29*(22), 2933–2935. https://doi.org/10.1093/bioinformatics/btt509

850   Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and*

852   *Evolution*, *32*(1), 268–274. https://doi.org/10.1093/molbev/msu300

Nixon, P. (1987). *The Waratah*. Kangaroo Press.

854   Nock, C. J., Baten, A., Barkla, B. J., Furtado, A., Henry, R. J., & King, G. J. (2016). Genome and

transcriptome sequencing characterises the gene space of *Macadamia integrifolia*

856   (Proteaceae). *BMC Genomics*, *17*(1), 937. https://doi.org/10.1186/s12864-016-3272-3

Nock, C. J., Baten, A., Mauleon, R., Langdon, K. S., Topp, B., Hardner, C., Furtado, A., Henry, R. J., &

858   King, G. J. (2020). Chromosome-scale assembly and annotation of the macadamia genome

(*Macadamia integrifolia* HAES 741). *G3: Genes, Genomes, Genetics*, *10*(10), 3497–3504.

860   https://doi.org/10.1534/g3.120.401326

Offord, C. A. (2003). Improvement of waratahs (*Telopea* spp.) through breeding. *Acta Horticulturae*,

862   *603*, 119–122.

Offord, C. A. (2006). Analysis of characters and germplasm of significance to improvement of

864   Australian native waratahs (*Telopea* spp., family Proteaceae) for cut flower production.

Genetic Resources and Crop Evolution, *53*(6), 1263–1272. https://doi.org/10.1007/s10722-

866   005-3487-7

Offord, C. A., Nixon, P., & Goodwin, P. B. (1987). Development of the waratah as a commercial crop.

868   *Journal International Protea Association*, *14*, 14–15.

Oliveira, L. C., & Torres, G. A. (2018). Plant centromeres: Genetics, epigenetics and evolution.

870   *Molecular Biology Reports*, *45*(5), 1491–1497. https://doi.org/10.1007/s11033-018-4284-7

Oxford Nanopore Technologies Ltd. (2018). *Medaka*. https://github.com/nanoporetech/medaka

872   Patil, I. (2021). Visualizations with statistical details: The "ggstatsplot" approach. *Journal of Open

Source Software*, *6*(61), 3167. https://doi.org/10.21105/joss.03167

874   Phase Genomics. (2019). *Hic_qc*. https://github.com/phasegenomics/hic_qc

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic

876   features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/bioinformatics/btq033

R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R Foundation for

878   Statistical Computing. https://www.R-project.org/

Radwan, J., & Babik, W. (2012). The genomics of adaptation. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1749), 5024–5028. https://doi.org/10.1098/rspb.2012.2322

Ramsay, H. (1963). Chromosome numbers in the Proteaceae. *Australian Journal of Botany*, *11*(1), 1–20.

Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2019). GenomeScope 2.0 and Smudgeplots: Reference-free profiling of polyploid genomes. *BioRxiv*, 747568. https://doi.org/10.1101/747568

Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M., Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L., Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D. (2021). Towards complete and error-free genome assemblies of all vertebrate species. *Nature*, *592*(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality, completeness, and phasing assessment for genome assemblies. *Genome Biology*, *21*(1), 1–27.

Roach, M. J., Schmidt, S. A., & Borneman, A. R. (2018). Purge Haplotigs: Allelic contig reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*, *19*(1), 460. https://doi.org/10.1186/s12859-018-2485-7

Rossetto, M., Allen, C. B., Thurlby, K. A. G., Weston, P. H., & Milner, M. L. (2012). Genetic structure and bio-climatic modeling support allopatric over parapatric speciation along a latitudinal gradient. *BMC Evolutionary Biology*, *12*, 149. https://doi.org/10.1186/1471-2148-12-149

Rossetto, M., Thurlby, K. A., Offord, C. A., Allen, C. B., & Weston, P. H. (2011). The impact of distance and a shifting temperature gradient on genetic connectivity across a heterogeneous landscape. *BMC Evolutionary Biology*, *11*, 126. https://doi.org/10.1186/1471-2148-11-126

Royal Botanic Gardens, Kew. (2017). *State of the World's Plants 2017* (No. 978-1-84246-647–6). Royal

904   Botanic Gardens, Kew.

Sauquet, H., Weston, P. H., Anderson, C. L., Barker, N. P., Cantrill, D. J., Mast, A. R., & Savolainen, V.

906   (2009). Contrasted patterns of hyperdiversification in Mediterranean hotspots. *Proceedings*

   of the National Academy of Sciences, *106*(1), 221–225.

908   https://doi.org/10.1073/pnas.0805607106

Schalamun, M., Nagar, R., Kainer, D., Beavan, E., Eccles, D., Rathjen, J. P., Lanfear, R., & Schwessinger,

910   B. (2019). Harnessing the MinION: An example of how to establish long-read sequencing in a

   laboratory using challenging plant tissue from *Eucalyptus pauciflora*. *Molecular Ecology*

912   *Resources*, *19*(1), 77–89. https://doi.org/10.1111/1755-0998.12938

Seehausen, O., Butlin, R. K., Keller, I., Wagner, C. E., Boughman, J. W., Hohenlohe, P. A., Peichel, C. L.,

914   Saetre, G.-P., Bank, C., Brännström, Å., Brelsford, A., Clarkson, C. S., Eroukhmanoff, F., Feder,

   J. L., Fischer, M. C., Foote, A. D., Franchini, P., Jiggins, C. D., Jones, F. C., … Widmer, A. (2014).

916   Genomics and the origin of species. *Nature Reviews Genetics*, *15*(3), 176–192.

   https://doi.org/10.1038/nrg3644

918 Seemann, T. (2018). *Barrnap*. https://github.com/tseemann/barrnap

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H., Remmert,

920   M., Söding, J., Thompson, J. D., & Higgins, D. G. (2011). Fast, scalable generation of high-

   quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems*

922   *Biology*, *7*, 539. https://doi.org/10.1038/msb.2011.75

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015). BUSCO:

924   Assessing genome assembly and annotation completeness with single-copy orthologs.

   Bioinformatics (Oxford, England), *31*(19), 3210–3212.

926   https://doi.org/10.1093/bioinformatics/btv351

Simon, L., Voisin, M., Tatout, C., & Probst, A. V. (2015). Structure and function of centromeric and

928   pericentromeric heterochromatin in *Arabidopsis thaliana*. *Frontiers in Plant Science*, *6*.

https://doi.org/10.3389/fpls.2015.01049

930 Soltis, P. S., & Soltis, D. E. (2014). Flower Diversity and Angiosperm Diversification. In J. L. Riechmann

& F. Wellmer (Eds.), *Flower Development: Methods and Protocols* (pp. 85–102). Springer New

932   York. https://doi.org/10.1007/978-1-4614-9408-9_4

Soltis, P. S., & Soltis, D. E. (2021). Plant genomes: Markers of evolutionary history and drivers of

934   evolutionary change. *PLANTS, PEOPLE, PLANET*, *3*(1), 74–82.

https://doi.org/10.1002/ppp3.10159

936 Stace, H. M., Douglas, A. W., & Sampson, J. F. (1998). Did 'Paleo-polyploidy' Really occur in

Proteaceae? *Australian Systematic Botany*, *11*(4), 613–629. https://doi.org/10.1071/sb98013

938 Stanke, M., & Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in eukaryotes

that allows user-defined constraints. *Nucleic Acids Research*, *33*(suppl_2), W465–W467.

940 Sterck, L., Rombauts, S., Vandepoele, K., Rouzé, P., & Van de Peer, Y. (2007). How many genes are

there in plants (… and why are they there)? *Current Opinion in Plant Biology*, *10*(2), 199–203.

942   https://doi.org/10.1016/j.pbi.2007.01.004

Stuart, K. C., Edwards, R. J., Cheng, Y., Warren, W. C., Burt, D. W., Sherwin, W. B., Hofmeister, N. R.,

944   Werner, S. J., Ball, G. F., Bateson, M., Brandley, M. C., Buchanan, K. L., Cassey, P., Clayton, D.

F., Meyer, T. D., Meddle, S. L., & Rollins, L. A. (2021). Transcript- and annotation-guided

946   genome assembly of the European starling. *BioRxiv*, 2021.04.07.438753.

https://doi.org/10.1101/2021.04.07.438753

948 Summerell, B. A. (1997). Pests and diseases. In *The Waratah* (2nd edition). Kangaroo Press.

Summerell, B. A., Nixons, P. G., & Burgess, L. W. (1990). Crown and stem canker of waratah caused

950   by *Cylindrocarpon destructans*. *Australasian Plant Pathology*, *19*(1), 13–15.

https://doi.org/10.1071/APP9900013

952   Tarailo-Graovac, M., & Chen, N. (2009). Using RepeatMasker to identify repetitive elements in

        genomic sequences. *Current Protocols in Bioinformatics*, *25*(1), 4.10.1-4.10.14.

954   https://doi.org/10.1002/0471250953.bi0410s25

        Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome assembly

956   from long uncorrected reads. *Genome Research*, *27*(5), 737–746.

                https://doi.org/10.1101/gr.214270.116

958   Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., & Schatz, M.

        C. (2017). GenomeScope: Fast reference-free genome profiling from short reads.

960   *Bioinformatics*, *33*(14), 2202–2204. https://doi.org/10.1093/bioinformatics/btx153

        Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A., Zeng, Q.,

962   Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: An integrated tool for comprehensive

                Microbial variant detection and genome assembly improvement. *PLOS ONE*, *9*(11), e112963.

964   https://doi.org/10.1371/journal.pone.0112963

        Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct determination of

966   diploid genome sequences. *Genome Research*, *27*(5), 757–767.

                https://doi.org/10.1101/gr.214874.116

968   Weston, P. H. (2006). Proteaceae. In K. Kubitzki (Ed.), *The Families and Genera of Vascular Plants.

        Volume IX* (pp. 364–404). Springer-Verlag.

970   Weston, P. H., & Crisp, M. D. (1994). Cladistic biogeography of waratahs (Proteaceae, Embothrieae)

        and their allies across the pacific. *Australian Systematic Botany*, *7*(3), 225–249.

972   https://doi.org/10.1071/sb9940225

        Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Completing bacterial genome assemblies

974   with multiplex MinION sequencing. *Microbial Genomics*, *3*(10).

                https://doi.org/10.1099/mgen.0.000132

Worrall, R., & Gollnow, B. (2013). *Growing waratahs for cut flowers—A guide for commercial growers* (No. 12/087). Rural Industries Research and Development Corporation.

Xu, L., Dong, Z., Fang, L., Luo, Y., Wei, Z., Guo, H., Zhang, G., Gu, Y. Q., Coleman-Derr, D., Xia, Q., & Wang, Y. (2019). OrthoVenn2: A web server for whole-genome comparison and annotation

of orthologous clusters across multiple species. *Nucleic Acids Research*, *47*(W1), W52–W58. https://doi.org/10.1093/nar/gkz333

Yadav, S., Dudchenko, O., Esvaran, M., Rosen, B. D., Field, M. A., Skvortsova, K., Edwards, R. J., Gopalakrishnan, S., Keilwagen, J., Cochran, B. J., Manandhar, B., Bucknall, M., Bustamante, S.,

Rasmussen, J. A., Melvin, R. G., Omer, A., Colaric, Z., Chan, E. K. F., Minoche, A. E., … Ballard, J. W. O. (2020). Desert Dingo (*Canis lupus dingo*) genome provides insights into their role in

the Australian ecosystem. *BioRxiv*, 2020.11.15.384057. https://doi.org/10.1101/2020.11.15.384057

Yang, X., Zhao, X.-G., Li, C.-Q., Liu, J., Qiu, Z.-J., Dong, Y., & Wang, Y.-Z. (2015). Distinct regulatory changes underlying differential expression of TEOSINTE BRANCHED1-CYCLOIDEA-

PROLIFERATING CELL FACTOR genes associated with petal variations in zygomorphic flowers of *Petrocosmea* spp. Of the family Gesneriaceae. *Plant Physiology*, *169*(3), 2138–2151.

Zheng, T., Li, P., Li, L., & Zhang, Q. (2021). Research advances in and prospects of ornamental plant genomics. *Horticulture Research*, *8*(1), 1–19. https://doi.org/10.1038/s41438-021-00499-x

Zhong, J., & Kellogg, E. A. (2015). Duplication and expression of CYC2-like genes in the origin and maintenance of corolla zygomorphy in Lamiales. *New Phytologist*, *205*(2), 852–868.

DATA ACCESSIBILITY

The Tspe_v1 genome was deposited to NCBI under BioProject PRJNA712988 and BioSample

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local

1000   alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410.

Andrews, S. (2010). *FastQC: a quality control tool for high throughput sequence data*.

1002   http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Boetzer, M., & Pirovano, W. (2014). SSPACE-LongRead: Scaffolding bacterial draft genomes

1004   using long read sequence information. *BMC Bioinformatics*, *15*(1), 211.

https://doi.org/10.1186/1471-2105-15-211

1006   Busch, A., & Zachgo, S. (2009). Flower symmetry evolution: Towards understanding the

abominable mystery of angiosperm radiation. *BioEssays*, *31*(11), 1181–1190.

1008   https://doi.org/10.1002/bies.200900081

Bushnell, B. (2014). *BBMap: A fast, accurate, splice-aware aligner*.

1010   https://sourceforge.net/projects/bbmap/

Carta, A., Bedini, G., & Peruzzi, L. (2020). A deep dive into the ancestral chromosome

1012   number and genome size of flowering plants. *New Phytologist*, *228*(3), 1097–1106.

https://doi.org/10.1111/nph.16668

1014   Chapman, M. A., Leebens-Mack, J. H., & Burke, J. M. (2008). Positive selection and

expression divergence following gene duplication in the sunflower *CYCLOIDEA* gene

1016   family. *Molecular Biology and Evolution*, *25*(7), 1260–1273.

Chen, Y., Nie, F., Xie, S.-Q., Zheng, Y.-F., Dai, Q., Bray, T., Wang, Y.-X., Xing, J.-F., Huang, Z.-J.,

1018   Wang, D.-P., He, L.-J., Luo, F., Wang, J.-X., Liu, Y.-Z., & Xiao, C.-L. (2021). Efficient

assembly of nanopore reads via highly accurate and intact error correction. *Nature*

1020   *Communications*, *12*(1), 60. https://doi.org/10.1038/s41467-020-20236-7

Choi, J. Y., Dai, X., Alam, O., Peng, J. Z., Rughani, P., Hickey, S., Harrington, E., Juul, S.,

1022   Ayroles, J. F., Purugganan, M. D., & Stacy, E. A. (2021). Ancestral polymorphisms

shape the adaptive radiation of Metrosideros across the Hawaiian Islands.

1024   *Proceedings of the National Academy of Sciences*, *118*(37).

https://doi.org/10.1073/pnas.2023801118

1026   Citerne, H. L., Luo, D., Pennington, R. T., Coen, E., & Cronk, Q. C. B. (2003). A Phylogenomic

investigation of *CYCLOIDEA*-like TCP genes in the Leguminosae. *Plant Physiology*,

1028   *131*(3), 1042–1053. https://doi.org/10.1104/pp.102.016311

Citerne, H. L., Reyes, E., Le Guilloux, M., Delannoy, E., Simonnet, F., Sauquet, H., Weston, P.

1030   H., Nadot, S., & Damerval, C. (2017). Characterization of *CYCLOIDEA*-like genes in

Proteaceae, a basal eudicot family with multiple shifts in floral symmetry. *Annals of*

1032   *Botany*, *119*(3), 367–378. https://doi.org/10.1093/aob/mcw219

Conway, J. R., Lex, A., & Gehlenborg, N. (2017). UpSetR: An R package for the visualization of

1034   intersecting sets and their properties. *Bioinformatics*, *33*(18), 2938–2940.

https://doi.org/10.1093/bioinformatics/btx364

1036   Crisp, M. D., & Weston, P. H. (1993). Geographic and ontogenetic variation in morphology of

Australian waratahs (*Telopea*: Proteaceae). *Systematic Biology*, *42*(1), 49–76. JSTOR.

1038   https://doi.org/10.2307/2992556

Damerval, C., Citerne, H., Conde e Silva, N., Deveaux, Y., Delannoy, E., Joets, J., Simonnet, F.,

1040   Staedler, Y., Schönenberger, J., Yansouni, J., Le Guilloux, M., Sauquet, H., & Nadot, S.

(2019). Unraveling the developmental and genetic mechanisms underpinning floral

1042   architecture in Proteaceae. *Frontiers in Plant Science*, *10*, 18.

https://doi.org/10.3389/fpls.2019.00018

1044 Darlington, C. D., & Wylie, A. P. (1956). *Chromosome atlas of flowering plants*. George Allen and Unwin Ltd.

1046 Davey, N. E., Shields, D. C., & Edwards, R. J. (2006). SLiMDisc: Short, linear motif discovery, correcting for common evolutionary descent. *Nucleic Acids Research*, *34*(12), 3546–

1048 3554. https://doi.org/10.1093/nar/gkl486

De Coster, W., D'Hert, S., Schultz, D. T., Cruts, M., & Van Broeckhoven, C. (2018). NanoPack:

1050 Visualizing and processing long-read sequencing data. *Bioinformatics*, *34*(15), 2666–

2669. https://doi.org/10.1093/bioinformatics/bty149

Dudchenko, O., Batra, S. S., Omer, A. D., Nyquist, S. K., Hoeger, M., Durand, N. C., Shamim, M. S., Machol, I., Lander, E. S., Aiden, A. P., & Aiden, E. L. (2017). De novo assembly of

the *Aedes aegypti* genome using Hi-C yields chromosome-length scaffolds. *Science*, *356*(6333), 92–95. https://doi.org/10.1126/science.aal3327

Dudchenko, O., Shamim, M. S., Batra, S. S., Durand, N. C., Musial, N. T., Mostofa, R., Pham, M., Hilaire, B. G. S., Yao, W., Stamenova, E., Hoeger, M., Nyquist, S. K., Korchina, V.,

Pletch, K., Flanagan, J. P., Tomaszewicz, A., McAloose, D., Estrada, C. P., Novak, B. J., … Aiden, E. L. (2018). The Juicebox Assembly Tools module facilitates de novo

assembly of mammalian genomes with chromosome-length scaffolds for under $1000. *BioRxiv*, 254797. https://doi.org/10.1101/254797

Durand, N. C., Shamim, M. S., Machol, I., Rao, S. S. P., Huntley, M. H., Lander, E. S., & Aiden, E. L. (2016). Juicer provides a one-click system for analyzing loop-resolution Hi-C

experiments. *Cell Systems*, *3*(1), 95–98. https://doi.org/10.1016/j.cels.2016.07.002

Eddy, S. R. (2011). Accelerated Profile HMM Searches. *PLOS Computational Biology*, *7*(10),

1066 e1002195. https://doi.org/10.1371/journal.pcbi.1002195

Edwards, R. J., Field, M. A., Ferguson, J. M., Dudchenko, O., Keilwagen, J., Rosen, B. D.,

1068   Johnson, G. S., Rice, E. S., Hillier, L. D., Hammond, J. M., Towarnicki, S. G., Omer, A.,

Khan, R., Skvortsova, K., Bogdanovic, O., Zammit, R. A., Aiden, E. L., Warren, W. C., &

1070   Ballard, J. W. O. (2021). Chromosome-length genome assembly and structural

variations of the primal Basenji dog (*Canis lupus familiaris*) genome. *BMC Genomics*,

1072   *22*(1), 188. https://doi.org/10.1186/s12864-021-07493-6

Edwards, R. J., Moran, N., Devocelle, M., Kiernan, A., Meade, G., Signac, W., Foy, M., Park, S.

1074   D. E., Dunne, E., Kenny, D., & Shields, D. C. (2007). Bioinformatic discovery of novel

bioactive peptides. *Nature Chemical Biology*, *3*(2), 108–112.

1076   https://doi.org/10.1038/nchembio854

Ellegren, H., Smeds, L., Burri, R., Olason, P. I., Backström, N., Kawakami, T., Künstner, A.,

1078   Mäkinen, H., Nadachowska-Brzyska, K., Qvarnström, A., Uebbing, S., & Wolf, J. B. W.

(2012). The genomic landscape of species divergence in *Ficedula* flycatchers. *Nature*,

1080   *491*(7426), 756–760. https://doi.org/10.1038/nature11584

Fambrini, M., & Pugliesi, C. (2017). *CYCLOIDEA* 2 clade genes: Key players in the control of

1082   floral symmetry, inflorescence architecture, and reproductive organ development.

Plant Molecular Biology Reporter, *35*(1), 20–36. https://doi.org/10.1007/s11105-016-

1084   1005-z

Feng, X., Zhao, Z., Tian, Z., Xu, S., Luo, Y., Cai, Z., Wang, Y., Yang, J., Wang, Z., Weng, L., Chen,

1086   J., Zheng, L., Guo, X., Luo, J., Sato, S., Tabata, S., Ma, W., Cao, X., Hu, X., … Luo, D.

(2006). Control of petal shape and floral zygomorphy in *Lotus japonicus*. *Proceedings*

1088   *of the National Academy of Sciences*, *103*(13), 4970–4975.

https://doi.org/10.1073/pnas.0600681103

1090   Field, M. A., Rosen, B. D., Dudchenko, O., Chan, E. K. F., Minoche, A. E., Edwards, R. J.,

Barton, K., Lyons, R. J., Tuipulotu, D. E., Hayes, V. M., D. Omer, A., Colaric, Z.,

1092   Keilwagen, J., Skvortsova, K., Bogdanovic, O., Smith, M. A., Aiden, E. L., Smith, T. P. L.,

Zammit, R. A., & Ballard, J. W. O. (2020). Canfam_GSD: De novo chromosome-length

1094   genome assembly of the German Shepherd Dog (*Canis lupus familiaris*) using a

combination of long reads, optical mapping, and Hi-C. *GigaScience*, *9*(giaa027).

1096   https://doi.org/10.1093/gigascience/giaa027

Forslund, K., Pereira, C., Capella-Gutierrez, S., da Silva, A. S., Altenhoff, A., Huerta-Cepas, J.,

1098   Muffato, M., Patricio, M., Vandepoele, K., Ebersberger, I., Blake, J., Fernández Breis,

J. T., Quest for Orthologs Consortium, Boeckmann, B., Gabaldón, T., Sonnhammer, E.,

1100   Dessimoz, C., Lewis, S., & Quest for Orthologs Consortium. (2018). Gearing up to

handle the mosaic nature of life in the quest for orthologs. *Bioinformatics (Oxford,*

1102   *England)*, *34*(2), 323–329. https://doi.org/10.1093/bioinformatics/btx542

Grabherr, M. G., Russell, P., Meyer, M., Mauceli, E., Alföldi, J., Di Palma, F., & Lindblad-Toh,

1104   K. (2010). Genome-wide synteny through highly sensitive sequence alignment:

Satsuma. *Bioinformatics*, *26*(9), 1145–1151.

1106   https://doi.org/10.1093/bioinformatics/btq102

Gu, Z., Gu, L., Eils, R., Schlesner, M., & Brors, B. (2014). Circlize implements and enhances

1108   circular visualization in R. *Bioinformatics*, *30*(19), 2811–2812.

https://doi.org/10.1093/bioinformatics/btu393

1110   Hibrand Saint-Oyant, L., Ruttink, T., Hamama, L., Kirov, I., Lakhwani, D., Zhou, N. N., Bourke,

P. M., Daccord, N., Leus, L., Schulz, D., Van de Geest, H., Hesselink, T., Van Laere, K.,

1112   Debray, K., Balzergue, S., Thouroude, T., Chastellier, A., Jeauffre, J., Voisine, L., …

Foucher, F. (2018). A high-quality genome sequence of *Rosa chinensis* to elucidate

1114   ornamental traits. *Nature Plants*, *4*(7), 473–484. https://doi.org/10.1038/s41477-

018-0166-1

1116   Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M. L.,

Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the genomic basis of local

1118   adaptation: Pitfalls, practical solutions, and future directions. *The American*

Naturalist, *188*(4), 379–397. https://doi.org/10.1086/688018

1120   Horn, S., Pabón-Mora, N., Theuß, V. S., Busch, A., & Zachgo, S. (2015). Analysis of the

CYC/TB1 class of TCP transcription factors in basal angiosperms and magnoliids. *The*

1122   *Plant Journal*, *81*(4), 559–571. https://doi.org/10.1111/tpj.12750

Howarth, D. G., & Donoghue, M. J. (2006). Phylogenetic analysis of the "ECE" (CYC/TB1)

1124   clade reveals duplications predating the core eudicots. *Proceedings of the National*

Academy of Sciences, *103*(24), 9101–9106.

1126   Inglis, P. W., Pappas, M. de C. R., Resende, L. V., & Grattapaglia, D. (2018). Fast and

inexpensive protocols for consistent extraction of high quality DNA and RNA from

1128   challenging plant and fungal samples for high-throughput SNP genotyping and

sequencing applications. *PLOS ONE*, *13*(10), e0206085.

1130   https://doi.org/10.1371/journal.pone.0206085

Jiang, J., Birchler, J. A., Parrott, W. A., & Kelly Dawe, R. (2003). A molecular view of plant

1132   centromeres. *Trends in Plant Science*, *8*(12), 570–575.

https://doi.org/10.1016/j.tplants.2003.10.011

1134   Johnson, L. A. S., & Briggs, B. G. (1963). Evolution in the Proteaceae. *Australian Journal of*

Botany, *11*(1), 21–61.

1136 Johnson, L. A. S., & Briggs, B. G. (1975). On the Proteaceae—The evolution and classification

of a southern family. *Botanical Journal of the Linnean Society*, *70*(2), 83–182.

1138 https://doi.org/10.1111/j.1095-8339.1975.tb01644.x

Jones, B. M., Edwards, R. J., Skipp, P. J., O'Connor, C. D., & Iglesias-Rodriguez, M. D. (2011).

1140 Shotgun proteomic analysis of *Emiliania huxleyi*, a marine phytoplankton species of

major biogeochemical importance. *Marine Biotechnology (New York, N.Y.)*, *13*(3),

1142 496–504. https://doi.org/10.1007/s10126-010-9320-0

Jordan, G. J., Carpenter, R. J., Koutoulis, A., Price, A., & Brodribb, T. J. (2015). Environmental

1144 adaptation in stomatal size independent of the effects of genome size. *New

Phytologist*, *205*(2), 608–617. https://doi.org/10.1111/nph.13076

1146 Kammonen, J. I., Smolander, O.-P., Paulin, L., Pereira, P. A. B., Laine, P., Koskinen, P., Jernvall,

J., & Auvinen, P. (2019). GapFinisher: A reliable gap filling pipeline for SSPACE-

1148 LongRead scaffolder output. *PLOS ONE*, *14*(9), e0216885.

https://doi.org/10.1371/journal.pone.0216885

1150 PVC (Research Infrastructure), UNSW Sydney. (2010). *Katana*.

https://doi.org/10.26190/669x-a286

1152 Keilwagen, J., Hartung, F., & Grau, J. (2019). GeMoMa: Homology-based gene prediction

utilizing intron position conservation and RNA-seq data. *Methods in Molecular

1154 Biology (Clifton, N.J.)*, *1962*, 161–177. https://doi.org/10.1007/978-1-4939-9173-0_9

Kolmogorov, M., Yuan, J., Lin, Y., & Pevzner, P. A. (2019). Assembly of long, error-prone

1156 reads using repeat graphs. *Nature Biotechnology*, *37*(5), 540–546.

https://doi.org/10.1038/s41587-019-0072-8

1158 Koren, S., Walenz, B. P., Berlin, K., Miller, J. R., Bergman, N. H., & Phillippy, A. M. (2017).

Canu: Scalable and accurate long-read assembly via adaptive k-mer weighting and

1160 repeat separation. *Genome Research*, gr.215087.116.

https://doi.org/10.1101/gr.215087.116

1162 Kundu, R., Casey, J., & Sung, W.-K. (2019). *HyPo: Super fast and accurate polisher for long*

read genome assemblies. https://doi.org/10.1101/2019.12.19.882506

1164 Lamesch, P., Berardini, T. Z., Li, D., Swarbreck, D., Wilks, C., Sasidharan, R., Muller, R.,

Dreher, K., Alexander, D. L., Garcia-Hernandez, M., Karthikeyan, A. S., Lee, C. H.,

1166 Nelson, W. D., Ploetz, L., Singh, S., Wensel, A., & Huala, E. (2012). The Arabidopsis

Information Resource (TAIR): Improved gene annotation and new tools. *Nucleic Acids*

1168 *Research*, *40*(D1), D1202–D1210. https://doi.org/10.1093/nar/gkr1090

Levy Karin, E., Mirdita, M., & Söding, J. (2020). MetaEuk—Sensitive, high-throughput gene

1170 discovery, and annotation for large-scale eukaryotic metagenomics. *Microbiome*,

8(1), 48. https://doi.org/10.1186/s40168-020-00808-x

1172 Lewin, H. A., Robinson, G. E., Kress, W. J., Baker, W. J., Coddington, J., Crandall, K. A., Durbin,

R., Edwards, S. V., Forest, F., Gilbert, M. T. P., Goldstein, M. M., Grigoriev, I. V.,

1174 Hackett, K. J., Haussler, D., Jarvis, E. D., Johnson, W. E., Patrinos, A., Richards, S.,

Castilla-Rubio, J. C., … Zhang, G. (2018). Earth BioGenome Project: Sequencing life for

1176 the future of life. *Proceedings of the National Academy of Sciences*, *115*(17), 4325–

4333. https://doi.org/10.1073/pnas.1720115115

1178 Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*,

34(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

1180   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G.,

Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence

1182   Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079.

https://doi.org/10.1093/bioinformatics/btp352

1184   Lindblad-Toh, K., Wade, C. M., Mikkelsen, T. S., Karlsson, E. K., Jaffe, D. B., Kamal, M., Clamp,

M., Chang, J. L., Kulbokas, E. J., & Zody, M. C. (2005). Genome sequence, comparative

1186   analysis and haplotype structure of the domestic dog. *Nature*, *438*(7069), 803–819.

Lowe, T. M., & Chan, P. P. (2016). tRNAscan-SE On-line: Integrating search and context for

1188   analysis of transfer RNA genes. *Nucleic Acids Research*, *44*(W1), W54-57.

https://doi.org/10.1093/nar/gkw413

1190   Lu-Irving, P., & Rutherford, S. (2021). *High molecular weight DNA extraction from leaf tissue*.

dx.doi.org/10.17504/protocols.io.bu9ynz7w

1192   Luo, D., Carpenter, R., Vincent, C., Copsey, L., & Coen, E. (1996). Origin of floral asymmetry in

Antirrhinum. *Nature*, *383*(6603), 794–799.

1194   Manni, M., Berkeley, M. R., Seppey, M., Simão, F. A., & Zdobnov, E. M. (2021). BUSCO

update: Novel and streamlined workflows along with broader and deeper

1196   phylogenetic coverage for scoring of eukaryotic, prokaryotic, and viral genomes.

Molecular Biology and Evolution, *38*(10), 4647–4654.

1198   https://doi.org/10.1093/molbev/msab199

Mapleson, D., Garcia Accinelli, G., Kettleborough, G., Wright, J., & Clavijo, B. J. (2017). KAT: A

1200   K-mer analysis toolkit to quality control NGS datasets and genome assemblies.

Bioinformatics, *33*(4), 574–576. https://doi.org/10.1093/bioinformatics/btw663

1202   Marçais, G., & Kingsford, C. (2011). A fast, lock-free approach for efficient parallel counting

of occurrences of k-mers. *Bioinformatics*, *27*(6), 764–770.

1204   https://doi.org/10.1093/bioinformatics/btr011

Mast, A. R., Willis, C. L., Jones, E. H., Downs, K. M., & Weston, P. H. (2008). A smaller

1206   *Macadamia* from a more vagile tribe: Inference of phylogenetic relationships,

divergence times, and diaspore evolution in *Macadamia* and relatives (tribe

1208   Macadamieae; Proteaceae). *American Journal of Botany*, *95*(7), 843–870.

Meyer, M., Munzner, T., & Pfister, H. (2009). MizBee: A multiscale synteny browser. *IEEE*

1210   *Transactions on Visualization and Computer Graphics*, *15*(6), 897–904.

https://doi.org/10.1109/TVCG.2009.167

1212   Ming, R., VanBuren, R., Liu, Y., Yang, M., Han, Y., Li, L.-T., Zhang, Q., Kim, M.-J., Schatz, M. C.,

Campbell, M., Li, J., Bowers, J. E., Tang, H., Lyons, E., Ferguson, A. A., Narzisi, G.,

1214   Nelson, D. R., Blaby-Haas, C. E., Gschwend, A. R., … Shen-Miller, J. (2013). Genome of

the long-living sacred lotus (*Nelumbo nucifera* Gaertn.). *Genome Biology*, *14*(5), 1–11.

1216   https://doi.org/10.1186/gb-2013-14-5-r41

Mirarab, S., Nguyen, N., & Warnow, T. (2011). SEPP: SATé-Enabled Phylogenetic Placement.

1218   In *Biocomputing 2012* (pp. 247–258). World Scientific.

https://doi.org/10.1142/9789814366496_0024

1220   Murat, F., Armero, A., Pont, C., Klopp, C., & Salse, J. (2017). Reconstructing the genome of

the most recent common ancestor of flowering plants. *Nature Genetics*, *49*(4), 490–

1222   496. https://doi.org/10.1038/ng.3813

Nawrocki, E. P., & Eddy, S. R. (2013). Infernal 1.1: 100-fold faster RNA homology searches.

1224   *Bioinformatics*, *29*(22), 2933–2935. https://doi.org/10.1093/bioinformatics/btt509

Nguyen, L.-T., Schmidt, H. A., von Haeseler, A., & Minh, B. Q. (2015). IQ-TREE: A fast and

1226   effective stochastic algorithm for estimating maximum-likelihood phylogenies.

Molecular Biology and Evolution, *32*(1), 268–274.

1228   https://doi.org/10.1093/molbev/msu300

Nixon, P. (1987). *The Waratah*. Kangaroo Press.

1230   Nock, C. J., Baten, A., Barkla, B. J., Furtado, A., Henry, R. J., & King, G. J. (2016). Genome and

transcriptome sequencing characterises the gene space of *Macadamia integrifolia*

1232   (Proteaceae). *BMC Genomics*, *17*(1), 937. https://doi.org/10.1186/s12864-016-3272-

1234   Nock, C. J., Baten, A., Mauleon, R., Langdon, K. S., Topp, B., Hardner, C., Furtado, A., Henry,

R. J., & King, G. J. (2020). Chromosome-scale assembly and annotation of the

1236   macadamia genome (*Macadamia integrifolia* HAES 741). *G3: Genes, Genomes,*

Genetics, *10*(10), 3497–3504. https://doi.org/10.1534/g3.120.401326

1238   Offord, C. A. (2003). Improvement of waratahs (*Telopea* spp.) through breeding. *Acta*

Horticulturae, *603*, 119–122.

1240   Offord, C. A. (2006). Analysis of characters and germplasm of significance to improvement of

Australian native waratahs (*Telopea* spp., family Proteaceae) for cut flower

1242   production. *Genetic Resources and Crop Evolution*, *53*(6), 1263–1272.

https://doi.org/10.1007/s10722-005-3487-7

1244   Offord, C. A., Nixon, P., & Goodwin, P. B. (1987). Development of the waratah as a

commercial crop. *Journal International Protea Association*, *14*, 14–15.

1246   Oliveira, L. C., & Torres, G. A. (2018). Plant centromeres: Genetics, epigenetics and evolution.

        Molecular Biology Reports, *45*(5), 1491–1497. https://doi.org/10.1007/s11033-018-

1248   4284-7

        Oxford Nanopore Technologies Ltd. (2018). *Medaka*.

1250   https://github.com/nanoporetech/medaka

        Patil, I. (2021). Visualizations with statistical details: The "ggstatsplot" approach. *Journal of*

1252   *Open Source Software*, *6*(61), 3167. https://doi.org/10.21105/joss.03167

        Phase Genomics. (2019). *Hic_qc*. https://github.com/phasegenomics/hic_qc

1254   Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing

        genomic features. *Bioinformatics*, *26*(6), 841–842.

1256   https://doi.org/10.1093/bioinformatics/btq033

        R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. R

1258   Foundation for Statistical Computing. https://www.R-project.org/

        Radwan, J., & Babik, W. (2012). The genomics of adaptation. *Proceedings of the Royal Society*

1260   *B: Biological Sciences*, *279*(1749), 5024–5028.

        https://doi.org/10.1098/rspb.2012.2322

1262   Ramsay, H. (1963). Chromosome numbers in the Proteaceae. *Australian Journal of Botany*,

        11(1), 1–20.

1264   Ranallo-Benavidez, T. R., Jaron, K. S., & Schatz, M. C. (2019). GenomeScope 2.0 and

        Smudgeplots: Reference-free profiling of polyploid genomes. *BioRxiv*, 747568.

1266   https://doi.org/10.1101/747568

        Rhie, A., McCarthy, S. A., Fedrigo, O., Damas, J., Formenti, G., Koren, S., Uliano-Silva, M.,

1268   Chow, W., Fungtammasan, A., Kim, J., Lee, C., Ko, B. J., Chaisson, M., Gedman, G. L.,

Cantin, L. J., Thibaud-Nissen, F., Haggerty, L., Bista, I., Smith, M., … Jarvis, E. D. (2021).

1270   Towards complete and error-free genome assemblies of all vertebrate species.

Nature, *592*(7856), 737–746. https://doi.org/10.1038/s41586-021-03451-0

1272   Rhie, A., Walenz, B. P., Koren, S., & Phillippy, A. M. (2020). Merqury: Reference-free quality,

completeness, and phasing assessment for genome assemblies. *Genome Biology*,

1274   *21*(1), 1–27.

Roach, M. J., Schmidt, S. A., & Borneman, A. R. (2018). Purge Haplotigs: Allelic contig

1276   reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*, *19*(1),

460. https://doi.org/10.1186/s12859-018-2485-7

1278   Rossetto, M., Allen, C. B., Thurlby, K. A. G., Weston, P. H., & Milner, M. L. (2012). Genetic

structure and bio-climatic modeling support allopatric over parapatric speciation

1280   along a latitudinal gradient. *BMC Evolutionary Biology*, *12*, 149.

https://doi.org/10.1186/1471-2148-12-149

1282   Rossetto, M., Thurlby, K. A., Offord, C. A., Allen, C. B., & Weston, P. H. (2011). The impact of

distance and a shifting temperature gradient on genetic connectivity across a

1284   heterogeneous landscape. *BMC Evolutionary Biology*, *11*, 126.

https://doi.org/10.1186/1471-2148-11-126

1286   Royal Botanic Gardens, Kew. (2017). *State of the World's Plants 2017* (No. 978-1-84246-647–

6). Royal Botanic Gardens, Kew.

1288   Sauquet, H., Weston, P. H., Anderson, C. L., Barker, N. P., Cantrill, D. J., Mast, A. R., &

Savolainen, V. (2009). Contrasted patterns of hyperdiversification in Mediterranean

1290   hotspots. *Proceedings of the National Academy of Sciences*, *106*(1), 221–225.

https://doi.org/10.1073/pnas.0805607106

1292   Schalamun, M., Nagar, R., Kainer, D., Beavan, E., Eccles, D., Rathjen, J. P., Lanfear, R., &

Schwessinger, B. (2019). Harnessing the MinION: An example of how to establish

1294   long-read sequencing in a laboratory using challenging plant tissue from *Eucalyptus*

pauciflora. *Molecular Ecology Resources*, *19*(1), 77–89.

1296   https://doi.org/10.1111/1755-0998.12938

Seehausen, O., Butlin, R. K., Keller, I., Wagner, C. E., Boughman, J. W., Hohenlohe, P. A.,

1298   Peichel, C. L., Saetre, G.-P., Bank, C., Brännström, Å., Brelsford, A., Clarkson, C. S.,

Eroukhmanoff, F., Feder, J. L., Fischer, M. C., Foote, A. D., Franchini, P., Jiggins, C. D.,

1300   Jones, F. C., … Widmer, A. (2014). Genomics and the origin of species. *Nature Reviews*

Genetics, *15*(3), 176–192. https://doi.org/10.1038/nrg3644

1302   Seemann, T. (2018). *Barrnap*. https://github.com/tseemann/barrnap

Sievers, F., Wilm, A., Dineen, D., Gibson, T. J., Karplus, K., Li, W., Lopez, R., McWilliam, H.,

1304   Remmert, M., Söding, J., Thompson, J. D., & Higgins, D. G. (2011). Fast, scalable

generation of high-quality protein multiple sequence alignments using Clustal

1306   Omega. *Molecular Systems Biology*, *7*, 539. https://doi.org/10.1038/msb.2011.75

Simão, F. A., Waterhouse, R. M., Ioannidis, P., Kriventseva, E. V., & Zdobnov, E. M. (2015).

1308   BUSCO: Assessing genome assembly and annotation completeness with single-copy

orthologs. *Bioinformatics (Oxford, England)*, *31*(19), 3210–3212.

1310   https://doi.org/10.1093/bioinformatics/btv351

Simon, L., Voisin, M., Tatout, C., & Probst, A. V. (2015). Structure and function of

1312   centromeric and pericentromeric heterochromatin in *Arabidopsis thaliana*. *Frontiers*

in Plant Science, *6*. https://doi.org/10.3389/fpls.2015.01049

1314   Soltis, P. S., & Soltis, D. E. (2014). Flower Diversity and Angiosperm Diversification. In J. L.

Riechmann & F. Wellmer (Eds.), *Flower Development: Methods and Protocols* (pp.

1316   85–102). Springer New York. https://doi.org/10.1007/978-1-4614-9408-9_4

Soltis, P. S., & Soltis, D. E. (2021). Plant genomes: Markers of evolutionary history and drivers

1318   of evolutionary change. *PLANTS, PEOPLE, PLANET*, *3*(1), 74–82.

https://doi.org/10.1002/ppp3.10159

1320   Stace, H. M., Douglas, A. W., & Sampson, J. F. (1998). Did 'Paleo-polyploidy' Really occur in

Proteaceae? *Australian Systematic Botany*, *11*(4), 613–629.

1322   https://doi.org/10.1071/sb98013

Stanke, M., & Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in

1324   eukaryotes that allows user-defined constraints. *Nucleic Acids Research*, *33*(suppl_2),

W465–W467.

1326   Sterck, L., Rombauts, S., Vandepoele, K., Rouzé, P., & Van de Peer, Y. (2007). How many

genes are there in plants (… and why are they there)? *Current Opinion in Plant*

1328   *Biology*, *10*(2), 199–203. https://doi.org/10.1016/j.pbi.2007.01.004

Stuart, K. C., Edwards, R. J., Cheng, Y., Warren, W. C., Burt, D. W., Sherwin, W. B.,

1330   Hofmeister, N. R., Werner, S. J., Ball, G. F., Bateson, M., Brandley, M. C., Buchanan, K.

L., Cassey, P., Clayton, D. F., Meyer, T. D., Meddle, S. L., & Rollins, L. A. (2021).

1332   Transcript- and annotation-guided genome assembly of the European starling.

BioRxiv, 2021.04.07.438753. https://doi.org/10.1101/2021.04.07.438753

1334   Summerell, B. A. (1997). Pests and diseases. In *The Waratah* (2nd edition). Kangaroo Press.

Summerell, B. A., Nixons, P. G., & Burgess, L. W. (1990). Crown and stem canker of waratah

1336   caused by *Cylindrocarpon destructans*. *Australasian Plant Pathology*, *19*(1), 13–15.

https://doi.org/10.1071/APP9900013

1338   Tarailo-Graovac, M., & Chen, N. (2009). Using RepeatMasker to identify repetitive elements

in genomic sequences. *Current Protocols in Bioinformatics*, *25*(1), 4.10.1-4.10.14.

1340   https://doi.org/10.1002/0471250953.bi0410s25

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate de novo genome

1342   assembly from long uncorrected reads. *Genome Research*, *27*(5), 737–746.

https://doi.org/10.1101/gr.214270.116

1344   Vurture, G. W., Sedlazeck, F. J., Nattestad, M., Underwood, C. J., Fang, H., Gurtowski, J., &

Schatz, M. C. (2017). GenomeScope: Fast reference-free genome profiling from short

1346   reads. *Bioinformatics*, *33*(14), 2202–2204.

https://doi.org/10.1093/bioinformatics/btx153

1348   Walker, B. J., Abeel, T., Shea, T., Priest, M., Abouelliel, A., Sakthikumar, S., Cuomo, C. A.,

Zeng, Q., Wortman, J., Young, S. K., & Earl, A. M. (2014). Pilon: An integrated tool for

1350   comprehensive Microbial variant detection and genome assembly improvement.

PLOS ONE, *9*(11), e112963. https://doi.org/10.1371/journal.pone.0112963

1352   Weisenfeld, N. I., Kumar, V., Shah, P., Church, D. M., & Jaffe, D. B. (2017). Direct

determination of diploid genome sequences. *Genome Research*, *27*(5), 757–767.

1354   https://doi.org/10.1101/gr.214874.116

Weston, P. H. (2006). Proteaceae. In K. Kubitzki (Ed.), *The Families and Genera of Vascular*

1356   *Plants. Volume IX* (pp. 364–404). Springer-Verlag.

Weston, P. H., & Crisp, M. D. (1994). Cladistic biogeography of waratahs (Proteaceae,

1358   Embothrieae) and their allies across the pacific. *Australian Systematic Botany*, *7*(3),

225–249. https://doi.org/10.1071/sb9940225

1360   Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Completing bacterial genome

assemblies with multiplex MinION sequencing. *Microbial Genomics*, *3*(10).

1362   https://doi.org/10.1099/mgen.0.000132

Worrall, R., & Gollnow, B. (2013). *Growing waratahs for cut flowers—A guide for commercial*

1364   *growers* (No. 12/087). Rural Industries Research and Development Corporation.

Xu, L., Dong, Z., Fang, L., Luo, Y., Wei, Z., Guo, H., Zhang, G., Gu, Y. Q., Coleman-Derr, D., Xia,

1366   Q., & Wang, Y. (2019). OrthoVenn2: A web server for whole-genome comparison and

annotation of orthologous clusters across multiple species. *Nucleic Acids Research*,

1368   *47*(W1), W52–W58. https://doi.org/10.1093/nar/gkz333

Yadav, S., Dudchenko, O., Esvaran, M., Rosen, B. D., Field, M. A., Skvortsova, K., Edwards, R.

1370   J., Gopalakrishnan, S., Keilwagen, J., Cochran, B. J., Manandhar, B., Bucknall, M.,

Bustamante, S., Rasmussen, J. A., Melvin, R. G., Omer, A., Colaric, Z., Chan, E. K. F.,

1372   Minoche, A. E., … Ballard, J. W. O. (2020). Desert Dingo (*Canis lupus dingo*) genome

provides insights into their role in the Australian ecosystem. *BioRxiv*,

1374   2020.11.15.384057. https://doi.org/10.1101/2020.11.15.384057

Yang, X., Zhao, X.-G., Li, C.-Q., Liu, J., Qiu, Z.-J., Dong, Y., & Wang, Y.-Z. (2015). Distinct

1376   regulatory changes underlying differential expression of TEOSINTE BRANCHED1-

CYCLOIDEA-PROLIFERATING CELL FACTOR genes associated with petal variations in

1378   zygomorphic flowers of *Petrocosmea* spp. Of the family Gesneriaceae. *Plant*

Physiology, *169*(3), 2138–2151.

1380 Zheng, T., Li, P., Li, L., & Zhang, Q. (2021). Research advances in and prospects of ornamental

plant genomics. *Horticulture Research*, *8*(1), 1–19. https://doi.org/10.1038/s41438-

1382 021-00499-x

Zhong, J., & Kellogg, E. A. (2015). Duplication and expression of CYC2-like genes in the origin

1384 and maintenance of corolla zygomorphy in Lamiales. *New Phytologist*, *205*(2), 852–

868.

1386 along with the raw data (ONT, 10x and Hi-C) to SRA as SRR14018636, SRR14018635 and

SRR14018634. The genome may be browsed via Apollo:

1388 https://edwapollo.babs.unsw.edu.au/apollo208/1468723/jbrowse/index.html. The NCBI

Annotation Release 100 is available at

1390 https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/018/873/765/GCF_018873765.1_Tspe_v1

and the annotation is available for browsing in

1392 GDV: https://www.ncbi.nlm.nih.gov/genome/gdv/browser/?acc=GCF_018873765.1&context

=genome.

1394

Supplementary data, was deposited to Dryad (https://doi.org/10.5061/dryad.12jm63xzt)

1396 and contains files for tracks available on the Apollo genome browser (genome, gaps,

mapped ONT and 10x reads and annotations) and the protein sequences from the GeMoMa

1398 genome annotation.

1400 Data for species used for genome annotation are available at the following repositories:

Macadamia integrifolia

1402 https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/013/358/625/GCA_013358625.1_SCU_Mint_

v3/ doi.org/10.25918/5e320fd1e5f06

1404 *Arabidopsis thaliana*

https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/001/735/GCF_000001735.4_TAIR10.1/

1406 *Rosa chinensis*

https://ftp.ncbi.nlm.nih.gov/genomes/all/GCA/002/994/745/GCA_002994745.2_RchiOBHm-

1408 V2/

Nelumbo nucifera

1410 https://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/000/365/185/GCF_000365185.1_Chinese_Lo

tus_1.1

 TABLES AND FIGURES

1414 Table 1. Library information of *Telopea speciosissima* reference genome (Tspe_v1).

Sequencing platform	Library	Median insert size (bp)	Mean read length (bp)	No. of reads	Sequence bases (Gb)
Oxford Nanopore Technologies[†]	Ligation (SQK-LSK109)	-	13,449	3,595,148	48.3
Illumina NovaSeq 6000	Paired-end 10x Chromium	336	2 x 150	822,558,750	123.4
Total gDNA	**-**	**-**	**-**	**826,153,898**	**171.7**
Illumina NextSeq 500[‡]	Phase Genomics Proximo Hi-C (Plant)	174	2 x 151	165,573,702	25.0

1416 [†] Two PromethION flow cells and two partial flow cells from a MinION pilot run
[‡] Includes a pilot iSeq run used to QC the library

1418

Table 2. Genome assembly and annotation statistics for the *Telopea speciosissima* reference
genome.

Statistic	Tspe_v1
Total length (bp)	**823,061,212**
No. of scaffolds	**1,289**
N50 (bp)	69,013,595
L50	6
No. of contigs	**1,452**
N50 (bp)	12,206,888
L50	21
N bases	18,174
GC (%)	40.11
BUSCO[†] complete (genome; n = 1,614)	**97.8 % (1,579)**
Single copy (genome)	86.7 % (1,399)
Duplicated (genome)	11.2 % (180)
BUSCO fragmented (genome)	1.7 % (27)
BUSCO missing (genome)	0.5 % (8)
Protein-coding genes	**40,158**
mRNAs	46,877
rRNAs	351
tRNAs	728
BUSCO[†] complete (proteome; n = 1,614)	**94.4 %** (1,524**)**
Single copy (proteome)	82.7 % (1,334)
Duplicated (proteome)	11.8 % (190)
BUSCO fragmented (proteome)	3.2 % (52)
BUSCO missing (proteome)	2.4 % (38)

[†] BUSCO v5 MetaEuk (embryophyta_odb10)

1422

Figure 1. New South Wales waratah (*Telopea speciosissima*). Photo taken by SH Chen.

1424

Figure 2. Assembly and annotation workflow for the *Telopea speciosissima* reference

genome Tspe_v1. Logos reproduced with permission. Waratah photo by SH Chen.

1428

Figure 5. BUSCOMP summary of BUSCO completeness rating compiled over different stages

1430 (see Figure 2) of the *Telopea speciosissima* genome assembly. The final BUSCOMP rating

uses the best rating per BUSCO gene across any of the assemblies.

1432

Figure 6. Hi-C contact matrices visualised in Juicebox.js in balanced normalisation mode a)

1434   before and b) after correction.

1436 Figure 7. DepthSizer *Telopea* assembly size prediction using read depth of BUSCO v5

MetaEuk genes a) sits between estimates from flow cytometry, Supernova and

1438 GenomeScope at mean of 904 Mb for the v1 final assembly and is b) robust to BUSCO

versions, with variation across the four adjustment methods. Dotted line represents the final

1440 assembly size.

1442   Figure 8. Genome-wide regional copy number analysis. Copy number (CN) is relative to a

single diploid (2n) copy in the genome, truncated at CN = 4. Violin plots and means

1444   generated with ggstatsplot. Each data point represents a different genomic region. BUSCO,

BUSCO v5 (MetaEuk) single-copy 'Complete' genes; Duplicated, BUSCO v5 'Duplicated'

1446   genes; Regions, NCBI gene annotations; Sequences, assembly scaffolds; Windows, 100 kb

non-overlapping windows across the genome.

1448

Figure 9. Features of the 11 chromosomes of the *Telopea speciosissima* reference genome.

1450 Concentric tracks from the outside inward represent: chromosomes, gaps (gaps of unknown

length appear as 100 bp in the assembly), GC content calculated using BEDTools v2.27.1

1452 (Quinlan & Hall, 2010), gene density and repeat density. The latter three tracks denote

values in 500 kb sliding windows. Density was defined as the fraction of a genomic window

1454 that is covered by genomic regions. Plots are white on a solid background coloured by

chromosome. Visualisation created using the R package circlize v0.4.12 (Gu et al., 2014).

1456

1458   Figure 10. Orthologous gene clusters shared among the three members of the order Proteales –

Telopea speciosissima, *Macadamia integrifolia* and *Nelumbo nucifera* – and the core eudicots –

1460   *Arabidopsis thaliana* (Brassicales) and *Rosa chinensis* (Rosales).

1462   Figure 11. Synteny between *Telopea speciosissima* (2*n* = 22) and *Macadamia integrifolia* (2*n*

= 28). Coloured squares for each species match painted chromosome regions in the other

1464   species. More detail of the underlying synteny and rearrangements can be found in Figure

S5.

1466

Figure 12. Phylogeny of *CYCLOIDEA* (*CYC*) proteins in Proteaceae, obtained from maximum-

1468   likelihood inference with IQ-TREE. Node numbers indicate bootstrap support expressed as

percentage. Scale bar represents 0.07 nucleotide substitutions per site. Branches terminate

1470   at circles; dotted extensions are for purporses only.