Research Article

Pathogenic neutrophilia drives acute respiratory distress syndrome
in severe COVID-19 patients

Running title: Pathogenic neutrophilia in severe COVID-19

Devon J. Eddins[1,2,3], Junkai Yang[1,9], Astrid Kosters[1,9], Vincent D. Giacalone[2,4], Ximo
Pechuan[5], Joshua D. Chandler[2,4], Jinyoung Eum[1,6], Benjamin R. Babcock[1], Brian S.
Dobosh[2,4], Mindy R. Hernández[7], Fathma Abdulkhader[1], Genoah L. Collins[2,4], Richard P.
Ramonell[7], Christine Moussion[5], Darya Y. Orlova[5], Ignacio Sanz[1,3,8], F. Eun-Hyung Lee[1,3,7],
Rabindra M. Tirouvanziam[2,4], Eliver E.B. Ghosn[1,2,3,6*]

[1]Lowance Center for Human Immunology, Department of Medicine, Division of Immunology and
Rheumatology, Emory University School of Medicine, Atlanta, GA 30322, USA

[2]Department of Pediatrics, Emory University School of Medicine, Atlanta, GA 30322, USA

[3]Emory Vaccine Center, Yerkes National Primate Research Center, Emory University School of
Medicine, Atlanta, GA 30322, USA

[4]Center for CF & Airways Disease Research, Children's Healthcare of Atlanta, Atlanta, 30322
GA, USA

[5]Genentech, Inc., South San Francisco, California 94080, USA

[6]Georgia Institute of Technology, Bioinformatics Graduate Program, Atlanta, GA 30332, USA

[7]Division of Pulmonary, Allergy, Critical Care & Sleep Medicine, Department of Medicine, Emory
University School of Medicine, Atlanta, GA 30322, USA

[8]Emory Autoimmunity Center of Excellence, Department of Medicine, Division of Rheumatology,
Emory University School of Medicine, Atlanta, GA 30322, USA

[9]These authors contributed equally: Junkai Yang and Astrid Kosters

__Corresponding Author Contact Info__
Eliver E.B. Ghosn (ORCID: 0000-0001-7258-906X): Lowance Center for Human Immunology,
Health Sciences Research Building, 1760 Haygood Dr. NE, E240, Atlanta, GA 30322, USA; tel:
+1-404-712-3211; email: eliver.ghosn@emory.edu

Abstract

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and the ensuing COVID-19 pandemic have caused ~40 million cases and over 648,000 deaths in the United States alone. Troubling disparities in COVID-19-associated mortality emerged early, with nearly 70% of deaths confined to Black/African-American (AA) patients in some areas, yet targeted studies within this demographic are scant. Multi-omics single-cell analyses of immune profiles from airways and matching blood samples of Black/AA patients revealed low viral load, yet pronounced and persistent pulmonary neutrophilia with advanced features of cytokine release syndrome and acute respiratory distress syndrome (ARDS), including exacerbated production of IL-8, IL-1β, IL-6, and CCL3/4 along with elevated levels of neutrophil elastase and myeloperoxidase. Circulating S100A12$^+$/IFITM2$^+$ mature neutrophils are recruited via the IL-8/CXCR2 axis, which emerges as a potential therapeutic target to reduce pathogenic neutrophilia and constrain ARDS in severe COVID-19.

Introduction

The coronavirus disease 2019 (COVID-19) pandemic, caused by SARS-CoV-2, is associated with high morbidity and mortality. To date, the pandemic has caused ~40 million cases and over 648,000 deaths in the United States alone, where communities of color are disproportionately burdened with disease severity and mortality[1, 2]. A hallmark of COVID-19 pathogenesis is the vast array of clinical presentations and outcomes, ranging from asymptomatic or mild, self-limiting disease to acute respiratory distress syndrome (ARDS), multiorgan failure, and death. Indeed, such diversity in COVID-19 pathogenesis poses challenges to identify processes that dictate progression to severe disease. Early reports highlighted systemic hyperinflammatory responses that are linked to disease severity[3]. Cytokine release syndrome, also called a cytokine storm, has been observed in many patients and is suspected of causing the detrimental progression of COVID-19 and sustained immune dysregulation[4].

Severe COVID-19 parallels the pathophysiology of sepsis[5], where clinical presentation often includes granulocytosis, elevated proinflammatory cytokine production, aberrant myeloid activation, altered dendritic cell (DC) population dynamics, and lymphopenia[6, 7]. Early single-cell analyses of bronchoalveolar lavage fluid (BALF) samples implicated dysregulated monocyte and macrophage responses as central features in poor outcomes[4, 8]. As such, most early efforts have been focused on characterizing and constraining aberrant monocyte/macrophage responses. Concurrently, reports of neutrophilia in the peripheral blood arose, and the neutrophil to lymphocyte ratio emerged as an independent risk factor for disease progression[9]. Interestingly, a prior investigation of ARDS following sepsis identified sustained neutrophilia associated with worsened prognosis and death compared to patients who resolved neutrophilia and showed an increase in tissue-resident alveolar macrophages[10], which may present another shared feature between COVID-19 and sepsis.

To date, there have been many studies of neutrophil responses in the blood of COVID-19 patients[11, 12, 13, 14, 15]. However, an in-depth analysis of the neutrophil activity in the lungs is lacking. Expressly, the extent to which neutrophils contribute to cytokine release syndrome, tissue damage, and ultimately ADRS in severe COVID-19 cases is incompletely understood. Our systems immunology approach, combining high-dimensional (Hi-D, 30-parameter) flow cytometry and multi-omics single-cell sequencing analyses of immune profiles from the airways and matching blood samples of Black/African-American (AA) patients, revealed pronounced and sustained pulmonary neutrophilia as a hallmark of severe disease. Furthermore, mature pulmonary neutrophils produce very high levels of the neutrophil chemotactic factor IL-8 in addition to IL-1β, IL-6, and CCL3/4 along with copious amounts of neutrophil elastase (NE) and myeloperoxidase (MPO). Altogether, our findings highlight that transcriptionally active and highly inflammatory neutrophils are sustained in the airways of severe patients and that reducing pathogenic neutrophilia may constrain ARDS in severe COVID-19 disease.

Results

Study Cohort

To better understand immune dynamics in Black/AA patients with severe COVID-19, we analyzed airway and matching blood samples from a cohort of 35 individuals presenting to Emory University Hospitals (severe) or the Emory Acute Respiratory Clinic (mild-acute) in Atlanta, GA, USA (Fig. 1 and Extended Data Table 1), including 8 demographic-matched healthy adults as controls. Of the 27 individuals who were confirmed positive by PCR from nasopharyngeal swabs, 18 had an NIH severity score of "critical" (https://www.covid19treatmentguidelines.nih.gov/overview/clinical-spectrum/; referred to as severe herein) and were admitted to the intensive care unit (ICU) requiring mechanical ventilator support, and 9 were mild (mild-acute) outpatients. All severe patients in our cohort received corticosteroids (dexamethasone or equivalent). Approximately half received one or more doses of the antiviral medication remdesivir, with an average ICU stay of 26 days (see Extended Data Tables 1 and 2).

Exacerbated neutrophilia in the airways and matching blood of severe COVID-19 patients

We first characterized major immune lineages in the airways (endotracheal aspiration, ETA) and matching blood samples by Hi-D flow cytometry and observed a pronounced circulating neutrophilia and lymphopenia (notably T and NK cells), which is similarly reflected in the airways (Fig. 2a-e and Extended Data Fig. 1). This is in line with recent reports showing lymphopenia associated with significant alterations in the myeloid compartment[15, 16]. Strikingly, in most cases of severe disease, ≥85% of all pulmonary leukocytes were neutrophils (Fig. 2b). This contrasts with other studies that have reported much more heterogeneity in neutrophil frequency in patients' blood and lungs[17, 18]. In addition, circulating T cells and NK cells decline with increased disease severity (Fig. 2d), which has recently been reported as a feature of COVID-19 pneumonia[19]. We also observed a notable decrease in pulmonary NK cells associated with disease severity (Fig. 2b). However, there were only subtle differences in the B-cell compartment (which were detected at very low levels in the airways) and myeloid-derived cells (MdCs) compartment compared to other reports[8, 20, 21], highlighting the importance of neutrophilia and neutrophil-to-lymphocyte ratio in our cohort. Interestingly, the patient with the lowest neutrophil frequency in the airways corresponds to the youngest patient in the severe disease group (Fig. 2b). However, since we only had a single patient under 35 years old in our severe group, we cannot conclude any associations between neutrophilia and patient age.

Since we identified prominent neutrophilia in our cohort, we developed a Hi-D, 30-parameter flow cytometry panel to interrogate inflammatory neutrophil phenotype in addition to broad leukocyte characterization (see "CoV-Neutrophil" panel in Extended Data Table 3). Here, we used extracellular staining of neutrophil elastase (NE) and CD63 (LAMP3) to assess primary granule release[22], coupled with intracellular staining for key effectors implicated in cytokine storm (IL-1β, IL-6, and IL-8). We observed that nearly all neutrophils recruited to the lung developed an inflammatory profile characterized by exacerbated levels of NE and elevated

production of IL-1β and IL-8 with a dramatic increase in primary granule release (Fig. 2c,e).
Indeed, most neutrophils in the airways were positive for all 3 intracellular cytokines and
express CD63 on the cell surface (Fig. 2c), suggesting that neutrophils are releasing cytokines
upon degranulation. However, this phenotype is diminished in the circulation. A smaller
proportion of blood neutrophils undergo granule release and express a cytokine signature
limited mainly to IL-1β production and less IL-6 and IL-8 (Fig. 2e). Similarly, we find a subset of
neutrophils in the lung that co-express the highest levels of surface CD10, CD184 (CXCR4) and
NE (Fig. 2c). Interestingly, neutrophils in the lung express lower levels of FcγRII (CD32) than
those in the blood (Extended Data Fig. 5a). Together, these results demonstrate systemic
neutrophilia in severe COVID-19 patients where neutrophils accumulating in the airways
produce exacerbated levels of potentially damaging enzymes (e.g., NE) and inflammatory
cytokines (IL-1β/6/8) while undergoing pronounced primary granule release.
144
145   **Neutrophil-secreted inflammatory molecules in the airways potentiate acute respiratory**
146   **distress syndrome in severe patients**
147
To better understand the inflammatory signaling milieu and relate the intracellular cytokine
staining to protein secretion, we measured 21 total analytes in the airways and plasma using the
Mesoscale UPLEX platform (Extended Data Table 4). We found that IL-8 is the most abundant
chemokine in the airways of severe patients accompanied by high levels of IL-1β and IL-6. This
is in line with our Hi-D flow cytometry data showing neutrophilia and pronounced intracellular
staining signal for IL-8 and IL-1β (Fig. 3a-i). Strikingly, IL-8 levels were ~100-fold higher in the
airways of severe versus mild patients and that of IL-1β and IL-6 levels in severe patients.
These findings are in stark contrast to an early report[23] that showed no differences between
circulating and pulmonary IL-6 and IL-8 secretion, reported in the <100 pg/mL range. However,
more recent studies in both ETA[24] and BALF[25] samples have noted similar findings that
corroborate our own. These data, therefore, implicate IL-8 signaling in the prominent
neutrophilia observed in severe patients in our cohort.
160
We noticed only minor differences in circulating levels of many targets between healthy
individuals and those affected by severe COVID-19 disease suggesting an uncoupling of local
versus systemic inflammatory responses[24], particularly for IL-8 and IL-1β, but also IL-6 to a
lesser extent (Fig. 3a,d,g). Conversely, we observed a rise in circulating IL-1RA levels with
increasing disease severity, similar to other reports[26]. However, we observed an opposite trend
in the airways, where IL-1RA levels appear to be decreased in severe patients (Fig. 3j). This is
in stark contrast to IL-1β levels, which were not markedly different in the circulation but
significantly increased in the airways with disease severity (Fig. 3j). We also found increased
CCL2 (MCP-1) in the airways of severe disease, and the same trend for M-CSF and TNF was
observed, albeit at much lower concentrations (Fig. 3j). Interferon (IFN)-γ was detected at the
highest levels in the airways of severe patients with no obvious differentiation in plasma across
all groups. IL-10 was also increased in the airways of severe patients with notable variation at
low levels (Fig. 3j). In contrast to other studies[27, 28], we did not observe appreciable differences
in IL-18 or CXCL10 (IP-10) levels across patient groups (Extended Data Fig. 2). Additionally,

CXCL12 was not detected in the airways of patients, and there were no notable differences in
plasma levels across patient groups (Extended Data Fig. 2).
177
We further measured the concentration of neutrophil-derived MPO and its enzymatic activity
across patient groups. MPO is an important neutrophil effector molecule (originating from
primary granules like NE) implicated in respiratory illnesses such as cystic fibrosis[29], and can
contribute to lung tissue damage during neutrophilic pneumonitis. MPO protein concentrations
increased stepwise with disease severity in the plasma and respiratory supernatant (Fig. 3k).
However, MPO activity was mostly detected in the airways, and rose with increasing disease
severity (Fig. 3k).
185
186 **Recruited airway neutrophils are mature, transcriptionally active, and further differentiate**
187 **into a highly inflammatory state**
188
To gain insight into the cellular states and transcriptional regulation in patients with severe
COVID-19 disease, we performed multi-omics scRNA-seq on cells from whole blood and ETA
samples of severe patients and whole blood from demographic-matched healthy control (Fig. 1).
We assessed immune features in data integrated from four or more patients from the same
cohort (i.e., healthy vs. severe; see Methods). First, major lineages from blood (Fig. 4a) and
ETA (Fig. 4g) were gated manually using the antibody-derived tag (ADT) data for surface
protein expression. Next, we used the gene expression (GEX) data to generate clusters for
each major lineage identified based on their surface markers, as per our previously described
SuPERR-seq pipeline[30] (Extended Data Fig. 3). For example, neutrophils identified by CD66b
and CD16 surface ADT in the blood (Fig. 4b) and ETA (Fig. 4h) were then selected and
clustered independently by GEX data (Fig. 4c,i) for further analyses.
200
A prior study had reported an increase in immature neutrophils in the lung concomitant to
increased circulating immature neutrophils from emergency hematopoiesis in severe COVID-19
patients[16]. We, therefore, assessed the immature neutrophil phenotype in the blood (Fig 4a-e)
and lung (Fig 4f-j) by scRNA-seq in our patient cohort. We readily identified a cluster of
neutrophils expressing *CAMP*, *LTF*, *RETN*, *OLFM4*, *DEFA3*, *CD24*, and *MMP8* (Fig. 4e and
Extended Data Fig. 4) in the blood of severe patients (cluster 3 in blood neutrophils) that also
has high expression of calprotectin (S100A8/9) and other calgranulins (Fig 4d). This is
consistent with immature neutrophil phenotypes described by others[7, 15] and was notably absent
in neutrophils from healthy blood, confirming an emergency hematopoiesis/granulopoiesis in our
patient cohort. However, in contrast to previous studies[16], we did not observe any signature of
immature neutrophils in the lungs of severe patients (Fig. 4j), suggesting that immature
neutrophils either are not directly recruited to the lung, or quickly differentiate upon infiltration
into the lung.
214
To determine which neutrophil subset in the blood can infiltrate the lung and further differentiate
into a pathogenic state, we explored cell-cell communication (CellChat[31]) of clustered cells from
blood (Fig. 4c) and lung (Fig. 4h) samples. We found significant communication through the
CXCL pathway between the lung neutrophil, myeloid, and non-immune populations (i.e.,

epithelial/stromal cells) with *CXCR2*-expressing neutrophils from the blood (blood cluster 2; Fig.
5a). The CXCL8 (IL-8)/CXCR2 pathway was identified as the primary recruitment axis for
circulating *CXCR2*[+] neutrophils (Fig. 5b). These data are in line with our finding that IL-8 was
increased in neutrophils by flow cytometry (Fig. 2) and secreted at very high levels in the
airways (Fig. 3) and suggests that the CXCL8/CXCR2 signaling axis is important for neutrophil
recruitment to the lungs during COVID-19 pathogenesis. Virtually all lung neutrophils
(particularly cluster 5), non-immune cells, and monocytes show the potential to recruit
circulating neutrophils (Fig. 5a,d), indicating a robust and redundant mechanism of neutrophil
recruitment to the airways via the CXCL8/CXCR2 axis. Surprisingly, the immature neutrophils
(blood cluster 3) lacked *CXCR2* (Fig. 5c), suggesting that immature neutrophils are unlikely to
infiltrate the lung via IL-8. In contrast, a defined subset of mature neutrophils in blood
expressing high levels of *CXCR2*, along with interferon-induced *IFITM2* and *S100A11/12*,
identify blood cluster 2 as the putative neutrophil subset that can infiltrate the lung via IL-8 (Figs.
4d and 5c-d). It is, therefore, probable that recent lung emigrants would still express detectable
levels of *CXCR2*, as well as *IFITM2* and *S100A11/12* (Fig. 5c).
234
After identifying the blood neutrophil cluster 2 as the putative source of lung-recruited
neutrophils, we sought to identify their cell trajectory/differentiation once they enter the inflamed
airway. Cell trajectory analyses (scVelo[32]) revealed two differentiation pathways (Trajectories 1
and 2) for the recently infiltrated neutrophils (Fig. 5e). Indeed, cells expressing the highest levels
of *CXCR2* and *S100A11/12* were at the beginning of the trajectory, further supporting these as
the putative emigrant population from the blood (Fig. 5f). Notably, neutrophils recruited to the
lung and differentiated along Trajectory 2 experience transcriptional reprogramming to acquire a
heightened inflammatory phenotype (Fig. 5e,f). In contrast to the canonical neutrophil
differentiation pathway, including a short half-life, the infiltrated neutrophils in severe COVID-19
patients are transcriptionally active (Fig. 5e-g) in comparison to blood neutrophils (Extended
Data Fig. 5b) and further differentiate into a hyperinflammatory state (Fig. 5f). This is consistent
with more recent reports of neutrophil transmigration in other respiratory illnesses such as cystic
fibrosis[33], where neutrophils undergo lung/condition-specific adaptations upon recruitment from
the circulation[34] instead of the canonical rapid and transient effector function proceeding cell
death. Indeed, we observed increased expression of the interferon-stimulated gene (ISG) *IFI30*
along with increased expression of *CCL3* (MIP-1α) and *CCL4* (MIP-1β) (Fig. 5f), which in turn
can promote the recruitment of inflammatory monocytes to the lung and/or proinflammatory
macrophage phenotypes in the airways.
253
Importantly, most neutrophils increase expression of *CXCL8* (IL-8) when recruited to the
airways (Fig. 5d), perpetuating neutrophil recruitment to the lung. These findings further
corroborate our Hi-D flow cytometry, gene expression, and secreted protein data analysis
identifying IL-8 as the most abundant neutrophil-derived chemokine (and neutrophil chemotactic
factor) in the airways of severe COVID-19 patients. Furthermore, most neutrophils increase the
expression of *CXCR4* when recruited to the airways (Fig. 5f). Interestingly, we find that *CXCR4*
expression, previously attributed to immature neutrophils[16], is increased in mature lung-recruited
neutrophils (Fig. 5f). This is consistent with *CXCR4* expression dynamics previously detected in
non-COVID lung inflammation, including in cystic fibrosis[35] and malaria[36]. Taken together, the

progressive increase in *IFI30, CCL3/4*, along with abundant *de novo CXCL8* and *CXCR4* mRNA
transcripts, reveal a transcriptionally active state in neutrophils that is poised to sustain a
hyperinflammatory milieu in the lung of severe COVID-19 patients.
266
267   **Viral load in the respiratory tract does not correlate with disease severity**
268
We then investigated whether the recruited neutrophils or other cell types (i.e., myeloid,
lymphoid, and non-immune cells) in the airways of severe patients were infected with SARS-
CoV-2. By including the SARS-CoV-2 genome sequence into our human reference
transcriptome in the multi-omics single-cell analysis, we were able to assess viral mRNA
(vRNA) transcripts at a single-cell level (Fig. 7a). Notably, we did not detect vRNA in any cell
types in the blood or airways of severe patients (Fig. 7b,c). Further, SARS-CoV-2-specific RT-
qPCR revealed that viral load was decreased in the upper airways of severe patients admitted
to the ICU compared to mild-acute patients (Fig. 7d,e). However, we did note a gene signature
in neutrophils associated with response to IFN-γ (Fig. 5g and Extended Data Fig. 5b), including
a pronounced increase in *IFITM2* (Extended Data Fig. 5c), which has been shown to promote
SARS-CoV-2 infection in human lung cells[37].
280
281
282   **Pulmonary TNF and IL-1β promote neutrophil reprogramming in the lungs**
283
To ascertain which ligand-receptor pairs are potentially responsible for the transcriptional
reprogramming of neutrophils in the lung, we first performed differential gene expression (DGE)
analysis between blood neutrophil cluster 2 and lung neutrophils to identify genes that are
upregulated in lung-recruited neutrophils (Fig. 6a). Next, we used the computational method
NicheNet[38] to identify the potential (prioritized) ligands that could induce the upregulation of the
genes identified by the DGE analysis, indicating their potential role in neutrophil reprogramming
in the lung. *TNF*, *IL1B*, and *APOE* were the highest prioritized ligands with high ligand activity
whose signaling axes have the regulatory potential to drive gene expression profiles observed in
lung-recruited neutrophils (Fig. 6b). Importantly, NicheNet analysis revealed *TNF* as the ligand
predicted to increase *BCL2A1* expression in pulmonary neutrophils, as well as *CCL4* (MIP-1β)
and *CXCL16*. Furthermore, both *TNF* and *IL1B* are the ligands predicted to induce *NFKBIA* and
CXCL8 (IL-8), and *IL1B* shows the greatest potential to induce *CCL3* (MIP-1α) expression in
recruited neutrophils. *APOE* is predicted to upregulate the expression of *FCER1G* in neutrophils
in the lung. Notably, *TNF, IL1B,* and *HMGB1* are the ligands that have the widest range of
regulatory potential, inducing neutrophil reprogramming by upregulating most genes that we
identified as differentially expressed (Fig. 6a) in lung-recruited neutrophils (Fig 6b).
300
Pulmonary T cells represent the primary cell population expressing *TNF* in the airways, where
MdCs have the highest *IL1B* and *APOE* transcripts (Fig. 6c,e). From the transcriptome data,
ligand-receptor pair analyses further identified putative signaling mediators for the top 15
prioritized ligands (Fig. 6d). *TNF* is predicted to signal through a receptor that has a *CALM1*
association (Fig. 6d). Intriguingly, it has been previously reported that CALM1 can bind other
transmembrane proteins, including ACE-2[39], and regulate their cell surface expression[40]. *IL1B* is
predicted to signal through the canonical IL1B/IL1R2 pathway, and *APOE* through the
APOE/SORL1 lipid pathway. *HMGB1* is predicted to signal through *LY96* (MD-2), which is
commonly associated with TLR4 (Fig. 6d)[41, 42, 43]. Accordingly, *IL1R2* and *CALM1* transcripts are

abundant in neutrophils from blood cluster 2, and their expression are sustained upon
recruitment to the airways (Fig. 6e).
312
Collectively, we show here that TNF- and IL-1β-mediated transcriptional reprogramming of lung
infiltrating neutrophils leads to induction of proinflammatory *NFKBIA*, *CCL3* (MIP-1α), and *CCL4*
(MIP-1β) along with *TNFAIP3/6* and *CXCL8* (Figs. 5 and 6). Of note, neutrophil-derived *CCL3*
and *CCL4* can attract inflammatory monocytes from the circulation to the airways. Interestingly,
lung-infiltrating monocytes also produce elevated levels of *CXCL8*, potentiating the recruitment
of circulating neutrophils via the CXCL8/CXCR2 axis (Fig. 5a,d). Hence, both neutrophils and
inflammatory monocytes in the airways exacerbate and potentiate pathogenic neutrophilia in the
lungs of severe patients in a positive feedback loop.

Discussion

Previous studies have reported increased neutrophilia in severe COVID-19, particularly in the circulation[14, 44, 45, 46]. In contrast, reports on exacerbated airway neutrophilia and the implication of lung neutrophils as the main cell type driving ARDS in severe patients have yielded inconclusive results[7, 8, 17, 47]. Here, we present the first comprehensive study of the lung immune response to SARS-CoV-2 in Black/AA individuals and unequivocally identify a robust and sustained airway neutrophilia associated with disease severity. The COVID-19 pandemic further highlighted some of the socio-economic and behavioral inequalities that may have contributed to troubling disparities in COVID-19-associated morbidity and mortality[48], with almost 70% of deaths being Black/AA patients in some areas[1, 2]. Although the socio-economic and behavioral differences indeed contribute to health disparities among demographics, a systematic investigation to determine the immunological features that characterize disease severity within Black/AA patients is lacking. Our systems biology approach addresses this knowledge gap and reveals new therapeutic targets to inhibit neutrophil migration, retention, and/or survival in the lung as potential effective interventions for individuals with severe disease that have been disproportionally affected by COVID-19.

Although previous studies on severe COVID-19 in humans have not been conclusive with regards to the role of airway neutrophilia in ARDS, studies in SARS-CoV-2-infected rhesus macaques and mice, where conditions and sample collection are more controlled, have identified exacerbated neutrophilia as a key immunological feature associated with disease severity[49, 50, 51]. Hence, taken together with previous studies on neutrophils[8, 14, 17, 45, 46, 47] along with animal models[49, 50, 51], our current findings provide compelling evidence that targeting exacerbated airway neutrophilia may constrain ARDS and prevent further lung damage in most patients requiring mechanical ventilator support. Although our studies represent one of the few targeted investigations in Black/AA patients, we believe our results have broader implications and may apply to most patients suffering from neutrophilic ARDS[11].

The heightened numbers of neutrophils in the lung are likely to induce and sustain inflammatory signatures by an autocrine/paracrine feedback loop among neutrophils, and paracrine signaling to other cell types that can potentiate disease severity. Notably, NE is a potent serine protease that we show is released from degranulating neutrophils in the lungs (see Fig. 2), and has potential to stimulate production of TNF, IL-1β, and IL-8[52, 53], and also abrogate protective effector functions of T cells and MdCs in the lung by cleaving cell surface receptors such as TLRs and Fc-receptors[54, 55]. Indeed, we demonstrate NE staining on the surface of T cells and MdCs in the lung (see Extended Data Fig. 5e) in addition to neutrophils, which may explain, in part, the reduction of FcγRII (CD32) expression on pulmonary neutrophils (see Extended Data Fig. 5a). Of note, the reduction in CD32 expression may prevent IgG-mediated suppression of ISG induction[14] in pulmonary neutrophils. Indeed, we demonstrate a pronounced signature for the ISG *IFI30* (see Fig. 5f). Similarly, we show by intracellular staining that pulmonary neutrophils are producing exacerbated levels of IL-1β protein that is likely released upon degranulation along with IL-8 and IL-6 (see Figs. 2 and 3). In addition to neutrophils, the myeloid lineage is also abundant in the airways and as such they are poised to influence disease

progression. For example, we and others[16] have shown that infiltrating neutrophils sustain local
production of calgranulins (S100A8/9/11/12), which can signal through and activate myeloid
cells via TLRs and RAGE receptors, compounding the already hyperinflammatory lung milieu.
Interestingly, the inflammatory monocytes that are likely recruited to the lungs via neutrophil-
secreted CCL3/4 also show elevated levels of neutrophil chemotactic factor *CXCL8* (IL-8),
which helps sustain recruitment of pathogenic neutrophils in a positive feedback loop.
371
Other immune and non-immune cell populations in the lung may contribute to COVID-19
pathogenesis in severe patients. Non-immune cells (e.g., stromal and epithelial cells) are known
targets of SARS-CoV-2 in the lung[56]. Therefore, these cells are also posited to influence
immune cell dynamics, especially at the outset of infection[57]. Indeed, our data indicate that non-
immune cells play a role in granulocyte activation (Extended Data Fig. 5d) and therefore may
initiate neutrophilia at the outset of infection[58]. Additionally, T cell-derived TNF and HMGB1,
along with myeloid-derived IL-1β, have regulatory potential to induce inflammatory gene
signatures observed in lung-recruited neutrophils. Strikingly, TNF is the ligand predicted to
increase *BCL2A1* expression in pulmonary neutrophils, which is an anti-apoptotic factor known
to regulate neutrophil survival[59, 60]. *TNF* signaling in recruited neutrophils also has the potential
to drive *CCL4* (MIP-1β) expression, along with *CXCL16*, which is another chemotactic factor
that can recruit neutrophils and macrophages via the CXCL16/CXCR6 pathway[61, 62]. In addition,
myeloid-derived *APOE* is predicted to regulate the expression of *FCER1G* in lung-recruited
neutrophils, which is associated with neutrophil activation during other viral respiratory
infections[63]. Further studies should investigate other effector functions of myeloid and lymphoid
subsets and how these cells interact with neutrophils to promote protection or pathology in the
lungs of severe COVID-19 patients.
389
Interestingly, *CXCR4* is upregulated in recruited neutrophils in severe patients (Figs. 4d,i and
5f), which may promote neutrophil survival and retention during pneumonitis and/or further
influence inflammatory neutrophil phenotype. Accordingly, CXCR4 was shown to promote
transcriptional reprogramming of neutrophils in pulmonary tissues[64]. Although we do not find
detectable levels of the canonical CXCR4 ligand (i.e., CXCL12) in the airways, we did observe
transcripts for *HMGB1* (Fig. 6), which is an alternative ligand for CXCR4. Since the IL-8/CXCR2
pathway is most notably increased and likely the primary neutrophil recruitment axis to the
airways, we speculate that CXCR4 signaling instead may promote neutrophil survival and
retention at the site of inflammation, which has been previously reported[65]. Additionally, CXCR4
signaling can stimulate de novo *CXCL8*/IL-8 production[66, 67], and has been shown to promote
neutrophil extracellular trap release during malaria disease progression[68]. Alternatively, CXCR4
is also associated with neutrophil aging and senescence[69, 70, 71]. As such, elevated CXCR4 may
be associated with prolonged neutrophil survival in COVID-19 pathogenesis. In a prior study[72],
we noted a similar pattern of CXCR4 expression on T cells from severe COVID-19 patients,
where a progressive decrease of surface CXCR4 is associated with recovery. In contrast,
patients that succumbed to the disease, showed a time-dependent escalation in CXCR4$^+$
circulating T cells concomitant to increased CXCR4$^+$ T cells in the lungs[72], implicating CXCR4
on the dysregulated lung-homing inflammatory T cells in COVID-19. Further investigation into
the potential role of CXCR4 in cell survival, retention, or recruitment may unravel novel

therapeutic targets to modulate inflammation and treat severe COVID-19. Notably, therapeutic
intervention with a CXCR4 antagonist during malaria-associated ARDS has shown significant
benefit in animal models[36].
412
Modulating inflammation through corticosteroids (particularly dexamethasone) has shown
clinical efficacy and is now the standard-of-care for patients progressing to severe COVID-19[73].
However, the pleiotropic effects of glucocorticoids and their propensity to cause neutrophilia are
well documented in asthma and chronic obstructive pulmonary disease (COPD)[74]. Although a
well-established mechanism of action for dexamethasone is via transcriptional repression of
proinflammatory cytokines, we observed very high *CXCL8* and *IL1B* transcripts concomitant
with elevated pulmonary IL-8 and IL-1β protein with prominent signatures by intracellular
staining—particularly in pulmonary neutrophils. In contrast, IL-6 levels were lower than that of
IL-8 and IL-1β in our patient cohort, and overly abundant neutrophils were not producing as
much IL-6 comparatively, which may explain, in part, why anti-IL-6/IL-6R studies failed to meet
primary endpoints[75]. Here, we provide evidence of an uncoupled cytokine profile in airway fluids
versus plasma where the lung microenvironment exhibits features of cytokine-induced ARDS
driven largely by a proinflammatory, neutrophilic feed-forward loop. Beyond COVID-19, it has
been shown that resolution of neutrophilia in ARDS has substantial prognostic benefit[10]. Taken
together, the CXCL8 (IL-8)/CXCR2 signaling axis emerges as a key potential target for next-
generation immunomodulatory therapy to reduce pathogenic neutrophilia and constrain severe
disease in patients in addition to corticosteroids.
430
We also contend that recognizing the lung pathology in severe COVID-19 to be a neutrophilic
and hyperinflammatory disease is paramount to achieve better outcomes in next-generation
therapies. Although the lung pathology in COVID-19 is initiated by a viral infection, severe
patients in the ICU no longer show signs of uncontrolled viral replication. In fact, not only did we
not detect viral transcripts by scRNA-seq within the cells in the airways, but we also noted
decreased viral burden in severe patients in the ICU versus mild-acute patients seen in the
outpatient clinic (Fig. 7). This is further supported by our previous study where we performed
plaque assays on the respiratory secretions from severe patients and revealed significantly
diminished, if any, viral plaques from the endotracheal aspirates[76]. This may explain, in part,
why antiviral drugs such as remdesivir are not able to prevent death when administered to
severe patients in ICU[77].
442
It is important to consider potentially confounding factors and limitations in our study. Though a
major strength of our study is the uniformity of sample collection in a single disease state, our
airway samples are limited to ETA, which contrasts with bronchoalveolar lavage fluid (BALF)
used in other studies[7, 17]. The ETA procedure can sample material from the medial airways—
i.e., an intermediate between the distal/lower airways (e.g., BALF) and the proximal/upper
airways (e.g., sputum or oro-/naso-pharyngeal samples). Although the relative abundance of
immune cells may vary across the upper and lower airways, previous studies observe a
correlation between paired ETA and BALF samples. In addition, these studies show that ETA
samples are not inherently neutrophilic[78], though neutrophilia/granulocytosis may be a shared
feature of non-COVID-19 and COVID-19 ARDS[10, 79]. In any event, our study is one of the only

known to date to employ integrated multi-omics single-cell investigation of immunity in
exclusively Black/AA subjects, which are disproportionately burdened with severe disease and
worse outcomes. Importantly, other studies of COVID-19 immune responses have noted similar
features of disease severity[7, 15, 16], which suggests that the findings in our target demographic
will be broadly applicable to other groups. Finally, our study was limited in scope to 35 total
subjects and single collection time points, limiting our abilities to interrogate correlations with
clinical outcomes, warranting further longitudinal studies in larger patient cohorts.
460
In conclusion, we present evidence that neutrophils are poised to be the leukocyte population
most responsible for the dysregulated hyperinflammatory response that drives ARDS in severe
COVID-19 patients. Neutrophil frequency and inflammatory profiles reveal that they are not only
the most abundant leukocyte population in the medial airways, but also major producers of
hallmark effector molecules associated with disease severity, including IL-8, IL-1β, and IL-6,
along with potent proteases such as NE and MPO, which are inherently inflammatory and
contribute to lung damage/pathology. Furthermore, we provide evidence for a neutrophil
feedforward loop where IL-8, produced by virtually all pulmonary neutrophils (and some myeloid
and stromal cells), is the primary chemokine recruiting circulating neutrophils and promoting
neutrophilia in the inflamed airways.
471
Collectively, our findings implicate neutrophilia in the immunopathophysiology of severe COVID-
19 disease where perpetual, transcriptionally active, and highly inflammatory pulmonary
neutrophils drive ARDS despite low viral burden. Thus, therapeutic intervention targeting
neutrophil recruitment/retention and/or survival/reprogramming at the site of inflammation has
the potential to constrain ARDS in severe patients, particularly those most vulnerable to
succumb to COVID-19 disease.

Methods

Ethics and biosafety. A total of 35 individuals were recruited for this study (Extended Data Tables 1 and 2). The 18 severe COVID-19+ patients were recruited from the Intensive Care Units of Emory University, Emory St. Joseph's, Emory Decatur, and Emory Midtown Hospitals. We also recruited 9 mild COVID-19-infected outpatients in the Emory Acute Respiratory Clinic and 8 healthy adults from the Emory University Hospital. All studies were approved by the Emory Institutional Review Board (IRB) under protocol numbers IRB00058507, IRB00057983, and IRB00058271. Informed consent was obtained from the patients when they had decision-making ability or from a legally authorized representative (LAR) if the patient was unable to provide consent. Blood and sputum or endotracheal aspirate (ETA) were obtained. Control blood samples were obtained from healthy adults matched by age and race. Study inclusion criteria included a confirmed COVID-19 diagnosis by PCR amplification of SARS-CoV-2 viral RNA obtained from nasopharyngeal or oropharyngeal swabs, age of 18 years or greater, and willingness to provide informed consent. Individuals with a confirmed history of COVID-19 diagnosis were excluded from the healthy donor group. All work with infectious virus and respiratory samples from COVID-19 patients was conducted inside a biosafety cabinet within the Emory Health and Safety Office (EHSO) and the United States Department of Agriculture (USDA)-approved BSL3 containment facility in the Health Sciences Research Building at Emory University following protocols approved by the Institutional Biosafety Committee (IBC) and Biosafety Officer (see ref[76]).

Patient sample collection and processing. Primary leukocytes from the airways of COVID-19 patients requiring mechanical ventilator support were collected bedside via endotracheal aspiration (ETA) and whole blood collected by standard venipuncture. Plasma from whole blood was isolated via centrifugation at 400 x g for 10 min at 4°C. To remove platelets, the isolated plasma was centrifuged at 4,000 x g for 10 min at 4°C. Untouched circulating leukocytes were isolated using the EasySep™ RBC Depletion Reagent (StemCell Technologies). ETA (from severe patients) or non-induced sputum (from mild patients) was mixed 1:1 with a 50mM EDTA solution (final concentration 25 mM EDTA) in custom RPMI-1640 media deficient in biotin, L-glutamine, phenol red, riboflavin, and sodium bicarbonate (defRPMI-1640), with 3% newborn calf serum (NBCS) and mechanically dissociated using a syringe to liberate leukocytes from mucins and other respiratory secretions. Supernatants were collected for further analysis, and then cells underwent an additional mechanical dissociation step using 1-3 mL of a 10 mM EDTA in defRPMI-1640 + 3% NBCS and a P1000 pipettor. Cells were then washed with 10 mL defRPMI-1640 + 3% NBCS, passed through a 70 μm nylon strainer, and pelleted through a 2 mL 100% NBCS layer prior to counting and downstream processing.

High-dimensional (Hi-D) 30-parameter flow cytometry. Cells (up to 10^7 total) were resuspended in defRPMI-1640 with 3% newborn calf serum and Benzonase™ (FACS buffer) in 5 mL FACS tubes and pre-incubated with GolgiStop™ (BD Biosciences) for ~60 min at 4°C. Human TruStain FcX™ was then added, followed by a 10 min incubation at RT. The 24-color extracellular staining master mix (Extended Data Table 2) was prepared 2X in BD Horizon™ Brilliant Stain Buffer to prevent staining artifacts from BD Horizon Brilliant dye interactions and

added 1:1 to cells, then incubated for 30 min at 4°C. Following staining (and total 1 h exposure
to GolgiStop™), cells were washed with ~4 mL FACS buffer. Next, the cells were resuspended
in 200 µL of BD Cytofix/Cytoperm™ fixation/permeabilization solution and incubated at 4°C for
30 min followed by a wash with ~4 mL BD Perm/Wash™ Buffer. The 4-color intracellular
staining (Extended Data Table 3) was prepared in BD Perm/Wash™ Buffer, and cells were
stained for 30 min at 4°C. Cells were washed with ~4 mL BD Perm/Wash™ Buffer, then
resuspended for a final 20 min incubation in 4% PFA and transported out of the BSL3
containment facility. Cells were washed in ~4 mL FACS buffer, then resuspended in 200-1000
µL FACS buffer for acquisition using BD FACSDiva™ Software on the Emory Pediatric/Winship
Flow Cytometry Core BD FACSymphony™ A5. To distinguish auto-fluorescent cells from cells
expressing low levels of a particular surface marker, we established upper thresholds for auto-
fluorescence by staining samples with fluorescence-minus-one (FMO) control stain sets in
which a reagent for a channel of interest is omitted. Data were analyzed with FlowJo™ v10.8
(FlowJo LLC).
536
Cell-surface antibody-derived tag (ADT) staining, single-cell encapsulation, and library
generation. Leukocytes from whole blood and ETA samples were incubated with oligo-
conjugated Ig-A/D/G/M for 10 min at 4°C followed by the addition of Human TruStain FcX™
(BioLegend) and 10 min incubation at RT. Cells were then surface stained with oligo-conjugated
monoclonal antibody panel (total 89 antibodies; Extended Data Table 5) for 30 min at 4°C,
followed by two washes in def-RPMI-1640/0.04% BSA. Cells were resuspended at a
concentration of 1200-1500 cells/µL in def-RPMI-1640/0.04% BSA and passed through a 20 or
40 µm cell strainer before loading onto a Chromium Controller (10X Genomics, Pleasanton,
CA). Cells were loaded to target encapsulation of 10,000 cells. Gene expression (GEX) and
antibody-derived tag (ADT) libraries were generated using the Chromium Single Cell 5′ Library
& Gel Bead Kit v1.1 with feature barcoding following the manufacturer's instructions. GEX
libraries were pooled and sequenced at a depth of approximately 540,000,000 reads per sample
in a single S4 flow cell and ADT libraries at a depth of approximately 79,000,000 reads per
sample in a single lane of an S4 flow cell on a NovaSeq™ 6000 (Illumina, San Diego, CA;
Extended Data Table 6)
552
Multi-omics single-cell RNA sequencing (scRNA-seq) analysis. Single-cell 5′ unique
molecular identifier (UMI) counting and barcode de-multiplexing were performed using
CellRanger Software (v.5.0.0). To detect SARS-CoV-2 viral RNA reads, we built a custom
reference genome from human GRCh38 and SARS-CoV-2 references (severe acute respiratory
syndrome coronavirus 2 isolate Wuhan-Hu-1, complete genome, GenBank MN908947.3).
Splicing-aware aligner STAR[80] was implemented to align FASTQ inputs to the reference
genome, and the resulting files are automatically filtered by CellRanger to include only cell
barcodes representing real cells. This determination is based on the distribution of UMI counts.
ADT reads were aligned to a feature reference file containing the antibody-specific barcode
sequences. To recover neutrophils, we applied our SuPERR-seq pipeline as previously
described[30]. Briefly, we recovered neutrophils from CellRanger unfiltered count matrices by
plotting surface CD16 ADT and CD66b ADT using the "FeatureScatter" function in Seurat v4.0[81]
(R version 4.0.2). The double-positive cell barcodes were then extracted and further evaluated

by GEX to confirm viable neutrophil identity. A threshold for mitochondrial content per barcode
was determined for each sample independently and applied as a cutoff to remove dead or dying
cells (Extended Data Table 7). Most samples show high cell viability with a minimal proportion of
dead cells.
570
The UMI counts of the GEX data were log-normalized by the "NormalizeData" function in Seurat
before downstream analysis, following the optimal workflow we previously described for sample
normalization and data integration[82]. Center log-ratio (CLR) transform in Seurat was performed
on ADT UMIs when recovering neutrophils from the unfiltered matrices. For surface protein
visualization to classify major lineages using our SuPERR-seq workflow[30], ADT UMIs were
normalized using the R package Denoised and Scaled by Background[83] (DSB) to remove
ambient UMI counts (i.e., background) prior to manual sequential gating by surface expression
(Extended Data Fig. 4) in SeqGeq v1.7 (FlowJo, LLC). DSB uses empty droplets to calculate
background expression, which was manually selected according to the distribution of total ADT
per cell in the raw count matrices (Extended Data Table 8). To minimize the influence from non-
informative empty droplets, we removed cell barcodes with less than 100 total ADT UMIs before
plotting the ADT distribution.
583
Before integrating the multiple datasets, we first classified major lineages in individual samples
based on a combination of gene transcript and surface protein markers (SuPERR-seq
workflow[30]) as in Fig. 4 for samples where the ADT library was of sufficient quality to allow
manual gating (Extended Data Fig. 4). Cell barcodes within each major lineage that co-
expressed markers exclusive to other major lineages were considered cell doublets and
removed (Extended Data Fig. 4). In addition, we removed cell barcodes with extremely high
total ADT UMIs, which we considered to be aggregated cells. To efficiently integrate replicate
samples, we concatenated major lineages derived from the same tissue in different donors. To
minimize batch effects and optimize data integration, we followed the data normalization and
merging strategies described previously[82]. Briefly, samples were first treated individually, and
log-normalized count matrices were scaled/Z-transformed, and the "vst" method of the Seurat
function "FindVariableFeatures" was utilized to select the top 1000 highly variable genes
(HVGs) of each sample. HVGs shared between replicate samples were used to perform
principal component analysis (PCA). To visualize the data, we performed UMAP reduction of
the first 30 PCs, and cell clustering was generated using the Leiden community detection
algorithm at a resolution of 0.8. UMAP visualizations for the integrated Blood and integrated
ETA were generated using Seurat v4 data integration workflow.
601
Receptor-ligand interaction analyses. Clustered cells from lung and blood samples from each
patient were investigated for evidence of intercellular communication using CellChat[31].
Clustered cell populations from the lung samples were combined with the blood neutrophils to
determine which lung cell populations could recruit circulating neutrophils. We utilized
NicheNet[38] to determine which ligand-receptor pairs could be responsible for the different
transcriptional states of the neutrophil populations in the lung and blood samples. We focused
on certain neutrophil clusters as the receiver populations, considering the remaining neutrophils
and other lung cell populations as the senders and thus potential interactors. The target set of

genes was determined using Seurat::FindMarkers(min.pct = 0.1), keeping only those genes with
an adjusted p-value lower than 0.05 and average log2-fold change of more than 0.2. To address
gene expression changes due to lung infiltration, we ran the algorithm using blood neutrophil
cluster 2 as the most likely candidate for lung infiltration. The genes considered were those
differentially expressed between blood cluster 2 and the lung neutrophils and genes
differentially expressed between blood cluster 2 and those cells that progress along "Trajectory
2" in the lung (see Fig.5).
617

Cell trajectory analyses. The Python toolkit scVelo[32] inferred the trajectories using biological
data of the ETA neutrophils. Input data for scVelo analysis was intron, exon, and spanning
count matrices estimated using the dropEST tool[84], then filtered with previously identified
neutrophil cell barcodes in R studio. Intron, exon, and spanning matrices were compared to
identify missing rows (genes) and were added to each matrix to equalize dimensions. The exon
matrix contained the spliced matrix, and the sum of the intron and spanning matrices constituted
the unspliced matrix. Spliced and unspliced matrices were imported with anndata library, and
pandas library was used to import gene names and cell barcodes. Raw count matrices were
added to anndata object layers as 'spliced' and 'unspliced.' Then, gene names and cell
barcodes were attached to variables and observations of anndata object, respectively. Anndata
object was transposed, followed by the regular scVelo analysis. The default parameters of
plotting velocity streams include vkey='velocity', colorbar=True, alpha=0.3, sort_order=True, and
legend_loc='on data'.
631

632   **Pathway and process enrichment analyses**. Differential gene expression (DGE) analyses
633   were performed in Seurat v4 and imported for gene annotation and further analysis using
634   Metascape[85]. DGE between neutrophils versus the total ETA were used to generate Fig. 5g and
635   non-immune cells versus total ETA were used for Extended Data Fig. 5d.
636

Mesoscale U-PLEX assays. U-PLEX Biomarker Group 1 Human Multiplex Assays (Meso
Scale Discovery) were used to evaluate levels of 21 analytes following the manufacturer's
protocol (Extended Data Table 4) in plasma and UVC-inactivated respiratory supernatants (see
ref[76]). Samples were diluted 1:5 for all assays except for IL-8, MCP-1, and IL-1RA, which were
above the upper limit of detection for the assay and were diluted 1:200 to acquire measurement
within the assay range. Final values were obtained by multiplying measurements by their
respective dilution factor.
644

Myeloperoxidase (MPO) content and activity. The abundance and activity of MPO were
quantified as previously described[29]. MPO activity and protein concentration were measured
sequentially following the immunocapture. On average, across six 96-well plate assays, lower
limits of quantification were 4.0 ng/mL (activity) and 0.84 ng/mL (protein). Samples above the
lower limit of detection but below the lower limit of quantification were imputed as half of the
latter, and those detected above the highest standard of 50 ng/mL (i.e., above the upper limit of
detection) at all dilutions were imputed as twice the standard concentration (Extended Data
Table 4).
653

SARS-CoV-2 quantitative reverse transcription PCR (RT-qPCR). Viral (v)RNA was
extracted from the respiratory secretions of COVID-19 patients using the *Quick*-RNA™ Viral Kit
(Zymo Research) following the manufacturer's protocol and complementary (c)DNA synthesized
using the High-Capacity cDNA Reverse Transcription Kit (Applied Biosystems™) per the
manufacturer's instructions, then diluted 1:5 in nuclease-free water. 10 µL diluted cDNA was
used with the NEB Luna® Universal Probe qPCR Master Mix (New England BioLabs® Inc.)
following the manufacturer's protocol and performed in 384-well plates using a QuantStudio™ 5
Real-Time PCR System (Applied Biosystems™). Primer/probe pairs were:
AGAAGATTGGTTAGATGATGATAGT (forward), TTCCATCTCTAATTGAGGTTGAACC
(reverse), and /56-FAM/TCCTCACTGCCGTCTTGTTGACCA/3IABkFQ/ (probe), which were
designed from sequences previously described[86] (Integrated DNA Technologies; IDT). To
generate a standard curve for the quantification of SARS-CoV-2 genome copies a gBlock from
IDT with the following sequence was used as a standard:
AATTAAGAACACGTCACCGCAAGAAGAAGATTGGTTAGATGATGATAGTCAACAAACTGTT
GGTCAACAAGACGGCAGTGAGGACAATCAGACAACTACTATTCAAACAATTGTTGAGGTTC
AACCTCAATTAGAGATGGAACTTACAGTTTCAGTGTTCAATTAA.
670
Statistical analyses. Statistical analyses were performed using GraphPad Prism9. Data were
analyzed for distribution (normal (Gaussian) vs. lognormal) independently using the D'Agostino
and Pearson test for normality in the untransformed and Log10-transformed data. In cases
where the sample size (N) was too small for D'Agostino and Pearson normality test, the
Shapiro-Wilk test was used to assess distribution. When data passed both distribution tests, the
likelihood of each distribution (normal vs. lognormal) was computed, and QQ-plots were
generated. When Log10 transformed data had a higher likelihood of a normal distribution
(passing normal distribution test) and/or failed lognormal distribution test, paired t-tests were
performed to compare matching blood and respiratory supernatant samples within a single
group. If the data had unequal variance (as determined by an F-test), a ratio paired t-test was
performed. All instances where lognormal distribution was likely non-parametric Wilcoxon
matched-pairs sign ranked tests were performed. For comparisons across the three patient
groups (i.e., healthy, mild-acute, severe), ordinary one-way ANOVA (if equal variance) or
Brown-Forsythe and Welch ANOVA (if unequal variance) tests were performed for data with a
normal distribution. Alternatively, data with a lognormal distribution were analyzed with a
Kruskal-Wallis test.
687
688 **Data availability.** Single-cell sequencing datasets presented here are available through NCBI
689 GEO, accession number XXX.

Author Contributions

Devon J. Eddins: Conceptualization, Methodology, Investigation, Formal Analysis, Data Curation, Writing–Original Draft, Writing–Review & Editing, Visualization. **Junkai Yang**: Methodology, Formal Analysis, Data Curation, Writing–Review & Editing, Visualization. **Astrid Kosters**: Methodology, Investigation, Data Curation, Writing–Review & Editing. **Vincent D. Giacalone**: Methodology, Formal Analysis, Investigation, Data Curation, Writing–Review & Editing. **Ximo Pechuan**: Methodology, Formal Analysis, Data Curation, Visualization, Writing– Review & Editing. **Joshua D. Chandler**: Methodology, Formal Analysis, Investigation, Data Curation, Writing–Review & Editing. **Jinyoung Eum**: Formal Analysis, Visualization. **Benjamin R. Babcock**: Investigation, Data curation. **Brian S. Dobosh**: Investigation, Data Curation, Formal Analysis. **Mindy R. Hernández**: Resources. **Fathma Abdulkhader**: Investigation. **Genoah L. Collins**: Investigation. **Richard P. Ramonell**: Resources, Data Curation, Writing– Review & Editing. **Christine Moussion**: Methodology, Formal Analysis. **Darya Y. Orlova**: Methodology, Data Curation. **Ignacio Sanz**: Resources. **F. Eun-Hyung Lee**: Methodology, Data Curation, Resources, Writing–Review & Editing. **Rabindra M. Tirouvanziam**: Methodology, Formal Analysis, Data Curation, Resources, Writing–Review & Editing. **Eliver E.B. Ghosn**: Conceptualization, Methodology, Formal Analysis, Resources, Data Curation, Writing–Original Draft, Visualization, Supervision, Project Administration**,** Funding Acquisition**.** All authors discussed the results and read and approved the final manuscript.

Competing interests

FEL is the founder of MicroB-plex, Inc., serves on the SAB of Be Bio Pharma, receives grants from BMGF and Genentech, and receives royalties from BLI, inc. All other authors have no competing interest to declare.

Acknowledgments

This study was funded by NIH/NIAID R01AI123126 (EEBG) and R01AI123126-05S1 (EEBG), the Program for Breakthrough Biomedical Research (EEBG), and the Lowance Center for Human Immunology (EEBG). DJE was partially supported by the Emory's Laney Graduate School Fellowship, and RPR was supported by the NIH T32-HL116271-07 Fellowship. The graphical abstract and Fig. 7d were generated in part using BioRender. We thank Keivan Zandi, Ann Chahroudi, Nils Schoof, Kira Moresco, and Stacy Heilman of the Department of Pediatrics (Emory University) along with the Emory Biosafety Officers Kalpana Rengarajan and Esmeralda Meyer for their assistance with setting up the BLS3 facility and kindly providing SARS-CoV-2 viral stocks. We thank Nadia Roan (Gladstone Institutes/UCSF) and Sulggi Lee (UCSF) for helpful discussions. Flow cytometry data were collected at the Emory's Pediatrics/Winship Flow Cytometry Core (access supported in part by Children's Healthcare of Atlanta). We acknowledge the Genomic Cores at Yerkes Non-Human Primate Research Center at Emory University (NIH P51 OD011132; NIH S10 OD026799), Baylor College of Medicine, and the Parker H. Petit Institute for Bioengineering and Bioscience at the Georgia Institute of Technology for the single-cell library sequences. We are grateful for the efforts of Sang N. Le, John Varghese, Anum Jalal, Saeyun Lee, and Rahul Patel, who also contributed to patient recruitment. We also thank the nurses, staff, and providers in the 71 ICU in Emory University Hospital Midtown, the medical ICU in Emory Decatur Hospital, the 5G/6G ICU in Emory University Hospital, and the ICU in Emory Saint Joseph's Hospital for their dedication and commitment during the COVID-19 pandemic. We thank all healthy individuals, patients, and their families for their participation in this study, without whom our work would not have been possible.

References

1.   Van Dyke, M.E. *et al.* Racial and ethnic disparities in COVID-19 incidence by age, sex, and period among persons aged< 25 years—16 US jurisdictions, January 1–December 31, 2020. *Morbidity and Mortality Weekly Report* **70**, 382 (2021).

2.   Romano, S.D. *et al.* Trends in Racial and Ethnic Disparities in COVID-19 Hospitalizations, by Region - United States, March-December 2020. *MMWR Morb Mortal Wkly Rep* **70**, 560-565 (2021).

3.   Lucas, C. *et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* **584**, 463-469 (2020).

4.   Merad, M. & Martin, J.C. Pathological inflammation in patients with COVID-19: a key role for monocytes and macrophages. *Nature Reviews Immunology* **20**, 355-362 (2020).

5.   Olwal, C.O. *et al.* Parallels in Sepsis and COVID-19 Conditions: Implications for Managing Severe COVID-19 Patients. *Frontiers in immunology* **12**, 91 (2021).

6.   Carvalho, T., Krammer, F. & Iwasaki, A. The first 12 months of COVID-19: a timeline of immunological insights. *Nat Rev Immunol* **21**, 245-256 (2021).

7.   Bost, P. *et al.* Deciphering the state of immune silence in fatal COVID-19 patients. *Nat Commun* **12**, 1428 (2021).

8.   Liao, M. *et al.* Single-cell landscape of bronchoalveolar immune cells in patients with COVID-19. *Nat Med* **26**, 842-844 (2020).

9.   Liang, W. *et al.* Development and Validation of a Clinical Risk Score to Predict the Occurrence of Critical Illness in Hospitalized Patients With COVID-19. *JAMA Intern Med* **180**, 1081-1089 (2020).

10.   Steinberg, K.P. *et al.* Evolution of bronchoalveolar cell populations in the adult respiratory distress syndrome. *Am J Respir Crit Care Med* **150**, 113-122 (1994).

11.   Barnes, B.J. *et al.* Targeting potential drivers of COVID-19: Neutrophil extracellular traps. *J Exp Med* **217** (2020).

12.   Kong, M., Zhang, H., Cao, X., Mao, X. & Lu, Z. Higher level of neutrophil-to-lymphocyte is associated with severe COVID-19. *Epidemiology & Infection* **148** (2020).

13.   Liu, Y. *et al.* Neutrophil-to-lymphocyte ratio as an independent risk factor for mortality in hospitalized patients with COVID-19. *Journal of Infection* (2020).

14.   Combes, A.J. *et al.* Global absence and targeting of protective immune states in severe COVID-19. *Nature* **591**, 124-130 (2021).

15.   Schulte-Schrepping, J. *et al.* Severe COVID-19 is marked by a dysregulated myeloid cell compartment. *Cell* **182**, 1419-1440 (2020).

16.   Silvin, A. *et al.* Elevated Calprotectin and Abnormal Myeloid Cell Subsets Discriminate Severe from Mild COVID-19. *Cell* **182**, 1401-1418 e1418 (2020).

17. Grant, R.A. *et al.* Circuits between infected macrophages and T cells in SARS-CoV-2
pneumonia. *Nature* **590**, 635-641 (2021).

18. Wilk, A.J. *et al.* A single-cell atlas of the peripheral immune response in patients with
severe COVID-19. *Nat Med* **26**, 1070-1076 (2020).

19. Kreutmair, S. *et al.* Distinct immunological signatures discriminate severe COVID-19
from non-SARS-CoV-2-driven critical pneumonia. *Immunity* **54**, 1578-1593 e1575
(2021).

20. Ren, X. *et al.* COVID-19 immune features revealed by a large-scale single-cell
transcriptome atlas. *Cell* **184**, 1895-1913 e1819 (2021).

21. Xu, G. *et al.* The differential immune responses to COVID-19 in peripheral and lung
revealed by single-cell RNA sequencing. *Cell Discov* **6**, 73 (2020).

22. Forrest, O.A. *et al.* Frontline Science: Pathological conditioning of human neutrophils
recruited to the airway milieu in cystic fibrosis. *J Leukoc Biol* **104**, 665-675 (2018).

23. Szabo, P.A. *et al.* Longitudinal profiling of respiratory and systemic immune responses
reveals myeloid cell-driven lung inflammation in severe COVID-19. *Immunity* **54**, 797-
814 (2021).

24. Jouan, Y., Baranek, T., Si-Tahar, M., Paget, C. & Guillon, A. Lung compartmentalization
of inflammatory biomarkers in COVID-19-related ARDS. *Critical Care* **25**, 1-3 (2021).

25. Zaid, Y. *et al.* Chemokines and Eicosanoids Fuel the Hyperinflammation Within the
Lungs of Patients with Severe COVID-19. *Journal of Allergy and Clinical Immunology*
(2021).

26. Jouan, Y. *et al.* Phenotypical and functional alteration of unconventional T cells in severe
COVID-19 patients. *J Exp Med* **217** (2020).

27. Satış, H. *et al.* Prognostic value of interleukin-18 and its association with other
inflammatory markers and disease severity in COVID-19. *Cytokine* **137**, 155302 (2021).

28. Yang, Y. *et al.* Plasma IP-10 and MCP-3 levels are highly associated with disease
severity and predict the progression of COVID-19. *Journal of Allergy and Clinical
Immunology* **146**, 119-127 (2020).

29. Chandler, J.D. *et al.* Myeloperoxidase oxidation of methionine associates with early
cystic fibrosis lung disease. *Eur Respir J* **52** (2018).

30. Xu, C. *et al.* Comprehensive multi-omics single-cell data integration reveals greater
heterogeneity in the human immune system. *bioRxiv*, 2021.2007.2025.453651 (2021).

31. Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nature
communications* **12**, 1-20 (2021).

32.   Bergen, V., Lange, M., Peidli, S., Wolf, F.A. & Theis, F.J. Generalizing RNA velocity to
transient cell states through dynamical modeling. *Nat Biotechnol* **38**, 1408-1414 (2020).
844

845   33.   Margaroli, C. *et al.* Transcriptional firing represses bactericidal activity in cystic fibrosis
846   airway neutrophils. *Cell Rep Med* **2**, 100239 (2021).
847

848   34.   Giacalone, V.D., Margaroli, C., Mall, M.A. & Tirouvanziam, R. Neutrophil Adaptations
849   upon Recruitment to the Lung: New Concepts and Implications for Homeostasis and
850   Disease. *International Journal of Molecular Sciences* **21**, 851 (2020).
851

852   35.   Makam, M. *et al.* Activation of critical, host-induced, metabolic and stress pathways
853   marks neutrophil entry into cystic fibrosis lungs. *Proc Natl Acad Sci U S A* **106**, 5779-
854   5783 (2009).
855

856   36.   Sercundes, M.K. *et al.* Targeting Neutrophils to Prevent Malaria-Associated Acute Lung
857   Injury/Acute Respiratory Distress Syndrome in Mice. *PLoS Pathog* **12**, e1006054 (2016).
858

859   37.   Prelli Bozzo, C. *et al.* IFITM proteins promote SARS-CoV-2 infection and are targets for
860   virus inhibition in vitro. *Nature Communications* **12**, 1-13 (2021).
861

862   38.   Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication
863   by linking ligands to target genes. *Nat Methods* **17**, 159-162 (2020).
864

865   39.   Chattopadhyay, S. *et al.* Calmodulin binds to the cytoplasmic domain of angiotensin-
866   converting enzyme and regulates its phosphorylation and cleavage secretion. *J Biol*
867   *Chem* **280**, 33847-33855 (2005).
868

869   40.   Lambert, D.W., Clarke, N.E., Hooper, N.M. & Turner, A.J. Calmodulin interacts with
870   angiotensin-converting enzyme-2 (ACE2) and inhibits shedding of its ectodomain. *FEBS*
871   *letters* **582**, 385-390 (2008).
872

873   41.   Shimazu, R. *et al.* MD-2, a molecule that confers lipopolysaccharide responsiveness on
874   Toll-like receptor 4. *J Exp Med* **189**, 1777-1782 (1999).
875

876   42.   Visintin, A., Mazzoni, A., Spitzer, J.A. & Segal, D.M. Secreted MD-2 is a large polymeric
877   protein that efficiently confers lipopolysaccharide sensitivity to Toll-like receptor 4. *Proc*
878   *Natl Acad Sci U S A* **98**, 12156-12161 (2001).
879

880   43.   Yang, H. *et al.* A critical cysteine is required for HMGB1 binding to Toll-like receptor 4
881   and activation of macrophage cytokine release. *Proc Natl Acad Sci U S A* **107**, 11942-
882   11947 (2010).
883

884   44.   Bohn, M.K. *et al.* Pathophysiology of COVID-19: Mechanisms Underlying Disease
885   Severity and Progression. *Physiology (Bethesda)* **35**, 288-301 (2020).
886

887   45.   Meizlish, M.L. *et al.* A neutrophil activation signature predicts critical illness and mortality
888   in COVID-19. *Blood Adv* **5**, 1164-1177 (2021).
889

890   46.   Metzemaekers, M. *et al.* Kinetics of peripheral blood neutrophils in severe coronavirus
891   disease 2019. *Clin Transl Immunology* **10**, e1271 (2021).
892

893   47.   Wauters, E. *et al.* Discriminating mild from critical COVID-19 by innate and adaptive
894   immune single-cell profiling of bronchoalveolar lavages. *Cell Res* **31**, 272-290 (2021).
895

896   48.   Abedi, V. *et al.* Racial, economic, and health inequality and COVID-19 infection in the
897   United States. *Journal of racial and ethnic health disparities* **8**, 732-742 (2021).
898

899   49.   Guo, Q. *et al.* Induction of alarmin S100A8/A9 mediates activation of aberrant
900   neutrophils in the pathogenesis of COVID-19. *Cell Host Microbe* **29**, 222-235 e224
901   (2021).
902

903   50.   Hoang, T.N. *et al.* Baricitinib treatment resolves lower-airway macrophage inflammation
904   and neutrophil recruitment in SARS-CoV-2-infected rhesus macaques. *Cell* **184**, 460-
905   475 e421 (2021).
906

907   51.   Vanderheiden, A. *et al.* CCR2-dependent monocyte-derived cells restrict SARS-CoV-2
908   infection. *bioRxiv* (2021).
909

910   52.   Krotova, K., Khodayari, N., Oshins, R., Aslanidi, G. & Brantly, M.L. Neutrophil elastase
911   promotes macrophage cell adhesion and cytokine production through the integrin-Src
912   kinases pathway. *Sci Rep* **10**, 15874 (2020).
913

914   53.   Towstyka, N.Y. *et al.* Modulation of γδ T-cell activation by neutrophil elastase.
915   *Immunology* **153**, 225-237 (2018).
916

917   54.   Domon, H. *et al.* Neutrophil Elastase Subverts the Immune Response by Cleaving Toll-
918   Like Receptors and Cytokines in Pneumococcal Pneumonia. *Front Immunol* **9**, 732
919   (2018).
920

921   55.   Kim, E. *et al.* Inhibition of elastase enhances the adjuvanticity of alum and promotes
922   anti-SARS-CoV-2 systemic and mucosal immunity. *Proc Natl Acad Sci U S A* **118**
923   (2021).
924

925   56.   Delorey, T.M. *et al.* COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular
926   targets. *Nature* **595**, 107-113 (2021).
927

928   57.   Chua, R.L. *et al.* COVID-19 severity correlates with airway epithelium-immune cell
929   interactions identified by single-cell analysis. *Nat Biotechnol* **38**, 970-979 (2020).
930

931   58.   Qi, F. *et al.* ScRNA-seq revealed the kinetic of nasopharyngeal immune responses in
932   asymptomatic COVID-19 carriers. *Cell Discov* **7**, 56 (2021).
933

934   59.   Jenal, M. *et al.* The anti-apoptotic gene BCL2A1 is a novel transcriptional target of PU.1.
935   *Leukemia* **24**, 1073-1076 (2010).
936

937   60.   Vier, J., Groth, M., Sochalska, M. & Kirschnek, S. The anti-apoptotic Bcl-2 family protein
938   A1/Bfl-1 regulates neutrophil survival and homeostasis and is controlled via PI3K and
939   JAK/STAT signaling. *Cell Death Dis* **7**, e2103 (2016).
940

941   61.   Woehrl, B. *et al.* CXCL16 contributes to neutrophil recruitment to cerebrospinal fluid in
942   pneumococcal meningitis. *J Infect Dis* **202**, 1389-1396 (2010).
943

944   62.   Zhang, L. *et al.* Chemokine CXCL16 regulates neutrophil and macrophage infiltration
945   into injured muscle, promoting muscle regeneration. *Am J Pathol* **175**, 2518-2527
946   (2009).
947

948   63.   Besteman, S.B. *et al.* Transcriptome of airway neutrophils reveals an interferon
949   response in life-threatening respiratory syncytial virus infection. *Clinical Immunology*
950   **220**, 108593 (2020).
951

952   64.   Ballesteros, I. *et al.* Co-option of Neutrophil Fates by Tissue Environments. *Cell* **183**,
953   1282-1297 e1218 (2020).
954

955   65.   Yamada, M. *et al.* The increase in surface CXCR4 expression on lung extravascular
956   neutrophils and its effects on neutrophils during endotoxin-induced lung injury. *Cell Mol*
957   *Immunol* **8**, 305-314 (2011).
958

959   66.   Bernhagen, J. *et al.* MIF is a noncognate ligand of CXC chemokine receptors in
960   inflammatory and atherogenic cell recruitment. *Nature medicine* **13**, 587-596 (2007).
961

962   67.   Pawig, L., Klasen, C., Weber, C., Bernhagen, J. & Noels, H. Diversity and Inter-
963   Connections in the CXCR4 Chemokine Receptor/Ligand Family: Molecular
964   Perspectives. *Front Immunol* **6**, 429 (2015).
965

966   68.   Rodrigues, D.A.S. *et al.* CXCR4 and MIF are required for neutrophil extracellular trap
967   release triggered by Plasmodium-infected erythrocytes. *PLoS pathogens* **16**, e1008230
968   (2020).
969

970   69.   Adrover, J.M. *et al.* A neutrophil timer coordinates immune defense and vascular
971   protection. *Immunity* **50**, 390-402 (2019).
972

973   70.   Casanova-Acebes, M. *et al.* Rhythmic modulation of the hematopoietic niche through
974   neutrophil clearance. *Cell* **153**, 1025-1035 (2013).
975

976   71.   Ng, L.G., Ostuni, R. & Hidalgo, A. Heterogeneity of neutrophils. *Nature Reviews*
977   *Immunology* **19**, 255-265 (2019).
978

979   72.   Neidleman, J. *et al.* Distinctive features of SARS-CoV-2-specific T cells predict recovery
980   from severe COVID-19. *medRxiv* (2021).
981

982   73.   Group, R.C. Dexamethasone in hospitalized patients with Covid-19. *New England*
983   *Journal of Medicine* **384**, 693-704 (2021).
984

985   74.   Ronchetti, S., Ricci, E., Migliorati, G., Gentili, M. & Riccardi, C. How glucocorticoids
986   affect the neutrophil life. *International journal of molecular sciences* **19**, 4090 (2018).
987

988   75.   Mullard, A. Anti-IL-6Rs falter in COVID-19. *Nat Rev Drug Discov* **19**, 577 (2020).
989

990   76.   Eddins, D.J. *et al.* Inactivation of SARS Coronavirus 2 and COVID-19 patient samples
991   for contemporary immunology studies. *bioRxiv* (2021).
992

993   77.   Beigel, J.H. *et al.* Remdesivir for the Treatment of Covid-19 - Final Report. *N Engl J Med*
994   **383**, 1813-1826 (2020).

996 78. Orlov, M. *et al.* Endotracheal aspirates contain a limited number of lower respiratory tract
997   immune cells. *Critical Care* **25**, 1-3 (2021).
998

999 79. Seren, S. *et al.* Proteinase release from activated neutrophils in mechanically ventilated
1000   patients with non-COVID-19 and COVID-19 pneumonia. *Eur Respir J* **57** (2021).
1001

1002 80. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
1003   (2013).
1004

1005 81. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *bioRxiv* (2020).
1006

1007 82. Babcock, B.R., Kosters, A., Yang, J., White, M.L. & Ghosn, E. Data Matrix Normalization
1008   and Merging Strategies Minimize Batch-specific Systemic Variation in scRNA-Seq Data.
1009   *bioRxiv* (2021).
1010

1011 83. Mulè, M.P., Martins, A.J. & Tsang, J.S. Normalizing and denoising protein expression
1012   data from droplet-based single cell profiling. *bioRxiv* (2020).
1013

1014 84. Petukhov, V. *et al.* dropEst: pipeline for accurate estimation of molecular counts in
1015   droplet-based single-cell RNA-seq experiments. *Genome Biol* **19**, 78 (2018).
1016

1017 85. Zhou, Y. *et al.* Metascape provides a biologist-oriented resource for the analysis of
1018   systems-level datasets. *Nat Commun* **10**, 1523 (2019).
1019

1020 86. Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus:
1021   implications for virus origins and receptor binding. *Lancet* **395**, 565-574 (2020).
1022
1023

1024

Graphical Abstract

The lung pathology due to severe COVID-19 is marked by a perpetual pathogenic neutrophilia,
leading to acute respiratory distress syndrome (ARDS) even in the absence of viral burden.
Circulating mature neutrophils are recruited to the airways via IL-8 (CXCL8)/CXCR2
chemotaxis. Recently migrated neutrophils further differentiate into a transcriptionally active and
hyperinflammatory state, with an exacerbated expression of IL-8 (*CXCL8*), IL-1β (*IL1B*), *CCL3*,
CCL4, neutrophil elastase (NE), and myeloperoxidase (MPO) activity. Airway neutrophils and
recruited inflammatory monocytes further increase their production of IL-8 (*CXCL8*),
perpetuating lung neutrophilia in a feedforward loop. MdCs and T cells produce IL-1β and TNF,
driving neutrophils reprogramming and survival.

1035

1036
1037

Figure 1. *Experimental design for the systems immunology approach (integrated multi-omics single-cell assays) to study COVID-19 patient samples*. Respiratory samples (sputum or endotracheal aspirates) and matching blood from all subjects were collected for analysis by 21-plex Mesoscale analysis, high-dimensional (Hi-D) 30-parameter flow cytometry, and multi-omics scRNA-seq. Cells from endotracheal aspirates (ETA) and blood of severe COVID-19 patients along with blood from healthy individuals were surface-stained with a panel of 89 oligo-conjugated monoclonal antibodies before single-cell encapsulation, and analyses were performed with a custom human reference genome that included the SARS-CoV-2 genome to simultaneously detect viral mRNA transcripts. Integrative multi-omics analyses were performed on the resulting data sets.

1048
1049

1050

1051
Figure 2. *Exacerbated neutrophilia in the airways and matching blood of severe COVID-*
19 patients. (a) Representative gating strategy for all samples (see Extended Data Fig. 1 for full
gating strategy). (b) Box plots show distributions of leukocytes isolated from endotracheal
aspirates (ETA). (c) Representative plots demonstrating inflammatory profile of pulmonary
neutrophils including neutrophil elastase (NE), CD184 (CXCR4), and intracellular staining of IL-
6, IL-8, and IL-1β including the full stain and fluorescence minus one (FMO) controls (d) Box
plots show distributions of leukocytes isolated from whole blood from severe patients. (e)
Representative histograms showing median fluorescence intensity (MFI) of key markers across
healthy blood (blue), severe blood (gray), and severe ETA (green) samples. MdCs = myeloid-
derived cells. For comparisons across the three patient groups (i.e., healthy, mild-acute,
severe), ordinary one-way ANOVA (if equal variance) or Brown-Forsythe and Welch ANOVA (if
unequal variance) tests were performed for data with a normal distribution. Data with a
lognormal distribution were analyzed with a Kruskal-Wallis test.
1065

1066

Figure 3. *Cytokine release syndrome is dominated by IL-8 and IL-1β with pronounced*
myeloperoxidase content and activity in the lung microenvironment. (a, d, g, j)
Concentration (pg/mL) of 15 analytes interrogated by Mesoscale analyses in plasma (gray
circles) and respiratory supernatant (Resp. SNT; green squares) from healthy control (HC),
mild-acute (MA), and severe COVID-19 patients. (b, e, h) Representative flow cytometric
intracellular staining for IL-8 (CXCL8), IL-1β, and IL-6, including the full stain and fluorescence
minus one (FMO) controls in both blood and ETA neutrophils (CD66b$^+$). (c, f, i) UMAP
visualizations of *CXCL8* (IL-8), *IL1B*, and *IL6*, which were also measured by intracellular flow
cytometry staining and Mesoscale in both blood and ETA. (k) Concentration (ng/mL) of
myeloperoxidase (MPO) protein and MPO activity in plasma vs. Resp. SNT. In (a, d, g, j, k),
black dotted line = median lower limit of detection (LLOD) for assays (see Extended Data Table
3). In (b, e, h) red dashed line indicates the median fluorescence intensity (MFI) of neutrophils in
the FMO control (value listed in the plot).
1080

1081

Figure 4. *Multi-omic single-cell RNA-seq reveals emergency granulopoiesis in the*
circulation and abundant heterogeneous populations of mature neutrophils in the
airways with distinct inflammatory states. UMAP visualization of scRNA-seq of total
integrated blood (a) and endotracheal aspirate (ETA) (f) cells. Neutrophils were identified based
on cell-surface markers (b and g), and total neutrophils were subclustered for further analysis (c
and h). Dot plots of the intersection of the top differentially expressed genes in neutrophil
clusters (d and i) sorted by average log-fold change for blood (d) and lung (i) neutrophils,
respectively. UMAP visualization of signature genes of immature neutrophils (highlighted in d
and i) in blood from severe patients compared to healthy individuals (e) and lungs of severe
patients (j).
1092

Figure 5. *Mature neutrophils are continuously recruited from circulation and progress*
toward a hyperinflammatory state. Cell receptor-ligand pair analyses (CellChat[31]) from
scRNA-seq data identify significant CXCL signaling pathway network enrichment between lung
cells and blood neutrophils (a and b). Blood neutrophil cluster 2 represents the main subset
potentially recruited to the lung (a). Recruitment of blood cluster 2 (a and c) is largely
orchestrated by the CXCL8 (IL-8)/CXCR2 axis, and to a lesser extent, by the CXCL2 and
CXCL3 (b and d). *S100A11/12, IFITM2,* and *CXCR2* mark a cluster of mature neutrophils in the
blood (c) that likely represents the neutrophil subset recruited to the lung (d). Cell trajectory
analysis (scVelo[32]) identifies two potential pathways (Trajectories 1 and 2) for recently migrated
neutrophils (e), that begin with a gene signature consistent with neutrophil blood cluster 2 (c and
f). Neutrophils recruited to the lung acquire a hyperinflammatory profile along Trajectory 2 (e
and f), characterized by high expression of interferon-stimulated gene (ISG) *IFI30* along with
macrophage inflammatory proteins *CCL3* (MIP-1α) and *CCL4* (MIP-1β), whereas *CSF3R* and
CXCR4 are increased in cells along both trajectories (f). Neutrophils along Trajectory 1 may
reflect cells progressing to apoptosis expressing higher levels of *HSPA1A* (HSP70) followed by
NEAT1 (f). Pathway and process enrichment analyses performed in Metascape[85] reveals that
myeloid-mediated immunity and platelet activation, signaling, and aggregation are significantly
enriched in neutrophils in the lungs of severe patients (g). MdCs: Myeloid-derived Cells.
1112

1113

Figure 6. *TNF and IL-1β drive inflammatory program in neutrophils recruited to the lung.*
(a) Dot plots showing 37 of the 67 differentially expressed genes in lung neutrophils versus
blood neutrophil cluster 2 sorted by average log-fold change used as input for NicheNet[38]
analyses (b) NicheNet[38] analyses identified the highest prioritized ligands (top 15) ordered by
ligand activity (y-axis) that best predict the pulmonary neutrophil gene signature (x-axis). The
predicted target genes represent the pulmonary neutrophil gene signature identified by
differential gene expression analysis between blood neutrophils from cluster 2 and ETA
neutrophils (see Extended Data Fig. 5d) (b) Dot plots of the intersection of the top 15 expressed
prioritized ligands from all cells in the ETA samples. (c) Ligand-receptor matrix of putative
signaling mediators for the top 15 prioritized ligands identified in (a). (d) UMAP visualizations of
TNF and *IL1B* transcripts in the total ETA along with expression of predicted signaling mediators
in blood (middle) and ETA (bottom) neutrophils from severe patients.

1126

1127
Figure 7. *Viral transcripts (scRNA-seq) and viral load in the airways of severe patients.*
Calu-3 cells infected with SARS-CoV-2 USA-WA1/2020 (MOI 0.4) were encapsulated for
scRNA-seq assays (a) following the same protocol used for patient cells from blood (b) and ETA
(c). Millions of viral transcripts (or unique molecular identifiers; UMIs) were detected in about
5,000 Calu-3 cells (a), but not in the cells from severe patients (total UMI = 0; b and c). RT-
qPCR (d) results of culture supernatant from Calu-3 infected cells and respiratory supernatants
from mild-acute and severe patients (e) show a lower viral load in severe patients than in mild-
acute patients and in the Calu-3-infected culture supernatant. ND = not detected; horizontal
dotted line = lower limit of detection (LLOD).

1137

1138 **Supplemental Material:**

#	Demographic	PID	Status	Sex	Age Group	Dexamethasone or equivalent	Remdesivir (No. of doses)	Date PCR positive Dx	Date intubated	Date sample collected	Length ICU stay (days)	Clinical Outcome
1	AA	PHA0001	Severe	Male	35-54	Yes	0	6/9/2020	6/8/2020	6/17/2020	32	Recovered
2	AA	PHA0004	Severe	Male	35-54	Yes	0	7/11/2020	7/18/2020	8/5/2020	37	Recovered
3	AA	PHA0006	Severe	Male	35-54	Yes	0	8/11/2020	8/16/2020	8/20/2020	8	Recovered
4	AA	PHA0018	Severe	Male	35-54	Yes	2	10/14/2020	9/29/2020	10/8/2020	13	Recovered
5	AA	PHA0041	Severe	Male	35-54	Yes	2	2/6/2021	2/6/2021	2/10/2021	14	Recovered
6	AA	PHA0003	Severe	Male	55+	Yes	2	6/23/2020	6/25/2020	6/30/2020	14	Succumbed
7	A	PHA0008	Severe	Male	55+	Yes	2	8/29/2020	8/30/2020	9/3/2020	19	Recovered
8	A	PHA0024	Severe	Male	55+	Yes	2	12/27/2020	12/31/2020	1/6/2021	9	Recovered
9	AA	PHA0025	Severe	Male	55+	Yes	0	12/20/2020	12/24/2020	1/8/2021	26	Recovered
10	AA	PHA0031	Severe	Male	55+	Yes	1	1/23/2021	1/24/2021	1/27/2021	11	Succumbed
11	AA	PHA0032	Severe	Male	55+	Yes	1	1/9/2021	1/14/2020	1/27/2021	40	Recovered
12	AA	PHA0035	Severe	Male	55+	Yes	0	1/18/2021	1/20/2021	2/2/2021	53	Recovered
13	AA	PHA0036	Severe	Female	18-34	Yes	2	1/10/2021	12/31/2020	2/2/2021	42	Recovered
14	AA	PHA0002	Severe	Female	55+	Yes	0	6/20/2020	6/20/2020	6/24/2020	25	Recovered
15	AA	PHA0005	Severe	Female	55+	Yes	0	7/16/2020	7/18/2020	8/12/2020	33	Recovered
16	AA	PHA0007	Severe	Female	55+	Yes	0	8/9/2020	8/10/2020	9/2/2020	60	Recovered
17	AA	PHA0010	Severe	Female	55+	Yes	2	9/17/2020	9/21/2020	9/24/2020	18	Recovered
18	AA	PHA0026	Severe	Female	55+	Yes	1	12/27/2020	12/28/2020	1/8/2021	16	Succumbed
19	A	PHA0040	Mild-Acute	Male	35-54	Yes	0	1/6/2021	N/A	2/8/2021	N/A	Recovered
20	AA	PHA0043	Mild-Acute	Male	35-54	Yes	0	Unclear	N/A	2/18/2021	N/A	Recovered
21	AA	PHA0023	Mild-Acute	Male	55+	Yes	0	11/6/2020	N/A	11/11/2020	N/A	Recovered
22	AA	PHA0047	Mild-Acute	Male	55+	Yes	0	2/8/2021	N/A	2/19/2021	N/A	Recovered
23	AA	PHA0012	Mild-Acute	Female	35-54	Yes	0	9/18/2020	N/A	9/29/2020	N/A	Recovered
24	AA	PHA0013	Mild-Acute	Female	35-54	Yes	0	9/21/2020	N/A	9/29/2020	N/A	Recovered
25	AA	PHA0044	Mild-Acute	Female	35-54	Yes	0	2/16/2021	N/A	2/19/2021	N/A	Recovered
26	AA	PHA0046	Mild-Acute	Female	35-54	Yes	0	2/12/2021	N/A	2/19/2021	N/A	Recovered
27	AA	PHA0045	Mild-Acute	Female	55+	Yes	0	2/12/2021	N/A	2/19/2021	N/A	Recovered
28	AA	PHA1009	Healthy	Female	18-34	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
29	AA	PHA1005	Healthy	Female	35-54	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
30	AA	PHA1006	Healthy	Female	35-54	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
31	AA	PHA1007	Healthy	Female	35-54	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
32	AA	PHA1008	Healthy	Female	35-54	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
33	AA	PHA1010	Healthy	Female	35-54	Yes	0	N/A	N/A	10/23/2020	N/A	N/A
34	AA	PHA1011	Healthy	Female	35-54	Yes	0	N/A	N/A	2/10/2021	N/A	N/A
35	AA	PHA1012	Healthy	Female	55+	Yes	0	N/A	N/A	2/17/2021	N/A	N/A

1139

Extended Data Table 1. *Demographic and clinical data from the 35 Black/African*
American individuals enrolled in our studies. Table describing demography, status, sex, and
age of study participants along with select details of clinical course including administration of
Dexamethasone and Remdesivir, date first intubated (severe, ICU patients only), date of PCR
positive diagnosis (Dx), date clinical specimens were collected, and total length ICU stay in
days, and clinical outcome (whether recovered or succumbed to disease).

1146

Characteristics	All subjects (N = 35)		
	Healthy (N = 8)	**Mild-Acute** (N=9)	**Severe** (N=18)
Age			
A. Mean age ± SD	46 ± 11	51 ± 12	56 ± 14
B. Age range	31 – 70	35 – 68	22 – 81
Gender			
A. Female	8 (100%)	5 (56%)	6 (33%)
B. Male	0 (0%)	4 (44%)	12 (67%)
Site of endotracheal aspiration			
A. Endotracheal tube	—	—	9 (50%)
B. Tracheostomy	—	—	9 (50%)
Medical History			
A. Smoking History			
i) Never	7 (87.5%)	7 (78%)	12 (67%)
ii) Former	1 (12.5%)	2 (22%)	2 (11%)
iii) Current	0 (0%)	0 (0%)	4 (22%)
B. Hypertension (HTN)	3 (37.5%)	3 (33%)	11 (61%)
C. Obesity	2 (25%)	1 (11%)	4 (22%)
D. Asthma	2 (25%)	2 (22%)	2 (11%)
E. COPD	0 (0%)	0 (0%)	2 (11%)
F. Diabetes mellitus (DM)			
i) Type I DM	0 (0%)	0 (0%)	0 (0%)
ii) Type II DM	1 (12.5%)	0 (0%)	8 (44%)
G. Deep vein thrombosis (DVT)	0 (0%)	0 (0%)	1 (5%)
H. Hypothyroidism	0 (0%)	0 (0%)	1 (5%)
I. Congestive Heart Failure (CHF)	1 (12.5%)	0 (0%)	5 (28%)
J. Gastroesophegeal reflux disease (GERD)	1 (12.5%)	1 (11%)	1 (5%)
K. Dyslipidemia	0 (0%)	1 (11%)	1 (5%)
L. Obstructive sleep apnea (OSA)	0 (0%)	0 (0%)	2 (11%)
M. Coronary artery disease (CAD)	0 (0%)	0 (0%)	1 (5%)
N. Chronic kidney disease	0 (0%)	1 (11%)	1 (5%)
O. Cancer	0 (0%)	0 (0%)	0 (0%)
P. Autoimmune hemolytic anemia	0 (0%)	0 (0%)	1 (5%)
Q. Thrombocytopenia	0 (0%)	0 (0%)	1 (5%)
R. Athritis/Rheumatoid arthritis (RA)	1 (12.5%)	0 (0%)	1 (5%)
Complications			
A. Acute respiratory distress syndrome (ARDS)	—	—	18 (100%)
B. Hypoxemic respiratory failure	—	—	13 (72%)
C. Extracorpeal membrane oxygenation (ECMO)	—	—	2 (11%)
D. Shock	—	—	15 (83%)
E. Nosocomial pneumonia	—	—	6 (33%)
F. Coagulopathy	—	—	3 (17%)
G. Seizures	—	—	3 (17%)
H. Cardiac arrythmia	—	—	3 (17%)
I. Renal failure	—	—	8 (44%)
J. Renal replacement therapy	—	—	6 (33%)

1147

1148 **Extended Data Table 2.** *Clinical characteristics of the 35 Black/African American*
1149 *individuals enrolled in our studies.* Table describing clinical characteristics of 35 patients
1150 across the 3 cohorts (healthy, mild-acute, and severe) including age, gender, site of
1151 endotracheal aspiration (ETA), medical history, along with complications for the severe patients
1152 admitted to the ICU.
1153

a

CoV-Neutrophil

	Laser	Detector	Antigen	Fluorophore	Clone	Isotype	Vendor	Cat. No.	Titration
		B 488/10	Side Scatter (SSC)	—	—	—	—	—	—
1		B 515/20	Neutrophil elastase (NE)	AF488	950318	Mouse IgG1, κ	R&D Systems	IC91671G	1:25
2		B 610/20	CD45-RA	BB630-P	HI-100	Mouse IgG2a, κ	BD Biosciences	624294	1:160
3	488 nm	B 670/30	CD11b	BB660-P	ICRF44	Mouse IgG1, κ	BD Biosciences	624295	1:50
4		B 710/50	CD45	BB700	HI-30	Mouse IgG1, κ	BD Biosciences	746090	1:640
5		B 750/30	CD4	BB755-P	RPA-T4	Mouse IgG1, κ	BD Biosciences	624391	1:100
6		B 780/60	HLA-DR (MHC II)	BB790-P	G46-6	Mouse IgG2a, κ	BD Biosciences	624296	1:50
7		V 431/28	CD163	BV421	GHI/6I	Mouse IgG1, κ	BioLegend	333612	1:25
8		V 470/15	IL-1β	Pacific Blue	H1b-98	Mouse IgG1, κ	BioLegend	511710	1:20
9		V 586/15	CD56	BV570	HCD56	Mouse IgG1, κ	BioLegend	318330	1:100
10	405 nm	V 610/20	CD86 *	BV605	FUN-1	Mouse IgG1, κ	BD Biosciences	562999	1:50
11		V 670/30	TNF	BV650	MAb11	Mouse IgG1, κ	BD Biosciences	563418	1:80
12		V 710/50	CD19	BV711	HIB19	Mouse IgG1, κ	BioLegend	302246	1:20
13		V 740/35	CD16 (FcγRIII)	BV750	3G8	Mouse IgG1, κ	BD Biosciences	747461	1:320
14		V 780/60	CD63 (LAMP-3)	BV785	H5C6	Mouse IgG1, κ	BioLegend	353044	1:80
15		YG 586/15	IL-6	PE	MQ2-13A5	Rat IgG1, κ	BD Biosciences	554545	1:20
16		YG 610/20	IL-8 (CXCL8)	PE-CF594	G265-8	Mouse IgG2b, κ	BD Biosciences	563531	1:20
17	561 nm	YG 670/30	Siglec-8 *	PE-Cy5	7C9	Mouse IgG1, κ	BioLegend	347114	1:100
18		YG 710/50	CD10	PE-Cy5.5	HI10a	Mouse IgG1, κ	Homemade	—	1:40
19		YG 780/60	CD184 *	PE-Cy7	12G5	Mouse IgG2a, κ	BioLegend	306514	1:40
20		UV 379/28	CD64 (FcγRI)	BUV395	10.1	Mouse IgG1, κ	BD Biosciences	740300	1:80
21		UV 450/50	GhostDye (Viability)	UV450	—	—	Tonbo Biosciences	13-0868-T500	1:100
22	355 nm	UV 586/15	CD172a/b (SIRPα/β1)	BUV563	SE5A5	Mouse IgG1, κ	BD Biosciences	748435	1:10
23		UV 670/30	CD123 (IL-3Rα)	BUV661	9F5	Mouse IgG1, κ	BD Biosciences	741628	1:40
24		UV 740/35	CD14	BUV737	M5E2	Mouse IgG2a, κ	BD Biosciences	612763	1:50
25		UV 820/60	CD3ε	BUV805	SK1	Mouse IgG1, κ	BD Biosciences	612889	1:40
26		R 670/30	CD66b	APC	QA17A51	Mouse IgG1, κ	BioLegend	396906	1:200
27	640 nm	R 710/50	CD11c	AF700	Bu15	Mouse IgG1, κ	BioLegend	337220	1:200
28		R 780/60	CD32 (FcγRII)	APC-Fire750	FUN-2	Mouse IgG2b, κ	BioLegend	303220	1:400

* = alternate reagents used in panel iterations (see c below); -P denotes prototype reagents

b

CoV-Global Distributions

	Laser	Detector	Antigen	Fluorophore	Clone	Isotype	Vendor	Cat. No.	Titration
	488 nm	B 488/10	Side Scatter (SSC)	—	—	—	—	—	—
1		B 515/20	CD45	AF488	HI-30	Mouse IgG1, κ	BioLegend	304017	1:160
2		V 431/28	CD56	BV421	HCD56	Mouse IgG1, κ	BioLegend	318328	1:40
3		V 610/20	CD8α	BV605	SK1	Mouse IgG1, κ	BioLegend	344742	1:20
4	405 nm	V 670/30	HLA-DR (MHC II)	BV650	L243	Mouse IgG2a, κ	BioLegend	307650	1:50
5		V 710/50	CD4	BV711	RPA-T4	Mouse IgG1, κ	BioLegend	300558	1:40
6		V 740/35	CD16 (FcγRIII)	BV750	3G8	Mouse IgG1, κ	BD Biosciences	747461	1:320
7		V 780/60	CD19	BV785	HIB19	Mouse IgG1, κ	BioLegend	302240	1:40
8		YG 586/15	CD123 (IL-3Rα)	PE	6H6	Mouse IgG1, κ	BioLegend	306006	1:400
9	561 nm	YG 670/30	Siglec-8	PE-Cy5	7C9	Mouse IgG1, κ	BioLegend	347114	1:100
10		YG 710/50	CD14	PE-Fire700	63D3	Mouse IgG1, κ	BioLegend	367158	1:100
11		YG 780/60	CD11c	PE-Cy7	Bu-15	Mouse IgG1, κ	BioLegend	337216	1:160
12	355 nm	UV 450/50	GhostDye (Viability)	UV450	—	—	Tonbo Biosciences	13-0868-T500	1:100
13		R 670/30	CD11b	APC	ICRF44	Mouse IgG1, κ	BioLegend	301350	1:400
14	640 nm	R 710/50	CD3ε	AF700	HIT3a	Mouse IgG2a, κ	BioLegend	300324	1:320
15		R 780/60	CD66b	APC-Fire750	QA17A51	Mouse IgG1, κ	BioLegend	396908	1:80

c

Alternate Reagents

	Laser	Detector	Antigen	Fluorophore	Clone	Isotype	Vendor	Cat. No.	Titration
1	405 nm	V670/30	CD169 (Siglec-1)	BV605	7-239	Mouse IgG1, κ	BioLegend	346010	1:40
2		V670/30	CD182 (CXCR2)	BV605	6C6	Mouse IgG1, λ	BD Biosciences	744197	1:40
3	561 nm	YG 670/30	CD86	PE-Cy5	IT2.2	Mouse IgG1, κ	BioLegend	305408	1:20
4		YG 780/60	CD206 (MMR)	PE-Cy7	15-2	Mouse IgG2a, κ	BioLegend	321124	1:10

Extended Data Table 3. *High-dimensional 30-parameter, including intracellular cytokine*
staining, and 17-parameter flow cytometry panels used for airway and blood cells. Flow
cytometer configuration and cytometry reagents used in the final panels to interrogate (a)
neutrophil phenotype and (b) global immune cell distributions and blood and ETA samples.
Channels marked with an asterisk (*) denote alternate reagents (c) used in earlier panel
iterations for some samples in this study. Monoclonal antibody (mAb) master mixes were
prepared in BD Horizon™ Brilliant Stain Buffer and samples stained as described in the
methods section. Titrations of all reagents were determined in house for each lot independently
prior to use. AF: AlexaFluor, APC: Allophycocyanin, BB: Brilliant Blue, BUV: Brilliant Ultraviolet,
BV: Brilliant Violet, FITC: Fluorescein isothiocyanate, PE: Phycoerythrin. Fluorophores marked
with -P denote prototype reagents and are custom conjugations from BD Biosciences.
1167

a

	Analyte	Median LLOD (pg/mL)	LLOD Range (pg/mL)
1	G-CSF (CSF3)	0.84 ± 0.88	0.15—2.12
2	GM-CSF (CSF2)	0.11 ± 0.13	0.01—0.31
3	IFN-γ	1.31 ± 1.12	0.16—2.79
4	IL-10	0.03 ± 0.02	0.01—0.06
5	IL-1α	0.13 ± 0.07	0.04—0.19
6	IL-1β	0.07 ± 0.05	0.01—0.12
7	IL-6	0.31 ± 0.39	0.03—0.89
8	IL-8 (CXCL8)	0.03 ± 0.03	0.02—0.08
9	TNF	0.24 ± 0.13	0.07—0.35
10	VEGFA	0.43 ± 0.44	0.06—1.06
11	ENA-78 (CXCL5)	0.15 ± 0.07	0.08—0.23
12	IL-18	1.02 ± 1.31	0.14—2.92
13	IL-1RA	1.08 ± 0.23	0.83—1.28
14	IL-29 (IFNλ1)	0.28 ± 0.11	0.13—0.38
15	IP-10 (CXCL10)	0.70 ± 0.99	0.18—2.19
16	I-TAC (CXCL11)	1.82 ± 0.23	1.59—2.13
17	MCP-1 (CCL2)	0.14 ± 0.05	0.09—0.21
18	M-CSF (CSF1)	0.10 ± 0.04	0.06—0.14
19	MIP-1β (CCL4)	5.17 ± 2.40	2.78—8.51
20	TRAIL (CD253)	0.28 ± 0.11	0.18—0.44
21	SDF1α (CXCL12)	30.13 ± 26.73	10.60—60.60

b

	Assay	Median LLOD (ng/mL)	LLOD Range (ng/mL)	Median LLOQ (ng/mL)	LLOQ Range (ng/mL)
1	Total MPO	0.15 ± 0.15	0.03—0.42	0.30 ± 0.30	0.05—0.84
2	MPO Activity	2.86 ± 1.98	0.38—6.03	5.72 ± 3.96	0.76—12.07

1168
Extended Data Table 4. *Mesoscale U-PLEX biomarker group 1 human and*
myeloperoxidase (MPO) assays. Median ± SD and range of lower limits of detection (LLOD)
for (a) Mesoscale UPLEX analytes and (b) MPO assays. Lower limits of quantification (LLOQ;
MPO assays) and LODs (determined by standard curve) were run for each assay plate
independently and the median LLOD plotted as horizontal dotted lines for each assay (see Fig.
3 and Extended Data Fig. 2).

	Antigen	Clone	Vendor	Cat. No.		Antigen	Clone	Vendor	Cat. No.
1	CD10	HI10a	Biolegend	312233	46	CD324	67A4	Biolegend	324127
2	CD101 (BB27)	BB27	Biolegend	331017	47	CD326 (EpCAM)	9C4	Biolegend	324247
3	CD103 (Integrin αE)	Ber-ACT8	Biolegend	350233	48	CD335 (NKp46)	9.00E+02	Biolegend	331941
4	CD107a (LAMP-1)	H4A3	Biolegend	328649	49	CD370 (CLEC9A)	8F9	Biolegend	353809
5	CD117 (c-kit)	104D2	Biolegend	313243	50	CD38	HIT2	Biolegend	303543
6	CD11b	ICRF44	Biolegend	301359	51	CD4	RPA-T4	Biolegend	300567
7	CD11c	S-HCL-3	Biolegend	371521	52	CD40	5C3	Biolegend	334348
8	CD123 (IL-3Rα)	6H6	Biolegend	306045	53	CD41	HIP8	Biolegend	303739
9	CD127 (IL-7Rα)	A019D5	Biolegend	351356	54	CD44	BJ18	Biolegend	338827
10	CD133	S16016B	Biolegend	394007	55	CD45	2D1	Biolegend	368545
11	CD137	4B4-1	Biolegend	309839	56	CD45R	RA3-6B2	Biolegend	103273
12	CD138 (Syndecan-1)	MI15	Biolegend	356539	57	CD45RA	HI100	Biolegend	304163
13	CD14	M5E2	Biolegend	301859	58	CD45RO	UCHL1	Biolegend	304259
14	CD15 (SSEA-1)	W6D3	Biolegend	323053	59	CD49a	TS2/7	Biolegend	328319
15	CD16 (FcγRIII)	3G8	Biolegend	302065	60	CD5	UCHT2	Biolegend	300637
16	CD161	HP-3G10	Biolegend	339947	61	CD56 (NCAM)	QA17A16	Biolegend	392425
17	CD163	GHI/61	Biolegend	333637	62	CD62L	DREG-56	Biolegend	304851
18	CD169	7-239	Biolegend	346021	63	CD64 (FcγRI)	10.1	Biolegend	305045
19	CD183 (CXCR3)	G025H7	Biolegend	353747	64	CD66b	6/40c	Biolegend	392909
20	CD184 (CXCR4)	12G5	Biolegend	306533	65	CD69	FN50	Biolegend	310951
21	CD185 (CXCR5)	J252D4	Biolegend	356939	66	CD80	2D10	Biolegend	305243
22	CD19	HIB19	Biolegend	302265	67	CD86	IT2.2	Biolegend	305447
23	CD196 (CCR6)	G034E3	Biolegend	353440	68	CD8α	RPA-T8	Biolegend	301071
24	CD197 (CCR7)	G043H7	Biolegend	353251	69	CD95 (Fas)	DX2	Biolegend	305651
25	CD1c	L161	Biolegend	331547	70	CX3CR1	K0124E1	Biolegend	355705
26	CD1d	51.1	Biolegend	350319	71	HLA-DR	L243	Biolegend	307663
27	CD20	2H7	Biolegend	302363	72	IgD	IA6-2	Biolegend	348245
28	CD206 (MMR)	15-2	Biolegend	321147	73	IgG	M1310G05	Biolegend	410727
29	CD209 (DC-SIGN)	9E9A8	Biolegend	330121	74	IgM	MHM-88	Biolegend	314547
30	CD21	Bu32	Biolegend	354923	75	Ig light chain κ	MHK-49	Biolegend	316533
31	CD23	EBVCS-5	Biolegend	338525	76	Ig light chain λ	MHL-38	Biolegend	316629
32	CD24	ML5	Biolegend	311143	77	Siglec-8	7C9	Biolegend	347115
33	CD25	BC96	Biolegend	302649	78	TCR Vα7.2	3C10	Biolegend	351735
34	CD269 (BCMA)	19F2	Biolegend	357523	79	TCR γ/δ	B1	Biolegend	331231
35	CD27	O323	Biolegend	302853	80	TIGIT (VSTM3)	A15153G	Biolegend	372729
36	CD274 (PD-L1)	29E.2A3	Biolegend	329751	81	XCR1	S15046E	Biolegend	372617
37	CD275 (ICOSL)	2D3	Biolegend	309419		*Custom oligo-conjugated in-house*			
38	CD279 (PD-1)	EH12.2H7	Biolegend	329963	82	ACE2	535919	Sino Biological	MAB9332-100
39	CD28	CD28.2	Biolegend	302963	83	CD160	BY55	Biolegend	341202
40	CD29 (Integrin β1)	TS2/16	Biolegend	303029	84	CD16b	2D2G5B9	Sino Biological	11046-MM01
41	CD3	UCHT1	Biolegend	300479	85	CD178	NOK-1	Biolegend	306402
42	CD307d (FcRL4)	413D12	Biolegend	340213	86	CD34	581	Biolegend	343602
43	CD307e (FcRL5)	509f6	Biolegend	340309	87	CD43	CD43-10G7	Biolegend	343202
44	CD309 (VEGFR2)	7D4-6	Biolegend	359921	88	CD9	HI9a	Biolegend	312102
45	CD32 (FcγRII)	FUN-2	Biolegend	303225	89	IgA	IS11-8E10	Myltenyi	130-093-073

Extended Data Table 5. *Panel of oligo-conjugated antibodies used in multi-omics scRNA-seq assays to measure surface protein markers on cells from airways and blood.* A total of 89 oligo-conjugated antibody-derived tags (ADT) were used to evaluate surface protein expression of target antigens via scRNA-seq. Reagents no. 82-89 were not commercially available in the TotalSeq-C™ format (BioLegend), so custom oligo-conjugated reagents were generated in-house using purified monoclonal antibodies and commercially available 5' Feature Barcode Antibody Conjugation - Lightning-Link® kits (abcam). Titrations of all reagents were determined in-house for each lot independently prior to use. Cells were surface stained prior to encapsulation and surface marker expression of major lineage markers were used to help cluster/validate gene expression data as described in the methods.

		PID	Total reads	Mean reads/cell
ETA	GEX	PHA0007	593,835,060	350,345
		PHA0008	555,825,992	2,241,234
		PHA0010	587,153,474	872,442
		PHA0018	541,561,148	459,729
	ADT	PHA0007	85,070,531	2,101
		PHA0008	78,647,311	7,514
		PHA0010	46,042,783	2,989
		PHA0018	73,265,461	5,399
Blood	GEX	PHA0007	518,672,044	273,129
		PHA0008	540,400,013	641,044
		PHA0010	592,394,112	543,980
		PHA0018	556,699,985	1,484,533
		PHA1005	541,306,254	158,787
		PHA1006	501,896,927	200,198
		PHA1007	512,999,099	172,902
		PHA1008	506,988,059	160,898
		PHA1009	527,201,752	130,463
		PHA1010	480,796,663	332,042
	ADT	PHA0007	74,597,277	2,115
		PHA0008	80,175,737	1,536
		PHA0010	62,927,249	1,067
		PHA0018	79,818,668	1,395
		PHA1005	77,435,258	1,145
		PHA1006	88,850,369	1,334
		PHA1007	88,329,209	1,169
		PHA1008	93,579,790	1,351
		PHA1009	91,162,676	1,281
		PHA1010	89,597,131	1,251

Extended Data Table 6. *Sequencing depth of independent samples.* All gene expression (GEX) libraries were pooled and run in a single S4 flow cell on a NovaSeq™ 6000. All antibody-derived tag (ADT) libraries were pooled and sequenced in a single lane of a S4 flow cell on a NovaSeq™ 6000. Total reads and mean number of reads per cell for GEX and ADT are reported independently for each sample, with an average read depth of 540,000,000 reads per sample (GEX) and 79,000,000 reads per sample (ADT).

a	PID	Blood (%)	ETA (%)
	PHA0007	30	30
	PHA0008	20	25
	PHA0010	10	25
	PHA0018	30	25
	PHA1005	30	
	PHA1006	30	
	PHA1007	30	
	PHA1008	20	
	PHA1009	20	
	PHA1010	25	

b		Total ADT UMI	
	PID	**Blood**	**ETA**
	PHA0007	125–500	125–1250
	PHA0008	125–1000	125–1250
	PHA0010	125–1500	125–750
	PHA0018	125–1500	125–1000
	PHA1005	125–1250	
	PHA1006	125–1250	
	PHA1007	125–1500	
	PHA1008	125–1500	
	PHA1009	125–1500	
	PHA1010	125–1500	

Extended Data Table 7. *Mitochondrial gene thresholds and total ADT UMIs of individual samples.* (a)Threshold for mitochondrial gene distribution (percentage) for each sample used to exclude potential dead cells in scRNA-seq data. Threshold was evaluated and set for each sample independently. (b) The distribution of total ADT UMIs was determined and recorded per each sample independently.

1201
1202 **Extended Data Figure 1.** *Representative gating strategy for Hi-D FACS data.*
1203 Representative gating of blood (a) and ETA (b) samples. Extracellular stain FMOs for markers
1204 used to interrogate neutrophil phenotype in blood (c) and ETA (d).

1205
1206 **Extended Data Figure 2.** *Additional cytokine assessment in blood and lungs*.
1207 Concentration (pg/mL) of remaining 6 analytes interrogated by Mesoscale analyses in plasma
1208 (gray circles) and respiratory supernatant (Resp. SNT; green squares) from healthy control
1209 (HC), mild-acute (MA), and severe COVID-19 patients. Dotted line = assay limit of detection
1210 (LOD).
1211

Extended Data Figure 3. *Multi-omic scRNA-seq analysis of leukocytes in the blood and lung.* Gating strategy (a and c) employed to classify major lineages of immune cells by surface antibody-derived tag (ADT) and dot plots of the intersection of the top differentially expressed genes sorted by average log-fold change across cell populations (b and d) from the blood (a-b) and ETA (c-d) of severe COVID-19 patients using the SuPPER-seq pipeline previously described[30].

1221

1222   **Extended Data Figure 4.** *Gene signature for immature neutrophils is lacking in healthy*
1223   *donors and lungs of COVID-19 patients.* UMAP visualizations of genes that identify immature
1224   neutrophils in the blood healthy individuals (a) and severe COVID-19 patients (b), along with
1225   cells from the lungs of severe COVID-19 patients (c).
1226

1227
1228
1229 **Extended Data Figure 5.** *Cellular signatures during SARS-CoV-2 pathogenesis.* (a)
1230 Pulmonary neutrophils have a reduction in CD32 (FcγRII) expression as compared to circulating

neutrophils. (b) Neutrophils undergo transcriptional reprogramming and increase de novo transcription (total RNA) upon migrating to the lung. (c) Neutrophils in the blood and lung increase expression of *IFITM2* during SARS-CoV-2 pathogenesis. (d) Metascape[85] pathway and process enrichment analyses reveal non-immune cells have an increased gene signature for granulocyte activation in the airways of severe COVID-19 patients. (e) Extracellular neutrophil elastase is also detected on the surface of pulmonary T cells and myeloid-derived cells (MdCs) in severe COVID-19 patients.