Non-Coding Genetic Analysis Implicates Interleukin 18 Receptor Accessory Protein 3'UTR in Amyotrophic Lateral Sclerosis

Authors:

Chen Eitan[1*], Aviad Siany[1*], Elad Barkan[2], Tsviya Olender[1], Kristel R. van Eijk[3], Matthieu Moisse[4,5], Sali M. K. Farhan[6,7], Yehuda M. Danino[1], Eran Yanowski[1], Hagai Marmor-Kollet[1], Natalia Rivkin[1], Nancy Yacovzada[1,2], Shu-Ting Hung[8-10], Johnathan Cooper-Knock[11], Chien-Hsiung Yu[12,13], Cynthia Louis[12,13], Seth L. Masters[12,13], Kevin P. Kenna[3], Rick A. A. van der Spek[3], William Sproviero[14], Ahmad Al Khleifat[14], Alfredo Iacoangeli[14], Aleksey Shatunov[14], Ashley R. Jones[14], Yael Elbaz-Alon[1], Yahel Cohen[1], Elik Chapnik[1], Daphna Rothschild[2,15,16], Omer Weissbrod[2], Gilad Beck[17], Elena Ainbinder[17], Shifra Ben-Dor[17], Sebastian Werneburg[18], Dorothy P. Schafer[18], Robert H. Brown Jr[19], Pamela J. Shaw[11], Philip Van Damme[4,5,20], Leonard H. van den Berg[3], Hemali P. Phatnani[21], Eran Segal[2], Justin K. Ichida[8-10], Ammar Al-Chalabi[14,22], Jan H. Veldink[3], Project MinE ALS Sequencing Consortium[23], NYGC ALS Consortium[23] and Eran Hornstein[1#]

Affiliations:

[1]Department of Molecular Genetics, Weizmann Institute of Science, Rehovot 7610001, Israel.

[2]Department of Computer Science And Applied Math, Weizmann Institute of Science, Rehovot 7610001, Israel.

[3]Department of Neurology, University Medical Center Utrecht Brain Center, Utrecht University, Utrecht, 3584 CG, The Netherlands.

[4]KU Leuven - University of Leuven, Department of Neurosciences, Experimental Neurology, B-3000 Leuven, Belgium.

[5]VIB, Center for Brain & Disease Research, Laboratory of Neurobiology, Leuven, Belgium.

[6]Analytic and Translational Genetics Unit, Center for Genomic Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA.

[7]Stanley Center for Psychiatric Research, Broad Institute of MIT and Harvard, Cambridge, MA 02142, USA.

[8]Department of Stem Cell Biology and Regenerative Medicine, Keck School of Medicine, University of Southern California, Los Angeles, CA 90033, USA.

[9]Eli and Edythe Broad CIRM Center for Regenerative Medicine and Stem Cell Research at USC, Los Angeles, CA 90033, USA.

[10]Zilkha Neurogenetic Institute, Keck School of Medicine of the University of Southern California, Los Angeles, CA 90033, USA.

[11]Sheffield Institute for Translational Neuroscience (SITraN), University of Sheffield, Sheffield S10 2HQ, UK.

[12]Inflammation Division, The Walter and Eliza Hall Institute of Medical Research, Parkville 3052, Australia.

[13]Department of Medical Biology, University of Melbourne, Parkville 3010, Australia.

[14]King's College London, Maurice Wohl Clinical Neuroscience Institute, Institute of Psychiatry, Psychology & Neuroscience, De Crespigny Park, London, SE5 8AF, United Kingdom.

[15]Department of Developmental Biology, Stanford University, Stanford, CA 94305, USA [16]Department of Genetics, Stanford University, Stanford, CA 94305, USA

[17]Department of Life Sciences Core Facilities, Weizmann Institute of Science, Rehovot 7610001, Israel.

[18]Department of Neurobiology, Brudnick Neuropsychiatric Research Institute, University of Massachusetts Chan Medical School, Worcester, MA 01605, USA.

[19]Department of Neurology, University of Massachusetts Medical School, Worcester, MA 01655, USA.

[20]University Hospitals Leuven, Department of Neurology, Leuven, Belgium.

[21]Center for Genomics of Neurodegenerative Disease, New York Genome Center.

[22]King's College Hospital, Denmark Hill, London, SE5 9RS, United Kingdom.

[23]A list of Consortiums PIs and affiliations appears in the Supplementary Information.

*These authors contributed equally to this work

#Corresponding author. Tel: +972 89346215; Fax: +972 89342108; E-mail: Eran.hornstein@weizmann.ac.il

Abstract:

The non-coding genome is substantially larger than the protein-coding genome but is largely unexplored by genetic association studies. Here, we performed region-based burden analysis of >25,000 variants in untranslated regions of 6,139 amyotrophic lateral sclerosis (ALS) whole-genomes and 70,403 non-ALS controls. We identified Interleukin-18 Receptor Accessory Protein (IL18RAP) 3'UTR variants significantly enriched in non-ALS genomes, replicated in an independent cohort, and associated with a five-fold reduced risk of developing ALS. Variant IL18RAP 3'UTR reduces mRNA stability and the binding of RNA-binding proteins. Variant IL18RAP 3'UTR further dampens neurotoxicity of human iPSC-derived C9orf72-ALS microglia that depends on NF-κB signaling. Therefore, the variant IL18RAP 3'UTR provides survival advantage for motor neurons co-cultured with C9-ALS microglia. The study reveals direct genetic evidence and therapeutic targets for neuro-inflammation, and emphasizes the importance of non-coding genetic association studies.

One Sentence Summary: Non-coding genetic variants in IL-18 receptor 3'UTR decrease ALS risk by modifying IL-18-NF-κB signaling in microglia.

[Main Text:]

Introduction

Genomic sequencing technologies facilitate the identification of variants in open reading frames (ORFs). Although

allelic variants in non-coding regions are expected to be numerous [1, 2] they are largely overlooked.

Amyotrophic lateral sclerosis (ALS) is a fatal neurodegenerative syndrome, primarily affecting the human motor

neuron system with a strong genetic predisposing component [3, 4]. Thus far, mutations in approximately 25 protein-

coding genes have been associated with ALS [3, 5-7]. Hexanucleotide repeat expansion in an intronic sequence of the

C9orf72 gene is the most common genetic cause of ALS [8-10] and enrichment of variants was recently discovered in

the CAV1 enhancer [11]. However, non-coding nucleotide variants in ALS have yet to be systematically explored.

Burden analysis is a genetics approach that is based on the rationale that different rare variants in the same gene

may have a cumulative contribution [12]. Therefore, burden analysis allows the identification of genes containing an

excess of rare and presumably functional variation in cases relative to controls. Although de novo mutations in

non-coding regions were recently shown in family-based autism studies [13], variants in non-coding regions are not

routinely included in rare-variant burden association studies. The application of burden analysis to non-coding

regulatory variation is constrained by the availability of whole-genome sequencing (WGS) data, and the ability to

recognize functional variants in non-coding regulatory regions, which is currently far less effective than for

protein-coding genes.

MicroRNAs (miRNAs) are endogenous posttranscriptional repressors that silence mRNA expression through

sequence complementarity. miRNA primarily act on 3' untranslated regions (3'UTRs) [14], which are non-coding

parts of messenger RNAs (mRNAs) and often regulate degradation and translation [15]. miRNA dysregulation has

been implicated in ALS pathogenesis, and ALS-associated RNA-binding proteins, TARDBP/TDP-43 and FUS,

regulate miRNA biogenesis [16-27].

Microglia are the resident immune cells of the central nervous system and are the primary mediators of

neuroinflammation in neurodegeneration [28, 29]. In ALS, microglia induce motor neuron death via the classical

nuclear factor-kappa B (NF-κB) pathway [29-35]. One suggested mechanism for microglia-induced cytotoxicity is

based on detection of extracellular TDP-43 aggregates and triggering of IL-1beta and Interleukin 18 (IL-18; also

known as: IGIF/IL1F4/IL-1g) signaling [31]. Accordingly, IL-18 levels are elevated in ALS patient tissues and biofluids

[36-39], supporting a role for IL-18 signaling in the disease's neuroinflammatory milieu [40]. IL-18 also induces NF-κB

signaling by binding and dimerising the two IL-18 receptor subunits, IL18RAP (also known as: AcPL/CD218b/IL-

18R-Beta) and IL18R1 [31, 40-48]. In turn, NF-κB contributes to microglial inflammation [49, 50], microglial-mediated motor

neuron death [51] and to disease progression [52, 53].

Here, we identified rare variants in miRNAs and 3'UTR of mRNAs, and performed collapsed genetic analysis [54] to

test if these regulatory RNAs are associated with ALS. We discovered an enrichment of rare variants in the IL18RAP

3'UTR and provide experimental evidence for their relevance to human ALS. Therefore, non-coding variant

analysis reveals a genetic and mechanistic link for the IL-18 pathway in ALS and encourages systematic exploration

of non-coding regions to uncover genetic mechanisms of disease.

Results

To test whether genetic variations in non-coding regulatory regions are associated with ALS, we analyzed regions

of interest in WGS data from the Project MinE ALS sequencing consortium [55] (Supplementary Fig. 1A,B and

Supplementary Tables 1,2). The discovery cohort consisted of 3,955 ALS patients and 1,819 age- and sex-matched

controls, for a total of 5,774 whole-genomes from the Netherlands, Belgium, Ireland, Spain, United Kingdom,

United States, and Turkey (Project MinE Datafreeze 1). We performed a region-based burden test, in which rare

genetic variants with minor allele frequencies (MAF) ≤0.01 are binned together, to weight their contribution to

disease, in 295 non-coding 3'UTRs of candidate genes, linked to sporadic ALS via GWAS [56] or genes encoding RNA-

binding proteins (Supplementary Table 3). In addition we tested all autosomal human-pre-miRNA genes (1,750

pre-miRNAs; miRBase v20 [57]).

As a positive control, we also performed an association analysis of rare variants in the open reading frames of

these 295 genes. For the proteins, we called variants that are predicted to cause frameshifting, alternative splicing,

an abnormal stop codon, or a deleterious non-synonymous amino acid substitution that were detected in ≥ 3 of 7

independent dbNSFP prediction algorithms [58] (Fig. 1A and Supplementary Table 3). In total, 30,721 rare qualifying

variants were identified (Supplementary Table 4). Optimized Sequence Kernel Association Test (SKAT-O) [59]

identified a significant excess of deleterious minor alleles in the ALS genes *NEK1* (127 cases; 19 controls [3.21%;

1.04%]: P = 8x10^{-7}; $P_{corrected}$= 2.3x10^{-4}), comparable with a reported prevalence of 3% [60], and in *SOD1* (36 cases

[0.91%]; 0 controls: P = 2.6x10^{-4}; $P_{corrected}$ = 3.73x10^{-2}) [61], which is below the reported 2% prevalence [5, 62] (Fig. 1B,

Supplementary Fig. 2A and Supplementary Table 5). Other known ALS genes did not reach statistical significance

(Supplementary Table 3), consistent with reported statistical power limitations of Project MinE WGS data in

assessing the burden of rare variants [63]. Our analysis did not consider the *C9orf72* hexanucleotide (GGGGCC)

repeat expansion region.

The burden of rare variants did not identify a disease association for any of the autosomal pre-miRNAs in the

human genome, nor for any of the predicted genetic networks based on variants aggregated over specific mature

miRNAs and their cognate downstream 3'UTR targets. This may be because the small size of miRNA genes makes

genetic aggregation studies particularly challenging (Supplementary Fig. 2B,C).

When we tested the burden of variants in 3'UTRs, the strongest association found was for the 3'UTR of IL18RAP

(Fig. 1B, Supplementary Fig. 2D and Supplementary Table 5). This association was higher than expected at random

(P = 1.93x10^{-5}, $P_{corrected}$ =5.41x10^{-3}) and from the association gained for all protein-coding ALS genes in this cohort,

with the exception of *NEK1*. Notably, the signal was more prevalent in controls [12/1819, 0.66%] relative to ALS

patients [6/3955, 0.15%], indicating that these variants might act as protective variants against ALS.

IL18RAP 3'UTR also ranked as the top hit by three other algorithms – the Sequence Kernel Association Test (SKAT,

P = 1.77×10^{-5}; permutated P-value < 10^{-4}), the Combined Multivariate and Collapsing (CMC, P = 8.78×10^{-4}) or

Variable Threshold (VT) with permutation analysis (permutated P-value = 1.75×10^{-3}, suggesting that the

association does not depend on a particular statistical genetics method (Supplementary Fig. 3A-C). Furthermore,

when we tested the burden of variants in miRNA recognition elements (MREs) in 3'UTRs (variants that are

potentially either abrogating conserved miRNA binding sites or creating new miRNA binding sites in 3'UTRs), the

strongest association was also gained for the 3'UTR of IL18RAP (SKAT-O P-value = 3.42×10^{-5}, Supplementary Fig.

3D, see Methods). A diagram of variants in IL18RAP 3'UTR is presented in Supplementary Fig. 3E and a description

of IL18RAP 3'UTR variants in Supplementary Table 6. The top 10 principal components (PCs) of common variant-

based ancestry information and sex were included as covariates in the SKAT-O, SKAT, CMC, and VT analyses.

In addition, genome-wide analysis of all known human 3'UTRs (16,809 3'UTRs from RefSeq [64]) identified IL18RAP

3'UTR as the most significant 3'UTR associated with ALS in the Project MinE cohort (Fig. 1C).

Finally, we tested if different functional genetic classes were enriched overall for ALS risk/protection variants by

testing the burden of rare variants in all genes pooled together. SKAT-O signal for open reading frames of 295

proteins, the 3'UTR of the same 295 genes, all autosomal pre-miRNA genes [miRBase v20; [57]] or networks

composed of all miRNA genes and their cognate set of downstream targets (TargetScan) were all not significant

(P-values of 0.024, 0.59, 0.08, 0.58, respectively). Therefore, results from these burden tests do not implicate any

of the functional class of genomic elements in ALS risk.

Because the number of ALS genomes was ~2.17-fold larger than the number of controls, the data depict a 4.35-

fold enrichment in the abundance of variants in controls over cases. IL18RAP 3'UTR potentially-protective variants

reduced the disease odds ratio by five-fold (OR = 0.23; Fig. 2A), and was consistent across independent population

strata (Fig. 2B), whereas *NEK1* and *SOD1* increased the disease odds ratio (OR = 3.14, 33.89, respectively; Fig. 2A).

To determine if the rare IL18RAP 3'UTR variants are depleted in another ALS cohort, we performed independent

replication studies. Similar results for rare IL18RAP 3'UTR variants were reproduced in the New York Genome

Center (NYGC) ALS Consortium cohort (2,184 ALS genomes), which was studied against: (i) 263 non-neurological

controls from the NYGC; (ii) 62,784 non-ALS genomes from NHLBI's Trans-Omics for Precision Medicine (TOPMed);

and (iii) 5,537 non-ALS genomes from gnomAD. This replication effort yielded a joint analysis P-value = 9.58×10^{-4}

(χ2 with Yate's correction; OR=0.32; 95% CI: 0.16 − 0.64; Fig. 2C and Supplementary Table 7). Combining this

cohort with our discovery cohort from Project MinE, yielded a superior joint P-value < 1.00×10^{-5} (χ^2 with Yate's

correction; OR=0.20; 95% CI: 0.12 – 0.34; Fig. 2C). A meta-analysis of Project MinE datafreeze 1 and 2 [7], which

consisted of 5,185 ALS patients and 2,262 age- and sex-matched controls, reproduced the initial signal (P-value =

7.6×10^{-4}).

Together, IL18RAP 3'UTR sequence variants are associated with a lower risk of suffering from ALS, which is

approximately one-fifth of the general population, although it did not reach conventional exome-wide multiplicity-

adjusted significance threshold ($\alpha \approx 2.6 \times 10^{-6}$, ref. [12]) in our study.

To investigate the source of the signal in the IL18RAP 3'UTR in a posthoc analysis, we divided the 11 variant

nucleotides into two synthetic sets, of either nine singleton variants (9 variants / 3 controls / 6 patients) or two

variants that were identified solely in controls (2 variants / 9 controls / 0 patients). While the signal of the nine

singleton variants was not statistically significant, analysis of the two control variants, which were identified in

multiple samples, derived an improved significance compared to the original signal (SKAT-O P-value = 4.36×10^{-6}).

Thus, these two rare variants (V1, Chr2:103068691 C>T; V3, Chr2:103068718 G>A) are likely central in generating

the genetic association signal in IL18RAP 3'UTR.

Because of the enrichment of V1 and V3 at the proximal (5') side of the IL18RAP 3'UTR, we tested if restricting

burden analysis to the 5' end of the 3'UTR, might boost the association signal. However, the P-values gained from

the 3'UTRs proximal quadrant were comparable to that of the full 3'UTRs in the cohort of 295 3'UTRs (Wilcoxon

matched-pairs P-value > 0.05, Cohen's d effect size = 0.1, Supplementary Fig. 4A,B), suggesting that the apparent

spatial distribution of variants in the case of IL18RAP 3'UTR is a particular case rather than part of a global pattern.

To determine the functional impact of the IL18RAP 3'UTR variants we analyzed IL18RAP expression in

lymphoblastoid cell lines (LCLs) from the UK MNDA DNA Bank [65] that were derived from twelve different

individuals: 4 healthy individuals (without ALS), carrying the canonical IL18RAP 3'UTR sequence (Control;

Canonical IL18RAP 3'UTR); 4 sporadic ALS patients, carrying the canonical IL18RAP 3'UTR sequence (sALS;

Canonical IL18RAP 3'UTR); two healthy individuals, carrying a variant form of IL18RAP 3'UTR (Control;

VariantIL18RAP 3'UTR) and two sporadic ALS patients carrying a variant form of IL18RAP 3'UTR (sALS;

VariantIL18RAP 3'UTR; see Supplementary Table 8 for list of variants).

ALS-derived LCLs carrying the canonical IL18RAP 3'UTR sequence expressed higher levels of IL18RAP (Fig. 3A,B).

In addition, LCLs from both healthy and ALS individuals harboring the IL18RAP 3'UTR variant significantly down-

regulated IL18RAP mRNA and protein expression (Fig. 3A,B and Data File S1). Phosphorylation of the nuclear factor

kappa-light-chain-enhancer of activated B cells (p-NF-κB), an established intracellular effector downstream of IL-

18 signaling, was similarly higher in the ALS LCLs with canonical IL18RAP 3'UTR and also significantly reduced in

control and ALS LCLs harboring IL18RAP variants (Fig. 3C,D and Data File S1). Consistent results were obtained

with C9orf72 hexanucleotide expansion ALS LCLs (Supplementary Fig. 5 and Data File S2). Accordingly, variants of

IL18RAP 3'UTR reduced NF-κB activity, relative to the canonical 3'UTR in an NF-κB reporter assay in U2OS cells

(Supplementary Fig. 6). Therefore, variant forms of IL18RAP 3'UTR correlate with reduced expression of the

endogenous IL18RAP and reduced NF-κB signaling.

To further establish the functional relevance of the IL18RAP 3'UTR variants, we edited the genome of human-

induced pluripotent cells (iPSCs) donated by ALS patients with C9orf72 repeat expansion ([66] NINDS/Coriell Code:

ND10689, ND12099, see Supplementary Table 8) to include two point mutations that recapitulate the most

prevalent variants (Chr2:103068691 C>T (V1) and Chr2:103068718 G>A (V3)) in the IL18RAP 3'UTR sequence (Fig.

4A). The resulting isogenic pair lines all carry C9orf72 repeat expansion and vary by only the presence of the

canonical or a variant IL18RAP 3'UTR.

We explored the receptive cell type involved in IL-18 receptor signaling by profiling dissociated mouse brain cells,

namely, neurons, microglia, and astrocytes. Fluorescence cytometric gating on CD11b+ and CD45+ and IL18RAP

(CD218b) revealed that IL18RAP is mainly expressed on microglia cells (Supplementary Fig. 7A-C). Although IL-18

and IL18RAP expression increases in ALS motor neurons (Supplementary Fig. 8A-C), our observations are

consistent with the accepted notion that the role of IL-18 and other cytokines in disease heavily rests on a chronic

inflammatory state established particularly by microglia [67].

Therefore, we next differentiated the isogenic IL18RAP 3'UTR lines into human microglia following the protocol

of Haenseler et al. [68] (Fig. 4A). iPSC-derived microglia differentiation was validated by immunofluorescence

staining of the microglial-specific marker, TMEM119 (Supplementary Fig. 9). In differentiated human microglia,

we detected a ~5-6 fold downregulation in the levels of the variant IL18RAP protein, as well as in the levels of the

IL18RAP mRNA, relative to the canonical sequence of the isogenic line (Fig. 4B,C and Data File S3). Therefore, the

variants at the 3'UTR regulate IL18RAP mRNA and protein expression and provide a conceivable explanation for

the variant function in human C9-ALS microglia. Next, we investigated the molecular mechanism that controls the

IL18RAP mRNA levels by performing an mRNA stability assay in human microglia. We measured an mRNA

degradation rate that is twice as fast with the rare 3'UTR variants, relative to the canonical sequence, after

inhibition of mRNA transcription by actinomycin D (Fig. 4D). Thus, the mechanism for reduced IL18RAP mRNA

levels is associated with destabilization of IL18RAP mRNA via variants in the 3'UTR.

We sought the potential trans-acting factors that might differentially bind to the canonical and variant 3'UTRs. To

this end, we performed RNA-pulldown assays and mass spectrometry on *in vitro* transcribed canonical and variant

forms of the IL18RAP 3'UTRs, V1 and V3 (Fig. 5A diagram of exp. design). Mass spectrometry after pull-down

identified 552 proteins with good confidence (passed all QC filters, found in 50% of the repeats in at least one

experimental group, and were represented by at least 2 unique peptides, Supplementary Table 9), that were

enriched in comparison to the negative control. Principal component analysis demonstrated a clear separation of

proteomes bound by the canonical and variant IL18RAP 3'UTRs (Fig. 5B and Supplementary Table 9). Gene set

enrichment analysis (GSEA) revealed a reduction in the association of double-stranded RNA (dsRNA) binding

proteins, to V1 IL18RAP 3'UTR, relative to the canonical 3'UTR (ELAVL1/Hur; PRKRA, EIF2AK2/PKR; ADAR; ADARB1;

ILF2; ILF3; DHX9; DHX58; DDX58, Fig. 5C,D,E and Supplementary Table 10). These dsRNA binding proteins were

reported in other contexts to play roles in controlling the stability of mRNA [69-75], consistent with the observed

changes to IL18RAP mRNA stability. A similar analysis of the V3 variant was unproductive (Supplementary Fig.

10A).

In accordance, RNA Fold analysis predicted that the canonical 3'UTR sequence consists of a more stable dsRNA

structure than the V1 variant sequence (minimum free energy (MFE) of canonical and variant IL18RAP 3'UTR, -

39.9 kcal/mol and -27.8 kcal/mol, respectively) (Fig. 5F and Supplementary Fig. 10B). In light of these results, we

propose that variant-dependent changes to the secondary structure of IL18RAP 3'UTR attenuate the binding of

one or more of the dsRNA proteins and may be involved in controlling the stability of IL18RAP mRNA.

To study the potential protective impact of IL18RAP 3'UTR mutations, we performed survival analyses in a

coculture system of human iPSC-derived isogenic IL18RAP 3'UTR microglia (on a *C9orf72 repeat expansion*

background) with human iPSC-derived lower motor neurons (i³LMNs; healthy, non-ALS, [76]). Time-lapse microscopy

was used to quantify motor neuron survival after microglia activation with a cocktail of LPS and the cytokine IL-18

(experimental design, Fig. 6A). Motor neuron survival was significantly improved in the presence of microglia

harboring the IL18RAP 3'UTR variants relative to microglia harboring the canonical IL18RAP 3'UTR (two

independent isogenic pairs, based on independent patient C9orf72 lines, n=3 independent differentiation

procedures from different passages per line, with 3-8 co-culture wells per passage; Fig. 6B-D, Supplementary

movie and Data File S4). Based on these studies, we conclude that rare variants of IL18RAP 3'UTR increase *C9orf72*

microglia-dependent motor neuron survival and hence convey a protective property.

To determine whether the mutant IL18RAP 3'UTR is also protective in human patients with ALS, we tested the

association between age of diagnosis and age of death in ALS patients harboring canonical or variants of the

IL18RAP 3'UTR. Of 4216 patients for whom data on the age of diagnosis was available (Project MinE and NYGC

cohorts), 8 harbored IL18RAP 3'UTR variants. Of 4263 patients for whom the age of death was available, 9 harbored IL18RAP 3'UTR variants. IL18RAP 3'UTR variants are expected to be depleted in ALS genomes, nonetheless, in those extremely rare patients harboring IL18RAP 3'UTR variants, these were associated with an older age of death and an older age of diagnosis. On average, the age of death was higher by 6.1 years after the average for patients with canonical Il18RAP 3'UTR (Permutation P-value = 0.02, Cohen's d effect size = 0.65; Fig. 6E and Supplementary Table 11), and the age of diagnosis was higher by 6.2 years after the average for patients with canonical IL18RAP 3'UTR (Permutation P-value = 0.05, Cohen's d effect size = 0.62; Fig.6F and Supplementary Table 11). Thus, variants in IL18RAP 3'UTR are protective against ALS in a tissue culture model and correlate with survival advantage for patients suffering from the disease.

To study the role of NF-κB signaling in our system, we analyzed NF-κB phosphorylation and the impact on the transcriptome after microglia activation (Fig. 7A). Western blot analysis revealed reduced levels of phospho-NF-κB in variant IL18RAP 3'UTR relative to isogenic control (Fig. 7B and Data File S5). Reduced phosphorylation is associated with decreased nuclear localization and transcriptional activity of NF-κB [77-80]. In parallel, we conducted a next-generation sequencing study (Supplementary Table 12, Gene Expression Omnibus accession number: GSE186757) of the differentially expressed transcriptomes in microglia harboring variant vs. canonical IL18RAP 3'UTR. Over-representation analysis (ORA) of differentially expressed genes (DEGs) revealed downregulation of the NF-κB signaling pathway in microglia harboring the variant IL18RAP 3'UTR (KEGG Pathway enrichment results: Ratio = 3.77, FDR P-value = 7.34×10^{-6}; Gene Ontology Biological Process enrichment results: Ratio = 3.48, FDR P-value = 3.70×10^{-12}, Fig. 7C,D and Supplementary Table 13). In addition, an unsupervised study of NF-κB pathway mRNAs (GO:0007249) demonstrated broad downregulation of pathway-associated mRNAs in microglia with the variant IL18RAP 3'UTR, relative to the isogenic control (Fig. 7E). Therefore, microglia's NF-κB transcriptomic signature depends on signaling via the IL-18 receptor and is attenuated by protective IL18RAP 3'UTR variants.

To test a plausible neurotoxic role for NF-κB downstream of the IL-18 receptor in this system, we next performed a co-culture survival assay with or without IKK16, a selective IκB kinase (IKK) inhibitor that inhibits NF-κB signaling [81]. In human microglia with the canonical IL18RAP 3'UTR, IKK16 significantly ameliorated motor neuron toxicity, relative to control (carrier alone, Fig. 7F). However, in human microglia with the protective variant IL18RAP 3'UTR, inhibition of NF-κB had no effect (two independent isogenic pairs, based on independent patient C9orf72 lines with 3-8 co-culture wells per line, Fig. 7F). This suggests that NF-κB neurotoxic function resides epistatically downstream of IL18RAP in human microglia. Together, rare variants in IL18RAP 3'UTR diminish NF-κB signaling, thus increasing *C9orf72* microglia-dependent motor neuron survival.

Discussion

Data from the Project MinE and NYGC ALS consortia provide unprecedented opportunities for investigating the role of the non-coding genome in ALS. By performing rare variant aggregation analysis in regulatory non-coding regions, we demonstrate that variants in the 3'UTR of IL18RAP are enriched in non-ALS genomes, indicating that these are relatively depleted in ALS. IL18RAP 3'UTR variants reduced the chance of developing ALS five-fold, and delayed onset and therefore age of death in people with ALS.

These protective variants recall other protective variants that have been reported in the past in protein-coding regions in Alzheimer's disease [82-85] and implicated in ALS as well [86, 87]. In addition, deleterious variants were suggested in VEGF promoter/5'UTR and within CAV1/CAV2 enhancers [11, 88]. However, the 3'UTR of IL18RAP is a protective non-coding allele associated with a neurodegenerative disease.

Elevated levels of the cytokine IL-18 were reported in tissues and biofluids of ALS patients [36-38]. Accordingly, we reveal the upregulation of endogenous IL18RAP in sporadic and C9orf72 lymphoblastoid cells. In addition, we demonstrate the downregulation of IL18RAP expression in lymphoblastoid cells harboring variant versions of the IL18RAP 3'UTR.

We elucidated the regulatory changes affected by the IL18RAP 3'UTR variants by showing destabilization of the IL18RAP mRNA and downregulation of IL18RAP mRNA levels. Sequence analysis suggests that at least one variant (V1) potentially reduced the propensity of the 3'UTR to form a double-stranded secondary structure. Accordingly, unbiased proteomics demonstrated that the 3'UTR harboring the variant fails to bind dsRNA binding proteins that are known to stabilize mRNAs. Together, this supports a mechanism for reduced IL18RAP signaling involving changes to mRNA stability and differential binding of stabilizing RNA-binding proteins.

Neuro-inflammation is prevalent in neurodegeneration, including in ALS [89], and is often characterized by the activation of microglia [29-35]. The cytokine, IL-18, is part of this neuro-inflammatory milieu, promoting receptor subunit (IL18RAP, IL18R1) dimerization on the membrane of cells [40], and activating intracellular signaling cascades, including NF-κB.

By CRISPR editing of two independent C9orf72 lines, from one female and one male patient, we created isogonic IL18RAP 3'UTR cell lines (canonical or harboring V1 and V3 variants), at the endogenous gene locus. By differentiating these lines to human microglia, we demonstrated that variants downregulated IL18RAP mRNA and

protein expression. In addition, tracking of human (wild-type) motor neuron survival, in co-culture with microglia,

over 21 days, demonstrated the neuroprotective effect of microglia carrying the variant IL18RAP 3'UTR.

Finally, we demonstrate that variant IL18RAP 3'UTR attenuates NF-κB signaling in lymphoblastoid cells and in

microglia. Unbiased next-generation sequencing of microglia RNA demonstrated broad transcriptomic changes,

typical of reduced NF-κB signaling. In addition, inhibition of NF-κB was able to ameliorate motor neuron death

when co-cultured with microglia harboring the canonical IL18RAP 3'UTR. However, inhibition of NF-κB was not

further protective if microglia with variant IL18RAP 3'UTR were present, suggesting an epistatic relationship,

whereby IL18RAP is upstream of NF-κB in this system. We conclude that IL18RAP acts in microglia and controls

the cytotoxicity conveyed to motor neurons, at least in human C9orf72 types of disease.

The discovery of functional, disease-modifying IL18RAP 3'UTR variants underscores the need to explore the role

of additional non-coding genomic regions in ALS. One limitation of our study is that IL18RAP 3'UTR signal did not

reach the conventional exome-wide multiplicity-adjusted significance threshold ($\alpha \approx 2.6 \times 10^{-6}$, ref. [12]). However,

IL18RAP 3'UTR signal is comparable to that of protein-coding ALS-causing genes, such as *SOD1* and *NEK1.*

Furthermore, the key findings were reproduced in a genome-wide study of all human 3'UTRs and in an

independent replication study. Limitations in the statistical power might have prevented the discovery of other

non-coding variants and may be overcome with larger ALS and control cohorts, which are not currently available.

Additionally, we have focused our tissue culture studies on human C9orf72 microglia. Therefore, the involvement

of IL18RAP 3'UTR in other ALS-associated genetic backgrounds remains to be experimentally explored, as is the

relevance to other neurodegenerative diseases. Finally, the mechanism underlying IL18RAP dose sensitivity is not

fully understood. While we provide evidence that variant IL18RAP 3'UTR endows neuroprotection via dampening

of microglia-dependent neurotoxicity, additional studies should explore the degree to which other cell types, such

as motor neurons and astroglia, are involved.

In summary, we have identified the IL18RAP 3'UTR as a non-coding genetic disease modifier by burden analysis of

WGS data using ALS case-control cohorts. We show that IL-18 signaling modifies ALS susceptibility and

progression, delineating a neuro-protective pathway and identifying potential therapeutic targets for ALS.

Whereas the 3'UTR of IL18RAP is a protective non-coding allele associated with a neurodegenerative disease, the

increasing wealth of WGS data in Project MinE, NYGC and elsewhere, indicates that the exploration of non-coding

regulatory genomic regions should reveal further disease-relevant genetic mechanisms.

Methods

Human genetic cohorts

All participants contributed DNA after signing informed consent at the submitting sites. Human materials were

studied under approval of the Weizmann Institute of Science Institutional Review Board (Weizmann IRB: 1039-1).

Discovery cohort: Project MinE ALS sequencing consortium Datafreeze 1 includes 3,955 ALS patients and 1,819

age- and sex-matched controls, free of any neurodegenerative disease, for a total of 5,774 quality control (QC)

passing whole-genomes, from the Netherlands, Belgium, Ireland, Spain, United Kingdom, United States, and

Turkey. Rare variant association in cases versus controls was evaluated for regions of interest, when we could

identify ≥2 variants per region, by SKAT-O, SKAT, CMC, and VT in RVTESTS environment [90], with sex and the top 10

principal components (PCs) as covariates. To construct the PCs of the population structure, an independent set of

~450,000 SNPs was sampled from WGS, (MAF ≥0.5%) followed by LD-pruning. Rare genetic variants were included

based on minor allele frequencies (MAF) ≤0.01 within the controls in the current data set.

Replication cohorts: Utilized for testing rare variant alleles (MAF < 0.01) in human IL18RAP 3'UTR (GRCh37/hg19

chr2:103068641-103069025 or GRCh38 chr2:102452181-102452565) from Project MinE datafreeze 2: ~1300

European heritage ALS genomes without middle eastern (Turkish and Israelis) genomes. The New York Genome

Center (NYGC) ALS Consortium (2,184 ALS Spectrum MND and 263 non-neurological control genomes from

European/Americas ancestries), NHLBI's Trans-Omics for Precision Medicine (TOPMed; 62,784 non-ALS genomes)

and gnomAD (5,537 non-ALS genomes; Europeans, non-Finnish, non-TOPMed). Joint analysis in replication cohort

was performed by Chi-square test with Yate's correction. Meta-analysis was not possible because TOPMed and

gnomAD covariate information is not available.

Quality control procedures in Project MinE genomics

Sample selection, data merging, and sample- and variant level quality control procedures for Project MinE ALS

sequencing consortium genomes are described in full previously [63]. Briefly, 6,579 Project MinE ALS sequencing

consortium whole genomes were sequenced on Illumina HiSeq2000 or HiSeqX platforms. Reads were aligned to

human genome build hg19 and sequence variants called with Isaac Genome Alignment Software and variant caller

[91]. Individual genomic variant call format files (GVCFs) were merged with 'agg' tool: a utility for

aggregating Illumina-style GVCFs. Following completion of the raw data merge, multiple QC filtering steps were

performed: (i) setting genotypes with GQ<10 to missing; (ii) removing low-quality sites (QUAL< 30 and QUAL< 20

for SNPs and indels, respectively); (iii) removing sites with missingness > 10%. (iv) Samples excluded if deviated

from mean by more than 6SD for total numbers of SNPs, singletons and indels, Ti/Tv ratio, het/hom-non-ref ratio,

and inbreeding (by cohort). (v) missingness > 5%, (vi) genotyping-sequence concordance (made possible by

genotyping data generated on the Illumina Omni 2.5M SNP array for all samples; 96% concordance), (vii) depth of

coverage, (viii) a gender check (to identify mismatches), (ix) relatedness (drop samples with >100 relatedness

pairs). (x) Related individuals were further excluded until no pair of samples had a kinship coefficient > 0.05. (xi)

missing phenotype information. Following QC, 312 samples with expended/inconsistent *C9orf72* status were

omitted from further analysis. A total of 5,774 samples (3,955 ALS patients and 1,819 healthy controls) passed all

QC and were included in downstream analysis. Per-nucleotide site QC was performed on QC-passing samples only,

for Biallelic sites: variants were excluded from analysis based on depth (total DP < 10,000 or > 226,000),

missingness > 5%, passing rate in the whole dataset < 70%, sites out of Hardy–Weinberg equilibrium (HWE; by

cohort, controls only, $p < 1 \times 10^{-6}$) and sites with extreme differential missingness between cases and control

samples (Overall and by cohort, $p < 1 \times 10^{-6}$). Non-autosomal chromosomes and multiallelic variants were excluded

from analysis.

Selection of regions of interest

Discontinuous regions of interest approximating in total ~5Mb, include coding sequences and 3' untranslated

regions (3'UTRs) of 295 genes (Supplementary Table 3) encoding for proteins that were: (i) previously reported to

be associated with ALS, (ii) RNA-binding proteins including miRNA biogenesis or activity factors [UCSC gene

annotation; [92]]. In addition to (iii) all 1,750 autosomal human pre-miRNA genes [miRBase v20; [57]]. In addition,

genome-wide analysis of all known human 3'UTRs (RefSeq [64]). Variants in regions of interest were extracted from

Project MinE ALS sequencing consortium genomes using vcftools [93] according to BED file containing genomic

coordinates of interest (hg19) ±300 bp that ensures covering splice junctions and sequence (Supplementary Table

14).

Annotation and burden analysis

After quality control and extraction of regions of interest, we performed functional annotation of all variants. Indels were left-aligned and normalized using bcftools and multiallelic sites were removed. For variant annotation we developed a pipeline that calculates the impact of genetic variation in coding regions as well as in 3'UTR and miRNA regions, using ANNOVAR [94]. The frequency of the variants in the general population was assessed by screening the 1000 Genomes Project, the Exome Aggregation Consortium (ExAC), and NHLBI Exome Sequencing Project (ESP). For protein-coding ORFs, association analysis of deleterious rare variants was performed, i.e., frameshift variants, deviation from canonical splice variant, stop gain/loss variants, or a non-synonymous substitution, as predicted by at least three prediction programs (SIFT, Polyphen2 HVAR, LRT, MutationTaster, MutationAssessor, FATHMM, MetaLR) in dbNSFP environment [v2.0; [58]].

Non-coding sequence burden analysis included (i) 3'UTRs, (ii) variants in miRNA recognition elements (MREs) in 3'UTRs (Supplementary Table 3): Variants that impaired conserved-miRNA binding sites in 3'UTRs (predicted loss of function) were called by TargetScan [v7.0; [95]]. Newly created miRNA binding sites in 3'UTRs (predicted gain of function) were called by textual comparison of all possible mutated seeds around a variant to all known miRNA seed sequences in the genome, (iii) all human pre-miRNAs (mirBase v20 [57]) and (iv) miRNAs:target gene networks: mature miRNA sequences (mirBase v20 [57]) and cognate targets within the 3'UTRs (Supplementary Table 3). Variant annotation scripts are available at GitHub: https://github.com/TsviyaOlender/Non-coding-Variants-in-ALS-genes-

Mammalian Cell Cultures

Lymphoblastoid cell lines (LCLs) from the UK MNDA DNA Bank [65] were originally derived from sixteen different individuals: 4 healthy individuals (without ALS), carrying the canonical IL18RAP 3'UTR sequence (Control; Canonical IL18RAP 3'UTR); 4 sporadic ALS patients, carrying the canonical IL18RAP 3'UTR sequence (sALS; Canonical IL18RAP 3'UTR); two healthy individuals, carrying a variant form of IL18RAP 3'UTR (Control; Variant IL18RAP 3'UTR); two sporadic ALS patients carrying a variant form of IL18RAP 3'UTR (sALS; Variant IL18RAP 3'UTR) and 4 C9orf72 ALS patients, carrying the canonical IL18RAP 3'UTR sequence (C9orf72; Canonical IL18RAP 3'UTR) (Cell lines listed in Supplementary Table 8; Weizmann IRB: 537-1). LCLs were cultured in RPMI-1640 (Gibco, 21875091) with 20% inactivated fetal bovine serum (FBS, Biological Industries, 04-001-1A), 1% L-glutamine and 1% penicillin-streptomycin (Biological Industries, 03-0311B) at 37°C, 5% CO_2. Human Bone Osteosarcoma Epithelial Cells (U2OS), were maintained in Dulbecco's Modified Eagle Medium (DMEM, Biological Industries, 01-050-1A) supplemented with 10% FBS, 1% penicillin-streptomycin at 37°C, 5% CO_2. Human iPSCs were cultured on Matrigel (Corning, 354277) coated plated in mTeSR1 medium (Stemcell technologies, 85850) according to the

361 manufacturer's instructions. Briefly, cells were passaged at 70–90% confluent with StemPro accutase (Gibco,

362 A11105-01) and seeded in mTeSR1 medium supplemented with 10 nM Y-27632 dihydrochloride (Tocris, 1254).

363 Cells were refreshed with mTeSR1 medium every 24 hours until passaged.

Isolation and Culture of Rat Cortical Astrocytes

All experiments were performed in accordance with relevant guidelines and regulations of the Institutional Animal

Care and Use Committee at Weizmann Institute of Science (IACUC 09491120-1). Primary cortical a**s**trocytes were

isolated and cultured as previously described [96] with several modifications. Briefly, the cerebral cortex of postnatal

day 1 (P1) Sprague-Dawley rat pups was dissected and placed in DMEM/F12 containing 0.5% trypsin (biological

industries, 03-046-5B). After 30 min incubation at 37 °C water bath, the cortical tissues were mechanically

dissociated with pipette into single cells and were seeded on poly-D-lysine (Sigma Aldrich, 7405) coated T75

culture flask in Astrocytes medium (DMEM/F12 (Gibco, 31330) supplemented with 10% FBS, 50U/mL Pen-strep

and 2Mm Glutamax (Gibco, 35050-038)). The confluent cultures were shaken for 4 hours at 200 rpm to remove

microglial cells. Each T75 flask was trypsinized and splited into three new T75 flasks. After 7-8 days the confluent

flasks were trypsinized and were frozen (in 90% FBS, 10% DMSO) until further use.

I^{3}LMNs neuronal differentiation and *Syn::GFP+* transduction

Differentiation of hiPSCs into lower motor neurons (i^{3}LMNs, iPSCs containing Doxycycline induced human NGN2,

ISL1, and LHX3 (hNIL)) was performed as described previously [76]. Briefly, iPSCs were seeded on day 0 into mTeSR1

medium supplemented with 10 nM Y-27632 dihydrochloride. Few hours after seeding, cells were transduced with

Syn::GFP lentivirus (pHR-hSyn-EGFP, Addgene #114215). 24 hrs. after seeding the cells medium was replaced with

differentiation medium (DMEM/F12 (Gibco, 31330-038) with 1× MEM non-essential amino acids (Gibco, 11140-

035), 2mM GlutaMAX (Gibco, 35050-038), 1× N-2 supplement (Gibco, 17502-048), 2 µg/mL Doxycycline (Sigma

Aldrich, D9891-1G.) and 10 nM Y-27632 dihydrochloride). On day 3, cells were split using accutase, counted and

re-seeded on poly-D-Lysine coated dishes containing Rat astrocytes in neuronal medium (B27 Electrophysiology

medium (Gibco, A14137-01) supplemented with 1× MEM non-essential amino acids, 2mM GlutaMAX, 1× N-2

supplement, and 1 µg/mL mouse laminin (Gibco, 23017-015)). Twice a week half of the media was removed, and

an equal volume of fresh media was added.

Generation of IL18RAP 3'UTR rare variant hiPSCs lines

iPSCs were generated by the Ichida lab from human lymphocytes from ALS patients obtained from the National Institute of Neurological Disorders and Stroke (NINDS) Biorepository at the Coriell Institute for Medical Research. lymphocytes were reprogrammed into iPSCs as previously described [66]. The NINDS Biorepository requires informed consent from patients.

Human iPSC lines were maintained on irradiated MEFs in hESCs medium [DMEM/F12 (Sigma-Aldrich, D6421) supplemented with 20% KO Serum Replacement (Gibco, 10828-028,), 1% GlutaMax (Gibco, 35050038), 1% MEM-NEAA (Biological Industries, 01-040-1A), 0.1mM 2-Mercaptoethanol (31350-010, Gibco), 10ng/ml hFGF (PeproTech, 100-18B)] and passaged twice a week with Collagenase IV (Worthington, LS004188).

CRISPR guides were chosen using several design tools, including: the MIT CRISPR design tool [97] and sgRNA Designer, Rule set 2 [98], in the Benchling implementations (www.benchling.com), SSC [99], and sgRNAscorer [100], in their websites.

Prior to CRISPR procedure iPSCs were passaged once in feeder-free condition [LDEV Free GelTrex matrix (Gibco, A1413202), mTESR1 medium (StemCell Technologies, 85850)], dislodged as single cells using StemPro Accutase (Gibco, A11105-01), washed twice with Opti-MEM (Gibco, 31985-047) and counted. 90ul cells suspension containing 1M cells was mixed with 10 uL DNA mix: 4 ug pSpCas9(BB)-2A-Puro (PX459) plasmid (Addgene #48139), 0.4 ug gRNA encoding plasmid (pKLV-U6gRNA(BbsI)-PGKzeo2ABFP, derived from pKLV-U6gRNA(BbsI)-PGKpuro2ABFP (Addgene)), 1 ug (8 pmol) ssODN repair template (Supplementary Table 15) (IDT, 400 bases Megamer DNA Oligonucleotide) and 2.6 ug carrier plasmid DNA. CRISPR reaction components were introduced to iPSCs by single round electroporation using Nepa21 system (NEPA GENE). 100 uL cells and DNA suspension was transferred to Nepa Electroporation Cuvette 2 mm gap (Nepa Gene, EC-002). Electroporation conditions: 150 V Poring pulse; 5 ms Pulse length; 20 V Transfer pulse; 50 ms Pulse length. Electroporated cells were transferred to two GelTrex coated 100 mm dishes (1K and 10K) in mTeSR medium supplemented with 10 uM ROCK inhibitor (PeproTech, 1293823) and placed into CO_2 incubator for 2 days. 48h past electroporation cells were treated with 0.5 ug/mL Puromycin (Sigma-Aldrich) for 2 consecutive days. Survived cells were maintained until clones development. Single clones were picked and transferred to 96 well plates. Matured clones were genotyped at the first passage. Additionally, the top five predicted off-target sites for the guide RNA were sequenced (Supplementary Table 16). Selected clones containing desired mutations were expanded, cryopreserved, and used for the downstream experiments.

Differentiation and culturing of human iPSC-derived microglia

hiPSCs were differentiated to microglia-like cells as previously described [68]. Briefly, to form embryoid bodies (EBs), iPSCs were seeded into 96 well suspension plates in mTeSR1 media supplemented with 50 ng/mL rhBMP4 (Peprotech, 314-BP), 50 ng/mL VEGF (Peprotech, 100-20), 20 ng/mL SCF (Peprotech, 300-07) and 10 nM Y-27632 dihydrochloride. Everyday half of the medium was removed, and an equal volume of fresh media was added. After four days 12 EBs were transferred into each well of 6 well plate in X-VIVO 15 (Lonza, BE02-060Q) containing 100 ng/mL M-CSF (Peprotech, 300-25), 25 ng/mL IL-3 (Peprotech, 200-03), 2 mM Glutamax, 55 uM 2-mercaptoethanol (Gibco, 31350-10) and 100 U/mL penicillin/streptomycin (Biological Industries, 03-031-1B). iPSC derived progenitor microglia (ipMG) were collected weekly from the supernatant and were co-cultured with iPSC derived neurons in 96 well plates (Greiner, 655090) in neuronal medium containing 10 ng/mL IL-34 (Peprotech,200-34). EB medium was refreshed weekly.

i³LMNs survival assay

Survival assay was conducted by monitoring eGFP signal of day 5 i³LMNs co-cultured with two independent CRISPR-edited isogenic iPSC-derived microglia (harboring canonical or variant IL18RAP 3'UTR), with C9orf72 genetic background. Cells were monitored for over 20 days using Incucyte® Live-Cell Analysis System (Sartorius). Daily longitudinal microscopic tracking was performed following Lipopolysaccharide (LPS, 100 ng/mL) and IL-18 treatment (100 ng/mL). i³LMNs survival assay was performed using three individual replicates for each line, with 3-8 co-culture wells per condition. Twice a week half of the media was removed, and an equal volume of fresh media containing LPS and IL-18 was added.

Cloning

Full IL18RAP coding sequence (CDS) and 3'UTR sequence (2223bp) in pMX vector was purchased from GeneArt (Invitrogen, Supplementary Table 15) and subcloned with V5 epitope into pcDNA3. Different mutants, including: WT IL18RAP CDS + mutant 3'UTR (V1 or V3), and a dominant-negative coding mutant E210A-Y212A-Y214A CDS + WT 3'UTR (3CDS) [41] created by Transfer-PCR mutagenesis[101]. Next, WT and mutants full IL18RAP were subcloned into pUltra vector (a gift from Malcolm Moore, Addgene plasmid #24130, for which mCherry was replaced with EGFP), downstream of the human Ubiquitin C promoter and EGFP-P2A. Cloning procedures were done via restriction-free cloning [102]. List of primers used for cloning and Transfer-PCR mutagenesis described in Supplementary Table 16.

Transfection

Transfection to U2OS cells at 1.9 cm^2 corning plates was performed at 70–80% confluence, 24 hrs. post-plating in antibiotic-free media, using Lipofectamine 2000, 0.5 µL per well (Thermo Fisher Scientific, Cat# 11668027). Each well was considered as a single replicate. For NF-κB reporter assay, U2OS cells were induced with/without recombinant IL-18 (5 ng/mL) 72 hrs. post-transfection with full coding sequence of IL18RAP coding region + 3'UTRs (pUltra vector 500 ng / 1.9 cm^2 plate), luc2P/NF-κB-RE (pGL4.32 100 ng) luciferase and Renilla luciferase (hRluc 10 ng). Following 6 hrs. cells were harvested for Dual-Luciferase® Reporter Assay (E1960) and luminescence was quantified using Veritas™ Microplate Luminometer.

RNA extraction, cDNA synthesis, and quantitative real-time PCR.

Total RNA from LCLs was extracted using Direct-Zol RNA MiniPrep (Zymo Research ,R2052) according to manufacturer instructions. Total RNA from ipMGs was extracted using miRNeasy micro Kit (QIAGEN, 217084) according to manufacturer instructions. Total RNA was reverse transcribed using High Capacity cDNA Reverse Transcription Kit (applied biosystem, 4368814) according to manufacturer instructions, except for the mRNA stability assay, where equal volume of RNA (and not equal amounts of RNA) from each sample was used to generate cDNA. Quantitative Real-time PCR was performed using TaqMan Universal PCR master Mix (applied biosystem, 4304437) or KAPA SYBR FAST (Roche, KK4605). Primers and TaqMan probes are shown in Supplementary Table 16.

Bulk MARS-Seq

200,000 ipMGs harboring variant or canonical IL18RAP 3'UTR (n=4) were treated with 100 ng/mL LPS + 100 ng/mL IL-18 for 6 hrs. in ipMG media (Advanced DMEM (Gibco, 12491-015) containing 1× N-2 supplement (Gibco, 17502-048), 2mM GlutaMAX (Gibco, 35050-038), 55 uM 2-mercaptoethanol (Gibco, 31350-10), 50 U/mL penicillin/streptomycin (Biological Industries, 03-031-1B) and 100 ng/mL IL-34 (Peprotech, 200-34). Following 6 hrs. RNA was extracted as described above and a bulk adaptation of the MARS-Seq protocol (Jaitin et al., Science 2014; Keren-Shaul et al., Nature Protocols, 2019) was used to generate 3' RNA-Seq libraries for expression profiling. Briefly, 50 ng of input RNA from each sample was barcoded during reverse transcription and pooled. Following Agencourt Ampure XP beads cleanup (Beckman Coulter), the pooled samples underwent second strand synthesis and were linearly amplified by T7 in-vitro transcription. The resulting RNA was fragmented and converted into a sequencing-ready library by tagging the samples with Illumina sequences during ligation, RT, and PCR. Libraries were quantified by Qubit and TapeStation as well as by qPCR for GAPDH gene as previously

described (Jaitin et al., Science 2014; Keren-Shaul et al., Nature Protocols, 2019). Sequencing was done on a
NovaSeq 6000 system, SP Reagent Kit, 100 cycles (Illumina; paired-end sequencing).

Analysis of the MARS-seq was done using the UTAP pipeline ([103]; the Weizmann Institute Bioinformatics Unit) to
map the reads to the human genome and to calculate Unique Molecule Identifier (UMI) counts per gene. Reads
were trimmed from their adapter using cutadapt (parameters: -a AGATCGGAAGAGCACACGTCTGAACTCCAGTCAC
-a "A(10)" –times 2 -u 3 -u –3 -q 20 -m 25) and mapped to hg38 genome (STAR v2.4.2a). The pipeline removes UMI
redundancy and quantifies the 3' of RefSeq annotated genes (1,000 bases upstream and 100 bases downstream
of the 3' end). Genes having a minimum of five reads in at least one sample were considered for further analysis.
Differentially expressed (DE) gene detection and count normalization analysis were performed by DESeq2. P-
values in the UTAP results were adjusted for multiple testing using the Benjamini and Hochberg procedure.
Thresholds for significant DE genes: padj < 0.01, |log2FoldChange| >= 0.585, baseMean > 20. This assay was done
with critical advice from Dr. Hadas Keren-Shaul from the Genomics Sandbox unit at the Life Science Core Facility
of Weizmann Institute of Science.

Cell lysis and Western blot
LCLs were washed in PBSx1, centrifuged at 800 × g for 5 min at 4°C, pelleted, and lysed in ice-cold RIPA buffer
(Supplementary Table 17) supplemented with cOmplete™ Protease Inhibitor Cocktail (Roche, 4693116001) and
PhosSTOP™ (Roche, 4906837001). The lysates were cleared by centrifugation at 15,000 × g for 10 min at 4°C.
Protein concentrations quantified with Protein Assay Dye Reagent (Bio-Rad, 500-0006), resolved at 30-50µg of
total protein/well by 8-10% polyacrylamide / SDS gel electrophoresis at 100-120 V for 70 min. After gel
electrophoresis proteins were transferred to nitrocellulose membrane (Whatmann, 10401383) at 250 mA for 70
min. Membranes were stained with Ponceau (Sigma, P7170), blocked for 1 hour at RT with 3% Bovine albumin
fraction V (MPBio 160069) or 5% milk protein in PBST (PBS containing 0.05% TWEEN-20), and then incubated with
primary antibodies (see Supplementary Table 18) O.N. at 4°C with rocking in antibody-Solution [5% albumin, 0.02%
sodium azide, 5 drops of phenol red in 0.05% PBST]. Following primary antibody incubation, membranes were
washed 3 times for 5 min at RT with 0.05% PBST then incubated for 1 hour at RT with horseradish peroxidase
(HRP)-conjugated species-specific secondary antibodies, washed 3 x 5 min in 0.05% PBST at RT, and visualized
using EZ-ECL Chemiluminescence (Biological Industries, 20500-120) by ImageQuant™ LAS 4000 (GE Healthcare Life
Sciences). Densitometric analysis was performed using ImageJ (NIH).

In-Vitro Transcription of biotinylated IL18RAP 3'UTR

To identify the potential trans-acting factors that might differentially bind to the canonical and variant 3'UTRs an

RNA-pulldown and mass spectrometry assay was performed on *in vitro* transcribed canonical and variant forms

of the IL18RAP 3'UTRs, V1 and V3. Briefly, The canonical, V1 and V3 biotinylated-IL18RAP 3'UTR sequences

(384nt), and the negative control (ultrapure water only), were produced by using in vitro transcription HiScribe™

T7 ARCA Kit (NEB, E2060S) following the manufacturer instructions. Briefly, 300 ng of purchased DNA template

(50 ng/uL) (Twist, Supplementary Table 15) was incubated with unlabeled ATP/GTP/CTP and 5% biotin-labelled

UTP, at 37°C for 3 hrs. Next, DNase treatment was performed by incubating the reactions at 37°C for 30 min and

was followed by incubation at 65°C for 10 min to terminate the reaction. The RNA products were purified by an

RNA cleanup purification kit (Zymo Research, R1015). The concentrations of the purified RNA samples were

measured by nanodrop and the expected length was analyzed by TapeStation.

Pull Down of IL18RAP 3'UTR RNA-associated proteins

LCL cell pellets were suspended and lysed in RIPA buffer followed by centrifugation at 15,000xg for 10 min at 4°C.

The concentrations of the cleared supernatants were measured by Bradford assay. 1 mg lysate per sample was

incubated with Pierce streptavidin magnetic beads (Thermo Scientific, 88817) for 30 min at 4°C in rotation, to pre-

clear the lysates from endogenous biotinylated-proteins. To bind IVT products (WT, V1, V3 and negative control;

n=6 repeats/group) to the beads, new prepared binding Pierce streptavidin magnetic beads were incubated by

rotation with equal amounts of IVT products for 30 min at 4°C (100 uL beads/10 pmol RNA product). After 30 min,

the tubes of incubated IVT products with beads were washed three times, and then the cleared lysate was added

equally to each tube and incubated for 30 min at 4°C. In the next step, the samples were washed three times by

magnetizing the beads and resuspended by vortex with a high salt buffer. The bound beads were magnetized and

suspended in 20 ul RNase-free PBSx1 for on-bead digestion procedure.

Liquid Chromatography and Mass Spectrometry

The resulting peptides were analyzed using nanoflow liquid chromatography (nanoAcquity) coupled to high

resolution, high mass accuracy mass spectrometry (Q-Exactive HF). Each sample was analyzed on the instrument

separately in a random order in discovery mode.

Raw proteomic data processing

Raw MS data were processed using MaxQuant version 1.6.6.0 (Cox and Mann, 2008). Database search was performed with the Andromeda search engine (Cox and Mann, 2011; Cox et al., 2011) using the human Uniprot database, appended with common lab protein contaminants. Forward/decoy approach was used to determine the false discovery rate and filter the data with a threshold of 1% false discovery rate (FDR) for both the peptide-spectrum matches and the protein levels. The label-free quantification (LFQ) algorithm in MaxQuant (Cox et al., 2014) was used to compare between experimental samples. Additional settings included the following modifications: Fixed modification- cysteine carbamidomethylation. Variable modifications- methionine oxidation, asparagine and glutamine deamidation, and protein N-terminal acetylation.

Proteomics statistical analysis

ProteinGroups output table was imported from MaxQuant to Perseus v.1.6.2.3 environment (Tyanova et al., 2016). Quality control excluded reverse proteins, proteins identified only based on a modified peptide, and contaminants. Non-specific streptavidin-bead binders were excluded by the following procedure: LFQ Intensity values were log2 transformed, and two outlier samples were excluded from further analysis. Missing values were imputed by creating an artificial normal distribution with a downshift of 1.8 standard deviations and a width of 0.4 of the original ratio distributions. Student's t-test with S0 = 0.1 was performed with FDR P-value ≤ 0.05 between the experimental groups (Canonical, V1 and V3) and the negative control group, which was defined as a single control group. Proteins that passed all QC filters were separated for each of the experimental groups and compared to the negative control samples (ultrapure water). The statistically significant-associated proteins were filtered to retain only proteins that were found in 50% of the repeats in at least one experimental group and were represented by at least one unique peptide. The enriched proteins were subjected to student's t test between every two groups (canonical vs. V1 and canonical vs. V3), with S0 = 0.1, FDR P-value ≤ 0.05 and fold-change threshold >2.

Processing of Mouse Brain Samples for Flow Cytometry

Wild-type C57BL/6 mice were euthanized with CO2 and perfused with PBS through the left ventricle of the heart. Dissected mouse cortex was cut into smaller pieces using scissors and digested in 0.5 mg/mL Collagenase IV (Worthington Biochemical), 10 μg Deoxyribonuclease (Sigma-Aldrich), 10% HI-FBS, RPMI1640 (Gibco) at 37^{o}C for 30 minutes with continuous agitation. Digested samples were gently triturated for 1 minute and the enzymatic reaction was stopped by adding 1 mM EDTA in PBS. The homogenate was filtered through a 100 μm cell strainer and centrifuged at 400 x g for 8 minutes at 4^{o}C to pellet the cells and myelin. This was followed by myelin removal

step by gradient centrifugation with 30% Percoll (Sigma-Aldrich) in PBS (700 x g for 20 minutes at 21oC; without

brakes during deceleration). After myelin (the top white layer) separation, the middle transparent layer was

collected, washed in PBS, and centrifuged at 400 x g for 8 minutes at 4°C to pellet the cells.

Cells pellets were incubated with Mouse Fc block (BD Biosciences 553142), Fixable Viability Stain 620 (BD

Biosciences 564996) and the following antibody mixture in PBS at 4°C for 30 minutes: BV421 Rat Anti-CD11b (BD

Biosciences 562605), BV510 Hamster Anti-Mouse TCR β Chain (BD Biosciences 563221), BV711 Rat Anti-Mouse

Ly-6G (BD Biosciences 563979), APC-Cy7 Rat Anti-Mouse CD45 (BD Biosciences 557659), and Polyclonal Goat IgG

Anti-Mouse IL-18Rβ (R&D Systems AF199). Samples were then washed with PBS and incubated with Alexa Fluor

647 Donkey Anti-Goat IgG (H+L) Cross-Adsorbed Secondary Antibody (Invitrogen A-21447) in PBS at 4°C for 30

minutes. Surface-stained samples were washed with PBS and fixed and permeabilized with BD

Fixation/Permeabilization solution (BD Biosciences 554714) at 4°C for 30 minutes, followed by intracellular

staining with Alexa Fluor 488 Anti-NeuN Antibody (EMD Millipore MAB377X) and eFluor 570 Anti-GFAP

(eBioscience 41-9892-82) in BD Perm/Wash Buffer (BD Biosciences 554714) at 4°C for 30 minutes. Cells were

washed with BD Perm/Wash Buffer and resuspended in PBS for analysis with a FACSymphony (BD Biosciences).

Data were collected as FCS files and analyzed with FlowJo v10 software (BD Biosciences). Antibody specificity was

assessed using relevant isotype control antibodies and fluorescence minus one. Compensation was adjusted using

single-stained samples.

The expression of IL-18RAP (IL-18Rβ) was expressed as Mean Fluorescence Intensity (MFI) or % frequency after

gating for the following cell types: immune cells (CD45^{hi}), microglia (MG: CD45^{int} CD11^{hi}), neurons (CD45^{-} CD11b^{-}

NeuN^{+}), and astrocytes (CD45^{-} CD11b^{-} GFAP^{+}). Animal procedures were approved by the Walter and Eliza Hall

Institute Animal Ethics Committee (Ethics application: 2020.017).

Statistical analysis

Statistics performed with Prism Origin (GraphPad). Shapiro-Wilk test was used to assess normality of the data.

Pairwise comparisons passing normality test were analyzed with Student's *t*-test, whereas the Mann-Whitney test

was used for pairwise comparison of nonparametric data. Multiple group comparisons were analyzed using

ANOVA with post hoc tests. For age of diagnosis and age of death a Permutation Test was used (a Monte-Carlo

simulation test on the t-test between ALS patients harboring canonical or variants of the IL18RAP 3'UTR).

Statistical P-values <0.05 were considered significant. Data are shown as scatter dot plot with mean and SEM, box

plots, or as noted in the text.

Supplementary Materials

Fig. S1. Study design.

Fig. S2. Region-based rare-variant association analyses.

Fig. S3. 3'UTR-based rare-variant association analysis, using different algorithms, and illustration of rare variants

identified in the IL18RAP 3'UTR.

Fig. S4. Restricting burden analysis to the proximal part of 3'UTRs does not improve the association signal.

Fig. S5. IL18RAP and p-NF-κB expression is elevated in lymphoblastoid cells from patients with the C9orf72

repeat expansion.

Fig. S6. IL18RAP 3'UTR variant attenuates IL-18 - NF-κB signaling in U2OS cells.

Fig. S7. IL18RAP is mainly expressed on mouse microglia cells.

Fig. S8. Evaluation of IL18RAP and IL-18 mRNA expression in motor neurons of patients with ALS.

Fig. S9. iPSC-derived microglia express the microglial-specific marker, TMEM119.

Fig. S10. Differentially bound RNA binding proteins to variant 3'UTR (V3) relative to canonical 3'UTR.

Table S1. Total number of samples before and after quality control procedures, stratified by country.

Table S2. Samples quality control procedures.

Table S3. Candidate genes list.

Table S4. Number of rare genetic variants identified.

Table S5. Detailed description of variants in protein-coding sequences of *NEK1* and *SOD1* and the IL18RAP 3'UTR,

in Project MinE discovery cohort.

Table S6. Identified IL18RAP 3'UTR variants in Project MinE discovery cohort.

Table S7. Identified IL18RAP 3'UTR variants in discovery and replication cohorts.

Table S8. Control and patient human cell lines information.

Table S9. Proteomics data from IL18RAP 3'UTR pull-down experiments.

Table S10. Gene set enrichment analysis of differentially bound proteins in canonical vs. V1 IL18RAP 3'UTR.

Table S11. Phenotypic and variant information for patients with protective 3'UTR variants in Project MinE and

NYGC cohorts.

Table S12. MARS-seq data for mRNAs measured in isogenic ipMG harboring variant vs. canonical IL18RAP 3'UTR.

Table S13. KEGG Pathway and Gene Ontology (GO) Biological Process Enrichment Results.

Table S14. BED file containing genomic coordinates of regions of interest.

Table S15. IL18RAP oligonucleotides.

Table S16. List of primers.

Table S17. Medium formulation.

Table S18. Materials and Antibodies.

Data File S1. Source data for IL18RAP and p-NF-κB western blot studies in LCLs (Figure 3D).

Data File S2. Source data for IL18RAP and p-NF-κB western blot studies in Control vs. C9orf72 LCLs (Supplementary
figure 5D).

Data File S3. Source data for IL18RAP western blot studies in isogenic microglia (Figure 4B).

Data File S4. Source data for motor neuron survival assays (Figure 6B,C).

Data File S5. Source data for pNF-κB western blot studies in isogenic microglia, following microglia activation
(Figure 7B).

Project MinE ALS Sequencing Consortium PI List

NYGC ALS Consortium PI List

Supplementary movie. Motor neuron survival was significantly improved in the presence of microglia harboring
variant IL18RAP 3'UTR relative to canonical IL18RAP 3'UTR.

Acknowledgments:

We gratefully acknowledge the contributions of all participants and the investigators who provided biological samples and data for Project Mine ALS sequencing consortium, the New York Genome Center (NYGC) ALS Consortium, the Genome Aggregation Database (gnomAD) and Trans-Omics for Precision Medicine (TOPMed) of the National Heart, Lung, and Blood Institute (NHLBI, https://www.nhlbiwgs.org/topmed-banner-authorship). We thank Michael Ward (NINDS, NIH) for sharing human inducible i^{3}LMN cells. Samples used in this research were in part obtained from the UK National DNA Bank for MND Research, funded by the MND Association and the Wellcome Trust. We acknowledge sample management undertaken by Biobanking Solutions funded by the Medical Research Council at the Centre for Integrated Genomic Medical Research, University of Manchester. The authors would like to thank the NINDS Biorepository at Coriell Institute for iPSC cell lines used in this study. We thank Bernardo Oldak and Prof. Jacob Hanna for microglia differentiation protocols, Dr. Noga Kozer and Dr. Haim Barr for assistance with live cell imaging, Dr. Alon Savidor and Dr. Yishai Levin for mass spectrometry, Dr. Merav Shmueli, Dr. Yifat Merbl and Dr. Ron Rotkof for advice and protocols. We thank LSE for language and scientific editing. Some illustrations were created with BioRender. Hornstein lab is supported by friends of Dr. Sydney Brenner. EH is Head of Andi and Larry Wolfe Center for Research on Neuroimmunology and Neuromodulation and incumbent of Ira & Gail Mondry Professorial chair. **Funding:** The work is funded by Legacy Heritage Fund, Bruno and Ilse Frick Foundation for Research on ALS, Teva Pharmaceutical Industries Ltd as part of the Israeli National Network of Excellence in Neuroscience (NNE) and Minna-James-Heineman Stiftung through Minerva, the European Research Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC grant agreement n° 617351. Israel Science Foundation, the ALS-Therapy Alliance, AFM Telethon (20576 to E.H.), Motor Neuron Disease Association (UK), The Thierry Latran Foundation for ALS research, ERA-Net for Research Programmes on Rare Diseases (FP7), A. Alfred Taubman through IsrALS, Yeda-Sela, Yeda-CEO, Israel Ministry of Trade and Industry, Y. Leon Benoziyo Institute for Molecular Medicine, Kekst Family Institute for Medical Genetics, David and Fela Shapell Family Center for Genetic Disorders Research, Crown Human Genome Center, Nathan, Shirley, Philip and Charlene Vener New Scientist Fund, Julius and Ray Charlestein Foundation, Fraida Foundation, Wolfson Family Charitable Trust, Adelis Foundation, MERCK (UK), Maria Halphen, Estates of Fannie Sherr, Lola Asseof, Lilly Fulop, Andi and Larry Wolfe Center for Research on Neuroimmunology and Neuromodulation and Benoziyo center for Neurological diseases. Weizmann - Brazil Center for Research on Neurodegeneration at The Weizmann Institute of Science, Redhill Foundation – Sam and Jean Rothberg Charitable Trust, Edward and Janie Moravitz, the Israeli Council for Higher Education (CHE) via the Weizmann Data Science Research Center, a research grant from the Estate of Tully and Michele Plesser and M. Judith Ruth Institute for Preclinical Brain Research. To A.A.-C. from Neurodegenerative Disease Research (JPND), Medical Research Council

(MR/L501529/1; STRENGTH, MR/R024804/1; BRAIN-MEND), Economic and Social Research Council (ES/L008238/1; ALS-CarE)), MND Association. National Institute for Health Research (NIHR) Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement n° 772376 - EScORIAL. The collaboration project is co-funded by the PPP Allowance made available by Health~Holland, Top Sector Life Sciences & Health, to stimulate public-private partnerships. This study was supported by the ALS Foundation Netherlands. To P.V.D.: Project MinE Belgium was supported by a grant from IWT (n° 140935), the ALS Liga België, the National Lottery of Belgium and the KU Leuven Opening the Future Fund. P.V.D. holds a senior clinical investigatorship of FWO-Vlaanderen and is supported by E. von Behring Chair for Neuromuscular and Neurodegenerative Disorders, the ALS Liga België and the KU Leuven funds "Een Hart voor ALS", "Laeversfonds voor ALS Onderzoek" and the "Valéry Perrier Race against ALS Fund". Several authors of this publication are members of the European Reference Network for Rare Neuromuscular Diseases (ERN-NMD). To P.J.S: from the Medical Research Council, MND Association, NIHR Senior Investigator Award, National Institute for Health Research (NIHR) Sheffield Biomedical Research Centre, and NIHR Sheffield Clinical Research Facility. To P.M.A.: Knut and Alice Wallenberg Foundation, the Swedish Brain Foundation, the Swedish Science Council, the Ulla-Carin Lindquist Foundation. H.P.P. and sequencing activities at NYGC were supported by the ALS Association (ALSA) and The Tow Foundation. C.E. was supported by scholarship from Teva Pharmaceutical Industries Ltd as part of the Israeli National Network of Excellence in Neuroscience (NNE). S.M.K.F. is supported by the ALS Canada Tim E. Noël Postdoctoral Fellowship. R. H. Brown Jr. was funded by ALS Association, ALS Finding a Cure, Angel Fund, ALS-One, Cellucci Fund and NIH grants (R01 NS104022, R01 NS073873 and NS111990-01 to R.H.B.J.). J.K.I. is a New York Stem Cell Foundation-Robertson Investigator. Work at J.K.I. lab was supported by NIH grants R01NS097850, U.S. Department of Defense grant W81XWH-19-PRARP-CSRA, and grants from the Tau Consortium, the New York Stem Cell Foundation, the ALS Association, and the John Douglas French Alzheimer's Foundation. To R.L.McL.: Science Foundation Ireland (17/CDA/4737). To A.N.B.: Suna and Inan Kirac Foundation. To J.E.L.: National Institute of Health/NINDS (R01 NS073873). **Author contributions:** C.E. and A.Si. led the project; C.E. and A.Si. contributed to research conception, design and interpretations and wrote the manuscript with E.H.; C.E., E.B., T.O., K.R.V.E., S.L.P., M.M., S.M.K.F., N.Y., J.C.-K., K.P.K., R.A.A.V.D.S., W.S., A.A.K., A.I., A.Sh., A.R.J., E.C., D.R., O.W., R.H.B.J., P.J.S., P.V.D., L.H.v.d.B., H.P.P., E.S., A.A.-C. and J.H.V. collected samples, were involved in the sequence analysis pipeline, phenotyping, variant calling, provided expertise or were involved in the genetic association analysis of rare non-coding variants in human patients with ALS; S.-T.H. and J.K.I. provided stem cells and initial data; S.B.-D., E.A., G.B. and H.M.-K. were involved in the design, generation and validation of CRISPR-edited IL18RAP isogenic iPSCs; H.M.-K. and Y.M.D. performed the pull down experiments of

IL18RAP 3'UTR RNA-associated proteins and analyzed the proteomic data; A.Si. and C.E. established human iPSC-

derived microglia differentiation and culturing protocols, performed motor neuron survival experiments and

interpreted data; A.Si., N.R. and C.E. performed molecular biology studies in LCLs and U2OS cell lines, including

reporter assays, qPCR and protein quantification by western blots; E.Y. performed Bulk MARS-Seq experiment; C.-

H.Y., C.L. and S.L.M. provided expertise and processed mouse cortex samples for Flow Cytometry; Y.C., Y.E.-A.,

S.W. and D.P.S. helped performing research; E.H. conceived and supervised the study and wrote the manuscript

with C.E. and A.Si. All co-authors provided approval of the manuscript.

Competing interests: J.K.I. is a co-founder of AcuraStem Incorporated. J.K.I. declares that he is bound by

confidentiality agreements that prevent him from disclosing details of his financial interests in this work. J.H.V.

and L.H.v.d.B. report to have sponsored research agreements with Biogen. E.H. is inventor on pending patent

family PCT/IL2016/050328 entitled "Methods of treating motor neuron diseases". All other authors declare that

they have no competing interests. **Data availability:** Human genetics data is publically available from the

sequencing consortia: Project Mine ALS sequencing consortium, the New York Genome Center (NYGC) ALS

Consortium, the Genome Aggregation Database (gnomAD), and NHLBI's Trans-Omics for Precision Medicine

(TOPMed). Gene Expression Omnibus accession number: GSE186757. All Other data used for this manuscript are

available in the manuscript. **Code availability:** Variant annotation scripts are available at GitHub:

https://github.com/TsviyaOlender/Non-coding-Variants-in-ALS-genes-.

FIGURES

Fig.1 - Eitan et al. (Hornstein)

Fig. 1. Region-based rare-variant association analysis reveals association of IL18RAP 3'UTR with ALS. (A) Diagram of study design. Collapsed region-based rare-variant (MAF ≤0.01) association analysis was performed on: (i) 295 candidate protein-coding genes (Supplementary Table 3), encoding for ALS-relevant proteins or proteins associated with miRNA biogenesis/activity. Variants were included if predicted to cause frameshifting, alternative splicing, an abnormal stop codon, or a deleterious non-synonymous amino acid substitution, in ≥ 3 of 7 independent dbNSFP prediction algorithms; (ii) variants in 3'-untranslated regions (3'UTRs) of the 295 genes (Supplementary Table 3); (iii) all known autosomal pre-miRNA genes in the human genome; and (iv) predicted networks, comprised of aggregated variants detected in a specific mature miRNA sequence and its cognate down-stream 3'UTR targets. **(B)** QQ plot of obtained and expected P-values for the burden of rare variants (log scale), gained by collapsed region-based association analysis of all genomic regions described in (A). Data were obtained from 3,955 ALS cases and 1,819 controls (Project MinE). Features positioned on the diagonal line represent results obtained under the null hypothesis. Open-reading frames of 10 known ALS genes (blue). IL18RAP 3'UTR (red). Genomic inflation λ = 1.2. **(C)** QQ plot of obtained and expected P-values for the burden of rare variants (log scale), gained by collapsed region-based association analysis for all known human 3'UTRs (RefSeq). The IL18RAP 3'UTR (red) is the most significant 3'UTR associated with ALS. P-values, calculated with Optimized Sequence Kernel Association Test, SKAT-O (genomic inflation λ = 0.97).

A

Gene	Region	ALS (3955)	Control (1819)	OR	OR 95% CI	P	P corrected
NEK1	Coding	127	19	3.14	1.93-5.11	8.00×10^{-7}	2.30×10^{-4}
SOD1	Coding	36	0	33.89	2.08-552.47	2.60×10^{-4}	3.73×10^{-2}
IL18RAP	3'UTR	*6*	*12*	0.23	0.09-0.61	1.93×10^{-5}	5.41×10^{-3}

B

Cohort	Cases	Control	OR	OR 95% CI
Ireland	0/239	1/136	0.19	0.008-4.662
Netherlands	2/1633	7/1004	0.17	0.036-0.842
Turkey	0/142	1/67	0.16	0.006-3.870
UnitedKingdom	2/1043	2/272	0.26	0.036-1.850
USA	2/398	1/68	0.34	0.030-3.784
Belgium	0/295	0/172	-	-
Spain	0/205	0/100	-	-
Total	6/3955	12/1819	0.23	0.086-0.611

C

Cohort	Cases	Control	OR	95% CI	P χ2	P SKAT-O
Discovery: Project MinE	6/3955	12/1819	0.23	0.086-0.611	3.00×10^{-3}	1.93×10^{-5}
Replication: NYGC, TOPMed & gnomAD	8/2184	786/68584	0.32	0.158-0.637	9.58×10^{-4}	-
Joint analysis: Discovery & replication	14/6139	798/70403	0.20	0.118-0.338	$< 1.00 \times 10^{-5}$	-

Fig.2 - Eitan et al. (Hornstein)

Fig 2. Odds of ALS are reduced with rare variants in the IL18RAP 3'UTR. (A) Odds ratio (OR) estimates with 95%

confidence intervals (CI) for NEK1 (coding), SOD1 (coding) and IL18RAP (3'UTR). P-values corrected for false

discovery rate (FDR). **(B)** Stratification of data pertaining to IL18RAP 3'UTR in seven geographically-based sporadic

ALS sub-cohorts and forest plot (OR on log scale with whiskers for 95% CI). *NEK1* (grey) and *SOD1* (blue) signals

are from combined data of all cohorts. Vertical dotted line denotes OR=0.23. **(C)** Stratification of IL18RAP 3'UTR

variants data across discovery and replication cohorts and joint analysis thereof; Forest plot (OR on log scale with

whiskers for 95% CI). Vertical dotted line denotes OR=0.2. P-values, calculated with SKAT-O or Chi-squared test

with Yate's correction.

Lymphoblastoid cell lines

Fig. 3. IL18RAP 3'UTR variant correlates with attenuated IL-18 - NF-κB signaling in human lymphoblastoid cells.

(A) IL18RAP mRNA expression (qPCR normalized to IPO8 mRNA levels) and **(B)** IL18RAP or **(C)** p-NF-κB protein

expression (Western blots, normalized to Tubulin). Scatter dot plot with mean and SEM. **(D)** Representative blots

processed with anti-IL18RAP, anti p-NF-κB and anti-Tubulin antibodies. Extracts from twelve different human

lymphoblastoid cell lines (listed in Supplementary Table 8): Four lines of healthy individuals (without ALS), carrying

the canonical IL18RAP 3'UTR sequence (Control; Canonical IL18RAP 3'UTR, black); Four sporadic ALS patients,

carrying the canonical IL18RAP 3'UTR sequence (sALS; Canonical IL18RAP 3'UTR, red); Two healthy individuals,

carrying a variant form of IL18RAP 3'UTR (Control; Variant IL18RAP 3'UTR, light blue) and two sporadic ALS

patients carrying a variant form of IL18RAP 3'UTR (sALS; Variant IL18RAP 3'UTR, navy blue). One-way ANOVA

followed by Newman-Keuls multiple comparisons test, was conducted based on the mean value of three

independent passages for each of the twelve human lymphoblastoid cell lines. * $P<0.05$; ** $P<0.01$; *** $P<0.001$.

A

① **Genome editing of C9-ALS iPSCs**

② **iPSC-derived microglia differentiation**

Fig. 4. IL18RAP 3'UTR variant destabilizes IL18RAP mRNA in CRISPR-edited isogenic iPSC-derived microglia, with

C9orf72 genetic background. (A) Diagram of experimental design. (1) Genome editing with CRISPR Cas9 of point

mutations that recapitulate the most prevalent variants (Chr2:103068691 C>T (V1) and Chr2:103068718 G>A (V3))

in the IL18RAP 3'UTR sequence in human induced pluripotent stem cells (iPSCs) donated by ALS patients with a

C9orf72 repeat expansion ([66] NINDS/Coriell Code: ND10689, ND12099, Supplementary Table 8). The two

independent isogenic pairs of cell lines both carry the C9orf72 repeat expansion and vary only by the presence of

the canonical or a variant IL18RAP 3'UTR. (2) The four IL18RAP 3'UTR lines (two isogenic pairs) were differentiated

into human microglia [68]. Dot plots of IL18RAP **(B)** protein levels (by Western blot analysis, normalized to GAPDH,

N=3, Data File S3) and **(C)** mRNA (by qPCR, normalized to IPO8 mRNA N=3) in differentiated human microglia. **(D)**

IL18RAP mRNA degradation rate studied in human isogenic microglia at 0, 4 and 6 hrs after introduction of a

transcriptional block with actinomycin D (7.5 µg/mL, Sigma-Aldrich A9415) (by qPCR, normalized to average of

IPO8 and GAPDH mRNA expression, n = 4 independent wells per time point with two technical duplicates). Variant

3'UTR destabilizes the IL18RAP mRNA relative to the canonical sequence. Scatter dot plot with mean and SEM.

Two sided t-test P- values * <0.05, ** <0.01, **** <0.0001.

Fig. 5. Reduced association of double-stranded RNA binding proteins to variant IL18RAP 3'UTR. (A) Diagram of

mass spectrometry of RNA binding proteins pulled-down by IL18RAP 3'UTR sequences (canonical, V1 and V3). **(B)**

Principal-component analysis (PCA) of IL18RAP 3'UTR-associated proteomes pulled down by the canonical (grey,

N=6 experimental repeats), V3 (orange, N=5), and V1 (blue, N=5) biotin-tagged, in-vitro transcribed oligos. **(C)**

Volcano plot of protein abundance associated with the canonical relative to variant (V1) IL18RAP 3'UTR (x-axis

log2 scale), analyzed by MS. Y-axis depicts P-values (−log10 scale). Proteins significantly enriched in association

with canonical/variant 3'UTR are colored (grey/blue). Double-stranded RNA-binding proteins (dsRBPs) are

demarcated by a purple oval. Features above the horizontal dashed line demarcate proteins with adjusted p <

0.05, in student's t-test with FDR correction to multiple hypotheses. Vertical dashed lines are of 2 or ½ fold change.

A non-significant data point of KIF13B (P-value = 0.08) is not shown for clarity of the illustration (Supplementary

Table 9) **(D)** Volcano plot of normalized enrichment score of the Gene Ontology (GO) molecular function gene sets

from GSEA analysis of differentially expressed proteins (canonical vs. V1 IL18RAP 3'UTR). Reduced association of

double-stranded RNA binding proteins (GO:0003725; purple) with V1 IL18RAP 3'UTR, relative to the canonical

3'UTR. All gene sets are described in Supplementary Table 10. **(E)** Profile of GSEA enrichment score and positions

of the 10 double-stranded RNA binding proteins (purple) within all differentially expressed proteins, ranked from

most enriched in canonical 3'UTR to most depleted protein (Supplementary Table 10; WebGestalt [104]). **(F)**

Prediction of 3'UTR secondary structure by RNA Fold [105] suggests a more stable dsRNA structure of canonical 3'UTR

(green), with lower minimum free energy (MFE) than that of the sequence harboring a V1 variant (red).

A Co-culture of isogenic IL18RAP 3'UTR C9-ALS microglia and WT iMNs

C9-ALS microglia line 2
C9-ALS microglia line 1

Canonical IL18RAP 3'UTR — Isogenic pair — Variant IL18RAP 3'UTR

+

WT i³LMNs
Syn::GFP+

Microglia activation
LPS+IL-18

Longitudinal tracking of neuron survival (20 days)

Survival — Variant / Canonical — Time

B

C9-ALS microglia line 1

Motor neuron survival (%)

Variant IL18RAP
Canonical IL18RAP

**

C9-ALS microglia line 2

Motor neuron survival (%)

Time (Days)

**

C

Motor neuron survival at day 20 (%)

**** **** ****

**** *** ****

Rep1 Rep2 Rep3

D

i³LMNs in co-culture at day 20

Variant IL18RAP 3'UTR

Canonical IL18RAP 3'UTR

200 µm

E

Human ALS patients

Age of death (years)

*

ALS Cohort   ALS with IL18RAP variants

F

Age of diagnosis (years)

*

ALS Cohort   ALS with IL18RAP variants

Fig. 6. Variant IL18RAP 3'UTR are protective in human microglia and in patients with ALS. (A) Diagram of experimental design. Co-culture of human iPSC-derived transcription factor-induced motor neurons (i[3]LMNs) that express GFP driven by the synapsin (Syn) promoter (healthy, non-ALS, [76]) and human iPSC-derived isogenic IL18RAP 3'UTR microglia (on a *C9orf72 repeat expansion background)*. Time-lapse microscopic analyses of i[3]LMNs survival, after microglia activation with a cocktail of LPS and the cytokine IL-18. **(B,C)** i[3]LMNs survival over 20 days in the presence of microglia harboring variant (blue) or canonical (grey) IL18RAP 3'UTR (two independent isogenic pairs, based on independent patient C9orf72 lines, n=3 independent differentiation procedures from different passages per line, with 3-8 co-culture wells per passage). **(B)** Survival plot of i[3]LMNs in a representative experiment for each isogenic pair (Two-way ANOVA) and **(C)** Box plot depicting the percentage of i[3]LMNs survival on day 20 of co-culture, median, upper and lower quartiles of all experiments. Two independent isogenic pairs, based on independent patient C9orf72 lines, n=3 independent differentiation procedures from different passages per line, with 3-8 co-culture wells per passage. Two-way ANOVA followed by Tukey's multiple comparison test. **(D)** Representative micrographs of fluorescent i[3]LMNs after 20 days of culture with C9-ALS microglia. **(E)**. Association of age of death (9 patients with protective 3'UTR variants /4263 patients with available phenotypic data in Project MinE and NYGC cohorts, or **(F)** age of diagnosis (8/4216 patients). IL18RAP variant is associated with delayed age of death (+6.1 years, Permutation P-value = 0.02, Cohen's d effect size = 0.65) and age of diagnosis (+6.2 years, Permutation P-value = 0.05, Cohen's d effect size = 0.62), relative to the mean age of all Project MinE and NYGC ALS patients. Box plots depicting median, upper and lower quartiles, and extreme points. * P<0.05, ** P<0.01, *** P<0.001, **** P<0.0001.

Fig. 7. Variant IL18RAP 3'UTR dampens neurotoxic NF-κB signaling in human microglia. (A) Diagram of
experimental design. Four IL18RAP 3'UTR lines (two isogenic pairs) were differentiated into human microglia [68]
and analyzed for phosphorylated NF-κB protein levels, transcriptomics, and neuronal survival in co-culture
with/without IKK16, following activation with a cocktail of LPS and the cytokine IL-18, for 0.5h, 6h and 20 days,
respectively. **(B)** Western blot analysis revealed reduced levels of phosphorylated NF-κB in variant IL18RAP 3'UTR
relative to isogenic control. Scatter dot plot with mean and SEM (Two sided t-test P-value *** <0.001, N=3, Data
File S5). mRNA extracted from human microglia was subjected to a next generation sequencing study with
downstream bioinformatics studies **(C-E)**. Over-representation analysis (ORA) within **(C)** KEGG Pathways and **(D)**
Gene Ontology biological processes, of the differentially expressed transcriptome in microglia harboring variant
vs. canonical IL18RAP 3'UTR. Bar graph depicting the Ratio of enrichment for significantly enriched pathways (FDR
≤ 0.05) are shown (Supplementary Table 13; WebGestalt [104]). **(E)** Unsupervised study of the NF-κB transcriptomic
signature (GO:0007249 pathway-associated mRNAs) in microglia with the variant relative to the isogenic canonical
IL18RAP 3'UTR. **(F)** Time-lapse microscopic analyses of co-cultured human i^{3}LMNs (healthy, non-ALS, [76]) with
human iPSC-derived isogenic IL18RAP 3'UTR microglia (on a *C9orf72 repeat expansion background)*, activated with
a cocktail of LPS and the cytokine IL-18, without (carrier alone, DMSO), or with IKK16 (200nM), a selective IκB
kinase (IKK) inhibitor that inhibits NF-κB signaling [81]. IKK16 significantly ameliorates motor neuron death, relative
to control only in the context of canonical IL18RAP 3'UTR, but did not further contribute to rescue in human
microglia with the protective variant IL18RAP 3'UTR (two independent isogenic pairs, based on independent
patient C9orf72 lines with 3-8 co-culture wells per line). Two-way ANOVA P-value * <0.05.

A

WGS sequencing, variant calling and QC

↓

Capture regions of interest from WGS

↓

Bioinformatics assay and functional annotation

↓

Region-based rare-variant association analysis

↓

Genetic and biological validation

B

ALS   Control

3'UTR and ORF of ALS-associated genes or genes encoding RBPs

SOD1   ORF   3'UTR   AAA

miRNA genes

Supplementary Fig.1 - Eitan et al. (Hornstein)

871   **Supplementary Fig. 1. Study design. (A)** Flow chart of approach for discovery of region-based rare-variants in

872   non-coding genomic regions via association studies and **(B)** diagram depicting regions of interest comprising of

873   1,750 autosomal human pre-miRNA genes, 295 open reading frames encoding for proteins of interest, and 295

874   3'UTRs.

Supplementary Fig.2 - Eitan et al. (Hornstein)

Supplementary Fig. 2. Region-based rare-variant association analyses. (A-D) QQ plot of obtained and expected

P-values for the burden of rare-variants (log scale) gained by collapsed region-based association analysis of

different genomic regions, comprised of (A) 295 candidate protein-coding regions listed in Supplementary Table

3. These ORFs encode for ALS-relevant proteins or proteins that are associated with miRNA biogenesis or activity.

Variants were depicted if predicted to cause frameshifting, alternative splicing, abnormal stop codon or a

deleterious non-synonymous amino acid substitution, in ≥ 3 of 7 independent dbNSFP prediction algorithms

(genomic inflation λ = 0.96), (B) All known pre-miRNA genes in the human genome (genomic inflation λ = 1.31),

(C) predicted networks, comprised of aggregated variants detected on a specific mature miRNA sequence and its

cognate downstream 3'UTR targets (genomic inflation λ = 1.16), and (D) variants in 3'UTRs of the same 295 genes

listed in Supplementary Table 3 (genomic inflation λ = 1.08). Data was obtained from 3,955 ALS cases and 1,819

controls (Project MinE). Features positioned on the diagonal line represent results obtained under the null

hypothesis. Open-reading frames of 10 known ALS genes (blue). IL18RAP 3'UTR (red). P-values, calculated with

SKAT-O.

Supplementary Fig. 3. 3'UTR-based rare-variant association analysis, using different algorithms, and illustration
of rare variants identified in the IL18RAP 3'UTR. (A-D) QQ plot of obtained and expected P-values for the burden
of rare variants (log scale) gained by collapsed region-based association analysis of genomic regions comprised of
295 3'UTRs listed in Supplementary Table 3, in the Project MinE cohort (3,955 ALS cases and 1,819 non-ALS
controls). Features positioned on the diagonal line represent results obtained under the null hypothesis. IL18RAP
3'UTR (red) is the most significant 3'UTR associated with ALS using different algorithms: **(A)** Sequence Kernel
Association Test, SKAT (genomic inflation λ = 1.02), **(B)** Combined Multivariate and Collapsing, CMC (genomic
inflation λ = 1.34), **(C)** Variable Threshold with permutation analysis, VT (genomic inflation λ = 1.03). **(D)** IL18RAP
3'UTR also ranked as the top hit when aggregating variants abrogating or gaining miRNA recognition elements
(MREs) in 3'UTRs (genomic inflation λ = 1.04). **(E)** Schematic of the IL18RAP transcript and 3'UTR (5' to 3') showing
the number of control (upper) or ALS (lower) samples in which variants (black arrow) were identified in the Project
MinE discovery cohort (Supplementary Table 6).

A

B

Differences plot

Supplementary Fig. 4. Restricting burden analysis to the proximal part of 3'UTRs does not improve the association signal. (A) Scatter plot with SKAT-O P-values (log scale) calculated for the burden of rare variants gained by collapsed region-based association analysis of the full 3'UTRs on the x-axis versus the 3'UTRs proximal quadrant on the y-axis, for the 295 3'UTRs listed in Supplementary Table 3, in the Project MinE cohort (3,955 ALS cases and 1,819 non-ALS controls) (Pearson correlation coefficient (r=0.61) and P-value ****<0.0001). The 45-degree diagonal line represents a perfect correlation of r=1. IL18RAP 3'UTR (red). **(B)** A Difference plot showing the difference between the two P-value measurements (3'UTRs proximal quadrant minus the full 3'UTRs). The bias (difference between means) is only 0.03. Overall the P-values gained from the 3'UTRs proximal quadrant were comparable to that of the full 3'UTRs in the cohort of 295 3'UTRs. Box plots depict median, upper and lower quartiles, and extreme points (Wilcoxon matched-pairs P-value > 0.05, Cohen's d effect size = 0.1). Hence, the apparent spatial distribution of variants in IL18RAP 3'UTR seems to be a particular case, rather than part of a global pattern.

Lymphoblastoid cell lines

A

IL18RAP mRNA

ns | ● Control;   Canonical IL18RAP 3'UTR
   | ● C9orf72;   Canonical IL18RAP 3'UTR

B

IL18RAP protein

p=0.056 | ● Control;   Canonical IL18RAP 3'UTR
        | ● C9orf72;   Canonical IL18RAP 3'UTR

C

pNF-κB protein

** | ● Control;   Canonical IL18RAP 3'UTR
   | ● C9orf72;   Canonical IL18RAP 3'UTR

D

Control;
Canonical
IL18RAP
3'UTR

C9orf72;
Canonical
IL18RAP
3'UTR

IL18RAP
68 kDa

Tubulin
52 kDa

pNF-κB
65 kDa

Tubulin
52 kDa

Supplementary Fig. 5. IL18RAP and p-NF-κB expression is elevated in lymphoblastoid cells from patients with

the C9orf72 repeat expansion. (A) IL18RAP mRNA expression (qPCR normalized to IPO8 mRNA levels) and **(B)**

IL18RAP or **(C)** p-NF-κB protein expression (Western blots, normalized to Tubulin). Extracts from eight different

human lymphoblastoid cell lines (listed in Supplementary Table 8): Four lines of healthy individuals (without ALS)

carrying the canonical IL18RAP 3'UTR sequence (Control; Canonical IL18RAP 3'UTR, black) and four C9orf72 ALS

patients carrying the canonical IL18RAP 3'UTR sequence (C9orf72; Canonical IL18RAP 3'UTR, red). **(D)**

Representative blots processed with anti-IL18RAP, anti p-NF-κB and anti-Tubulin antibodies. Mann-Whitney test

(A) or one-sided student's t-test with Welch's correction on log-transformed data (B,C), was conducted based on

the mean value of three independent passages for each of the eight human lymphoblastoid cell lines. Scatter dot

plot with mean and SEM. ** P<0.01.

A Biosensor for NF-κB pathway activity

B U2OS cells

Supplementary Fig. 6. IL18RAP 3'UTR variant attenuates IL-18 - NF-κB signaling in U2OS cells. Diagram **(A)** and

quantification **(B)** of NF-κB reporter assays in human U2OS cell line. To determine the ability of the IL18RAP

variants V3 and V1 to induce NF-κB activity, U2OS cells were co-transfected with different IL18RAP coding region

(CDS) and 3'UTR constructs (GFP, Canonical, V3, V1, n=9; 3CDS, *n*=4), along with an NF-κB activity reporter that

drives luciferase (Luc2P) transcription via five copies of the NF-κB response element. NF-κB signaling was induced

by adding human recombinant IL-18 to the medium. Variants V3 and V1 of the IL18RAP 3'UTR reduced NF-κB

activity by ~10% and ~21%, respectively, relative to the WT IL18RAP 3'UTR. GFP vector and a dominant-negative

coding mutant E210A-Y212A-Y214A CDS + WT 3'UTR (3CDS) [41], served as controls. Luciferase expression was

normalized to transfected U2OS cells that were not induced with human recombinant IL-18. One-way ANOVA

followed by Dunnett's multiple comparison test was performed on square root-transformed data. Box plots depict

median, upper and lower quartiles, and extreme points. * $P<0.05$; *** $P<0.001$. The experiment was repeated

independently three times with similar results.

A

Supplementary Fig. 7. IL18RAP is mainly expressed on mouse microglia cells. (A-C) Flow cytometry was used to

characterize IL18RAP expression levels in dissociated wild-type mouse cortex cells. The expression of IL-18RAP (IL-

18Rβ) was expressed as Mean Fluorescence Intensity (MFI) and % frequency after gating for the following cell

types: immune cells (CD45hi), microglia (MG: CD45int CD11hi), neurons (CD45- CD11b- NeuN+), and astrocytes

(CD45- CD11b- GFAP+). FACS analysis reveals that IL18RAP is mainly expressed on microglia cells. A scatter dot

plot with mean and SEM values for the median fluorescence intensity (MFI) and percentage of IL18RAP+ cells is

shown. One-way ANOVA followed by Tukey's multiple comparison test. ** $P<0.01$, *** $P<0.001$, **** $P<0.0001$.

1000

1001

1002

1003

1004

1005

1006

1007

1008

1009

1010

1011

1012

Supplementary Fig. 8. Evaluation of IL18RAP and IL-18 mRNA expression in motor neurons of patients with ALS. **(A-B)** mRNA expression of IL18RAP **(A)** and IL-18 **(B)**, as reads per kilobase million (RPKM), from NGS study of laser capture microdissection–enriched surviving motor neurons from lumbar spinal cords of patients with sALS with rostral onset and caudal progression (n = 12) and non-neurodegeneration controls (n = 9; [106] GSE76220). Two-sided Student's t test with Welch's correction on log-transformed data. **(C)** IL-18 mRNA expression, as log2-normalized counts, from NGS study of induced ALS motor neurons (n = 4 different donors in duplicates) or non-neurodegeneration controls (n=3 different donors in duplicates; [107] DESeq analysis). Box plots depict median, upper and lower quartiles, and extreme points. *P < 0.05; **P < 0.01.

C9-ALS microglia line 1

C9-ALS microglia line 2

1037   **Supplementary Fig. 9. iPSC-derived microglia express the microglial-specific marker, TMEM119.**

1038   Immunofluorescence staining of TMEM119 (green) and DAPI (blue), in two different C9orf72 iPSC-derived

1039   progenitor microglia lines. Lens, ×20; scale bar, 100 μm.

1040

1041

1042

1043

1044

1045

1046

1047

1048

1049

1050

1051

1052

1053

1054

1055

1056

1057

1058

Supplementary Fig. 10. Differentially bound RNA binding proteins to variant 3'UTR (V3) relative to canonical

3'UTR. (A) Volcano plot of protein abundance associated with the canonical relative to variant (V3) IL18RAP 3'UTR

(x-axis log2 scale), analyzed by MS. Y-axis depicts P-values (−log10 scale). Proteins significantly enriched in

association with canonical/variant 3'UTR are colored (grey/orange). Features above the horizontal dashed line

demarcate proteins with adjusted p < 0.05, in student's t-test with FDR correction to multiple hypotheses. Vertical

dashed lines are of 2 or ½ fold change (Supplementary Table 9). **(B)** Prediction of 3'UTR secondary structure by

RNA Fold [105], suggests a minor change to the structure of the sequence harboring a V3 variant (red), relative to

the canonical 3'UTR (green).

1067

1068

1069

1070

1071

1072

1073

1074

1075

1076

1077

1078

1079

1080

1081

1082

1083   **References**

1084

1.   Cookson, W., Liang, L., Abecasis, G., Moffatt, M. & Lathrop, M. Mapping complex disease traits with
global gene expression. *Nat Rev Genet* **10**, 184-194 (2009).

2.   Knight, J.C. Regulatory polymorphisms underlying complex disease traits. *J Mol Med (Berl)* **83**, 97-109
(2005).

3.   Brown, R.H. & Al-Chalabi, A. Amyotrophic Lateral Sclerosis. *N Engl J Med* **377**, 162-172 (2017).

4.   Taylor, J.P., Brown, R.H., Jr. & Cleveland, D.W. Decoding ALS: from genes to mechanism. *Nature* **539**,
197-206 (2016).

5.   Renton, A.E., Chio, A. & Traynor, B.J. State of play in amyotrophic lateral sclerosis genetics. *Nat Neurosci*
17, 17-23 (2014).

6.   Al-Chalabi, A., van den Berg, L.H. & Veldink, J. Gene discovery in amyotrophic lateral sclerosis:
implications for clinical management. *Nat Rev Neurol* **13**, 96-104 (2017).

7.   van Rheenen, W.*, et al.* Common and Rare Variant Association Analyses in Amyotrophic Lateral Sclerosis
Identify 15 Risk Loci with Distinct Genetic Architectures and Neuron-Specific Biology. *MedRxiv* (2021).

8.   DeJesus-Hernandez, M.*, et al.* Expanded GGGGCC hexanucleotide repeat in noncoding region of
C9ORF72 causes chromosome 9p-linked FTD and ALS. *Neuron* **72**, 245-256 (2011).

9.   Renton, A.E.*, et al.* A hexanucleotide repeat expansion in C9ORF72 is the cause of chromosome 9p21-
linked ALS-FTD. *Neuron* **72**, 257-268 (2011).

10.   La Spada, A.R. & Taylor, J.P. Repeat expansion disease: progress and puzzles in disease pathogenesis.
Nat Rev Genet **11**, 247-258 (2010).

11.   Cooper-Knock, J.*, et al.* Rare Variant Burden Analysis within Enhancers Identifies CAV1 as an ALS Risk
Gene. *Cell Rep* **33**, 108456 (2020).

12.   Povysil, G.*, et al.* Rare-variant collapsing analyses for complex traits: guidelines and applications. *Nat Rev*
Genet **20**, 747-759 (2019).

13.   An, J.Y.*, et al.* Genome-wide de novo risk score implicates promoter variation in autism spectrum
disorder. *Science* **362** (2018).

14.   Bartel, D.P. MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215-233 (2009).

15.   Mayr, C. Regulation by 3'-Untranslated Regions. *Annu Rev Genet* **51**, 171-194 (2017).

16.   Haramati, S.*, et al.* miRNA malfunction causes spinal motor neuron disease. *Proc Natl Acad Sci U S A* **107**,
13111-13116 (2010).

17.   Emde, A*., et al.* Dysregulated miRNA biogenesis downstream of cellular stress and ALS-causing

mutations: a new mechanism for ALS. *EMBO J* **34**, 2633-2651 (2015).

18.   Eitan, C. & Hornstein, E. Vulnerability of microRNA biogenesis in FTD-ALS. *Brain Res* (2016).

19.   Campos-Melo, D., Droppelmann, C.A., He, Z., Volkening, K. & Strong, M.J. Altered microRNA expression

profile in Amyotrophic Lateral Sclerosis: a role in the regulation of NFL mRNA levels. *Mol Brain* **6**, 26 (2013).

20.   Buratti, E*., et al.* Nuclear factor TDP-43 can affect selected microRNA levels. *FEBS J* **277**, 2268-2281

(2010).

21.   Kawahara, Y. & Mieda-Sato, A. TDP-43 promotes microRNA biogenesis as a component of the Drosha

and Dicer complexes. *Proc Natl Acad Sci U S A* **109**, 3347-3352 (2012).

22.   Morlando, M*., et al.* FUS stimulates microRNA biogenesis by facilitating co-transcriptional Drosha

recruitment. *EMBO J* **31**, 4502-4510 (2012).

23.   Hoye, M.L*., et al.* MicroRNA Profiling Reveals Marker of Motor Neuron Disease in ALS Models. *J Neurosci*

37, 5574-5586 (2017).

24.   Rotem, N*., et al.* ALS Along the Axons - Expression of Coding and Noncoding RNA Differs in Axons of ALS

models. *Sci Rep* **7**, 44500 (2017).

25.   Butovsky, O*., et al.* Modulating inflammatory monocytes with a unique microRNA gene signature

ameliorates murine ALS. *J Clin Invest* **122**, 3063-3087 (2012).

26.   Figueroa-Romero, C*., et al.* Expression of microRNAs in human post-mortem amyotrophic lateral

sclerosis spinal cords provides insight into disease mechanisms. *Mol Cell Neurosci* **71**, 34-45 (2016).

27.   Williams, A.H*., et al.* MicroRNA-206 delays ALS progression and promotes regeneration of

neuromuscular synapses in mice. *Science* **326**, 1549-1554 (2009).

28.   Hickman, S., Izzy, S., Sen, P., Morsett, L. & El Khoury, J. Microglia in neurodegeneration. *Nat Neurosci* **21**,

1359-1369 (2018).

29.   Beers, D.R. & Appel, S.H. Immune dysregulation in amyotrophic lateral sclerosis: mechanisms and

emerging therapies. *Lancet Neurol* **18**, 211-220 (2019).

30.   Lall, D. & Baloh, R.H. Microglia and C9orf72 in neuroinflammation and ALS and frontotemporal

dementia. *J Clin Invest* **127**, 3250-3258 (2017).

31.   Zhao, W*., et al.* TDP-43 activates microglia through NF-kappaB and NLRP3 inflammasome. *Exp Neurol*

273, 24-35 (2015).

32.   Vahsen, B.F*., et al.* Non-neuronal cells in amyotrophic lateral sclerosis - from pathogenesis to

biomarkers. *Nat Rev Neurol* **17**, 333-348 (2021).

33.   McCauley, M.E. & Baloh, R.H. Inflammation in ALS/FTD pathogenesis. *Acta Neuropathol* **137**, 715-730
(2019).

34.   Motataianu, A., Barcutean, L. & Balasa, R. Neuroimmunity in amyotrophic lateral sclerosis: focus on
microglia. *Amyotroph Lateral Scler Frontotemporal Degener* **21**, 159-166 (2020).

35.   Philips, T. & Robberecht, W. Neuroinflammation in amyotrophic lateral sclerosis: role of glial activation
in motor neuron disease. *Lancet Neurol* **10**, 253-263 (2011).

36.   Kadhim, H., Deltenre, P., Martin, J.J. & Sebire, G. In-situ expression of Interleukin-18 and associated
mediators in the human brain of sALS patients: Hypothesis for a role for immune-inflammatory mechanisms.
Med Hypotheses **86**, 14-17 (2016).

37.   Johann, S.*, et al.* NLRP3 inflammasome is expressed by astrocytes in the SOD1 mouse model of ALS and
in human sporadic ALS patients. *Glia* **63**, 2260-2273 (2015).

38.   Italiani, P.*, et al.* Evaluating the levels of interleukin-1 family cytokines in sporadic amyotrophic lateral
sclerosis. *J Neuroinflammation* **11**, 94 (2014).

39.   Huang, F.*, et al.* Longitudinal biomarkers in amyotrophic lateral sclerosis. *Ann Clin Transl Neurol* **7**, 1103-
1116 (2020).

40.   Alboni, S., Cervia, D., Sugama, S. & Conti, B. Interleukin 18 in the CNS. *J Neuroinflammation* **7**, 9 (2010).

41.   Tsutsumi, N.*, et al.* The structural basis for receptor recognition of human interleukin-18. *Nature*
communications **5**, 1-13 (2014).

42.   Adachi, O.*, et al.* Targeted disruption of the MyD88 gene results in loss of IL-1- and IL-18-mediated
function. *Immunity* **9**, 143-150 (1998).

43.   Kato, Z.*, et al.* The structure and binding mode of interleukin-18. *Nat Struct Biol* **10**, 966-971 (2003).

44.   Matsumoto, S.*, et al.* Interleukin-18 activates NF-kappaB in murine T helper type 1 cells. *Biochem*
Biophys Res Commun **234**, 454-457 (1997).

45.   Robinson, D.*, et al.* IGIF does not drive Th1 development but synergizes with IL-12 for interferon-gamma
production and activates IRAK and NFkappaB. *Immunity* **7**, 571-581 (1997).

46.   Kojima, H.*, et al.* An essential role for NF-kappa B in IL-18-induced IFN-gamma expression in KG-1 cells. *J*
Immunol **162**, 5063-5069 (1999).

47.   Morel, J.C., Park, C.C., Kumar, P. & Koch, A.E. Interleukin-18 induces rheumatoid arthritis synovial
fibroblast CXC chemokine production through NFkappaB activation. *Lab Invest* **81**, 1371-1383 (2001).

48.   Miyoshi, K., Obata, K., Kondo, T., Okamura, H. & Noguchi, K. Interleukin-18-mediated
microglia/astrocyte interaction in the spinal cord enhances neuropathic pain processing after nerve injury. *J*
Neurosci **28**, 12775-12787 (2008).

49.   Kaltschmidt, B. & Kaltschmidt, C. NF-kappaB in the nervous system. *Cold Spring Harb Perspect Biol* **1**,
a001271 (2009).

50.   Mattson, M.P. & Meffert, M.K. Roles for NF-kappaB in nerve cell survival, plasticity, and disease. *Cell*
Death Differ **13**, 852-860 (2006).

51.   Frakes, A.E.*, et al.* Microglia induce motor neuron death via the classical NF-kappaB pathway in
amyotrophic lateral sclerosis. *Neuron* **81**, 1009-1023 (2014).

52.   Uranishi, H.*, et al.* Involvement of the pro-oncoprotein TLS (translocated in liposarcoma) in nuclear
factor-kappa B p65-mediated transcription as a coactivator. *J Biol Chem* **276**, 13395-13401 (2001).

53.   Swarup, V.*, et al.* Deregulation of TDP-43 in amyotrophic lateral sclerosis triggers nuclear factor kappaB-
mediated pathogenic pathways. *J Exp Med* **208**, 2429-2447 (2011).

54.   Lee, S., Abecasis, G.R., Boehnke, M. & Lin, X. Rare-variant association analysis: study designs and
statistical tests. *Am J Hum Genet* **95**, 5-23 (2014).

55.   Project MinE Consortium, Van Rheenen, W. & et al. Project MinE: study design and pilot analyses of a
large-scale whole-genome sequencing study in amyotrophic lateral sclerosis. (2017).

56.   Dunckley, T.*, et al.* Whole-genome analysis of sporadic amyotrophic lateral sclerosis. *N Engl J Med* **357**,
775-788 (2007).

57.   Griffiths-Jones, S., Grocock, R.J., van Dongen, S., Bateman, A. & Enright, A.J. miRBase: microRNA
sequences, targets and gene nomenclature. *Nucleic Acids Res* **34**, D140-144 (2006).

58.   Liu, X., Jian, X. & Boerwinkle, E. dbNSFP v2.0: a database of human non-synonymous SNVs and their
functional predictions and annotations. *Hum Mutat* **34**, E2393-2402 (2013).

59.   Lee, S.*, et al.* Optimal unified approach for rare-variant association testing with application to small-
sample case-control whole-exome sequencing studies. *Am J Hum Genet* **91**, 224-237 (2012).

60.   Kenna, K.P.*, et al.* NEK1 variants confer susceptibility to amyotrophic lateral sclerosis. *Nat Genet* **48**,
1037-1042 (2016).

61.   Rosen, D.R.*, et al.* Mutations in Cu/Zn superoxide dismutase gene are associated with familial
amyotrophic lateral sclerosis. *Nature* **362**, 59-62 (1993).

62.   Chio, A.*, et al.* Prevalence of SOD1 mutations in the Italian ALS population. *Neurology* **70**, 533-537
(2008).

63.   van der Spek, R.A.A.*, et al.* The project MinE databrowser: bringing large-scale whole-genome
sequencing in ALS to researchers and the public. *Amyotroph Lateral Scler Frontotemporal Degener* **20**, 432-440
(2019).

64.   O'Leary, N.A*., et al.* Reference sequence (RefSeq) database at NCBI: current status, taxonomic

expansion, and functional annotation. *Nucleic Acids Res* **44**, D733-745 (2016).

65.   Smith, L*., et al.* Establishing the UK DNA Bank for motor neuron disease (MND). *BMC genetics* **16**, 84-84

(2015).

66.   Shi, Y*., et al.* Haploinsufficiency leads to neurodegeneration in C9ORF72 ALS/FTD human induced motor

neurons. *Nature medicine* **24**, 313 (2018).

67.   Haukedal, H. & Freude, K. Implications of Microglia in Amyotrophic Lateral Sclerosis and Frontotemporal

Dementia. *J Mol Biol* **431**, 1818-1829 (2019).

68.   Haenseler, W*., et al.* A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a

Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports* **8**, 1727-1742

(2017).

69.   Peng, S.S., Chen, C.Y., Xu, N. & Shyu, A.B. RNA stabilization by the AU-rich element binding protein, HuR,

an ELAV protein. *EMBO J* **17**, 3461-3470 (1998).

70.   Fan, X.C. & Steitz, J.A. Overexpression of HuR, a nuclear-cytoplasmic shuttling protein, increases the in

vivo stability of ARE-containing mRNAs. *EMBO J* **17**, 3448-3460 (1998).

71.   Stellos, K*., et al.* Adenosine-to-inosine RNA editing controls cathepsin S expression in atherosclerosis by

enabling HuR-mediated post-transcriptional regulation. *Nat Med* **22**, 1140-1150 (2016).

72.   Brennan, C.M. & Steitz, J.A. HuR and mRNA stability. *Cell Mol Life Sci* **58**, 266-277 (2001).

73.   Garcia-Dominguez, D.J., Morello, D., Cisneros, E., Kontoyiannis, D.L. & Frade, J.M. Stabilization of Dll1

mRNA by Elavl1/HuR in neuroepithelial cells undergoing mitosis. *Mol Biol Cell* **22**, 1227-1239 (2011).

74.   Rothamel, K*., et al.* ELAVL1 primarily couples mRNA stability with the 3' UTRs of interferon-stimulated

genes. *Cell Rep* **35**, 109178 (2021).

75.   Mukherjee, N*., et al.* Integrative regulatory mapping indicates that the RNA-binding protein HuR couples

pre-mRNA processing and mRNA stability. *Mol Cell* **43**, 327-339 (2011).

76.   Fernandopulle, M.S*., et al.* Transcription Factor-Mediated Differentiation of Human iPSCs into Neurons.

Curr Protoc Cell Biol **79**, e51 (2018).

77.   Christian, F., Smith, E.L. & Carmody, R.J. The Regulation of NF-kappaB Subunits by Phosphorylation. *Cells*

5 (2016).

78.   Zhong, H., May, M.J., Jimi, E. & Ghosh, S. The phosphorylation status of nuclear NF-kappa B determines

its association with CBP/p300 or HDAC-1. *Mol Cell* **9**, 625-636 (2002).

79.   Zhong, H., Voll, R.E. & Ghosh, S. Phosphorylation of NF-kappa B p65 by PKA stimulates transcriptional

activity by promoting a novel bivalent interaction with the coactivator CBP/p300. *Mol Cell* **1**, 661-671 (1998).

80. Oeckinghaus, A. & Ghosh, S. The NF-kappaB family of transcription factors and its regulation. *Cold Spring*

Harb Perspect Biol **1**, a000034 (2009).

81. Waelchli, R*., et al.* Design and preparation of 2-benzamido-pyrimidines as inhibitors of IKK. *Bioorg Med*

Chem Lett **16**, 108-112 (2006).

82. Ayers, K.L*., et al.* A loss of function variant in CASP7 protects against Alzheimer's disease in homozygous

APOE epsilon4 allele carriers. *BMC Genomics* **17 Suppl 2**, 445 (2016).

83. Benitez, B.A*., et al.* Missense variant in TREML2 protects against Alzheimer's disease. *Neurobiol Aging*

35, 1510 e1519-1526 (2014).

84. Jonsson, T*., et al.* A mutation in APP protects against Alzheimer's disease and age-related cognitive

decline. *Nature* **488**, 96-99 (2012).

85. Sims, R*., et al.* Rare coding variants in PLCG2, ABI3, and TREM2 implicate microglial-mediated innate

immunity in Alzheimer's disease. *Nat Genet* (2017).

86. Landers, J.E*., et al.* Reduced expression of the Kinesin-Associated Protein 3 (KIFAP3) gene increases

survival in sporadic amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A* **106**, 9004-9009 (2009).

87. Farhan, S.M.K*., et al.* Exome sequencing in amyotrophic lateral sclerosis implicates a novel gene,

DNAJC7, encoding a heat-shock protein. *Nat Neurosci* **22**, 1966-1974 (2019).

88. Lambrechts, D*., et al.* VEGF is a modifier of amyotrophic lateral sclerosis in mice and humans and

protects motoneurons against ischemic death. *Nat Genet* **34**, 383-394 (2003).

89. Beers, D.R. & Appel, S.H. Immune dysregulation in amyotrophic lateral sclerosis: Mechanisms and

emerging therapies. *The Lancet Neurology* **18**, 211-220 (2019).

90. Zhan, X., Hu, Y., Li, B., Abecasis, G.R. & Liu, D.J. RVTESTS: an efficient and comprehensive tool for rare

variant association analysis using sequence data. *Bioinformatics* **32**, 1423-1426 (2016).

91. Raczy, C*., et al.* Isaac: ultra-fast whole-genome secondary analysis on Illumina sequencing platforms.

Bioinformatics **29**, 2041-2043 (2013).

92. Tyner, C*., et al.* The UCSC Genome Browser database: 2017 update. *Nucleic Acids Res* **45**, D626-D634

(2017).

93. Danecek, P*., et al.* The variant call format and VCFtools. *Bioinformatics* **27**, 2156-2158 (2011).

94. Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants from high-

throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010).

95. Agarwal, V., Bell, G.W., Nam, J.W. & Bartel, D.P. Predicting effective microRNA target sites in mammalian

mRNAs. *Elife* **4** (2015).

96.   Schildge, S., Bohrer, C., Beck, K. & Schachtrup, C. Isolation and culture of mouse cortical astrocytes. *J Vis*
Exp (2013).
97.   Hsu, P.D.*, et al.* DNA targeting specificity of RNA-guided Cas9 nucleases. *Nat Biotechnol* **31**, 827-832
(2013).
98.   Doench, J.G.*, et al.* Optimized sgRNA design to maximize activity and minimize off-target effects of
CRISPR-Cas9. *Nat Biotechnol* **34**, 184-191 (2016).
99.   Xu, H.*, et al.* Sequence determinants of improved CRISPR sgRNA design. *Genome Res* **25**, 1147-1157
(2015).
100.   Chari, R., Mali, P., Moosburner, M. & Church, G.M. Unraveling CRISPR-Cas9 genome engineering
parameters via a library-on-library approach. *Nat Methods* **12**, 823-826 (2015).
101.   Erijman, A., Dantes, A., Bernheim, R., Shifman, J.M. & Peleg, Y. Transfer-PCR (TPCR): a highway for DNA
cloning and protein engineering. *J Struct Biol* **175**, 171-177 (2011).
102.   Peleg, Y. & Unger, T. Application of the Restriction-Free (RF) cloning for multicomponents assembly.
Methods Mol Biol **1116**, 73-87 (2014).
103.   Kohen, R.*, et al.* UTAP: User-friendly Transcriptome Analysis Pipeline. *BMC Bioinformatics* **20**, 154
(2019).
104.   Liao, Y., Wang, J., Jaehnig, E.J., Shi, Z. & Zhang, B. WebGestalt 2019: gene set analysis toolkit with
revamped UIs and APIs. *Nucleic Acids Res* **47**, W199-W205 (2019).
105.   Lorenz, R.*, et al.* ViennaRNA Package 2.0. *Algorithms Mol Biol* **6**, 26 (2011).
106.   Krach, F.*, et al.* Transcriptome–pathology correlation identifies interplay between TDP-43 and the
expression of its kinase CK1E in sporadic ALS. *Acta neuropathologica* **136**, 405-423 (2018).
107.   Thompson, L. iMN (Exp 2)—ALS. *SMA and control (unaffected) iMN cell lines differentiated from iPS cell*
lines using a long differentiation protocol—RNA-seq., LINCS (collection) (2017).

1294