Nitrogen fixation in mesoscale eddies of the North Pacific Subtropical Gyre: patterns and mechanisms

Mathilde Dugenne[a,b*†], Mary R. Gradoville[c*†],

Matthew J. Church[d], Benedetto Barone[a,b], Samuel T. Wilson[a,b], Uri Sheyn[a,b], Matthew J. Harke[e,f], Karin M. Björkman[a,b], Nicholas J. Hawco[a], Annette M. Hynes[g], François Ribalet[g], David M. Karl[a,b], Edward F. Delong[a,b], Sonya T. Dyhrman[e,h], E. Virginia Armbrust[g], Seth John[i], John M. Eppley[a,b], Katie Harding[c], Brittany Stewart[c,i], Ana M. Cabello[c,j], Kendra A. Turk-Kubo[c], Mathieu Caffin[a,b], Angelicque E. White[a,b], Jonathan P. Zehr[c]

[a] Department of Oceanography, University of Hawai'i at Mānoa, Honolulu, HI, USA
[b] Daniel K. Inouye Center for Microbial Oceanography: Research and Education, University of Hawai'i at Mānoa, Honolulu, HI, USA
[c] Ocean Sciences Department, University of California Santa Cruz, Santa Cruz, CA, USA
[d] Flathead Lake Biological Station, University of Montana, Polson, MT, USA
[e] Lamont-Doherty Earth Observatory, Columbia University, Palisades, NY, USA
[f] Present Address: Gloucester Marine Genomics Institute, Gloucester, MA, USA
[g] School of Oceanography, University of Washington, Seattle, WA, USA
[h] Department of Earth and Environmental Sciences, Columbia University, New York, NY, USA
[i] Department of Earth Sciences, University of Southern California, Los Angeles, CA, USA
[j] Present address: Centro Oceanográfico de Málaga, Instituto Español de Oceanografía, Fuengirola, Málaga 29001, Spain

*Contributed equally

†Corresponding authors: mdugenne@hotmail.com, rosiegradoville@gmail.com

Running title: Nitrogen fixation in mesoscale eddies

Keywords: Cyanobacterial diazotrophs / nitrogen fixation / mesoscale eddies / frontal accumulation / P-limitation / grazing rate estimates

1. Abstract

Mesoscale eddies have been shown to support elevated dinitrogen (N_2) fixation rates

(NFRs) and abundances of N_2-fixing microorganisms (diazotrophs), but the mechanisms

underlying these observations are not well understood. We explored relationships among NFRs

and cyanobacterial diazotroph abundances in eddy pairs of opposite polarity sampled in the

North Pacific Subtropical Gyre and compared our observations to seasonal trends from the

Hawaii Ocean Time-series (HOT) program. Consistent with previous reports, we found that

NFRs were anomalously high for this region (up to 3.7-fold above previous monthly HOT

observations) in the centers of the sampled anticyclones, coinciding with elevated abundances of

Crocosphaera in the summertime. We then coupled our field-based observations, together with

transcriptomic analyses of nutrient stress marker genes and ecological models, to evaluate

potential mechanisms controlling diazotroph abundance and activity at the mesoscale.

Specifically, we evaluated the role of biological (via estimates of growth and grazing rates) and

physical controls on populations of *Crocosphaera*, *Trichodesmium,* and diatom symbionts. Our

results suggest that increased *Crocosphaera* abundances associated with summertime

anticyclones resulted from the alleviation of phosphate limitation, allowing cells to grow at rates

exceeding grazing losses. In contrast, distributions of larger, buoyant taxa (*Trichodesmium* and

diatom symbionts) appeared less affected by eddy-driven biological controls. Instead, they

appeared driven by physical dynamics along frontal boundaries that separate cyclonic and

anticyclonic eddies. Together, the interplay of eddy-specific changes in bottom-up control, top-

down control, and the physical accumulation of cells likely explains the elevated diazotroph

abundances and NFRs associated with anticyclones and eddy fronts.

2. Introduction

In the oligotrophic North Pacific Subtropical Gyre (NPSG), the supply of growth-limiting

nutrients influences rates of photosynthesis in the sunlit surface ocean, determining the

magnitude and variability of trophic transfer and the export of carbon (C) from the euphotic zone

(Karl et al., 1996). Microorganisms with the ability to fix dinitrogen (N_2) into bioavailable

nitrogen (N), termed diazotrophs, introduce newly fixed N to otherwise N-limited phytoplankton

(Karl et al., 2002). At Station ALOHA (A Long-term Oligotrophic Habitat Assessment, 22.75°N,

158.00°W), the field site of the Hawaii Ocean Time-series (HOT) program, N_2 fixation fuels up

to half of new production and N export (Böttjer et al., 2017; Karl et al., 1997). Most N_2 fixation

is controlled by unicellular diazotrophs (typically smaller than 10 μm), including the

cyanobacteria UCYN-A (symbionts of haptophytes), UCYN-B (*Crocosphaera*), and UCYN-C

(*Cyanothece*-like organisms) (Church et al., 2009; Zehr et al., 2001). Other, larger cyanobacterial

diazotrophs including filamentous *Trichodesmium* and diatom-diazotroph associations (DDAs)

with the symbiotic, heterocyst-forming *Richelia* and *Calothrix* can also form episodic blooms

and contribute appreciably to N_2 fixation. Finally, non-cyanobacterial diazotrophs are also

present in the NPSG (Church, Short, et al., 2005; Farnelid et al., 2011), but the contribution of

these organisms to N_2 fixation is uncertain (Moisander et al., 2017).

N_2 fixation rates (NFRs) at Station ALOHA are highly variable, ranging from ~0.3 to 21

nmol N L^{-1} d^{-1} in near-surface waters (Böttjer et al., 2017). Some of this variability is seasonal:

NFRs generally peak during the late summer (Böttjer et al., 2017), when an annual pulse of

particle export to the deep ocean has also been putatively linked to blooms of DDAs (Karl et al.,

2012; Poff et al., 2021). However, several studies have also highlighted patchiness in both NFRs

and diazotroph abundances over relatively small temporal (<2 days) and spatial (<30 km) scales

(Gradoville et al., 2020; Robidart et al., 2014), suggesting that submesoscale and mesoscale

processes may influence diazotroph distributions in the NPSG. Such physical disturbances of the

upper pelagic zone are regular features of this habitat (Rii et al., 2021); for example, over a 23-

year period of near-monthly sampling by the HOT program at Station ALOHA, 31% of the

sampling occasions coincided with the presence of a trackable mesoscale eddy (Barone et al.,

2019).

Through uplift or depression of the pycnocline, eddies perturb microbial populations by

altering nutrient and light fields. As a result, mesoscale eddies, especially cyclonic eddies that

drive upward displacement of isopycnal surfaces (together with associated nutrients) across the

upper ocean light gradient, have been linked to elevated phytoplankton biomass and high rates of

primary production in the open ocean (McGillicuddy Jr & Robinson, 1997). There is growing

evidence that diazotrophic community structure and NFRs are also modified by mesoscale

eddies. High NFRs have been observed in anticyclonic eddies in many ocean regions, including

the South Pacific Ocean, Indian Ocean, South China Sea, and Mediterranean Sea (Holl et al.,

2007; Liu et al., 2020; Löscher et al., 2016; Rahav et al., 2013). At Station ALOHA, the highest

NFRs have been observed under conditions of elevated sea surface height (characteristic of

anticyclonic eddies) in summer months (Böttjer et al., 2017; Church et al., 2009). Elevated

abundances of *Trichodesmium*, DDAs, and *Crocosphaera* have been reported associated with

anticyclonic eddies in both the North Atlantic and North Pacific Oceans (Cheung et al., 2020;

Davis & McGillicuddy, 2006; Fong et al., 2008; Olson et al., 2015; Taboada et al., 2010; Wilson

et al., 2017).

These studies have highlighted multiple processes that could influence N_2 fixation in

eddies of different polarities. For example, elevated NFRs and *Crocosphaera* abundances were

linked to severe N-depletion in the surface waters of anticyclones, presumably driven by the

depression of isopycnal surfaces (Liu et al., 2020; Wilson et al., 2017). In addition, although

poorly characterized, high rates of grazing on diazotrophs and their hosts have been reported

within mesoscale eddies of both polarities (Dugenne et al., 2020; Landry et al., 2008; Wilson et

al., 2017). Finally, the physical movement of water, both vertically and horizontally, due to

mesoscale circulation, can lead to the physical accumulation of buoyant diazotrophs (e.g.

Trichodesmium) (Guidi et al., 2012; Olson et al., 2015). All of these mechanisms are modulated

by specific life traits (e.g. buoyancy, size, symbiosis) and potential adaptations of individual

taxa, suggesting that understanding diazotrophic diversity and activity within eddies is

prerequisite to identifying the mechanisms driving variability in bulk NFRs. Effectively,

mesoscale and sub-mesoscale features can impact both bottom-up and top-down processes

controlling diazotroph diversity and activity. The relative influence of these controls is important

in determining diazotroph abundance as well as the magnitude of new N delivered to the system.

In this study, we examined cyanobacterial diazotroph abundances and NFRs associated

with two pairs of mesoscale eddies (cyclones and anticyclones) sampled in the NPSG and

compared these data to trends from the HOT program for regional and seasonal context. We used

these observations, together with environmental data, metatranscriptomes, and ecological

models, to evaluate the importance of bottom-up, top-down, and physical control mechanisms in

driving the mesoscale distribution and activity of diazotroph populations.

3.   Material and Methods

3.1 Characterization and sampling of eddy pairs

In 2017 and 2018, two field expeditions sampled mesoscale eddies in the region north of

the Hawaiian Islands. Observations took place 26 June–15 July 2017 during the Microbial

Ecology of the Surface Ocean-Simons Collaboration on Ocean Processes and Ecology cruise

(MESO-SCOPE, KM1709, *R/V Kilo Moana*) and 27 March–10 April 2018 during the Hawaiian

Eddy Experiment (FK180310, *R/V Falkor*). Prior to each cruise, cyclonic and anticyclonic eddies

were identified via satellite-derived sea level anomalies (SLA) distributed by the Copernicus

Marine Service (marine.copernicus.eu/). Daily SLA maps (Fig. 1) were corrected for the multi-

year trend and seasonal cycle following Barone et al. (2019). Eddies were tracked back in time to

determine the stage, age, and average SLA during the sampling period (Supporting Information

S1 and Table S1).

Figure 1: Sampling locations for *nifH* gene abundances (purple circles), hydrographical stations
(blue circles), a *Trichodesmium* bloom (yellow star), and N_2 fixation rates (triangles) across each
pair of eddies during the 2017 (A) and 2018 (B) cruises. Note that surface diazotroph cell
abundances were measured by autonomous flow cytometry at high spatial resolution along each
cruise transect. Sampled cyclones are indicated by negative sea level anomalies (-0.1 m blue
contour) and anticyclones by positive sea level anomalies (0.1 m red contour). Scale bars: 100
km.

Both cruises included an eddy mapping phase, where sampling occurred along a transect

spanning one eddy to the other to characterize the water column, quantify diazotrophs via *nifH*

gene abundances and automated imaging flow cytometry cell counts, and measure NFRs. They

also included time-resolved, Lagrangian surveys that sampled along the drift trajectories of

Surface Velocity Program (Pacific Gyre) drifters in one or both eddy centers (drogue centered at

15 m), with continuous imaging of near surface plankton communities to assess high resolution

temporal dynamics.

Photosynthetically active radiation (PAR) at the sea-surface was measured using a

shipboard cosine collector LI-190 (LI-COR Environmental, Lincoln, NE, USA). Downwelling

PAR was measured at 1 m-intervals using a free-falling optical profiler (Satlantic HyperPro, Sea-

Bird Scientific, Bellevue, WA, USA), and corrected by a factor of 1.2 to convert downwelling

irradiance into scalar irradiance (Letelier et al., 2017; Wozniak et al., 2003). Both sea surface and

downwelling PAR measurements were used compute the fraction of incident PAR (%) at

discrete depths.

A Conductivity, Temperature, and Depth (CTD, Sea-Bird Scientific) sensor attached to a

rosette sampler was used to measure depth profiles of temperature, salinity, and derived potential

density. Mixed layer depth (MLD) was calculated using a 0.03 potential density offset relative to

10 m (de Boyer Montégut et al., 2004). Seawater samples collected from discrete depths using 24

$\times$ 10 L Niskin® bottles mounted to the rosette frame were used to measure soluble reactive

phosphorus (hereafter phosphate, PO_4^{3-}) concentrations (precision of $\pm$ 1 nmol L^{-1} and detection

limit of 3 nmol L^{-1}) (Karl & Tien, 1992) and nitrate + nitrite ($NO_3^-+NO_2^-$) concentrations

(detection limit of 1 nmol L^{-1}) (Foreman et al., 2016).

Dissolved Fe (dFe) for the 2017 cruise was collected using a trace metal clean sampling

rosette with external spring bottles (Ocean Test Equipment). For the 2018 cruise, samples for

dFe were collected in Go-Flo® bottles mounted on a non-metallic line and triggered with a

PTFE messenger. Dissolved iron concentrations were determined by inductively coupled plasma

mass spectrometry (ICP-MS) using isotope dilution (Hawco et al., 2021; Pinedo-González et al.,

2020). The accuracy of these measurements was confirmed by analyzing reference seawater

samples, GS (0.55 ± 0.01 nM Fe, n = 3) and GSP (0.19 ± 0.03 nM Fe, n= 3), provided by the

GEOTRACES community (www.geotraces.org/standards-and-reference-materials).

3.2 N_2 fixation rates

NFRs were measured on all cruises using a modification of the $^{15}N_2$ uptake method of

Montoya et al. (1996), in which the $^{15}N_2$ tracer was added as $^{15}N_2$-enriched seawater to avoid

incomplete dissolution of $^{15}N_2$ gas (Mohr et al., 2010; Wilson et al., 2012). Prior to each cruise,

surface seawater was collected from Station ALOHA, filtered through a 0.2 μm in-line filter

(AcroPak 1000 capsule, Pall Corporation, Port Washington, NY, USA) and transported in the

dark to the laboratory. There, the seawater was degassed, injected with ~13 mL $^{15}N_2$ gas (99

atom %, Cambridge Scientific, Watertown, MA, USA) per L seawater, and manually agitated to

dissolve the gas bubble, as described by Wilson et al. (2012). The resulting $^{15}N_2$-enriched

seawater was dispensed into crimp-sealed, glass serum bottles, stored at 4°C for less than one

week, and brought to sea. To initiate incubations, 100 mL of seawater was withdrawn from each

full 4.4 L incubation bottle and replaced with 100 mL of $^{15}N_2$-enriched seawater. The $^{15}N/^{14}N$

ratio of each batch of $^{15}N_2$-enriched seawater was measured using membrane inlet mass

spectrometry according to Ferrón et al. (2016). These values were used to calculate the initial

atom % of $^{15}N_2$ for each incubation. A recent study by White et al. (2020) indicates that

measuring the $^{15}N_2$ atom % of the enriched seawater rather than the final inoculum may lead to

inaccurate initial $^{15}N_2$ atom % values, which for the enriched seawater preparation method used

for the eddy and HOT observations reported in this study could have led to a <20%

underestimation of final calculated rates.

NFRs were measured using free-drifting, in situ arrays deployed at eddy centers (Fig. 1).

Seawater was collected pre-dawn from 5, 25, 45, 75, 100, and 125 m depths using the CTD

rosette Niskin® sampling bottles. Seawater was subsampled into triplicate 4.4 L acid-washed,

seawater-rinsed polycarbonate bottles which were spiked with $^{15}N_2$ tracer (see above), attached

to the free-floating arrays, and incubated for 24 h at the depth from which they had been

collected. Incubations were terminated by filtering (via a peristaltic pump or vacuum filtration)

through pre-combusted (5 h at 450°C) 25 mm glass fiber filters (GF/F, Whatman, Maidstone,

UK). Additional 4.4 L seawater samples were collected from each sampling depth for $\delta^{15}N$

natural abundance measurements; these samples were immediately filtered at the start of each

incubation. All filters were frozen at -20°C or -80°C and transported to the laboratory, where

they were dried at 60°C overnight and pelleted into tin capsules (Costech Analytical Tech Inc,

Valencia, CA, USA). Concentrations and isotopic composition ($\delta^{15}N$) of particulate N were

analyzed via continuous-flow isotope ratio mass spectrometry using a Carlo-Erba EA NC2500

coupled with Thermo Finnigan DeltaPlus XP at the University of Hawai'i Biogeochemical

Stable Isotope Facility. NFRs and detection limits were calculated according to Montoya et al.

(1996) (Table S2).

Time-series NFRs measured at Station ALOHA are also presented to provide context for

the two eddy cruises. A subset of the time-series (2012-2014) has previously been published by

Böttjer et al. (2017); for this study, we extend that time series through 2019. These extended

time-series measurements followed procedures identical to those described in Böttjer et al.

(2017).

3.3 Diazotroph abundances

3.3.1. DNA extraction and nifH *gene quantification*

The *nifH* genes of seven major cyanobacterial diazotroph groups were quantified using

digital droplet polymerase chain reaction (ddPCR). During the 2018 cruise, seawater for

subsequent ddPCR analyses was collected from depth profiles at the same locations sampled for

$^{15}N_2$ incubations. During the 2017 cruise, seawater for ddPCR was collected from depth profiles

within each eddy, but from separate locations than those used for the $^{15}N_2$ arrays (Fig. 1). For

both cruises, seawater samples were collected from a rosette sampler and subsampled into

duplicate (2018) or single (2017) 4.4 L acid-washed, milliQ-rinsed polycarbonate bottles. A

peristaltic pump was used to filter 4 L of seawater through 0.2 µm pore-size Supor® membranes

(Pall), which were placed into microcentrifuge tubes containing mixtures of 0.1- and 0.5-mm

glass beads (Biospec Products, Bartlesville, OK, USA), flash-frozen in liquid N_2, and stored at -

80°C until analysis.

DNA was extracted from filters with a QIAcube instrument (Qiagen, Venlo, Netherlands)

using the DNeasy Plant Mini Kit. Extractions followed the manufacturer's protocol with

additional steps of three flash freeze/thaw cycles, bead-beating for two min, and a Proteinase K

treatment, as described by Moisander et al. (2008). The final DNA elution volume was 100 μL.

DNA extracts were stored at -20°C prior to ddPCR. Duplicate ddPCR reactions were performed

for each sample using primers and probes targeted the following groups: *Trichodesmium*

(Church, Jenkins, et al., 2005), UCYN-A (targeting the small UCYN-A1 sublineage, Church,

Jenkins, et al., 2005), *Crocosphaera* (UCYN-B, Moisander et al., 2010), *Cyanothece*-like

organisms, (UCYN-C, Foster et al., 2007), *Richelia* associated with *Rhizosolenia* (Het-1, Church,

Short, et al., 2005), *Richelia* associated with *Hemiaulus* (Het-2, Foster et al., 2007), and

Calothrix associated with *Chaetoceros* (Het-3, Foster et al., 2007). Descriptions of ddPCR

reaction setup, droplet generation, thermocycling, thresholding, and detection limit calculations

are provided in Gradoville et al. (2021).

For temporal context, we also report *nifH* gene abundances of UCYN-A, UCYN-B,

Trichodesmium, Het-1, Het-2, and Het-3, measured at Station ALOHA between 2004 and 2017

(dataset doi: 10.5281/zenodo.4477269). For these measurements, *nifH* gene abundances were

determined using quantitative PCR (qPCR); details on sample collection, DNA extraction, and

qPCR are provided in Church et al. (2009). All qPCR assays except for Het-2 and Het-3 used the

same primer/probe sets as the corresponding ddPCR assays for the eddy cruises. Het-2 was

quantified using the primer/probes sets of Church et al. (2008), while Het-3 used the same

forward and reverse primers used for ddPCR, but the probe sequence contained an extra adenine

on the 3' end.

3.3.2. Diazotroph cell and filament enumeration by autonomous flow cytometry

Concentrations of large (>4 μm in diameter or length, imposed by the size detection limit

of the instrument) free-living and filamentous diazotrophs, including *Trichodesmium* (single

filaments), *Richelia* (free and symbiotic heterocyst-containing filaments), *Calothrix* (unattached

and epiphytic heterocyst-containing filaments), and large-size *Crocosphaera-like* organisms

(free-living, hereafter referred as *Crocosphaera*), were estimated by automated imaging flow

cytometry equipped with a 635 nm red laser (Imaging FlowCytobot [IFCb], McLane, East

Falmouth, MA, USA) during the 2017 and 2018 cruises. Five mL samples were collected every

20 min from the ship's uncontaminated underway system (~7 m depth). Discrete bucket samples

from a visual *Trichodesmium* bloom were also analyzed on the IFCb in 2017 (Fig. 1). During the

analysis, individual images of single or colonial cells were collected along with their optical

fingerprints: light scattering and red fluorescence (680±30 nm), emitted by chlorophyll and

phycocyanin pigments. The latter is characteristic of N_2-fixing cyanobacteria (Boatman et al.,

2018; Webb et al., 2009; Zeev et al., 2008). A training set of images was used to classify

organisms to the genera level based on morphological traits, as described in Dugenne et al.

(2020). The output of the random forest classifier (Sosik & Olson, 2007) was corrected by

manually annotating misclassified images to provide accurate estimates of cell and filament

concentration of individual genera. To compare the cell counts of rare filamentous diazotrophs

with *nifH* abundances, we binned consecutive samples in 2-h-intervals (in order to integrate a

larger volume and lower the detection limit) and manually estimated the number of cells per

filament.

Small-size *Crocosphaera* (2-4 μm) were enumerated using an autonomous SeaFlow flow

cytometer (Ribalet et al., 2019). Briefly, the SeaFlow was used to continuously sample surface

seawater from the ship's uncontaminated underway system, acquiring optical measurements of

light scatter, red fluorescence, and orange fluorescence (characteristic of the phycoerythrin-

containing *Crocosphaera*) at 3 min intervals. The population of small-size *Crocosphaera* was

gated over time to estimate its abundance during time points close (less than 2 h) to collection

times of the discrete *nifH* gene abundance samples.

3.4. Metatranscriptomic analysis of stress-marker genes

For insight into possible diazotroph growth limitation by P and Fe in eddy centers, we

examined transcriptional patterns in P- and Fe-stress marker genes from samples collected during

the 2017 cruise. Metatranscriptome samples were collected within the mixed layer (15 m) and

the deep chlorophyll maximum (DCM, located at depths of 106±5 m and 123±5 m in the cyclone

and anticyclone, respectively) at ~4 h intervals for three days during the Lagrangian sampling of

the cyclonic and anticyclonic eddies (n=18 for each eddy). Seawater samples were collected

using the rosette sampler and seawater (1 L) was filtered through 25 mm diameter, 0.2 μm

Supor® membranes (Pall) housed in Swinnex™ units (MilliporeSigma) using a peristaltic pump.

Filtration times were limited to 15-20 min. Immediately following filtration, filters were placed

in RNALater (Invitrogen, Waltham, MA, USA) and stored at -80°C until processed.

Sample and data processing were performed using the methods described by Gifford et

al. (2016) and Wilson et al. (2017). Briefly, sample processing steps included RNA extraction,

spiking with internal controls, cDNA synthesis, and sequencing using Illumina NextSeq 500 v2.

Approximately two million 150 bp paired-end reads were produced for each sample; raw

sequences are available on NCBI SRA under project number PRJNA596510. Reads were

screened for quality, trimmed, and assembled after removing sequencing primers. Genes were

mapped to the ALOHA 2.0 gene catalog (Luo et al., 2020), and annotated using the Genome

Taxonomy Database (GTDB, Parks et al., 2018). Raw sequence reads are available on NCBI

SRA under project number PRJNA596510. Transcript counts were normalized to internal

standards to account for methodological biases among samples, including library preparation and

sequencing, and also normalized to the volume of water filtered for each sample as previously

described (Wilson et al., 2017). Values above detection limits were further normalized to the

total sample expression sum within each genus-level annotation. This step was required in order

to compare eddy differential gene expression based on cellular regulation rather than based on

the total number of cells. Transcripts of P- and Fe-stress marker genes (genes with higher

expression levels under low P and Fe concentrations, as reviewed by Stenegren (2020) and Snow

et al. (2015) were extracted from the dataset according to functional annotations assigned by

comparison to the Kyoto Encyclopedia of Genes and Genomes.

Several genes exhibited diel expression patterns. To account for the expression

periodicity and evaluate the differences in gene transcript abundances between eddies, we

normalized the abundance of transcripts above detection limits within each eddy to the overall

average of each gene transcript from the entire time-series (time points with undetected gene

transcripts were excluded from this analysis). This step facilitated comparisons of different gene

transcripts having large variations in their diel baseline and maximum transcript expression

levels.

The statistical significance of the relative change in expression was tested using a

Kruskal-Wallis test (R function 'kruskal.test'). All gene transcripts from UCYN-A and

Calothrix, as well as the transcripts of several genes from *Richelia* and *Trichodesmium*, were

excluded from analyses as their abundances always fell below the detection limit in at least one

eddy, especially in samples collected at the DCM.

3.5. Diazotroph populations models

3.5.1. Bottom-up control on diazotroph populations

We used an ecophysiological model (Follett et al., 2018; McGillicuddy Jr, 2014; Stukel et

al., 2014) to evaluate how measured abiotic factors (nutrient concentrations, light, and

temperature) may have influenced diazotroph intrinsic growth rates (μ; per day) across the pairs

of eddies. Predictions are based on the ecophysiological parameters of cyanobacterial diazotroph

taxa derived from laboratory culture studies examining the effect of temperature (T), average

instantaneous scalar PAR (E), PO_4^{3-}, and dFe concentrations on growth rates (Table S3). Model

assumptions and caveats are described in Supporting Information S2.

We integrated the effects of abiotic factors on population growth rates by scaling the

theoretical maximum growth rate of each taxon (μ_{max}) to the relative change in growth rate (f_μ)

predicted by the main limiting factor, according to Liebig's law of the minimum:

$(Eq.\ 1)\ \mu(T, E,\ PO_4^{3-}, dFe) = \mu_{max} \times min(f_\mu(T), f_\mu(E), f_\mu(PO_4^{3-}), f_\mu(dFe))$

where f_μ designate unitless limiting functions (Eq. 2-3, 5) representing the changes of diazotroph

growth rates relative to their maximum. The change in relative growth rate as a function of T

followed Eq. 2:

$$(Eq.\ 2)\ f_\mu(T) = Q_{10}^{\frac{T - T_{opt}}{10}}$$

with T_{opt} (°C) the optimal temperature for which f_μ (T_{opt})=1, and Q_{10}, the Arrhenius coefficient

corresponding to the change of growth rate over a 10°C temperature increase (reported in Table

S3 for individual diazotroph taxa).

The change in relative growth rate as a function of E (μmol quanta m^{-2} s^{-1}) was predicted

using a modification of the hyperbolic tangent function described in Jassby and Platt (1976):

$$(Eq.\ 3)\ f_\mu(E) = tanh\left(\frac{\alpha(E - E_c)}{\mu_{max}}\right)$$

with E_c (μmol quanta m^{-2} s^{-1}), the compensation light intensity. The light saturation parameter, E_k

(μmol quanta m^{-2} s^{-1}), is then derived using the maximum growth rate, μ_{max}, reported in Table S3

and the initial slope of μ-E curve presented in the literature, α, as:

$$(Eq.\ 4)\ E_k = \frac{\mu_{max}}{\alpha} + E_c$$

Finally, Michaelis-Menten kinetics were used to express the change in relative growth

rate as a function of both PO_4^{3-} and dFe concentration (nmol L^{-1}):

$$(Eq.\ 5)\ f_\mu(PO_4^{3-}) = \frac{PO_4^{3-}}{PO_4^{3-} + k_{PO_4^{3-}}}, \qquad f_\mu(dFe) = \frac{dFe}{dFe + k_{Fe}}$$

All models were fitted to published culture data using the nonlinear least-squares R

function 'nls'. We tested the differences in intrinsic growth rate estimates between eddies of

opposite polarity by generalized linear regression, assuming a quasi-Binomial distribution

(which is typically assumed for biological rates), weighted by the uncertainties of the estimates

with the R function glm(μ~Eddy, family=quasibinomial(link = 'logit'),weights=1/ σ^2(μ(T,E,

PO_4^{3-},dFe))), where the explanatory variable 'Eddy' corresponds to eddy polarity.

3.5.2. Predation control on diazotroph populations

We explored how mesoscale eddies affected estimates of diazotroph grazing rates using a

standard predator-prey model. The predation model relied on co-occurrence interactions between

a given diazotroph (i.e. *Crocosphaera, Richelia, Calothrix, Trichodesmium*) and potential

predators, based on abundance data from the IFCb (Dugenne et al., 2020). More specifically, we

leveraged the IFCb images to (1) determine significant interactions within a generalized

predator-prey model (Eq. 6) and (2) visualize ingested diazotrophic prey. We relied on the

temporal dynamics of diazotrophic prey (x_i, cells or filaments L^{-1}) and putative predators (x_j,

cells L^{-1}), measured in the eddy centers throughout the Lagrangian surveys (hence assuming that

population dynamics are only driven by growth and loss rates), to identify significant interactions

(a_{ij}):

$$(Eq.6) \quad \frac{dx_i}{dt} = x_i(r_i + \sum_j a_{ij}x_j)$$

The term r comprised a linear intrinsic growth rate and a linear loss rate due to processes not

explicitly accounted for with the biological interactions tested (e.g. viral lysis, programmed cell

death, additional grazers not imaged by the IFCb, vertical migration, mixed layer entrainment,

and/or sinking). A similar model has previously been used to assess the short-term variability of

grazing rate estimates (Dugenne et al., 2020); however, here we simplified this model to have

fixed parameters in each eddy center (e.g. throughout each Lagrangian survey) in order to

compare grazing rates between eddies.

In the current model, the strength of the interaction between populations i and j is

determined by the coefficient a_{ij} and by population abundances; a_{ij} can be interpreted as different

biological rates depending on the type of the interaction. When coefficients a_{ij} and a_{ji} are

negative and positive respectively, j grows at i's expense, as would a putative grazer of the

diazotroph i. All pairs of interaction coefficients were estimated by a linear multi-regression

model fitted to the changes in population abundances throughout the Lagrangian sampling period

within individual eddies as follows:

346
$$(Eq.\,7)\ \frac{\Delta \ln(x_i)}{\Delta t} = y = r_i + \sum_j a_{ij} x_j$$

with Δt, a 1 h interval based on the sensitivity analysis presented in Supporting Information S3,

yielding an average uncertainty, calculated as the ratio of the standard error over the estimate for

the regression coefficient, of $12\pm6\%$ of $a_{ij,}$ the linear coefficient for the interaction with

population j, and $r_{i,}$ the intercept of the linear regression. Since the regression included a large

number of coefficients, accommodating 103 populations in 2017 and 89 populations in 2018, we

used a regularized Lasso regression implemented in the 'glmnet' R package to penalize non-

significant interactions (Tibshirani et al., 2012). We also looked for visual evidence of

diazotrophic prey ingestion using the IFCb images from 2017 and 2018 to confirm the significant

interaction between diazotrophs and their putative grazers. Significant coefficients were sorted to

determine the importance of individual protists based on g_j (d^{-1}), the product of coefficient a_{ij} and

the abundance of grazer population j. The overall rates of grazing on a diazotroph taxon, g (d^{-1}),

was calculated as the sum of all the individual grazing rates:

359
$$(Eq.8) \quad g = \sum_j g_j = \sum_j a_{ij} x_j$$

with a_{ij} negative and a_{ji} positive. The uncertainty of g was estimated by propagating errors from

a_{ij} and x_j.

4. Results

4.1 Mesoscale patterns in environmental conditions, NFRs, and diazotroph abundances

4.1.1 Physical and biogeochemical setting

Sampled eddies varied by age and phase, as well as by in situ light, temperature, and

nutrient concentrations. At the time of sampling, both anticyclones were in a stable phase

(defined by the fluctuations of SLA amplitude in the eddy centers, Table S1) and both cyclones

were in a weakening phase. The two pairs of eddies were sampled in different seasons: summer

(July) in 2017 and early spring (March-April) in 2018. Physical variables generally followed

expected seasonal trends, with higher sea surface temperature and daily incident light in the 2017

cruise (Fig. 2) and the deepest mixed layer observed in the 2018 cruise (2018 anticyclone MLD

= 50.3 ± 17.8 m; mean MLD in other eddies ranged 17.8-30.6 m, Table S1). Temperature and in

situ light flux decreased with depth, except in the 2018 anticyclone, where temperature was

Figure 2: Depth profiles of temperature (A-B, linear scale), PAR (C-D, linear scale), PO_4^{3-} concentrations (E-F, log scale), and dFe concentrations (G-H, log scale) on the 2017 (left) and 2018 (right) cruises. Red and blue indicate observations from the anticyclonic and cyclonic eddies, respectively. Profiles of temperature and PAR are from the days and locations of the N_2 fixation measurements (note that N_2 fixation rates were measured twice within the 2018 cyclone); PO_4^{3-} and dFe concentrations were measured at nearby stations within the same eddy.

nearly constant within the upper 150 m (Fig. 2). Near-surface temperatures were similar between

eddies within each cruise; however, temperatures were lower in the lower euphotic zone of the

cyclonic eddies (Fig. 2), consistent with the uplift of deeper isopycnals. Daily integrated light did

not vary between eddies during the 2017 cruise, while in the 2018 cruise, lower daily-integrated

light was observed in the anticyclone due to cloud cover on the day of sampling (Fig. 2).

PO_4^{3-} concentrations varied with depth and eddy polarity. PO_4^{3-} concentrations were

lowest at the surface and increased with depth, except for the 2018 anticyclone, in which

concentrations were relatively stable within the upper 150 m, consistent with the near-

homogenous temperature profile (Fig. 2) and deep downwelling intensified by winter mixing

(Table S1). At depth (100-150 m), PO_4^{3-} concentrations were always higher in cyclones than in

anticyclones (Fig. 2) due to eddy-induced changes in the depths of isopycnal surfaces. In

contrast, during the 2017 cruise, PO_4^{3-} concentrations in the upper 45 m were greater in the

anticyclone (42 ± 2 nmol L^{-1}) than in the cyclone (21 ± 4 nmol L^{-1}) (Fig. 2). Consequently, the

surface $[NO_3^- + NO_2^-]:[PO_4^{3-}]$ ratio was lower in anticyclone (0.12 ± 0.03 mol:mol at 15 m, n=3)

than in the cyclone (0.37 ± 0.4 mol:mol at 15 m, n=2) during 2017; all surface ratios were well

below Redfield stoichiometry (N:P <1). During the 2018 cruise, PO_4^{3-} concentrations (range: 35-

42 nmol L^{-1}) and inorganic $[NO_3^- + NO_2^-]:[PO_4^{3-}]$ ratios (0.15-0.34 mol:mol) in the upper 45 m

did not differ between the cyclonic and anticyclonic eddies (Fig. 2).

Dissolved Fe concentrations ranged from 0.06 to 1.36 nmol L^{-1} within the upper 150 m

across all eddies (Fig. 2). Concentrations were typically highest at the surface and decreased with

depth into the lower euphotic zone; in the 2017 cyclone, dFe concentrations reached a minimum

near the deep chlorophyll maximum and increased in deeper waters alongside the increase in

macronutrients (Hawco et al., 2021). In the 2018 anticyclone, dFe concentrations did not vary

within the upper 150 m, consistent with the deeper mixed layer and downwelling of surface

waters in this eddy. Concentrations were generally higher in the 2018 cruise than in the 2017

cruise, and near-surface dFe patterns between eddies also varied on the two cruises. During the

2017 cruise, dFe in the upper 45 m ranged from 0.17 to 0.31 nmol L^{-1} and were slightly higher in

the anticyclone (0.30 ± 0.08 nmol L^{-1}) than in the cyclone (0.23 ± 0.06 nmol L^{-1}). During the

2018 cruise, elevated dFe concentrations were observed in the mixed layer of the cyclone ($1.24 \pm$

0.11 nmol L^{-1} in the upper 45 m) but not in the anticyclone (0.39 nmol L^{-1} at 40 m, within the

mixed layer). Concentrations greater than 1 nmol L^{-1} in the 2018 cyclone are anomalously high

compared to the existing measurements for Station ALOHA (typically between 0.2–0.5 nmol L^{-1}

in the mixed layer (Fitzsimmons et al., 2015)). In the weeks prior to sampling, the path of 2018

cyclonic eddy deflected off the northern coast of Maui, possibly leading to input of dFe from

coastal sources along the Hawaiian Islands.

4.1.2. N_2 fixation rates and diazotroph abundances

We measured NFRs using *in situ* arrays deployed for 24 h in each eddy center. The

highest NFRs were observed in the anticyclones during both cruises, with the highest rates

occurring in the 2017 anticyclone (up to 18.6 nmol N L^{-1} d^{-1}, Fig. 3). There were strong

differences in NFRs between eddies sampled during the 2017 cruise: depth-integrated rates (0-

125 m) were ~6 times higher in the anticyclonic eddy (670 μmol N m^{-2} d^{-1}) than in the cyclonic

eddy (115 μmol N m^{-2} d^{-1}). During the 2018 cruise, depth-integrated NFRs were ~1.7-fold higher

in the anticyclone (428 μmol N m^{-2} d^{-1}) than the cyclone (240-267 μmol N m^{-2} d^{-1}) (Fig. 2). In all

four sampled eddies, NFRs were highest at the surface and decreased with depth.

Figure 3: N₂ fixation rates (NFRs, A-B, linear scale) and *nifH* gene abundances (C-N, log-scale), measured in the center of eddies during the 2017 (left) and 2018 (right) cruises. Note that NFRs and *nifH* gene abundances were measured twice within the 2018 cyclone. × represent rates below detection limits; shaded areas indicate the detection limits of each ddPCR assay. Error bars represent the standard deviation among biological replicates (n=3 for NFR data, n=2 for 2018 ddPCR data, not available for 2017 ddPCR data). *nifH* gene abundances of UCYN-C were relatively low (maximum 2.2 × 10⁴ L⁻¹) and are not presented here; they can be found in Supporting Information Table S4.

We estimated diazotroph abundances by quantifying the *nifH* genes of seven

cyanobacterial taxa and by enumerating *Trichodesmium*, *Calothrix*, *Richelia*, and *Crocosphaera*

cells/filaments using automated flow cytometry. Depth profiles of *nifH* gene abundances were

measured in eddy centers (Fig. 3, Table 1, Table S4) and along a high-resolution sampling

transect spanning the two eddies during the 2017 cruise (Fig. 4). Automated flow cytometry cell

counts were measured continuously in the surface layer and four consecutive samples (~2 h

sampling time) corresponding to the nearest *nifH* samples were binned to provide estimates of

cell abundances in eddy centers (Table 1). Together, the *nifH* gene- and cell-based datasets

indicate that the small diazotrophs UCYN-A and UCYN-B (*Crocosphaera*) were numerically

dominant over large diazotroph taxa in all sampled eddies.

Table 1: Average *nifH* gene (genes L^{-1}) and cell abundances (cells L^{-1}) of diazotroph taxa

measured by ddPCR (15 m for 2017 cruise; 25 m for 2018 cruise) and automated flow cytometry

(7 m) in eddy centers. NA: data not available. BD: below detection limits.

	UCYN-A1	UCYN-B/ *Crocosphaera*		*Trichodesmium*	Het-1&2 *Richelia*	Het-3 *Calothrix*	Units
		(Small-size class)	(Large-size class)				
2017:							
Anticyclone	1.3×10^4	3.9×10^5		2.2×10^3	1.5×10^3	BD	genes L^{-1}
	NA	(2.1×10^5)	(2.9×10^4)	2.2×10^3	6.3×10^2	BD	cells L^{-1}
Cyclone	1.6×10^4	1.1×10^5		3.3×10^3	2.4×10^3	1.4×10^2	genes L^{-1}
	NA	(1.2×10^5)	(8.9×10^2)	1.5×10^3	6.9×10^2	1.4×10^2	cells L^{-1}
2018:							
Anticyclone	3.1×10^5	2.9×10^2		1.5×10^3	9.7×10^2	5.5×10^1	genes L^{-1}
	NA	(BD)	(2.2×10^2)	BD	BD	2.5×10^2	cells L^{-1}
Cyclone (array 1)	3.9×10^5	3.2×10^3		7.7×10^2	2.1×10^2	2.5×10^3	genes L^{-1}
	NA	(BD)	(BD)	7.1×10^2	2.7×10^2	1.9×10^3	cells L^{-1}
Cyclone (array 2)	3.4×10^5	9.4×10^2		1.4×10^2	3.3×10^1	2.8×10^3	genes L^{-1}
	NA	(BD)	(9.3×10^2)	2.2×10^2	BD	1.6×10^3	cells L^{-1}

During the 2017 cruise, *nifH* gene abundances of UCYN-A (up to 1.4×10^6 *nifH* genes L⁻

^{1}) and UCYN-B (up to 6.2×10^5 *nifH* genes L^{-1}) were higher in the anticyclone center than in the

cyclone center (Table 1, Fig. 3). Independent measurements of *Crocosphaera* cells via flow

cytometry likewise showed higher average concentrations in the anticyclone (2.4×10^5 cells L^{-1})

than in the cyclone (1.2×10^5 cells L^{-1}) and included the presence of both small and large sub-

populations (Table 1). Abundances of other taxa were lower (maximum 5.1×10^3 *nifH* genes L^{-1}

and 2.2×10^3 cells L^{-1} in surface waters of eddy centers, Table 1) and did not differ between

eddies, except for Het-3, which had higher abundances in the cyclone than in the anticyclone.

Abundances of most groups decreased with depth, although UCYN-A, UCYN-B, UCYN-C, Het-

2, and Het-3 all displayed subsurface maxima (Fig. 3, Table S4).

On the transect spanning the two 2017 eddies, large and small diazotroph taxa displayed

different spatial patterns relative to isopycnal displacements (Fig. 4). Abundances of

Trichodesmium and Het-1&2 (symbionts of diatoms) were maximal in surface waters on the

northern edge of the anticyclone and in surface waters on the cyclonic side of the front separating

the two eddies, where a visual bloom of *Trichodesmium* was also observed (Fig. 4). Bucket

samples were analyzed with the IFCb to measure the concentration of filaments at the bloom

location (148 ± 44 filaments mL^{-1}). Het-3 abundances were highest in subsurface (45 m) waters

of the cyclonic eddy, at the base of the isopycnal uplift. In contrast, abundances of UCYN-B and

UCYN-A were highest in the anticyclonic eddy, with maximal abundances from 15-45 m for

UCYN-B and 45-75 m for UCYN-A (Fig. 4).

During the 2018 cruise, UCYN-A had the highest *nifH* gene abundances of all groups

quantified (maximum abundance of 6.2×10^5 *nifH* genes L^{-1}, Table 1). UCYN-A abundances

within the upper 75 m did not vary between eddy centers, while in the lower euphotic zone (100-

125 m), abundances decreased but were higher in the anticyclone ($4.6 \times 10^3 \pm 5.6 \times 10^3$ *nifH*

genes L^{-1}) than in the cyclone ($1.2 \times 10^2 \pm 4.7 \times 10^1$ *nifH* genes L^{-1}) (Fig. 3).

Figure 4: Interpolated *nifH* gene abundances of UCYN-A1 (A), UCYN-B (*Crocosphaera*, B), Tricho (*Trichodesmium*, C), Het-1&2 (*Richelia*, D), and Het-3 (*Calothrix*, E) across the high resolution 2017 eddy transect. The contours of potential density anomaly (kg m^{-3}) in A-E depict the vertical displacement of isopycnal surfaces associated with mesoscale eddies. A surface bloom of *Trichodesmium* (yellow star, positioned in the contour based on its latitude between Stations 10 and 11) was visually detected and sampled with imaging flow cytometry.

Abundances of other groups were lower (maximum 3.8×10^4 *nifH* genes L^{-1} and 1.9×10^3 cells L^{-1}

in surface waters of eddy centers, Table 1) and generally decreased with depth (Fig. 3). *nifH*

abundances of Het-1 did not display consistent differences between eddies (Fig. 3, Table S4). In

comparison, UCYN-B, UCYN-C, *Trichodesmium*, Het-2, and Het-3 were all higher in the

cyclonic eddy than the anticyclonic eddy, despite higher NFRs in the latter. IFCb cell count data

from 2018 likewise show that abundances of large diazotrophs were generally lower than in

2017, and more abundant in the cyclone. Notably, *Crocosphaera* cells all belonged exclusively

to the large-size phenotype and *Calothrix* was mainly found free-living and not attached to their

diatom hosts.

4.2 Growth and grazing of diazotrophs at the mesoscale

We investigated potential bottom-up and top-down controls on diazotroph populations to

understand the mechanisms driving observed differences in abundances and NFRs between

eddies. Bottom-up control mechanisms were explored by modeling diazotroph growth rates as a

function of limiting factors (Section 4.2.1) and examining patterns in diazotroph P- and Fe-stress

marker gene expression in eddies (Section 4.2.2). Top-down controls were explored by

comparing the community structure of putative diazotroph grazers and modeled grazing rates in

the sampled eddies (Section 4.2.3).

4.2.1. Modeled diazotroph growth rates and limiting factors

Diazotroph growth rates and the factors limiting growth were predicted for the euphotic

zone along eddy transects. This was accomplished using a physiological model forced with mean

in situ PAR, temperature, PO_4^{3-}, and dFe concentrations and parameterized using taxa-specific

ecophysiological parameters gleaned from culture studies in the existing literature (Table S3).

Due to their dependence on light and/or temperature, growth rate predictions were generally

higher within the mixed layer and decreased with depth, with the exception of DDAs, for which

the effect of temperature has not been tested in cultures (Supporting Information S2). Growth

rates of UCYN-A could not be modeled due to a lack of available culture data and growth

parameterization.

Figure 5: Estimates of the growth rates and limiting factors for *Crocosphaera* (A, D, G, J), *Trichodesmium* (B, E, H, K), and *Richelia/Calothrix* (C, F, I, L) in eddy cruises. Predictions of growth rates (A-C, G-I) and limiting factors (D-F, J-L) across the eddy transect are based on parameters provided in Table S3 combined with interpolated cross-sections of temperature (Temp.), average scalar irradiance (PAR), and inorganic nutrient concentrations (PO_4^{3-}, dFe). Since parameters for limiting resources were identical for *Richelia* and *Calothrix*, with the exception of respective maximum growth rates reflected in the top (*Richelia*) and bottom (*Calothrix*) scales, predictions showed similar patterns. The locations of the observations with the lowest resolution (nutrient concentrations) are indicated by the black dots.

Table 2: Predictions of diazotroph biological rates in eddy centers during the 2017 and 2018 cruises. Biological rates were derived from population models applied to diazotrophs imaged by the IFCb (large *Crocosphaera*, *Trichodesmium*, *Richelia*, and *Calothrix*). Intrinsic growth rates, $\mu(T,E, PO_4^{3-},dFe)$, were predicted across the euphotic zone based on culture adaptations to temperature (T), instantaneous light (E), phosphate (PO_4^{3-}) concentration, and dissolved iron (dFe) concentration (Fig. 5). Here, we report growth rates predictions ($\mu(T,E, PO_4^{3-},dFe) \pm$ standard deviation derived from the uncertainty of parameters) matching estimates of surface (7 m) diazotroph growth rates (r $\pm$ standard deviation) and grazing rates (g $\pm$ standard deviation) based on Lagrangian time-series of diazotroph and protistan populations. The grazing pressure (g:μ) is reported to indicate whether a diazotroph population grew faster than it was grazed upon (g<μ), allowing for potential accumulation. Note that r is generally lower than the intrinsic growth rates $\mu(T,E, PO_4^{3-},dFe)$, as this parameter accounts for additional losses from other grazers, viruses, programmed cell death, vertical migration, and/or sinking. NA: Not Available

Eddy	Rates (d^{-1})	*Crocosphaera*	*Trichodesmium*	*Richelia*	*Calothrix*
2017:	$\mu(T,E,PO_4^{3-},dFe)$	0.64 ± 0.07	0.29 ± 0.07	0.09 ± 0.00	0.04 ± 0.00
	r	0.24 ± 0.06	0.18 ± 0.03	0.12 ± 0.05	0.11 ± 0.07
Cyclone	g	0.50 ± 0.14	NA	0.08 ± 0.02	0.02 ± 0.01
	g :μ (%)	78 ± 3	NA	88 ± 25	50 ± 2
	$\mu(T,E,PO_4^{3-},dFe)$	0.81 ± 0.20	0.28 ± 0.05	0.13 ± 0.00	0.05 ± 0.00
Anticyclone	r	0.42 ± 0.10	0.16 ± 0.01	0.16 ± 0.04	0.15 ± 0.10
	g	0.39 ± 0.11	NA	0.09 ± 0.04	0.18 ± 0.03
	g :μ (%)	48 ± 7	NA	69 ± 44	360 ± 16
2018:	$\mu(T,E,PO_4^{3-},dFe)$	0.43 ± 0.07	0.18 ± 0.07	0.25 ± 0.00	0.13 ± 0.00
	r	0.19 ± 0.60	0.09 ± 0.00	0.06 ± 0.01	0.18 ± 0.06
Cyclone	g	0.40 ± 0.16	NA	0.12 ± 0.02	0.04 ± 0.01
	g :μ (%)	93 ± 43	NA	48 ± 17	31 ± 25
	$\mu(T,E,PO_4^{3-},dFe)$	0.35 ± 0.08	0.18 ± 0.01	0.24 ± 0.00	0.13 ± 0.00
Anticyclone	r	0.26 ± 0.10	0.002 ± 0.001	0.02 ± 0.00	0.20 ± 0.07
	g	0.24 ± 0.09	NA	NA	0.10 ± 0.01
	g :μ (%)	68 ± 43	NA	NA	77 ± 10

In general, the compilation of growth parameters highlighted large differences in

theoretical maximum growth rates among taxa, with values of 0.86 d^{-1}, 0.6 d^{-1}, 0.26 d^{-1}, and 0.33

d^{-1} for large-size *Crocosphaera*, *Richelia*, *Calothrix*, and *Trichodesmium*, respectively. Growth

rate predictions across eddy transects were always highest for *Crocosphaera*, with maximum

rates of 0.81 ± 0.20 d^{-1} and 0.43 ± 0.07 d^{-1} within the mixed layer in 2017 and 2018 respectively

(Fig. 5A, G, Table 2). *Trichodesmium* appeared to grow at rates that were near its theoretical

maximum in 2017 (0.28-0.29 d^{-1}) and slower rates in 2018 presumably due to lower observed

temperatures (Fig. 2, Fig. 5K, L). In comparison, predictions for DDAs were consistently lower

than their theoretical maximum, with maximum growth rates of 0.25 d^{-1} and 0.13 d^{-1} for *Richelia*

and *Calothrix* respectively. DDA growth rates were generally higher in 2018, when nutrient

concentrations were elevated, and at depth (~50-75 m) in 2017, when subsurface nutrient

concentrations were well above that of the stratified surface layer (Fig. 2).

Predicted dominant limiting factors varied among taxa and between sampling periods

(Fig. 5D-F, J-L). Given the relatively low affinity of *Crocosphaera* for PO$_4$$^{3-}$ ($k_{PO_4^{3-}}$=75 ± 12

nmol L^{-1}) and the large differences in surface concentrations observed in 2017 (Fig. 2), predicted

surface growth rates (≤50 m) appeared PO$_4$$^{3-}$-limited in the cyclone and not limited by any factor

in the anticyclone (based on a limitation threshold of 80% of the maximum theoretical growth

rate). Consequently, predicted *Crocosphaera* growth rates were significantly higher in the 2017

anticyclone (0.81 ± 0.20 d^{-1}) than in the 2017 cyclone (0.64 ± 0.07 d^{-1}). This represents the only

significant difference in predicted surface growth rates between eddies across all taxa examined.

However, predicted dominant limiting factors did vary by taxa and year. In 2017, *Trichodesmium*

appeared non-limited at the surface and equally limited by PO$_4$$^{3-}$ between 50-100 m and by PAR

below 100 m, while *Richelia/Calothrix* appeared PO$_4$$^{3-}$-limited at the surface (≤50 m) and dFe-

limited at depth ($\geq$75 m) in both eddy centers. In 2018, the dominant limiting factors for

Crocosphaera and *Trichodesmium* were predicted to be temperature above 100 m and PAR at

deeper depths, while DDAs were predicted to be PO_4^{3-}-limited throughout the water column.

None of the surface or deep growth rate estimates differed between eddies, as all abiotic factors

examined remained similar in the cyclone and anticyclone (Fig 2).

4.2.2. Diazotroph expression of P- and Fe-stress marker genes

We analyzed patterns of diazotroph expression of known P- and Fe-stress marker genes

using metatranscriptomes collected within the mixed layer and deep chlorophyll maximum

during the 2017 cruise. Transcripts of six P-stress genes and three Fe-stress genes were detected

for *Crocosphaera* in surface waters (Fig. 6). Fewer gene transcripts were detected for

Trichodesmium and *Richelia* in surface waters, and transcript abundance for all these

diazotrophic cyanobacterial groups were minimal at the DCM, as expected given their lower

abundances (Fig. 6). No genes examined in UCYN-A or *Calothrix* had transcript levels above

detection limits. *Crocosphaera* transcript abundances of all six P-stress marker genes were

significantly higher within the mixed layer of the cyclonic eddy, where PO_4^{3-} concentrations

were significantly lower (Fig. 2), as compared to the anticyclonic eddy (Fig. 6). Transcripts for

two *Richelia* P-stress genes were detected at the surface: the sulfolipid biosynthase (*sqdB*)

abundances were significantly higher in the cyclonic eddy, while the high-affinity phosphate

transporter (*pstS)* abundances did not differ between eddies. The only *Trichodesmium* P-stress

gene transcript detected, a phosphonate transporter (*phnE)*, did not differ in abundance between

eddies.

Figure 6: Relative expression of P- (A-F) and Fe-stress (G-L) marker genes (according to Stenegren (2020) and Snow et al. (2015)) of *Crocosphaera*, *Trichodesmium*, and *Richelia* within the mixed layer (15 m) and at the deep chlorophyll maximum (DCM) in 2017. Individual bars represent the average (± standard errors) standardized transcript abundances for each stress marker gene in a specific eddy. P-stress markers include genes related to dissolved inorganic (*ppk: polyphosphate kinase, sqdB: sulfolipid biosynthase (substitution of P-lipids), pstS/sphX: high-affinity phosphate transporter*) and organic (*phoD: Alkaline phosphatase, phnE: phosphonate transporter*) P acquisition/transport. Fe-stress markers include genes related to dissolved inorganic iron acquisition/transport (*feoB: ferrous iron transporter, idiA: iron deficiency-induced protein*) and the substitution of enzymes involved in electron transport (*isiA: flavodoxin*). *P*-values: ** (<0.005) * (<0.01) † (not significant)

Of the three *Crocosphaera* Fe-stress genes with transcripts detected in the mixed layer,

only one (*feoB*, related to Fe(II) acquisition/transport) had higher transcripts levels in the

cyclonic eddy (Fig. 6G). The other two detected gene transcripts (*idiA*, the putative periplasmic

Fe(III) binding protein, and *isiA*, encoding the iron-free electron transfer protein flavodoxin) had

no significant difference in transcript abundance between eddies (Fig. 6G, J). The only additional

diazotroph Fe-stress gene transcript detected was the iron deficiency-induced protein (*idiA)* gene

for *Trichodesmium*, which had no significant difference in abundance between eddies (Fig.

6H,K).

4.2.3. Diazotroph grazer community structure and estimated grazing rates

We tracked the abundances of several putative grazers of cyanobacterial diazotrophs

using the IFCb. Diazotroph grazers were identified by analyzing protists co-occurring with

Crocosphaera, *Trichodesmium*, or DDAs, and via microscopic visual evidence of diazotroph

ingestion in IFCb images (Fig. 7). Putative grazers included the dinoflagellates *Gymnodinium,*

Cochlodinium, *Protoperidinium, Katodinium,* and *Prorocentrum*, the ciliates *Strombidium* and

Uronema, as well as large copepods and nauplii, most of which were imaged with ingested

diazotrophs during the eddy cruises (Fig. 7A-I). Results of a predator-prey model provide

estimates of grazing rates on specific diazotrophs during the two eddy cruises (Fig. 7J-K). The

grazing rates of copepods were not accounted for in this analysis, as their low abundance

precluded the predator-prey model from accurately estimating grazing rates (Supporting

Information S3). Also, grazing by small protists (<4 μm) or upon small diazotrophs (<4 μm,

including UCYN-A) could not be assessed using this method due to the size constraints of IFCb

imaging.

The community structure of putative grazers varied across cruises and between eddies.

Abundances of *Gymnodinium* (435 ± 100 cells L^{-1} in 2017 and 202 ± 29 cells L^{-1} in 2018),

Uronema (239 ± 40 cells L^{-1} in 2017 and 22 ± 0 cells L^{-1} in 2018), *Cochlodinium* (123 ± 4 cells

L^{-1} in 2017 and 31 ± 4 cells L^{-1} in 2018), *Amphidinium* (560 ± 78 cells L^{-1} in 2017 and 364 ± 29

cells L^{-1} in 2018), and copepods/nauplii (8 ± 3 cells L^{-1} in 2017 and 1.6 cells L^{-1} in 2018) were

higher in the 2017 cruise, while abundances of *Strombidium* (162 ± 46 cells L^{-1} in 2018 and $95 \pm$

27 cells L^{-1} in 2017), *Katodinium* ($2{,}958 \pm 373$ cells L^{-1} in 2018 and 438 ± 15 cells L^{-1} in 2017),

and *Gyrodinium* ($1{,}994 \pm 430$ cells L^{-1} in 2018 and 379 ± 90 cells L^{-1} in 2017) were higher in the

2018 cruise (Fig. 7J-K). Abundances of some taxa also varied with eddy polarity: *Prorocentrum*,

Uronema, *Gymnodinium*, and *Strombidium* were generally more abundant in cyclones while

Gyrodinium, *Amphidinium*, *Protoperidinium*, and copepods were more abundant in both

anticyclones (Fig. 7). During the 2017 cruise, the community structure of grazers was also

assessed by analyzing sequences (ribotags) from near the V4 region of the18S rRNA gene from

metatranscriptomes (methods and results described in Supporting Information S5). These data

likewise show different community structures of putative diazotroph grazers between eddies; for

instance, the relative abundance of *Protoperidinium* ribotags was higher in the anticyclonic eddy,

while *Strombidium* and *Cochlodinium* ribotags had higher relative abundances in the cyclone

(Supporting Information Fig. S3).

Results of the predator-prey model provide estimates of grazing rates on specific

diazotrophs during the two eddy cruises (Fig. 7). Maximum grazing rates ranged between 0.08-

0.23 d^{-1} for heterotrophic dinoflagellates, between 0.12-0.22 d^{-1} for mixotrophic dinoflagellates,

and 0.02-0.16 d^{-1} for ciliates (Fig. 7J-K). Predicted grazing rates varied between eddies

according to grazer abundances, with the exception of *Cochlodinium*, *Katodinium*, and

Gyrodinium, whose apparent abundances determined with the IFCb and metatranscriptome

sequencing did not differ significantly (Fig. 7 J-K, Supporting Information S5). A few taxa had

higher grazing rates in cyclones, including *Gymnodinium* (0.12 ± 0.04 d^{-1} in the cyclone and

0.001 ± 0.04 d^{-1} in the anticyclone) and *Prorocentrum* (0.18 ± 0.06 d^{-1} in the cyclone and $0.05 \pm$

0.06 d^{-1} in the anticyclone). IFCb analysis indicated that these appeared to graze mainly upon

Crocosphaera, with a mean grazing pressure (the ratio of predicted grazing rate to growth rate)

of 85 ± 46 % and higher grazing pressures in cyclonic eddies than anticyclonic eddies (Table 2).

Grazers responsible for higher grazing rates in anticyclones included *Protoperidinium* ($0.20 \pm$

0.04 d^{-1} in the anticyclone and 0.004 ± 0.006 d^{-1} in the cyclone), *Gyrodinium* (0.05 ± 0.06 d^{-1} in

the anticyclone and 0.02 ± 0.03 d^{-1} in the cyclone), and *Amphidinium* (0.07 ± 0.004 d^{-1} in the

anticyclone and 0.007 ± 0.01 d^{-1} in the cyclone). These grazers were associated with a high

grazing pressure (mean 218 ± 13 %) in anticyclonic eddies upon *Calothrix* (Table 2). Predicted

losses due to grazing of the large/endosymbiotic diazotroph *Richelia* were always below 0.1 d^{-1},

resulting in a grazing pressure of 68 ± 45% in cyclones and 69% in anticyclones (Table 2).

Figure 7: Evidence of diazotroph grazing in eddy cruises from images of co-occurring protists with ingested prey (A-I) and grazing rate estimates based on predator-prey dynamics (J-K). Grazing rates were predicted from protist abundances and interaction coefficients. Diazotrophic prey (indicated by arrows) were imaged inside copepods/nauplii (A), *Katodinium* (B), *Strombidium* (C), *Amphidinium* (D), *Prorocentrum* (E), *Gymnodinium* (F), *Gyrodinium* (H), *Uronema* (I), and *Cochlodinium* (J). Scale bars: 10 *μ*m.

5. Discussion

5.1 NFRs and diazotroph abundances at the mesoscale

Mesoscale eddies add variability to the biogeochemistry and microbial community

structure in the NPSG (Barone et al., 2022; Barone et al., 2019), but the effects of eddies on N_2

fixation are not fully understood. We sampled two pairs of eddies in spatial proximity, providing

a unique opportunity to explore the effects of anticyclonic and cyclonic eddies on diazotrophy

independent of seasonal changes. Since samples were collected in the vicinity of Station

ALOHA, we also leveraged data from the HOT program (Karl & Church, 2014) to give

historical and seasonal context to our observations.

NFRs in both anticyclone centers examined here were anomalously high relative to

previous measurements at Station ALOHA. Surface (25 m) rates in these eddies exceeded all

previous NFRs from the time-series (2012-2019), with values 3.7- and 1.8-fold above the highest

previous HOT observations from the months of sampling (July and April, respectively) (Fig. 8).

Between 2012 and 2019, 13 trackable eddies occupied Station ALOHA during a HOT cruise, yet

no NFR measurements from these cruises occurred close to the eddy center (data not shown);

hence, it is possible that we observed high NFRs in anticyclones because we specifically targeted

the center of mesoscale eddies. In contrast, rates in both cyclones were within the range of

previous observations. During the 2017 cruise, the range in depth-integrated NFRs between

eddies, sampled 5 days and ~2° latitude apart, exceeded the range in monthly mean NFRs from

the time-series measurements (~100-400 μmol N m^{-2} d^{-1}, Böttjer et al. 2017). Our results are

similar to the studies of Fong et al. (2008) and Wilson et al. (2017), which both sampled

anticyclonic eddies in the NPSG during July and observed anomalously high NFRs (surface rates

of 8.6 and 10.9 nmol N L^{-1} d^{-1}, respectively). The March-April 2018 cruise data also highlights

that elevated NFRs in anticyclones are not necessarily restricted to warm (>25°C) summer

months, as previously observed by Church et al. (2009).

Figure 8: Eddy observations (cyclones [blue] and anticyclones [red]) of N_2 fixation rates at 25 m depth (A, linear scale) and depth-integrated (0-125 m) *nifH* gene abundances (B, log scale) superimposed on historical HOT data. Boxplots illustrate ~monthly measurements observed at Station ALOHA from 2012-2019 for N_2 fixation rates (Böttjer et al., 2017) and from 2004-2017 for qPCR-derived *nifH* gene abundances (dataset doi: 10.5281/zenodo.4477269). Medians are shown as thick horizontal lines, 25-75% quantiles as boxes, the minimum and maximum values (up to 1.5 times the interquartile range) as whiskers, and outliers as black dots. Observations below detection limits are represented by colored ×. Shading is used to differentiate vertical panels.

The high NFRs observed in the 2017 and 2018 anticyclones appear to be driven by

different diazotroph taxa. UCYN-A *nifH* gene abundances were the highest of all groups

quantified during the spring (2018) cruise and UCYN-A and *Crocosphaera* (UCYN-B) were

most abundant in the summer (2017) cruise (Fig. 3, Fig. 8). This is consistent with the known

seasonal succession of *nifH* gene-based diazotroph abundances at Station ALOHA: abundances

of *Trichodesmium*, *Crocosphaera*, and DDA *nifH* genes peak in the summer or early fall, while

abundances of UCYN-A peak during the spring (Böttjer et al., 2014; Church et al., 2009).

However, our only observation of both anomalously high diazotroph abundances and high

anticyclone NFRs was during the 2017 cruise, when *Crocosphaera* abundances in the

anticyclone exceeded all previous measurements in the region from the months of June and July

(Fig. 8). The high abundances of *Crocosphaera* in the 2017 anticyclone were verified by

independent cell counts via automated flow cytometry (Table 1). Although UCYN-A abundances

were also high in the 2017 anticyclone, they only slightly exceeded the seasonal mean (Fig. 8).

We infer that *Crocosphaera* likely played a major role in driving the high NFRs observed in the

2017 anticyclone. High *Crocosphaera* abundances have been recently linked to high NFRs in

another anticyclonic eddy of the NPSG during July (Wilson et al., 2017).

It is less clear which diazotroph taxa drove the high NFRs observed in the 2018

anticyclone. In 2018, UCYN-A had the highest *nifH* gene abundances of all diazotrophs

quantified; however, abundances were near the monthly mean and did not differ between eddies

(Fig. 3, Fig. 8). No diazotroph taxa quantified via *nifH* gene abundances or automated flow

cytometry had higher abundances in the anticyclonic eddy, where the highest rates were

measured. There are several possible explanations for this apparent mismatch between

diazotroph abundances and NFRs. NFRs vary as a function of both diazotroph abundance and

cell-specific activity; hence, difference in rates between eddies could be due to differences in

cell-specific activity rather than abundance. These differences were investigated for known

cyanobacterial diazotrophs and are discussed in Section 5.2.1 below. Our ddPCR assays may

also miss important diazotrophic taxa, such as non-cyanobacterial diazotrophs, whose

importance in marine ecosystems are not well-understood (Moisander et al., 2017), or could

misrepresent organismal abundances due to polyploidy (Sargent et al., 2016), low DNA

extraction efficiency (Boström et al., 2004), or other methodological biases. Finally, the pattern

observed in 2018 may be driven by patchiness of diazotroph communities (e.g. Robidart et al.,

2014). We next examine evidence for eddy modulation of bottom-up, top-down, and physical

control on diazotroph abundance and activity to better understand the mechanistic underpinnings

of the observed patterns (Section 5.2). Finally, we address the challenges in generalizing the

effects of eddies on N_2 fixation due to confounding effects of season, diazotroph community

structure, and eddy age/stage (Section 5.3).

5.2 Mechanisms leading to high NFRs in anticyclonic eddies

5.2.1. Bottom-up control

Mesoscale eddies can modulate several factors that control diazotroph growth rates. Eddy

dynamics can affect sea surface temperature via isopycnal uplift, resulting in cold core cyclones

(McGillicuddy Jr & Robinson, 1997), and can affect light attenuation via nutrient injection and

subsequent particle formation (Benitez-Nelson et al., 2007). Temperature and light both control

cyanobacterial diazotroph growth rates (Supporting Information Table S3), and global NFRs

correlate positively to both parameters (Luo et al., 2014). Eddies also influence nutrient

availability (Hawco et al., 2021). Diazotrophs in the NPSG oscillate between P and Fe limitation

(Letelier et al., 2019) and are able to coexist with non-diazotrophic phytoplankton under low N:P

and N:Fe supply ratios (Ward et al., 2013). The uplift of isopycnals in cyclonic eddies can inject

N, P, and Fe into the lower euphotic zone, where most of these nutrients are rapidly assimilated

by the plankton community (McGillicuddy Jr & Robinson, 1997). Conversely, isopycnal

downwelling in anticyclonic eddies can result in extreme N-depletion in surface waters, which

has previously been used to explain observations of high NFRs in a NPSG anticyclone (Wilson

et al., 2017).

In the present study, we have employed a suite of techniques that allowed us to address

the potential limiting factors of diazotroph growth and modulation of these factors by mesoscale

physical forcing. We found that strong differences in NFRs between the anticyclone and cyclone

during the 2017 cruise may be partially driven by differences in PO_4^{3-} availability. Surface PO_4^{3-}

concentrations and NFRs were significantly higher in the anticyclonic eddy than in the cyclonic

eddy (Fig. 2). This pattern is consistent with a recent analysis of long-term trends at Station

ALOHA documenting the depletion in PO_4^{3-} concentrations in surface waters under low SLA

(Barone et al., 2019). In our 2017 (summertime) cruise, surface PO_4^{3-} in both 2017 eddies fell

below 50 nmol L^{-1}, the empirical threshold for P-limitation of diazotrophs in the NPSG reported

by Letelier et al. (2019), and PO_4^{3-} concentrations in the cyclone were below 30 nmol L^{-1}, an

apparent threshold for severe diazotroph P-limitation (Letelier et al., 2019). We hypothesize that

the higher PO_4^{3-} in the 2017 anticyclone partially relieved P-limitation, allowing for higher rates

of diazotroph growth and N_2 fixation by diazotrophic taxa, specifically *Crocosphaera*.

Eddy-driven reduction in P-limitation in the 2017 anticyclone is further supported by

results from our ecophysiological model, patterns in nutrient stress marker gene expression, and

previous studies. The ecophysiological model predicted that during the 2017 cruise,

Crocosphaera was PO_4^{3-}-limited in the cyclone but not in the anticyclone, driving higher

predicted growth rates in the anticyclone (Fig. 5). Indeed, in a complementary study that

surveyed the 2017 eddies on the same expedition, *Crocosphaera* populations had higher intrinsic

growth rates in the anticyclonic eddy than in the cyclonic eddy (Dugenne et al., 2020). Another

study from the same 2017 expedition reported lower bulk alkaline phosphatase activity in the

2017 anticyclone (Harke et al., 2021), supporting the postulated reduced P-limitation there.

The metatranscriptome data provide additional evidence that PO_4^{3-} availability helped

drive the difference in *Crocosphaera* growth rates between eddies: all six *Crocosphaera* genes

related to P acquisition and transport (commonly used as P-stress markers, Dyhrman & Haley,

2006; Pereira et al., 2016) had lower normalized transcript abundances in the 2017 anticyclone,

consistent with a reduction in P-limitation relative to the cyclonic eddy (Fig. 6). We view the

combined datasets presented here and in previous studies of the 2017 eddies as strong evidence

that increased PO_4^{3-} availability resulted in higher *Crocosphaera* growth rates in the anticyclone.

This may ultimately help explain the higher *Crocosphaera nifH* gene abundances, biomass, and

bulk NFRs observed in the anticyclone relative to the cyclone (Table 1, Fig. 3-4).

The higher concentrations of nutrients measured in 2018 did not seem to affect NFRs

between cyclonic and anticyclonic eddies. Our ecophysiological model predicted that most

diazotrophs were limited by low temperature, with the exception of DDAs, for which the thermal

adaptation has not been tested in cultures. While data on the physiological adaptations of UCYN-

A in culture are not available, a complementary study by Gradoville et al. (2021), which sampled

eddies on the same 2018 expedition as the present study, reported no significant difference

UCYN-A cell-specific NFRs between eddies, suggesting that bottom-up forcing did not differ

between the 2018 eddies for this organism.

755   It is less likely that the differences in NFRs between cyclonic and anticyclonic eddies

756 were driven by light, temperature, or dFe concentrations on either cruise. There were only small

757 differences in near-surface light flux or temperatures between cyclones and anticyclones, except

758 for the 2018 cruise, where lower daily-integrated light was observed in the anticyclone due to

759 cloud cover (Fig. 2). Since higher NFRs were observed in the eddy with lower light levels, the

760 opposite trend as expected (Luo et al., 2014), it is very unlikely that temperature or light drove

761 the observed differences in NFRs between eddies, though they may help explain observed

762 differences in NFRs between seasons (see Section 5.3) as well as variation in growth rates of

763 distinct taxa (Fig. 5). Likewise, based on existing parameterization of Fe-uptake and Fe-stress

764 marker genes expression, patterns in NFRs between eddies did not appear to be primarily driven

765 by dFe. In the 2018 cruise, surface dFe concentrations were anomalously high in the cyclonic

766 eddy, where lower NFRs were observed. In the 2017 cruise, dFe concentrations were slightly

767 higher in the anticyclone, where rates were higher (Fig. 2). However, ecophysiological model

768 results suggest that Fe was less limiting than PO_4^{3-} at the surface for any diazotroph group on

769 either cruise (Fig. 5). In addition, all surface dFe concentrations fell above the hypothesized Fe-

770 limitation threshold for diazotrophs (0.13 nmol L^{-1}), estimated from the empirical P-limitation

771 threshold for diazotrophs in the NPSG (Letelier et al., 2019) coupled to diazotroph cell

772 stoichiometry. There are caveats linked to testing the effect of iron concentration in culture and

773 extrapolating these experiments to field populations, as described in Supporting Information S2.

774 Nevertheless, expression patterns of Fe-stress marker genes (available for 2017 only) support our

775 model results since most transcripts did not differ between cyclonic and anticyclonic eddies (Fig.

776 6). Collectively, these data are not consistent with an overriding control by Fe on NFR in our

777 experiments.

We cannot exclude the possibility that there are other bottom-up control factors not

analyzed here that could have potentially driven differences in diazotroph growth, and ultimately

in bulk NFRs, between eddies. For example, some diazotrophs, such as *Trichodesmium,* use

dissolved organic P (DOP) in P-depleted environments (Dyhrman et al., 2006; Orchard et al.,

2010; White et al., 2010). Growth of symbiotic diazotrophs can also depend on the bottom-up

control factors of their hosts, such as silica for the diatom hosts of *Richelia* and *Calothrix*. We

did not report concentrations of DOP, silica, or any micronutrients other than dFe in this study

due to a lack of data linking growth rates of cultured diazotrophs to the availability of these

nutrients. As such, nutrients/micronutrients not assessed in our model could theoretically have

driven differences in NFRs. In sum, of those potentially limiting abiotic factors measured, we

find the strongest evidence of differential P limitation. While not exhaustive, our combined

datasets, along with model results and long-term trends at station ALOHA (Barone et al. 2019),

suggest that P may be the main abiotic factor driving the differences of NFRs in mesoscale

eddies of the NPSG.

5.2.2. Top-down control

Little is known about the identity of diazotrophic grazers, or the extent to which grazing

controls diazotroph abundances and NFRs in the NPSG (Landolfi et al., 2021). Early reports

identified mesoplanktonic copepods grazing on *Trichodesmium* (Azimuddin et al., 2016; O'Neil

et al., 1996; O'Neil, 1998), and later on UCYN-C (Hunt et al., 2016), UCYN-A (Scavotto et al.,

2015), UCYN-B, Het-1, and Het-2 (Conroy et al., 2017). Recently, small nano/microplanktonic

species (e.g. dinoflagellates and ciliates) were shown to prey on the large population of

Crocosphaera in the NPSG (Dugenne et al., 2020). These small grazers generally dominate

nano/microzooplankton biomass most of the year (Pasulka et al., 2013).

We identified numerous putative predators of diazotrophs using a predator-prey model

and direct IFCb images of ingested prey: putative grazers spanned various sizes, from large

copepods and nauplii imaged after having ingested *Trichodesmium* filaments, to nanoplanktonic

ciliates and dinoflagellates mostly preying on *Crocosphaera* or *Calothrix*. Predicted rates varied

between cyclonic and anticyclonic eddies (Fig. 7). Most of the grazers known to be mixotrophs

(Jeong et al., 2010), such as *Prorocentrum*, *Gymnodinium*, and *Strombidium*, were more

abundant in cyclones, while the heterotrophs *Protoperidinium*, *Amphidinium*, and *Katodinium*

had higher abundances in anticyclones. Although understanding the drivers of grazer

communities at the mesoscale is beyond the scope of this study, these differences in grazer

abundances resulted in clear differences in predicted grazing rates upon specific diazotrophs in

eddies of specific polarity and may help explain the observed patterns in diazotroph abundances

and NFRs. During the 2017 cruise, Het-3 (*Calothrix*) *nifH* gene abundances measured within the

mixed layer were 2-4 orders of magnitude greater in the cyclone then the anticyclone (Fig. 3) yet

predicted growth rates did not differ between eddies (Fig. 5). Our predator-prey model predicts

that rates of grazing upon *Calothrix* for the 2017 cruise were significantly higher in the

anticyclone (0.18 ± 0.03 d^{-1}, Table 2) than the cyclone (0.02 ± 0.004 d^{-1}); thus, the observed

differences in *Calothrix* abundance between eddies can simply be explained by differences in

grazing pressures. In contrast, estimated rates of grazing upon *Crocosphaera* were higher in the

cyclonic eddies for both cruises (Table 2). Therefore, the high abundance of *Crocosphaera* in the

2017 anticyclone appears driven by both enhanced growth rates and reduced grazing pressure,

which ultimately helped to drive higher bulk NFRs (Fig. 3).

Results presented here and in previous studies (Dugenne et al., 2020; Landry et al., 2008;

Wilson et al., 2017) show that grazing can be a significant loss term for small unicellular

diazotrophs, but less so for larger diazotrophs such as *Richelia* and *Trichodesmium* (Table 2).

Indeed, *Trichodesmium* has few known grazers (O'Neil & Roman, 1992), and losses are more

often thought to be driven by viral lysis (Mulholland, 2007). While we did not directly assess

viral lysis in this study, visual inspection of *Trichodesmium* filaments from the bloom at the edge

of the 2017 cyclone (Fig. 4) suggests that *Trichodesmium* losses in the bloom may have been

driven by programmed cell death or viral lysis (Berman-Frank et al., 2004; Hewson et al., 2004),

as a majority of the filaments contained cells which were shrunken or lacking thylakoids and

bloom filaments had significantly lower chlorophyll per cell than filaments imaged within the

2017 eddy centers (Supporting Information S4, Fig. S2). The fraction of dimly-fluorescing

Trichodesmium filaments did not differ between eddies of specific polarity. We view this

fraction as a proxy for senescence but other sources of diazotroph mortality, including viral lysis

and programmed cell death, should be assessed directly in future studies.

5.2.3. Physical control

837   Buoyant plankton can be concentrated in areas of surface convergence, where cells are

horizontally advected to the site of convergence and then acted upon by the opposing forces of

downwelling and upward buoyancy (D'Asaro et al., 2018; Yoder et al., 1994). The buoyancy of

diazotrophic taxa varies greatly: smaller diazotrophs (e.g. *Crocosphaera* and *Calothrix-*

Chaetoceros) are negatively buoyant and sink (Bach et al., 2012; Tuo, 2015) while large

diazotrophs (*Trichodesmium* and *Richelia* associations) can regulate their buoyancy, sometimes

becoming positively buoyant and ascending the water column at high rates (Hoppe, 2013;

Villareal & Carpenter, 2003; Villareal & Carpenter, 1989; Walsby, 1978). Elevated surface

abundances of *Trichodesmium* and *Richelia* have been observed at numerous frontal features

(Aldeco-Ramírez et al., 2009; Benavides et al., 2011; Guidi et al., 2012; Yoder et al., 1994),

presumably due to frontal downwelling and strong positive cell buoyancy. In mesoscale eddies,

downwelling can occur in intensifying anticyclones and weakening cyclones (McGillicuddy Jr,

2016), cyclones under constant wind (Gaube et al., 2015), and at the cyclonic side of eddy fronts

(Mahadevan, 2016). The downwelling-induced physical accumulation of cells has been used to

explain the occurrence of surface *Trichodesmium* slicks (Olson et al., 2015) and *Rhizosolenia*

mats (Villareal & Carpenter, 1989) in eddies of both polarities.

The fine-scale distribution of diazotrophs across the pair of adjacent eddies measured in

2017 shows evidence for the physical accumulation of *Trichodesmium* and *Richelia* (Fig. 4). The

highest *nifH* gene abundances of *Trichodesmium* and *Richelia* were observed near the surface at

the cyclonic side of the front separating the two eddies, consistent with accumulation via surface

downwelling caused by frontal dynamics. A surface bloom of *Trichodesmium* was also visually

observed at the cyclonic side of the front, likely also driven by frontal downwelling. These

results are consistent with a previous survey of *Trichodesmium* in an eddy dipole of the NPSG,

which also reported maximal abundances at the front between eddies (Guidi et al., 2012).

Our observations are insufficient for determining the importance of physical

accumulation on NFRs across the full eddy-eddy transect. NFR measurements were conducted in

the center of each eddy but not on the cyclonic side of the eddy-eddy front where *Trichodesmium*

and DDAs appeared to be physically accumulated during the 2017 cruise. Thus, while the eddy-

induced physical accumulation of large diazotrophs likely explains elevated NFRs and

abundances of large diazotrophs in some previous studies (e.g. Fong et al., 2008; Olson et al.,

2015), our eddy observations were likely driven by differences in biological control—the

balance of bottom-up and top-down control—between eddies.

5.3 Challenges in generalizing eddy effects on diazotrophs

Our finding of high NFRs in anticyclonic eddies agrees with other recent observations of

high NFRs in anticyclones from many ocean regions (Fong et al., 2008; Holl et al., 2007; Liu et

al., 2020; Löscher et al., 2016; Rahav et al., 2013; Wilson et al., 2017). However, there have also

been reports of high diazotroph abundances associated with cyclonic eddies (Olson et al., 2015),

and on a different cruise that sampled a cyclone-anticyclone eddy pair in the NPSG, surface

NFRs were significantly higher in the cyclonic eddy than in the anticyclonic eddy (dataset doi:

10.5281/zenodo.5565560). At Station ALOHA, the correlation between NFR and SLA measured

between 2012-2019 has remained non-significant (Spearman correlation, r=0.08, *p*-value=0.58,

n=48, data not shown), yet high rates of N_2 fixation are more prevalent in warm summer months

often coincident with high SLA.

Despite this relative predictability, several taxa showed inconsistent patterns with eddy

polarity during our cruises. Our data suggest that effects of eddies vary strongly among

diazotroph taxa. Our ecophysiological model predicted bottom-up effects of mesoscale eddies to

vary among diazotrophs, driven by the different adaptations of taxa to light, temperature, and

nutrient concentrations. Likewise, eddy-specific grazing varied among diazotroph taxa, with

some taxa being grazed upon more in cyclones and others being grazed upon more in

anticyclones. Physical accumulation via frontal downwelling appeared to drive the elevated

abundance of *Trichodesmium* and *Richelia* at the cyclonic side of the eddy-eddy front in 2017,

but apparently did not affect the abundances of smaller, less buoyant taxa. Unfortunately, we

were not able to assess eddy-driven mechanisms for some potentially important diazotroph taxa,

including UCYN-A, whose growth and grazing could not be predicted due to the lack of culture

data and the size constraints of IFCb imaging. Nevertheless, our results suggest that eddy effects

on small diazotrophs like *Crocosphaera* are mostly due to biological effects (the net effects of

growth and grazing), while eddy effects on large diazotrophs (*Trichodesmium* and DDAs) may

be more affected by physical mechanisms.

Finally, changes in the properties of eddies through age and stage further complicates

generalizing eddy effects. Mesoscale eddies are evolving features that intensify (with

increasingly positive or negative SLA), reach a stable phase, and finally decay over the

timescales of months (Sweeney et al., 2003). Though cyclonic and anticyclonic eddies are

commonly associated with isopycnal uplift/depression and nutrient injection/dilution in the lower

euphotic zone, respectively, in reality, vertical motions and nutrient fluxes vary with eddy age

and stage (McGillicuddy Jr & Robinson, 1997). Our eddy observations, sampled over the

timescales of days, represent snapshots of evolving features; we expect the eddy-induced

mechanisms discussed above to vary through the lifetime of an eddy. At the time of sampling,

both anticyclones were in a stable phase and both cyclones were in a weakening phase (Table

S1), so these four observations are insufficient to deduce effects of eddy stage on NFRs.

6. Conclusions

Here we report anomalously high NFRs in two anticyclonic eddies in the NPSG. We

coupled diverse datasets and ecological models to explore the mechanisms driving this pattern.

Our analyses suggest that the 2017 anticyclone modulated bottom-up (via a reduction in PO_4^{3-}

limitation) and top-down (via eddy-specific grazing) control, allowing for the net accumulation

of *Crocosphaera* cells, and ultimately, high NFRs. This mechanism may drive high NFRs in

other anticyclonic eddies in the NPSG, where surface PO_4^{3-} concentrations are significantly

higher in anticyclones than in cyclones (Barone et al., 2019), especially during the summer

months of peak *Crocosphaera* abundance. However, no drivers we tested appear to explain the

results of our 2018 expedition, for which the high NFR observed in the anticyclonic eddy

remains enigmatic. The biological and physical mechanisms through which eddies influence

NFRs vary with season, diazotroph community composition, and eddy age/stage. Future efforts

may benefit from assessing the ecological controls of uncultivated diazotrophs and from

temporal measurements throughout the lifetime of eddies to better understand how eddies might

affect successional patterns of diazotrophic communities and measured bulk NFRs.

7.   Acknowledgements

This work was funded by the Simons Foundation (Award # 721252 to DMK, 721256 to AEW,

721223 to EFD, 721221 to MJC, 721244 to EVA, 721225 to STD, 329108 to SJ, and 724220 to

JPZ) and expedition funding from the Schmidt Ocean Institute for R/V *Falkor* Cruise FK180310

in 2018. We are thankful to the chief scientists, including Tara Clemente (2017) and Steve

Poulos (2018), as well as the captains and crew members of the eddy cruises. We are also

grateful to Eric Shimabukuro, Ryan Tabata, and Tim Burrell for their help with field operations,

as well as Katie Watkins-Brandt for valuable support with the IFCb on the 2017 expedition.

8.   Data availability statement

NFRs and diazotroph abundances, measured by ddPCR and automated flow cytometry, along

with environmental data and model output from the eddy cruises can be accessed at

https://doi.org/10.5281/zenodo.6342202. Metatranscriptome sequences are available on the

NCBI SRA under project numbers PRJNA596510 and PRJNA515070. NFRs and diazotroph

nifH gene abundances from the Hawaii Ocean Time-series program can be accessed under the

doi 10.5281/zenodo.3718435, 10.5281/zenodo.6341629, and 10.5281/zenodo.4477269.

9. References

Aldeco-Ramírez, J., Monreal-Gómez, M., Signoret, M., Salas-de-León, D., & Hernández-

Becerril, D. (2009). Occurrence of a subsurface anticyclonic eddy, fronts, and

Trichodesmium spp. *Ciencias Marinas, 35*(4), 333-344.

Azimuddin, K. M., Hirai, J., Suzuki, S., Haider, M. N., Tachibana, A., Watanabe, K., Kitamura,

M., Hashihama, F., Takahashi, K., & Hamasaki, K. (2016). Possible association of

diazotrophs with marine zooplankton in the Pacific Ocean. *MicrobiologyOpen, 5*(6),

1016-1026.

Bach, L. T., Riebesell, U., Sett, S., Febiri, S., Rzepka, P., & Schulz, K. G. (2012). An approach

for particle sinking velocity measurements in the 3–400 μm size range and considerations

on the effect of temperature on sinking rates. *Marine Biology, 159*(8), 1853-1864.

Barone, B., Church, M. J., Dugenne, M., Hawco, N. J., Jahn, O., White, A. E., John, S. G.,

Follows, M. J., DeLong, E. F., & Karl, D. M. (2022). Biogeochemical dynamics in

adjacent mesoscale eddies of opposite polarity. *Global Biogeochemical Cycles, 36*(2),

e2021GB007115.

Barone, B., Coenen, A. R., Beckett, S. J., McGillicuddy, D. J., Weitz, J. S., & Karl, D. M.

(2019). The ecological and biogeochemical state of the North Pacific Subtropical Gyre is

linked to sea surface height. *Journal of Marine Research, 77*(2), 215-245.

Benavides, M., Agawin, N. S., Arístegui, J., Ferriol, P., & Stal, L. J. (2011). Nitrogen fixation by

Trichodesmium and small diazotrophs in the subtropical northeast Atlantic. *Aquatic*

Microbial Ecology, 65(1), 43-53.

Benitez-Nelson, C. R., Bidigare, R. R., Dickey, T. D., Landry, M. R., Leonard, C. L., Brown, S.

L., Nencioli, F., Rii, Y. M., Maiti, K., & Becker, J. W. (2007). Mesoscale eddies drive

increased silica export in the subtropical Pacific Ocean. *Science, 316*(5827), 1017-1021.

Berman-Frank, I., Bidle, K. D., Haramaty, L., & Falkowski, P. G. (2004). The demise of the

marine cyanobacterium, *Trichodesmium* spp., via an autocatalyzed cell death pathway.

Limnology and Oceanography, 49(4), 997-1005.

Boatman, T. G., Davey, P. A., Lawson, T., & Geider, R. J. (2018). The physiological cost of

diazotrophy for *Trichodesmium erythraeum* IMS101. *PloS One, 13*(4), e0195638.

Boström, K. H., Simu, K., Hagström, Å., & Riemann, L. (2004). Optimization of DNA

extraction for quantitative marine bacterioplankton community analysis. *Limnology and*

Oceanography: Methods, 2(11), 365-373.

Böttjer, D., Dore, J. E., Karl, D. M., Letelier, R. M., Mahaffey, C., Wilson, S. T., Zehr, J., &

Church, M. J. (2017). Temporal variability of nitrogen fixation and particulate nitrogen

export at Station ALOHA. *Limnology and Oceanography, 62*, 200-216.

Böttjer, D., Karl, D. M., Letelier, R. M., Viviani, D. A., & Church, M. J. (2014). Experimental

assessment of diazotroph responses to elevated seawater pCO_2 in the North Pacific

Subtropical Gyre. *Global Biogeochem. Cycles, 28*(6), 601-616.

Cheung, S., Nitanai, R., Tsurumoto, C., Endo, H., Nakaoka, S. i., Cheah, W., Lorda, J. F., Xia,

X., Liu, H., & Suzuki, K. (2020). Physical forcing controls the basin-scale occurrence of

nitrogen-fixing organisms in the North Pacific Ocean. *Global Biogeochemical Cycles*,

e2019GB006452.

Church, M. J., Björkman, K. M., Karl, D. M., Saito, M. A., & Zehr, J. P. (2008). Regional

distributions of nitrogen-fixing bacteria in the Pacific Ocean. *Limnology and*

Oceanography, 53, 63-77.

Church, M. J., Jenkins, B. D., Karl, D. M., & Zehr, J. P. (2005). Vertical distributions of

nitrogen-fixing phylotypes at Stn ALOHA in the oligotrophic North Pacific Ocean.

Aquatic Microbial Ecology, 38(1), 3-14.

1003

1004   Church, M. J., Mahaffey, C., Letelier, R. M., Lukas, R., Zehr, J. P., & Karl, D. M. (2009).

1005   Physical forcing of nitrogen fixation and diazotroph community structure in the North

1006   Pacific subtropical gyre. *Global Biogeochemical Cycles, 23*(2), GB2020.

1007

1008   Church, M. J., Short, C. M., Jenkins, B. D., Karl, D. M., & Zehr, J. P. (2005). Temporal patterns

1009   of nitrogenase gene (*nifH*) expression in the oligotrophic North Pacific Ocean. *Applied*

1010   *and Environmental Microbiology, 71*(9), 5362-5370.

1011

1012   Conroy, B. J., Steinberg, D. K., Song, B., Kalmbach, A., Carpenter, E. J., & Foster, R. A. (2017).

1013   Mesozooplankton graze on cyanobacteria in the Amazon River plume and western

1014   tropical North Atlantic. *Frontiers in Microbiology, 8*, 1436.

1015

1016   D'Asaro, E. A., Shcherbina, A. Y., Klymak, J. M., Molemaker, J., Novelli, G., Guigand, C. M.,

1017   Haza, A. C., Haus, B. K., Ryan, E. H., & Jacobs, G. A. (2018). Ocean convergence and

1018   the dispersion of flotsam. *Proceedings of the National Academy of Sciences USA, 115*(6),

1019   1162-1167.

1020

1021   Davis, C. S., & McGillicuddy, D. J. (2006). Transatlantic abundance of the N_2-fixing colonial

1022   cyanobacterium *Trichodesmium*. *Science, 312*(5779), 1517-1520.

1023

1024   de Boyer Montégut, C., Madec, G., Fischer, A. S., Lazar, A., & Iudicone, D. (2004). Mixed layer

1025   depth over the global ocean: An examination of profile data and a profile-based

1026   climatology. *Journal of Geophysical Research: Oceans, 109*, C12003.

1027

1028   Dugenne, M., Henderikx Freitas, F., Wilson, S. T., Karl, D. M., & White, A. E. (2020). Life and

1029   death of *Crocosphaera* sp. in the Pacific Ocean: Fine scale predator–prey dynamics.

1030   *Limnology and Oceanography, 65*, 2603-2617.

1031

1032   Dyhrman, S., Chappell, P., Haley, S., Moffett, J., Orchard, E., Waterbury, J., & Webb, E. (2006).

1033   Phosphonate utilization by the globally important marine diazotroph *Trichodesmium*.

1034   *Nature, 439*(7072), 68-71.

1035

1036   Dyhrman, S. T., & Haley, S. T. (2006). Phosphorus scavenging in the unicellular marine

1037   diazotroph *Crocosphaera watsonii*. *Applied and Environmental Microbiology, 72*(2),

1038   1452-1458.

1039

1040   Farnelid, H., Andersson, A. F., Bertilsson, S., Al-Soud, W. A., Hansen, L. H., Sørensen, S.,

1041   Steward, G. F., Hagström, Å., & Riemann, L. (2011). Nitrogenase gene amplicons from

1042   global marine surface waters are dominated by genes of non-cyanobacteria. *PloS One,*

1043   *6*(4), e19223.

1044

1045   Ferrón, S., del Valle, D. A., Björkman, K. M., Quay, P. D., Church, M. J., & Karl, D. M. (2016).

1046   Application of membrane inlet mass spectrometry to measure aquatic gross primary

1047   production by the ^{18}O *in vitro* method. *Limnology and Oceanography: Methods, 14*(9),

1048   610-622.

1049

1050   Fitzsimmons, J. N., Hayes, C. T., Al-Subiai, S. N., Zhang, R., Morton, P. L., Weisend, R. E.,

1051   Ascani, F., & Boyle, E. A. (2015). Daily to decadal variability of size-fractionated iron

1052   and iron-binding ligands at the Hawaii Ocean Time-series Station ALOHA. *Geochimica*

1053   *et Cosmochimica Acta, 171*, 303-324.

1054

1055   Follett, C. L., Dutkiewicz, S., Karl, D. M., Inomura, K., & Follows, M. J. (2018). Seasonal

1056   resource conditions favor a summertime increase in North Pacific diatom–diazotroph

1057   associations. *The ISME journal, 12*(6), 1543-1557.

1058

1059   Fong, A. A., Karl, D. M., Lukas, R., Letelier, R. M., Zehr, J. P., & Church, M. J. (2008).

1060   Nitrogen fixation in an anticyclonic eddy in the oligotrophic North Pacific Ocean. *The*

1061   *ISME journal, 2*(6), 663–676.

1062

1063   Foreman, R. K., Segura-Noguera, M., & Karl, D. M. (2016). Validation of Ti (III) as a reducing

1064   agent in the chemiluminescent determination of nitrate and nitrite in seawater. *Marine*

1065   *Chemistry, 186*, 83-89.

1066

1067   Foster, R., Subramaniam, A., Mahaffey, C., Carpenter, E., Capone, D., & Zehr, J. (2007).

1068   Influence of the Amazon River plume on distributions of free-living and symbiotic

1069   cyanobacteria in the western tropical north Atlantic Ocean. *Limnology and*

1070   *Oceanography, 52*(2), 517-532.

1071

1072   Gaube, P., Chelton, D. B., Samelson, R. M., Schlax, M. G., & O'Neill, L. W. (2015). Satellite

1073   observations of mesoscale eddy-induced Ekman pumping. *Journal of Physical*

1074   *Oceanography, 45*(1), 104-132.

1075

1076 Gifford, S. M., Becker, J. W., Sosa, O. A., Repeta, D. J., & DeLong, E. F. (2016). Quantitative

1077   transcriptomics reveals the growth-and nutrient-dependent response of a streamlined

1078   marine methylotroph to methanol and naturally occurring dissolved organic matter.

1079   *MBio, 7*(6), e01279-01216.

1080

1081 Gradoville, M. R., Cabello, A. M., Wilson, S. T., Turk-Kubo, K. A., Karl, D. M., & Zehr, J. P.

1082   (2021). Light and depth dependency of nitrogen fixation by the non-photosynthetic,

1083   symbiotic cyanobacterium UCYN-A. *Environmental Microbiology, 23*(8), 4518-4531.

1084

1085 Gradoville, M. R., Farnelid, H., White, A. E., Turk-Kubo, K. A., Stewart, B., Ribalet, F., Ferrón,

1086   S., Pinedo-Gonzalez, P., Armbrust, E. V., & Karl, D. M. (2020). Latitudinal constraints

1087   on the abundance and activity of the cyanobacterium UCYN-A and other marine

1088   diazotrophs in the North Pacific. *Limnology and Oceanography, 65*, 1858-1875.

1089   https://doi.org/10.1002/lno.11423

1090

1091 Guidi, L., Calil, P. H., Duhamel, S., Björkman, K. M., Doney, S. C., Jackson, G. A., Li, B.,

1092   Church, M. J., Tozzi, S., & Kolber, Z. S. (2012). Does eddy-eddy interaction control

1093   surface phytoplankton distribution and carbon export in the North Pacific Subtropical

1094   Gyre? *Journal of Geophysical Research: Biogeosciences, 117*, G02024.

1095

1096 Harke, M. J., Frischkorn, K. R., Hennon, G. M., Haley, S. T., Barone, B., Karl, D. M., &

1097   Dyhrman, S. T. (2021). Microbial community transcriptional patterns vary in response to

1098   mesoscale forcing in the North Pacific Subtropical Gyre. *Environmental Microbiology,*

1099   *23*(8), 4807-4822.

1100

1101   Hawco, N. J., Barone, B., Church, M. J., Babcock-Adams, L., Repeta, D. J., Wear, E. K.,

1102   Foreman, R. K., Björkman, K. M., Bent, S., & Van Mooy, B. A. (2021). Iron depletion in

1103   the deep chlorophyll maximum: mesoscale eddies as natural iron fertilization

1104   experiments. *Global Biogeochemical Cycles, 35*(12), e2021GB007112.

1105

1106   Hewson, I., Govil, S. R., Capone, D. G., Carpenter, E. J., & Fuhrman, J. A. (2004). Evidence of

1107   *Trichodesmium* viral lysis and potential significance for biogeochemical cycling in the

1108   oligotrophic ocean. *Aquatic Microbial Ecology, 36*(1), 1-8.

1109

1110   Holl, C. M., Waite, A. M., Pesant, S., Thompson, P. A., & Montoya, J. P. (2007). Unicellular

1111   diazotrophy as a source of nitrogen to Leeuwin Current coastal eddies. *Deep Sea*

1112   *Research Part II: Topical Studies in Oceanography, 54*(8-10), 1045-1054.

1113

1114   Hoppe, K. S. (2013). The sinking rate and transparent exopolymer particle (TEP) production of

1115   *Hemiaulus hauckii.* Doctoral dissertation. University of Texas at Austin.

1116

1117   Hunt, B. P., Bonnet, S., Berthelot, H., Conroy, B. J., Foster, R. A., & Pagano, M. (2016).

1118   Contribution and pathways of diazotroph-derived nitrogen to zooplankton during the

1119   VAHINE mesocosm experiment in the oligotrophic New Caledonia lagoon.

1120   *Biogeosciences, 13*(10), 3131.

1121

1122   Jassby, A. D., & Platt, T. (1976). Mathematical formulation of the relationship between

1123   photosynthesis and light for phytoplankton. *Limnology and Oceanography, 21*(4), 540-

1124   547.

1125

1126   Jeong, H. J., Yoo, Y. D., Kim, J. S., Seong, K. A., Kang, N. S., & Kim, T. H. (2010). Growth,

1127   feeding and ecological roles of the mixotrophic and heterotrophic dinoflagellates in

1128   marine planktonic food webs. *Ocean science journal, 45*(2), 65-91.

1129

1130   Karl, D., Christian, J., Dore, J., Hebel, D., Letelier, R., Tupas, L., & Winn, C. (1996). Seasonal

1131   and interannual variability in primary production and particle flux at Station ALOHA.

1132   *Deep Sea Research Part II: Topical Studies in Oceanography, 43*(2-3), 539-568.

1133

1134   Karl, D., Letelier, R., Tupas, L., Dore, J., Christian, J., & Hebel, D. (1997). The role of nitrogen

1135   fixation in biogeochemical cycling in the subtropical North Pacific Ocean. *Nature, 388*,

1136   533-538.

1137

1138   Karl, D., Michaels, A., Bergman, B., Capone, D., Carpenter, E., Letelier, R., Lipschultz, F.,

1139   Paerl, H., Sigman, D., & Stal, L. (2002). Dinitrogen fixation in the world's oceans.

1140   *Biogeochemistry, 57*(1), 47-98.

1141

1142   Karl, D. M., & Church, M. J. (2014). Microbial oceanography and the Hawaii Ocean Time-series

1143   programme. *Nature Reviews: Microbiology, 12*(10), 699-713.

1144

1145   Karl, D. M., Church, M. J., Dore, J. E., Letelier, R. M., & Mahaffey, C. (2012). Predictable and

1146   efficient carbon sequestration in the North Pacific Ocean supported by symbiotic nitrogen

1147   fixation. *Proceedings of the National Academy of Sciences USA, 109*(6), 1842-1849.

1148

1149   Karl, D. M., & Tien, G. (1992). MAGIC: A sensitive and precise method for measuring

1150   dissolved phosphorus in aquatic environments. *Limnology and Oceanography, 37*(1),

1151   105-116.

1152

1153   Landolfi, A., Prowe, A., Pahlow, M., Somes, C. J., Chien, C.-T., Schartau, M., Koeve, W., &

1154   Oschlies, A. (2021). Can top-down controls expand the ecological niche of marine N_2

1155   fixers? *Frontiers in Microbiology, 12*, 690200.

1156

1157   Landry, M. R., Brown, S. L., Rii, Y. M., Selph, K. E., Bidigare, R. R., Yang, E. J., & Simmons,

1158   M. P. (2008). Depth-stratified phytoplankton dynamics in Cyclone Opal, a subtropical

1159   mesoscale eddy. *Deep Sea Research Part II: Topical Studies in Oceanography, 55*(10-

1160   13), 1348-1359.

1161

1162   Letelier, R. M., Björkman, K. M., Church, M. J., Hamilton, D. S., Mahowald, N. M., Scanza, R.

1163   A., Schneider, N., White, A. E., & Karl, D. M. (2019). Climate-driven oscillation of

1164   phosphorus and iron limitation in the North Pacific Subtropical Gyre. *Proceedings of the*

1165   *National Academy of Sciences USA, 116*(26), 12720-12728.

1166

1167 Letelier, R. M., White, A. E., Bidigare, R. R., Barone, B., Church, M. J., & Karl, D. M. (2017).

1168   Light absorption by phytoplankton in the North Pacific Subtropical Gyre. *Limnology and*

1169   *Oceanography, 62*(4), 1526-1540.

1170

1171 Liu, J., Zhou, L., Li, J., Lin, Y., Ke, Z., Zhao, C., Liu, H., Jiang, X., He, Y., & Tan, Y. (2020).

1172   Effect of mesoscale eddies on diazotroph community structure and nitrogen fixation rates

1173   in the South China Sea. *Regional Studies in Marine Science, 35*, 101106.

1174

1175 Löscher, C., Bourbonnais, A., Dekaezemacker, J., Charoenpong, C. N., Altabet, M. A., Bange,

1176   H. W., Czeschel, R., Hoffmann, C., & Schmitz, R. (2016). N_2 fixation in eddies of the

1177   eastern tropical South Pacific Ocean. *Biogeosciences, 13*, 2889-2899.

1178

1179 Luo, E., Eppley, J. M., Romano, A. E., Mende, D. R., & DeLong, E. F. (2020). Double-stranded

1180   DNA virioplankton dynamics and reproductive strategies in the oligotrophic open ocean

1181   water column. *The ISME journal, 14*(5), 1304-1315.

1182

1183 Luo, Y.-W., Lima, I., Karl, D., Deutsch, C., & Doney, S. (2014). Data-based assessment of

1184   environmental controls on global marine nitrogen fixation. *Biogeosciences, 11*(3), 691-

1185   708.

1186

1187 Mahadevan, A. (2016). The impact of submesoscale physics on primary productivity of

1188   plankton. *Annual Review of Marine Science, 8*, 161-184.

1189

1190 McGillicuddy Jr, D. (2016). Mechanisms of physical-biological-biogeochemical interaction at

1191   the oceanic mesoscale. *Annual Review of Marine Science, 8*, 125-159.

1192

1193 McGillicuddy Jr, D., & Robinson, A. (1997). Eddy-induced nutrient supply and new production

1194   in the Sargasso Sea. *Deep Sea Research Part I: Oceanographic Research Papers, 44*(8),

1195   1427-1450.

1196

1197 McGillicuddy Jr, D. J. (2014). Do *Trichodesmium* spp. populations in the North Atlantic export

1198   most of the nitrogen they fix? *Global Biogeochemical Cycles, 28*(2), 103-114.

1199

1200 Mohr, W., Großkopf, T., Wallace, D. W. R., & LaRoche, J. (2010). Methodological

1201   underestimation of oceanic nitrogen fixation rates. *PloS One, 5*(9), e12583.

1202

1203 Moisander, P. H., Beinart, R. A., Hewson, I., White, A. E., Johnson, K. S., Carlson, C. A.,

1204   Montoya, J. P., & Zehr, J. P. (2010). Unicellular cyanobacterial distributions broaden the

1205   oceanic N_2 fixation domain. *Science, 327*(5972), 1512-1514.

1206

1207 Moisander, P. H., Beinart, R. A., Voss, M., & Zehr, J. P. (2008). Diversity and abundance of

1208   diazotrophic microorganisms in the South China Sea during intermonsoon. *The ISME*

1209   *journal, 2*(9), 954-967.

1210

1211   Moisander, P. H., Benavides, M., Bonnet, S., Berman-Frank, I., White, A. E., & Riemann, L.

1212   (2017). Chasing after non-cyanobacterial nitrogen fixation in marine pelagic

1213   environments. *Frontiers in Microbiology, 8*, 1736.

1214

1215   Montoya, J. P., Voss, M., Kahler, P., & Capone, D. G. (1996). A simple, high-precision, high-

1216   sensitivity tracer assay for N_2 fixation. *Applied and Environmental Microbiology, 62*(3),

1217   986-993.

1218

1219   Mulholland, M. (2007). The fate of nitrogen fixed by diazotrophs in the ocean. *Biogeosciences,*

1220   *4*(1), 37-51.

1221

1222   O'Neil, J., Metzler, P., & Glibert, P. (1996). Ingestion of $^{15}N_2$-labelled *Trichodesmium* spp. and

1223   ammonium regeneration by the harpacticoid copepod *Macrosetella gracilis*. *Marine*

1224   *Biology, 125*(1), 89-96.

1225

1226   O'Neil, J. M. (1998). The colonial cyanobacterium *Trichodesmium* as a physical and nutritional

1227   substrate for the harpacticoid copepod *Macrosetella gracilis*. *Journal of Plankton*

1228   *Research, 20*(1), 43-59.

1229

1230   O'Neil, J. M., & Roman, M. R. (1992). Grazers and associated organisms of *Trichodesmium*.

1231   *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs. Kluwer*, 61-73.

1232

1233 Olson, E. M., McGillicuddy, D. J., Flierl, G. R., Davis, C. S., Dyhrman, S. T., & Waterbury, J.

1234   B. (2015). Mesoscale eddies and *Trichodesmium* spp. distributions in the southwestern

1235   North Atlantic. *Journal of Geophysical Research: Oceans, 120*(6), 4129-4150.

1236

1237 Orchard, E. D., Ammerman, J. W., Lomas, M. W., & Dyhrman, S. T. (2010). Dissolved

1238   inorganic and organic phosphorus uptake in *Trichodesmium* and the microbial

1239   community: The importance of phosphorus ester in the Sargasso Sea. *Limnology and*

1240   *Oceanography, 55*(3), 1390-1399.

1241

1242 Parks, D. H., Chuvochina, M., Waite, D. W., Rinke, C., Skarshewski, A., Chaumeil, P.-A., &

1243   Hugenholtz, P. (2018). A standardized bacterial taxonomy based on genome phylogeny

1244   substantially revises the tree of life. *Nature Biotechnology, 36*(10), 996-1004.

1245

1246 Pasulka, A. L., Landry, M. R., Taniguchi, D. A., Taylor, A. G., & Church, M. J. (2013).

1247   Temporal dynamics of phytoplankton and heterotrophic protists at station ALOHA. *Deep*

1248   *Sea Research Part II: Topical Studies in Oceanography, 93*, 44-57.

1249

1250 Pereira, N., Shilova, I. N., & Zehr, J. P. (2016). Molecular markers define progressing stages of

1251   phosphorus limitation in the nitrogen-fixing cyanobacterium, *Crocosphaera*. *Journal of*

1252   *Phycology, 52*(2), 274-282.

1253

1254 Pinedo-González, P., Hawco, N. J., Bundy, R. M., Armbrust, E. V., Follows, M. J., Cael, B.,

1255   White, A. E., Ferrón, S., Karl, D. M., & John, S. G. (2020). Anthropogenic Asian

aerosols provide Fe to the North Pacific Ocean. *Proceedings of the National Academy of Sciences USA, 117*(45), 27862-27868.

Poff, K. E., Leu, A. O., Eppley, J. M., Karl, D. M., & DeLong, E. F. (2021). Microbial dynamics of elevated carbon flux in the open ocean's abyss. *Proceedings of the National Academy of Sciences USA, 118*(4), e2018269118.

Rahav, E., Herut, B., Stambler, N., Bar-Zeev, E., Mulholland, M. R., & Berman-Frank, I. (2013). Uncoupling between dinitrogen fixation and primary productivity in the eastern Mediterranean Sea. *Journal of Geophysical Research: Biogeosciences, 118*, 195-202.

Ribalet, F., Berthiaume, C., Hynes, A., Swalwell, J., Carlson, M., Clayton, S., Hennon, G., Poirier, C., Shimabukuro, E., White, A., & Armbrust, E. V. (2019). SeaFlow data v1, high-resolution abundance, size and biomass of small phytoplankton in the North Pacific. *Scientific data, 6*(1), 1-8.

Rii, Y. M., Peoples, L. M., Karl, D. M., & Church, M. J. (2021). Seasonality and episodic variation in picoeukaryote diversity and structure reveal community resilience to disturbances in the North Pacific Subtropical Gyre. *Limnology and Oceanography*.

Robidart, J. C., Church, M. J., Ryan, J. P., Ascani, F., Wilson, S. T., Bombar, D., Marin III, R., Richards, K. J., Karl, D. M., Scholin, C. A., & Zehr, J. P. (2014). Ecogenomic sensor

1278   reveals controls on N_2-fixing microorganisms in the North Pacific Ocean. *The ISME*

1279   *journal, 8*(6), 1175.

1280

1281   Sargent, E. C., Hitchcock, A., Johansson, S. A., Langlois, R., Moore, C. M., LaRoche, J.,

1282   Poulton, A. J., & Bibby, T. S. (2016). Evidence for polyploidy in the globally important

1283   diazotroph *Trichodesmium*. *FEMS Microbiology Letters, 363*(21), fnw244.

1284

1285   Scavotto, R. E., Dziallas, C., Bentzon-Tilia, M., Riemann, L., & Moisander, P. H. (2015).

1286   Nitrogen-fixing bacteria associated with copepods in coastal waters of the North Atlantic

1287   Ocean. *Environmental Microbiology, 17*(10), 3754-3765.

1288

1289   Snow, J. T., Polyviou, D., Skipp, P., Chrismas, N. A., Hitchcock, A., Geider, R., Moore, C. M.,

1290   & Bibby, T. S. (2015). Quantifying integrated proteomic responses to iron stress in the

1291   globally important marine diazotroph *Trichodesmium*. *PloS One, 10*(11), e0142626.

1292

1293   Sosik, H. M., & Olson, R. J. (2007). Automated taxonomic classification of phytoplankton

1294   sampled with imaging-in-flow cytometry. *Limnology and Oceanography: Methods, 5*(6),

1295   204-216.

1296

1297   Stenegren, M. (2020). *Significance of N_2 fixing planktonic symbioses for open ocean ecosystems*

1298   [Doctoral dissertation, Stockholm University].

1299

1300   Stukel, M. R., Coles, V. J., Brooks, M., & Hood, R. R. (2014). Top-down, bottom-up and

1301   physical controls on diatom-diazotroph assemblage growth in the Amazon River plume.

1302   *Biogeosciences, 11*(12), 3259-3278.

1303

1304   Sweeney, E. N., McGillicuddy Jr, D. J., & Buesseler, K. O. (2003). Biogeochemical impacts due

1305   to mesoscale eddy activity in the Sargasso Sea as measured at the Bermuda Atlantic

1306   Time-series Study (BATS). *Deep Sea Research Part II: Topical Studies in*

1307   *Oceanography, 50*(22-26), 3017-3039.

1308

1309   Taboada, F. G., Gil, R. G., Höfer, J., González, S., & Anadón, R. (2010). *Trichodesmium* spp.

1310   population structure in the eastern North Atlantic subtropical gyre. *Deep Sea Research*

1311   *Part I: Oceanographic Research Papers, 57*(1), 65-77.

1312

1313   Tibshirani, R., Bien, J., Friedman, J., Hastie, T., Simon, N., Taylor, J., & Tibshirani, R. J. (2012).

1314   Strong rules for discarding predictors in lasso-type problems. *Journal of the Royal*

1315   *Statistical Society, Series B (Statistical Methodology), 74*(2), 245-266.

1316

1317   Tuo, S. (2015). *The distributions and the mechanisms of the cyanobionts and their host diatoms*

1318   *in the northern South China Sea and the Kuroshio* [Doctoral dissertation, National Sun

1319   Yat-sen University].

1320

1321   Villareal, T., & Carpenter, E. (2003). Buoyancy regulation and the potential for vertical

1322   migration in the oceanic cyanobacterium *Trichodesmium*. *Microbial Ecology, 45*(1), 1-

1323   10.

1324

1325   Villareal, T. A., & Carpenter, E. J. (1989). Nitrogen fixation, suspension characteristics, and

1326   chemical composition of *Rhizosolenia* mats in the central North Pacific gyre. *Biological*

1327   *Oceanography, 6*(3-4), 327-345.

1328

1329   Walsby, A. (1978). The properties and buoyancy-providing role of gas vacuoles in

1330   *Trichodesmium* Ehrenberg. *British Phycological Journal, 13*(2), 103-116.

1331

1332   Ward, B. A., Dutkiewicz, S., Moore, C. M., & Follows, M. J. (2013). Iron, phosphorus, and

1333   nitrogen supply ratios define the biogeography of nitrogen fixation. *Limnology and*

1334   *Oceanography, 58*(6), 2059-2075.

1335

1336   Webb, E. A., Ehrenreich, I. M., Brown, S. L., Valois, F. W., & Waterbury, J. B. (2009).

1337   Phenotypic and genotypic characterization of multiple strains of the diazotrophic

1338   cyanobacterium, *Crocosphaera watsonii*, isolated from the open ocean. *Environmental*

1339   *Microbiology, 11*(2), 338-348.

1340

1341   White, A. E., Granger, J., Selden, C., Gradoville, M. R., Potts, L., Bourbonnais, A., Fulweiler, R.

1342   W., Knapp, A. N., Mohr, W., & Moisander, P. H. (2020). A critical review of the $^{15}N_2$

1343   tracer method to measure diazotrophic production in pelagic ecosystems. *Limnology and*

1344   *Oceanography: Methods, 18*(4), 129-147.

1345

1346   White, A. E., Karl, D. M., Bjorkman, K. M., Beversdorf, L. J., & Letelier, R. M. (2010).

1347   Production of organic matter by *Trichodesmium* IMS101 as a function of phosphorus

1348   source. *Limnology and Oceanography, 55*(4), 1755-1767. <Go to

1349   ISI>://WOS:000283657000023

1350

1351   Wilson, S. T., Aylward, F. O., Ribalet, F., Barone, B., Casey, J. R., Connell, P. E., Eppley, J. M.,

1352   Ferrón, S., Fitzsimmons, J. N., & Hayes, C. T. (2017). Coordinated regulation of growth,

1353   activity and transcription in natural populations of the unicellular nitrogen-fixing

1354   cyanobacterium *Crocosphaera*. *Nature Microbiology, 2*(9), 17118.

1355

1356   Wilson, S. T., Böttjer, D., Church, M. J., & Karl, D. M. (2012). Comparative assessment of

1357   nitrogen fixation methodologies, conducted in the oligotrophic North Pacific Ocean.

1358   *Applied and Environmental Microbiology, 78*(18), 6516-6523.

1359

1360   Wozniak, B., Dera, J., Ficek, D., Majchrowski, R., Ostrowska, M., & Kaczmarek, S. (2003).

1361   Modelling light and photosynthesis in the marine environment. *Oceanologia, 45*(2), 171-

1362   245.

1363

1364   Yoder, J. A., Ackleson, S. G., Barber, R. T., Flament, P., & Balch, W. M. (1994). A line in the

1365   sea. *Nature, 371*(6499), 689-692.

1366

1367   Zeev, E. B., Yogev, T., Man-Aharonovich, D., Kress, N., Herut, B., Béja, O., & Berman-Frank,

1368   I. (2008). Seasonal dynamics of the endosymbiotic, nitrogen-fixing cyanobacterium

1369   *Richelia intracellularis* in the eastern Mediterranean Sea. *The ISME journal, 2*(9), 911-

1370   923.

1371

1372   Zehr, J. P., Waterbury, J. B., Turner, P. J., Montoya, J. P., Omoregie, E., Steward, G. F., Hansen,

1373   A., & Karl, D. M. (2001). Unicellular cyanobacteria fix N_2 in the subtropical North

1374   Pacific Ocean. *Nature, 412*(6847), 635-637.

1375

1376