Efficacy of heat-killed and formalin-killed vaccines against *Tilapia tilapinevirus* in juvenile

Nile tilapia (*Oreochromis niloticus*)

Thao Thu Mai[1,2,3], Pattanapon Kayansamruaj[4], Suwimol Taengphu[5,6], Saengchan Senapin[5,6],

Janina Z. Costa[7], Jorge del-Pozo[8], Kim D. Thompson[7], Channarong Rodkhum[1,2]*, Ha Thanh

Dong[9]*

[1]Fish Infectious Diseases Research Unit (FID RU), Department of Veterinary Microbiology,

Faculty of Veterinary Science, Chulalongkorn University, Bangkok, Thailand

[2]The International Graduate Program of Veterinary Science and Technology (VST), Faculty of

Veterinary Science, Chulalongkorn University, Bangkok, Thailand

[3]Division of Aquacultural Biotechnology, Biotechnology Center of Ho Chi Minh city, Vietnam.

[4]Center of Excellence in Aquatic Animal Health Management, Faculty of Fisheries, Kasetsart

University, Bangkok, Thailand

[5]Fish Health Platform, Center of Excellence for Shrimp Molecular Biology and Biotechnology

(Centex Shrimp), Faculty of Science, Mahidol University, Bangkok, Thailand

[6]National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and

Technology Development Agency (NSTDA), Khlong Nueng, Pathum Thani, Thailand

[7]Aquaculture Research Group, Moredun Research Institute, Edinburgh, UK

[8]Infection and Immunity Division, Roslin Institute, Edinburgh, UK

[9]Faculty of Science and Technology, Suan Sunandha Rajabhat University, Bangkok, Thailand

Running head: Vaccines against *Tilapia tilapinevirus*

***Corresponding authors**:

C. Rodkhum, Channarong.R@chula.ac.th

H.T. Dong, hathanh.do@ssru.ac.th

Acknowledgements

This study was financially supported by the GCRF Networks in Vaccines Research and Development, which was co-funded by the MRC and BBSRC and is supported by the International Veterinary Vaccinology Network. Thao Thu Mai acknowledges the PhD scholarship program of Chulalongkorn University for ASEAN and Non-ASEAN countries and the 90th year anniversary of Chulalongkorn University Fund. The authors thank Ms. Putiya Sangpo and Mr. Dinh Hung Nguyen for their technical assistances.

Data Availability Statement

The data that support the findings of this study are available on request.

Conflict of interests

The authors declare no conflict of interest.

Author Contributions

Conceptualization, H.T.D., P.K., K.D.T.; investigation, T.T.M., S.T.; formal analysis, T.T.M., P.K., H.T.D.; methodology, S.S., S.T., J.Z.C, J.D.P, K.D.T, T.T.M., H.T.D; supervision, H.T.D., C.R.; writing - original draft, T.T.M, H.T.D., S.S.; review & editing, C.R., S.S., P.K., J.Z.C, J.D.P, K.D.T. All authors have read and agreed to the current version of the manuscript.

Abstract

Tilapia tilapinevirus (also known as tilapia lake virus, TiLV) is considered to be a new threat to the global tilapia industry. The objective of this study was to develop simple cell culture-based heat-killed (HKV) and formalin-killed (FKV) vaccines for the prevention of disease caused by TiLV. The fish were immunized with 100 µL of either HKV or FKV by intraperitoneal injection with each vaccine containing 1.8×10^6 TCID$_{50}$ inactivated virus. A booster vaccination was carried out at 21-day post vaccination (dpv) using the same protocol. The fish were then challenged with a lethal dose of TiLV at 28 dpv. The expression of five immune genes (*IgM, IgD, IgT*, *CD4* and *CD8*) in the head kidney and spleen of experimental fish was assessed at 14 and 21 dpv and again after the booster vaccination at 28 dpv. TiLV-specific IgM responses were measured by ELISA at the same time points. The results showed that both vaccines conferred significant protection, with relative percentage survival (RPS) of 71.3% and 79.6% for HKV and FKV, respectively. Significant up-regulation of *IgM* and *IgT* was observed in the head kidney of fish vaccinated with HKV at 21 dpv, while *IgM*, *IgD* and *CD4* expression increased in the head kidney of fish receiving FKV at the same time point. After booster vaccination, *IgT* and C*D8* transcripts were significantly increased in the spleen of fish vaccinated with the HKV, but not with FKV. Both vaccines induced a specific IgM response in both serum and mucus. In summary, this study showed that both HKV and FKV are promising injectable vaccines for the prevention of disease caused by TiLV in Nile tilapia.

KEY WORDS

Tilapia lake virus, Nile tilapia, inactivated vaccine, protection, immune responses

1. Introduction

Tilapia (*Oreochromis* sp.) is the second most farmed fish species worldwide after carps, reaching 6 million tons in 2020 (Fletcher, 2020). As the demand for animal protein as a food source increases, tilapia has been considered an important freshwater fish for low- and middle-income countries (LMICs) due to its inexpensive price, high adaptability to various environmental conditions and ease to culture (Prabu et al., 2019). Intensification of tilapia farming systems has occurred as a result of this growing demand. This has led to an increased risk of emerging infectious diseases caused by bacteria, viruses, parasites and fungi (Dong et al., 2015; Mesalhy, 2013). In 2013, a new disease with a suspected viral etiology emerged in Ecuador (Ferguson et al., 2014), and was named syncytial hepatitis of tilapia (SHT) based on its characteristic histopathological features. Around the same time, a novel RNA virus causing mass mortalities in tilapia was discovered in Israel, termed tilapia lake virus (TiLV) (Eyngor et al., 2014). Subsequent studies, supported with molecular analysis, revealed that the disease episodes in Ecuador and Israel shared the same causative virus, TiLV (Bacharachet al., 2016a; del-Pozo et al., 2014). The virus has recently been classified as *Tilapia tilapinevirus,* in the *Tilapinevirus* genus, within the *Amnoonviridae* family (Bacharach et al., 2016b).

TiLV has subsequently been reported in 14 additional countries/region worldwide, including Egypt (Fathi et al., 2017), Colombia (Kembou Tsofack et al., 2017), Thailand (Dong et al., 2017b; Surachetpong et al., 2017), India (Behera et al., 2017), Malaysia (Amal et al., 2018), Peru (Pulido, et al., 2019), Chinese Taipei, Philippines, Mexico and the United States of America (OIE, 2020), Tanzania (Mugimba et al., 2018), Uganda (Mugimba et al., 2018), Indonesia (Koesharyani et al., 2018) and Bangladesh (Chaput et al., 2020). Active movement of live fish for aquaculture trade without appropriate control measures is believed to have contributed to transboundary spreading of TiLV (Dong et al., 2017a; OIE, 2020).

Tilapia lake virus is a negative sense, segmented, single-stranded RNA virus. The virus has a diameter ranging from 55 – 90 nm, containing 10 genome segments with predicted 14 encoded proteins (Acharya et al., 2019; Bacharach et al., 2016a; del-Pozo et al., 2017; Eyngor et al., 2014). It has been demonstrated that TiLV can be horizontally transmitted from fish to fish through cohabitant studies (Eyngor et al., 2014). Intragastric challenge suggested that TiLV can be transmitted via the digestive system (Pierezan et al., 2020). Experimental infection in tilapia

broodstock has demonstrated that TiLV is present in the gonads of infected fish. Furthermore, detection of TiLV in fertilized eggs obtained from TiLV-positive broodstock and in 2-day old fry suggested that TiLV can also be vertically transmitted (Dong et al, 2020; Yamkasem et al., 2019). Current knowledge indicates that TiLV can infect all stages of fish development, including fertilized eggs, larvae, fry, fingerlings, juveniles and large-size fish (Dong et al., 2017c; Senapin et al., 2018) although fingerlings and juveniles appear to be more vulnerable to infection with the virus (Amal et al., 2018; Dong et al., 2017c; Ferguson et al., 2014; Surachetpong et al., 2017). Cumulative mortalities of up to 80% have been reported for farmed tilapia in Israel, while in a report from Ecuador the percentage of mortalities appeared to fluctuate from 10-20% up to 80% depending on the fish strain when tilapia fish were transferred to grow out cages, with fish dying within 4-7 days of transfer (Eyngor et al., 2014; Ferguson et al., 2014). The mortality levels caused by TiLV infection in Thailand were also variable, ranging from 20-90% (Dong et al., 2017c), and experimental infections also tended to result in high levels of mortality (66-100%) (Behera et al., 2017; Dinh-Hung et al., 2021; Eyngor et al., 2014; Tattiyapong et al., 2017). All of these reports suggest that TiLV is highly virulent and will cause significant mortality loss if introduced to a production site.

Vaccines are an effective way to prevent diseases caused by either bacteria or viruses in farmed fish. Many vaccines traditionally formulated from inactivated bacteria or viruses, have been licensed and are commercially available for a variety of fish species, mainly salmon, trout and carp (Ma et al., 2019). The few studies that have been reported relating to the development of a vaccine to prevent TiLV infections in tilapia as summarized in Table 1. The first TiLV vaccine was developed in Israel using strains of TiLV that were attenuated by 17-20 subsequent passages in cell culture. The prototype for these vaccines had relative percentage survival (RPS) values of over 50% (Bacharach et al., 2016c). More recently, a cell-culture derived vaccine containing virus inactivated with β-propiolactone and adjuvant Montanide IMS 1312 VG, with a virus titer of 10^8 50% tissue culture infectious dose per milliliter (TCID$_{50}$ mL^{-1}) was developed in China. The vaccine gave a relatively high level of protection, with the RPS value of 85.7 %. This vaccine was able to induce specific IgM, as well as upregulate a variety of immune genes (Zeng et al., 2021a). In another study, a DNA vaccine consisting of a pVAX1 DNA vector containing the sequence for TiLV's segment 8, encoding an immunogenic protein VP20, was used for the primary immunization and a recombinant VP20 (rVP20) protein was used as a booster vaccine given at 3-

week post-vaccination (wpv). This vaccine combination resulted in a RPS value of 72.5 %,
compared to 50 % and 52.5 % respectively for the DNA vaccine or rVP20 alone (Zeng et al.,
2021b). In the present study, we investigated whether simple cell culture-based vaccines (water-
based with no adjuvant), containing either heat-killed or formalin-killed virus, were able to
provoke a specific immune response in vaccinated fish and if the vaccines protected them from
TiLV infection.

2. Materials and methods

2.1. Fish

Juvenile Nile tilapia (*Oreochromis niloticus*) (body weight, 7.3 ± 1.2 g) were obtained from a
commercial tilapia hatchery with no previous record of TiLV infection. The fish were placed in
100-liter containers at a density of 60 fish per tank at around 28°C and fed with a commercial diet
daily at 3% of body weight for 15 days before performing the vaccination trial. Prior to the
experiment, 5 fish were randomly selected and screened for the presence of TiLV using a semi-
nested PCR (Taengphu et al., 2020). Water quality parameters (pH, NH_3, NO_2) was monitored
every 3 days using a standard Aqua test kit (Sera, Germany), and water was changed twice per
week. The vaccination study was approved by Kasetsart University Institutional Animal Care and
Use Committee (ACKU62-FIS-008).

2.2. Virus preparation

TiLV strain TH-2018-K was isolated from Nile tilapia during a TiLV outbreak in Thailand in 2018
using E11 cell line following the protocol described previously by Eyngor et al., (2014). The virus
was cultured in 75 cm^2 flasks containing confluent E11 cells and 15 ml of L15 medium at 25°C
for 5-7 days or until the cytopathic effect (CPE) of around 80 % was obtained in the cell monolayer.
The culture supernatant containing the virus was centrifuged at 4,500 g for 5 min at 4°C (Eppendorf
5810R), and stored at -80°C. The concentration of the virus was determined by calculating the
virus titre as 50% tissue culture infective dose per milliliter ($TCID_{50}$ mL^{-1}) (Reed & Muench,
1938).

2.3. Vaccine preparation

TiLV TH-2018-K (1.8×10^7 TCID$_{50}$ ml^{-1}) was used to prepare both HKV and FKV. Viral

inactivation was performed at 60°C for 2, 2.5, and 3 h or with formalin (QReC) at a final

concentration of 0.002%, 0.004%, 0.006%, 0.008% and 0.01% for 24 h at 25°C. Viral infectivity

was then checked on E11 cells. Successful inactivation of the virus was confirmed by the absence

of a CPE after 7 days with all inactivation conditions tested. Subsequently, inactivation of the virus

was performed at 60°C for 2.5 h for HKV, while incubation of 0.006% formalin at 25°C for 24 h

was used for FKV. The inactivated viral solutions were used as vaccine preparations and were not

adjuvanted. These were stored at 4°C until used.

2.4. Immunization, sampling and challenge test

Before immunization, 6 fish was chosen randomly from the fish population for blood and mucus

sampling. The vaccine study comprised of three groups (HKV, FKV and control). Each group

consisted of two 100-L replicate tanks with 25 fish each. Fish in the vaccine groups were

immunized with either HKV or FKV by intraperitoneal (IP) injection using 100 µL of vaccine.

Booster immunization was carried out at 21 dpv with the same dose of vaccine. Fish in the control

group were treated the same, except L15 medium was used in place of the virus solution. Three

fish from each tank were randomly collected at 14, 21 and 28 dpv for blood, mucus and tissue

sampling (6 biological replicates per treatment). Before sampling, fish were anaesthetized with

clove oil at 200 ppm. Mucus samples were collected from each fish by placing the fish into a

plastic bag containing 1 mL phosphate-buffered saline (PBS, 137 mM NaCl, 2.7 mM KCl, 10 mM

Na$_2$HPO$_4$, and 1.8 mM KH$_2$PO$_4$) followed by gentle rubbing for 30s. These were then centrifuged

at 4,000 g for 10 min. The mucus supernatant samples were collected and stored at -20°C until

used. Blood (~ 200 µL) was withdrawn from caudal vein using a 25G $\times$ 16 mm needle and allowed

to clot for 2 h at 4°C. Serum was collected after centrifugation the blood at 4,000 g for 10 min

(Thermo Scientific, UK) and then stored at -20°C. Tissues (head kidney and spleen) were collected,

immediately placed in Trizol solution (Invitrogen, UK), and kept at -20°C until RNA extraction.

For the challenge test, a viral stock of TiLV strain TH-2018-K (1.8×10^7 TCID$_{50}$ mL^{-1}) was diluted

2 times with sterile distilled water. Each fish was injected IP with 0.1 mL of the diluted TiLV

solution (9×10^5 TCID$_{50}$ fish^{-1}) at 28 dpv, and mortalities were monitored daily for 21 days.

2.5. Immune-related gene expression by RT-qPCR

RNA was extracted using Trizol (Invitrogen, UK) following the protocol recommended by the manufacturer. Genomic DNA contamination was removed using DNase I (Ambion, UK) according to the manufacturer's instructions. After DNase I treatment, RNA samples were re-purified using an equal volume of acid phenol:chloroform (5:1, pH 4.7) (Green & Sambrook, 2019) before checking quality and quantity of extracted RNA with Nanodrop ND-1000 Spectrophotometer (Thermo Scientific, UK). DNA contamination in the treated RNA samples was assessed by performing a qPCR cycling with tilapia elongation factor 1α (*EF-1α*) primers using No-RT master mix (absence of reverse transcriptase enzyme provided in iScript[TM] Reverse Transcription kit, Bio-Rad, USA). The cDNA synthesis (20 µL reactions) was performed using an iScript[TM] Reverse Transcription Supermix (Bio-Rad, USA) containing 100 ng RNA and incubated at 25^oC for 5 min for priming, followed by 46^oC for 20 min for reverse transcription and then 95^oC for 1 min for inactivation of the reverse transcriptase. Immune-related gene expression in the head kidney and spleen were analyzed using a quantitative real-time PCR, with specific primers as listed in Table 2 and iTaq Universal SYBR Supermix (Bio-Rad, USA). The 10 µL reaction consisted of 5.0 µL 2X Supermix, 0.5 µL forward and reverse primers (10 µM each), 1.0 µL cDNA and 3.0 µL distilled water. The reaction consisted of an initial activation at 95^oC for 2 min, followed by 40 amplification cycles of denaturation at 95^oC for 30 s, annealing at the optimal temperature of each primer pair (as shown in Table 2), and extension at 72^oC for 30 s. Gene expression data for the immune-related genes of vaccinated and control fish were normalized with that of *EF-1α* gene amplification using the $2^{-\triangle\triangle Ct}$ method (Livak & Schmittgen, 2001).

2.6. Measurement of antibody response by ELISA

Polystyrene 96 well ELISA plates were coated with 0.01% poly-L-lysine solution for 1 h. The plates were then rinsed 3 times with low salt wash buffer (LSWB, 2 mM Tris; 38 mM NaCl; 0.005% Tween 20, pH 7.3) before the addition of 100 µL of either heat- or formalin-inactivated TiLV (1.8×10^7 $TCID_{50}$ mL^{-1}) overnight at 4^oC. The plates were washes 3 times with LSWB, followed by a blocking step with PBS + 1% bovine serum albumin (BSA, Sigma) for 2 h at room temperature (around 28^oC). Then, 100 µL mucus (undiluted) or sera (diluted 1:512 in PBS) were added to each well and incubated overnight at 4^oC. The following day, the plates were washed 5 times with high salt wash buffer (HSWB, 2 mM Tris; 50 M NaCl; 0.01% Tween 20, pH 7.7) and incubated with anti-tilapia IgM (Soonthonsrima et al., 2019) diluted at the ratio 1:200 in PBS + 1% BSA for 2 h at around 28^oC. The plates were then washed 5 times with HSWB followed by

incubation of goat anti-mouse antibody (Merck, Germany) conjugated with HRP (diluted 1:3000

in LSWB + 1% BSA) for 1 h at around 28°C. The plates were finally washed 5 times with HSWB

before adding 100 µL of TMB (Merck, Germany) to each well. Color was developed in the dark

for 5-10 min before adding 50 µL of 2 M H_2SO_4 stop solution (Merck, Germany). Optical density

was read at wavelength 450 nm using the microplate reader (SpectraMax ID3, USA).

2.7. Statistical analysis

GraphPad Prism 6 was used to generate the graphs. Kaplan-Meier analysis was performed and the

log-rank test was used to compare the survival curves between vaccinated and control groups. The

relative percentage survival (RPS) was calculated using following equation:

$$RPS = 1 - \left[\frac{\text{average \% mortality of vaccinated fish}}{\text{average \% mortality of unvaccinated fish}} \right] \times 100$$

The differences in relative fold change of immune-related gene expression and specific antibody

IgM level were compared using two-way ANOVA followed by the LSD post hoc test. The

differences are considered at different levels of significance *p <0.05*, *p<0.01*, *p<0.001* and

p<0.0001.

3. Results

3.1. Efficacy of vaccine

In the challenge experiment, the first mortality occurred at 3-day post challenge (dpc) in the non-

vaccinated group (control) and at 5 and 7 dpc in the HKV and FKV groups, respectively (Fig. 1).

Mortalities continued until 13-15 dpc. Moribund fish showed gross signs of TiLV infection

including abdominal distension, skin erosion, exophthalmos and pale liver (figures not shown).

The survival rates were 81.3 ± 0.0 % and 86.3 ± 0.0% for HKV and FKV groups, respectively,

compared to 28.13 ± 30.9% for the control (*p<0.0001*). The survival percentage were analysed

using Kaplan-Meier curves with the log rank test (Fig. 1). Average RPS values were 71.3 % for

the HKV and 79.6 % for the FKV vaccine (Table 3).

3.2. Immune-related gene expression

The relative fold changes of five immune genes (*IgM*, *IgT*, *IgD*, *CD4*, *CD8α*) were compared to

that of the control group (Fig. 2). In the head kidney, a non-significant increase of *IgM* mRNA

relative to the control was noted at 14 dpv, which was followed by significant increase relative to

the control at 21 dpv for both HKV and FKV groups (Fig. 2A, *p<0.05*). A similar trend was observed for *IgT* at 14 dpv for both vaccine groups, which was followed by significantly higher expression levels at 21 dpv for the HKV group only (Fig. 2B, *p<0.05*). Regarding mRNA levels of *IgD*, there was significant up-regulation of *IgD* in the FKV group only at 21 dpv (Fig. 2C, *p<0.01*). The *CD4* gene was significantly upregulated at 14 dpv in the HKV only (Fig. 2D, *p<0.05*) and at 21 dpv in the FKV (*p < 0.001*). No statistical difference was observed in *CD8α* expression between the vaccinated and control groups at the time point examined (Fig. 2E).

In the spleen, non-significant, relative up-regulation of *IgM* expression was noted in both HKV and FKV groups compared to the control at 14 dpv. (Fig. 2F). There was a slight increase of *IgM* mRNA level relative to the control in the HKV group after booster (28 dpv), which were not significant. Also at 28 dpv, *IgT* expression was over 25 times higher in the HKV group (*p<0.05*) and almost 20 times higher in the FKV group (Fig. 2G). A slight significant increase in *IgD* expression was seen in the HKV group at 14-dpv (Fig. 2H, *p<0.05*). No significant increase of *CD4* expression was found at any time point (Fig. 2I); meanwhile, an approximately tenfold increase of *CD8α* expression was observed at 28 dpv in the HKV group (Fig. 2J, *p<0.05*).

3.3. Detection of antibody IgM against TiLV in serum and mucus.

Systemic TiLV-specific antibody IgM (anti-TiLV IgM) levels pre-vaccination (0 dpv) and at 14, 21 and 28 dpv, as indicated by optical density (OD) at 450 nm, were determined by ELISA (Fig. 3A). Before immunization, the average OD value of the fish sera was 0.096 ± 0.009. The OD readings for HKV, FKV and control groups were 0.254 ± 0.053, 0.363 ± 0.09 and 0.096 ± 0.015 at 14 dpv, respectively. The OD values showed an increase in antibody levels in both groups of vaccinated fish, but were only statistically different in the FKV group (*p<0.01*). A slight decrease was seen in OD readings at 3 wpv in both the HKV and FKV groups relative to the control group (0.249 ± 0.049, 0.317 ± 0.043 and 0.128 ± 0.017, respectively). One week after the booster vaccination at 28 dpv, the anti-TiLV IgM levels had increased considerably in both the HKV (*p<0.001*) and the FKV (*p<0.0001*) groups, reaching the highest values obtained between the different sampling points, compared to that of the non-vaccinated group (average OD readings were 0.438 ± 0.127, 0.483 ± 0.088, and 0.081 ± 0.01 respectively) (Fig. 3A).

A similar pattern was observed with the mucosal anti-TiLV IgM response (Fig. 3B). Before vaccination, the average OD value of fish mucus was 0.068 ± 0.003. At 14 dpv, the TiLV-specific

antibody IgM rose in both of the vaccinated groups, HKV and FKV, compared to the non-

vaccinated group (0.251 ± 0.104, 0.404 ± 0.142, and 0.07 ± 0.005, respectively), but a significant

difference was only noted for the FKV group (*p<0.01*). At 3 wpv, the antibody levels were not

significantly differ between groups, with OD values of 0.159 ± 0.031 (HKV), 0.290 ± 0.064

(FKV), and 0.083 ± 0.007 (control) being recorded. At 4 wpv (after administering the booster

vaccination), a considerable increase in anti-TiLV IgM levels was seen in the mucus of the FKV

group (*p<0.001*) (0.585 ± 0.145), whereas the increase measured in HKV fish (0.235 ± 0.044) was

not statically different to that of the control group (0.107 ± 0.018). No significant changes in

average OD readings was seen between the non-vaccinated group and pre-immunized fish in either

sera or mucus (Fig. 3 A-B).

4. Discussion

4.1. Both simple HKV and FKV were effective in protecting tilapia from TiLV infection

Although many different types of vaccines have been developed for aquaculture in recent years,

whole-cell inactivated vaccines remain the major type of vaccine licensed for use by the

aquaculture industry (Kayansamruaj et al., 2020; Ma et al., 2019). They are safe, relatively simple

to produce, and are affordable for farmers, especially for species that are intensively cultured, but

low in price like tilapia in LMICs. In this study, we prepared two versions of simple water-based

inactivated vaccine (HKV and FKV) for TiLV and assessed the ability of both to protective tilapia

against the virus. Both HKV and FKV were able to confer relatively high levels of protection

(RPS, 71.3% vs. 79.6%) in vaccinated fish. Differences in methods used to inactivate the virus,

vaccine formulation, viral strains, antigen concentration, route of vaccine administration and the

population of fish can all contribute to the level of protection obtained from a vaccine (Table 1).

Despite this, vaccination is still considered as a promising strategy to protect tilapia from TiLV

infection, although the design of the vaccine should be carefully considered to optimize the level

of protection obtained. Other inactivated vaccines have shown relatively high levels of protection

in fish. For example, other formalin-killed vaccines resulted in RPS values of 79%, 81.9% and

74% for infectious hematopoietic necrosis virus in rainbow trout (*Oncorhynchus mykiss*) (Tang et

al., 2016), *Betanodavirus* in European sea bass (*Dicentrarchus labrax*) (Nuñez-Ortiz et al., 2016),

and scale drop disease virus (SDDV) in barramundi (*Lates calcarifer*) (de Groof et al., 2015),

respectively. In addition, a heat-killed *Aeromonas hydrophila* vaccine gave 84% protection in rain-
bow trout (Dehghani et al., 2012). Although the efficacy of these and the current vaccines were
not tested against heterologous strains of TiLV, the high level of protection elicited against the
homologous strain suggests that autogenous inactivated vaccines may be effective as an
emergency vaccine to reduce the risk of production losses in affected tilapia farms.

4.2. Immunization with HKV or FKV activated both branches of the tilapia's specific immune system

Upregulation in the expression of *IgM*, *IgD* and *IgT* and *CD4* (genes encoding proteins involved
in humoral immunity) and *CD8* (cell-mediated immunity) following immunization with HKV and
FKV suggests that the vaccines are able to activate both arms of the specific immune response in
Nile tilapia. Protection from these vaccines is, therefore, likely to result from a synergistic effect
of humoral (B cell) and cellular immune (T cell) responses. This is similar to the recent report by
Zeng et al (2021a), showing that β-propiolactone-inactivated TiLV vaccines induced up-regulation
of *MHC-I* and *MHC-II/CD4,* which belong to different arms of the immune system.

The increase in *CD4* transcripts at 14 and 21 dpv in fish vaccinated with HKV or FKV may reflect
activated naïve CD4+ cells differentiating into helper T-cell subsets, Th1 and Th2. The Th1 cells
produce cytokines that stimulate the expression of anti-viral and inflammatory genes, whereas
cytokines secreted by Th2 cells stimulate the differentiation of B-cells into plasma cells to produce
specific antibody (Secombes & Wang, 2012; Secombes & Belmonte; 2016; Smith et al., 2019).
On the other hand, CD8α transcription was only seen to be significantly up-regulated in the spleen
of the HKV group after booster vaccination, indicating that the HKV may stimulate CD8+ cell
activation, which then differentiate into cytotoxic T-cells. These cells play a crucial role in cell-
mediated immunity (Bo et al., 2012; Somamoto et al., 2002; Smith et al., 2019).

As well as assessing the expression of *IgM* transcripts, this study also examined the expression of
two additional immunoglobulins IgD and IgT. Similar patterns of up-regulation were found in
head kidney of fish after the primary immunization, suggesting that all three antibodies may be
involved in the protective response elicited by the vaccines. Interestingly, significant increases in
mRNA *IgT* levels were seen in the head kidney before booster vaccination and in the spleen after
the booster vaccination for both the HKV and FKV groups, suggesting that IgT may be strongly
associated with the protective response against TiLV. Unfortunately, the function of IgT in tilapia

remains poorly understood. Functional localization studies in other fish species have shown that
IgT plays an important role against infectious pathogens on mucosal surfaces, such as skin, gills
and gut (Smith et al., 2019; Salinas et al., 2021; Zhang et al., 2011). Nevertheless, further studies
are required to gain a better understanding on the role of IgT in tilapia's defense system, especially
in response to infection.

4.3. HKV and FKV induce both systemic and mucosal IgM

In present study, HKV and FKV were shown to trigger both systemic and mucosal IgM responses,
with similar patterns observed between the two vaccines. The increase in systemic and mucosal
IgM in teleost is usually derived from the major lymphoid organs, such as head kidney and spleen
(Zapata et al., 2006), but also from the mucosa-associated lymphoid organs located in the skin,
gills, gut, or nasopharynx (not investigated in this study) (Smith et al., 2019; Salinas et al., 2021).
In the present study, up-regulation of IgM expression occurred mainly in the head kidney, and to
a less extent in the spleen, suggesting head kidney to be one of the main organs for IgM production
in response to the TiLV vaccines. Although the pathway of IgM secretion in the mucosal
compartment (mucus) is unclear, it is possible that mucosal antibodies are produced locally in the
mucosa-associated lymphoid organs and/or by the systemic immune system (Ángeles Esteban &
Cerezuela, 2015; Koppang et al., 2015; Salinas et al., 2011; Salinas & Parra, 2015; Salinas et al.,
2021). In other research using Asian seabass, monovalent and bivalent bacterial vaccines induced
both systemic and mucosal IgM (Thu-Lan et al., 2021). Similar kinetics have been reported for
IgM secretion in the serum of red hybrid tilapia, infected IP with TiLV (Tattiyapong et al., 2020).
The levels of serum IgM increased significantly in Nile tilapia after immunization with β-
propiolactone-inactivated virus (Zeng et al. 2021a) or with a recombinant vaccine based on
segment 8 of TiLV (Zeng et al., 2021b). Mucosal IgM was not investigated in these studies,
however. The presence of TiLV-specific IgM in the mucus of vaccinated fish suggests that these
vaccines may be able to generate a primary immune response in multiple mucosal organs such as
skin and gills, which are crucial sites to prevent the initial invasion of pathogenic agents (Esteban
& Cerezuela, 2015; Koppang et al., 2015). The IgM levels produced by FKV was always slightly
higher than HKV in both serum and mucus at all sampling points analyzed, indicating that FKV
induces stronger systemic and mucosal IgM responses than HKV. This could be one of the factors
explaining for slightly higher level of protection conferred by FKV.

In this study, increased levels of TiLV specific IgM after booster vaccination in both serum and

mucus indicate successful induction of specific immune memory after first immunization. B-cells

are the major component involved in humoral adaptive immunity. They are activated by specific

antigen binding to the B-cell receptors on the cell, followed by presentation of processed antigens

to naïve CD4-Tcells, which then differentiate into helper T-cells. With T cell help, B-cells

differentiate into plasma cells and memory B-cells. Plasma cells are committed to antibody

secretion, whereas memory B-cells are responsible for the long-lasting protection from subsequent

exposure to the same pathogens (Secombes & Belmonte, 2016; Smith et al., 2019).

Although systemic and mucosal IgM levels were assessed in the study, we were unable to measure

levels of other antibodies i.e. IgD and IgT by ELISA due to a lack of monoclonal antibodies for

these immunoglobulin classes in tilapia. Further studies should investigate the persistence of the

immune response in vaccinated fish, duration of protection and efficacy testing these vaccines in

a commercial setting.

In conclusion, this study reported on the efficacy of two simple TiLV inactivated vaccines without

adjuvant (HKV and FKV) in preventing TiLV infection in Nile tilapia. The vaccines activated both

branches of adaptive immunity, triggered expression of three immunoglobulin classes and elicited

both systemic and mucosal IgM responses. Most importantly, these vaccines showed relatively

high levels of protection against TiLV infection, and therefore seem very promising for the

prevention of disease associated with TiLV.

References

Acharya, V., Jyoti, H., Ajaya, C., Rout, K., Balabantaray, S., & Kumar, B. (2019). Structural

Characterization of Open Reading Frame - Encoded Functional Genes from Tilapia Lake

Virus (TiLV). *Molecular Biotechnology*, *61*(12):945-957. doi: org/10.1007/s12033-019-

00217-y

Amal, M. N. A., Koh, C. B., Nurliyana, M., Suhaiba, M., Nor-Amalina, Z., Santha, S., … Zamri-

Saad, M. (2018). A case of natural co-infection of Tilapia Lake Virus and *Aeromonas*

veronii in a Malaysian red hybrid tilapia (*Oreochromis niloticus* × *O. mossambicus*) farm

experiencing high mortality. *Aquaculture*, *485,* 12–16. doi:

org/10.1016/j.aquaculture.2017.11.019

Ángeles Esteban, M., & Cerezuela, R. (2015). Fish mucosal immunity: skin BT - Mucosal

Health in Aquaculture. In *Mucosal Health in Aquaculture*. doi: org/10.1016/B978-0-12-
417186-2/00004-2

Bacharach, E., Mishra, N., Briese, T., Zody, M. C., Tsofack, J. E. K., Zamostiano, R., … Lipkin,
W. I. (2016a). Characterization of a novel orthomyxo-like virus causing mass die-offs of
Tilapia. *MBio*, *7*(2), 1–7. doi: org/10.1128/mBio.00431-16

Bacharach, E., Mishra, N., Briese, T., Eldar, A., Lipkin, W. I., Kuhn, J. H., & Lipkin, W. I.
(2016b). ICTV Taxonomic Proposal 2016.016a-dM.A.v2.Tilapinevirus. *Create the*
Unassigned Genus Tilapinevirus. Retrieved from:
https://talk.ictvonline.org/ICTV/proposals/2016.016a-dM.A.v2.Tilapinevirus.pdf

Bacharach, E., Eldar, A. (2016c). Tilapia lake virus vaccines. *US Patent Application Publication*
no.US2016/0354458A1. Retrieved from:
https://patents.google.com/patent/US20160354458A1/en

Behera, B. K., Pradhan, P. K., Swaminathan, T. R., Sood, N., & Paria, P. (2017). Emergence of
Tilapia Lake Virus associated with mortalities of farmed Nile Tilapia *Oreochromis niloticus*
(Linnaeus 1758) in India Emergence of Tilapia Lake Virus associated with mortalities of
farmed Nile Tilapia *Oreochromis niloticus* (Linnaeus 1758) . *Aquaculture*, *484*, 168–174.
doi: org/10.1016/j.aquaculture.2017.11.025

Bo, A., O, O., Po, O., & Io, O. (2012). Immune Response of Fish to Viral Infection. *Nature and*
Science, *10*(8):70-76.

Chaput, D. L., Bass, D., Alam, M., Hasan, N. Al, Stentiford, G. D., Aerle, R. Van, … Tyler, C.
R. (2020). The Segment Matters : Probable Reassortment of Tilapia Lake Virus (TiLV)
Complicates Phylogenetic Analysis and Inference of Geographical Origin of New Isolate
from Bangladesh. *Viruses*, *12*(3), 258. doi: org/10.3390/v12030258

De Groof, A., Guelen, L., Deijs, M., van der Wal, Y., Miyata, M., Ng, K. S., … van der Hoek, L.
(2015). A Novel Virus Causes Scale Drop Disease in *Lates calcarifer*. *PLoS Pathogens*,
11(8). doi: org/10.1371/journal.ppat.1005074

Dehghani, S., Akhlaghi, M., & Dehghani, M. (2012). Efficacy of formalin-killed, heat-killed and
lipopolysaccharide vaccines against motile aeromonads infection in rainbow trout
(*Oncorhynchus mykiss*). *Global Veterinaria*, *9*(4), 409–415. doi:
org/10.5829/idosi.gv.2012.9.4.6591

Del-Pozo, J., Mishra, N., Kabuusu, R., Cheetham, S., Eldar, A., Bacharach, E., … Ferguson, H.

W. (2017). Syncytial Hepatitis of Tilapia (*Oreochromis niloticus* L.) is Associated With

Orthomyxovirus-Like Virions in Hepatocytes. *Veterinary Pathology*, *54*(1), 164–170. doi:

org/10.1177/0300985816658100

Dinh-Hung, N., Sangpo, P., Kruangkum, T., Kayansamruaj, P., Rung-ruangkijkrai, T., Senapin,

S., … Dong, H. T. (2021). Dissecting the localization of Tilapia tilapinevirus in the brain of

the experimentally infected Nile tilapia, *Oreochromis niloticus* (L.). *Journal of Fish*

Diseases, *00*, 1– 12. doi: org/10.1111/jfd.13367

Dong, H.T., Senapin, S., Gangnonngiw, W., Nguyen, V.V., Rodkhum, C., Debnath, P.P.,

…Mohan, C. V. (2020). Experimental infection reveals transmission of tilapia lake virus (

TiLV) from tilapia broodstock to their reproductive organs and fertilized eggs.

Aquaculture, *515*. doi: org/10.1016/j.aquaculture.2019.734541

Dong, H., Rattanarojpong, T., & Senapin, S. (2017a). *Urgent update on possible worldwide*

spread of tilapia lake virus (TiLV) Urgent update on possible worldwide spread of tilapia

lake virus (TiLV). Retrieved from: http://www.fao.org/fi/static-

media/MeetingDocuments/TiLV/d28.pdf

Dong, H. T., Ataguba, G. A., Khunrae, P., Rattanarojpong, T., & Senapin, S. (2017b). Evidence

of TiLV infection in tilapia hatcheries in Thailand from 2012 to 2017 reveals probable

global spread of the disease. *Aquaculture*, *479*, 579–583. doi:

org/10.1016/j.aquaculture.2017.06.035

Dong, H. T., Siriroob, S., Meemetta, W., Santimanawong, W., Gangnonngiw, W., Pirarat, N., …

Senapin, S. (2017c). Emergence of tilapia lake virus in Thailand and an alternative semi-

nested RT-PCR for detection. *Aquaculture*, *476*, 111–118. doi:

org/10.1016/j.aquaculture.2017.04.019

Dong, Ha Thanh, Nguyen, V. V., Le, H. D., Sangsuriya, P., Jitrakorn, S., Saksmerprome, V., …

Rodkhum, C. (2015). Naturally concurrent infections of bacterial and viral pathogens in

disease outbreaks in cultured Nile tilapia (*Oreochromis niloticus*) farms. *Aquaculture*, *448*,

427–435. doi: org/10.1016/j.aquaculture.2015.06.027

Esteban, M.A., & Cerezuela, R. (2015). Fish mucosal immunity: Skin. In *Mucosal Health in*

Aquaculture (pp 67-93). doi: org/10.1016/B978-0-12-417186-2.00004-2

Eyngor, M., Zamostiano, R., Tsofack, J. E. K., Berkowitz, A., Bercovier, H., Tinman, S., …

Eldar, A. (2014). Identification of a novel RNA virus lethal to tilapia. *Journal of Clinical*

Microbiology, *52*(12), 4137–4146. doi: org/10.1128/JCM.00827-14

Fathi, M., Dickson, C., Dickson, M., Leschen, W., Baily, J., Muir, F., … Weidmann, M. (2017).
Identification of Tilapia Lake Virus in Egypt in Nile tilapia affected by 'summer mortality'
syndrome. *Aquaculture*, *473*, 430–432. doi: org/10.1016/j.aquaculture.2017.03.014

Ferguson, H. W., Kabuusu, R., Beltran, S., Reyes, E., Lince, J. A., & del Pozo, J. (2014).
Syncytial hepatitis of farmed tilapia, *Oreochromis niloticus* (L.): A case report. *Journal of
Fish Diseases*, *37*(6), 583–589. doi: org/10.1111/jfd.12142

Fletcher, R. (2020, October 6). 2020 tilapia production figures revealed. Retrieved from:
https://thefishsite.com/articles/2020-tilapia-production-figures-revealed

Green MR, Sambrook J. (2019). Removing DNA Contamination from RNA Samples by
treatment with RNase-Free DNase I. *Cold Spring Harb Protoc* (pp. 709-710). doi:
10.1101/pdb.prot101725.

Japhette Esther Kembou Tsofack, a Rachel Zamostiano, a Salsabeel Watted, b Asaf Berkowitz,
B., Ezra Rosenbluth, B., Nischay Mishra, C., Thomas Briese, C., W. Ian Lipkin, c Richard
M. Kabuusu, D., Hugh Ferguson, D., … Bacharach, E. (2017). Detection of Tilapia Lake
Virus in Clinical Samples by Culturing and Nested. *Journal of Clinical Microbiology*,
55(3), 759–767. doi: 10.1128/JCM.01808-16

Kayansamruaj, P., Areechon, N., & Unajak, S. (2020). Development of fish vaccine in Southeast
Asia: A challenge for the sustainability of SE Asia aquaculture. *Fish and Shellfish
Immunology*, *103*, 73–87. doi: org/10.1016/j.fsi.2020.04.031

Koesharyani, I., Gardenia, L., Widowati, Z., Khumaira, K., & Rustianti, D. (2018). Studi kasus
infeksi tilapia lake virus (TiLV) pada ikan nila (*Oreochromis niloticus*). *Jurnal Riset
Akuakultur*, *13*(1), 85. doi: org/10.15578/jra.13.1.2018.85-92

Koppang, E. O., Kvellestad, A., & Fischer, U. (2015). Mucosal Health in Aquaculture Fish
mucosal immunity: gill (pp 93 -133). In *Mucosal Health in Aquaculture*. doi:
org/10.1016/B978-0-12-417186-2/00005-4

L.L.H. Pulido, M.M. Chìo, A.L.H. Chaparro, et al.. (2019). Tilapia lake virus (TiLV) from Peru
is genetically close to the Israeli isolates. *Aquaculture, 510*, 61-65. doi:
org/10.1016/j.aquaculture.2019.04.058

Ma, J., Bruce, T. J., Jones, E. M., & Cain, K. D. (2019). A review of fish vaccine development
strategies: Conventional methods and modern biotechnological approaches.

Microorganisms, *7*(11). doi: org/10.3390/microorganisms7110569

Mesalhy, S. (2013, July). A Review of Fish Diseases in the Egyptian Aquaculture. *Cgiar*.

Retrieved from:

https://cgspace.cgiar.org/bitstream/handle/10568/34870/EgyptAquacultureSectorDiseaseRe

view.pdf?sequence=1

Mugimba, K. K., Chengula, A. A., Wamala, S., Mwega, E. D., Kasanga, C. J., Byarugaba, D. K.,

… Munang'andu, H. M. (2018). Detection of tilapia lake virus (TiLV) infection by PCR in

farmed and wild Nile tilapia (*Oreochromis niloticus*) from Lake Victoria. *Journal of Fish*

Diseases, *41*(8), 1181–1189. doi: org/10.1111/jfd.12790

Nuñez-Ortiz, N., Pascoli, F., Picchietti, S., Buonocore, F., Bernini, C., Toson, M., … Toffan, A.

(2016). A formalin-inactivated immunogen against viral encephalopathy and retinopathy

(VER) disease in European sea bass (*Dicentrarchus labrax*): Immunological and protection

effects. *Veterinary Research*, *47*(1). doi:org/10.1186/s13567-016-0376-3

OIE - World Organisation for Animal Health. (2020, May). Infection with Tilapia lake virus

(TiLV) - A novel Orthomyxo-like virus. *Disease Information Cards*. Retrieved from

https://www.oie.int/fileadmin/Home/eng/Internationa_Standard_Setting/docs/pdf/Aquatic_

Commission/A_TiLV_disease_card.pdf

Pierezan, F., Yun, S., Piewbang, C., Surachetpong, W., & Soto, E. (2020). Pathogenesis and

immune response of Nile tilapia (*Oreochromis niloticus*) exposed to Tilapia lake virus by

intragastric route. *Fish and Shellfish Immunology*, *107*, 289–300. doi:

10.1016/j.fsi.2020.10.019

Prabu, E., Rajagopalsamy, C. B. T., Ahilan, B., Jeevagan, I. J. M. A., & Renuhadevi, M. (2019).

Tilapia – An Excellent Candidate Species for World Aquaculture: A Review. *Annual*

Research & Review in Biology (pp. 1–14). doi: org/10.9734/arrb/2019/v31i330052

Reed & Muench. (1938). The American. *American Journal of Epidemiology*, *27*(3), 493–497.

doi: org/10.7723/antiochreview.72.3.0546

Salinas I., Zhang Y. A., & Sunyer. J. O. (2011). Mucosal immunoglobulins and B cells of

Teleost fish Irene. NIH Public Access. *Developmental & Comparative Immunology*,

35(12), 1346–1365. doi: org/10.1016/j.dci.2011.11.009.Mucosal

Salinas, I., Fernández-Montero, Á., Ding, Y., & Sunyer, J. O. (2021). Mucosal immunoglobulins

of teleost fish: a decade of advances. *Developmental & Comparative Immunology*, *121*:

104079. doi: org/10.1016/j.dci.2021.104079

Salinas, I., & Parra, D. (2015). Fish mucosal immunity: Intestine. In *Mucosal Health in*

Aquaculture (pp 135-170). doi: org/10.1016/B978-0-12-417186-2.00006-6

Secombes, C.J., & Wang, T. (2012). The innate and adaptive immune system of fish. In

Infectious Disease in Aquaculture. doi: 10.1533/9780857095732.1.3

Secombes, Chris J., & Belmonte, R. (2016). Overview of the fish adaptive immune system. In

Fish vaccines (pp. 35–52). doi: 10.1007/978-3-0348-0980-1_2

Senapin, S., Shyam, K. U., Meemetta, W., Rattanarojpong, T., & Dong, H. T. (2018). Inapparent

infection cases of tilapia lake virus (TiLV) in farmed tilapia. *Aquaculture*, *487*, 51–55. doi:

10.1016/j.aquaculture.2018.01.007

Smith, N. C., Rise, M. L., & Christian, S. L. (2019). A Comparison of the Innate and Adaptive

Immune Systems in Cartilaginous Fish, Ray-Finned Fish, and Lobe-Finned Fish. *Frontiers*

in Immunology, *10*, 2292. doi: org/10.3389/fimmu.2019.02292

Somamoto, T., Nakanishi, T., & Okamoto, N. (2002). Role of specific cell-mediated cytotoxicity

in protecting fish from viral infections. *Virology*, *297*(1), 120–127. doi:

10.1006/viro.2002.1486

Soonthonsrima, T., Wangman, P., Chaivisuthangkura, P., Pengsuk, C., Sithigorngul, P., &

Longyant, S. (2019). Generation of mouse monoclonal antibodies specific to tilapia

immunoglobulin using fish immunoglobulin/BSA complex for monitoring of the immune

response in Nile tilapia (*Oreochromis niloticus*). *Aquaculture Research*, *50*(1), 277–283.

doi: 10.1111/are.13894

Surachetpong, W., Janetanakit, T., Nonthabenjawan, N., Tattiyapong, P., Sirikanchana, K., &

Amonsin, A. (2017). Outbreaks of Tilapia Lake Virus Infection. Emerging Infectious

Diseases, *23*(6), 1031–1033. doi: 10.3201/eid2306.161278

Taengphu, S., Sangsuriya, P., Phiwsaiya, K., & Pratim, P …Senapin, S (2020). Genetic diversity

of tilapia lake virus genome segment 1 from 2011 to 2019 and a newly validated semi-

nested RT-PCR method. *Aquaculture, 526*. doi: 10.1016/j.aquaculture.2020.735423

Tang, L., Kang, H., Duan, K., Guo, M., Lian, G., Wu, Y., … Liu, M. (2016). Effects of Three

Types of Inactivation Agents on the Antibody Response and Immune Protection of

Inactivated IHNV Vaccine in Rainbow Trout. *Viral Immunology*, *29*(7), 430–435.

doi:org/10.1089/vim.2016.0035

Tattiyapong, P., Dachavichitlead, W., & Surachetpong, W. (2017). Experimental infection of

Tilapia Lake Virus (TiLV) in Nile tilapia (*Oreochromis niloticus*) and red tilapia

(Oreochromis spp.). *Veterinary Microbiology*, *207*, 170–177. doi:

org/10.1016/j.vetmic.2017.06.014

Tattiyapong, P., Dechavichitlead, W., Waltzek, T. B., & Surachetpong, W. (2020). Tilapia

develop protective immunity including a humoral response following exposure to tilapia

lake virus. *Fish and Shellfish Immunology*, *106*, 666–674. doi: 10.1016/j.fsi.2020.08.031

Thu Lan, N. G., Salin, K. R., Longyant, S., Senapin, S., & Dong, H. T. (2021). Systemic and

mucosal antibody response of freshwater cultured Asian seabass (*Lates calcarifer*) to

monovalent and bivalent vaccines against *Streptococcus agalactiae* and *Streptococcus*

iniae. *Fish and Shellfish Immunology*, *108*, 7–13. doi: org/10.1016/j.fsi.2020.11.014

Velázquez, J., Acosta, J., Lugo, J. M., Reyes, E., Herrera, F., González, O., … Estrada, M. P.

(2018). Discovery of immunoglobulin T in Nile tilapia (*Oreochromis niloticus*): A potential

molecular marker to understand mucosal immunity in this species. *Developmental and*

Comparative Immunology, *88*, 124–136. doi: org/10.1016/j.dci.2018.07.013

Wang, B., Wang, P., Wu, Z. H., Lu, Y. S., Wang, Z. L., & Jian, J. C. (2016). Molecular cloning

and expression analysis of IgD in nile tilapia (*Oreochromis niloticus*) in response to

Streptococcus agalactiae stimulus. *International Journal of Molecular Sciences*, *17*(3). doi:

org/10.3390/ijms17030348

Yamkasem, J., Tattiyapong, P., Kamlungdee, A., & Surachetpong, W. (2019). Evidence of

potential vertical transmission of tilapia lake virus. *Journal of Fish Diseases, 42* (9), 1123-

1331. doi: org/10.1111/jfd.13050

Zapata, A., Diez, B., Cejalvo, T., Gutiérrez-De Frías, C., & Cortés, A. (2006). Ontogeny of the

immune system of fish. *Fish and Shellfish Immunology*, *20*(2), 126–136. doi:

org/10.1016/j.fsi.2004.09.005

Zeng, W., Wang, Y., Hu, H., Wang, Q., Bergmann, S. M., Wang, Y., … Li, Y. (2021a). Cell

Culture-Derived Tilapia Lake Virus-Inactivated Vaccine Containing Montanide Adjuvant

Provides High Protection against Viral Challenge for Tilapia. *Vaccines*, *9*(2), 86. doi:

org/10.3390/vaccines9020086

Zeng, W., Wang, Y., Chen, X., Wang, Q., Bergmann, S. M., Yang, Y., … Lan, W. (2021b).

Potency and efficacy of VP20-based vaccine against tilapia lake virus using different prime-

boost vaccination regimens in tilapia. *Aquaculture*, *539*. doi:

10.1016/j.aquaculture.2021.736654

Zhang, Y. A., Salinas, I., & Oriol Sunyer, J. (2011). Recent findings on the structure and

function of teleost IgT. *Fish and Shellfish Immunology*, *31*(5), 627–634. doi:

org/10.1016/j.fsi.2011.03.021

Tables and figures

Table 1. Summary of TiLV vaccines and their efficacy

Vaccine type	Description	Country of origin	Dose ($TCID_{50}$ fish^{-1})	Administration route	Adjuvant	% RPS or survival rate	Reference
Live attenuated	17 and 20 passages (P17 & P20) on cell culture	Israel	1.2×10^7 (P17) and 8.9×10^6 (P20)	IP	No	> 50[a]	Bacharach et al., 2016c
DNA vaccine	Segment 8 (VP20)	China	5 µg	IM	No	50[b]	Zeng et al., 2021b
Recombinant vaccine			20 µg	IM	M402 (China)	52.5[b]	
DNA + recombinant vaccine			5 µg of DNA vaccine (prime) + 20 µg of recombinant vaccine (booster)	IM	M402 for booster	72.5[b]	
Inactivated	β-propiolactone	China	10^5; 10^6; 10^7	IM	Montanide IMS 1312 VG (Seppic, France)	32.1[a] - 85.7[a]	Zeng et al., 2021a
				IM	No	14.3[a] - 42.9[a]	
	Heat	Thailand	1.8×10^6	IP	No	71.3[a]	This study
	Formalin		1.8×10^6	IP	No	79.6[a]	

589 IP, intraperitoneal injection; IM: intramuscular injection; a: Relative percentage survival (RPS); b: survival rate

590   **Table 2.** Details of primers used for immune-related gene expression in this study.

Gene	Oligo sequences	Annealing temperature (^{o}C)	Product size (bp)	Reference
EF-1α	F-5'-CTACAGCCAGGCTCGTTTCG-3'	56	139	(Velázquez et al., 2018)
	R-5'-CTTGTCACTGGTCTCCAGCA-3'			
IgM	F-5'-GGATGACGAGGAAGCAGACT-3'	53	122	(Velázquez et al., 2018)
	R-5'-CATCATCCCTTTGCCACTGG-3'			
IgT	F-5'-TGACCAGAAATGGCGAAGTCTG-3'	53	163	(Velázquez et al., 2018)
	R-5'-GTTATAGTCACATTCTTTAGAATTACC-3'			
IgD	F-5'-AACACCACCCTGTCCCTGAAT-3'	61	127	(Wang et al., 2016)
	R-5'-GGGTGAAAACCACATTCCAAC-3'			
CD4	F-5'- GCTCCAGTGTGACGTGAAA-3'	61	106	XM_025911776.1, designed in this study
	R-5'- TACAGGTTTGAGTTGAGCTG-3'			
CD8α	F-5'- GCTGGTAGCTCTGGCCTTT-3'	49.5	91	XM_005450353.3*, designed in this study
	R'-5'-TGTGATGGTGTGGGCATCTC-3'			

591   *homolog (98% nucleotide sequence identity) of *Oreochromis aureus CD8α* (XM_031747820.2).

593   **Table 3.** Details of experimental groups and challenge results

Treatment	Administration route	No of fish challenged	Primary vaccination ($TCID_{50}$ fish^{-1})	Booster vaccination ($TCID_{50}$ fish^{-1})	Challenge ($TCID_{50}$ fish^{-1})	Survival rate (%)	RPS (%)	Significant level
			Day 0	Day 21	Day 28			
Control (L15 media)	IP	16 ($\times$ 2 rep.)	0	0	9 x 10^5	28.13 ± 30.9	NA	
HKV	IP	16 ($\times$ 2 rep.)	1.8×10^6	1.8×10^6	9 x 10^5	81.3 ± 0.0	71.3	*p<0.0001*
FKV	IP	15 ($\times$ 2 rep.)	1.8×10^6	1.8×10^6	9 x 10^5	86.3 ± 0.0	79.6	*p<0.0001*

594   HKV, heat-killed vaccine group; FKV, formalin killed vaccine group; rep, replicate; NA, not applicable; IP, intraperitoneal injection

Figure 1. Average percent survival of heat-killed and formaldehyde-killed vaccinated groups (HKV vs. FKV) compared to the non-vaccinated group (Control) during 21 days post challenge. with TiLV (strain TH-2018-K). Statistical analysis of cumulative survival between both vaccinated groups and the control were analyzed using Kaplan-Meier curve with log-rank test (*p < 0.0001*).

Figure 2. Fold change in gene expressions between non-vaccinated and vaccinated fish at 14, 21

and 28 days post vaccination. Data are presented as the mean ± SE (n=6). Control, non-vaccinated

group; HKV, heat-killed vaccine group; FKV, formalin-killed vaccine group. Asterisks show

significant levels between groups. * *p<0.05*, ** *p< 0.01*, *** *p<0.001*

Figure 3. Optical Density (OD) at 540 nm for IgM levels against TiLV in fish sera (diluted 1:512) (A) and mucus (undiluted) (B). Data are presented as the mean ± SE (n=6). Control, non-vaccinated group; HKV, heat-killed vaccine group; FKV, formalin-killed vaccine group. Asterisks indicate significant levels between groups. * *p<0.05*, ** *p< 0.01*, *** *p<0.001*, **** *p<0.0001*.