Physical basis for distinct basal and mechanically-gated activity of the human K$^+$ channel TRAAK

Robert A. Rietmeijer[1,2,3,4,#], Ben Sorum[1,2,4,#], Baobin Li[1,2,4] & Stephen G. Brohawn[1,2,4,*]

Affiliations
[1]Department of Molecular & Cell Biology, University of California Berkeley, Berkeley, California 94720, USA.
[2]Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley, California 94720, USA.
[3]Biophysics Graduate Program, University of California Berkeley, Berkeley, California 94720, USA.
[4]California Institute for Quantitative Biology (QB3), University of California, Berkeley, CA 94720, USA.

[#]equal contributions
*Correspondence and lead contact: brohawn@berkeley.edu

Summary

        TRAAK is a mechanosensitive two-pore domain K$^+$ (K2P) channel localized to nodes of Ranvier in myelinated neurons. TRAAK deletion in mice results in mechanical and thermal allodynia and gain-of-function mutations cause the human neurodevelopmental disorder FHEIG. TRAAK displays basal and stimulus-gated activities typical of K2Ps, but the mechanistic and structural differences between these modes are unknown. Here, we demonstrate that basal and mechanically-gated openings are distinguished by their conductance, kinetics, and structure. Basal openings are low conductance, short duration, and occur through a channel with an interior cavity exposed to the surrounding membrane. Mechanically-gated openings are high conductance, long duration, and occur through a channel that is sealed to the surrounding membrane. Our results explain how dual modes of activity are produced by a single ion channel and provide a basis for the development of state-selective pharmacology with the potential to treat disease.

Introduction

Two-pore domain K^+ (K2P) channels display low basal activity (also referred to as background, leak-type, flux-gated, or resting activity) that is important for setting and maintaining the resting membrane potential in many cells (Enyedi and Czirják, 2010; Renigunta et al., 2015). In addition, K2Ps are further activated by diverse physical and chemical stimuli including membrane tension, temperature, voltage, pH, signaling lipids, and calcium to regulate cellular electrical excitability. K2Ps are physiologically relevant targets of volatile anesthetics and antidepressants and their dysregulation has been implicated in pathologies including migraine, depression, pulmonary hypertension, atrial fibrillation, diabetes, Birk-Barel syndrome, and FHEIG syndrome (Facial Dysmorphism, Hypertrichosis, Epilepsy, Intellectual disability, and Gingival outgrowth) (Enyedi and Czirják, 2010; Barel et al., 2008; Bauer et al., 2018; Heurteaux et al., 2006; Lafrenière et al., 2010; Ma et al., 2013; Royal et al., 2019; Schmidt et al., 2015; Vierra et al., 2015)

TRAAK is a mechanosensitive K2P expressed in the nervous system of jawed vertebrates and localized to nodes of Ranvier, the gaps in myelinated axons where the action potential is regenerated during saltatory conduction. TRAAK accounts for ~25% of nodal background K^+ conductance and maintains the resting membrane potential and voltage gated-Na^+ channel availability required for high frequency spiking (Brohawn et al., 2019; Kanda et al., 2019; Maingret et al., 1999). Altering TRAAK activity in animals has physiological consequences; TRAAK knockout mice display mechanical and thermal allodynia and mechanical hyperalgesia while gain-of-function mutations in humans (TRAAK$_{A198E}$ and TRAAK$_{A270P}$) underlie the neurodevelopmental disorder FHEIG (Bauer et al., 2018; Noël et al., 2009).

TRAAK basal activity can be activated ~100-fold by increased membrane tension (Brohawn et al., 2014a, 2014b; Sorum et al., 2021). A model for mechanical gating primarily involves conformational changes in transmembrane helix 4 (TM4) (Brohawn et al., 2014b). In this model, nonconductive (closed) TRAAK adopts a TM4 "down" conformation and conductive (open) TRAAK adopts a TM4 "up" conformation. When TM4 is down, a lipid acyl chain can enter the channel cavity below the selectivity filter through membrane facing openings and sterically block the conduction path. Movement of TM4 up seals the membrane facing openings, prevents lipid access, and permits ion conduction. Structures of these two conformations provide a biophysical explanation for mechanosensitivity: shape changes upon opening, including an expansion in cross sectional area and increase in cylindricity, are energetically favored by membrane tension.

Notably, single channel recordings of TRAAK, and the closely related mechanosensitive K2P channels TREK1 and TREK2, suggest the presence of multiple distinct open and closed states (Bang et al., 2000; Kang et al., 2005; Maingret et al., 1999; Patel et al., 1998; Sorum et al., 2021; Xian Tao Li et al., 2006; Clausen et al. 2020). In addition, alternative models for gating mechanosensitive K2Ps posit closures involving C-type gating at the selectivity filter (Chatelain et al., 2012; González et al., 2013; Li et al., 2020; Lolicato et al., 2020; Schewe et al., 2016), closures involving dewetting of the channel cavity (Aryal et al., 2014, 2015), and openings that do not involve upward movement of TM4 (Lolicato et al., 2014). For TRAAK, and other K2Ps, basally open and stimulus-activated currents display different voltage dependence (Schewe et al., 2016), consistent with structurally distinct open states. Here, we show basal and mechanically-gated

TRAAK open states are distinct in their conductance, kinetics, and structure and present a model for channel gating that associates conformational and functional states.

Results and Discussion

Basal and mechanically-gated TRAAK openings are physically distinct

A macroscopic recording from a patch containing wild-type TRAAK (TRAAK$_{WT}$) illustrates basal activity and robust activation by membrane tension (generated by negative pressure applied through the patch pipette) that are characteristic of the channel (Fig. 1A,B). We performed single channel recordings to determine whether basally open and mechanically activated TRAAK have different biophysical characteristics consistent with distinct channel open states. By varying applied pressure over approximately an hour of single channel records, we sampled a wide range of TRAAK$_{WT}$ activity ($0.01 \leq$ open probability (P_O) ≤ 0.98) (Fig. 1C-F, Table S1). We analyzed records in three bins according to P_O (Fig. 1C). We found that in addition to a simple increase in P_O, mechanically activated TRAAK openings are distinct from basal openings in two ways.

First, mechanical force favors higher conductance openings (Figs. 1G-I, S1). Two conductance states are observed in TRAAK$_{WT}$: a low conductance opening with a unitary current of ~1 pA and a high conductance opening with a unitary current of ~2 pA. In unstretched patches at low P_O ($0.01 \leq P_O \leq 0.07$), the two conductance states are observed at roughly similar frequency (Fig. 1G,M). At intermediate P_O (mid P_O, $0.38 \leq P_O \leq 0.69$), more high conductance openings are observed (Fig. 1H,M). In records mechanically activated to high P_O ($0.93 \leq P_O \leq 0.98$), almost all openings are high conductance (Fig. 1I,M). Notably, unitary currents of these openings do not change significantly as a function of P_O, but the relative proportion of each conductance state does (Fig. 1G-J,M).

Second, mechanical force favors longer duration openings. Two kinetically distinct open states are observed in TRAAK$_{WT}$ records: a short duration (~1 ms) opening and a long duration (~3 ms or longer) opening (Figs. 1K, S2). In unstretched patches at low P_O, almost all openings are short duration (Fig. 1M). In contrast, when mechanically activated to high P_O, almost all openings are long duration. At intermediate P_O, both short and long duration openings are observed. Open duration and unitary current are positively correlated: low conductance openings have a narrow duration distribution around 1 ms while high conductance openings have a wider duration distribution ≥2 ms, with longer openings observed at higher P_O (Fig. 1M).

In addition, mechanical force favors shorter duration closures. TRAAK$_{WT}$ accesses two kinetically distinct closed states: a short duration (~0.5 ms) closure and a long duration (~3 ms or longer) closure (Figs. 1L, S3). Short duration closures of comparable length are observed in all records. However, long duration closures are only observed in low and intermediate P_O records and their prevalence and duration decrease as P_O increases (Figs. 1L, S3).

Single channel TRAAK$_{WT}$ records can be fit with a linear kinetic model relating the two closed states (long duration (C1) and short duration (C2)) and two open states (short duration/low conductance (O1) and long duration/high conductance (O2)) (Fig. 1N, Table S1)(Sorum et al., 2021). The distribution of states and the duration of C1 and

O2 are dependent on membrane tension. When mechanically activated to high P_O, long duration closures and short duration/low conductance openings are almost never observed, so data can be fit with a two-state equilibrium model between C2 and O2 states (Figs. 1I,K-M, S1-3, Table S1). At intermediate P_O, the duration of long C1 closures is shorter than those at low P_O and the duration of O2 openings is shorter than those at high P_O (Figs. 1H, K-M, S1-3, Table S1). We conclude that mechanically activated openings correspond to the O2 state while basal openings correspond to the O1 state. We speculate weak mechanical activation by resting membrane tension in the patch results in low P_O O2 openings with a conductance distinct from O1 openings (Brohawn et al., 2014b; Opsahl and Webb, 1994). However, the duration of low P_O O1 and O2 openings is similarly short and therefore indistinguishable kinetically (Fig. 1D,K).

We then asked whether we could identify the structural basis for these two distinct TRAAK open states. We reasoned that gain-of-function mutations could differentially stabilize particular open states and investigated three point mutants reported to dramatically increase channel activity: TRAAK$_{A198E}$, TRAAK$_{A270P}$, and TRAAK$_{G158D}$. TRAAK$_{A198E}$ and TRAAK$_{A270P}$ were recently identified in human patients with the neurological and developmental disorder FHEIG while TRAAK$_{G158D}$ was identified as a pan-K2P gain-of-function mutation (Bauer et al., 2018; Ben Soussia et al., 2019).

Mechanical force and FHEIG mutations promote a TM4 up open state

We first consider the FHEIG mutations TRAAK$_{A198E}$ and TRAAK$_{A270P}$ (Figs. 2A, 3A). Macroscopic recordings of TRAAK$_{A198E}$ or TRAAK$_{A270P}$ channels display currents that are only subtly activated by mechanical force, consistent with the initial characterization of these mutations (Figs. 2B,C, 3B,C) (Bauer et al., 2018). Single channel records reveal a simple explanation for reduced mechanical activation of TRAAK$_{A198E}$ and TRAAK$_{A270P}$ relative to TRAAK$_{WT}$. TRAAK$_{A198E}$ and TRAAK$_{A270P}$ open probability is nearly one at rest and mechanical force does not change channel conductance, so stimulation can only slightly increase channel activity (Figs. 2D-G, 3D-G).

TRAAK$_{A198E}$ and TRAAK$_{A270P}$ single channel behavior closely matches TRAAK$_{WT}$ when mechanically activated to high P_O. TRAAK$_{A198E}$ and TRAAK$_{A270P}$ openings are nearly all high conductance (with a unitary current ~2 pA) and long duration (~3 ms or longer), while closures are short duration (~0.5 ms) (Figs. 2F-J, 3F-J). The modest mechanical activation of TRAAK$_{A198E}$ and TRAAK$_{A270P}$ is due to an increase in long duration openings without significantly changing closed duration.

We determined crystal structures of TRAAK$_{A198E}$ and TRAAK$_{A270P}$ in the presence of K$^+$ and in complex with a mouse monoclonal antibody Fab fragment to 2.5 Å and 3.0 Å resolution, respectively (Figs. 2A, 3A, S4). As previously reported for this crystal form, the conformation of one side of the TRAAK dimer is restricted because it is involved in forming contacts that propagate the lattice (Fig. S5). We therefore focus our discussion on the opposing side of the TRAAK dimer (including TM4 from protomer B), which is conformationally unrestricted. In the same crystallization conditions, TRAAK$_{WT}$ adopts a TM4 down nonconductive conformation; the membrane-facing lateral opening above TM4 permits hydrophobic acyl chains to access the channel cavity and block ion passage (Fig. 2K) (Brohawn et al., 2014b). TRAAK$_{A198E}$ and TRAAK$_{A270P}$, however, adopt TM4 up conformations (Figs. 2L, 3K). Movement of TM4 up seals the membrane-facing

lateral openings, preventing acyl chain access to the cavity and creating an unobstructed path for ion movement through the channel. This TM4 up conformation is similar to a TRAAK$_{WT}$ TM4 up conductive conformation captured in the presence of a small molecule activator trichloroethanol (Fig. S7B,E) (Brohawn et al., 2014b). As observed in the TRAAK$_{WT}$ TM4 up conductive structure, density consistent with a K$^+$ ion is present in the cavities of TRAAK$_{A198E}$ and TRAAK$_{A270P}$ (Figs. 3L, S6A).

To confirm that the TRAAK$_{A198E}$ TM4 up structure represents an open state, we determined a second structure in the presence of Tl$^+$ to 3.0 Å resolution and used anomalous diffraction of Tl$^+$ ions to unambiguously identify ion binding sites. Indeed, a Tl$^+$ ion is identified in the channel cavity in addition to four sites in the selectivity filter (S1-S4) and one on the extracellular mouth of the pore (S0), consistent with a conductive conformation (Fig. 2M).

How do TRAAK$_{A198E}$ and TRAAK$_{A270P}$ promote TM4 up open states? Comparing TRAAK$_{A198E}$ and TRAAK$_{WT}$ suggests two mechanisms in this mutant (Figs. 2N, S6D&E). The first is a steric relay from A198E through F201 and Y271 that favors TM4 up. In TRAAK$_{WT}$, the TM4 down conformation involves movement of Y271 1 Å towards the cytoplasm relative to TM4 up. This requires the intracellular TM2-TM3 linkage to rotate 15° away from TM4 to prevent a clash between Y271 on TM4 and F201 on TM3. In TRAAK$_{A198E}$, TM2-TM3 is similarly rotated out, but A198E pushes F201 closer to TM4 where it would clash with Y271 in a TM4 down conformation. A second possibility is that A198E causes local thinning of the lipid bilayer to favor TM4 up. While A198 is predicted to be embedded in the membrane inner leaflet facing hydrophobic residues on TM4 down, the negatively charged carboxylic acid of A198E likely favors hydration of this pocket, promoting TM4 up to maintain nonpolar interactions between its hydrophobic residues and lipid acyl chains (Brooks, B. R. and Brooks, III, C. L. and Mackerell, Jr. et al., 2009; Jo et al., 2008).

In TRAAK$_{A270P}$, the introduced proline kinks TM4 to closely approximate the TRAAK$_{WT}$ TM4 up conformation (Fig 3M). In TRAAK$_{WT}$, movement of TM4 up depends on a hinge at a conserved glycine G268. The A270P mutation instead disrupts backbone hydrogen bonding in TM4 and creates a ~27° bend and ~9.5 Å upward movement of TM4 relative to TRAAK$_{WT}$ TM4 down.

We conclude that the TRAAK$_{A198E}$, TRAAK$_{A270P}$, and TRAAK$_{WT}$ TM4 up structures correspond to the high conductance/long duration mechanically activated open state O2. The following lines of evidence support this assignment: (i) TRAAK$_{A198E}$ and TRAAK$_{A270P}$ have an open probability ≥ 0.9 (Figs. 2F,H, 3F,H), (ii) the structures are similar to a TRAAK$_{WT}$ mechanically activated conductive conformation (Figs. 2K,L, 3K), (iii) the structures show unobstructed conduction paths with ions in the channel cavities (Figs. 2M, 3L, S6A), (iv) the predominant TRAAK$_{A198E}$ and TRAAK$_{A270P}$ unitary current is indistinguishable from TRAAK$_{WT}$ O2 (Figs. 1I, 2G, 3G), and (v) the mean duration of the TRAAK$_{A198E}$ and TRAAK$_{A270P}$ open states are indistinguishable from TRAAK$_{WT}$ O2 (Figs. 1K, 2I, 3I, S2).

A TM4 down open state promoted by a pan-K2P activating mutation underlies basal activity

We next consider TRAAK$_{G158D}$ (Fig. 4A). In macroscopic recordings, mechanical force activates TRAAK$_{G158D}$ ~2-fold; more than TRAAK$_{A198E}$ and TRAAK$_{A270P}$, but less

than $TRAAK_{WT}$ (Fig. 4B-D). Single channel analysis shows that, similar to $TRAAK_{WT}$, mechanical activation of $TRAAK_{G158D}$ involves an increase in both open probability and conductance (Fig. 4E-H). However, the resting open probability of $TRAAK_{G158D}$ (P_O ~0.7) is much higher than $TRAAK_{WT}$ (P_O ~0.04) such that mechanical stimulation can only increase $TRAAK_{G158D}$ P_O ~1.25 fold (Fig. 4E,F). Basal openings of $TRAAK_{G158D}$ are low conductance with a unitary current ~1 pA while mechanically activated openings are higher conductance with a unitary current ~2 pA (Figs. 4G-I, S1).

Kinetic analysis shows that $TRAAK_{G158D}$ closely resembles $TRAAK_{WT}$ except that the long duration closed state C1 is absent in the mutant (Figs. 4J, S2, S3). $TRAAK_{G158D}$ accesses only a single short closed state (Fig. 4K). Like $TRAAK_{WT}$, $TRAAK_{G158D}$ has two open states, one short duration/low conductance (~1 ms / 1 pA) and one long duration/high conductance (~3 ms or longer / 2 pA) (Fig. 4I,J). Mechanical stimulation favors longer duration/high conductance openings. (Figs. 4L, S1, S2).

We determined the crystal structure of $TRAAK_{G158D}$ to 3.0 Å resolution in the presence of K^+ (Fig. 4A). $TRAAK_{G158D}$ adopts a TM4 down conformation like $TRAAK_{WT}$ crystallized under the same conditions (Fig. 4M). However, density in the $TRAAK_{G158D}$ cavity is consistent with a K^+ ion rather than the hydrophobic acyl chain present in the $TRAAK_{WT}$ TM4 down structure (Fig. 4N). This suggests that the $TRAAK_{G158D}$ structure represents a TM4 down open state and is consistent with the high resting open probability of the mutant channel.

$TRAAK_{G158D}$ likely promotes a TM4 down open state through a simple electrostatic consequence of the mutation (Fig. 4O). G158D projects into the channel cavity ~5 Å underneath the selectivity filter and increases hydrophilicity of the ion conduction path close to where acyl chains are observed in nonconductive TM4 down $TRAAK_{WT}$ structure. The electronegative cavity of $TRAAK_{G158D}$ is expected to disfavor acyl chain access and channel block, instead promoting ion occupancy in the cavity and a conductive open state.

We conclude that the TM4 down open state captured in the $TRAAK_{G158D}$ structure corresponds to the low conductance/short duration basal open state O1 in $TRAAK_{WT}$. The following data support this assignment: (i) $TRAAK_{G158D}$ has a high resting open probability (Fig. 4E), (ii) a K^+ ion rather than an acyl chain is observed in the $TRAAK_{G158D}$ cavity (Fig. 4N), (iii) unitary current of the $TRAAK_{G158D}$ basal open state is indistinguishable from $TRAAK_{WT}$ O1 (Fig. 4I), (iv) mean dwell time of the $TRAAK_{G158D}$ short duration opening is statistically indistinguishable from $TRAAK_{WT}$ O1 (Fig. 4J), and (v) like $TRAAK_{WT}$, the basal low conductance/short duration $TRAAK_{G158D}$ O1-like open state can be mechanically activated to a high conductance/long duration O2-like open state (Fig. 4F-J). Consistent with (v), $TRAAK_{G158D}$ can adopt a TM4 up-like conformation (observed on the side of the channel involved in forming crystal contacts) and $TRAAK_{G158D}$ TM4 up is expected to be energetically favored in the presence of membrane tension relative to $TRAAK_{G158D}$ TM4 down because it has an increased cross-sectional area (Fig. S7C). The lower conductance of TRAAK O1 openings may be due to reduced cavity polarity or a longer path length through the channel in a TM4-down conformation relative to the TM4-up conformation of O2 openings.

An integrated structural and functional model for TRAAK gating

We further conclude that the TM4 down lipid-blocked closed state observed in the TRAAK$_{WT}$ structures corresponds to the long duration closed state C1. Analogous long duration closures are not observed in in TRAAK$_{G158D}$, TRAAK$_{A198E,}$ or TRAAK$_{A270P}$ (Figs. 1L, 2J, 3J, 4K). In TRAAK$_{A198E}$ and TRAAK$_{A270P}$, this is likely because TM4s are favored in an up conformation that seals lateral membrane openings, preventing lipid access to the cavity (Figs. 2L, 3K). In TRAAK$_{G158D}$, this is likely because the mutation increases polarity of the channel cavity to disfavor occupancy of hydrophobic acyl chains (Fig. 4O).

Together, these data support a model for TRAAK gating shown in Figure 5 that maps structural conformations to the linear four-state kinetic model derived from single-channel records (Fig. 1N). Long duration closures (C1) correspond to the TM4 down, lipid blocked conformation captured in TRAAK$_{WT}$ structures(Brohawn et al., 2014b). Low conductance/short duration basal openings (O1) correspond to the TM4 down, conductive conformation captured in the TRAAK$_{G158D}$ structure (Fig. 4). High conductance/long duration mechanically activated openings (O2) correspond to the TM4 up, conductive conformation captured in structures of TRAAK$_{A198E}$, TRAAK$_{A270P}$, and TRAAK$_{WT}$ with activating small molecules (Fig. 2,3). Structures of two other gain-of-function mutations, TRAAK$_{G124I}$ and TRAAK$_{W262S}$, have been previously reported(Lolicato et al., 2014), although their open probability, single channel behavior, and cavity ion occupancy is not known. However, both adopt TM4 down-like conformations and may promote O1 like openings similarly to TRAAK$_{G158D}$.

TRAAK$_{WT}$ under basal conditions transitions primarily between C1, O1, and C2. Mechanical force favors the TM4 up open state O2 because it involves shape changes (relative to TM4 down) that are energetically favored in the presence of membrane tension, including an increase in cross-sectional area and cylindricity(Brohawn et al., 2014b). Mechanical force disfavors the TM4 down states C1 and O1 because the relatively smaller cross-sectional area and more wedge-shaped structure is disfavored in the presence of membrane tension. The gain of function mutations TRAAK$_{A198E}$ and TRAAK$_{A270P}$ mimic mechanically activated TRAAK$_{WT}$ by promoting a TM4 up conductive conformation. As a consequence, TRAAK$_{A198E}$ and TRAAK$_{A270P}$, like mechanically activated TRAAK$_{WT}$, primarily transition between O2 and C2 states. Gain of function mutant TRAAK$_{G158D}$ mimics basally open TRAAK$_{WT}$, promoting a conductive TM4 down conformation by disfavoring lipid entry and channel block at low membrane tension. As a consequence, TRAAK$_{G158D}$ primarily transitions between O1, C2, and O2 states. Like TRAAK$_{WT}$, mechanical stimulation of TRAAK$_{G158D}$ disfavors O1 to favor transitions between O2 and C2 states.

The short duration closure (C2) does not yet have a defined structural correlate. We observe C2-like closures of similar duration in all TRAAK variants analyzed (Figs. 1L, 2J, 3J, 4K), suggesting they may occur independently of TM4 position or cavity environment. One possibility is that C2 closures represent gating at the selectivity filter. While gating conformational changes at the selectivity filter have not been observed in TRAAK, they have been hypothesized from functional data and visualized in other K2Ps including the mechanosensitive TREK1 in low [K$^+$] and pH-gated TASK2 in physiological [K$^+$] and low pH (Li et al., 2020; Lolicato et al., 2020). Alternatively, C2 closures could correspond to cavity dewetting that has been proposed to occur based on molecular

dynamics simulations (Aryal et al., 2014, 2015) or other, yet to be visualized, conformational rearrangements.

Notably, some evidence suggests that the physical basis for distinct modes of TRAAK activity shown here could be shared among K2P channels. Differences between basal and stimulus-activated openings, including their kinetics and voltage dependence (Schewe et al., 2016), have been reported for other K2Ps, suggesting that physically distinct open states could similarly underlie different modes of channel activity (Bang et al., 2000; Gnatenco et al., 2002; Kang et al., 2005; Maingret et al., 1999; Clausen et al 2020; Sorum et al., 2021). Mutations analogous to TRAAK$_{G158D}$ were found to activate all K2P channels, with the extent of activation correlated with the polarity of the introduced amino acid change (Ben Soussia et al., 2019). Membrane-facing openings have been observed in TM4-down-like structures of TREK2 (Dong et al., 2015), TWIK1 (Miller and Long, 2012), TASK1 (Rödström et al., 2020), and TASK2 (Li et al., 2020) in addition to TRAAK(Brohawn et al., 2012) (Fig. S7D-L). Acyl chains that could block conduction were modeled in TWIK1 and TRAAK structures (Brohawn et al., 2012; Miller and Long, 2012), though their absence in other K2Ps may be due to limited resolution of, or partial occupancy in, structures determined to date. This raises the possibility that lipid-mediated block could be more common among K2Ps than currently appreciated. Like TRAAK, basal activity of other K2Ps could result from periodic unbinding of lipid from channel cavities and ion passage through a transiently unobstructed conduction path, while stimulus gating could involve conformational changes that prevent lipid entry into the channel entirely (Brohawn et al., 2012, 2014b; Dong et al., 2015; Li et al., 2020; Miller and Long, 2012; Rödström et al., 2020).

Methods

Electrophysiology

Full length hsTRAAK (Uniprot Q9NYG8-2) was codon optimized for eukaryotic expression, synthesized (Genewiz), and cloned into a modified pGEMHE vector using XhoI and EcoRI restriction sites such that the mRNA transcript encodes hsTRAAK 1-419 with an additional "SNS" at the C-terminus. Point mutants G158D, A198E, and A270P were generated by inverse PCR of the parent construct and verified by Sanger sequencing. cRNA was transcribed from linearized plasmids *in vitro* using T7 polymerase, and 0.1–10 ng cRNA was injected into *Xenopus laevis* oocytes extracted from anaesthetized frogs.

Currents were recorded at 25°C from inside-out patches excised from oocytes 1–5 days after mRNA injection. Pipette solution contained (in mM) 15 KCl, 135NaCl, 2MgCl$_2$, 10 HEPES, pH = 7.4 with NaOH. The solution in the bath contained 150 KCl, 2 MgCl$_2$, 10 HEPES, 1 EGTA, pH = 7.1 with KOH. All single channel records were made at a holding voltage of 0 mV and presented in physiological convention.

Currents were recorded using an Axopatch 200B Patch Clamp amplifier at a bandwidth of 1 kHz and digitized with an Axon Digidata 1550B at 50 to 500 kHz. To maintain the integrity of brief TRAAK dwell times, records were not further filtered unless explicitly stated. Baseline correction was performed manually based on the observable closed state for each recording; since there was no solution exchange in our experiments, no associated large drifts in baseline were observed. Portions of TRAAK$_{WT}$ records were assigned as low, mid, or high P$_O$ for analysis. Low P$_O$ was observed prior to pressure application, while mid and high P$_O$ behavior was observed after pressure application. Portions of mutant records during pressure application were analyzed separately from mutant records prior to pressure application. High P$_O$ TRAAK$_{WT}$ and mutant records with pressure were made with the highest achievable pressure prior to patch rupture. Single channel open-close transitions were idealized by half-amplitude threshold crossing and dwell time event lists were generated for each baseline corrected record. Dwell time histograms were generated from the dwell time event lists using custom built software (Sorum et al., 2015), while the fits for these histograms were generated in GraphPad Prism version 9.0 using single, double, or triple Gaussian fits.

For unitary current analysis, event lists for TRAAK$_{WT}$ low P$_O$, TRAAK$_{WT}$ mid P$_O$, TRAAK$_{G158D}$, and TRAAK$_{G158D}$ with pressure were first generated in Clampfit 10.7, applying a 0.8 pA event detection cutoff. Currents were filtered to 10 kHz and were analyzed with a custom script in Python 3.7 using the pyABF module (Harden, 2019). Each point in the filtered record was annotated as closed (C1/C2), open low conductance (O1), or open high conductance (O2) using the events list. To account for the sharp detection cutoff, openings were padded by an additional point on either side of the cutoff (these points corresponded to 100 us of ~0.5-1 pA current during openings and closings). Open only event lists were then generated by removing all closed points. Open event only lists were binned by 0.2 pA in current and 0.2 ms in time to generate bubble plots relating unitary current and mean open duration for TRAAK$_{WT}$ and TRAAK$_{G158D}$. Open only event lists were also used to generate open event current histograms to distinguish relatively rare low conductance openings from prevalent closed events. Mean unitary currents for TRAAK$_{WT}$ low P$_O$ and TRAAK$_{WT}$ mid P$_O$ were derived from Gaussian fits to these open event current histograms. Mean unitary currents for TRAAK$_{WT}$ high P$_O$ and all mutants were derived from Gaussian fits to square root total event current histograms in order to distinguish conductance states when one state is present at higher frequency.

TRAAK expression and purification

Human TRAAK UniProt Q9NYG8-2 was cloned for expression in *Pichia pastoris* cells as previously described (Brohawn et al., 2012) with modifications described here. The construct used for purification included an additional 26 amino acid N-terminal sequence compared to Q9NYG8-1 that improved heterologous expression. The final construct is C-terminally truncated by 119 amino acids, incorporates two mutations to remove N-linked glycosylation sites (N104Q/N108Q), and is expressed as a C-terminal PreScission protease-cleavable EGFP-10xHis fusion protein. As a result, there is an additional amino acid sequence of "SNSLEVLFQ" at the C-terminus of the final purified protein after protease cleavage. Recombinant *Pichia* were grown in a 4 liter fermenter and harvested 48-72h after induction with methanol.

30-120g of frozen *Pichia* cells expressing TRAAK were disrupted by milling (Retsch model MM301) 5 times for 3 minutes at 25 Hz. All subsequent purification steps were carried out at 4 ºC. Milled cells were resuspended in buffer A (in mM) 50 Tris pH 8.0, 150 KCl, 1 EDTA 0.1 mg/mL DNase1, 1 mg/mL pepstatin, 1 mg/mL leupeptin, 1 mg/mL aprotinin, 10 mg/mL soy trypsin inhibitor, 1 mM benzamidine, 100 µM AEBSF, 1 µM E-64, and 1 mM phenylmethysulfonyl fluoride added immediately before use) at a ratio of 1 g cell pellet per 4 mL lysis buffer and sonicated for 16 minutes with a duty cycle of 15 seconds of sonication per minute. The solution was ultracentrifuged at 150,000 xg for 1 hour at 4 ºC. Pellets were transferred to a Dounce homogenizer in buffer B (buffer A + 60 mM DM). Following homogenization, solutions were stirred for 3 hours at 4 ºC followed by centrifugation at 33,000g for 45 minutes. Anti-GFP nanobody resin (1 mg purified anti-GFP nanobody conjugated to 1 mL resin) was washed in Buffer B and added to the supernatant at a ratio of 1 mL resin / 15 g *Pichia* cells. The solution was gently stirred for 3 hours. Resin was collected on a column and washed in 15 column volumes (CV) of Buffer C (buffer A +6 mM DM+ 150 mM KCl), followed by 2 CV Buffer D (buffer A +6 mM DM). The resin was resuspended in ~3 CV of Buffer D with 1 mg purified Precission protease and gently rocked in column overnight. Cleaved TRAAK was eluted in ~4 CV of Buffer D, concentrated (50 kDa MWCO), and applied to a Superdex 200 SEC column (GE Healthcare) equilibrated in Buffer E (20 mM Tris pH 8.0, 150 mM KCl, 1 mM EDTA, 4 mM DM). Peak fractions were pooled and concentrated to 200-300 µL for incubation with Fab 13E9 and subsequently applied to a Superdex 200 column (GE Healthcare) equilibrated in Buffer E. Fab 13E9 was prepared as described previously(Brohawn et al., 2014b). TRAAK-Fab complexes were pooled and concentrated to 25-33 mg/mL for hanging drop crystallization.

For crystallization in Tl^+, all purification steps were conduction as with KCl, with two exceptions: KCl is substituted for KNO_3 for all buffers except Buffer E, where KCl is substituted with $TlNO_3$.

Protein Crystallization

Crystals were grown in drops of 0.125–0.250 µL protein added to an equal volume of reservoir, in hanging drops over a 100 µL reservoir at 4 ºC. Reservoir for each mutant in KCl was 50mM Tris pH 8.8, 64-200 mM $CaCl_2$, 27–33%(vol/vol) PEG400. Reservoir for each mutant in $TlNO_3$ was 50 mM Tris pH 8.8, 64-200 mM $Ca(NO3)_2$, 27–33%(vol/vol)PEG400. Lower Calcium concentrations were observed to correspond to increased nucleation and rapid growth of large, well-ordered crystals. Crystals grew to ~100 µm x 100 µm x 200 µm in 1–5 weeks. TRAAK A270P crystals rarely achieved sizes larger than 70 µm x 70 µm x 100 µm after 5 weeks.

For cryoprotection, an approximately equal volume of mother liquor supplemented to be 30%(vol/vol) PEG400 was added to one side of the drop and crystals were moved through this solution with a cryoloop before being plunged into

liquid nitrogen.

X-ray Data Collection, Model Building & Refinement

Data were collected at beamline 8.3.1 at the ALS, or 24-IDC & 24-IDE at the APS. Thallium-containing crystals were collected at 12680 eV, and Potassium-containing crystals were collected at 12663 eV. Were rotated 360 degrees, and data were collected in 0.20 degree wedges with a 0.25 second exposure time per wedge. Datasets were processed as individual wedges in XDS (Kabsch, 2010; Kabsch et al., 2010). Some datasets were truncated to eliminate redundant data displaying beam-induced damage prior to scaling with XSCALE, and merging with XDSCONV (Kabsch et al., 2010). Due to the anisotropy of the data, elliptically truncated resolution cutoffs generated by STARANISO (Tickle et al., 2018) were used to increased overall resolution of the maps. Structures were solved by molecular replacement using Phaser (McCoy et al., 2007) with an input model of nonconductive hsTRAAK for G158D (PDB ID 4WFF) and conductive hsTRAAK for A198E and A270P mutants (PDB ID 4WFE). Structure refinement was carried out in Coot (Emsley and Cowtan, 2004) and Refmac5 (Murshudov et al., 2011). Jelly body and automatically generated local NCS constraints were used throughout refinement. For the final round of refinement, three TLS groups per protein chain were incorporated. Molprobity was used to assess model geometry in later stages of refinement (Chen et al., 2010).

Ion occupancy in the conduction axis was determined by calculating either a model phased anomalous difference map for A198E crystals grown in Thallium, or a Polder map for G158D and A270P crystals grown Potassium. In all maps, density in the filter fits well to ions occupying the S1, 2, 3, and 4 positions. Similarly, densities above the filter (S0) are well fit with one or two ions. Density in the cavity below the filter is also fit with an ion. All ion occupancies are set to 1.

The angle and orientation of transmembrane helix 4 in each model was compared using UCSF Chimera (Pettersen et al., 2004). Cross-sectional area calculations were performed with CHARMM (Brooks, B. R. and Brooks, III, C. L. and Mackerell, Jr. et al., 2009; Jo et al., 2007, 2008). The area calculation used a surface calculated with 1.4 Å added to the van der Waals radii of protein atoms, and a probe radius of 3.5 Å to approximate lipid-accessible surface area. A water cylinder with a 6 Å radius was used to fill the cavity of TRAAK channels to exclude its contribution from the calculation. All calculations are made with channels containing symmetric protomers of the B chain, which does not make a contact in the crystal. Symmetric molecules were generated by rotation of the B protomer about the conduction axis, using pore helix and selectivity filter residues 119-133 and 228-242. Pore radii and pore-lining residue hydropathy were calculated with CHAP (Klesse et al., 2019) using a pathway lining residue margin value of 0.55 and a hydrophobicity kernel bandwidth of 0.45.

Author contributions

S.G.B., R.A.R., and B.S. conceived of the project, analyzed data, and wrote the manuscript. R.A.R. performed all aspects of the biochemistry and structural biology. B.S. performed all aspects of the electrophysiology. R.A.R. and B.S. generated constructs for recording. B.L. cultured the hybridoma for antibody purification. S.G.B. supervised the project and secured funding.

Data Availability

The X-ray crystallographic coordinates and structure factors for $TRAAK_{A198E}$ in K^+ (7LJ5), $TRAAK_{A198E}$ in Tl^+ (7LJA), $TRAAK_{A270P}$ in K^+ (7LJ4), and $TRAAK_{G158D}$ (7LJB) are available at the Protein Data Bank.

Acknowledgements

We thank R. MacKinnon for the 13E9 anti-human TRAAK hybridoma line. We thank staff at ALS beamline 8.3.1, especially J. Holton and G. Meigs, at APS beamline 24-IDC/E, especially I. Kourinov, for assistance at the synchrotron. We thank all members of the Brohawn laboratory for discussions. SGB is a New York Stem Cell Foundation-Robertson Neuroscience Investigator. This work was supported by the New York Stem Cell Foundation, NIGMS grant DP2GM123496, a McKnight Foundation Scholar Award, a Klingenstein-Simons Foundation Fellowship Award, a Sloan Research Fellowship, and a Rose Hill Innovator Award to SGB.

Declaration of Interests

The authors declare no competing interests.

References

Aryal, P., Abd-Wahab, F., Bucci, G., Sansom, M.S.P., and Tucker, S.J. (2014). A hydrophobic barrier deep within the inner pore of the TWIK-1 K2P potassium channel. Nat. Commun. 10.1038/ncomms5377.

Aryal, P., Sansom, M.S.P., and Tucker, S.J. (2015). Hydrophobic gating in ion channels. J. Mol. Biol. 10.1016/j.jmb.2014.07.030.

Bang, H., Kim, Y., and Kim, D. (2000). TREK-2, a new member of the mechanosensitive tandem-pore K+ channel family. J. Biol. Chem. 10.1074/jbc.M000445200.

Barel, O., Shalev, S.A., Ofir, R., Cohen, A., Zlotogora, J., Shorer, Z., Mazor, G., Finer, G., Khateeb, S., Zilberberg, N., et al. (2008). Maternally Inherited Birk Barel Mental Retardation Dysmorphism Syndrome Caused by a Mutation in the Genomically Imprinted Potassium Channel KCNK9. Am. J. Hum. Genet. 10.1016/j.ajhg.2008.07.010.

Bauer, C.K., Calligari, P., Radio, F.C., Caputo, V., Dentici, M.L., Falah, N., High, F., Pantaleoni, F., Barresi, S., Ciolfi, A., et al. (2018). Mutations in KCNK4 that Affect Gating Cause a Recognizable Neurodevelopmental Syndrome. Am. J. Hum. Genet. 10.1016/j.ajhg.2018.09.001.

Brohawn, S.G., del Marmol, J., and MacKinnon, R. (2012). Crystal Structure of the Human K2P TRAAK, a Lipid- and Mechano-Sensitive K+ Ion Channel. Science (80-.). *335*, 436–441, 10.1126/science.1213808.

Brohawn, S.G., Su, Z., and MacKinnon, R. (2014a). Mechanosensitivity is mediated directly by the lipid membrane in TRAAK and TREK1 K+ channels. Proc. Natl. Acad. Sci. U. S. A. *111*, 3614–3619, 10.1073/pnas.1320768111.

Brohawn, S.G., Campbell, E.B., and MacKinnon, R. (2014b). Physical mechanism for gating and mechanosensitivity of the human TRAAK K+ channel. Nature *516*, 126–130, 10.1038/nature14013.

Brohawn, S.G., Wang, W., Handler, A., Campbell, E.B., Schwarz, J.R., and MacKinnon, R. (2019). The mechanosensitive ion channel traak is localized to the mammalian node of ranvier. Elife 10.7554/eLife.50403.

Brooks, B. R. and Brooks, III, C. L. and Mackerell, Jr., A.D. and, Nilsson, L. and Petrella, R. J. and Roux, B. and Won, Y. and A., G. and Bartels, C. and Boresch, S. and Caflisch, A. and Caves, L. and, Cui, Q. and Dinner, A. R. and Feig, M. and Fischer, S. and Gao, J. and, Hodoscek, M. and Im, W. and Kuczera, K. and Lazaridis, T. and Ma, J., and Ovchinnikov, V. and Paci, E. and Pastor, R. W. and Post, C.B. and, Pu, J. Z. and Schaefer, M. and Tidor, B. and Venable, R.M. and, and Woodcock, H. L. and Wu, X. and Yang, W. and York, D. M. and Karplus, M. (2009). CHARMM: The Biomolecular Simulation Program. J. Comput. Chem. 10.1002/jcc.20945

Chatelain, F.C., Bichet, D., Douguet, D., Feliciangeli, S., Bendahhou, S., Reichold, M., Warth, R., Barhanin, J., and Lesage, F. (2012). TWIK1, a unique background channel with variable ion selectivity. Proc. Natl. Acad. Sci. U. S. A. 10.1073/pnas.1201132109.

Chen, V.B., Arendall, W.B., Headd, J.J., Keedy, D.A., Immormino, R.M., Kapral, G.J., Murray, L.W., Richardson, J.S., and Richardson, D.C. (2010). MolProbity: All-atom structure validation for macromolecular crystallography. Acta Crystallogr. Sect. D Biol. Crystallogr. 10.1107/S0907444909042073.

Clausen, M.V., Ulstrup, J., Poulsen, H. and Nissen, P. (2020) The main activatory and tension-sensitive transitions occur within Prozac sensitive down-states of the potassium selective TREK-2 channel. bioRxiv https://doi.org/10.1101/2020.10.22.351205

Dong, Y.Y., Pike, A.C.W.W., Mackenzie, A., McClenaghan, C., Aryal, P., Dong, L., Quigley, A., Grieben, M., Goubin, S., Mukhopadhyay, S., et al. (2015). K2P channel gating mechanisms revealed by structures of TREK-2 and a complex with Prozac. Science (80-.). *347*, 1256–1259, 10.1126/science.1261512.

Emsley, P., and Cowtan, K. (2004). Coot: Model-building tools for molecular graphics. Acta Crystallogr. Sect. D Biol. Crystallogr. 10.1107/S0907444904019158.

Enyedi, P., and Czirják, G. (2010). Molecular background of leak K+ currents: two-pore domain potassium channels. Physiol. Rev. *90*, 559–605, 10.1152/physrev.00029.2009.

Gnatenco, C., Han, J., Snyder, A.K., and Kim, D. (2002). Functional expression of TREK-2 K+ channel in cultured rat brain astrocytes. Brain Res. 10.1016/S0006-8993(02)02261-8.

González, W., Zúniga, L., Cid, L.P., Arévalo, B., Niemeyer, M.I., and Sepúlveda, F. V. (2013). An extracellular ion pathway plays a central role in the cooperative gating of a K2P K+ channel by extracellular pH. J. Biol. Chem. 10.1074/jbc.M112.445528.

Harden, S. (2019). Analysis of electrophysiological recordings was performed with custom software written for this project using Python 3.7 and the pyABF module.

Heurteaux, C., Lucas, G., Guy, N., El Yacoubi, M., Thümmler, S., Peng, X.D., Noble, F., Blondeau, N., Widmann, C., Borsotto, M., et al. (2006). Deletion of the background potassium channel TREK-1 results in a depression-resistant phenotype. Nat. Neurosci. 10.1038/nn1749.

Jo, S., Kim, T., and Im, W. (2007). Automated builder and database of protein/membrane complexes for molecular dynamics simulations. PLoS One 10.1371/journal.pone.0000880.

Jo, S., Kim, T., Iyer, V.G., and Im, W. (2008). CHARMM-GUI: A web-based graphical user interface for CHARMM. J. Comput. Chem. 10.1002/jcc.20945.

Kabsch, W. (2010). Integration, scaling, space-group assignment and post-refinement. Acta Crystallogr. Sect. D Biol. Crystallogr. 10.1107/S0907444909047374.

Kabsch, W., T., B.A., K., D., A., K.P., K., D., S., M., G., R.R.B., P., E., S., F., K., W., et al. (2010). *XDS*. Acta Crystallogr. Sect. D Biol. Crystallogr. 10.1107/S0907444909047337.

Kanda, H., Ling, J., Tonomura, S., Noguchi, K., Matalon, S., and Gu, J.G. (2019). TREK-1 and TRAAK Are Principal K+ Channels at the Nodes of Ranvier for Rapid Action Potential Conduction on Mammalian Myelinated Afferent Nerves. Neuron 10.1016/j.neuron.2019.08.042.

Kang, D., Choe, C., and Kim, D. (2005). Thermosensitivity of the two-pore domain K + channels TREK-2 and TRAAK. J Physiol *5641*, 103–116, 10.1113/jphysiol.2004.081059.

Klesse, G., Rao, S., Sansom, M.S.P., and Tucker, S.J. (2019). CHAP: A Versatile Tool for the Structural and Functional Annotation of Ion Channel Pores. J. Mol. Biol. 10.1016/j.jmb.2019.06.003.

Lafrenière, R.G., Cader, M.Z., Poulin, J.F., Andres-Enguix, I., Simoneau, M., Gupta, N., Boisvert, K., Lafrenière, F., McLaughlan, S., Dubé, M.P., et al. (2010). A dominant-negative mutation in the TRESK potassium channel is linked to familial migraine with aura. Nat. Med. 10.1038/nm.2216.

Li, B., Rietmeijer, R.A., and Brohawn, S.G. (2020). Structural basis for pH gating of the two-pore domain K+ channel TASK2. Nature 10.1038/s41586-020-2770-2.

Lolicato, M., Riegelhaupt, P.M., Arrigoni, C., Clark, K.A., and Minor, D.L. (2014). Transmembrane helix straightening and buckling underlies activation of mechanosensitive and thermosensitive K_{2P} channels. Neuron *84*, 1198–1212, 10.1016/j.neuron.2014.11.017.

Lolicato, M., Natale, A.M., Abderemane-Ali, F., Crottès, D., Capponi, S., Duman, R., Wagner, A., Rosenberg, J.M., Grabe, M., and Minor, D.L. (2020). K2P channel C-type gating involves asymmetric selectivity filter order-disorder transitions. Sci. Adv. 10.1126/sciadv.abc9174.

Ma, L., Roman-Campos, D., Austin, E.D., Eyries, M., Sampson, K.S., Soubrier, F., Germain, M., Treġouët, D.A., Borczuk, A., Rosenzweig, E.B., et al. (2013). A novel channelopathy in pulmonary arterial hypertension. N. Engl. J. Med. 10.1056/NEJMoa1211097.

Maingret, F., Fosset, M., Lesage, F., Lazdunski, M., and Honoré, E. (1999). TRAAK is a mammalian neuronal mechano-gated K+ channel. J. Biol. Chem. *274*, 1381–1387, 10.1074/jbc.274.3.1381.

McCoy, A.J., Grosse-Kunstleve, R.W., Adams, P.D., Winn, M.D., Storoni, L.C., and Read, R.J. (2007). Phaser crystallographic software. J. Appl. Crystallogr. 10.1107/S0021889807021206.

Miller, A.N., and Long, S.B. (2012). Crystal structure of the human two-pore domain potassium channel K2P1. Science *335*, 432–436, 10.1126/science.1213274.

Murshudov, G.N., Skubák, P., Lebedev, A.A., Pannu, N.S., Steiner, R.A., Nicholls, R.A., Winn, M.D., Long, F., and Vagin, A.A. (2011). REFMAC5 for the refinement of macromolecular crystal structures. Acta Crystallogr. Sect. D Biol. Crystallogr. 10.1107/S0907444911001314.

Noël, J., Zimmermann, K., Busserolles, J., Deval, E., Alloui, A., Diochot, S., Guy, N., Borsotto, M., Reeh, P., Eschalier, A., et al. (2009). The mechano-activated K+ channels TRAAK and TREK-1 control both warm and cold perception. EMBO J. 10.1038/emboj.2009.57.

Opsahl, L.R., and Webb, W.W. (1994). Lipid-glass adhesion in giga-sealed patch-clamped membranes. Biophys J *66*, 75–79, 10.1016/S0006-3495(94)80752-0.

Patel, A.J., Honoré, E., Maingret, F., Lesage, F., Fink, M., Duprat, F., and Lazdunski, M. (1998). A mammalian two pore domain mechano-gated S-like K+ channel. EMBO J. 10.1093/emboj/17.15.4283.

Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera - A visualization system for exploratory research and analysis. J. Comput. Chem. 10.1002/jcc.20084.

Renigunta, V., Schlichthörl, G., and Daut, J. (2015). Much more than a leak: Structure and function of K2P-channels. Pflugers Arch. Eur. J. Physiol. 867–894, 10.1007/s00424-015-1703-7.

Rödström, K.E.J., Kiper, A.K., Zhang, W., Rinné, S., Pike, A.C.W., Goldstein, M., Conrad, L.J., Delbeck, M., Hahn, M.G., Meier, H., et al. (2020). A lower X-gate in TASK channels traps inhibitors within the vestibule. Nature 10.1038/s41586-020-2250-8.

Royal, P., Andres-Bilbe, A., Ávalos Prado, P., Verkest, C., Wdziekonski, B., Schaub, S., Baron, A., Lesage, F., Gasull, X., Levitz, J., et al. (2019). Migraine-Associated TRESK Mutations Increase Neuronal Excitability through Alternative Translation Initiation and Inhibition of TREK. Neuron 10.1016/j.neuron.2018.11.039.

Schewe, M., Nematian-Ardestani, E., Sun, H., Musinszki, M., Cordeiro, S., Bucci, G., De Groot, B.L., Tucker, S.J., Rapedius, M., and Baukrowitz, T. (2016). A Non-canonical Voltage-Sensing Mechanism Controls Gating in K2P K+ Channels. Cell *164*, 937–949, 10.1016/j.cell.2016.02.002.

Schmidt, C., Wiedmann, F., Voigt, N., Zhou, X.B., Heijman, J., Lang, S., Albert, V., Kallenberger, S., Ruhparwar, A., Szabó, G., et al. (2015). Upregulation of K2P 3.1 K + Current Causes Action Potential Shortening in Patients with Chronic Atrial Fibrillation. Circulation 10.1161/CIRCULATIONAHA.114.012657.

Sorum, B., Czégé, D., and Csanády, L. (2015). Timing of CFTR Pore Opening and Structure of Its Transition State. Cell 10.1016/j.cell.2015.09.052.

Sorum, B., Rietmeijer, R.A., Gopakumar, K., Adesnik, H., and Brohawn, S.G. (2021). Ultrasound activates mechanosensitive TRAAK K+ channels directly through the lipid membrane. Proceedings of the National Acadamy of Sciences https://doi.org/10.1073/pnas.2006980118.

Ben Soussia, I., El Mouridi, S., Kang, D., Leclercq-Blondel, A., Khoubza, L., Tardy, P., Zariohi, N., Gendrel, M., Lesage, F., Kim, E.J., et al. (2019). Mutation of a single residue promotes gating of vertebrate and invertebrate two-pore domain potassium channels. Nat. Commun. 10.1038/s41467-019-08710-3.

Tickle, I.J., Flensburg, C., Keller, P., Paciorek, W., Sharff, A., Vonrhein, C., and Bricogne, G. (2018). STARANISO. Cambridge, United Kingdom Glob. Phasing Ltd.

Vierra, N.C., Dadi, P.K., Jeong, I., Dickerson, M., Powell, D.R., and Jacobson, D.A. (2015). Type 2 diabetes-associated K+ channel TALK-1 modulates β-cell electrical excitability, second-phase insulin secretion, and glucose homeostasis. Diabetes 10.2337/db15-0280.

Xian Tao Li, Dyachenko, V., Zuzarte, M., Putzke, C., Preisig-Müller, R., Isenberg, G., and Daut, J. (2006). The stretch-activated potassium channel TREK-1 in rat cardiac ventricular muscle. Cardiovasc. Res. 10.1016/j.cardiores.2005.08.018.

Figure 1 – Distinct basal and mechanically-activated open states in TRAAK_WT.
(A) Macroscopic currents recorded across an inside-out patch pulled from a TRAAK-expressing cell in response to a voltage step protocol (V_{hold} = 0, V_{test} = -80 to 80, ΔV = 10 mV, 20 mV intervals displayed) with a pressure step applied at each voltage (purple bar). (B) Current-voltage relationship from (A). (C) Open probability calculated from single channel records (0.04 ± 0.02, 0.47 ± 0.07, and 0.96 ± 0.01 for low P_O, mid P_O, and high P_O, respectively (mean ± sem, n = 3, 4, and 4 patches)). All single channel data in the paper were recorded at V_{hold} = 0 mV in a ten-fold gradient of [K⁺] and are presented in physiological convention. (D-F) 1 s portion from representative low P_O, mid P_O, and high P_O recordings, respectively. (G-I) All event (left) and open-only or square root all event (right) current histograms from representative low P_O, mid P_O, and high P_O recordings, respectively. (J) Unitary currents of O1 and O2 states (0.81 ± 0.03 pA and 1.80 ± 0.16 pA, 0.99 ± 0.03 pA and 2.14 ± 0.09 pA, and 1.07 ± 0.07 pA and 2.41 ± 0.13 pA for low P_O, mid P_O, and high P_O recordings, respectively (mean ± sem, n= 3, 4, and 4 patches)). (K) Open dwell time of O1 and O2 states (0.80 ± 0.15 ms, 0.91 ± 0.05 ms and 3.52 ± 0.51 ms, and 9.60 ± 1.46 ms for low P_O, mid P_O, and high P_O recordings, respectively (mean ± sem, n= 3, 4, and 4 patches)). (L) Closed dwell times of C1 and C2 states (28.70 ± 13.50 and 0.77 ± 0.03 ms, 3.59 ± 0.67 and 0.58 ± 0.05 ms, and 0.35 ± 0.01 ms for low P_O, mid P_O, and high P_O recordings, respectively (mean ± sem, n=3, 4, and 4 patches, n.d. not determined, n.s. not significant, ***p = 0.0006, ****p < 0.0001 (one-way ANOVA with Tukey correction)). (M) Unitary current-open dwell time relationships for low P_O, mid P_O, and high P_O open events and an overlay of low P_O and high P_O relationships at expanded time scale. Bubble size is proportional to percentage of open events. (N) Four state TRAAK gating model.

Figure 2 – The FHEIG mutation TRAAK_{A198E} promotes a long duration, high conductance TM4-up open state.
(A) Crystal structure of TRAAK_{A198E}. Side view from the membrane plane with one protomer gray and the second protomer colored according to the key below. A198E is shown as a yellow sphere and K+ ions are colored purple. (B) Macroscopic currents from a TRAAK_{A198E}-containing patch in response to a voltage step protocol (V_{hold} = 0, V_{test} = -80 to 80, ΔV = 10 mV, 20 mV intervals displayed) with a pressure step applied at each voltage (dark blue bar). (C) Current-voltage relationship from (B). (D,E) Open probability calculated from all TRAAK_{A198E} and TRAAK_{A198E} with pressure (+P) records (left, P_O = 0.90 ± 0.04 and P_O = 0.97 ± 0.003 (mean ± sem, n = 4 patches) and 1 s portion from representative recordings. (F,G) All event (left) and square root all event (right) current histograms from representative recordings. (H) Unitary currents of TRAAK_{WT} HP_O, TRAAK_{A198E}, and TRAAK_{A198E}+P O1 and O2 states (1.07 ± 0.07 and 2.41 ± 0.13 pA, 1.44 ± 0.14 and 2.57 ± 0.10 pA, and 1.35 ± 0.06 and 2.72 ± 0.09 pA, respectively). (I) Open dwell times of TRAAK_{WT} HPO, TRAAK_{A198E}, and TRAAK_{A198E}+P O2 states (9.60 ± 1.46 ms, 4.63 ±1.03 ms, 12.46 ±1.02 ms, respectively). (J) Closed dwell times of TRAAK_{WT} HPO, TRAAK_{A198E}, and TRAAK_{A198E}+P C2 (0.35 ± 0.01 ms, 0.42 ± 0.02 ms, and 0.39 ± 0.01 ms, respectively). For H-J, data are mean ± sem, n = 4 patches; n.d., not determined; n.s., not significant (one-way Anova with Dunnett correction). (K) View of the membrane-facing lateral opening in a TRAAK_{WT} TM4-down structure (PDB 4WFF). A cavity-bound lipid acyl chain blocks conduction. (L) TRAAK_{A198E} in the same view as (K). A TM4-up conformation seals the membrane opening. (M) Ions in a TRAAK_{A198E}-Tl+ structure. Anomalous density (grey) around Tl+ ions (green) displayed at 2.5 σ (extracellular / selectivity filter / cavity ions). (N) Overlay of the TM2-TM3-TM4 interaction from TRAAK_{A198E} and TRAAK_{WT} structures. A198E sterically promotes a TM4-up open state.

Figure 3 – The FHEIG mutation TRAAK$_{A270P}$ promotes a long duration, high conductance TM4-up open state.
(A) Crystal structure of TRAAK$_{A270P}$. Side view from the membrane plane with one protomer gray and the second protomer colored according to the key below. A270P is shown as a yellow sphere and K$^+$ ions are colored purple. (B) Macroscopic currents from a TRAAK$_{A270P}$-containing patch in response to a voltage step protocol (V$_{hold}$ = 0, V$_{test}$ = -80 to 80, ΔV = 10 mV, 20 mV intervals displayed) with a pressure step applied at each voltage (dark green bar). (C) Current-voltage relationship from (B). (D,E) Open probability calculated from all TRAAK$_{A270P}$ and TRAAK$_{A270P}$ with pressure (+P) records (left, P$_O$ = 0.94 ± 0.01 and P$_O$ = 0.96 ± 0.01 (mean ± sem, n = 6 and 5 patches) and 1 s portion from representative recordings. (F,G) All event (left) and square root all event (right) current histograms from representative recordings. (H) Unitary currents of TRAAK$_{WT}$ HP$_O$, TRAAK$_{A270P}$, and TRAAK$_{A270P}$+P O1 and O2 states (1.07 ± 0.07 and 2.41 ± 0.13 pA, 1.34 ± 0.15 and 2.65 ± 0.16 pA, and 1.21 ± 0.09 and 2.57 ± 0.12 pA, respectively). (I) Open dwell times of TRAAK$_{WT}$ HP$_O$, TRAAK$_{A270P}$, and TRAAK$_{A270P}$+P O2 states (9.60 ± 1.46 ms, 8.56 ± 1.69 ms, and 14.47 ± 3.30 ms, respectively). (J) Closed dwell times of TRAAK$_{WT}$ HP$_O$, TRAAK$_{A270P}$, and TRAAK$_{A270P}$+P C2 (0.35 ± 0.01 ms, 0.43 ± 0.01 ms, and 0.39 ± 0.01 ms, respectively). For H-J, data are mean ± sem, n = 4, 6, and 5 patches; n.d., not determined; n.s., not significant (one-way Anova with Dunnett correction). (K) View of the membrane-facing cytoplasmic half of TM4 in TRAAK$_{A270P}$. A TM4-up conformation seals the membrane opening. (L) Ions in the TRAAK$_{A270P}$-K$^+$ structure. Polder omit F$_o$-F$_c$ density (grey) around K$^+$ ions (purple) displayed at 5 and 5.5 σ for extracellular and selectivity filter and cavity ions, respectively. (M) Overlay of TM4 from TRAAK$_{A270P}$ and TRAAK$_{WT}$ structures. A270P kinks TM4 to promote a TM4-up open state.

Figure 4 – The gain-of-function mutation TRAAK_G158D promotes a short duration, low conductance TM4-down open state.
(A) Crystal structure of TRAAK_G158D. Side view from the membrane plane with one protomer gray and the second protomer colored according to the key below. G158D is shown as a yellow sphere and K^+ ions are colored purple. (B) Macroscopic currents from a TRAAK_G158D-containing patch in response to a voltage step protocol (V_{hold} = 0, V_{test} = -80 to 80, ΔV = 10 mV, 20 mV intervals displayed) with a pressure step applied at each voltage (dark orange bar). (C) Current-voltage relationship from (B). (D) Maximum fold-activation by pressure of macroscopic TRAAK_WT, TRAAK_G158D, TRAAK_A198E, and TRAAK_A270P currents with expanded scale in inset (59.33 ± 10.73, 1.75 ± 0.11, 1.11 ± 0.03, and 1.10 ± 0.03, respectively, mean ± sem, n = 3, 4, 3, and 3). (E,F) Open probability calculated from all TRAAK_G158D and TRAAK_G158D with pressure (+P) records (left, P_O = 0.71 ± 0.04 and P_O = 0.89 ± 0.02 (mean ± sem, n=3 patches) and 1s portion from representative recordings. (G,H) All event (left) and square root all event (right) current histograms from representative recordings. (I) Unitary currents of TRAAK_WT LP_O, MP_O, HP_O, TRAAK_G158D, and TRAAK_G158D+P O1 and O2 states (0.81 ± 0.03 pA and 1.80 ± 0.16 pA, 0.99 ± 0.03 pA and 2.14 ± 0.09 pA, 1.07 ± 0.07 pA and 2.41 ± 0.13 pA, 0.99 ± 0.01, and 1.06 ± 0.02 and 2.02 ± 0.08 pA, respectively). (J) Open dwell times of TRAAK_WT LP_O, MP_O, HP_O, TRAAK_G158D, and TRAAK_G158D+P O1 and O2 states (0.80 ± 0.15 ms, 0.91 ± 0.05 ms and 3.52 ± 0.51 ms, 9.60 ± 1.46 ms, 0.88 ± 0.07 and 3.58 ± 0.46 ms, and 6.63 ± 0.83 ms, respectively). (K) Closed dwell times of TRAAK_WT LP_O, MP_O, HP_O, TRAAK_G158D, and TRAAK_G158D+P C1 and C2 states (28.70 ± 13.50 and 0.77 ± 0.03 ms, 3.59 ± 0.67 and 0.58 ± 0.05 ms, 0.35 ± 0.01 ms, 0.42 ± 0.01 ms, and 0.76 ± 0.03 ms, respectively). For I-K, data are mean ± sem, n = 3, 4, 4, 3 and 3 patches; n.d., not determined; n.s., not significant, ****p < 0.0001 (one-way Anova with Tukey correction). (L) Unitary current-open dwell time relationships for TRAAK_G158D, and TRAAK_G158D+P open events and an overlay at expanded time scale. Bubble size is proportional to percentage of open events. (M) View of the membrane-facing cytoplasmic half of TM4 in TRAAK_G158D. (N) Ions in the TRAAK_G158D-K^+ structure. Polder omit F_o-F_c density (grey) around K^+ ions (purple) displayed at 6.5 and 6 σ for selectivity filter and extracellular and cavity ions, respectively. (O) The conduction path in TRAAK_G158D colored by hydrophobicity. G158D increases cavity electronegativity to promote a TM4-down open state.

Figure 5 – An integrated model for TRAAK gating with distinct basal and mechanically-gated open states.
Known structures are mapped to a linear-four state model for TRAAK gating. Basal activity corresponds to a TM4-down, low conductance, short duration open state O1. Mechanically-gated activity corresponds to a TM4-up, high conductance, long duration open state O2. Long duration closures correspond to a TM4-down lipid-blocked state C1. The unknown structure of the short duration closed state C2 is drawn without ions and the position of TM4 is indicated with dashed lines.

Figure S1 - Unitary current histograms (continued on next page)

Open event histograms with individual Gaussian fits (dashed lines) and summed fits (solid lines) for each record analyzed. (A) TRAAK$_{WT}$ low P$_O$, (B) TRAAK$_{WT}$ mid P$_O$, (C) TRAAK$_{WT}$ high P$_O$, (D) TRAAK$_{G158D}$, (E) TRAAK$_{G158D}$ with pressure, (F) TRAAK$_{A198E}$, (G) TRAAK$_{A198E}$ with pressure, (H) TRAAK$_{A270P}$, and (I) TRAAK$_{A270P}$ with pressure.

A

B

C

D

E

F

G

H

I

Figure S1 - Unitary current histograms (continued from previous page)
Open event histograms with individual Gaussian fits (dashed lines) and summed fits (solid lines) for each record analyzed. (A) TRAAK$_{WT}$ low P$_O$, (B) TRAAK$_{WT}$ mid P$_O$, (C) TRAAK$_{WT}$ high P$_O$, (D) TRAAK$_{G158D}$, (E) TRAAK$_{G158D}$ with pressure, (F) TRAAK$_{A198E}$, (G) TRAAK$_{A198E}$ with pressure, (H) TRAAK$_{A270P}$, and (I) TRAAK$_{A270P}$ with pressure.

Figure S2 - Open event dwell time histograms

Open event histograms with individual Gaussian fits (dashed lines) and summed fits (solid lines) for each record analyzed. (A) TRAAK$_{WT}$ low P$_O$, (B) TRAAK$_{WT}$ mid P$_O$, (C) TRAAK$_{WT}$ high P$_O$, (D) TRAAK$_{G158D}$, (E) TRAAK$_{G158D}$ with pressure, (F) TRAAK$_{A198E}$, (G) TRAAK$_{A198E}$ with pressure, (H) TRAAK$_{A270P}$, and (I) TRAAK$_{A270P}$ with pressure.

Figure S3 - Closed event dwell time histograms

Closed event histograms with individual Gaussian fits (dashed lines) and summed fits (solid lines) for each record analyzed. (A) $TRAAK_{WT}$ low P_O, (B) $TRAAK_{WT}$ mid P_O, (C) $TRAAK_{WT}$ high P_O, (D) $TRAAK_{G158D}$, (E) $TRAAK_{G158D}$ with pressure, (F) $TRAAK_{A198E}$, (G) $TRAAK_{A198E}$ with pressure, (H) $TRAAK_{A270P}$, and (I) $TRAAK_{A270P}$ with pressure.

Figure S4 - Purification and crystallization of TRAAK_WT, TRAAK_A198E, TRAAK_A270P, and TRAAK_G158D
(A) Chromatograms from purifications of TRAAK constructs over a Superdex 200 column. (B) Chromatograms purifications of 13E9 Fab – TRAAK construct complexes over a Superdex 200 column. (C) Coomassie-stained SDS-PAGE of fractions from TRAAK_G158D-13E9 Fab complex purification from (B). (D-G) Representative crystals grown of TRAAK_A198E, TRAAK_A270P, and TRAAK_G158D.

Figure S5 - Crystal packing and contacts between TRAAK and monoclonal Fab 13E9
(A) Crystal lattice and (B) zoomed in view of contacts with TRAAK$_{G158D}$ colored orange and Fabs blue and red.

Figure S6 - Electron density maps of TRAAK point mutations

(A) Ions in the TRAAK$_{A198E}$-K$^+$ structure. Polder omit F$_o$-F$_c$ density (grey) around K$^+$ ions (purple) displayed at 5 and 5.5 σ for selectivity filter and extracellular and cavity ions, respectively. (B-G) TRAAK$_{G158D}$ (orange), TRAAK$_{A198E}$ (blue), and TRAAK$_{A270P}$ (green) models and 2F$_o$-F$_c$ electron density displayed at 1 σ. Position of mutations are indicated.

Figure S7 - Lateral openings in K2P family ion channels
(A) Overlays of TM4A and (B) TM4B from TRAAK$_{WT}$ (gray), TRAAK$_{G158D}$ (orange), TRAAK$_{A198E}$ (blue), and TRAAK$_{A270P}$ (green) structures crystallized under the same conditions. (C) Change in cross sectional area between symmetric TRAAK$_{G158D}$ TM4-up and TRAAK$_{G158D}$ TM4-down conformations. (D-L) View of the membrane-facing cytoplasmic half of TM4 from K2Ps indicated.

		Record Properties		State Properties						Transition Rates (1/s)					
		Record length (s)	Open Probability	Duration (ms)	Duration (ms)	Unitary Current (pA)	Duration (ms)	Duration (ms)	Unitary Current (pA)	C1-O1	O1-C1	O1-C2	C2-O1	C2-O2	O2-C2
				C2	O1		C2	O2							
TRAAK WT low Po	1	134	0.01	54.93	0.62	0.77	0.79	n.d.	1.63	11	995	810	294	81	477
	2	235	0.07	10.02	0.68	0.79	0.80	n.d.	1.64	43	635	402	543	67	395
	3	1627	0.04	21.14	1.11	0.86	0.71	n.d.	2.12	11	427	305	339	144	243
	total/mean	1996	0.04	28.70	0.80	0.81	0.77	n.d.	1.80	22	686	506	392	97	372
	s.e.m.		0.02	13.50	0.15	0.03	0.03	n.d.	0.16	11	166	155	77	24	69
TRAAK WT mid Po	1	13	0.41	2.97	0.87	0.91	0.48	2.95	1.90	168	226	914	344	253	282
	2	125	0.40	4.36	0.87	0.98	0.57	3.48	2.21	145	308	743	441	338	249
	3	402	0.69	2.03	0.86	1.04	0.55	2.68	2.11	283	53	348	424	825	478
	4	629	0.38	5.02	1.06	1.04	0.73	4.97	2.34	55	137	523	492	231	192
	total/mean	1170	0.47	3.59	0.91	0.99	0.58	3.52	2.14	163	181	632	425	412	300
	s.e.m.		0.07	0.67	0.05	0.03	0.05	0.51	0.09	47	55	124	31	140	62
TRAAK WT high Po	1	6	0.97	n.d.	n.d.	1.18	0.34	9.22	2.67	n.d.	n.d.	n.d.	n.d.	2681	90
	2	20	0.98	n.d.	n.d.	1.19	0.38	12.96	2.57	n.d.	n.d.	n.d.	n.d.	2329	60
	3	6	0.97	n.d.	n.d.	0.99	0.34	10.31	2.22	n.d.	n.d.	n.d.	n.d.	2512	73
	4	14	0.93	n.d.	n.d.	0.91	0.35	5.88	2.16	n.d.	n.d.	n.d.	n.d.	1703	124
	total/mean	46	0.96	n.d.	n.d.	1.07	0.35	9.59	2.40	n.d.	n.d.	n.d.	n.d.	2306	87
	s.e.m.		0.01	n.d.	n.d.	0.07	0.01	1.47	0.13	n.d.	n.d.	n.d.	n.d.	214	14
TRAAK G158D	1	90	0.75	n.d.	0.75	0.99	0.41	4.18	n.d.	420	124	991	893	1168	400
	2	278	0.74	n.d.	0.99	1.00	0.43	3.89	n.d.	288	206	609	604	1204	307
	3	95	0.64	n.d.	0.89	0.98	0.43	2.68	n.d.	323	427	471	850	1438	381
	total/mean	463	0.71	n.d.	0.88	0.99	0.42	3.58	n.d.	344	252	690	782	1270	363
	s.e.m.		0.04	n.d.	0.07	0.01	0.01	0.46	n.d.	39	90	156	90	85	28
TRAAK G158D + pressure	1	5	0.84	n.d.	n.d.	1.10	0.74	7.87	2.16	n.d.	n.d.	n.d.	n.d.	1001	97
	2	7	0.91	n.d.	n.d.	1.02	0.81	5.05	2.04	n.d.	n.d.	n.d.	n.d.	1060	105
	3	56	0.91	n.d.	n.d.	1.07	0.73	6.97	1.87	n.d.	n.d.	n.d.	n.d.	631	119
	total/mean	68	0.89	n.d.	n.d.	1.06	0.76	6.63	2.02	n.d.	n.d.	n.d.	n.d.	897	107
	s.e.m.		0.02	n.d.	n.d.	0.02	0.03	0.83	0.08	n.d.	n.d.	n.d.	n.d.	134	6
TRAAK A198E	1	82	0.95	n.d.	n.d.	1.68	0.39	6.47	2.85	n.d.	n.d.	n.d.	n.d.	2693	108
	2	451	0.92	n.d.	n.d.	1.66	0.43	4.44	2.43	n.d.	n.d.	n.d.	n.d.	1782	162
	3	91	0.79	n.d.	n.d.	1.28	0.48	1.81	2.57	n.d.	n.d.	n.d.	n.d.	1476	356
	4	119	0.95	n.d.	n.d.	1.13	0.40	5.78	2.43	n.d.	n.d.	n.d.	n.d.	2006	112
	total/mean	743	0.90	n.d.	n.d.	1.44	0.42	4.63	2.57	n.d.	n.d.	n.d.	n.d.	1989	185
	s.e.m.		0.04	n.d.	n.d.	0.14	0.02	1.03	0.10	n.d.	n.d.	n.d.	n.d.	259	58
TRAAK A198E + pressure	1	51	0.97	n.d.	n.d.	1.50	0.38	9.79	2.93	n.d.	n.d.	n.d.	n.d.	2295	76
	2	45	0.97	n.d.	n.d.	1.21	0.43	12.01	2.52	n.d.	n.d.	n.d.	n.d.	1762	62
	3	4	0.98	n.d.	n.d.	1.39	0.36	13.57	2.66	n.d.	n.d.	n.d.	n.d.	2427	55
	4	20	0.98	n.d.	n.d.	1.31	0.39	14.45	2.79	n.d.	n.d.	n.d.	n.d.	2289	52
	total/mean	119	0.97	n.d.	n.d.	1.35	0.39	12.46	2.72	n.d.	n.d.	n.d.	n.d.	2193	61
	s.e.m.		0.00	n.d.	n.d.	0.06	0.01	1.02	0.09	n.d.	n.d.	n.d.	n.d.	147	5
TRAAK A270P	1	22	0.95	n.d.	n.d.	1.06	0.48	9.03	2.24	n.d.	n.d.	n.d.	n.d.	1576	81
	2	23	0.95	n.d.	n.d.	0.80	0.44	6.86	2.07	n.d.	n.d.	n.d.	n.d.	1939	110
	3	87	0.98	n.d.	n.d.	1.49	0.36	15.41	2.95	n.d.	n.d.	n.d.	n.d.	1862	45
	4	28	0.94	n.d.	n.d.	1.41	0.44	6.63	2.86	n.d.	n.d.	n.d.	n.d.	1685	111
	5	150	0.95	n.d.	n.d.	1.38	0.44	10.26	3.02	n.d.	n.d.	n.d.	n.d.	1298	70
	6	240	0.90	n.d.	n.d.	1.90	0.43	3.18	2.78	n.d.	n.d.	n.d.	n.d.	1597	197
	total/mean	551	0.94	n.d.	n.d.	1.34	0.43	8.56	2.65	n.d.	n.d.	n.d.	n.d.	1660	102
	s.e.m.		0.01	n.d.	n.d.	0.15	0.02	1.69	0.16	n.d.	n.d.	n.d.	n.d.	93	22
TRAAK A270P + pressure	1	27	0.96	n.d.	n.d.	0.98	0.41	11.65	2.25	n.d.	n.d.	n.d.	n.d.	1646	62
	2	5	0.99	n.d.	n.d.	1.10	0.40	22.44	2.50	n.d.	n.d.	n.d.	n.d.	2341	34
	3	5	0.93	n.d.	n.d.	1.17	0.38	7.39	2.39	n.d.	n.d.	n.d.	n.d.	1987	96
	4	5	0.98	n.d.	n.d.	1.33	0.39	21.85	2.90	n.d.	n.d.	n.d.	n.d.	1926	33
	5	59	0.95	n.d.	n.d.	1.50	0.41	8.02	2.80	n.d.	n.d.	n.d.	n.d.	1757	84
	total/mean	101	0.96	n.d.	n.d.	1.22	0.40	14.27	2.57	n.d.	n.d.	n.d.	n.d.	1931	62
	s.e.m.		0.01	n.d.	n.d.	0.09	0.01	3.30	0.12	n.d.	n.d.	n.d.	n.d.	119	13

Table S1 – Summary of all single channel recording data

	TRAAK A198E K+ PDB xxxx		TRAAK A198E Tl+ PDB xxxx		TRAAK A270P K+ PDB xxxx		TRAAK G158D K+ PDB xxxx	
Data Collection								
Beamline	ALS 8.3.1 and APS 24-IDC		APS 24-IDC		APS 24-IDC		APS 24-IDC	
Wavelength (Å)	1.115820, 0.979100, 0.979180		0.9778		0.9791		0.9791	
Number of crystals	4		4		2		2	
Space group	$P2_1$		$P2_1$		$P2_1$		$P2_1$	
Cell dimensions								
a,b,c (Å)	80.16, 136.98, 95.74		80.23, 136.53, 95.28		80.280, 136.880, 95.580		80.144, 137.308, 96.298	
α=γ, β (°)	90, 94.472		90, 94.541		90, 94.243		90, 94.72	
Resolution (Å)	95.50-2.26 (2.50-2.26)*		47.49-2.78 (3.02-2.78)*		47.57-2.78 (3.09-2.78)*		48.03-2.97 (3.23-2.97)*	
Unique reflections	95738		51728		51681		42845	
Wilson B-factor	55.04		67.69		60.08		91.95	
	Anisotropic cutoff	Spherical cutoff	Anisotropic cutoff	Spherical cutoff	Anisotropic cutoff	Spherical cutoff	Anisotropic cutoff	Spherical cutoff
R_{merge}	0.246 (2.827)	0.3263 (12.75)	0.307 (3.288)	0.4317 (10.23)	0.277 (2.167)	0.3906 (8.413)	0.168 (2.023)	0.2037 (5.504)
R_{meas}	0.251 (2.884)	0.3325 (12.99)	0.318 (3.412)	0.4397 (10.41)	0.289 (2.254)	0.4065 (8.744)	0.176 (2.109)	0.2132 (5.765)
R_{pim}	0.049 (0.567)	0.06387 (2.46)	0.085 (0.908)	0.08328 (1.946)	0.079 (0.617)	0.1116 (2.374)	0.052 (0.597)	0.06254 (1.703)
$CC_{1/2}$	0.971 (0.578)	0.999 (0.134)	0.995 (0.581)	0.999 (0.176)	0.985 (0.554)	0.992 (0.131)	0.999 (.567)	0.999 (0.174)
I / σI	10.2 (1.5)	7.43 (0.20)	9.8 (1.3)	7.43 (0.22)	6.9 (1.3)	4.69 (0.18)	8.5 (1.2)	7.03 (0.35)
Completeness (%)	93.8 (75.4)	60.31 (4.26)	93.4 (66.5)	68.62 (7.48)	93.4 (69.0)	63.22 (4.09)	93.6 (64.2)	74.03 (8.25)
Multiplicity	27.2 (25.7)	27.5 (27.7)	27.7 (27.9)	27.8 (28.4)	13.2 (13.2)	13.2 (13.5)	11.6 (12.4)	11.5 (11.3)
Refinement								
No. of reflections used†	54920		33755		31089		20170	
R_{work} (%)	22.00		19.60		21.66		19.39	
R_{free} (%)	25.21		24.12		25.96		24.13	
No. of total atoms	10660		10651		10596		10592	
No. of protein residues	1351		1351		1351		1347	
No. of K+ / Tl+ ions	7		7		7		7	
Average B factor (Å2)	69.5		82.02		71.14		110.93	
Clashscore	6.39		9.63		8.52		7.33	
Molprobity score	1.84		1.92		2.02		2.03	
Ramachandran plot								
favored (%)	96.69		96.92		96.39		96.38	
allowed (%)	3.31		3.08		3.61		3.62	
disallowed (%)	0		0		0		0	
R.m.s. deviations								
Bond lengths (Å)	0.007		0.011		0.0087		0.0088	
Bond angles (°)	1.4061		1.6072		1.5105		1.558	

*Values in parentheses are for the highest resolution shell.

5% of reflections were used to calculate Rfree

Table S2 – X-ray crystallographic data collection and model refinement statistics