Heterogeneous mutation rates and spectra within and between species of yeast and their hybrids

Anna Fijarczyk[1,2,3,4,5*], Mathieu Hénault[1,2,3,4,5], Souhir Marsit[1,2,3,4,5,6], Guillaume Charron[1,2,3,4,5,7], Christian R. Landry[1,2,3,4,5*]

[1]Département de Biologie, Université Laval, Québec, G1V 0A6 Québec, Canada; [2]Institut de Biologie Intégrative et des Systèmes (IBIS), Université Laval, Québec, G1V 0A6 Québec, Canada; [3]Département de Biochimie, Microbiologie et Bioinformatique, Université Laval, Québec, G1V 0A6 Québec, Canada; [4]PROTEO, Le réseau québécois de recherche sur la fonction, la structure et l'ingénierie des protéines, Université Laval, Québec, G1V 0A6 Québec, Canada; [5]Centre de Recherche en Données Massives (CRDM), Université Laval, Québec, G1V 0A6 Québec, Canada; [6]Current address: Département de Biologie Chimie et Géographie, Université du Québec à Rimouski, Rimouski, G5L 3A1, Québec, Canada; [7]Current address: Laurentian Forestry Center, Québec, G1V 4C7 Québec, Canada

*Authors for correspondence:
anna.fijarczyk.1@ulaval.ca and christian.landry@bio.ulaval.ca

Abstract

Mutation rates and spectra vary between species and among populations. Hybridization can contribute to this variation, but its role remains poorly understood. Estimating mutation rates requires controlled conditions where the effect of natural selection can be minimized. One way to achieve this is through mutation accumulation experiments coupled with genome sequencing. Here we investigate 400 mutation accumulation lines initiated from 11 genotypes spanning intra-lineage, inter-lineage and interspecific crosses of the yeasts *Saccharomyces paradoxus* and *S. cerevisiae* and propagated for 770 generations. We find significant differences in mutation rates and spectra among crosses, which are not related to the level of divergence of parental strains but are genotype specific. We find that departures from neutrality, differences in growth rate, ploidy and loss of heterozygosity only play a minor role as a source of variation and conclude that unique combinations of parental genotypes drive distinct rates and spectra in some crosses.

Introduction

Mutations generate genome variation, which in turn fuels evolution. They can be a source of adaptations (Venkataram et al. 2016), though more often they contribute to increased genetic load and play an important role in disease development (Deciphering Developmental Disorders Study 2017), in particular the evolution of cancer (ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium 2020). Mutations result from DNA replication errors and DNA damage, from the activity of transposable elements and other physical alterations of DNA molecules. The balance between mechanisms generating and diminishing mutations results in largely conserved mutation rates within large taxonomic units (Drake 1991; Lynch 2007), but to what extent rates of mutations and different mutation types (mutation spectra) differ among populations and between closely related species is only beginning to be understood.

Whole genome population data can provide estimates of mutation rates; however, whether novel mutations become observable variants in natural populations depends on environmental variation, selection, recombination, and demographic history. Largely unbiased mutation rates can be measured in microorganisms using fluctuation assays (Luria and Delbrück 1943) or mutation accumulation (MA) experiments followed by phenotypic assays using reporters or genome sequencing. In MA experiments, the strains are first propagated through sequential

bottlenecks for many generations and then sequenced to detect mutations. Both approaches have been used in different contexts with model organisms. In yeast, mutation rates and spectra have been shown to differ within and between species (Nguyen et al. 2020; Jiang et al. 2021), between environmental factors such as mild (Liu and Zhang 2019) and severe stressors (Shor et al. 2013) and vary depending on ploidy (Sharp et al. 2018) or sequence context (Ma et al. 2012). A wide array of genes, notably involved in DNA repair, have been shown to influence the mutation landscape (Demogines et al. 2008; Lang et al. 2013; Serero et al. 2014; Stirling et al. 2014; Gou et al. 2019; Loeillet et al. 2020).

Hybridization could play a role in shaping mutation rates as it could compound the effects of different mutator alleles (Demogines et al. 2008), introduce mutations through mutagenic crossing-over in high diversity regions (Yang et al. 2015) or disrupt mechanisms repressing transposable elements (Petrov et al. 1995; Serrato-Capuchina and Matute 2018). Alternatively, parental effects can dominate in shaping mutation rates in hybrids (Bashir et al. 2014). MA experiments by Tattini et al (Tattini et al. 2019) showed that an interspecific hybrid between one *S. cerevisiae* and *S. paradoxus* strain has mutation rates similar to *S. cerevisiae* diploid strain. However, to better understand the role of hybridization and genotype effects on mutation rate, different genetic backgrounds need to be considered.

Here we analyzed whole genome sequences from our previous MA experiments (Charron et al. 2019; Hénault et al. 2020; Marsit et al. 2021) to estimate and compare mutation rates and spectra in nine intraspecific crosses of *Saccharomyces paradoxus* and two interspecific crosses between *S. paradoxus* and *S. cerevisiae* yeast. We find that parental sequence divergence does not explain differences in mutation rate, and there is a minor role of variation in growth rate and ploidy in influencing mutation rate. Our results suggest that combinations of individual genotypes determine mutation rates and spectra in analyzed crosses.

Results and Discussion

Mutation rate and spectrum

We analyzed 400 lines from 11 crosses, including 5 intra-lineage crosses of *S. paradoxus SpC* (CC) and *SpB* (BB), 4 inter-lineage crosses between *S. paradoxus SpB* and *SpC* (BC) and

between *SpB* and *SpA* (BA) and 2 interspecific crosses between *S. paradoxus SpB* and *S. cerevisiae* (BSc) (fig. 1A,B, supplementary table S1). After ~770 generations of mutation accumulation (MA), lines accumulated a total of 1040 single and multiple nucleotide substitutions (supplementary table S2). Mean per-cross mutation rates range from 7.88×10^{-11} in the inter-lineage cross BA2 (4% parental divergence) to 2.64×10^{-10} in the intra-lineage cross BB2 (0.3% parental divergence) and they are generally consistent with mutation rates found previously in *S. cerevisiae* and *S. paradoxus* (fig. 1C, supplementary tables S2-3). Mutation rates vary significantly among crosses (supplementary table S4, negative binomial regression, cross factor at *P*-value = 1.11×10^{-6}). BC crosses (2% parental divergence) have mutation rates close to CC crosses (Mann Whitney U test, *P*-value = 0.334) but lower than BB crosses (Mann Whitney U test, 1.04×10^{-6}), suggesting that inter-lineage hybrids do not have an increased or decreased mutation rate compared to intra-lineage crosses.

There are small but insignificant differences (chi-square contingency test, *P*-value > 0.07 before BH correction) in frequency spectra of 6 mutation types (fig. 1D, after excluding 2 lines, see Methods). Mutations of type C>T (G>A) are most frequent in all crosses except BB2, which also has the lowest transition to transversion (Ts/Tv) ratio close to 0.5 (supplementary fig. S1). Unlike *S. cerevisiae,* most *S. paradoxus* crosses show in general a high frequency of A>G (T>C) mutations, which results in a smaller fraction of G/C positions being hit by a mutation (fig. 1E). Similar to *S. cerevisiae* we observe a bias of G/C to T/A mutations in hybrid crosses (fig. 1E). Again, we found no differences between intra- and inter-lineage crosses, and BB2 stands out in terms of both mutation rate and spectrum.

We investigated several biological factors which could explain differences in mutation rate and spectra among crosses and lines including differences in their growth rate, deviation from neutral evolution, whole genome or chromosome ploidy changes, and the presence of loss of heterozygosity (LOH) events.

Fig. 1. Mutation rates differ among crosses. A. Schematic phylogenetic relationships among the lineages. **B.** Crosses analyzed. Numbers within cells indicate two *S. paradoxus SpB* and two *S. paradoxus SpC* strains used in multiple crosses (details in supplementary table S1). Percentages under the crosses indicate % of sequence divergence between parental genomes. Parental strains, and crosses at T_{ini} and T_{end} were used for whole genome sequencing and growth rate was measured at three timepoints (T_{ini}, T_{mid}, T_{end}). **C.** Black points show mean mutation rates per haploid position per generation with standard errors estimated over 770 generations of MA experiment. Remaining points indicate rates with generation time adjusted for growth rates including all analyzed lines (grey) and lines with no missing replicates (white). Number of lines is depicted in parentheses. **D.** Frequencies of 6 nucleotide changes including their complementary changes. Scer MA stands for mutation spectrum of *S. cerevisiae* from the mutation accumulation experiment from Zhu et al. 2014. The three Spar columns correspond to a population dataset of 3 lineages of *S. paradoxus*: *SpA, SpB* and *SpC*. Scer 1011 corresponds to the population dataset from Peter et al. 2018. **E.** All crosses show higher frequency of G/C to A/T than expected from the proportion of mutated G/C positions. The dotted line represents the expected proportion of mutations G/C to A/T if mutations were random.

Growth rate

Mutation rate estimates calculated with an average expectation of generation time can be biased by variation in actual generation time. We sought to correct for this by measuring the growth of individual lines at three timepoints of the MA experiment by fitting the maximal slope of growth curves in liquid medium (fig. 2A) and using these values as a proxy of generation time. Overall, growth rates showed no significant differences between timepoints (mixed linear model, *P*-value = 0.63), which is consistent with reduced selection in favor of faster growth rate. There were significant differences between crosses and timepoints within individual crosses (fig. 2A, two-way ANOVA, *P*-values = 9.8×10^{-108} and 2.5×10^{-13} respectively). Neither mean growth rate (across three timepoints) nor mean growth rate by cross interaction have a significant effect on mutation rate (fig. 2B, supplementary fig. S2, supplementary table S4, negative binomial regression, *P*-values = 0.658 and 0.097 for growth and interaction respectively). At the initial timepoint of measuring growth, growth rate by cross interaction is significant, but the effects are small, and correlations are weak (supplementary fig. S2, supplementary table S4). We find that growth rates do not explain differences in mutation rates between crosses, and at most they decrease and increase slightly the mean mutation rate in BB and BSc crosses, respectively (fig. 1C).

Fig. 2. Growth rate does not explain mutation rate. A. Growth rates differ significantly between crosses. The maximum slope of individual lines at the initial, median, and final timepoints of the MA experiment are shown with sample sizes in parentheses. **B.** Mean growth rate (across three timepoints per cross) vs. mean number of mutations per cross with errors depicting standard deviation with sample sizes in parentheses.

Functional properties of *de novo* mutations

The demographic regimen of MA experiments is designed to minimize the efficiency of natural selection on spontaneous mutations, which should result in accumulated mutations that are randomly distributed in genomes. We tested this expectation by classifying the *de novo* mutations according to three functional properties expected to impact their average fitness effect: whether they occur in protein-coding sequences, and if so, whether they lead to non-synonymous substitutions or fall into essential genes. Assuming that spontaneous mutations are deleterious on average, efficient purifying selection would leave fewer mutations than expected by chance in essential genes, coding regions, and fewer non-synonymous mutations. We examined these predictions while accounting for the genomic composition and substitution spectrum of each cross (supplementary figs. S3-6; supplementary table S5). Only BA1 (4% parental divergence) exhibits a pattern consistent with purifying selection, with more mutations in non-essential genes than expected. In the BA1 lines, only three mutations were identified in essential genes. Despite all three being non-synonymous, they are not associated with growth rate decrease (supplementary fig. S7). Although growth differences may fall below our method's detection limit, this suggests that the scarcity of mutations in essential genes is not linked to increased fitness. On the other hand, four crosses exhibit opposite patterns for at least one property, with less non-coding mutations (BB1, BSc1), less mutations in non-essential genes (BB2) and more non-synonymous

mutations (BB2, BA1). Nevertheless, no pattern was consistent with increased fitness (supplementary fig. S7). Overall, these results indicate that accumulated mutations are largely consistent with neutral evolution.

Loss of heterozygosity

Crosses between closely related yeast strains are more likely to undergo homologous recombination (Tattini et al. 2019; Marsit et al. 2021), a process which can generate errors (Hicks et al. 2010; Deem et al. 2011) and potentially lead to higher mutation rates. Using data from (Marsit et al. 2021) we found no difference in the number of *de novo* mutations around the breaks of LOH segments (+/-2 kbp) compared to the rest of chromosomes carrying LOH (binomial test, 2 out of 109 mutations, expected frequency = 0.012, *P*-value = 0.395) in all crosses with *S. paradoxus SpB* parent. Chromosomes carrying LOH are not enriched for *de novo* mutations, except in BB2 (fig. 3A). The number of LOH events does not explain the number of mutations within lines (supplementary table S4, negative binomial regression, *P*-value = 0.3029) and the sum of lengths of LOH events has a positive but very small effect (supplementary table S4, negative binomial regression, *P*-value = 0.0019). In BB2, mutations on chromosomes carrying vs. devoid of LOH events do not differ in terms of Ts/Tv ratio and mutation spectrum (contingency chi square tests, P-value > 0.56). Overall, the results do not support the hypothesis that LOH is a source of mutations or small number of LOH events and uncertainty in detecting LOH breakpoints in crosses with low heterozygosity does not allow us to identify the relationship.

Fig. 3. Mutation rate is higher for some crosses with frequent LOH and lines with lower ploidy. A. Fraction of mutations on chromosomes with LOH compared to expected (diagonal), estimated as the fraction of the genome length comprising chromosomes with LOH. BB2 has significantly more mutations on LOH-carrying chromosomes than expected from the number or length of LOH-carrying chromosomes. *P*-value from binomial test after BH correction at FDR = 0.05. LOH was not calculated in CC crosses due to low genetic diversity. **B**. Differences in mutation rates between diploid and triploid MA lines. Wilcoxon rank sum test of greater mutation rate in diploids (W = 159 and W = 158 for BC1 and BC2). Significant *P*-values are shown.

Ploidy

Differences in mutation rates have been observed between haploid and diploid *S. cerevisiae* (Sharp et al. 2018). We used the BC crosses (2% parental divergence), which contain a substantial number of triploid strains from the beginning of the experiment (Charron et al. 2019), to compare mutation rates between strains that had 2 or 3 genome copies during the entire MA experiment. Mutation rates are significantly lower in BC1 triploids than in diploids, but there is no difference between the two groups in BC2 (fig. 3B). We confirmed that differences are not caused by uncalled mutations in low covered regions (supplementary fig. S8). Differences in mutation rates could be exaggerated or masked by change in fitness of some polyploid strains and consequently their higher or lower growth rate, however adjusting for growth rate does not change the results (supplementary fig. S8). Gains or losses of chromosomes during the experiment, as well as whole genome duplications do not have a measurable effect on mutation rate (supplementary fig. S9). Our results suggest that strains with higher ploidy in some crosses can experience lower mutation rates (BC1) but even if ploidy affects the mutation rate it does not influence the overall differences between crosses.

DNA repair genes

Observed differences in mutation rates and spectra are mainly driven by BB2, which has the highest mutation rate and the lowest Ts/Tv ratio and these differences are not caused by outlier lines. One possible explanation is the disruption of genes involved in DNA repair appearing just after formation of crosses. We found no common non-synonymous or loss of function variants at the beginning of MA in any gene in BB2. We also found that none of the 202 genes in BB2 involved in DNA repair lost a copy (supplementary fig. S10), and lines with lost, gained, or unbalanced number of copies of DNA repair genes do not have an increased mutation rate (supplementary fig. S11). Yeast with disrupted DNA mismatch repair processes tend to accumulate spontaneous mutations in regions with repeated sequence patterns (Lang et al. 2013). We observe that BB crosses have indeed the largest (although not significant) departure from the expected sequence

complexity around *de novo* mutations towards lower values (supplementary fig. S11). Given no obvious candidate mutator allele, increased mutation rates and biased spectra in BB2 are likely caused by unidentified interactions between parental genotypes.

In this study we demonstrate significant, though small differences in mutation rates between 11 types of hybrid crosses of yeast. Low mutation rates in *S. paradoxus SpA* reported previously (Tattini et al. 2019) are closest to BA2 (4% parental divergence), which has the lowest mutation rate among all crosses. Including genetically diverse backgrounds is therefore essential for uncovering species diversity in mutation rates. Out of several potential factors leading to such variation, differences in growth rate or changes of ploidy play a negligible role in our experiment. Estimates of mutation rates for the same cross types are largely consistent, and inter-lineage crosses (BC, 2% parental divergence) have mutation rates similar to intra-lineage crosses (BB and CC), suggesting that the level of divergence between parental genomes does not influence mutation rate. This is consistent with the results of (Hénault et al. 2020) who found no relationship between the activity of transposable elements and the level of divergence between parental genomes in the same crosses. Overall, our results suggest that lineage-specific mutation rates and genotype-specific factors contribute most to differences in mutation rates and spectra among hybrid crosses of different levels of parental divergence.

Materials & Methods

Hybrid crosses, mutation accumulation experiments and genome sequencing
The detailed methods for the generation of hybrid crosses, the mutation accumulation experiments and the whole-genome sequencing are described in (Charron et al. 2019; Hénault et al. 2020; Marsit et al. 2021). Briefly, hybrids were generated by crossing haploid strains of *S. paradoxus* or *S. cerevisiae* of genotype MATa or MATα, *ade2Δ*::hphNT1, and *hoΔ*::kanMX or *hoΔ*::natMX, and by selecting for dual resistance to geneticin and nourseothricin. From 48 to 96 lines (each derived from independent mating events) were initiated for each of a set of 11 different crosses, spanning a range of genetic divergence levels: within *S. paradoxus SpC* (CC1, CC2, CC3); within *S. paradoxus SpB* (BB1, BB2); between *S. paradoxus SpB* and *SpC* (BC1, BC2); between *S. paradoxus SpB* and *SpA* (BA1, BA2); and between *S. paradoxus SpB* and *S. cerevisiae* (BSc1, Bsc2). All the lines were evolved at room temperature on YPD solid medium (1% yeast extract, 2% tryptone, 2% dextrose, 2% agar) for 35 passages. Each passage was performed by streaking one randomly selected single colony (the closest to a predefined position

on the plate) for single colonies every three days, thus creating periodic single-cell bottlenecks in each line. Glycerol stocks of all the lines were sampled at intervals of three passages. From 23 to 40 lines were selected for whole-genome sequencing of the stocks at the initial and final timepoints. Shotgun genomic DNA libraries were prepared with the Illumina Nextera kit and a modified protocol (Baym et al. 2015). Libraries were sequenced in paired-end 150 bp mode on HiSeqX or NovaSeq 6000 systems. Short read sequence data is available from NCBI under accession PRJNA515073.

Growth rate measurements

Maximal growth rates in liquid cultures were measured to approximate the number of generations for which each MA line was evolved in the evolution experiments. For each sequenced line, the stocks from the initial, median and final timepoints were selected for growth measurement. Growth measurements were performed following a randomized design. Two technical replicates of each line at each timepoint were included. For each combination of (cross, timepoint), all lines were randomly assigned to one of seven measurement experiments. Within each experiment, lines and replicates were randomly assigned to a position in a 16 by 24 array. Glycerol stocks were thawed and spotted on YPD solid medium (1% yeast extract, 2% tryptone, 2% dextrose, 2% agar). Plates were incubated for 72 hrs at room temperature. Precultures were inoculated in 0.5 ml of YPD liquid medium and incubated at room temperature for 24 hrs. 50 μl of preculture was diluted in 1 ml of fresh YPD liquid medium. Two 80 μl aliquots of this dilution were distributed into sterile 384-well polystyrene microplates (BRANDplates pureGrade S 384, BRAND GMBH + CO KG, Wertheim, Germany) using an Eppendorf epMotion 5075 liquid handling robotic platform (Eppendorf, Hamburg, Germany). Incubation at a controlled temperature of 25°C and optical density (OD_{595}) measurements at 15 min intervals for 48 hrs were performed in a Tecan Spark microplate reader (Tecan Group Ltd, Männedorf, Switzerland) equipped with a cooling module. Replicates with no growth or abnormally high initial OD_{595} values were discarded. Values were normalized by subtracting the timewise average of empty wells and log_2 transformed. Maximum growth rate of each line at each timepoint of the experiment was inferred by fitting a Gaussian process to the curves, using the empirical variance of replicates to estimate technical error (Swain et   al.   2016).

***De novo* mutation rates**

All reads were trimmed for Illumina Nextera adapters using Trimmomatic v0.33 (Bolger et al. 2014) with options ILLUMINACLIP:nextera.fa:6:20:10 MINLEN:40. Reads from lines derived from

S. paradoxus crosses were mapped to *S. paradoxus* assembly from the *SpB* lineage (MSH-604, crosses BB1, BB2, BC1, BC2, BA1 and BA2) or *SpC* lineage (LL2011_012, 3 *SpC* x *SpC* crosses CC1, CC2, CC3). Read mapping was done with bwa mem v0.7.17 (Li 2013). To minimize the effect of divergence between the parents of *S. cerevisiae* and *S. paradoxus* on mapping accuracy, reads coming from crosses BSc1 and BSc2 were mapped to concatenated assemblies of *S. paradoxus SpB* and *S. cerevisiae* (YPS128, (Yue et al. 2017)). Picard tools v2.18 (http://broadinstitute.github.io/picard/) were used to correct ReadGroup IDs and mark duplicate reads. Nanopore assemblies of *S. paradoxus* are available from NCBI under accession PRJNA514804.

Overall sample ploidy was taken from (Marsit et al. 2021). Chromosome copy number was determined using Control-FREEC v11.5 (Boeva et al. 2012), by estimating copy number in 250 bp non-overlapping windows, and choosing one supported by most windows. Variants were called for each cross separately using Freebayes v1.3.1 (Garrison and Marth 2012) with a ploidy preset for each sample and chromosome. Options -q 20 (minimum base quality) --use-best-n-alleles 4 (evaluate only 4 best SNP alleles) --limit-coverage 20000 (maximum read depth of 20000) -F 0.02 (at least 0.02 reads supporting alternate allele) were used. Variants overlapping repeats, telomeres and centromeres were excluded. Variants were filtered based on given criteria: minimum variant quality of 30 (QUAL>30) and quality by the count of alternate alleles higher than 10 (QUAL/AO>10), each alternate allele supported by at least 3 reads on forward and 3 on reverse strand (SAF>3 & SAR>3), at least one read placed to the right and one read placed to the left of the alternate variant (RPL>1 & RPR>1) and mean mapping quality for alternate alleles vs. reference allele between 0.9 and 1.05 (0.9<MQM/MQMR<1.05). On the sample level, variant alleles had to be supported by at least 4 reads and all genotypes (including parents) had to be supported by at least 10 reads. Finally, *de novo* mutations had to be absent in both parents and absent at time T_{ini} but present in T_{end} in at most one sample from each cross.

14 samples were identified as contaminants with Principal Component Analysis and were subsequently discarded. Despite stringent variant calling, we observed reads from different crosses (parents) mapping with high quality in very low frequencies. To exclude the possibility of calling variants from these reads, we identified all parental variants separately from crosses mapped to *SpB* (BB1, BB2, BC1, BC2, BA1, BA2, BSc1 and BSc2) or *SpC* (CC1, CC2, CC3) and did not consider these positions for *de novo* variant detection.

Crosses BSc1 and BSc2 were processed with the same criteria, with an exception that *de novo* mutations on either of the genomes were considered conditional on the absence of the mutation in T_{ini} and one parental strain only. Also, small copy number variants were excluded. To check if some of the *de novo* mutations were shared between the two genomes e.g. due to loss of heterozygosity, two reference genomes were aligned with each other using minimap2 v2.17-r941 (Li 2018) and paftools.js liftover options was used to identify corresponding positions of mutations. None of them were overlapping.

De novo mutation rate per line was estimated as the number of *de novo* mutations acquired between T_{ini} and T_{end}, divided by the number of generations times the haploid genome size. The haploid genome size was summed across chromosomes where haploid chromosome length equaled length of an assembled chromosome excluding repeats, telomeres, centromeres, positions with coverage less than 10x for a line and positions with parental variants divided by the ploidy of the chromosome. Ploidy per line was estimated as an average ploidy of samples from T_{ini} and T_{end}. To adjust for the differences in growth rate in each line, generation time (770) was multiplied by the ratio: (growth rate - median growth rate)/median growth rate, where growth rate is mean growth rate estimated over 3 timepoints, and median corresponds to median growth rate over all lines from the cross.

Statistical tests were performed with R v3.6.3 or Python v3.6.8. To test different factors on mutation rates, generalized linear models were fitted to mutation counts per line. Mutation counts per 2 x 10 Mbp genome were derived from per line mutation rates. As mutation counts showed overdispersion we used negative binomial regression with glm.nb() function from MASS package in R to fit cross, growth rate, LOH count and LOH length. *P*-values for each factor or interaction were obtained with the chi-square test using anova() function in R.

Mutation spectra

Inspection of mutation spectra across all experimental lines showed that 2 outlier lines with a substantial number of mutations have highly skewed mutation spectra. CC1_48 has 14 A>T and 7 A>G nucleotide substitutions out of a total of 22 and BC2_75 has 11 A>G mutations out of a total of 17 mutations. The two samples were excluded from the mutation spectra analyses.

To obtain mutation spectra for natural populations of *S. paradoxus* we took a population dataset of biallelic single nucleotide variants from (Eberlein et al. 2019). Genome of *S. cerevisiae*

(YPS128) was aligned to determine the ancestral states. Hybrid lineages (*S. paradoxus SpCi* and *SpD*) were removed. Derived variants private to *S. paradoxus SpA, SpB* and *SpC*, present in at least 75% of strains were selected. Singletons and variants segregating at frequencies above 0.5 were removed to reduce the impact of misassignment of ancestral base. Mutation spectra were obtained from 34, 20321 and 3633 variants in *SpA, SpB* and *SpC*. A mutation spectrum for *S. cerevisiae* was obtained from a combined dataset of 1011 yeast genomes (Peter et al. 2018), with *S. paradoxus* genome (CBS432) used to assign ancestral alleles. Variants were selected with the same criteria for the combined set of all strains and the mutation spectrum was obtained from 944984 variants.

Functional analysis of *de novo* mutations

The properties of *de novo* mutations were compared to null distributions expected if mutations arose in a random manner. The properties analyzed were whether mutations arise in coding or non-coding sequences, in essential or non-essential genes, and if they were synonymous or non-synonymous substitutions. For each cross and property, the null distribution accounted for the genomic composition of the parental lineages involved and the empirical mutation spectrum. For *S. paradoxus SpA*, *SpB* and *SpC* parents, the gene predictions, and annotations of representative strains (LL2012_001, MSH-604 and LL2011_012 respectively) from (Hénault et al. 2020) were used. For *S. cerevisiae*, genome annotations of the strain YPS128 from (Yue et al. 2017) were used. Coding sequences were defined as any sequence annotated as a protein-coding gene. Essential gene annotations were taken from the Saccharomyces Genome Deletion Project at http://www-sequence.stanford.edu/group/yeast_deletion_project/Essential_ORFs.txt. Synonymous and non-synonymous substitutions were scored by translating the sequence of annotated genes.

For each property, a null distribution was derived for each cross as follows. The GC content of coding vs. non-coding regions and essential vs. non-essential genes was calculated for the representative reference genome of each subgenome of a given cross. The GC contents were used to derive the probability that a substitution occurs in a non-coding, or a non-essential region given that its initial state is an A/T or G/C base pair. Similarly, the codon frequency distribution of each reference subgenome was calculated for each cross. The per-codon count of possible synonymous and non-synonymous substitutions for each substitution type (A_C, A_G, A_T, C_A, C_G, C_T) was derived from the standard genetic code. The codon frequencies and substitution type counts per codon were used to derive the probability that a substitution is synonymous given

its substitution type. Mutations were simulated to yield the same mutation count and spectrum as determined by the empirical de novo mutations for each cross. Mutations were sampled from a multinomial distribution corresponding to the mutation spectrum, and functional properties of each mutation were sampled from binomial distributions with the previously described probabilities. The random sampling procedure was repeated for 10000 iterations to produce distributions of functional properties (proportion of non-coding, non-essential and synonymous substitutions) from which 95% confidence intervals were derived.

Analysis of DNA repair genes

In total, 202 genes present in either of the three assemblies used for mapping were selected including genes involved in DNA repair (GO:0006281), genes known to be part of the DNA mismatch repair pathway (Phillips et al. 2021) and few other genes known to affect mutation rate or spectrum in yeast (Gou et al. 2019; Loeillet et al. 2020). Variant effects were annotated with SnpEff (Cingolani et al. 2012). We searched for derived variants relative to parental genotypes, present uniquely in one cross and segregating in high frequencies. To identify loss of genes through sequence deletions, we used previously identified copy number estimates in windows to identify significant copy number losses and gains (Wilcoxon Rank Sum test P-value <0.05) overlapping genes of interest.

Data Availability

The data underlying this article are available in NCBI under project numbers: PRJNA515073 (short-sequence data) and PRJNA514804 (genome assemblies).

Acknowledgments and funding information

We thank M. Drouin, J. Hallin, D. Biot-Pelletier and Carla Bautista for their comments on the manuscript. C.R.L. holds the Canada Research Chair in Cellular Systems and Synthetic Biology. This work was supported by the Natural Sciences and Engineering Research Council of Canada (RGPIN-2020-04844 to C.R.L., Alexander Graham Bell doctoral scholarship to M.H. and G.C.), Genome Canada Grant (LSARP BIOSAFE 10106 to A.F.), Fonds de Recherche du Québec - Santé (post-doctoral scholarship to S.M.), and Fonds de recherche du Québec – Nature et technologies (doctoral scholarship to G.C.).

References

Bashir T, Sailer C, Gerber F, Loganathan N, Bhoopalan H, Eichenberger C, Grossniklaus U, Baskar R. 2014. Hybridization alters spontaneous mutation rates in a parent-of-origin-dependent fashion in Arabidopsis. *Plant Physiol.* 165:424–437.

Baym M, Kryazhimskiy S, Lieberman TD, Chung H, Desai MM, Kishony RK. 2015. Inexpensive multiplexed library preparation for megabase-sized genomes. *PLoS One* 10:1–15.

Boeva V, Popova T, Bleakley K, Chiche P, Cappo J, Schleiermacher G, Janoueix-Lerosey I, Delattre O, Barillot E. 2012. Control-FREEC: a tool for assessing copy number and allelic content using next-generation sequencing data. *Bioinformatics* 28:423–425.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120.

Charron G, Marsit S, Hénault M, Martin H, Landry CR. 2019. Spontaneous whole-genome duplication restores fertility in interspecific hybrids. *Nat. Commun.* [Internet] 10. Available from: http://dx.doi.org/10.1038/s41467-019-12041-8

Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM. 2012. A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly* 6:80–92.

Deciphering Developmental Disorders Study. 2017. Prevalence and architecture of de novo mutations in developmental disorders. *Nature* 542:433–438.

Deem A, Keszthelyi A, Blackgrove T, Vayl A, Coffey B, Mathur R, Chabes A, Malkova A. 2011. Break-induced replication is highly inaccurate. *PLoS Biol.* 9:e1000594.

Demogines A, Wong A, Aquadro C, Alani E. 2008. Incompatibilities involving yeast mismatch repair genes: a role for genetic modifiers and implications for disease penetrance and variation in genomic mutation rates. *PLoS Genet.* 4:e1000103.

Drake JW. 1991. A constant rate of spontaneous mutation in DNA-based microbes. *Proc. Natl. Acad. Sci. U. S. A.* 88:7160–7164.

Eberlein C, Hénault M, Fijarczyk A, Charron G, Bouvier M, Kohn LM, Anderson JB, Landry CR. 2019. Hybridization is a recurrent evolutionary stimulus in wild yeast speciation. *Nat. Commun.* 10:923.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing. *arXiv [q-bio.GN]* [Internet]. Available from: http://arxiv.org/abs/1207.3907

Gou L, Bloom JS, Kruglyak L. 2019. The Genetic Basis of Mutation Rate Variation in Yeast. *Genetics* 211:731–740.

Hénault M, Marsit S, Charron G, Landry CR. 2020. The effect of hybridization on transposable element accumulation in an undomesticated fungal species. *Elife* [Internet] 9. Available

from: http://dx.doi.org/10.7554/eLife.60474

Hicks WM, Kim M, Haber JE. 2010. Increased mutagenesis and unique mutation signature associated with mitotic gene conversion. *Science* 329:82–85.

ICGC/TCGA Pan-Cancer Analysis of Whole Genomes Consortium. 2020. Pan-cancer analysis of whole genomes. *Nature* 578:82–93.

Jiang P, Ollodart AR, Sudhesh V, Herr AJ, Dunham MJ, Harris K. 2021. A modified fluctuation assay reveals a natural mutator phenotype that drives mutation spectrum variation within Saccharomyces cerevisiae. *bioRxiv* [Internet]:2021.01.11.425955. Available from: https://www.biorxiv.org/content/10.1101/2021.01.11.425955v1

Lang GI, Parsons L, Gammie AE. 2013. Mutation rates, spectra, and genome-wide distribution of spontaneous mutations in mismatch repair deficient yeast. *G3* 3:1453–1465.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. *arXiv [q-bio.GN]* [Internet]. Available from: http://arxiv.org/abs/1303.3997

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* 34:3094–3100.

Liu H, Zhang J. 2019. Yeast Spontaneous Mutation Rate and Spectrum Vary with Environment. *Curr. Biol.* 29:1584–1591.e3.

Loeillet S, Herzog M, Puddu F, Legoix P, Baulande S, Jackson SP, Nicolas AG. 2020. Trajectory and uniqueness of mutational signatures in yeast mutators. *Proc. Natl. Acad. Sci. U. S. A.* 117:24947–24956.

Luria SE, Delbrück M. 1943. Mutations of Bacteria from Virus Sensitivity to Virus Resistance. *Genetics* 28:491–511.

Lynch M. 2007. The Origins of Genome Architecture. Sinauer Associates

Marsit S, Hénault M, Charron G, Fijarczyk A, Landry CR. 2021. The neutral rate of whole-genome duplication varies among yeast species and their hybrids. *Nat. Commun.* 12:1–11.

Ma X, Rogacheva MV, Nishant KT, Zanders S, Bustamante CD, Alani E. 2012. Mutation hot spots in yeast caused by long-range clustering of homopolymeric sequences. *Cell Rep.* 1:36–42.

Nguyen DT, Wu B, Long H, Zhang N, Patterson C, Simpson S, Morris K, Thomas WK, Lynch M, Hao W. 2020. Variable spontaneous mutation and loss of heterozygosity among heterozygous genomes in yeast. *Mol. Biol. Evol.* [Internet]. Available from: https://academic.oup.com/mbe/article-pdf/doi/10.1093/molbev/msaa150/33399075/msaa150.pdf

Peter J, De Chiara M, Friedrich A, Yue J-X, Pflieger D, Bergström A, Sigwalt A, Barre B, Freel K, Llored A, et al. 2018. Genome evolution across 1,011 Saccharomyces cerevisiae isolates. *Nature* 556:339–344.

Petrov DA, Schutzman JL, Hartl DL, Lozovskaya ER. 1995. Diverse transposable elements are mobilized in hybrid dysgenesis in Drosophila virilis. *Proc. Natl. Acad. Sci. U. S. A.* 92:8050–

8054.

Phillips MA, Steenwyk JL, Shen X-X, Rokas A. 2021. Examination of gene loss in the DNA mismatch repair pathway and its mutational consequences in a fungal phylum. *bioRxiv* [Internet]:2021.04.13.439724. Available from: https://www.biorxiv.org/content/10.1101/2021.04.13.439724v1.full

Serero A, Jubin C, Loeillet S, Legoix-Né P, Nicolas AG. 2014. Mutational landscape of yeast mutator strains. *Proc. Natl. Acad. Sci. U. S. A.* 111:1897–1902.

Serrato-Capuchina A, Matute DR. 2018. The role of transposable elements in speciation. *Genes* [Internet] 9. Available from: http://dx.doi.org/10.3390/genes9050254

Sharp NP, Sandell L, James CG, Otto SP. 2018. The genome-wide rate and spectrum of spontaneous mutations differ between haploid and diploid yeast. *Proc. Natl. Acad. Sci. U. S. A.* 115:E5046–E5055.

Shor E, Fox CA, Broach JR. 2013. The yeast environmental stress response regulates mutagenesis induced by proteotoxic stress. *PLoS Genet.* 9:e1003680.

Stirling PC, Shen Y, Corbett R, Jones SJM, Hieter P. 2014. Genome destabilizing mutator alleles drive specific mutational trajectories in Saccharomyces cerevisiae. *Genetics* 196:403–412.

Swain PS, Stevenson K, Leary A, Montano-Gutierrez LF, Clark IBN, Vogel J, Pilizota T. 2016. Inferring time derivatives including cell growth rates using Gaussian processes. *Nat. Commun.* 7:13766.

Tattini L, Tellini N, Mozzachiodi S, D'Angiolo M, Loeillet S, Nicolas A, Liti G. 2019. Accurate Tracking of the Mutational Landscape of Diploid Hybrid Genomes. *Mol. Biol. Evol.* 36:2861–2877.

Venkataram S, Dunn B, Li Y, Agarwala A, Chang J, Ebel ER, Geiler-Samerotte K, Hérissant L, Blundell JR, Levy SF, et al. 2016. Development of a Comprehensive Genotype-to-Fitness Map of Adaptation-Driving Mutations in Yeast. *Cell* 166:1585–1596.e22.

Yang S, Wang L, Huang J, Zhang X, Yuan Y, Chen J-Q, Hurst LD, Tian D. 2015. Parent-progeny sequencing indicates higher mutation rates in heterozygotes. *Nature* 523:463–467.

Yue J-X, Li J, Aigrain L, Hallin J, Persson K, Oliver K, Bergström A, Coupland P, Warringer J, Lagomarsino MC, et al. 2017. Contrasting evolutionary genome dynamics between domesticated and wild yeasts. *Nat. Genet.* 49:913–924.