FINE-SCALE   ADAPTATIONS   TO   ENVIRONMENTAL   VARIATION   AND   GROWTH

STRATEGIES DRIVE PHYLLOSPHERE *METHYLOBACTERIUM* DIVERSITY.

Jean-Baptiste Leducq*[1,2,3], Émilie Seyer-Lamontagne[1], Domitille Condrain-Morel[2], Geneviève

Bourret[2], David Sneddon[3], James A. Foster[3], Christopher J. Marx[3], Jack M. Sullivan[3], B. Jesse

Shapiro*[1,4] & Steven W. Kembel*[2]

1 - Université de Montréal

2 - Université du Québec à Montréal

3 – University of Idaho

4 – McGill University

* Corresponding authors. Email: jeanbaptiste@uidaho.edu, jesse.shapiro@mcgill.ca,

kembel.steven_w@uqam.ca

Abstract

Methylobacterium is a prevalent bacterial genus of the phyllosphere. Despite its ubiquity, little is

known about the extent to which its diversity reflects neutral processes like migration and drift,

versus environmental filtering of life history strategies and adaptations. In two temperate forests,

we investigated how phylogenetic diversity within *Methylobacterium* was structured by

biogeography, seasonality, and growth strategies. Using deep, culture-independent barcoded

marker gene sequencing coupled with culture-based approaches, we uncovered a considerable

diversity of *Methylobacterium* in the phyllosphere. We cultured different subsets of

Methylobacterium lineages depending upon the temperature of isolation and growth (20 °C or 30

°C), suggesting long-term adaptation to temperature. To a lesser extent than temperature

adaptation, *Methylobacterium* diversity was also structured across large (>100km; between

forests) and small geographical scales (<1.2km within forests), among host tree species, and was

dynamic over seasons. By measuring growth of 79 isolates at different temperature treatments,

we observed contrasting growth performances, with strong lineage- and season-dependent

variations in growth strategies. Finally, we documented a progressive replacement of lineages

with a high-yield growth strategy typical of cooperative, structured communities, in favor of

those characterized by rapid growth, resulting in convergence and homogenization of community

structure at the end of the growing season. Together our results show how *Methylobacterium* is

phylogenetically structured into lineages with distinct growth strategies, which helps explain their

differential abundance across regions, host tree species, and time. This works paves the way for

further investigation of adaptive strategies and traits within a ubiquitous phyllosphere genus.

Importance

Methylobacterium is a bacterial group tied to plants. Despite its ubiquity and importance to their

hosts, little is known about the processes driving *Methylobacterium* community dynamics. By

combining traditional culture-dependent and –independent (metabarcoding) approaches, we

monitored *Methylobacterium* diversity in two temperate forests over a growing season. On the

surface of tree leaves, we discovered remarkably diverse and dynamic *Methylobacterium*

communities over short temporal (from June to October) and spatial scales (within 1.2 km).

Because we cultured different subsets of *Methylobacterium* diversity depending on the

temperature of incubation, we suspected that these dynamics partly reflected climatic adaptation.

By culturing strains in lab conditions mimicking seasonal variations, we found that diversity and

environmental variations were indeed good predictors of *Methylobacterium* growth

performances. Our findings suggest that *Methylobacterium* community dynamics at the surface of

tree leaves results from the succession of strains with contrasted growth strategies in response to

environmental variations.

Acknowledgments: FRQNT (funding), NSERC (funding), Canada Research Chairs (funding),

NSF (DEB-1831838), Geneviève Lajoie, Dominique Tardif, Ariane Lafrenière, Hélène Dion-

Phénix, Yves Terrat, Kenta Araya (help for sampling), Sylvain Dallaire (help for phenotyping

screen), Sergey Stolyar (discussions), Gault Natural Reserve (McGill University), Station

Biologique des Laurentides (UdeM), Centre d'Étude de la Forêt (CEF), QCBS and two

anonymous reviewers for their helpful suggestions.

Author contributions : J.B.L., E.S.L., D.C.M., G.B., B.J.S. and S.W.K. planned fieldwork and

the experiments. J.B.L. E.S.L., D.C.M. and G.B. performed experiments. J.B.L., E.S.L., S.W.K. ,

D.S. and J.M.S performed bioinformatic analyses. J.A.F., C.J.M. and J.M.S. provided discussion

in the early stages of this study. J.B.L. B.J.S. and S.W.K. drafted the manuscript with

contributions from C.J.M.

Data accessibility: raw reads for *16S rRNA* and *rpoB* barcoding on phyllosphere communities

(BioProject PRJNA729807; BioSamples SAMN19164946-SAMN19165146) were deposited in

NCBI under SRA Accession Numbers SRR14532212-SRR14532451. Partial nucleotide

sequences from marker genes obtained by SANGER sequencing on *Methylobacterium* isolates

(BioProject PRJNA730554 ; Biosamples SAMN19190155-SAMN19190401) were deposited in

NCBI under GenBank Accession Numbers MZ268514-MZ268593 (*16S rRNA* gene),

MZ330152-MZ330358 (*rpoB* gene) and MZ330130-MZ330151 (*sucA*). Bioprojects, Biosamples,

SRA and GenBank accession numbers are listed in **Dataset S1**. R codes and related data were

deposited on Github (https://github.com/JBLED/methylo-phyllo-diversity).

Introduction

The phyllosphere, the aerial parts of plants including leaves, is a microbial habitat estimated to be as vast as twice the surface of the earth (1). Although exposed to harsh conditions including UV radiation, temperature variation, and poor nutrient availability, the phyllosphere harbors a diverse community of microorganisms, of which bacteria are the most abundant (1). A key challenge in microbial ecology and evolution is understanding the evolutionary and ecological processes that maintain diversity in habitats such as the phyllosphere. Bacteria living in the phyllosphere carry out key functions including nitrogen fixation, growth stimulation and protection against pathogens (1–3). At broad spatial and temporal scales, bacterial diversity in the phyllosphere varies as a function of geography and host plant species, potentially due to restricted migration and local adaptation to the biotic and abiotic environment (4–6), leading to patterns of cophylogenetic evolutionary association between phyllosphere bacteria and their host plants (7). Whether those eco-evolutionary processes are important at the scale of several days to several years, as microbes and their host plants migrate and adapt to changing climates, is still an open question (8). Another challenge is to link seasonal variation with plant-associated microbial community dynamics, as shifts in microbial community composition are tighly linked with host plant carbon cycling (9) and ecosystem functions including nitrogen fixation (10). More generally, we understand very little about how the ecological strategies of phyllosphere bacteria vary among lineages and in response to variation in environmental conditions throughout the growing season (9, 11).

Phenotypic traits are often phylogenetically conserved in microbes (12), and these traits influence the assembly of ecological communities through their mediation of organismal interactions with

the abiotic and biotic environment (13). Recent work has shown that many microbial traits

exhibit phylogenetic signal, with closely related lineages possessing more similar traits, although

the phylogenetic depth at which this signal is evident differs among traits (14). Most comparative

studies of microbial trait evolution have focused on broad patterns across major phyla and classes

(14), although some studies have found evidence for complex patterns of biotic and abiotic niche

preferences evolving within genus-level phylogenies (15, 16). Furthermore, to date the majority

of studies of the diversity of plant-associated microbes have been based on the use of universal

marker genes such as the bacterial *16S rRNA* gene, providing a global picture of long-term

bacterial adaptation to different biomes and host plants at broad phylogenetic scales (17).

However, these studies lack sufficient resolution to assess the evolutionary processes at finer

spatial and temporal scales that lead to the origin of adaptations within microbial genera and

species (18, 19).

The   Rhizobiales   genus   *Methylobacterium*   (*Alphaproteobacteria*,   *Rhizobiales*,

Methylobacteriaceae) is one of the most prevalent bacterial genera of the phyllosphere, present

on nearly every plant (20–22). Characterized by pink colonies due to carotenoid production,

Methylobacterium are facultative methylotrophs, able to use one-carbon compounds, such as

methanol excreted by plants, as sole carbon sources (23, 24). Experimental studies have shown

the important roles of *Methylobacterium* in plant physiology, including growth stimulation

through hormone secretion (25–27), heavy metal sequestration (27), anti-phytopathogenic

compound secretion, and nitrogen fixation in plant nodules (28), sparking increasing interest in

the use of *Methylobacterium* in plant biotechnology applications (27, 29, 30). Although up to 64

Methylobacterium species have been described (31–39), genomic and phenotypic information

was until recently limited to a small number of model species: *M. extorquens, M. populi*, *M.*

nodulans, *M. aquaticum* and *M radiotolerans*, mostly isolated from anthropogenic environments,

and only rarely from plants (40–44). Aditionaly, *Methylobacterium* was mostly isolated assuming

its optimal growth was in the range 25-30 °C (45), an approach that could bias strain collections

toward mesophylic isolates to the exclusion of isolates from temperate forests where

temperatures typically range from 10 to 20 °C during the growing season (46). Newly available

genomic and metagenomic data now allow a better understanding of the distribution of

Methylobacterium diversity across biomes (31) and suggest that they represent a stable and

diverse fraction of the phyllosphere microbiota (22). However, we still understand relatively little

about the drivers of the evolution and adaptation of *Methylobacterium* in natural habitats.

In this study, we assessed the diversity of *Methylobacterium* in temperate forests and asked

whether *Methylobacterium* associated with tree leaves act as a single unstructured population, or

if their diversity is structured by regional factors (*e.g.* a combination of isolation by distance and

regional environmental variation) or by niche adaptation (*e.g.* host tree or temperature adaptation)

(12). First, we assessed *Methylobacterium* diversity by combining culturing and metabarcoding

approaches along with phylogenetic analysis and quantified how this diversity varied across

space, time, and environment in the phyllosphere. Second, we quantified the extent of

phylogenetic niche differentiation within the genus, with a focus on quantifying the evidence for

adaptation to local environmental variation at different spatial, temporal and phylogenetic scales.

We hypothesized that distinct phylogenetic lineages would be associated with distinct

environmental niches. Third, we quantified *Methylobacterium* growth performance under fine-

scale environmental variations, with a focus on temperature, to determine whether fine-scale

changes in diversity over space and time might result from environmental filtering of isolates

with contrasting growth strategies under local environmental conditions. We found that

Methylobacterium phyllosphere diversity consisted of deeply branching phylogenetic lineages

associated with distinct growth phenotypes, isolation temperatures, and large-scale spatial effects

(forest of origin), while finer-scale spatial effects, host tree species, and time of sampling were

more weakly and shallowly phylogenetically structured. Over the course of a year, from spring to

fall, we observed a homogenization of *Methylobacterium* community structure coinciding with

the progressive replacement of isolates with high yield strategy by isolates with rapid growth.

Together our results show that this ubiquitous phyllosphere genus is structured into lineages with

distinct growth strategies, which helps explain their differential abundance across space and time.

Methods

Phylogenetics of plant-associated Methylobacterium diversity.

We evaluated the known *Methylobacterium* diversity and its distribution across biomes, with a

special emphasis on the phyllosphere. First, we constructed a phylogeny of *Methylobacteriaceae*

from the complete nucleotide sequence of *rpoB*, a highly polymorphic housekeeping gene

commonly used to reconstruct robust phylogenies in bacteria, because unlikely to experience

horizontal gene transfer or copy number variation (47, 48). We retrieved *rpoB* sequences from

genomes publicly available in September 2020, including 153 *Methylobacteria*, 30 *Microvirga*

and 2 *Enterovirga* (**Dataset S1a**), performed alignment and inferred a consensus phylogeny with

MrBayes v. 3.2.7a ((49); **Supplementary Materials and Methods S1)**. For each

Methylobacterium reference genome, we retrieved the species name and the sampling origin,

when available. Additionally, we assigned each genome to a group (A, B, C) according to

previously proposed subdivisions (31). We subdivided group A in nine clades (A1-A9;

Supplementary Materials and Methods S1).

Study sites and sample collection

The two study forests were located at the Gault Nature Reserve (Mont Saint-Hilaire, Quebec,

Canada; 45.54 N 73.16 W), here referred as MSH, an old forest occupying Mount Saint-Hilaire,

and the *Station Biologique des Laurentides* (Saint-Hippolyte, Quebec, Canada; 45.99 N 73.99

W), here referred to as SBL, a mosaic of natural wetlands, xeric and mesic forests (**Figure 1**,

Dataset S1b). In August 2017, for the purpose of a pilot survey, we collected leaves from the

subcanopy (3-5m) of 19 trees among dominant species in MSH (*Fagus grandifolia*, *Acer*

saccharum, *Acer pensylvanicum* and *Ostrya virginiana*). In 2018, we realized a time series

survey in MSH and SBL. In each forest, we marked and collected leaf samples from the

subcanopy of 40 trees (representative of local tree species diversity) in 4-6 plots distributed along

a 1.2 km transect (**Supplementary Materials and Methods S1**). In MSH, the transect followed

an elevation and floristic gradient dominated by tree species *F. grandifolia (FAGR)*, *A.*

saccharum (ACSA), *O. virginiana (OSVI)* and *Quercus rubra (QURU)*. In SBL, the transect

followed a constant environnement dominated by *A. saccharum*, *F. grandifolia*, *A.*

pensylvanicum (ACPE), *Abies balsamea* (*ABBA*) and *Acer rubrum* (*ACRU*). For this time series,

each tree was sampled 3-4 times from June to October 2018. For each sampled plot and time

point, we also sampled a negative control consisting of empty sterile bags opened and sealed on

site. The leaf surface microbial community from each sample was collected with phosphate

buffer and split in two equal volumes for microbial community DNA extraction and

Methylobacterium isolation, respectively (**Supplementary Materials and Methods S1**).

Methylobacterium *isolation and development of a fine-scale single-copy molecular marker*
specific to Methylobacterium

For both pilot and time series surveys, we performed *Methylobacterium* isolation on MMS
synthetic solid media with 0.1% methanol supplemented with yeast extract and vitamins
(**Supplementary Materials and Methods S1**). For each leaf sample, isolation was replicated at
20 °C and 30 °C to minimize biases toward mesophylic strains. Isolates from the 2017 pilot
survey (*n*=80; **Dataset S1c**) were identified by PCR amplification and partial sequencing of the
16S rRNA ribosomal gene and assigned to *Methylobacterium* clades (**Supplementary Materials**
and Methods S1; Dataset S1d,e). As an alternative to the *16S rRNA* gene, we developed a
highly polymorphic marker targeting the *Methylobacteriaceae* family. We tested two candidate
genes, *rpoB* (47, 48, 50, 51) and *sucA* (51–53), which, contrary to *16S rRNA*, were single-copy in
Methylobacterium genomes and were polymorphic enough to distinguish among
Methylobacterium groups and clades (**Supplementary Materials and Methods S1; Figure S1**).
In 20 representative *Methylobacterium* isolates from the 2017 pilot survey (**Dataset S1c,d,e**), we
successfully amplified a *rpoB* hypervariable region (targeted by primers Met02-352-F and
Met02-1121-R), that we choose as a specific marker for *Methylobacteriaceae* (**Supplementary**
Materials and Methods S1; **Table S1**). Isolates from the 2018 timeline survey (*n*=167; **Dataset**
S1e,f,) were assigned to *Methylobacterium* clades using a consensus phylogenetic tree inferred
with MrBayes v. 3.2.7a (49) from nucleotide sequences of the *rpoB* marker obtained for these
isolates, aligned together with *rpoB* complete nucleotide sequences available from 188
Methylobacteriaceae genomes (**Dataset S1a**) and partial nucleotide sequences obtained from 20
representative isolates from the pilot survey (**Supplementary Materials and Methods S1**).

Culture-based assessment of Methylobacterium *diversity in the tree phyllosphere.*

We tested for associations between *Methylobacterium* culture-based diversity at different

phylogenetic depths, with isolate characteristics as proxy for adaptive response to environmental

variables through their evolution, using the *rpoB* phylogenetic tree built from timeline survey

isolates as a guide. We assigned *Methylobacterium* isolates according to their phylogenetic

placement. After excluding nodes supported by less than 30% of bootstraps, the tree was

converted into an ultrametric tree scaled proportionally to pairwise nucleotide similarity (*PS;*

Supplementary Materials and Methods S1). First, for each *PS* value in the tree in the range

0.926-1.000 (corresponding to *PS* range within clades), we classified isolates into discrete taxa

and performed a PERMANOVA (10,000 permutations) on *Methylobacterium* community

dissimilarity using the *Bray-Curtis* index (*BC*) based on taxa absolute abundance (Hellinger

transformation) using the R package *vegan* (54). We tested for the relative contribution of four

factors and their interactions on taxon frequency: sampling forest (*F*); temperature of isolation

(*T*); sampling time (*D*) and host tree species (*H*). Second, we asked specifically which nodes

within the tree were associated with *F* and *T*. For each node with at least 30% of support, and

each factor, we tested for the association between embedded taxa and *F* (SBL and MSH) or *T* (20

and 30 °C) by permutation of factors between embedded nodes (100,000 permutations per node;

Supplementary Materials and Methods S1).

Culture-free assessment of Methylobacterium *diversity in the tree phyllosphere (barcoding)*

We evaluated the bacterial phyllosphere diversity through barcoding and sequencing of phyllosphere samples from the 2018 timeline survey. First, we evaluated the bacterial diversity targeting the *16S rRNA* gene (55) in 46 phyllosphere samples from 13 trees from both forests sampled 3-4 times throughout the 2018 growth season. We included one negative control, and one positive control consisting of mixed DNAs of *Methylobacterium* isolates typical of the phyllosphere (METH community; **Supplementary Materials and Methods S1**). Second, we evaluated the *Methylobacteriaceae* phyllosphere diversity targeting the *rpoB* marker (see above) in 184 phyllosphere samples from 53 trees representative of diversity found in MSH (n=26) and SBL (n=27), sampled 3-4 times throughout the 2018 growth season. We included four negative controls and four positive controls (METH community). Library preparation and sequencing were performed as described in **Supplementary Materials and Methods S1.** For each phyllosphere sample and controls, we estimated bacterial diversity based on Amplicon Sequence Variants (ASVs) using package *dada2* in R (56). We assessed ASV taxonomy using SILVA v.138 database for *16S rRNA* gene (57) and a *rpoB* nucleotide sequence database available for *Bacteria* (48), curated by a ML phylogenetic tree (200 permutations**; Supplementary Materials and Methods S1**). Taxonomy for *Methylobacterium* ASVs (at the clade level) was refined using blast against NCBI databases for *16S rRNA* gene (58) and using phylogenetic placement for *rpoB* (**Supplementary Materials and Methods S1**). To validate the *rpoB* barcoding accuracy in estimating *Methylobacterium* diversity, we compared *Methylobacterium* clade relative abundances estimated from *16S rRNA* and *rpoB* barcoding in a heatmap (**Supplementary Materials and Methods S1**). We also compared *Methylobacterium* diversity estimations from *rpoB* barcoding and culture-dependant approaches by matching *rpoB* partial nucleotide sequences obtained from isolates with those obtained from ASVs (**Supplementary Materials and Methods S1**). We evaluated relative contributions of sampling forest (*F*), plot within forest (*P*), host tree

species (*H*), time of sampling (*T*) and their interactions on bacteria (*16S rRNA* barcoding) and

Methylobacteriaceae (*rpoB* barcoding) community dissimilarity among phyllosphere samples

(*BC* index, Hellinger transformation on ASV relative abundance), using PERMANOVA (10,000

permutations; **Supplementary Materials and Methods S1**) and principal component analysis

(PCA). For *rpoB* barcoding, specifically, we reported *Methylobacterium* ASV significantly

associated with the aforementioned factors (*F*, *P*, *H*, *T;* ANOVA) into the PCA (**Supplementary**

Materials and Methods S1).

Spatial and temporal dynamics of Methylobacterium *communities*

We evaluated the spatial and temporal dynamics of *Methylobacterium* communities in the

timeline survey (*rpoB* barcoding) using autocorrelation analyses. In order to remove potential

differences in community composition between forests, we analyzed samples from MSH and

SBL separately. For each pairwise comparison between two samples from the same forest, we

evaluated the effects of spatial distance (*pDist*) separating trees sampled at the same date (*spatial*

autocorrelation analyses) and time (*pTime*) separating dates at which trees were sampled

(*temporal autocorrelation analyses*) on *BC* dissimilarity among samples (see above). We

evaluated the effects of *pDist* and *pTime* on *BC* under linear models by ANOVA

(**Supplementary Materials and Methods S1**).

Ecophylogenetic structure of Methylobacterium *communities*

We quantified the ecophylogenetic structure of *Methylobacterium* communities by comparing the

phylogenetic dissimilarity of co-occurring *rpoB* ASVs with the dissimilarity expected under a

null model of stochastic community assembly from the pool of all ASVs, in order to quantify the

evidence for different community assembly processes (59) as a function of forest, host tree

species, and time of sampling. For each community of *Methylobacterium* ASVs, we calculated a

measure of phylogenetic dissimilarity among co-occurring ASVs (mean nearest taxon distance

(MNTD)) and compared observed MNTD to that expected under a null model of stochastic

community assembly from the pool of all ASVs. We calculated the standardized effect size (SES)

of MNTD (60), which expresses the difference between the observed MNTD value versus the

mean and standard deviation of MNTD values obtained across 999 random draws of ASVs from

the pool of observed ASVs across all samples while maintaining observed sample ASV richness

(61). We evaluated the effects of forest, host tree species, and time of sampling on SES(MNTD)

by ANOVA.

Monitoring of Methylobacterium *growth performance*

We evaluated the growth abilities of 79 *Methylobacterium* isolates from the timeline survey for

four temperature treatments mimicking temperature variations during the growing season. Each

treatment consisted of an initial pre-conditioning step (*P*) during which each isolate was

incubated on solid MMS media with methanol as sole carbon source for 20 days at either 20 °C

(*P20*) or 30 °C (*P30*), and a second monitoring step (*M*) during which pre-conditioned isolates

were incubated on the same media and their growth monitored for 24 days at 20 °C (*P20M20* and

P20M30) or 30 °C (*P30M20* and *P30M30*; **Figure S2***).* Treatments *P20M20* and *P30M30*

mimicked stable thermal environments, and treatments *P20M30* and *P30M20* mimicked variable

thermal environments. For each isolate and each combination of treatments (*PXXMXX*), we

realized 5 replicates, randomly spotted on 48 petri dishes according to a 6×6 grid. During the

monitoring step, we took photographs of each petri dish at days 7, 13 and 24 after inoculation

(**Figure S2**). Photos were converted to pixel intensities with ImageJ 1.52e and processed in R for

background correction, measurement of spot intensities and correction for position-dependant

competition effects (**Figure S3**; **Supplementary Materials and Methods S1).** For each isolate

and temperature treatment, logistic growth curves were inferred from bacteria spot intensity

variation observed over three time points during the monitoring step. From growth curves, we

estimated maximum growth intensity, or yield (Y) and growth rate (r) as the inverse of lag+log

time necessary to reach Y (**Figure S4** (62, 63); **Supplementary Materials and Methods S1**). We

evaluated the effects of following factors on *Methylobacterium* growth abilities (Y and r) under

different temperature treatments: isolate assignement to clades (C), forest of origin (F), host tree

species (H), time of sampling (D), temperature of isolation (T_I; at which each isolate was

isolated), temperature of incubation during pre-conditioning (T_P) and monitoring (T_M) steps, and

all possible interactions between those factors (ANOVA; **Supplementary Materials and**

Methods S1).

Results

Phylogenetics of plant-associated Methylobacterium diversity.

A phylogeny of 153 *Methylobacterium* isolates built from available genomic databases showed

that plants (65% of strains) and especially the phyllosphere compartment (41% of strains) were

the most prevalent source of *Methylobacterium* sampled to date (**Figure 2; Dataset S1a**).

Phyllosphere-associated diversity was not randomly distributed in the *Methylobacterium*

phylogenetic tree. Isolates from the phyllosphere represented the largest part of diversity within

group A (56% of isolates) but not in groups B and C (17 and 12% of isolates, respectively).

Group A was paraphyletic and most of its diversity consisted of undescribed taxa falling outside

previously well-described linages. Accordingly, we subdivided *Methylobacterium* group A into 9

monophyletic clades (A1-A9).

16S rRNA *community analyses of the tree phyllosphere.*

We focused on *Methylobacterium* phyllosphere diversity variation observable at the scale of

seasonal variation (within year 2018) on individual trees within two temperate forests of

northeastern North America (**Figure 1a,b; Dataset S1b,g**): Mont Saint Hilaire (MSH**; Figure 1c**)

and Station biologique des Laurentides (SBL**; Figure 1d**). The distribution of the phyllosphere

bacterial community assessed in 46 leal samples by bacterial *16S rRNA* gene amplicon sequence

variants (ASVs) was mostly explained by differences among forests (31.6% of variation

explained; p<0.001; PERMANOVA), host tree species (15.6% of variation; p<0.001) and time of

sampling (12.0%; p<0.05; **Table 1**). Although representing only 1.3% (0.0-3.2% per sample) of

total *16S rRNA* sequence diversity, *Methylobacterium* was present in almost all analyzed samples

(45 out of 46; **Dataset S1h**). We assigned the 15 *Methylobacterium* ASVs identified by *16S*

rRNA sequencing to clades from *Methylobacterium* group A: A9 (related to *M.*

phyllosphaerae/M. mesophilicum/M. phyllostachyos/ M. pseudosasicola/M. organophilum;

0.87% of total diversity, nine ASVs), A6 (related to *M. cerastii*, 0.29%; one ASV) and A1

(related to *M. gossipicola*; 0.13%, 3 ASVs; **Table S2**; **Dataset S1i**). With two rare ASVs

(<0.01% of relative abundance) related to *M. komagatae*, belonging to group A (31) but unrelated

to any aforementioned clade, we defined a new clade (A10). No ASVs from MSH or SBL were

assigned to group B or group C.

Culture-based assessment of Methylobacterium diversity in the tree phyllosphere.

We evaluated the culturable part of *Methylobacterium* diversity from a subsample of 36 trees (18 per forest). Using *rpoB* gene partial nucleotide sequences as a marker, we identified 167 pink isolates that we assigned to *Methylobacterium* based upon their phylogenetic placement (**Dataset S1e,f; Figure 3**). As observed for *16S rRNA* ASVs, most isolates were assigned to clades from group A typical of the phyllosphere: A9 (59.9% of isolates), A6 (24.6%), A1 (5.4%), A10 (3.6%) and A2 (related to *M. bullatum* and *M. marchantiae*; **Dataset S1d** 1.8%). Few isolates were assigned to group B (4.2% of isolates, related to *M. extorquens*) and none to group C (**Table S2**). The higher polymorphism in the *rpoB* marker revealed a considerable diversity within clades, as we identified 71 unique *rpoB* sequences, in contrast to the smaller number obtained with *16S rRNA* barcoding (15 ASVs). We determined that *Methylobacterium* diversity assessed at varying depths in the *rpoB* phylogeny was systematically explained by forest of origin (4.5±1.0% of variance explained; PERMANOVA; $p<0.001$; **Figure 3a; Dataset S1j**) and temperature of isolation (5.9±2.1% of variance explained; $p<0.001$). Temperature of isolation was the most important factor distinguishing deep phylogenetic divergences (pairwise nucleotide similarity range: 0.948-0.993), while forest of origin was slightly more important in structuring more recently diverged nodes (pairwise nucleotide similarity >0.993). Time of sampling had a slight but significant effect on diversity (2.1±0.2% of variance explained; $p<0.05$) and it was only observed for higher pairwise nucleotide similarity values (range 0.994-1.000). We did not observe any significant effects of host tree species on *Methylobacterium* isolate diversity, at any level of the phylogeny. In the phylogeny, we identified two nodes strongly associated with temperature of isolation, corresponding to clades A6 (20 °C; $p<0.001$; permutation test) and

A9+A10 (30 °C; *p*<0.001; **Figure 3b**). Other clades were evenly isolated at 20 and 30 °C and we

observed no significant association between temperature of isolation and nodes embedded within

clades. Nodes associated with forest of origin also roughly corresponded to certain major clades,

with clades A1+A2 almost exclusively sampled in MSH (*p*<0.01). Overall, clade A9 was isolated

significantly more often at SBL (*p*<0.001) but at least three of its subclades were significantly

associated with either MSH or SBL (*p*<0.05).

Comparison of Methylobacterium diversity assessed by rpoB barcoding and isolation

We performed culture-independent *rpoB* amplicon sequencing from 179 leaf samples from 53

trees in both forests, allowing a monthly monitoring for most trees (**Dataset S1b,g**). We

identified 283 *Methylobacteriaceae rpoB* ASVs in these samples (**Dataset S1k,l**), representing

24.6% of all sequences. Non-*Methylobacteriaceae* ASVs were mostly assigned to other

Rhizobiales families (850 ASVs, 70.33% of sequence abundance) and to Caulobacterales (209

ASVs, 4.42% of sequence abundance) typical of the phyllosphere (**Supplementary Materials**

and Methods S1), indicating that the *rpoB* marker can potentially be used at a broader taxonomic

scale (**Figure S5a**). Within *Methylobacteriaceae*, ASVs were mostly classified as

Methylobacterium (200 ASVs, 23.05% of sequence relative abundance), and *Enterovirga* (78

ASVs, 1.56%; **Dataset S1k**). We assigned most of *Methylobacterium* ASVs to previously

cultured clades A9 (45.2% of *Methylobacterium* sequence abundance), A6 (24.3%), A1 (6.1%)

and A10 (1.0%; **Dataset S1k; Table S2; Figure S5b**). Estimates of *Methylobacterium* diversity

based on *rpoB* sequences from culture-independent sequencing were generally concordant with

estimates based on *16S rRNA* barcoding (**Figure S5c; Table S2**) and estimates from cultured

isolates (**Figure S5d; Table S2**). The major exception was group B, representing 19.1% of

Methylobacterium sequence abundance (*rpoB* barcoding) but not detected by *16S rRNA*

barcoding, and representing 4.2% of isolates (**Table S2**). Clade A4 (related to *M. gnaphalii* and

M. brachytecii) represented 1.7% of *Methylobacterium* sequence abundance (*rpoB* barcoding) but

was not detected by *16S rRNA* barcoding, nor was it isolated. Other clades could be detected by

rpoB barcoding with low sequence abundance (<0.3%) but not by *16S rRNA* barcoding, and were

unevenly isolated (<1.8% of isolates).

Fine-scale temporal and spatial distribution of Methylobacterium diversity assessed by rpoB

barcoding

The community composition of the 200 *Methylobacterium* ASVs was mostly explained by spatial

variation at both large (distance between forests: 100 km) and local scales (distance between

plots within forest: 150-1,200 m), as well as sampling date during the growing season (1-5

months; proportion of variation explained: 32.4%, 8.0% and 4.8%, respectively; p<0.001;

PERMANOVA; **Table 1**). We observed slight but significant effects of host tree species, and of

the interaction between host tree species and plots within forests, on *Methylobacterium*

community composition (explaining 7.1% and 4.3% of variation in community composition;

p<0.001 and p<0.01, respectively; PERMANOVA; **Table 1**). A large proportion of

Methylobacterium ASVs (83 out of 200) were significantly associated with one or either forest

(ANOVA; **Figure 4a; Dataset S1m**), regardless their clade membership. The only exception was

clade A1, which was almost exclusively observed (and isolated; see **Figure 3b**) in the MSH

forest. We found 25 ASVs whose relative abundance significantly increased throughout the

growing season (ANOVA; p<0.05), mostly belonging to clades A1 (n=11). Four ASVs increased

significantly in frequency over time in both forests, and mostly belonged to group B (n=3),

(**Dataset S1m**). We found no clear association between ASV or clade with host tree species, nor

plots within forests (data not shown). *Methylobacterium* diversity was heterogeneously

distributed at local spatial scale, as we observed a significant increase of community dissimilarity

(Bray-Curtis index; *BC*) with geographical distance separating two samples within MSH (spatial

autocorrelation analysis; ANOVA; $p<0.001$) but not SBL ($p>0.05$, **Table 2, Figure 4b**). We also

observed a significant increase of community dissimilarity over time separating two sampling

dates in both forests (temporal autocorrelation analysis; $p<0.001$; **Table 2**), indicating that

community composition changed during the growing season. This effect was more marked in

MSH than in SBL (**Figure 4c**). The overall community *BC* dissimilarity consistently decreased

from June to October in both MSH (from 0.624 to 0.297) and SBL (from 0.687 to 0.522; **Table 2,**

Figure 4d), indicating that the observed change of diversity over time resulted from a progressive

homogeinization of *Methylobacterium* community between the beginning and the end of the

growing season at the scale of a forest, although without affecting its heterogeneous spatial

distributions in MSH (**Table 2, Figure 4e**). *Methylobacterium* communities were strongly

phylogenetically clustered (**Figure 4f**), with all communities containing ASVs that were much

more closely related than expected by chance (mean SES(MNTD) (± standard deviation) = -4.8 ±

0.9, all SES(MNTD) *p*-values <0.05 compared with null model of random community assembly).

While all communities were strongly phylogenetically clustered, SES(MNTD) differed among

host tree species (ANOVA; F=6.4, $p<0.001$) and forests (ANOVA; F=10.9, P<0.001), and

decreased during the growing season (ANOVA; F=95.2, $p<0.001$).

Effect of short scale temperature variation in combination with other environmental and genetic

factors on Methylobacterium growth performances

We measured growth of 79 *Methylobacterium* isolates (sampled in 2018 in both forests; MSH:

$n=32$, SBL: $n=47$) in conditions mimicking temperature variations during the growing season

(**Figures S2-S4** ; **Dataset S1n**). Clade membership explained a large part of variation in growth

rate (r) and yield ($Y;$ 7.6 and 30.6% of variation explained, respectively; ANOVA; $p<0.001$;

Figures 5a,b, Table 3; Dataset S1o). Group B isolates ($Y = 12.2 \pm 5.0$) have higher yield than

group A ($Y = 5.4 \pm 3.5$). Isolates from clades A1, A2 and B had the highest growth rate (r range:

$0.101\pm0.032 – 0.121\pm0.031$). Other clades (A6, A9 and A10) had on average slower growth (r

range: $0.082\pm0.021 – 0.088\pm0.024$). Time of sampling, host tree species and forest also explained

significant variation in growth rate (5.4%, $p<0.001$; 2.2%, $p<0.01$ and 1.5%, $p<0.05$,

respectively; ANOVA) and limited or no significant variation in yield (1.3%; $p<0.001$; 1.3%;

$p<0.01$; 0.2%; $p>0.05$, respectively; **Table 3**). Among the aforementioned factors only the

interaction between time of sampling and clade membership explained significant variation in

growth rate (2.9%; $p<0.001$), while all possible pairwise interactions between these factors

explained significant variation in yield (range 1.4 – 5.9%; $p<0.01$; **Table 3**). In both SBL and

MSH, growth rate increased consistently from June ($r = 0.075\pm0.018$ and 0.085 ± 0.033,

respectively) to September/October ($r = 0.097\pm0.031$ and 0.103 ± 0.027, respectively; **Figure 5c**).

The temperature of isolation (at which each isolate was originally isolated) had very limited

effect on growth rate (1.0%; $p<0.01$) and yield (0.6%; $p<0.05$). These effects were independent

of temperatures during pre-conditioning and monitoring steps (no significant interaction in the

ANOVA). Temperature of incubation had significant effects on growth performance.

Temperature during the monitoring step explained respectively 2.0% and 15.8% of variation in

yield and growth rate ($p<0.01$ and $p<0.001$, respectively; ANOVA; **Table 3**), regardless of clade

membership, time of sampling, and other environmental factors (no significant interaction in the

ANOVA). Isolates incubated at 20 °C had on average higher yield ($Y=6.9\pm5.4$) but slower growth

($r=0.077\pm0.022$) than isolates incubated at 30 °C ($Y=4.9\pm3.6$; $r=0.100\pm0.030$; **Figure 5d**).

Temperature during the pre-conditioning step had no effect on growth rate ($p>0.05$; ANOVA),

and limited effect on yield (1.4%; $p<0.05$; ANOVA; **Table 3**).

Discussion

Methylobacterium is ubiquitous on leaves in the temperate forests of Québec and its diversity in

this habitat is quite similar to what has been described in the phyllosphere throughout the world,

with three main clades A9 (*M. brachiatum*, *M. pseudosasicola*), A6 (related to *M. cerastii*) and

A1 (related to *M. gossipicola*) dominating diversity. Our barcoding approach based on a clade-

specific *rpoB* marker revealed previously undocumented diversity within these clades, as well as

within several other clades that were not detected by a classical *16S rRNA* marker: B (related to

M. extorquens), A2 (related to *M. bullatum* and *M. marchantiae*), A4 (related to *M. gnaphalii* and

M. brachytecii) and A10 (related to *M. komagatae*). This diversity, like that of the overall

phyllosphere community, was mostly determined by differences between forests, with barcoding

approaches suggesting combined effects of restricted migration, local adaptation to host tree

species, and climatic conditions at large geographical scales (>100km). With higher molecular

resolution, we observed that *Methylobacterium* diversity was spatially structured even at the scale

of a forest (within 1.2 km), and also showed a clear pattern of temporal dynamics and succession

over the course of a growing season. This result indicates that, although representing a stable

proportion of the plant leaf microbiota between years (22) *Methylobacterium* diversity is highly

dynamic within the course of a season. A finer analysis of *Methylobacterium* diversity suggested

that clade identity partly explained *Methylobacterium* geographical distribution at large scales

(between forests) but not at finer scales (plots), nor was it an indicator of adaptation to a

particular host tree species, nor a determinant of temporal dynamics. These results are consistent

with previous observations that geographic origin is a stronger driver of phyllosphere

Methylobacterium diversity than host identity (22). The distribution of *Methylobacterium*

diversity at small temporal and geographical scales likely resulted from more contemporaneous

community assembly events selecting for phenotypic traits that evolved among deeply diverging

lineages of *Methylobacterium*, as has been observed in other bacterial (16) and plant clades (64).

We found further evidence for deterministic community assembly as *Methylobacterium*

communities were strongly phylogenetically clustered compared to the expectation under a

stochastic model of community assembly, indicating that the leaf habitat acts as an ecological

filter selecting for a non-random subset of *Methylobacterium* diversity.

We explored mechanisms explaining the temporal dynamics of *Methylobacterium* diversity at the

scale of a growing season. Because we observed contrasting *Methylobacterium* culturable

diversity between 20 and 30 °C, we suspected that adaptation to temperature variation during the

growing season could explain part of these temporal dynamics. By monitoring *Methylobacterium*

isolate growth under different temperature treatments, we confirmed that temperature affected

isolate growth performances but interestingly, independantly from the temperature at which

isolates were obtained. The fact that most tested isolates also grew slower but more efficiently at

20 °C than at 30 °C (**Figure 5d**), regardless of their phylogenetic and environmental

characteristics, is in line with a temperature-dependent trade-off between growth rate and yield

described in many bacteria (reviewed in (63)). High yield strategies are typical of cooperative

bacterial populations, while fast growth-strategies are typical of competitive populations (63).

These observations also stress the importance of considering incubation temperature when

interpreting results from previous culture-based assessments of *Methylobacterium* diversity.

We provide two lines of evidence that factors other than direct adaptation to temperature drive

Methylobacterium responses to temperature variation, by affecting their growth strategy in

different competitive conditions rather than by affecting their metabolism directly. First, clade

identity was one of the main predictors of overall isolate performance, with some clades (A1, A2,

B) possessing a rapid growth strategy under all temperature conditions, while others (clades A6,

A9, A10) had systematically slower growth. These clade-specific growth strategies could explain

why certain *Methylobacterium* isolates are less competitive and less frequently isolated at higher

temperatures. Still, we cannot rule out that clade-specific growth strategy also reflect

experimental conditions. Second, we observed strong associations between isolate growth

performance and time of sampling, regardless of clade membership, suggesting that growth

strategies also respond to seasonal variations in environmental conditions, and to the level of

establishment and competition in the phyllosphere community (63). These associations are

unlikely to be driven by the direct effects of temperature on metabolic rates because isolation

temperature had little effect on growth strategies, in contrast to clade identity and time of

sampling which had more significant effects. Together, these observations could explain why

isolates from clades A1 and B with fast-growth strategies consistently increase in frequency

during this period due to changes in selection for different ecological strategies, leading to the

homogeneization of the community.

Taken together, our temporal survey of diversity dynamics and screening for growth performance

suggest the following timeline of the dynamics of the *Methylobacterium* phyllosphere

community. At the very beginning of the growing season, a pool of bacteria with mixed

ecological strategies and genotypes colonizes newly emerging leaves. Due to the stochasticity of

this colonization, we initially observe strong dissimilarity among phyllosphere communities,

regardless of their spatial position. During the summer, conditions allow the progressive

establishment of a diverse *Methylobacterium* community with a high yield strategies (63),

dominated by increasingly closely phylogenetically related strains. At the end of the growing

season, with migration, environmental conditions shifting and leaves senescing, isolates with a

fast-growth strategy are able to grow rapidly, dominating the phyllosphere community and

leading to its further homogeneization before leaves fully senesce. This scenario provides an

explanation for the observation of community convergence and increasing homogeneity of

phyllosphere communities throughout the growing season (65, 66).

Our study illustrates that *Methylobacterium* is a complex group of divergent lineages with

different ecological strategies and distributions, reflecting long-term adaptation to contrasting

local environments. Based upon a similar observation, some authors recently proposed to

reclassify *Methylobacterium* group B within a new genus (*Methylorubrum*) that they argue is

ecologically and evolutionarily distinct from other *Methylobacterium* clades (31). Although clade

B was well supported as a distinct clade in our analyses, our results suggest that it is in fact

embedded within clade A, which would render the genus *Methylobacterium* paraphyletic if clade

B is defined as a distinct genus (**Figure S1**). Furthermore, group B was not particularly

ecologically distinct in comparison with other major clades (**Figure 2**). Our results emphasize the

fact that thorough genomic investigations are needed to clarify the taxomonic status of

Methylobacterium. Beyond any taxonomic considerations, neither clade identity assessed by

individual genetic markers nor the tremendous ecological diversity among *Methylobacterium*

clades can predict all of the spatial and temporal variation in *Methylobacterium* diversity in

nature. In order to define the niches of *Methylobacterium* clades and to understand the metabolic

mechanisms underlying their contrasting life strategies, future characterization of their functions

and genome structure will be required using phylogenomic approaches.

In conclusion, we find that *Methylobacterium* adaptive responses to local environmental variation

in the phyllosphere are driven by both long-term inherited ecological strategies that differ among

major clades within the genus, as well by seasonal changes affecting habitat characteristics and

community structure in the phyllosphere habitat. Overall, our study combining culture-free and

culture-based approaches provides novel insights into the factors driving fine-scale adaptation of

microbes to their habitats. In the case of *Methylobacterium,* our approach revealed the particular

importance of considering organismal life-history strategies to help understand the fine-scale

diversity and dynamics of this ecologically important taxon.

References

1.   Vorholt JA. 2012. Microbial life in the phyllosphere. 12. Nature Reviews Microbiology

10:828–840.

2.   Lindow SE, Brandl MT. 2003. Microbiology of the Phyllosphere. Appl Environ

Microbiol 69:1875–1883.

3.   Fürnkranz M, Wanek W, Richter A, Abell G, Rasche F, Sessitsch A. 2008. Nitrogen

fixation by phyllosphere bacteria associated with higher plants and their colonizing epiphytes of a

tropical lowland rainforest of Costa Rica. 5. The ISME Journal 2:561–570.

4.   Laforest-Lapointe I, Messier C, Kembel SW. 2016. Host species identity, site and time

drive temperate tree phyllosphere bacterial community structure. Microbiome 4:27.

5.   Laforest-Lapointe I, Messier C, Kembel SW. 2016. Tree phyllosphere bacterial

communities: exploring the magnitude of intra- and inter-individual variation among host

species. PeerJ 4:e2367.

6.   Noble AS, Noe S, Clearwater MJ, Lee CK. 2020. A core phyllosphere microbiome exists

across distant populations of a tree species indigenous to New Zealand. PLOS ONE

15:e0237079.

7.   Kembel SW, O'Connor TK, Arnold HK, Hubbell SP, Wright SJ, Green JL. 2014.

Relationships between phyllosphere bacterial communities and plant functional traits in a

neotropical forest. PNAS 111:13715–13720.

8.   Shade A, Gregory Caporaso J, Handelsman J, Knight R, Fierer N. 2013. A meta-analysis

of changes in bacterial and archaeal communities with time. 8. The ISME Journal 7:1493–1506.

9.   Malik AA, Martiny JBH, Brodie EL, Martiny AC, Treseder KK, Allison SD. 2020.

Defining trait-based microbial strategies with consequences for soil carbon cycling under climate

change. 1. The ISME Journal 14:1–9.

10.   Moyes AB, Kueppers LM, Pett☐Ridge J, Carper DL, Vandehey N, O'Neil J, Frank AC.

2016. Evidence for foliar endophytic nitrogen fixation in a widely distributed subalpine conifer.

New Phytologist 210:657–668.

11.   Lajoie G, Kembel SW. 2019. Making the Most of Trait-Based Approaches for Microbial

Ecology. Trends in Microbiology 27:814–823.

12.   Nemergut DR, Schmidt SK, Fukami T, O'Neill SP, Bilinski TM, Stanish LF, Knelman

JE, Darcy JL, Lynch RC, Wickey P, Ferrenberg S. 2013. Patterns and Processes of Microbial

Community Assembly. Microbiol Mol Biol Rev 77:342–356.

13.   Cavender☐Bares J, Kozak KH, Fine PVA, Kembel SW. 2009. The merging of

community ecology and phylogenetic biology. Ecology Letters 12:693–715.

14.   Martiny JBH, Jones SE, Lennon JT, Martiny AC. 2015. Microbiomes in light of traits: A

phylogenetic perspective. Science 350.

15.   Tromas N, Taranu ZE, Castelli M, Pimentel JSM, Pereira DA, Marcoz R, Shapiro BJ,

Giani A. 2020. The evolution of realized niches within freshwater Synechococcus.

Environmental Microbiology 22:1238–1250.

16.   Tromas N, Taranu ZE, Martin BD, Willis A, Fortin N, Greer CW, Shapiro BJ. 2018.

Niche Separation Increases With Genetic Distance Among Bloom-Forming Cyanobacteria. Front

Microbiol 9.

17.   Thompson LR, Sanders JG, McDonald D, Amir A, Ladau J, Locey KJ, Prill RJ, Tripathi

A, Gibbons SM, Ackermann G, Navas-Molina JA, Janssen S, Kopylova E, Vázquez-Baeza Y,

González A, Morton JT, Mirarab S, Zech Xu Z, Jiang L, Haroon MF, Kanbar J, Zhu Q, Jin Song

S, Kosciolek T, Bokulich NA, Lefler J, Brislawn CJ, Humphrey G, Owens SM, Hampton-

Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman JA, Clauset A, Stevens RL, Shade A,

Pollard KS, Goodwin KD, Jansson JK, Gilbert JA, Knight R. 2017. A communal catalogue

reveals Earth's multiscale microbial diversity. 7681. Nature 551:457–463.

18. Poretsky R, Rodriguez-R LM, Luo C, Tsementzi D, Konstantinidis KT. 2014. Strengths

and Limitations of 16S rRNA Gene Amplicon Sequencing in Revealing Temporal Microbial

Community Dynamics. PLOS ONE 9:e93827.

19. Ranjan R, Rani A, Metwally A, McGee HS, Perkins DL. 2016. Analysis of the

microbiome: Advantages of whole genome shotgun versus 16S amplicon sequencing.

Biochemical and Biophysical Research Communications 469:967–977.

20. Corpe WA, Rheem S. 1989. Ecology of the methylotrophic bacteria on living leaf

surfaces. FEMS Microbiol Ecol 5:243–249.

21. Keppler F, Hamilton JTG, Braß M, Röckmann T. 2006. Methane emissions from

terrestrial plants under aerobic conditions. 7073. Nature 439:187–191.

22. Knief C, Ramette A, Frances L, Alonso-Blanco C, Vorholt JA. 2010. Site and plant

species are important determinants of the Methylobacterium community composition in the plant

phyllosphere. ISME J 4:719–728.

23. Clarke PH. 1983. The biochemistry of Methylotrophs by C. Anthony Academic Press;

London, New York, 1982 xvi + 432 pages. £24.00, $49.50. FEBS Letters 160:303–303.

24. Anthony C. 1991. Assimilation of carbon by methylotrophs. Biotechnology 18:79–109.

25. Ivanova EG, Doronina NV, Trotsenko YuA. 2001. Aerobic Methylobacteria Are Capable

of Synthesizing Auxins. Microbiology 70:392–397.

26. Madhaiyan M, Poonguzhali S, Lee HS, Hari K, Sundaram SP, Sa TM. 2005. Pink-

pigmented facultative methylotrophic bacteria accelerate germination, growth and yield of

sugarcane clone Co86032 (Saccharum officinarum L.). Biol Fertil Soils 41:350–358.

27. Madhaiyan M, Poonguzhali S, Sa T. 2007. Metal tolerating methylotrophic bacteria

reduces nickel and cadmium toxicity and promotes plant growth of tomato (Lycopersicon

esculentum L.). Chemosphere 69:220–228.

28.   Dourado MN, Aparecida Camargo Neves A, Santos DS, Araújo WL. 2015.

Biotechnological and Agronomic Potential of Endophytic Pink-Pigmented Methylotrophic

Methylobacterium spp. Biomed Res Int 2015.

29.   Ryu JH (Chungbuk NU, Madhaiyan M (Chungbuk NU, Poonguzhali S (Chungbuk NU,

Yim WJ (Chungbuk NU, Indiragandhi P (Chungbuk NU, Kim KA (Chungbuk NU, Anandham R

(Chungbuk NU, Yun JC (National I of AS and T, Kim KH (The U of S, Sa TM (Chungbuk NU.

2006. Plant Growth Substances Produced by Methylobacterium spp. and Their Effect on Tomato

(Lycopersicon esculentum L.) and Red Pepper (Capsicum annuum L.) Growth. Journal of

Microbiology and Biotechnology.

30.   Lee HS, Madhaiyan M, Kim CW, Choi SJ, Chung KY, Sa TM. 2006. Physiological

enhancement of early growth of rice seedlings (Oryza sativa L.) by production of phytohormone

of N2-fixing methylotrophic isolates. Biol Fertil Soils 42:402–408.

31.   Green PN, Ardley JK. 2018. Review of the genus Methylobacterium and closely related

organisms: a proposal that some Methylobacterium species be reclassified into a new genus,

Methylorubrum gen. nov. International Journal of Systematic and Evolutionary Microbiology

68:2727–2748.

32.   Chen W-M, Cai C-Y, Li Z-H, Young C-C, Sheu S-Y. 2019. Methylobacterium

oryzihabitans sp. nov., isolated from water sampled from a rice paddy field. International Journal

of Systematic and Evolutionary Microbiology, 69:3843–3850.

33.   Feng G-D, Chen W, Zhang X-J, Zhang J, Wang S-N, Zhu H. 2020. Methylobacterium

nonmethylotrophicum sp. nov., isolated from tungsten mine tailing. International Journal of

Systematic and Evolutionary Microbiology, 70:2867–2872.

34.   Jia L juan, Zhang K shuai, Tang K, Meng J yu, Zheng C, Feng F ying. 2020.

Methylobacterium crusticola sp. nov., isolated from biological soil crusts. International Journal of

Systematic and Evolutionary Microbiology, 70:2089–2095.

35.   Kim J, Chhetri G, Kim I, Lee B, Jang W, Kim MK, Seo T. 2020. Methylobacterium

terricola sp. nov., a gamma radiation-resistant bacterium isolated from gamma ray-irradiated soil.

International Journal of Systematic and Evolutionary Microbiology, 70:2449–2456.

36.   Kim J, Chhetri G, Kim I, Kim MK, Seo T. 2020. Methylobacterium durans sp. nov., a

radiation-resistant bacterium isolated from gamma ray-irradiated soil. Antonie van Leeuwenhoek

113:211–220.

37.   Ten LN, Li W, Elderiny NS, Kim MK, Lee S-Y, Rooney AP, Jung H-Y. 2020.

Methylobacterium segetis sp. nov., a novel member of the family Methylobacteriaceae isolated

from soil on Jeju Island. Arch Microbiol 202:747–754.

38.   Jiang L, An D, Wang X, Zhang K, Li G, Lang L, Wang L, Jiang C, Jiang Y. 2020.

Methylobacterium planium sp. nov., isolated from a lichen sample. Arch Microbiol 202:1709–

1715.

39.   Pascual JA, Ros M, Martínez J, Carmona F, Bernabé A, Torres R, Lucena T, Aznar R,

Arahal DR, Fernández F. 2020. Methylobacterium symbioticum sp. nov., a new species isolated

from spores of Glomus iranicum var. tenuihypharum. Curr Microbiol 77:2031–2041.

40.   Marx CJ, Bringel F, Chistoserdova L, Moulin L, Farhan Ul Haque M, Fleischman DE,

Gruffaz C, Jourand P, Knief C, Lee M-C, Muller EEL, Nadalig T, Peyraud R, Roselli S, Russ L,

Goodwin LA, Ivanova N, Kyrpides N, Lajus A, Land ML, Médigue C, Mikhailova N, Nolan M,

Woyke T, Stolyar S, Vorholt JA, Vuilleumier S. 2012. Complete Genome Sequences of Six

Strains of the Genus Methylobacterium. J Bacteriol 194:4746–4748.

41.   Tani A, Ogura Y, Hayashi T, Kimbara K. 2015. Complete Genome Sequence of

Methylobacterium aquaticum Strain 22A, Isolated from Racomitrium japonicum Moss. Genome

Announc 3.

42.   Minami T, Ohtsubo Y, Anda M, Nagata Y, Tsuda M, Mitsui H, Sugawara M,

Minamisawa K. 2016. Complete Genome Sequence of Methylobacterium sp. Strain AMS5, an

Isolate from a Soybean Stem. Genome Announc 4.

43.   Morohoshi T, Ikeda T. 2016. Complete Genome Sequence of Methylobacterium populi P-

1M, Isolated from Pink-Pigmented Household Biofilm. Genome Announc 4.

44.   Belkhelfa S, Labadie K, Cruaud C, Aury J-M, Roche D, Bouzon M, Salanoubat M,

Döring V. 2018. Complete Genome Sequence of the Facultative Methylotroph Methylobacterium

extorquens TK 0001 Isolated from Soil in Poland. Genome Announc 6.

45.   Green PN. 2015. Methylobacterium, p. 1–8. *In* Bergey's Manual of Systematics of

Archaea and Bacteria. American Cancer Society.

46.   Charron G, Leducq JB, Bertin C, Dube AK, Landry CR. 2014. Exploring the northern

limit of the distribution of Saccharomyces cerevisiae and Saccharomyces paradoxus in North

America. FEMS Yeast Res 14:281–8.

47.   Vos M, Quince C, Pijl AS, Hollander M de, Kowalchuk GA. 2012. A Comparison of

rpoB and 16S rRNA as Markers in Pyrosequencing Studies of Bacterial Diversity. PLOS ONE

7:e30600.

48.   Ogier J-C, Pagès S, Galan M, Barret M, Gaudriault S. 2019. rpoB, a promising marker for

analyzing the diversity of bacterial communities by amplicon sequencing. BMC Microbiol

19:171.

49.   Ronquist F, Teslenko M, van der Mark P, Ayres DL, Darling A, Höhna S, Larget B, Liu

L, Suchard MA, Huelsenbeck JP. 2012. MrBayes 3.2: Efficient Bayesian Phylogenetic Inference

and Model Choice Across a Large Model Space. Systematic Biology 61:539–542.

50.   Küpfer M, Kuhnert P, Korczak BM, Peduzzi R, Demarta A. 2006. Genetic relationships

of Aeromonas strains inferred from 16S rRNA, gyrB and rpoB gene sequences. International

Journal of Systematic and Evolutionary Microbiology, 56:2743–2751.

51.   Ee R, Madhaiyan M, Ji L, Lim Y-L, Nor NM, Tee K-K, Chen J-W, Yin W-F. 2016.

Chania multitudinisentens gen. nov., sp. nov., an N-acyl-homoserine-lactone-producing

bacterium in the family Enterobacteriaceae isolated from landfill site soil. International Journal of

Systematic and Evolutionary Microbiology, 66:2297–2304.

52.   Frank RAW, Price AJ, Northrop FD, Perham RN, Luisi BF. 2007. Crystal Structure of the

E1 Component of the Escherichia coli 2-Oxoglutarate Dehydrogenase Multienzyme Complex.

Journal of Molecular Biology 368:639–651.

53.   Bell RL, González-Escalona N, Stones R, Brown EW. 2011. Phylogenetic evaluation of

the 'Typhimurium' complex of Salmonella strains using a seven-gene multi-locus sequence

analysis. Infection, Genetics and Evolution 11:83–91.

54.   Dixon P. 2003. VEGAN, a package of R functions for community ecology. Journal of

Vegetation Science 14:927–930.

55.   Redford AJ, Bowers RM, Knight R, Linhart Y, Fierer N. 2010. The ecology of the

phyllosphere: geographic and phylogenetic variability in the distribution of bacteria on tree

leaves. Environmental Microbiology 12:2885–2893.

56.   Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. 2016.

DADA2: High resolution sample inference from Illumina amplicon data. Nat Methods 13:581–

583.

57.   Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO.

2013. The SILVA ribosomal RNA gene database project: improved data processing and web-

based tools. Nucleic Acids Res 41:D590–D596.

58.   Boratyn GM, Camacho C, Cooper PS, Coulouris G, Fong A, Ma N, Madden TL, Matten

WT, McGinnis SD, Merezhuk Y, Raytselis Y, Sayers EW, Tao T, Ye J, Zaretskaya I. 2013.

BLAST: a more efficient report with usability improvements. Nucleic Acids Res 41:W29-33.

59.   Stegen JC, Lin X, Fredrickson JK, Chen X, Kennedy DW, Murray CJ, Rockhold ML,

Konopka A. 2013. Quantifying community assembly processes and identifying features that

impose them. ISME J 7:2069–2079.

60.   Kembel SW, Cowan PD, Helmus MR, Cornwell WK, Morlon H, Ackerly DD, Blomberg

SP, Webb CO. 2010. Picante: R tools for integrating phylogenies and ecology. Bioinformatics

26:1463–1464.

61.   Kembel SW. 2009. Disentangling niche and neutral influences on community assembly:

assessing the performance of community phylogenetic structure tests. Ecology Letters 12:949–

960.

62.   Zwietering MH, Jongenburger I, Rombouts FM, Riet K van 't. 1990. Modeling of the

Bacterial Growth Curve. Appl Environ Microbiol 56:1875–1881.

63.   Lipson DA. 2015. The complex relationship between microbial growth rate and yield and

its implications for ecosystem processes. Front Microbiol 6.

64.   Cavender☐Bares J, Ackerly DD, Baum DA, Bazzaz FA. 2004. Phylogenetic

Overdispersion in Floridian Oak Communities. The American Naturalist 163:823–843.

65.   Copeland JK, Yuan L, Layeghifard M, Wang PW, Guttman DS. 2015. Seasonal

Community Succession of the Phyllosphere Microbiome. MPMI 28:274–285.

66.   Maignien L, DeForce EA, Chafee ME, Eren AM, Simmons SL. Ecological Succession

and Stochastic Variation in the Assembly of Arabidopsis thaliana Phyllosphere Communities.

mBio 5:e00682-13.

Figure 1 - Sampling design. a) Locations of the two sampled forests MSH (green) and SBL

(orange) in the province of Québec (Canada). b) Time line survey in each forest in 2018 (2-4 time

points available per tree). c-d). Detailed map of each forest and each plot within forests (squares;

6 to 10 trees were sampled per plot; see **Dataset S1b**). For each plot, trees are indicated by points

colored according to their taxonomy (color code on bottom left): ABBA (*Abies balsamea*),

ACRU (*Acer rubrum*). ACSA (*Acer saccharum*), OSVI (*Ostrya virginiana*), QURU (*Quercus*

rubra), FAGR (*Fagus grandifolia*), ASPE (*Acer Pennsylvanicum*). Shades of grey indicate

elevation (50 m elevation scale)

Figure 2 - *Methylobacterium* **phylogeny and ecology.** Most of *Methylobacterium* diversity is found

in association with plants, especially in the phyllosphere. Phylogenetic consensus tree (nodal

posterior probabilities indicated next to the branches) from *rpoB* complete nucleotide sequences

available for 153 *Methylobacterium* genomes and rooted on 32 *Methylobacteriaceae* outgroups

(*Microvirga*, *Enterovirga*; no shown; see **Dataset S1a**). For each genome, species name, the

anthropogenic origin (black squares) and/or environmental origin (color code on top right) are

indicated. Groups A, B, C adapted from Green *et* Ardley (31).

Figure 3 - Tests for phylogenetic association of traits with culture-based estimation of

Methylobacterium **diversity.** a) Part of variance (PERMANOVA; x-axis) in *Methylobacterium*

isolated diversity explained by forest of origin, host tree species, sampling date, temperature of

isolation and their interactions (see Venn diagram on top left for color code) in function of

pairwise nucleotide similarity (PS; y-axis; see **Dataset S1j**) in a phylogenetic tree (partial *rpoB*

nucleotide sequences of 187 isolates and 188 *Methylobacteriaceae* reference sequences). b)

Permutation test for node association with forest of origin and temperature of isolation (color

code on top) mapped on the *rpoB* phylogeny (scaled on PS values). Frames in the tree indicate

nodes significantly associated with at least one factor (ANOVA; Bonferroni correction; $p<0.001$:

"***"; $p<0.01$:"**"; $p<0.05$:"*"). For each isolate (names in bold), colored boxes at the tip of the

tree indicate forest of origin and temperature of isolation.

Figure 4 - Short-scale spatial and temporal dynamics of *Methylobacterium* communities

assessed by *rpoB* barcoding. a) A principal component analysis (PCA) on *Methylobacterium*

ASVs relative abundance shows that 179 phyllosphere samples cluster according to forest of

origin (MSH: open triangles, SBL: full triangles) and date of sampling (detail showed only for

MSH). The significant association of 83 and 25 ASVs with forest of origin and/or sampling date,

respectively is shown (points colored according to clade assignation; legend on top right). b)

Spatial and c) temporal autocorrelation analyses conducted in each forest separately. Points

represent Bray-Curtis (*BC*) dissimilarity in function of pairwise geographic (*pDist*; b) or pairwise

time (*pTime*; c) distance separating two communities. For each forest and variable, the predicted

linear regression is indicated (full line: $p<0.001$; dotted line: $p>0.05$; ANOVA). d) *BC* in function

of sampling time for each forest. e) Detail of spatial autocorrelation analyzes in MSH, conducted

for each sampling time point separately. f) Standardized effect size of mean nearest taxon

phylogenetic distance (SES(MNTD)) between forests and across sampling dates. Negative values

of SES(MNTD) indicate communities contain ASVs that are phylogenetically clustered

compared to a null model of stochastic community assembly.

Figure 5 - Analysis of 79 *Methylobacterium* isolate growth performances under 4 different

temperature treatments. a) Average growth curves (growth intensity in function of time) for

each clade (line: mean value; frame: 1/3 of standard deviation; point: average maximal growth).

b) Growth rate (r) in function of yiel (Y). Each point represents the average r/Y values for an

isolate and a temperature treatment (79 isolates x 4 treatments), colored according to clade

membership. Ellipsoides are centered on average values per clade and represent 30% of

confidence interval (standard deviation). c) r (log scale) in function of time at which samples

strains were isolated from were collected, colored according to the forest of origin. Points: real

data; bars: average r value per forest (n=2) and time (n=4) category. d) r in function of Y,

corrected for clade assignement (residuals of the r~Clade and Y~Clade linear regressions). Each

point represents the average r/Y residual values for an isolate and a temperature, colored

according to the monitoring temperature (legend on top right).

Table 1 - PERMANOVA analysis of variance in *Bacteria* and *Methylobacterium* community

diversity. Part of variance in dissimilarity (R^2; Bray-Curtis index) among samples associated

with four factors and their possible interactions (*F*: forest of origin; *D*: date of sampling; *H*: host

tree species; *P*: plot within forest) and their significance are shown (10,000 permutations on ASV

relative abundance, Hellinger transformation; "***": $p<0.00l$; "**": $p<0.01$; "*": $p<0.05$.). For

16S rRNA, *P* was omitted to conserve degrees of freedom.

	Bacteria (*16S rRNA*)		*Methylobacterium* (*rpoB*)	
Samples	**46**		**179**	
Factor	R^2	Pr(>F)	R^2	Pr(>F)
Forest of origin (F)	0.316***	<0.000	0.324***	<0.001
Host tree specie (H)	0.156***	<0.001	0.071***	<0.001
Time of sampling (D)	0.120*	0.016	0.048***	<0.001
Plot within forests (P)	-	-	0.080***	<0.001
F:H	0.020	0.080	0.004	0.110
H:D	0.239	0.217	0.074**	0.028
H:P	-	-	0.043**	0.007
D:P	-	-	0.058	0.455
H:D:P	-	-	0.085	0.052
Residuals	0.150	-	0.213	-

Table 2 - Summary of statistics from autocorrelation analyzes on 179 phyllosphere *Methylobacterium* samples assessed by *rpoB* barcoding (200 ASVs). *Spatial autocorrelation general models* : pairwise dissimilarity between two communities (Bray-Curtis index; *BC*) as a function of pairwise spatial distance separating two sampled trees (*pDist*) and date of sampling (*Date*) and their interaction (*pDist:Date*). *Spatial autocorrelation models per date*: *BC* as a function of pairwise spatial distance (*pDist*). *Temporal autocorrelation: general models*: *BC* as a function of pairwise spatial time separating two sampled trees (*pTime*). For each model, the average and standard deviation of the intercept (mean *BC* value) are indicated. For each factor (*pDist*, *Date*, *pDist:Date* and *pTime*), the average and standard deviation of estimates (slope) are indicated. Significance of estimates was assessed by ANOVA ("***": p<0.00l; "**": p<0.01; "*": p<0.05).

Categories (*n*)	Intercept (sd)		Estimates*10^{-3} (sd)	
Spatial autocorrelation general models: lm(BC~pDist*D)				
Site (within dates)	*BC*	*pDist*	*D*	*pDist:Date*
MSH	0.5965 (0.0107)	**-0.0041 (0.0192)***	**-2.7648 (0.1313)***	**0.0007 (0.0002)**
SBL	0.6493 (0.0097)	0.0157 (0.0145)	**-1.5575 (0.1646)***	0.0000 (0.0002)
Spatial autocorrelation models per date: lm(BC~pDist)				
Site **Date**	*BC*	*pDist*		
MSH 27 Jun.	0.6237 (0.0340)	-0.0425 (0.0725)		
6 Aug.	0.4919 (0.0112)	**0.0503 (0.0192)**		
7 Sept.	0.3746 (0.0059)	**0.0313 (0.0099)**		
18 Oct.	0.2966 (0.0045)	**0.0795 (0.0073)***		
SBL 20 Jun.	0.6868 (0.0146)	0.0082 (0.0216)		
16 Jul.	0.5819 (0.0113)	0.0215 (0.0174)		
16 Aug.	0.5415 (0.0105)	0.0114 (0.0150)		
20 Sept.	0.5222 (0.0089)	0.0145 (0.0130)		
Temporal autocorrelation general models (BC~pTime)				
Site	*BC*	*pTime*		
MSH	0.4086 (0.0032)	**1.0786 (0.0607)***		
SBL	0.5789 (0.0030)	**0.3012 (0.0617)***		

Table 3 - Variance in yield (*Y*) and growth rate (*r*) measured in 79 *Methylobacterium* isolates

grown under 4 temperature treatments. *Y* and *r* values were transformed in log to meet normal

distribution. For each factor following factors: clade (*C*), forest of origin (*F*), host tree species

(*H*), time of sampling (*D*), temperature of incubation during pre-conditioning (T_P) and monitoring

(T_M) steps, temperature of isolation (T_I) and their interactions, significance of *Y* and *r* responses

are shown ("***": p<0.00l; "**": p<0.01; "*": p<0.05; see **Dataset S1o** for details).

	Rate	Yield
Forest (*F*)	0.015**	0.002
Host tree species (*H*)	0.022***	0.013***
Date of sampling (*D*)	0.054***	0.013***
Pre-conditioning temperature (T_P)	0.001	0.014***
Monitoring temperature (T_M)	0.158***	0.020***
Temperature of isolation (T_I)	0.010**	0.006*
Clade (*C*)	0.076***	0.306***
F:D	0.005	0.035***
H:D	0.003	0.019***
H:C	0.005	0.059***
D:C	0.029**	0.028***
F:C	0.011	0.014**
F:TI	0.003	0.021***
H:TI	0.000	0.021***
C:TI	0.024***	0.007
F:H:D	0.012**	0.023***
F:H:C	0.010**	0.002
F:D:C	0.001	0.008**
H:D:C	0.000	0.013***
H:D:TI	0.016***	0.019***
other interactions (sum)	0.177	0.079
Residuals	0.371	0.279

Supplemental material legends

Supplementary Materials and Methods S1 – Detailed materials and methods and

supplementary references.

Figure S1 - ML phylogenetic trees from *sucA* (a) and *rpoB* (b) concatenated hypervariable (HV)

regions and consensus clade tree (c).

Figure S2 - Experimental design of *Methylobacterium* monitoring for growth performance under

four temperature treatments.

Figure S3 - Example of image analysis of *Methylobacterium* monitoring for growth performance

under four temperature treatments.

Figure S4 - Prediction of log normal growth curve, growth rate and yield for 79 isolates

incubated under four temperature treatments.

Figure S5 – *Alphaproteobacteria* (a) and *Methylobacterium* (b) diversity assessed by *rpoB*

barcoding, comparison of *Methylobacterium* diversity assessement from *rpoB* barcoding and *16s*

rRNA barcoding (c) and comparison of *Methylobacterium* diversity assessment from *rpoB*

barcoding and isolation (d).

Table S1 - Primers used to amplify hyper variable regions in genes *sucA* and *rpoB* and sequence

amplification success in 20 *Methylobacterium* isolates from a pilot survey in MSH in august

2017.

Table S2 - *Methylobacterium* diversity assessed by culture-dependant (isolates; *rpoB* sanger

sequencing) and culture-free approaches (*16S rRNA* and *rpoB* barcoding) and comparison

between different methods.

Dataset S1 – a) List of reference *Methylobacteriaceae* genomes used in this study ; b) List of

phyllosphere samples and their deposited accession numbers ; c) List of 80 methylotrophic

isolates from MSH (pilot survey in august 2017) ; d) Clade assignment of 76 *Methylobacterium*

isolates from the 2017 pilot survey based on BLAST ; e) List of isolates and nucleotide sequence

obtained from pilot and timeline surveys, and they deposited accession numbers ; f) List of 167

Methylobacterium isolates from timeline survey (2018) ; g) List of *16s rRNA* and *rpoB* barcoding

libraries and they deposited accession numbers ; h) List of *16s rRNA* ASV nucleotide sequences,

taxonomy and their absolute abundance in 46 phyllosphere samples ; i) Summary of *16s rRNA*

ASV taxonomic assignation ; j) Tests for phylogenetic association of traits with culture-based

estimation of *Methylobacterium* diversity for different phylogenetic depth ; k) List of *rpoB* ASV

nucleotide sequences, taxonomy and their absolute abundance in 184 phyllosphere samples ; l)

Summary of *rpoB* ASV taxonomic assignation ; m) Detail of ANOVA analysis for each

Methobacterium ASV relative abundance ; n) Average rate and yield values for 79

Methylobacterium isolates monitored under four temperature treatments ; o) Detailled ANOVA

results for yield and growth rate.