Title:

A cross-biomes bacterial diversity shed light on ocean-atmosphere microbial transmission.

Authors:

Naama Lang-Yona[1,6], J. Michel Flores[2], Rotem Haviv[1], Adriana Alberti[3,4,7], Julie Poulain[3,4], Caroline Belser[3,4], Miri Trainic[2], Daniella Gat[2], Hans-Joachim Ruscheweyh[5], Patrick Wincker[3,4], Shinichi Sunagawa[5], Yinon Rudich[2], Ilan Koren[2,*], Assaf Vardi[1,*]

Affiliations:

[1]Department of Plant and Environmental Science, Weizmann Institute of Science, 7610001 Rehovot, Israel; [2]Department of Earth and Planetary Sciences, Weizmann Institute of Science, 7610001 Rehovot, Israel; [3]Génomique Métabolique, Genoscope, Institut François Jacob, CEA, CNRS, Univ Evry, Université Paris-Saclay, 91057 Evry, France; [4]Research Federation for the study of Global Ocean Systems Ecology and Evolution, FR2022/Tara Oceans GOSEE, 3 rue Michel-Ange, 75016 Paris, France; [5]Department of Biology and Swiss Institute of Bioinformatics, ETH Zürich, Vladimir-Prelog-Weg 4, 8093 Zürich, Switzerland; [6]Now at: Civil and Environmental Engineering, Technion - Israel Institute of Technology, 3200003 Haifa, Israel. [7]Now at: Université Paris-Saclay, CEA, CNRS, Institute for Integrative Biology of the Cell (I2BC), 91198, Gif-sur-Yvette, France.

*Corresponding Authors:

Assaf Vardi, Department of Plant and Environmental Science, Weizmann Institute of Science, 7610001 Rehovot, Israel, Tel: +972 8 934 2914; assaf.vardi@weizmann.ac.il.

Ilan Koren, Department of Earth and Planetary Sciences, Weizmann Institute of Science, 7610001 Rehovot, Israel, Tel: +972 8 934 2522; Ilan.Koren@weizmann.ac.il.

Abstract

Microbes are ubiquitous in the oceans and the atmosphere, playing essential roles in biogeochemical processes. The bio-exchanges between the two environments can provide important insights into microbial distribution and diversity but are still not well understood. We simultaneously surveyed the genomic diversity of airborne and marine bacterial communities across 15 000 kilometers in the Atlantic and Pacific oceans. Higher variability of microbial community composition was observed in the atmosphere than in the ocean surface waters. In addition, a greater similarity was observed between oceans than their overlaying atmosphere, and between atmospheric samples than with the ocean beneath. We additionally detected a higher coverage rate and relative abundance of marine bacteria in the Pacific atmosphere as compared to the Atlantic, while the dominant fraction in the Atlantic atmosphere was annotated as soil-associated bacteria. This study advances our understanding of microbial dispersion in the ocean, the atmosphere, and the exchange between them, as well as their potential impact on microbial composition, ecology, and biogeochemistry.

Keywords: *Tara* Pacific Aerobiome; Marine bioaerosols; Atmospheric microbiome; Dust-borne bacteria; Microbial spatial distribution.

Introduction

Microbes are ubiquitous in oceanic and atmospheric environments. Within the ocean, they account for approximately 70% of the total marine biomass (1), playing a crucial role in biogeochemical cycles (such as the carbon, nitrogen and sulfur cycles) (2). In the atmosphere, the microbial community comprises a major part of atmospheric bioaerosols (3), but little is known about the factors that affect their diversity, abundance (4), and role in this environment (5).

Understanding the ocean role as a source and sink of microorganisms and the transport of airborne bacteria can provide important insights into microbial spatial distribution, diversity, and the interplay between terrestrial communities and their transmission over oceanic regions. Over land, airborne bacteria are emitted from a wide range of sources, from anthropogenic to natural ecosystems. Over the oceans, bacteria can be locally emitted as sea spray aerosols, generated at the ocean's surface by wind-driven processes, or transferred into the marine atmosphere through long-range transport from terrestrial sources (4). Due to their aerodynamic sizes, it has been hypothesized that all airborne bacteria can disperse globally and may proliferate in any habitat with suitable environmental conditions (6, 7). Thus, it can be expected that the airborne bacterial community aloft a given marine environment would exhibit a large fraction of the local marine microbiome.

Indeed, biodiversity studies indicated a link between microbial community composition and geographic locations (7, 8). Nevertheless, the geo-distribution of microorganisms in the interface between the ocean and the atmosphere is still underexplored, as the type, amount, and efficiency of particle emissions from the ocean, or atmosphere deposition hold large uncertainties. Exploring such distributions and microbial fluxes can improve our understanding of the metabolic capabilities introduced into the ocean and the enrichment of local marine diversity (9, 10). They may also affect

biogeochemical processes upon deposition of bacteria into the ocean or their release to the

atmosphere.

In this study, we present the spatially resolved composition of bacterial communities in the

atmospheric marine boundary layer (AMBL) and the ocean surface along two basin-scale oceanic

transects: The North Atlantic Ocean and the Western Pacific Ocean, covering approximately

15 000 kilometers. The similar sampling times, both placed in the north hemisphere (except for a

few sampling points in the Pacific Ocean, southern to the equatorial), allow us to focus on non-

seasonal changes in the microbial communities across these transects. We first explore microbial

community compositions in the atmospheric and oceanic environments, revealing larger

similarities between different oceans, in contrast to their overlaying atmospheres. By focusing on

specific genera, we track terrestrial-associated bacteria in the airborne community and show distinct

patterns of marine bacterial emission into the atmosphere. We thus suggest that bacterial geo-

distribution theories and models should include constraints between the hydrosphere and

atmosphere.

Materials and Methods

Sampling. Details of the expedition and sampling system used are described extensively in Flores

et al. (11). In short, marine aerosols were collected aboard the *R/V* Tara during the first year of the

Tara Pacific Expedition (12). Airborne particles were collected at ~15 m above sea level (ASL)

during the Atlantic crossing from Lorient, France, to Miami, USA. After Miami, the inlet was

relocated to ~27 m ASL. The inlet was constructed out of conductive tubing of 1.9 cm inner

diameter and a funnel (allowing the collection of all diameters) and mounted on the rear backstay

of Tara. A custom-made aerosol filter system consisting of four 47 mm filter holders and one

vacuum pump (Diaphragm pump ME 16 NT, VACUUBRAND BmbH & Co KG, Wertheim,

Germany) was installed at the end of the inlet and used to collect the marine particles.

The flow through the filter system was 80 lpm (20 lpm for each filter) during the Atlantic crossing

and 120 lpm (30 lpm for each filter) for the Pacific crossing. Three of the four filter holders were

loaded with PVDF filters (47 mm diameter, 0.45 µm pore size, PAL, Port Washington, New York,

USA), and were used for the current study. The flow rates of each filter holder were monitored

continuously and recorded at the beginning and the end of each sampling event. Aerosols were

collected for periods between approximately 12 – 24 hours (see Table S1 for the exact times). The

filters were folded into a 2 ml cryotube and immediately dropped into liquid nitrogen, and these

conditions were maintained while on board. Blank filters were collected by placing filters on the

filter holders, closing the system for a few seconds, reopening the holders, folding the filters into

cryotubes, and dropping into liquid nitrogen. These conditions were maintained while on board.

Thorough validation tests were conducted to verify no boat-originated contamination masked the

airborne bacterial population composition (See SI).

The surface water sample collection is described in detail in Gorsky *et al.* (12). In short, a "Dolphin

sampler", collecting < 2000 µm size particles, and connected to a large volume peristaltic pump

installed on the deck (max flow rate = $3 \, m^3 \, h^{-1}$), was used for water sampling. Each sample endured

for ~ 120 min. Water serial filtration (<0.22, 0.22–3, and 3–20 µm) was performed using 142 mm

diameter polycarbonate filters (Millipore, Burlington, Massachusetts, USA), and the 0.22-3 µm

fraction, where free bacteria are expected to be concentrated, was used for the current study. The

filters were folded into a 5 ml cryotube and immediately dropped into liquid nitrogen, and these

conditions were maintained while on board. The samples were shipped to the laboratory on dry ice

and kept at -80°C until DNA extraction was carried in the laboratory.

Back-Trajectory Analysis. The air mass origins were tracked using the National Oceanic and

Atmospheric Administration HYSPLIT trajectory model and Global Data Assimilation System

meteorological data (13). Although the average residence time for 3 µm particles is of about 4.7

days (5), as considerable terrestrial influence was observed, the model was run to obtain the 240-

hour (10 days) back trajectories using the "Ensemble option" at an endpoint height of 250 m, which

is the minimum height for an optimal configuration of the ensemble. The presented back trajectories

in Fig. 1 are the average of the 27 back trajectories produced by the Ensemble option.

DNA extraction and sequencing. DNA extraction from 0.22-3 μm water filters was performed as

described in Alberti *et al.* (14). Extraction of DNA from the air filters, with minute amounts of

DNA, was carried out using the DNeasy PowerWater Kit (Qiagen, Hilden, Germany), following

thorough optimization procedure for the extraction of low DNA concentrations from air filters (see

SI text and Fig. S8). The processing of the water and air samples was performed separately, with

random DNA extraction order for the different transects, thus preventing cross-contaminations

between the different environments, and geographic location.

The DNA concentrations were evaluated with a Qubit® 3.0 Fluorometer (ThermoFischer,

Massachusetts, USA), using the DeNovix (Wilmington, Delaware, USA) dsDNA Ultra High

Sensitivity Assay. For DNA sequencing, the bacterial V4–V5 region of the 16S rRNA gene (515Fa:

5'–GTGYCAGCMGCCGCGGTAA–3', and 926R: 5'–CCGYCAATTYMTTTRAGTTT–3')(15),

was amplified. A PCR mix of 25 μl was prepared in triplicate using 1X Mytaq mix (Bioline,

London, UK), 0.2 μM primers, 4 μl DNA extract, and PCR-grade water (Merck, Darmstadt,

Germany). A no-template control was included in all runs, as well as DNA from a mock community

(ZymoBIOMICS Microbial Community DNA Standard; Zymo, Irvine, CA, USA). The PCR

products were validated on 1% agarose gel, and triplicates were pooled and sent to Genoscope, the

French National DNA Sequencing Facility. The PCR products of the water samples from the

Pacific transect were sent to the DNA Sequencing Facility at the University of Illinois at Chicago.

DNA sequencing in both facilities was conducted using Illumina MiSeq sequencing technology

(maximum read length of 2×300 base pairs). No batch effects were observed, as validated by mock

community positive control sequencing.

Sequences Processing. Raw amplicon reads provided from the Genoscope sequencing facility

were processed using the Tara Pacific metabarcoding (16S) pipeline up to the read margin stage

using usearch v11(16), followed by the DADA2 pipeline (version 1.12) (17), using R (dada2

package). The merged reads (provided as .fastq files) were trimmed and filtered by removing reads

exceeding the maximum expected error of 2 bp or an ambiguous read. The reads were dereplicated

to acquire unique sequences, which were used to infer sequence variants with the trained error

model. After chimeric sequence removal, amplicon sequence variants (ASVs) were used to assign

taxa. ASVs were taxonomically assigned using dada2, with two steps: classification of sequences

against the SILVA nr version 132 training dataset, followed by an exact matching between ASVs

and the SILVA species assignment dataset (version 132) providing species-level assignment. Raw

reads provided from the sequencing facility of the University of Illinois were also processed using

the dada2 pipeline. Forward and reversed reads were merged after trimming and filtering steps.

Dereplication and annotation processes were carried out as described above. A total of 11 577

bacterial ASVs were obtained after error correction, chimera, Archean, chloroplast, and

mitochondrial DNA removal. Possible shifts in microbial composition due to differences in

analyses were excluded by comparing mock community positive controls sequenced and analyzed

using both methods. The microbial population sequencing analysis was performed only after a

thorough decontamination procedure, to mitigate putative contamination in sequence libraries from

the air samples with low microbial biomass (18), as described in the SI. The raw 16S amplicon

sequences were deposited in the European Nucleotide Archive at EMBL-EBI (accession numbers:

PRJEB39048, and PRJEB38899).

Environmental ontology. Environmental descriptive terms were extracted from the closest

matches (97% identity) using the SEQenv pipeline for Python (version 1.3.0) with default

parameters (19) and ENVO terms. The input data included FASTA files of sequences after quality

control check and removal of blank contaminants per each sample, to be compared to highly similar

sequences from public repositories (such as GenBank, using the NCBI nucleotide data base).
Detailed description of the pipeline flow is given in the SI file. In the current study, the terms were
further clustered into five main groups: marine, terrestrial, fresh water, anthropogenic, and
unclassified, as detailed in Table S7.

Statistical analysis. Statistical significance was analyzed using R (3.5.2) and Origin 2019.
Analysis of similarities (ANOSIM) was used to verify the significance of the nonmetric
multidimensional scaling (NMDS) ordination for taxonomic grouping (using the Bray-Curtis
dissimilarity score, vegan package), and differences in phyla composition between environments.
Analysis of molecular variance (AMOVA) was used to verify significance of the Bray-Curtis
dissimilarity differences of bacterial composition between biomes. Alpha diversity values were
represented by the Shannon index and were tested for differences using analysis of variance
(ANOVA) after normal distribution was verified.

Results

**Regional distribution of airborne and surface water bacterial phyla in the Pacific and Atlantic
oceans.** The two open ocean sailing transects examined in this study were as follows: western
Pacific, sampled in June 2016 (Table S1), starting from Keelung, Taiwan, towards Fiji (Fig. 1A),
and the Atlantic crossing, sampled in May 2017 (Table S1) from Lorient, France, to Miami, USA.
(Fig. 1B). In the water, we found a relatively homogeneous phyla distribution within each transect,
with Proteobacteria dominating both oceans ($58 \pm 3\%$ in the Pacific and $66 \pm 4\%$ in the Atlantic;
Fig. 1C and D, respectively). Cyanobacteria ($29 \pm 2\%$ and $9 \pm 4\%$) and Bacteroidetes ($9 \pm 1\%$ and
$17 \pm 2\%$) were the next two most abundant phyla. When comparing to other marine microbiome
studies, the phyla distribution of the near-surface water environment is similar (2, 20). For example,
the Cyanobacteria to Proteobacteria ratios in our study are 0.49 ± 0.06 and 0.14 ± 0.07 in the Pacific

and Atlantic surface water, respectively. Similarly-calculated ratios characterized in these oceanic

regions are 0.43 and 0.16, for the Pacific (20) and Atlantic (2) regions, respectively.

In the AMBL, Proteobacteria was also the most dominant airborne bacterial phylum in both the

Pacific and Atlantic oceans, with $69 \pm 12\%$ and $64 \pm 8\%$ average percentile abundance, respectively

(Fig. 1E and F). However, we found a more heterogeneous distribution of bacterial phyla than that

in ocean surface water samples (Fig. S1), even when considering an air mass presence of at least

120 hours over an oceanic path prior to sampling (Fig. 1A and 1B, colored lines). Other abundant

phyla in the Pacific AMBL included Cyanobacteria ($11 \pm 9\%$), Bacteroidetes ($8 \pm 3\%$), Firmicutes

($8 \pm 3\%$), Actinobacteria ($4 \pm 3\%$), and Planctomycetes ($2 \pm 1\%$). In the Atlantic AMBL, the

abundant phyla included Actinobacteria ($11 \pm 5\%$), Firmicutes ($10 \pm 6\%$), and Bacteroidetes ($6 \pm$

3%; Fig. 1F), while Cyanobacteria was observed with an average abundance less than 0.5%.

Firmicutes, Actinobacteria, Fusobacteria, Deinococcus-Thermos and Acidobacteria were higher in

abundance in the atmospheric samples of both oceans compared to the waters. Firmicutes were

predominantly observed in the airborne samples ($8 \pm 3\%$ and $10 \pm 6\%$ in the Pacific and Atlantic

AMBL, respectively), with low (<1% in average) to non-significant abundance in the water

samples. Firmicutes were previously detected in the oceanic environment, specifically the Bacillus

genus (21), and Sul *et al.* found in low relative abundance (< 6% on average) across latitudes with

little latitudinal dependence (22). Actinobacteria abundance was also significantly higher in the

Atlantic AMBL compared to the water (*t*-test, *p*-value < 0.0001). Airborne Actinobacteria and

Firmicutes have been previously connected to desert dust samples in the Eastern Mediterranean

(23-25), and detected in different studies of airborne bacteria (26-28). In addition, Firmicutes are

usually more abundant in soils than in marine surface water (2, 21). Therefore, these groups may

represent a terrestrial source.

Members of the airborne-detected Fusobacteria, Deinococcus-Thermos and Acidobacteria phyla

exhibit high physiological diversity in cell shapes and sizes (21, 29). Their presence in remote

locations, such as Antarctica (30), airborne dust (23), and precipitation over the alpine (31),

together with their detection above the western Pacific Ocean, after 120 h transport above the ocean,

suggests that they are ubiquitous in the atmospheric environment.

The local primary production imprint in the AMBL was estimated by calculating the ratio of

autotrophic to heterotrophic bacteria in the atmospheric and oceanic samples (Fig. S2A and b). The

ratios in the Pacific atmosphere were more than an order of magnitude higher than those measured

in the Atlantic atmosphere (mean values: 0.113 ± 0.109 and 0.006 ± 0.009, respectively, two sample

t-test p-value <0.001; Fig. S2A and B). Additionally, the average ratio in the Pacific water is

approximately four times higher than that in the Atlantic (0.392 ± 0.083, compared to $0.103 \pm$

0.052, respectively; two sample t-test p-value < 0.001). Similarly, a significantly higher relative

abundance of cyanobacterial 16S rRNA gene was observed in the Pacific compared to the Atlantic

atmosphere (Fig. S2C and D, Table S3) based on qPCR analysis. The observed difference in

oceanic cyanobacterial abundance is consistent with results reported by Flombaum *et al.*, showing

a higher abundance of marine Synechococcales in the Pacific compared to the Atlantic (33).

In general, the atmospheric phyla composition showed high variations between samples, whereas

in the marine microbiome, a more homogeneous and stable community structure was observed.

This may relate to the lower biomass sampled in the air, thus more prone to changes in community

composition. Furthermore, while the wind-driven surface water currents show connectivity

between oceans with time scales of years, the atmospheric circulation time scales are in the range

of days to weeks (34, 35).

Similarities and differences in the atmospheric and oceanic microbiomes. The marine and

atmospheric microbiomes were further analyzed at the ASV level. We found the variability in the

airborne bacterial diversity and composition to be significantly higher compared to the water

samples (Fig. 2A and B; Two-sample t-test, p-value < 0.001, and ANOSIM, p-value = 0.043,

respectively), with a distinct difference between the air and water ASV composition (Fig. 2C;

AMOVA, *p*-value < 0.01 between the atmosphere and oceanic samples of both Atlantic and Pacific

environments).

The surface water environments shared 166 taxa (comprising 28% of all ocean taxa; Fig. 2D), and

the AMBL biomes shared 341 taxa (comprising 25% of all airborne taxa). Only 78 taxa in the

Atlantic (7% of all Atlantic taxa) and 134 in the Pacific (15% of all Pacific taxa) were shared

between an ocean and its corresponding atmosphere. We found different oceans to have a greater

resemblance to one another than to their overlaying AMBL, and atmospheric samples from distinct

locations (at least 13 000 km apart from each other) share more common taxa than with the ocean

beneath. This suggests that the proximity of the sampled biomes is second in significance to the

type of sampled environment.

A phylogenetic tree based on the bacterial 16S amplicon sequences provides an overview of the

bacterial community and genetic distances between them in the observed marine environment (Fig.

S3). Among the shared groups in the Atlantic and Pacific atmospheric biomes, the main phyla

occurrences included 44% Proteobacteria, 19% Actinobacteria, 19% Firmicutes, and 10%

Bacteroidetes. The high diversity in the common airborne bacterial taxa in both the Atlantic and

Pacific is most likely a consequence of a short turnover time of the air mass, leading to continuously

changing and dynamic community composition in the AMBL.

Spatial distributions of bacteria across the Pacific and Atlantic environments. The spatial

distribution of specific bacteria in the AMBL allows the identification of transported bacteria and

indicates a nonrandom contribution from the marine environment. The bacterial ASVs were

clustered into taxonomic groups, and >5% taxon occurrences were ranked based on the averaged

abundance of the different phyla (Fig. 3 and listed in Tables S4-S6). The most abundant

Proteobacteria detected only in the AMBL was *Paracoccus* (Table S4). *Paracoccus* strains have

been isolated from different environments including soil (36), marine sediments (37), sewage (38),

and have been detected in other atmospheric bacterial studies (39, 40). Some Proteobacteria,

abundant in the air samples, were continuously detected in the surface water (*e.g., Halomonas*,

Pseudomonas, *Idiomarina*, and *Methylobacterium*; Table S5), and thus assumed to be emitted from

the local marine environment.

The most abundant genus in the phylum Actinobacteria was *Actinomarina*, which appeared in both

oceans and in the Pacific air samples (Table S5). The main Firmicutes included Bacillus, known

for their endospores that can remain dormant for years. *Bacillus* is a common bacterium found in

transported desert dust (23), and the deep marine environment (41). Some *Bacillus* species are

known for their unique metabolites and antagonistic activity against pathogens(42). We further

explored the differences between the surface waters and the atmospheric Firmicutes population,

which was dramatically under-represented in the water samples, with a focused phylogenetic tree

targeting only Firmicutes ASVs (Fig. S4). Firmicutes ASVs were rarely detected in the water

samples and differed phylogenetically from the atmospheric ASVs. The rare abundance of marine

bacilli in the water may result from the preferential growth environment of the deep sea, coral and

sediments (41), and their copiotrophic property (*i.e.,* flourishing in environments with high nutrient

availability) (43). Notably, *Tumebacillus*, and *Clostridium saccharobutylicum*, spore forming

Firmicutes, were detected in the Atlantic AMBL. Endospores can survive harsh and dry conditions

and thus might be transported through the air at higher survival rates than others. We also found

ASVs assigned to genera that are known to include human-associated microbes (i.e., *Micrococcus*,

Actinomyces; Table S4). Although not detected in the blank filters, we cannot exclude the

possibility that those taxa may originate from the human activity onboard Tara. Nevertheless, the

abundance of these genera is low, ranging on average between ~0.1 – ~2.7 % of the ASVs per filter.

The most abundant genus in the phylum Bacteroidetes found in the atmospheric samples was

Sediminibacterium (Table S5). This genus was previously found to contribute to the coral

microbiome (44) and detected in air samples over the Great Barrier Reef (26) and the Mediterranean

Sea (45). In this study, it was detected in the water samples of the Pacific Ocean solely, with low

relative abundance ($< 0.01\%$) and spatial coverage.

We continuously detected water-originated species in the air samples, with higher relative

abundance in the Pacific AMBL. One such genus is the Cyanobacteria *Prochlorococcus*,

considered a key and most abundant autotroph (46), found mainly in oligotrophic oceans (47). A

notable difference in the relative abundance and appearance of marine bacteria in the Pacific

compared to the Atlantic AMBL is also seen for different phyla, including Proteobacteria,

Bacteroidetes, Verrucomicrobia, Planctomycetes, etc. (Fig. 3, and Table S5). While the increased

fraction of dust-borne bacteria could partially explain the reduction in relative abundance of the

local marine bacteria, marine-associated taxa were absent from a significantly high fraction of the

sampled Atlantic atmosphere, suggesting other factors may also play a role in the observed

difference between the two AMBLs. A clear case of such difference is seen for the Pelagibacterales

(SAR-11 clade), representing approximately one-third of the oceanic surface water microbial

community (48), and highly abundant in both oceans' samples. Their appearance in the atmospheric

samples is significantly lower (Two-sample *t*-test, *p*-value < 0.0001 in both tests; Fig. S4B).

However, while in the Atlantic AMBL, the spatial coverage is minimal, the Pacific AMBL show

almost a full spatial coverage of SAR-11 ASVs in these samples, with significantly higher relative

abundance than the Atlantic (two-sample *t*-test, *p*-value < 0.001; Tables S5). A reduced aerosolized

fraction of SAR-11 compared to the seawater was also observed in the Arctic Sea by Fahlgren *et*

al. (49). The difference in abundance between the Atlantic and Pacific AMBLs could be related to

properties of the sea-surface microlayer (SML), including thickness, concentration, and chemical

composition, which was shown to differ according to changes in heat exchanges, microbial

composition, oceanic waves, pollution and dust storms (50, 51). However, to fundamentally

characterize the causing factors for the reduced detection of marine bacteria in the Atlantic AMBL,

further investigation is required.

Relationships between bacterial taxa in the marine boundary layer and other ecosystems. To

better understand the environmental affiliation of the sampled microbiome we performed an

environmental ontology (ENVO) analysis on all bacterial ASVs (52). The ENVO analysis allows

a comparison of our dataset to published microbiomes found in other environments, by grouping

all environments where the ASVs were previously detected (Fig. 4 and Table S7 detailing the

annotations grouped into the five environments presented).

The water samples showed 7% nonoceanic-annotated bacterial ASVs in the Pacific compared to

16% in the Atlantic (Fig. S5). Of these, 0.5%, in both environments, were from anthropogenic-

annotated bacteria. In the Atlantic water samples, 9% were annotated as terrestrial bacteria, and

might originate from sedimentation of dust particles or high maritime transportation activity in this

environment (53).

The microbiome in the Pacific AMBL exhibited a higher fraction of marine-annotated bacteria, 47

$\pm 20\%$ compared to $12 \pm 5\%$ in the Atlantic (Fig. 4A), while the Atlantic aerobiome was dominantly

annotated as terrestrial ($59 \pm 16\%$ compared to $35 \pm 18\%$ in the Pacific; Fig. 4B). A previous study

by Mayol *et al.* (54) determined that overall, 25% of the airborne bacteria over the ocean originate

from the marine environment, and 42% originate from terrestrial sources based on

parameterizations of sea spray and deposition flux calculations. Higher soil-borne ENVO

annotation was also observed in Firmicutes-targeted analysis (Fig. S6). Thus, airborne Firmicutes

may originate from terrestrial long-range transport, but the extent in which their sedimentation and

proliferation in the ocean is yet to be determined. Anthropogenic-annotated ASVs were higher in

the Atlantic AMBL, with $19 \pm 6\%$, compared to $9 \pm 6\%$ in the Pacific).

Both environments presented freshwater annotations ($6 \pm 3\%$ and $5 \pm 3\%$ in the Pacific and Atlantic

AMBL, respectively). The contribution of bacteria to the formation of water precipitation is of high

interest, and studies revealed bacterial proteins from *e.g., Pseudomonas* and *Pantoea* sp., identified

in the air samples (Table S4), can promote droplet freezing (55) and even detected bacterial activity

in clouds (28). The freshwater annotation of the AMBL bacteria may indicate the presence of such

bacteria in this environment.

Flores *et al.* (11, 56) have found higher concentrations of larger particles related to the deposition

of mineral dust in aerosol sampled in the Atlantic compared to the Pacific transect. Other studies

report massive dust quantities crossing over the Atlantic Ocean (53), which may introduce bacteria

into this environment. The ENVO annotation retrieved from genomic databases corroborate these

findings and emphasize the vast contribution of terrestrial dust-borne bacteria into the Atlantic

Ocean. Additionally, we detected significantly higher DNA biomass in the Atlantic air filters

compared to the Pacific (average of 639.6 ± 468.2 and 128.4 ± 54.4 pg m^{-3}, respectively, *t*-test, *p*-

value < 0.0001), implying a higher concentration of microbial cells per air volume in this region.

It has been reported that the imprint of dust transport and human activity on the Pacific Ocean is

relatively small (57), and that this area is considered a pristine and remote environment, while the

Atlantic Ocean experiences high loads of dust, and anthropogenic impact, with a fast pace of change

(58). Although with lower rates, the terrestrial annotations observed in the Pacific environment

emphasize the long-range transport and dispersion of bacteria in this remote environment, implying

that it cannot be considered a pristine environment, as observed in the Southern Ocean of Antarctica

(59).

Environmental ontology analysis was further conducted for the shared and unique ASVs of the

Atlantic and Pacific AMBLs (Fig. 4C-E). The unique ASVs detected in the Pacific samples were

mainly annotated as marine (37%) and terrestrial (34%) (1 703 ASVs in total; Fig. 4C). The airborne

Pacific-Atlantic shared bacterial ASVs were also composed mostly of marine (47%) and terrestrial

(32%) annotated bacteria (220 ASVs in total; Fig. 4D). A large percentage of unique Atlantic

airborne bacteria (62%) were annotated as terrestrial, whereas anthropogenic- (16%) and marine-

(15%) annotated bacteria were less abundant (1 511 ASVs in total; Fig. 4E). The ubiquity of

shared species found only in the atmospheric samples of the Atlantic and Pacific oceans

suggest a potentially higher pool of air-resident bacteria with efficient long-range transport

in the atmosphere (7).

It seems that boundaries might be drawn between the atmosphere and hydrosphere, allowing a

nonrandom distribution of species between them. One case is the underrepresentation of marine

bacteria in the Atlantic air, and others are airborne taxa (*e.g.,* Firmicutes species) not detected in

the water samples. Thus, we propose another constraint to the hypothesis, referring to the geo-

distribution of bacteria depending not only on distance and time but also on the chemical and

physical differences between these ecosystems, dictating a selective transport of different bacteria.

Discussion

This study illustrates microbial landscape in the interface between the ocean and the atmospheric

boundary layer and their spatial diversity and suggests mechanisms of dispersal at both local and

global scales (Fig. 5). We observed a clear difference between the atmospheric and oceanic

microbiomes, across thousands of kilometers in the North Atlantic and western Pacific oceans, with

a highly variable airborne microbial composition, even with air masses spending more than 120

hours over the open ocean (Fig. 5A), compared to a more stable, and homogeneous composition of

surface-water microbiome, even across latitudes and different oceans (Fig. 5B). This contrast can

be attributed to the orders of magnitude differences in the characteristic advection and mixing

scales in the two media, with higher stability and longer mixing cycles in the ocean and rapid

changes of days to weeks in the atmosphere.

Furthermore, the Atlantic and Pacific water samples showed greater resemblance to one another

than to the atmosphere atop, and thousands of kilometer distant atmospheric samples shared more

common taxa than with the ocean beneath (Fig. 5C). This suggests that the proximity of the sampled

biomes might be less influential compared to the type of sampled environment (*i.e.,* water *vs.* air).

We additionally detected differences in the relative abundance of marine bacteria in the Pacific

compared to the Atlantic atmosphere, with a significantly reduced spatial coverage in the Atlantic's

(Fig. 5D). This observation may be linked to the dilution effect in the Atlantic AMBL due to the

relatively high terrestrial-associated bacteria in this environment. Nevertheless, the consistent low

appearance of these taxa in the Atlantic air, together with the high sequencing coverage (Fig. S13)

of the air samples, suggests an additional mechanism. Differences in the properties of these

environments and of their SMLs could impact the aerosolization efficiency of local marine bacteria,

and explain the biases in their atmospheric abundance, but this hypothesis requires further

validation. Although reduced in the Atlantic air, the airborne autotrophic cyanobacteria maintained

a similar ratio *vs.* heterotrophs as in the surface waters in both environments (Fig. 5E).

While we found bacterial taxa associated with terrestrial environment (*e.g.,* Firmicutes) to be

significantly present in both oceans' AMBL, we did not detect them in the oceanic surface waters,

suggesting no preferential proliferation there (Fig. 5F). Since prokaryote concentrations in the

atmosphere are orders of magnitude smaller compared to the ocean ($\sim 10^3$-$10^4 \, \mathrm{m}^{-3}$ in the atmosphere

(4, 54) *vs.* $\sim 10^8$-$10^{12} \, \mathrm{l}^{-1}$ in the surface waters (21, 54)), sedimentation of terrestrial-originated

bacteria to the ocean are not expected to induce significant change in the water composition, unless

proliferating. Nevertheless, it cannot be excluded that difference in atmospheric microbial

composition could impact surface water composition and function.

In addition, this study suggests that the local marine bacterial community emitted into the AMBL

is enriched by long-range transported bacteria, associated with particle transport from terrestrial

and anthropogenic environments (Fig. 5G). Thus, even remote locations such as the atmospheric

environment of the western Pacific Ocean cannot be defined as pristine.

Finally, we associated the microbial composition of Atlantic aerosols with dust sources, making

these bacteria suitable biomarkers to trace dust transport (Fig. 5H).

This study depicts a high-resolution spatial diversity of airborne bacteria and their shifts between

the ocean and the atmosphere, at both local and global scales. This interplay between the ocean

surface and atmospheric feedback provides new opportunities for future studies to further explore

the selective properties of marine microbes within both environments and how they in turn may

affect biogeochemical cycles.

Acknowledgments

This research was supported by a research grant from Scott Jordan and Gina Valdez, the De Botton

for Marine Science, the Yeda-Sela Center for Basic Research, and the Sustainability and Energy

Research Initiative (SAERI). We are grateful to the following institutions for their financial and

scientific support in Tara Pacific expedition: CNRS, PSL, CSM, EPHE, Genoscope/CEA, and

France Génomique funding (ANR-10-INBS-09-08), Inserm, Université Côte d'Azur, ANR, agnès

b., UNESCO-IOC, the Veolia Environment Foundation, Région Bretagne, Serge Ferrari,

Billerudkorsnas, Amerisource Bergen Company, Lorient Agglomeration, Oceans by Disney, the

Prince Albert II de Monaco Foundation, L'Oréal, Biotherm, France Collectivités, Kankyo Station,

Fonds Français pour l'Environnement Mondial (FFEM), Etienne BOURGOIS, and the Tara

Foundation teams and crew. The authors particularly thank Dr. Serge Planes, Prof. Denis Allemand,

and the Tara Pacific consortium for their management and coordination. The authors gratefully

acknowledging Prof. Daniel Sher for providing Prochlorococcus cultures for qPCR quantification,

and Dr. Flora Vincent for her helpful, critical feedback on the manuscript. NLY acknowledges

support from the Women Bridging position and the Sustainability and Energy Research Initiative

(SAERI), Weizmann Institute of Science. YR acknowledges support from the Israel Science

Foundation (grant #236/16). SSu acknowledges support from the ETH and Helmut Horten

Foundation. This is publication number # 15 of the Tara Pacific Consortium.

Conflict of interest: Authors declare no conflict of interests.

References

1. Bar-On YM, Phillips R, Milo R. The biomass distribution on Earth. Proc Natl Acad Sci U S A. 2018;115(25):6506-11.

2. Sunagawa S, Coelho LP, Chaffron S, Kultima JR, Labadie K, Salazar G, et al. Structure and function of the global ocean microbiome. Science. 2015;348(6237):1261359.

3. Ruiz-Gil T, Acuña JJ, Fujiyoshi S, Tanaka D, Noda J, Maruyama F, et al. Airborne bacterial communities of outdoor environments and their associated influencing factors. Environment International. 2020;145:106156.

4. Fröhlich-Nowoisky J, Kampf CJ, Weber B, Huffman JA, Pöhlker C, Andreae MO, et al. Bioaerosols in the Earth system: Climate, health, and ecosystem interactions. Atmos Res. 2016;182:346-76.

5. Burrows SM, Elbert W, Lawrence MG, Pöschl U. Bacteria in the global atmosphere – Part 1: Review and synthesis of literature data for different ecosystems. Atmos Chem Phys. 2009;9(23):9263-80.

6. Baas Becking LGM. Geobiologie of inleiding tot de milieukunde. 1934.

7. Martiny JBH, Bohannan BJM, Brown JH, Colwell RK, Fuhrman JA, Green JL, et al. Microbial biogeography: putting microorganisms on the map. Nat Rev Microbiol. 2006;4(2):102-12.

8. Martin K, Schmidt K, Toseland A, Boulton CA, Barry K, Beszteri B, et al. The biogeographic differentiation of algal microbiomes in the upper ocean from pole to pole. Nature Com. 2021;12(1):5483.

9. Rahav E, Paytan A, Mescioglu E, Galletti Y, Rosenfeld S, Raveh O, et al. Airborne Microbes Contribute to N2 Fixation in Surface Water of the Northern Red Sea. Geophys Res Lett. 2018;45(12):6186-94.

10. Tang W, Llort J, Weis J, Perron MMG, Basart S, Li Z, et al. Widespread phytoplankton blooms triggered by 2019–2020 Australian wildfires. Nature. 2021;597(7876):370-5.

11. Flores JM, Bourdin G, Altaratz O, Trainic M, Lang-Yona N, Dzimban E, et al. Tara Pacific expedition's atmospheric measurements. Marine aerosols across the Atlantic and Pacific Oceans Overview and Preliminary results. Bull Am Meteorol Soc. 2020;101(5):536-54.

12. Gorsky G, Bourdin G, Lombard F, Pedrotti ML, Audrain S, Bin N, et al. Expanding Tara Oceans Protocols for Underway, Ecosystemic Sampling of the Ocean-Atmosphere Interface During Tara Pacific Expedition (2016–2018). Front Mar Sci. 2019;6(750).

13. Stein AF, Draxler RR, Rolph GD, Stunder BJB, Cohen MD, Ngan F. NOAA's HYSPLIT Atmospheric Transport and Dispersion Modeling System. Bull Am Meteorol Soc. 2016;96(12):2059-77.

14. Alberti A, Poulain J, Engelen S, Labadie K, Romac S, Ferrera I, et al. Viral to metazoan marine plankton nucleotide sequences from the Tara Oceans expedition. Sci Dat. 2017;4(1):170093.

15. Parada AE, Needham DM, Fuhrman JA. Every base matters: assessing small subunit rRNA primers for marine microbiomes with mock communities, time series and global field samples. Environ Microbiol. 2016;18(5):1403-14.

16. Edgar RC. Search and clustering orders of magnitude faster than BLAST. Bioinformatics. 2010;26(19):2460-1.

17. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJA, Holmes SP. DADA2: High-resolution sample inference from Illumina amplicon data. Nat Methods. 2016;13(7):581-3.

18. Eisenhofer R, Minich JJ, Marotz C, Cooper A, Knight R, Weyrich LS. Contamination in Low Microbial Biomass Microbiome Studies: Issues and Recommendations. Trends Microbiol. 2019;27(2):105-17.

19. Ijaz AZ, Jeffries TC, Ijaz UZ, Hamonts K, Singh BK. Extending SEQenv: a taxa-centric approach to environmental annotations of 16S rDNA sequences. PeerJ. 2017;5:e3827-e.

20. DeLong EF, Preston CM, Mincer T, Rich V, Hallam SJ, Frigaard N-U, et al. Community Genomics Among Stratified Microbial Assemblages in the Ocean's Interior. Science. 2006;311(5760):496.

21. Rodrigues TB, Silva AET. Molecular Diversity of Environmental Prokaryotes: CRC Press; 2016.

22. Sul WJ, Oliver TA, Ducklow HW, Amaral-Zettler LA, Sogin ML. Marine bacteria exhibit a bipolar distribution. Proc Natl Acad Sci U S A. 2013;110(6):2342-7.

23. Gat D, Mazar Y, Cytryn E, Rudich Y. Origin-Dependent Variations in the Atmospheric Microbiome Community in Eastern Mediterranean Dust Storms. Environ Sci Technol. 2017;51(12):6709-18.

24. Rahav E, Belkin N, Paytan A, Herut B. The Relationship between Air-Mass Trajectories and the Abundance of Dust-Borne Prokaryotes at the SE Mediterranean Sea. Atmosphere. 2019;10(5):280.

25. Lang-Yona N, Öztürk F, Gat D, Aktürk M, Dikmen E, Zarmpas P, et al. Links between airborne microbiome, meteorology, and chemical composition in northwestern Turkey. Sci Total Environ. 2020;725:138227.

26. Archer SDJ, Lee KC, Caruso T, King-Miaow K, Harvey M, Huang D, et al. Air mass source

determines airborne microbial diversity at the ocean–atmosphere interface of the Great Barrier Reef

marine ecosystem. Isme j. 2019;14:871–6.

27. Maki T, Lee KC, Kawai K, Onishi K, Hong CS, Kurosaki Y, et al. Aeolian Dispersal of

Bacteria Associated With Desert Dust and Anthropogenic Particles Over Continental and Oceanic

Surfaces. J Geophys Res- Atmos. 2019;124(10):5579-88.

28. Amato P, Besaury L, Joly M, Penaud B, Deguillaume L, Delort A-M. Metatranscriptomic

exploration of microbial functioning in clouds. Sci Rep. 2019;9(1):4383.

29. Quaiser A, Ochsenreiter T, Lanz C, Schuster SC, Treusch AH, Eck J, et al. Acidobacteria form

a coherent but highly diverse group within the bacterial domain: evidence from environmental

genomics. Mol Microbiol. 2003;50(2):563-75.

30. Archer SDJ, Lee KC, Caruso T, Maki T, Lee CK, Cary SC, et al. Airborne microbial transport

limitation to isolated Antarctic soil habitats. Nature Microbiol. 2019;4(6):925-32.

31. Caliz J, Triado-Margarit X, Camarero L, Casamayor EO. A long-term survey unveils strong

seasonal patterns in the airborne microbiome coupled to general and regional atmospheric

circulations. Proc Natl Acad Sci U S A. 2018;115(48):12229-34.

32. Oh KH, Jeong DH, Shin SH, Cho YC. Simultaneous quantification of cyanobacteria and

Microcystis spp. using real-time PCR. J Microbiol Biotechnol. 2012;22(2):248-55.

33. Flombaum P, Gallegos JL, Gordillo RA, Rincón J, Zabala LL, Jiao N, et al. Present and future

global distributions of the marine Cyanobacteria *Prochlorococcus* and *Synechococcus*. Proc Natl

Acad Sci U S A. 2013;110(24):9824-9.

34. DeVries T, Primeau F. Dynamically and Observationally Constrained Estimates of Water-

Mass Distributions and Ages in the Global Ocean. J Phys Oceanogr. 2011;41(12):2381-401.

35. Seinfeld JH, Pandis SN. Atmospheric chemistry and physics : from air pollution to climate

change2016.

36. Siller H, Rainey FA, Stackebrandt E, Winter J. Isolation and characterization of a new gram-

negative, acetone-degrading, nitrate-reducing bacterium from soil, Paracoccus solventivorans sp.

nov. Int J Syst Bacteriol. 1996;46(4):1125-30.

37. Lee JH, Kim YS, Choi TJ, Lee WJ, Kim YT. Paracoccus haeundaensis sp. nov., a Gram-

negative, halophilic, astaxanthin-producing bacterium. Int J Syst Evol Microbiol. 2004;54(Pt

5):1699-702.

38. Liu XY, Wang BJ, Jiang CY, Liu SJ. Paracoccus sulfuroxidans sp. nov., a sulfur oxidizer from

activated sludge. Int J Syst Evol Microbiol. 2006;56(Pt 11):2693-5.

39. DeLeon-Rodriguez N, Lathem TL, Rodriguez-R LM, Barazesh JM, Anderson BE, Beyersdorf
AJ, et al. Microbiome of the upper troposphere: Species composition and prevalence, effects of
tropical storms, and atmospheric implications. Proc Natl Acad Sci U S A. 2013;110(7):2575-80.
40. Mazar Y, Cytryn E, Erel Y, Rudich Y. Effect of Dust Storms on the Atmospheric Microbiome
in the Eastern Mediterranean. Environ Sci Technol. 2016;50(8):4194-202.
41. Ivanova EP, Vysotskii MV, Svetashev VI, Nedashkovskaya OI, Gorshkova NM, Mikhailov
VV, et al. Characterization of Bacillus strains of marine origin. Int Microbiol. 1999;2(4):267-71.
42. Mondol MAM, Shin HJ, Islam MT. Diversity of secondary metabolites from marine Bacillus
species: chemistry and biological activity. Mar Drugs. 2013;11(8):2846-72.
43. Fierer N, Bradford MA, Jackson RB. Toward an ecological classification of soil bacteria.
Ecology. 2007;88(6):1354-64.
44. Pootakham W, Mhuantong W, Yoocha T, Putchim L, Sonthirod C, Naktang C, et al. High
resolution profiling of coral-associated bacterial communities using full-length 16S rRNA
sequence data from PacBio SMRT sequencing system. Sci Rep. 2017;7(1):2774.
45. Mescioglu E, Rahav E, Belkin N, Xian P, Eizenga JM, Vichik A, et al. Aerosol Microbiome
over the Mediterranean Sea Diversity and Abundance. Atmosphere. 2019;10(8):440.
46. Chisholm SW, Olson RJ, Zettler ER, Goericke R, Waterbury JB, Welschmeyer NA. A novel
free-living prochlorophyte abundant in the oceanic euphotic zone. Nature. 1988;334(6180):340-3.
47. Chisholm SW, Frankel SL, Goericke R, Olson RJ, Palenik B, Waterbury JB, et al.
Prochlorococcus marinus nov. gen. nov. sp.: an oxyphototrophic marine prokaryote containing
divinyl chlorophyll a and b. Arch Microbiol. 1992;157(3):297-300.
48. Morris RM, Rappe MS, Connon SA, Vergin KL, Siebold WA, Carlson CA, et al. SAR11 clade
dominates ocean surface bacterioplankton communities. Nature. 2002;420(6917):806-10.
49. Fahlgren C, Gomez-Consarnau L, Zabori J, Lindh MV, Krejci R, Martensson EM, et al.
Seawater mesocosm experiments in the Arctic uncover differential transfer of marine bacteria to
aerosols. Environ Microbiol Rep. 2015;7(3):460-70.
50. Wurl O, Ekau W, Landing W, Zappa C. Sea surface microlayer in a changing ocean – A
perspective. Elementa-Sci Anthrop. 2017;5.
51. Michaud JM, Thompson LR, Kaul D, Espinoza JL, Richter RA, Xu ZZ, et al. Taxon-specific
aerosolization of bacteria and viruses in an experimental ocean-atmosphere mesocosm. Nature
Com. 2018;9(1):2017.
52. Buttigieg PL, Morrison N, Smith B, Mungall CJ, Lewis SE, the EC. The environment
ontology: contextualising biological and biomedical entities. J Biomed Semant. 2013;4(1):43.

53.   Kaufman YJ, Koren I, Remer LA, Tanré D, Ginoux P, Fan S. Dust transport and deposition

observed from the Terra-Moderate Resolution Imaging Spectroradiometer (MODIS) spacecraft

over the Atlantic Ocean. J Geophys Res- Atmos. 2005;110(D10).

54.   Mayol E, Arrieta JM, Jiménez MA, Martínez-Asensio A, Garcias-Bonet N, Dachs J, et al.

Long-range transport of airborne microbes over the global tropical and subtropical ocean. Nature

Com. 2017;8(1):201.

55.   Huang S, Hu W, Chen J, Wu Z, Zhang D, Fu P. Overview of biological ice nucleating particles

in the atmosphere. Environment International. 2021;146:106197.

56.   Flores JM, Bourdin G, Kostinski A, Altaratz O, Guy D, Lombard F, et al. Diel cycle of sea

spray aerosol concentration over vast areas of the tropical Pacific Ocean and the Caribbean Sea.

Nature Com. Accepted.

57.   Jones KR, Klein CJ, Halpern BS, Venter O, Grantham H, Kuempel CD, et al. The Location

and Protection Status of Earth's Diminishing Marine Wilderness. Curr Biol. 2018;28(15):2506-

12.e3.

58.   Halpern BS, Frazier M, Afflerbach J, Lowndes JS, Micheli F, O'Hara C, et al. Recent pace of

change in human impact on the world's ocean. Sci Rep. 2019;9(1):11609.

59.   Uetake J, Hill TCJ, Moore KA, DeMott PJ, Protat A, Kreidenweis SM. Airborne bacteria

confirm the pristine nature of the Southern Ocean boundary layer. Proc Natl Acad Sci U S A.

2020:202000134.

60.   de Leeuw G, Guieu Cc, Arneth A, Bellouin N, Bopp L, Boyd PW, et al. Ocean–Atmosphere

Interactions of Particles.   Ocean-Atmosphere Interactions of Gases and Particles. Berlin,

Heidelberg: Springer Earth System Sciences; 2014. p. 171–246.

 Figures legends

Figure 1. Regional distribution of airborne bacterial phyla above the Pacific and Atlantic

oceans. Air-sampling locations and back trajectories, with height above mean sea levels (MSL)

indicated with color scale for the Pacific (A) and Atlantic (B) atmospheric samples. Phyla relative

abundance in the water (C and D) and the air (E and F) for taxa observed in > 5% of the samples

in the Atlantic and Pacific transects, respectively.

Figure 2. Similarities and differences in bacterial microbiomes of different oceanic

environments. Biome-based diversity (based on the Inverse Simpson index) of amplicon sequence

variants (ASVs) detected in more than 5% of the samples (A) and clusters represented by nonmetric

multidimensional scaling (NMDS) ordination, based on Bray-Curtis dissimilarity metrics (B), with

95% confidence ellipses. The distance in ASV composition is presented in a heatmap of the Bray-

Curtis dissimilarity between all analyzed samples (Bray-Curtis index varies between 0, for an

identical ASV composition, and 1, for most distant ASVs on the samples) (C). A Venn diagram

(*VennDiagram* 1.6.20) summarizes the number and the percentage out of the total number of

bacterial taxa that were observed in the different biomes (D).

Figure 3. Spatial distribution of specific airborne microorganisms across the Atlantic and

Pacific transects. The 16S amplicon sequence variants (ASVs) of Pacific air, Atlantic Air, Pacific

water, and Atlantic water samples are categorized based on abundance for each phylum. ASVs

were aggregated according to sequences and taxa annotation, and ASVs covering > 5% of all four

environments are presented. Phyla are distinguished based on color code.

Figure 4. Association of airborne bacterial communities with other environments. The relative

environmental ontology (ENVO) annotation of all amplicon sequence variants (ASVs) identified

in the air samples clustered into five main groups (detailed terms are listed in Table S6) for the

Pacific (A) and Atlantic (B) transects, as well as for Pacific air-unique ASVs (C), Pacific and

Atlantic air-shared ASVs (D), and Atlantic air-unique ASVs (E).

Figure 5. Trends and transitions in bacterial abundance and richness between marine and

atmospheric environments. The main findings of this study are presented as a conceptual model:

Residence time induces fluctuations in microbial composition rather than physical mixing (A, and

B). The microbial compositions of similar environments (*e.g.,* Atlantic and Pacific Oceans) are

more similar than geographical relations (*e.g.,* Pacific atmosphere and surface water; C). The

marine bacterial aerosolization efficiency is different for different oceans (D), and aerosolized

autotrophs kept a higher ratio *vs*. heterotrophs in the Pacific compared to the Atlantic, similarly as

in the water (E). The atmospheric contribution to marine ecology was not detected in the framework

of this study (F), but it cannot be excluded that difference in atmospheric microbial composition

could impact deposition and surface water composition and function. A terrestrial signature was

detected in a remote oceanic environment (G), with a significant contribution of dust-associated

bacteria in the Atlantic atmospheric marine boundary layer (H).

Pacific

Atlantic

Relative abundance

Phyla
- Rare
- Lentisphaerae
- Chloroflexi
- Acidobacteria
- Fusobacteria
- Deinococcus Thermus
- Kiritimatiellaeota
- Planctomycetes
- Verrucomicrobia
- Cyanobacteria
- Bacteroidetes
- Firmicutes
- Actinobacteria
- Proteobacteria

A Higher fluctuations in atmospheric diversity

F Proliferation of Air-borne bacteria in the oceans

G Terrestrial signature of bacterial taxa over oceans

H Dust-associated bacteria

E Airborne cyanobacterial fraction similar to marine borne

B More homogeneous marine compared to atmospheric diversity

D Selectivity in emission of marine bacteria

C Community composition between oceans is closely similar compared to the overlaying atmospheres

Pacific

Atlantic