Hepatic MIR20B promotes nonalcoholic fatty liver disease by suppressing *PPARA*

Yo Han Lee[1,5], Hyun-Jun Jang[1,5], Sounkou Kim[1,5], Sun Sil Choi[1], Keon Woo Khim[1], Hye-Jin Eom[1], Jimin Hyun[1], Kyeong Jin Shin[1], Young Chan Chae[1], Hongtae Kim[1], Jiyoung Park[1], Neung Hwa Park[2], Chang-Yun Woo[3], Chung Hwan Hong[4], Eun Hee Koh[3], Dougu Nam[1,*], Jang Hyun Choi[1,*]

[1]Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan, 44919, Republic of Korea

[2]Department of Internal Medicine, University of Ulsan College of Medicine, Ulsan University Hospital, Ulsan, 44033, Republic of Korea

[3]Department of Internal Medicine, Asan Medical Center, University of Ulsan College of Medicine, Seoul, 05505, Republic of Korea

[4]Department of Medical Science, Asan Medical Center, University of Ulsan College of Medicine, Seoul, 05505, Republic of Korea.

[5]These authors contributed equally to this work.

[*]**Corresponding Author**

Dougu Nam, Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan, Republic of Korea. Tel.: +82 52 217 2525; Fax: +82 52 217 2639

E-mail: dougnam@unist.ac.kr (D. Nam)

Jang Hyun Choi, Department of Biological Sciences, Ulsan National Institute of Science and Technology (UNIST), Ulsan, Republic of Korea. Tel.: +82 52 217 2543; Fax: +82 52 217 3219

E-mail: janghchoi@unist.ac.kr (J.H. Choi)

Abstract

Background:

Non-alcoholic fatty liver disease (NAFLD) is characterized by excessive lipid accumulation and

imbalances in lipid metabolism in the liver. Although nuclear receptors (NRs) play a crucial role in

hepatic lipid metabolism, the underlying mechanisms of NR regulation in NAFLD remain largely

unclear.

Methods:

Using network analysis and RNA-seq to determine the correlation between NRs and microRNA in

human NAFLD patients, we revealed that MIR20B specifically targets *PPARA*. MIR20B mimic and

anti-MIR20B were administered to human HepG2 and Huh-7 cells and mouse primary hepatocytes as

well as high fat diet (HFD)- or methionine-deficient diet (MCD)-fed mice to verify the specific function

of MIR20B in NAFLD. We tested the inhibition of the therapeutic effect of a PPARα agonist,

fenofibrate, by MIR20B and the synergic effect of combination of fenofibrate with anti-MIR20B.

Results:

We revealed that MIR20B specifically targets *PPARA* through miRNA regulatory network analysis of

nuclear receptor genes in NAFLD. The expression of *MIR20B* was upregulated in free fatty acid (FA)-

treated hepatocytes and the livers of both obesity-induced mice and NAFLD patients. Overexpression

of *MIR20B* significantly increased hepatic lipid accumulation and triglyceride levels. Furthermore,

MIR20B significantly reduced FA oxidation and mitochondrial biogenesis by targeting *PPARA*. In

MIR20B-introduced mice, the effect of fenofibrate to ameliorate hepatic steatosis was significantly

suppressed. Finally, inhibition of MIR20B significantly increased FA oxidation and uptake, resulting in

improved insulin sensitivity and a decrease in NAFLD progression. Moreover, combination of

fenofibrate and anti-*MIR20B* exhibited the synergic effect on improvement of NAFLD in MCD-fed mice.

Conclusions:

Taken together, our results demonstrate that the novel MIR20B targets *PPARA*, plays a significant

role in hepatic lipid metabolism, and present an opportunity for the development of novel therapeutics

for NAFLD.

Funding:

This research was funded by Korea Mouse Phenotyping Project (2016M3A9D5A01952411), the

National Research Foundation of Korea (NRF) grant funded by the Korea government

(2020R1F1A1061267, 2018R1A5A1024340, NRF-2021R1I1A2041463, 2020R1I1A1A01074940), and

the Future-leading Project Research Fund (1.210034.01) of UNIST.

Introduction

Obesity has emerged as a host of metabolic disorders, such as non-alcoholic fatty liver disease

(NAFLD). Many reports have demonstrated that 90% of obese patients in the United States have

NAFLD ranging from hepatic steatosis to much more severe forms of non-alcoholic steatohepatitis

(NASH), which can induce fibrosis, cirrhosis, and hepatocellular carcinoma (HCC)(Corey & Kaplan,

2014). NAFLD is associated with hepatic metabolic reprogramming that leads to excessive lipid

accumulation and imbalances in lipid metabolism in the liver(de Alwis & Day, 2008). Hepatic lipid

homeostasis is appropriately described as a complex machinery involving signaling and

transcriptional pathways, as well as targeted genes associated with fatty acid (FA) oxidation and

lipogenesis(Fabbrini, Sullivan, & Klein, 2010). Although the pathogenesis of NAFLD has been widely

studied for years, the molecular mechanisms underlying its complicated disorder are still being

investigated.

Nuclear receptors (NRs) are a superfamily of ligand-activated transcription factors that regulate

biological homeostasis(McKenna et al., 2009). Recent evidence suggests that NRs are key regulators

in the progression of diverse hepatic diseases associated with glucose and lipid metabolism,

inflammation, bile acid homeostasis, fibrosis, and cancer development in the liver(Lopez-Velazquez,

Carrillo-Cordova, Chavez-Tapia, Uribe, & Mendez-Sanchez, 2012). Among them, growing evidence

suggests a link between PPARα and obesity-induced NAFLD. Hepatic PPARα plays an important role

in energy homeostasis by regulating the expression of genes required for FA uptake, FA oxidation,

and triglyceride (TG) hydrolysis in the liver(Chakravarthy et al., 2005). The decreased expression of

PPARα is significantly associated with severity in NAFLD patients(Francque et al., 2015). Therefore,

understanding the molecular mechanism underlying PPARα regulation is critical for understanding the

pathogenesis of NAFLD.

MicroRNAs (miRNAs) are short, non-coding RNA molecules with a length of 18-25 nucleotides that

play an important role in regulating the expression of target genes in a post-transcriptional manner by

targeting base-pairing with the 3'UTR of specific target mRNAs, inhibiting translation, or mRNA

degradation(Bartel, 2004). These miRNAs contribute to the regulation of a wide variety of cellular

functions and metabolic homeostasis, including fatty acid metabolism. Recent studies have suggested

that miRNAs significantly regulate the pathogenesis of NAFLD by targeting the nuclear

receptors(Lopez-Sanchez, Dominguez-Perez, Uribe, Chavez-Tapia, & Nuno-Lambarri, 2021).

Previous report has demonstrated that MIR20B, a member of the MIR17 family, presents in the

circulating plasma of NAFLD patients and has been highlighted as a novel biomarker of NAFLD and

type 2 diabetes mellitus (T2DM) for the diagnosis and risk estimation of NAFLD(Ye et al., 2018).

However, the mechanisms underlying the involvement of MIR20B in the occurrence and progression

of NAFLD remain unknown. In this study, we analyzed the regulatory networks of miRNAs for NR

genes and RNA-seq data in NAFLD patients, which prioritized MIR20B as a key regulator in NAFLD.

Materials and Methods

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
Genetic reagent (*Mus. Musculus*)	C57BL/6JbomTac	DBL	RRID:IMSR_TAC:b6jbom	
cell line (*Homo sapiens*)	HepG2	ATCC	HB-8065, RRID:CVCL_0027	
cell line (*Homo sapiens*)	Huh-7	Dr. Yoshiharu Matsuura; originally from Japanese Collection of Research	JCRB0403, RRID: CVCL_0336	

		Bioresources Cell Bank		
Antibody	Anti-PPARα (Rabbit Polyclonal)	abcam	Cat# ab24509, RRID: AB_448110	WB (1:1000)
Antibody	Anti-HSP90 (Rabbit Polyclonal)	Cell Signaling Technology	Cat# 4877S, RRID: AB_2233307	WB (1:1000)
Sequence-based reagent	MIR20B/Mir20b (MIR20B/Mir20b mimic)	GenePharma	N/A	Sequence: CAAAGUGCUCAUAGUGCAGGUAG
Sequence-based reagent	anti-MIR20B/Mir20b (MIR20B/Mir20b inhibitor)	GenePharma	N/A	Sequence: CUACCUGCACUAUGAGCACUUUG
Sequence-based reagent	*PPARA* siRNA	GenePharma	N/A	Sequence: CGGCGAGGATAGTTCTGGAAGCTTT
Sequence-based reagent	*PPARA* shRNA	Sigma-Aldrich	N/A	Sequence: GAACAGAAACAAATGCCAGTA Sequence: GTAGCGTATGGAAATGGGTTT
Sequence-based reagent	Primers for qPCR	This paper	N/A	
Recombinant DNA reagent	psiCHECK-2-*PPARA*-WT (plasmid)	This paper	N/A	
Recombinant DNA reagent	psiCHECK-2-*PPARA*-Mut (plasmid)	This paper	N/A	

Recombinant DNA reagent	pOTTC385-pAAV CMV-IE IRES EGFP-*Mir20b*	This paper	N/A	
Recombinant DNA reagent	pOTTC385-pAAV CMV-IE IRES EGFP-anti-*Mir20b*	This paper	N/A	
commercial assay or kit	Dual-Luciferase kit	Promega	Cat# E1910	
commercial assay or kit	RNeasy mini kit	Qiazen	Cat# 74004	
commercial assay or kit	QuickChange II Site-Directed Mutagenesis Kit	Agilent	Cat# 200521	
commercial assay or kit	AAVpro® Purification Kit	Takara Bio.	Cat# 6675	
commercial assay or kit	Triglyceride Colorimetric Assay Kit	Cayman Chemical	Cat# 10010303	
commercial assay or kit	Alanine Transaminase Colorimetric Activity Assay Kit	Cayman Chemical	Cat# 700260	
commercial assay or kit	Aspartate Aminotransferase Colorimetric Activity Assay Kit	Cayman Chemical	Cat# 701640	
commercial assay or kit	Mouse Insulin ELISA Kit	Crystal Chem	Cat# 90080	
commercial assay or kit	PicoSens™ Free Fatty Acid Quantification Kit	BioMax	Cat# BM-FFA-100	
commercial assay or kit	B-hydroxybutyrate assay Kit	abcam	Cat# ab83390	

Chemical Compound, drug	Oleic acid	Sigma-Aldrich	Cat# O1008	
Chemical Compound, drug	Fenofibrate	Santa Cruz biotechnology	Cat# sc-204751	- HepG2 cells were treated with fenofibrate (100 µM) - mice injected with fenofibrate (100 mg/kg)

Cell Culture

Human liver cells, HepG2 and Huh-7 cells were purchased from American Type Culture Collection (ATCC, Manassas, VA) and cultured in Dulbecco's Modified Eagle's Medium (DMEM) containing 10 % fetal bovine serum (Gibco, BRL, Grand Island, NY) and 1 % penicillin/streptomycin (ThermoFisher Scientific, Waltham, MA). miRNAs and siRNA were obtained from GenePharma (Shanghai, China). The miRNAs and siRNA used in this study are listed in the table below. HepG2 and Huh-7 cells were transfected with miRNA or siRNA using Lipofectamine™ RNAiMAX transfection reagent (ThermoFisher Scientific) according to the manufacturer's instructions and following experiments were performed 48 h after transfection. For intracellular lipid accumulation, cells were cultured in a medium with the addition of 1 mmol/L sodium oleic acid for 24 hours and then cells were harvested for further analysis. Images of cells stained with Oil Red O were obtained with EVOS FL (Thermo Fisher Scientific). For quantification, Oil Red O was extracted with isopropanol and absorbance was measured at 520 nm (Lim et al., 2012).

Name	Sequence
MIR NC	UCACAACCUCCUAGAAAGAGUAGA
MIR20B/Mir20b (Mimic)	CAAAGUGCUCAUAGUGCAGGUAG
anti-MIR20B/Mir20b (Inhibitor)	CUACCUGCACUAUGAGCACUUUG
siPPARA	CGGCGAGGATAGTTCTGGAAGCTTT

Human patients

Human liver tissue samples of 21 patients were acquired from the BioResource Center (BRC) of Asan Medical Center, Seoul, Republic of Korea. The process of 21 human tissue samples was

officially approved by the Institutional Review Board of Asan Medical Center (IRB approval number:

2018-1512). Human samples used in this study were from hepatocellular carcinomas (HCC) liver

resection. All samples were taken from non-tumor parts of the resected tissue and selected as 11

normal liver samples, 4 simple steatosis, and 6 NASH samples through histological analysis of

pathologists using the grading and staging system for NASH (Brunt, Janney, Di Bisceglie,

Neuschwander-Tetri, & Bacon, 1999). Although limitations in completely excluding the effect of HCC,

histologically normal tissue adjacent to the tumor has obvious distinctions in global transcriptome

patterns for the tumor and can be used as a control (Aran et al., 2017). There were no significant

differences in age and male/female ratio among the three groups. Specifically, serum AST, ALT, and

fasting glucose levels were significantly increased with progression from normal to NASH, but total

cholesterol was comparable as previously reported (Chung et al., 2020). In addition, all patients

diagnosed with alcoholic liver disease, viral infected hepatitis and toxic hepatitis were excluded.

Library preparation for transcriptome sequencing

RNA-seq was performed on triplicate sample from HepG2 cell with or without overexpression of

MIR20B. Total RNA was isolated using the RNeasy mini kit (Qiagen, Hilden, Germany) according to

the manufacturer's instructions. Library prep and RNA-seq were performed by Novogene (Hong

Kong). A total amount of 1 μg RNA per sample was used as input material for the RNA sample

preparations. Sequencing libraries were generated using NEBNext® UltraTM RNA Library Prep Kit for

Illumina® (NEB, Ipswich, MA) following manufacturer's recommendations and index codes were

added to attribute sequences to each sample. PCR products were purified (AMPure XP system,

Beckman Coulter Life Sciences, Indianapolis, IN) and library quality was assessed on the Agilent

Bioanalyzer 2100 system (Agilent Technologies, Inc., Santa Clara, CA). The clustering of the index-

coded samples was performed on a cBot Cluster Generation System using PE Cluster Kit cBot-HS

(Illumina, San Diego, CA) according to the manufacturer's instructions. After cluster generation, the

library preparations were sequenced on an Illumina platform (NovaSeq 6000 PE150) and paired-end

reads were generated.

miRNA regulatory network analysis of nuclear receptor genes

The RNA-seq fold change data for 16,010 genes were obtained from the Supplementary file 1 of a

previous work (Hoang et al., 2019). The miRNA target genes that shared evolutionarily conserved

binding sites for 353 miRNAs (≥ 10 genes) were downloaded from TargetScan database (Ver.7.2)

(Agarwal, Bell, Nam, & Bartel, 2015). We used the 50 genes that belonged to the nuclear receptor

transcription pathway in REACTOME database (Jassal et al., 2020) (Liberzon et al., 2015) to analyze

how miRNAs regulate the transcription of NR genes. Among them 17 genes were significantly

downregulated in NASH patients (adjusted *p*-value < 0.1). The enrichment of these downregulated

NR genes in the targets of the 353 miRNAs were assessed using hypergeometric distribution. The *p*-

value of a miRNA indicates the probability of having downregulated nuclear receptors among the

miRNA targets. The *p*-value is given as follows:

$$p - \text{value (miRNA)} = \sum_{i=O}^{M \wedge D} \frac{\binom{D}{i}\binom{N-D}{M-i}}{\binom{N}{M}}$$

where *N* is the total number of genes analyzed, *M* is the number of candidate target genes of the

miRNA, *D* is the downregulated NR genes, and *O* is the observed overlap between miRNA targets

and the downregulated NR genes.

The miRNAs whose target genes were enriched in the downregulated NR genes with adjusted *p*-

value $\leq$ 0.05 were used to construct the regulatory networks of the NR transcription pathway (Figure

1A).

Differential expression and gene-set enrichment analysis (GSEA)

The differential expression analysis of RNA-seq data were performed using limma package(Ritchie

et al., 2015) where moderated *t*-test was applied for voom-transformed read counts. The resulting

fold-change values between the test and control conditions were used for the pathway analysis. The

preranked GSEA (R package)(Subramanian et al., 2005) was used for the pathway analysis of gene

sets from WikiPathway(Martens et al., 2021), REACTOME(Jassal et al., 2020), KEGG(Kanehisa,

Furumichi, Tanabe, Sato, & Morishima, 2017) databases (MSigDB) and the enrichment score plots.

Immunoblotting

Supernatants containing protein contents were determined by Bradford Protein Assay (Bio-Rad

Laboratories, Hercules, CA). Proteins were immunoblotted with anti-PPARα (ab24509, Abcam,

Cambridge, MA) and anti-HSP90 (4877S, Cell Signaling Technology, Danvers, MA).

Quantitative PCR

Total mRNAs were isolated using TRIzol reagent purchased from Thermo Fisher Scientific.

Reverse-transcription of the RNA was performed with ABI Reverse Transcription Kit (Thermo Fisher

Scientific). Quantitative PCR was performed using 7900HT Fast Real-Time PCR System (Life

Technologies, Carlsbad, CA) following the manufacturer's instructions. Relative mRNA expression

levels of each gene were normalized to TATA-binding protein TBP. The mtDNA copy number was

evaluated based on the ratio of mtDNA to nuclear DNA by quantitative PCR. The mtDNA was

quantified based on the mitochondrial gene, *VIPR1*, and *MT-ATP6*, respectively. The relative amounts

of mtDNA were normalized to nuclear DNA, *B2M*. The primer pairs used in this study are listed in the

table below.

Human Gene	Sequences of the primers for qPCR	
	Forward primer	Reverse primer
MIR20B	GCAAAGTGCTCATAGTGCAGGTAG	TCGCACTTGTCATACACCAG
MIR129	CTTTTTGCGGTCTGGGCTTGC	AGTGCAGGGTCCGAGGTATT
MIR219A	TGATTGTCCAAACGCAATTC	GAACATGTCTGCGTATCTC
MIR4782	TTCTGGATATGAAGACAATCAA	CAAGATCTATACTTCTGTTAGT
MIR6766	CGGGTGGGAGCAGATCTTAT	GAACATGTCTGCGTATCTC
MIR20A	TGCGCTAAAGTGCTTATAGTGC	CCAGTGCAGGGTCCGAGGTATT
MIR106A	GATGCTCAAAAAGTGCTTACAGTGCA	TATGGTTGTTCTGCTCTCTGTCTC
MIR106B	CAAGTACCCACAGTGCGGT	CTCGCTTCGGCAGCACA
MIR17	TCTAGATCCCGAGGACTG	ATCGTGACCTGAACC
MIR519D	AGTGCCTCCCTTTAGAG	GAACATGTCTGCGTATCTC
U6	CTCGCTTCGGCAGCACA	AACGCTTCACGAATTTGCGT
PPARA	GCTATCATTACGGAGTCCACG	TCGCACTTGTCATACACCAG
CPT1A	AGGCGACATCAATCCGAAC	AAAGGCTACGAATGGGAAGG
ACOX1	CCACGTATGACCCTGAAACC	TCCATAGCATTTCCCCTTAGTG
SREBF1	CAACACAGCAACCAGAAACTC	CTCCACCTCAGTCTTCACG
FASN	CAAGCTGAAGGACCTGTCTAG	CGGAGTGAATCTGGGTTGATG
CD36	GCCAGGTATTGCAGTTCTTTTC	TGTCTGGGTTTTCAACTGGAG
FABP1	GCAGAGCCAGGAAAACTTTG	AGCGGTGATGGTGAACTTG
PPARGC1A	ACCAAACCCACAGAGAACAG	GGGTCAGAGGAAGAGATAAAGTTG
SIRT1	CCCTCAAAGTAAGACCAGTAGC	CACAGTCTCCAAGAAGCTCTAC
VIPR1	CTCCACCATTAGCACCCAAAGCTAAG	GATATTGATTTCACGGAGGATGGTGGTC
MT-ATP6	AACGAAAATCTGTTCGCTTCAT	ATGTGTTGTCGTGCAGGTAGAG
HMGCL	GAGTTTTCAGAGGTTTGACGC	CAAGAGCACAGGAGACGTAC
ACAT1	CGGGCTAACTGATGTCTACAA	CAAATTTCCCAGCTTCCCATG

Gene	Forward primer	Reverse primer
ACAT2	CCCAGAACAGGACAGAGAATG	AGCTTGGACATGGCTTCTATG
RORA	GGTGATGCTTTTGTTCTTACTGG	TGTCTCCACAGATCTTGCATG
RORC	TGGTGCTGGTTAGGATGTG	GGAGTGGGAGAAGTCAAAGATG
THRB	CATCAAAACTGTCACCGAAGC	TCCAAGTCAACCTTTCCACC
NRBP1	GTTCCACCCAGCATTGTTTG	CAGGGATTTCAGCCAGTACG
NAGK	TATTTCCAGGTGCCAGATCG	CTGAAGATATAGCGGGAAAGGG
USP46	ATACACCAAGCTGTCTTACCG	ATATAATGCCCACGATTAGGACC
ITGB8	CGTCTCATCTCGCTCTTGATAG	TTCTCTGAAAGTTGGCCTAGTG
BMPR2	GGCTGACTGGAAATAGACTGG	CACAGTCCCTCAAGTTCACAG
ZNFX1	CCGAGGATTGTCATAGTGGAAG	AGATCATACACGTTGGCACTG
EPHA5	AGATTGAGGCAGTGAATGGAG	GCCAAGACAAAGAGATGCTG
E2F1	TCTCCGAGGACACTGACAG	ATCACCATAACCATCTGCTCTG
AGPAT1	AGGACGCAACGTCGAGAAC	GCAGTACCTCCATCATCCCAAG
AGPAT2	GCCGAGTTCTACGCCAAGG	CGAACCAGCCGATGATGCT
GPAT1	GATGTAAGCACACAAGTGAGGA	TCCGACTCATTAGGCTTTCTTTC
GPAT2	TGTGGTCGTCAGGCTTTGG	GGTCCGTTATGCTTCTGTGGA
DGAT1	TATTGCGGCCAATGTCTTTGC	CACTGGAGTGATAGACTCAACCA
DGAT2	ATTGCTGGCTCATCGCTGT	GGGAAAGTAGTCTCGAAAGTAGC
FATP2	TTTCCGCCATCTACACAGTCC	CGTAGGTGAGAGTCTCGTCG
FATP3	CCCTGCTGGAATTAGCGATTT	GGGCGAGGTAGATCACATCTT
FATP4	CTCAGGTCTTGGAGAAGGAAC	ACAGCGGGTCTTTCACAATAG
FATP5	TGGAGGAGATCCTTCCCAAGC	TGGTCCCCGAGGTATAGATGAA
FATP6	CTTCTGTCATGGCTAACAGTTCT	AGGTTTCCGAGGTTGTCTTTTG
CPT1B	ATCCTACTCCTATGACCCCG	TCTGCATTGAGACCCAACTG
CPT2	TTGAGTGCTCCAAGTACCATG	GCAAACAAGTGTCGGTCAAAG
ACADM	TCATTGTGGAAGCAGATACCC	CAGCTCCGTCACCAATTAAAAC
FATP1	ATGAGGACACAATGGAGCTG	TGACATAGCCATCGAAGCG
ACAA1	GCGGTTCTCAAGGACGTGAAT	GTCTCCGGGATGTCACTCAGA
ACSL1	CCATGAGCTGTTCCGGTATTT	CCGAAGCCCATAAGCGTGTT
TBP	CCACTCACAGACTCTCACAAC	CTGCGGTACAATCCCAGAACT
B2M	TGCTGTCTCCATGTTTGATGTATCT	TCTCTGCTCCCCACCTCTAAGT

	Sequences of the primers for qPCR	
Mouse Gene	**Forward primer**	**Reverse primer**
Mir20b	GCAAAGTGCTCATAGTGCAGGTAG	TCGCACTTGTCATACACCAG
Mir129	CTTTTTGCGGTCTGGGCTTGC	AGTGCAGGGTCCGAGGTATT
Mir219a	TGATTGTCCAAACGCAATTC	GAACATGTCTGCGTATCTC
Mir20a	TGCGCTAAAGTGCTTATAGTGC	CCAGTGCAGGGTCCGAGGTATT
Mir106a	GATGCTCAAAAAGTGCTTACAGTGCA	TATGGTTGTTCTGCTCTCTGTCTC
Mir106b	CAAGTACCCACAGTGCGGT	CTCGCTTCGGCAGCACA
Mir17	TCTAGATCCCGAGGACTG	ATCGTGACCTGAACC
Mir519d	AGTGCCTCCCTTTAGAG	GAACATGTCTGCGTATCTC
U6	CTCGCTTCGGCAGCACA	AACGCTTCACGAATTTGCGT
Ppara	TCAGGGTACCACTACGGAGT	CTTGGCATTCTTCCAAAGCG
Cpt1a	AGTTCCATGACCCATCTCTGTC	TTCTTCTTCCAGAGTGCAGC
Acox1	TAACTTCCTCACTCGAAGCCA	AGTTCCATGACCCATCTCTGTC
Srebf1	GGAGCCATGGATTGCACATT	CTTCCAGAGAGGAGGCCAG
Fasn	GGAGGTGGTGATAGCCGGTAT	TGGGTAATCCATAGAGCCCAG
Cd36	GCGACATGATTAATGGCACAG	GATCCGAACACAGCGTAGATAG
Fabp1	TCTCCGGCAAGTACCAATTG	TTGATGTCCTTCCCTTTCTGG
Tnf	CCCTCACACTCAGATCATCTTCT	GCTACGACGTGGGCTACAG
Ccl2	TTAAAAACCTGGATCGGAACCAA	GTTCACCGTAAGCCCAATTT
Il6	TAGTCCTTCCTACCCCAATTTCC	TTGGTCCTTAGCCACTCCTTC

Il1b	GCACTACAGGCTCCGAGATGAAC	TTGTCGTTGCTTGGTTCTCCTTGT
Acta2	GTGAAGAGGAAGACAGCACAG	GCCCATTCCAACCATTACTCC
Col1a1	CATAAAGGGTCATCGTGGCT	TTGAGTCCGTCTTTGCCAG
Col3a1	GAAGTCTCTGAAGCTGATGGG	TTGCCTTGCGTGTTTGATATTC
Fn	CTTTGGCAGTGGTCATTTCAG	ATTCTCCCTTTCCATTCCCG
Timp1	CTCAAAGACCTATAGTGCTGGC	CAAAGTGACGGCTCTGGTAG
Vim	CGTCCACACGCACCTACAG	GGGGGATGAGGAATAGAGGCT
E2f1	TGCAGAACAGATGGTCATAGTG	GGGCACAGGAAAACATCAATG
Rora	GTGGAGACAAATCGTCAGGAAT	TGGTCCGATCAATCAAACAGTTC
Rorc	TCCACTACGGGGTTATCACCT	AGTAGGCCACATTACACTGCT
Thrb	GGACAAGCACCCATCGTGAAT	CTCTGGTAATTGCTGGTGTGAT
Cpt1b	GACTTCCGGCTTAGTCGGG	GAATAAGGCGTTTCTTCCAGGA
Cpt2	CAGCACAGCATCGTACCCA	TCCCAATGCCGTTCTCAAAAT
Acadm	AACACAACACTCGAAAGCGG	TTCTGCTGTTCCGTCAACTCA
Fatp1	CTG GGA CTT CCG TGG ACC T	TCT TGC AGA CGA TAC GCA GAA
Acaa1	TCT CCA GGA CGT GAG GCT AAA	CGC TCA GAA ATT GGG CGA TG
Acsl1	TGC CAG AGC TGA TTG ACA TTC	GGC ATA CCA GAA GGT GGT GAG
Atgl	ATGTTCCCGAGGGAGACCAA	GAGGCTCCGTAGATGTGAGTG
Hsl	GATTTACGCACGATGACACAGT	ACCTGCAAAGACATTAGACAGC
Mgl	ACCATGCTGTGATGCTCTCTG	CAAACGCCTCGGGGATAACC
Tbp	ACCCTTCACCAATGACTCCTATG	TGACTGCAGCAAATCGCTTGG

Cellular Oxygen Consumption Rate (OCR)

OCR of HepG2 cells were analyzed by Seahorse XF24 extracellular flux analyzer (Seahorse

Bioscience, North Billerica, MA) following the manufacturer's instruction. The results were normalized

with the protein quantity of each corresponding well.

Measurement of FA β-oxidation and uptake

FA β-oxidation were measured by the conversion of [9,10-^{3}H(N)]-Palmitic Acid (PerkinElmer,

Waltham, MA) to ^{3}H$_2$O. Cells were incubated with 1.25 mCi/L [9,10-^{3}H(N)]-Palmitic Acid with cold

palmitic acid in a final concentration of 200μM for 4 hours. After incubation, medium were recovered

and precipitated with an equal volume of 10 % tricholoroacetic acid. The supernatant was transferred

to new- and capless-microtube and the capless-tube was inserted into D.W-added Scintillation tube

and incubated at 60 °C for 12 hours. The capless tube was removed from scintillation tube and

measured the CPMA with scincillation counter oil using Tri-carb 2910TR liquid scintillation counter

(PerkinElmer).

For FA uptake measurement, cells were incubated with 0.5 μCi/L [9,10-^{3}H(N)]-Palmitic Acid with

cold palmitic acid in a final concentration of 200μM for 2 hours. Uptake was stopped by addition of

200 μM phloretin in 0.1 % BSA and lysed in 0.1 N NaOH / 0.03 % SDS buffer. The radioactivity of

each lysate was counted using Tri-carb 2910TR liquid scintillation counter. The β-oxidation and

uptake was normalized to lysate protein concentration determined by BSA assay.

Metabolites assay

HepG2 cells were transfected with MIR20B, anti-MIR20B, or MIR NC, respectively. After 48 h,

HepG2 cells (20,000 cells per well) were seeded in 96-well MitoPlate S-1 plates and examined in

mitochondrial metabolites activity following the manufacturer's instructions (Biolog, Hayward, CA).

Mice

All animal experiments were performed according to procedures approved by the Ulsan National

Institute of Science and Technology's Institutional Animal Care and Use Committee (UNISTIACUC-

19-04). Mice were maintained in a specific pathogen–free animal facility under a 12-h light/dark cycle

at a temperature of 21°C and allowed free access to water and food. Seven-week-old male C57BL/6J

mice (DBL, Chungbuk, Republic of Korea) were fed a HFD (60% kcal fat, D12492, Research Diets

Inc., New Brunswick, NJ) for 12 weeks or a MCD (A02082002BR, Research Diets Inc., New

Brunswick, NJ, USA) for 4 weeks. Fenofibrate (100 mg/kg, sc-204751, Santa Cruz biotechnology,

Dallas, TX) was administered orally for 4 weeks before mice were sacrificed.

Hepatocyte isolation

Briefly, mice were anesthetized with isoflurane, and 24-gauge needle was inserted into the portal

vein. Then the inferior vena cava was cut, and the mouse liver was perfused sequentially with solution

I (142 µM NaCl, 6.7 µM KCl, 10 µM HEPES, and 2.5 mM EGTA), and solution II (66.7 mM NaCl, 6.7

mM KCl, 50mM HEPES, 4.8 mM $CaCl_2\cdot2H_2O$, and 0.01 % Type IV collagenase (Sigma- Aldrich, St.

Louis, MO)). After digestion, the liver was disrupted over a 70-µm cell strainer, and cell suspension

was spun at 50 x g for 5min at 4 °C. The supernatant was gently aspirated and the cells were

resuspended in M199 with EBSS (M199/EBSS) medium and gently mixed with equal volume of

Percoll working solution (48.6 % Percoll). The cell suspension was spun at 100 x g for 5 min at 4 °C,

and the pellet washed once with M199/EBSS. After viable cells were counted with trypan blue, the

isolated hepatocytes were seed in M199/EBSS medium supplemented with 10 % FBS, 1 %

penicillin/streptomycin, and 10 nM dexamethasone.

Production of AAV

The *Mir20b* and anti-*Mir20b* were cloned into the pOTTC385-pAAV CMV-IE IRES EGFP vector (Addgene plasmid # 102936)(Nelson et al., 2019) and co-transfected with pAAV-DJ vector and pAAV-Helper vector into HEK 293T cells to generate recombinant adeno-associated virus expressing *Mir20b* or anti-*Mir20b* according to the manufacturer's protocol (Cell Biolabs, San Diego, CA). The AAVs were purified with AAVpro® Purification Kit (Takara Bio, Shiga, Japan). When feeding HFD was started, purified AAV-*Mir20b*, AAV-anti-*Mir20b*, or AAV-Control (1 X 10^{10} PFU) was injected into mice *via* tail-vein. AAV-Control or AAV-anti-*Mir20b* was injected into mice *via* tail-vein before the initiation of MCD diet.

Lentivirus preparation, transduction, and stable cell line generation

All lentivirus-based shRNA clones used for making the viral transduction particles were purchased from Sigma-Aldrich. pLKO.1-Puro vector targeting human *PPARA* or non-target vector as a control was used. In short, 293T cells were plated at a density of 12×10^6 293T cells in a 150-mm dish containing 20 mL of media. After 24 hours, the cells were overlaid with a complex containing a three-plasmid system (pLKO.1 shRNA, pCMV-VSVG, and pCMV-Δ8.9) at the ratio of 4:2:3 using lipofectamine (Jang et al., 2020). The supernatant was collected starting from 48 to 72 hours post-transfection. The virus particles were concentrated by ultracentrifugation at 25,000 rpm for 1 hour 30 minutes with a Beckman ultracentrifuge using the SW28 rotor, resuspended in PBS, and stored until use at –70 °C. Harvested virus was used to transduce target cells for 24 hours. Stable cell lines were selected in the presence of 3 μg/mL puromycin for 5 days. To minimize batch-to-batch differences, we compared only cells made from the same batch of cells.

Name	Sequence
shPPARA #1	GAACAGAAACAAATGCCAGTA
shPPARA #2	GTAGCGTATGGAAATGGGTTT

Metabolic analysis

Mice were fasted overnight (18 h) before intraperitoneal injection of D-glucose (2 g/kg body weight) for glucose tolerance test. For insulin tolerance test, mice were fasted for 4 hours before intraperitoneal injection of insulin (0.75 U/kg body weight). Every glucose was examined with tail-vein

blood at indicated intervals after injection using a glucometer. For analyzing metabolic parameters,

insulin (90080, Crystal Chem, Elk Grove Village, IL), ALT (K752, Biovision Inc., Milpitas, CA), AST

(K753, Biovision Inc.), cholesterol (K603, BioVision Inc.), TG (10010303, Cayman Chemical, Ann

Arbor, MI), and FFA (BM-FFA-100, BioMax, Republic of Korea) were determined. Body composition

of mice was measured using an EchoMRI100V, quantitative nuclear resonance system (Echo Medical

Systems, Houston, TX).

274 Histological analysis

Liver tissues were isolated from mice and immediately fixed with 4% formalin (Sigma- Aldrich, St.

Louis, MO). Histological changes of lipid droplets were examined by H&E staining and Oil Red O

staining. As counterstain, Mayer's hematoxylin was used for every slide. Liver fibrosis was further

examined by Sirius red with liver section. Images were obtained with Olympus BX53 microscope and

DP26 camera. NAFLD activity score (NAS) and fibrosis score were measured by a pathologist

according to the NASH Clinical Research Network scoring system (Brunt et al., 2011). For statistical

analysis of fibrosis score, each grade (0~4) was calculated by giving a score from 1 to 6 points.

283 Statistical analysis

284   All data are represented as mean ± SEM. Statistically significant differences were assessed by the

285 Student's t-test. Statistical analyses and Pearson correlation analyses were performed using

286 Microsoft Excel or GraphPad Prism 9.3.0 (RRID:SCR_002798). All of the significance are expressed

287 as $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, $^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.001$, $^{\$}P < 0.05$, $^{\$\$}P < 0.01$, and

288 $^{\$\$\$}P < 0.001$. ns, not significant.

290 Data Availability

Except for RNA-Seq data, all data generated and analyzed during the current study are included in

the manuscript and supporting files. The RNA-Seq data have been deposited at the Gene Expression

Omnibus (GEO) website: https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE168484

(accession number GSE168484). Source data files are provided for Figure 1, Figure 1-figure

supplement 3, Figure 1-figure supplement 4, Figure 2, Figure 2-figure supplement 2, Figure 2-figure

supplement 3, Figure 2-figure supplement 4, Figure 2-figure supplement 5, Figure 3, Figure 4, Figure

4-figure supplement 1, Figure 4-figure supplement 2, Figure 4-figure supplement 3, Figure 4-figure

supplement 4, Figure 4-figure supplement 5, Figure 4-figure supplement 6, Figure 5, Figure 5-figure

supplement 1, Figure 5-figure supplement 2, Figure 5-figure supplement 3, Figure 5-figure

supplement 4, Figure 6, Figure 6-figure supplement 1, Figure 6-figure supplement 2, Figure 6-figure

supplement 3, Figure 6-figure supplement 4, Figure 7, Figure 8, and Figure 9 and Figure 10.

Results

MIR20B significantly increases in the livers of dietary obese mice and human.

We constructed a regulatory network of NRs that were differentially expressed in NAFLD

patients(Hoang et al., 2019) and microRNA targeting NRs based on miRNA target prediction(Agarwal

et al., 2015), to identify the correlation between NR and microRNA in the development of NAFLD. As

shown in Figure 1A, the top ten miRNAs were found to be highly correlated with the modulation of NR

expression in NAFLD. To further prioritize key miRNAs, we assessed the expression of miRNA

candidates in NASH, simple steatosis, and normal patient liver samples (Figure 1B, Figure 1-figure

supplement 1). Among the selected miRNAs, the expression of *MIR20B* was significantly increased in

simple steatosis and NASH compared to that in normal individuals, and the extent of increase in

NASH was higher than that in simple steatosis. In particular, the expression of *MIR20B* showed a

significant correlation with the BMI, AST, ALT, and fasting glucose during NAFLD progression (Figure

1-figure supplement 2).

MIR20B expression was also most significantly increased in both oleic acid (OA)-treated HepG2

and Huh-7 cells (Figure 1C, D, Figure 1-figure supplement 4A and B). Moreover, the expression of

MIR20B was significantly upregulated in the fatty livers of high-fat diet (HFD)-fed mice, *Lep*$^{ob/ob}$ mice,

and methionine-deficient diet (MCD)-fed mice compared to that in the liver of normal chow diet (NCD)-

fed wild mice (Figure 1E-G, Figure 1-figure supplement 3, and Figure 1-figure supplement 4C).

Together, our results indicate that *MIR20B* expression is increased in NAFLD and is highly associated

with the regulation of NRs in NAFLD.

PPARA **is a direct target of MIR20B.**

Next, we characterized the physiological roles of MIR20B in NAFLD. Oil Red O staining showed

that *MIR20B* expression increased intracellular lipid content, and this lipid accumulation was

increased with OA treatment in HepG2 cells (Figure 2A). As expected, overexpression of *MIR20B*

significantly upregulated TG content and cholesterol in OA-treated HepG2 cells (Figure 2B, C). To

further investigate the functions and targets of MIR20B, HepG2 cells were transfected with MIR20B

and two separate RNA-Seq libraries of non-targeting control (MIR NC) and forced *MIR20B* expression

(MIR20B) were constructed. The six RNA-seq samples were clearly separated between the forced

MIR20B expression and control conditions, implying a significant impact on gene expression in

HepG2 cells (Figure 2D, Figure 2-figure supplement 1A). The NR metapathway was detected as the

most significant pathways (adjusted *p*-value = 2.47-E8) by the gene-set enrichment analysis (GSEA)

(Subramanian et al., 2005), indicating that NRs and related genes are the major targets of MIR20B

(Figure 2E, F). The heatmap of RNA-seq data for the NR transcription pathway is shown in Figure 2G.

Five NRs (*PPARA*, *RORA*, *RORC*, *THRB*, and *NRBP1*) were downregulated (adjusted *p*-value $\leq$

0.05), and *PPARA* was the most significantly downregulated NR (adjusted *p*-value = 3.25-E5).

Furthermore, GSEA showed that PPARA pathways and PPAR signaling pathways were significantly

decreased in *MIR20B* overexpressed cells (Figure 2-figure supplement 1B, C). We validated the

expression of these NRs and the *PPARA*, *RORC*, *THRB*, and *NRBP1* expression was decreased by

MIR20B in human liver cells and mouse primary hepatocytes. Consistent with the RNA-seq data, the

expression change of *PPARA* at both the protein and mRNA levels with MIR20B transfection was the

most distinct compared to the control (Figure 2H, I). Moreover, among candidate targets from RNA-

seq analysis, only *PPARA* was selected as an overlapped predicted target of MIR20B between

various miRNA target prediction programs, including miRDB, picTAR, TargetSCAN, and miRmap

(Figure 2J, Figure 2-figure supplement 2). While the expression of other candidates, including *RORA*,

RORC, and *THRB*, remained unchanged, that of *PPARA* significantly decreased as NAFLD

progressed to steatosis and NASH, and showed a negative correlation with that of *MIR20B* (Figure 2-

figure supplement 3). In addition, *PPARA* decreased in diverse diets-induced NAFLD conditions with

MIR20B (Figure 1C-E, Figure 2-figure supplement 4)

Notably, the 3'UTR of *PPARA* mRNA includes MIR20B binding sites that are well conserved

between humans and mice, suggesting that MIR20B may have a direct inhibitory effect on *PPARA*

expression (Figure 2K). Using a luciferase reporter construct including the 3'UTR of *PPARA*, we

revealed that MIR20B suppressed the luciferase activity in both HepG2 and Huh-7 cells in a dose-

dependent manner. Furthermore, we built the mutant construct of the predicted MIR20B binding sites

within the 3'UTR of *PPARA*; the inhibitory effect of MIR20B on luciferase activity was completely

blunted (Figure 2L, M). To further validate the functional role of 3'UTR regulated by MIR20B, we

introduced the wild type *PPARA* open reading frame (ORF) followed by either the wild type (WT) or

mutant (Mut) 3'UTR of *PPARA* in HepG2 cells (Figure 2-figure supplement 5A). As shown in Figure 2-

figure supplement 5C, MIR20B significantly suppressed the expression of *PPARA* and its target

genes in *PPARA*-3'UTR WT expressing cells. Furthermore, Oil Red O staining showed that MIR20B

increased the intracellular lipid content in these cells (Figure 2-figure supplement 5B). However,

MIR20B did not have an effect on either the expression of *PPARA* and its target genes or the

intracellular lipid content in *PPARA*-3'UTR Mut expressing cells (Figure 2-figure supplement 5B-D).

Taken together, these results indicate that MIR20B inhibits the expression of *PPARA* by interacting

with its 3'UTR.

The effects of MIR20B are mediated by PPARα.

Next, we confirmed the effect of MIR20B on function of PPARα. Since PPARα is a master regulator

of lipid metabolism such as FA utilization and oxidation, we investigated the effects of MIR20B on

expression of genes involved in lipid metabolism. Transfection of MIR20B into HepG2 cells reduced

the expression and activity of PPARα, but co-transfected *PPARA* expression vector restored them

(Figure 3A, B). Overexpression of *MIR20B* reduced the expression of genes involved in FA β-

oxidation and FA uptake, including *CPT1A, ACOX1, CD36,* and *FABP1,* but not *de novo* lipogenesis

such as *SREBF1* and *FASN*, which was significantly restored by the forced expression of PPARα

(Figure 3C). Next, we tested whether anti-MIR20B enhanced activity of PPARα and the effects of anti-

MIR20B was specific to PPARα. Ectopic expression of anti-*MIR20B* increased the expression and

activity of PPARα (Figure 3D, E), and consequently enhanced the expression of genes involved in FA

utilization and oxidation (Figure 3F). *PPARA*-targeting siRNA (*siPPARA*) suppressed the expression

of *PPARA* and its activity (Figure 3D, E). The increased expression of genes by anti-MIR20B was also

suppressed by *siPPARA* (Figure 3F). In addition, fenofibrate, a PPARα agonist, increased the

expression of PPARα and its transcriptional activity in HepG2 cells transfected with MIR20B, but

could not restore as much on its own effects (Figure 3G, H). Interestingly, fenofibrate treatment

increased the expression of genes involved in FA β-oxidation and FA uptake which are regulated by

PPARα, but could not overcome the inhibitory effect of MIR20B (Figure 3I). Taken together, these

results indicate that MIR20B affects lipid metabolism through direct inhibition of *PPARA*.

MIR20B regulates fatty acid metabolism.

We investigated the effects of MIR20B on lipid metabolism to reveal the functional contribution of

increased MIR20B to NAFLD. Unlike in HepG2 cells (Figure 2A-C), MIR20B alone did not induce lipid

accumulation in primary hepatocytes without OA treatment, but MIR20B significantly increased lipid

accumulation in the presence of OA (Figure 4-figure supplement 2A-D). The mRNA levels of genes

involved in FA β-oxidation, FA uptake, including *CPT1A, ACOX1, CD36,* and *FABP1*, and PPARα

target genes were decreased by overexpression of *MIR20B* compared to the control in HepG2 cells

and primary hepatocytes (Figure 4A, B, Figure 4-figure supplement 1A), whereas the genes

associated with *de novo* lipogenesis and ketogenesis were not affected by MIR20B (Figure 4B, Figure

4-figure supplement 2E, 3A and B). Interestingly, *DGAT1*, rate limiting enzyme for TG synthesis, was

increased in HepG2 cells upon OA treatment (Figure 4C). *MIR20B* overexpressed HepG2 cells

showed reduced levels of palmitoyl-carnitine, a substrate of β-oxidation, and acetyl-CoA, a product of

β-oxidation. Subsequently, TCA cycle intermediate levels, including citrate and succinate, also

decreased (Figure 4D).

Enforced expression of *MIR20B* in HepG2 cells under both basal and OA treatments decreased the

expression of *PPARGC1A* and *SIRT1*, which are involved in mitochondrial biogenesis (Figure 4E).

The copy number of two mtDNA genes, *VIPR1* and *MT-ATP6*, was decreased by *MIR20B*

overexpression following OA treatment (Figure 4F). Consistently, mitochondrial function that was

analyzed *via* OCR (oxygen consumption rate) was reduced by MIR20B under both basal and OA

treatment conditions compared to the control (Figure 4G). In particular, the basal respiration and

maximal respiratory capacity were significantly suppressed by MIR20B (Figure 4H). Furthermore, the

level of ATP production, FA uptake, and FA oxidation was reduced in *MIR20B* overexpressed cells

compared with that in the control under both basal and OA-treated conditions (Figure 4I-K).

To further clarify the role of MIR20B in hepatic fatty acid metabolism, anti-MIR20B was delivered into

OA-treated HepG2 cells and primary hepatocytes, and which reduced the expression of *MIR20B*

(Figure 4L, Figure 4-figure supplement 4G). Oil red O staining showed that anti-MIR20B remarkably

decreased intracellular lipid accumulation upon OA treatment (Figure 4-figure supplement 4A, D). As expected, MIR20B inhibition reduced the levels of both TG and cholesterol under OA conditions compared to the control (Figure 4-figure supplement 4B, C, E, and F). Lipid consumption-associated genes and PPARα target genes, not *de novo* lipogenic genes, were significantly upregulated in MIR20B inhibited HepG2 cells and primary hepatocytes compared to those in the control under both basal and OA conditions (Figure 4M, Figure 4-figure supplement 1B, 4H and I). TG synthesis enzymes, such *GPAT1* and *AGPAT1*, were decreased by anti-MIR20B in HepG2 cells under OA-treated condition, however *DGAT1* and *DGAT2* were reduced under both basal and OA-treated conditions (Figure 4N). Inhibition of MIR20B increased the levels of palmitoyl-carnitine, acetyl-CoA, and TCA cycle intermediates (Figure 4O), whereas ketogenesis was not affected (Figure 4-figure supplement 3C, D). Furthermore, the expression of mitochondrial biogenesis genes (Figure 4P) and the copy number of mitochondrial DNA genes were increased in both basal and OA conditions (Figure 4Q). Consequently, anti-MIR20B treatment significantly upregulated the mitochondrial activity, FA uptake, and FA oxidation (Figure 4R-V). Taken together, these results demonstrated that MIR20B contributes to hepatic lipid accumulation by controlling lipid oxidation, mitochondrial function and TG synthesis through changes in gene expression, further contributing to the progression of NAFLD.

MiR20b promotes hepatic steatosis in HFD-fed mice

To confirm the *in vivo* roles of MiR20b in obesity model mice, we introduced MiR20b using an adenovirus-associated vector (AAV), referred to as AAV-*Mir20b*, into C57BL/6 mice that had been fed a normal chow diet (NCD) or a high-fat diet (HFD). Administration of AAV- *Mir20b* led to high expression levels of *Mir20b* in the livers of NCD- and HFD-fed mice compared to AAV-Control injection (Figure 5A). However, the expression level of *Mir20b* was not changed in adipose tissues including white and brown adipose tissues compared to NCD- or HFD-fed AAV-Control injected mice, respectively (Figure 5-figure supplement 1). Consequently, AAV-*Mir20b* injected mice exhibited a reduction in the expression of *Ppara* and PPARα target genes compared with AAV-Control injected mice on both NCD and HFD (Figure 5B, R, and Figure 5-figure supplement 2).

Alterations in body weight were not detected in NCD-fed mice after AAV-*Mir20b* administration; however, AAV-*Mir20b* led to a significant increase in the body weight of HFD-induced obese mice

(Figure 5C). The ratio of fat mass to body weight in AAV-*Mir20b* administration HFD-fed mice was

higher than that in AAV-Control treated mice (Figure 5D and Figure 5-figure supplement 3); however,

the ratio of lean mass to body weight was decreased (Figure 5E). Consistently, AAV-*Mir20b*

administration increased liver weight, steatosis, and NAFLD activity score (NAS) in HFD-fed mice

(Figure 5F-H). The hepatic TG level, serum activities of aspartate aminotransferase (AST) and alanine

aminotransferase (ALT), markers of liver injury, were significantly increased with AAV-*Mir20b*

administration compared with AAV-Control administration in HFD-fed mice (Figure 5I-K).

Additionally, we observed that delivery of AAV-*Mir20b* to HFD-fed mice significantly impaired

glucose tolerance and insulin sensitivity compared to the AAV-Control (Figure 5L, M). Fasting glucose,

insulin, and homeostasis model assessment of insulin resistance (HOMA-IR) levels were also

increased in AAV-*Mir20b* administrated HFD-fed mice (Figure 5N-P). We observed that genes

involved in FA β-oxidation and FA uptake pathways were downregulated by AAV-*Mir20b* compared to

AAV-Control in both NCD- and HFD-fed mice, whereas lipogenesis genes were not altered in AAV-

Mir20b administrated mice (Figure 5Q). These results suggest that MiR20b could aggravate NAFLD

by dysregulating lipid metabolism in a HFD-induced obesity model.

Inhibition of MiR20b alleviates hepatic steatosis in HFD-fed mice.

Next, we introduced anti-*Mir20b* into HFD-fed mice. Administration of AAV-anti-*Mir20b* led to

decrease of *Mir20b* in the livers of NCD- and HFD-fed mice compared to AAV-Control injection

(Figure 6A and Figure 6-figure supplement 1). AAV-anti-*Mir20b* significantly increased expression of

Ppara and its target genes in the livers of both NCD- and HFD-fed mice (Figure 6B, R and Figure 6-

figure supplement 2). Administration of AAV-anti-*Mir20b* in HFD-fed mice reduced the body weight

compared to that of AAV-Control administrated mice (Figure 6C). We further determined that

alterations in body weight were associated with fat mass loss and reduced trend of epididymal white

adipose tissue (Figure 6D and Figure 6-figure supplement 3). While the ratio of lean mass to body

weight of AAV-anti-*Mir20b* administrated HFD-fed mice was increased, the lean mass was

comparable to that of the control (Figure 6E). We next observed that AAV-anti-*Mir20b* administration

reduced liver weight and hepatic steatosis in HFD-fed mice than in AAV-Control mice (Figure 6F).

H&E and Oil Red O staining demonstrated that delivery of AAV-anti-*Mir20b* significantly attenuated

the size and number of lipid droplets in the liver and NAS compared to AAV-Control administration in

HFD-fed mice (Figure 6G, H). In accordance with histological changes, metabolic parameters were

reduced in AAV-anti-*Mir20b* administrated mice compared with the AAV-Control administrated mice

(Figure 6I-K). Furthermore, AAV-anti-*Mir20b* significantly improved glucose tolerance (Figure 6L) and

insulin sensitivity (Figure 6M) compared to the AAV-Control in HFD-fed mice. Consistently, we

determined that fasting glucose, fasting insulin, and HOMA-IR levels were decreased by AAV-anti-

Mir20b (Figure 6N-P). Delivery of AAV-anti-*Mir20b* increased the expression of genes associated with

FA β-oxidation and FA uptake compared with the administration of AAV-Control (Figure 6Q). Together,

these results suggest that suppression of MiR20b could ameliorate NAFLD by recovering lipid

metabolism in a HFD-induced obesity model.

The effects of fenofibrate are limited in *Mir20b*-introduced mice.

Next, we tested whether NAFLD treatment with fenofibrate was affected by *Mir20b* expression *in*

vivo. Administration of AAV-*Mir20b* led to elevated hepatic levels of *Mir20b* compared to AAV-Control

injection in HFD-fed mice, and the level was slightly decreased by fenofibrate treatment (Figure 7A).

Interestingly, we observed that administration of AAV-Control with fenofibrate increased the level of

Ppara; however, fenofibrate could not restore the reduced *Ppara* expression by AAV-*Mir20b* (Figure

7B). Administration of fenofibrate reduced the body and liver weights of AAV-Control injected mice;

however, AAV-*Mir20b* injected mice exhibited no significant differences by fenofibrate (Figure 7C, D).

The ratio of fat to body weight also displayed no alterations between AAV-*Mir20b* and AAV-*Mir20b*

with fenofibrate (Figure 7E). While the ratio of lean mass to body weight was increased by fenofibrate

in AAV-*Mir20b* injected mice, the lean mass was comparable (Figure 7F). H&E staining, Oil Red O

staining, NAS and hepatic TG levels demonstrated that fenofibrate significantly attenuated lipid

accumulation in the liver of HFD-fed mice, but the effect of fenofibrate was suppressed by AAV-

Mir20b (Figure 7G-I). Serum AST and ALT levels were decreased by fenofibrate, but this benefit was

did not detected in AAV-*Mir20b* injected mice (Figure 7J, K). We further observed that blood glucose

tolerance and insulin sensitivity were improved by fenofibrate; however, AAV-*Mir20b* offset the

improvement by fenofibrate (Figure 7L, M). Fasting glucose, fasting insulin, and HOMA-IR levels were

markedly decreased by fenofibrate in HFD-fed mice (Figure 7N-P). In AAV-*Mir20b* injected mice,

fenofibrate did not reduce fasting insulin levels, but decreased fasting glucose and HOMA-IR levels.

Fenofibrate also did not restore the suppressed expression of genes regulating FA β-oxidation in

AAV-*Mir20b*-injected mice (Figure 7Q). Taken together, the effect of fenofibrate to ameliorate NAFLD-

like symptoms was limited in AAV-*Mir20b* administrated HFD-fed mice because of the targeting

PPARα by MiR20b.

MiR20b promotes liver inflammation and fibrosis in MCD-Fed Mice.

We further demonstrated that RNA-seq data have a significant correlation of fold change values

with previously published RNA-seq data under both NASH and liver fibrosis conditions within the NR

transcription(Hoang et al., 2019) (Figure 8A, B). This implies that MIR20B is able to set up an NR

transcription program similar to that of NASH and liver fibrosis. To test this hypothesis, AAV-Control or

AAV-anti-*Mir20b* was administered to C57BL/6 mice placed on a methionine/choline-deficient diet

(MCD), which is the most widely used diet to induce NAFLD/NASH. Administration of AAV-anti-*Mir20b*

led to decrease of *Mir20b* in the livers of NCD- and MCD-fed mice compared to AAV-Control injection

(Figure 8C). We observed that the expression of *Ppara* was increased in MCD-fed mice and

administration of AAV-anti-*Mir20b* displayed an elevation of PPARα, both at the mRNA and protein

levels (Figure 8D-F). We next observed that AAV-anti-*Mir20b* administration significantly reduced

hepatic steatosis in MCD-fed mice than in AAV-Control mice (Figure 8G). Liver sections clearly

showed a decrease in both steatosis and fibrosis with AAV-anti-*Mir20b* administration in MCD-fed

mice (Figure 8H-J). Consistently, AAV-anti-*Mir20b* administration decreased the levels of hepatic TG,

AST, and ALT activity compared to AAV-Control injection (Figure 8K-M). Moreover, AAV-anti-*Mir20b*

significantly reduced the expression of genes related to hepatic inflammation, including *Tnf* and *Il6*

(Figure 8N), and fibrosis, including the NASH-relevant genes, such as *Acta2, Col1a1, Col3a1* and

Timp1 (Figure 8O), in MCD-fed mice. Taken together, these results indicate that MiR20b plays an

important role in the development of fibrosis, inflammation, and hepatic steatosis in NAFLD

progression.

Anti-MiR20b enhances efficacy of fenofibrate in MCD-fed mice.

Recently, drug development strategies for NAFLD/NASH are moving toward combination therapies

(Dufour, Caussy, & Loomba, 2020). However, the efficacy of developing drugs, including fenofibrate,

against NAFLD/NASH is limited (Fernandez-Miranda et al., 2008). Thus, we tested whether the

combination of anti-MiR20b and fenofibrate would improve NAFLD in MCD-fed mice. The levels of

hepatic *Mir20b* were reduced after administration of AAV-anti-*Mir20b* in MCD-fed mice compared to

those in mice administered with AAV-Control, and this reduction was also observed after fenofibrate

treatment (Figure 9A). Interestingly, the combination of AAV-anti-*Mir20b* and fenofibrate increased the

levels of *Ppara* to a greater extent than AAV-anti-*Mir20b* alone (Figure 9B, C). AAV-anti-*Mir20b* or

fenofibrate administration significantly reduced the liver weight and hepatic TG levels, and co-

administration further reduced hepatic steatosis (Figure 9D, E). Histological sections showed that the

combination of AAV-anti-*Mir20b* and fenofibrate improved NAFLD, as evidenced by the effects on

both lipid accumulation and fibrosis in the liver (Figure 9F-H). Consistently, the levels of AST and ALT

were significantly lower after combined treatment with AAV-anti-*Mir20b* and fenofibrate than after a

single treatment (Figure 9I, J). In addition, the expression of genes related to hepatic inflammation,

such as *Tnf* and *Il6* (Figure 9K), and fibrosis, such as *Acta2*, *Col1a1*, *Fn*, and *Timp1*, (Figure 9L), was

further decreased by the combination of AAV-anti-*Mir20b* and fenofibrate. These results suggest that

AAV-anti-*Mir20b* may increase the efficacy of fenofibrate, especially its effect on fibrosis, and provide

a more effective option for improving NAFLD/NASH.

Discussion

Obesity has been widely demonstrated to be central to the pathogenesis of NAFLD. Among other

peripheral tissues, the liver plays a dominant role in the regulation of lipid homeostasis(Pawlak,

Lefebvre, & Staels, 2015). Although abnormal regulation of metabolic homeostasis in the liver has

been recognized in diabetes and NAFLD, the underlying molecular mechanisms remain to be

elucidated. Growing evidence has demonstrated that MIR20B levels are significantly upregulated in

the plasma miRNA profiles of NAFLD patients(Jin et al., 2012). Moreover, plasma MIR20B levels

were highly elevated in T2DM/NAFLD patients compared to those in T2DM patients(Ye et al., 2018).

However, the molecular mechanism through which MIR20B regulates NAFLD progression remains

unknown. In this study, we demonstrated that MIR20B promotes NAFLD progression by modulating

lipid metabolism, including FA β-oxidation, FA uptake, and TG synthesis, as well as ATP production

by mitochondrial biogenesis. Our data clearly showed the regulatory mechanism of PPARα by

MIR20B, and MIR20B may serve as a novel biological marker in NAFLD.

Decreased PPARα is contributed to development of NAFLD(Francque et al., 2015). A few miRNAs

were reported to regulate *PPARA* expression in NAFLD. miR-34a targets *PPARA* and *SIRT1*,

associating with FA oxidation and cholesterol synthesis. However the effect of MIR34A on

inflammation and fibrosis is not clear(Ding et al., 2015). MIR21 also decreases the expression of

PPARA in NASH, however, activated PPARα by MIR21 suppression reduces inflammation, liver injury,

and fibrosis without improvement in FA β-oxidation and lipid accumulation(Loyer et al., 2016). In the

present study, we demonstrate the novel miRNA which has different mode of action. MIR20B showed

the deteriorating effects on FA oxidation, steatosis, inflammation, and fibrosis in HFD- or MCD-fed

mice. How these miRNAs targeting *PPARA* have different regulatory mechanisms should be further

studied.

A previous study demonstrated that upregulated MIR20B levels in obesity-induced metabolic

disorders such as T2DM were considered to prevent several targets, such as *STAT3*, *CD36*, and

PTEN, which are involved in glucose and lipid homeostasis. The MIR20B/STAT axis, which is

involved in the insulin signaling pathway, alters glycogen synthesis in human skeletal muscle

cells(Katayama et al., 2019). Moreover, MIR20B directly targets *PTEN* involved in PI3K/Akt/mTOR

signaling pathway that modulates glucose metabolism in gastric cells(Streleckiene et al., 2020). Hosui

et al. (2017) proposed a model in which the fatty acid transporter *CD36* is also a potential target of

MIR20B, which crucially regulates hepatic lipid metabolism in *STAT5* KO mice models. But, it has

been reported that *CD36* and *FABP1* are direct PPARα target genes(Rakhshandehroo, Knoch, Muller,

& Kersten, 2010). In this study, we observed that MIR20B downregulated *PPARA* and suppressed the

expression of *CD36* and *FABP1*. Thus, *PPARA* is the primary target of MIR20B in regulating hepatic

lipid metabolism.

MIR20B was induced by OA, but FA uptake was decreased by overexpression of *MIR20B*, and was

accompanied by a considerable decrease in *CD36* expression (Figure 4B, J). However, other lipid

transporters such as FATPs, except for PPARα target *FATP1*, were not significantly altered (Figure 4-

figure supplement 1, 5), suggesting that FA uptake is continued by these transporters. The expression

of CD36 is relatively low in normal hepatocytes, and the molecule may not be the primary fatty acid

transporter in these cells(Wilson et al., 2016). Furthermore, the decrease in FA uptake upon *CD36*

KO is modest even during a HFD(Wilson et al., 2016). In addition, we observed that the expression of

MIR20B is induced and increased for up to 24 hours by OA treatment (Figure 4-figure supplement 6).

This is followed by a slight decrease, remaining at a constant elevated level. Together, the findings

indicated that other fatty acid transporters contributing to FA uptake account for the entry of OA into

cells and MIR20B induction.

Hepatic steatosis could be affected by adipose tissue through FFA release and hepatic uptake of

circulating FFAs(Rasineni et al., 2021). Our results showed that the epididymal adipose tissue of

HFD-fed mice was enlarged upon AAV-*mir20b* treatment; however, the serum FFA levels in these

mice were comparable to those in mice treated with the AAV-Control (Figure 5-figure supplement 4B).

Of note, the expression of genes related to lipolysis did not change in adipose tissues, and that of

hepatic FA transporter, CD36, was decreased by AAV-*mir20b* treatment (Figure 5Q and Figure 5-

figure supplement 4A). In addition, excess hepatic TGs are secreted as very low density lipoproteins

(VLDLs), and the secretion rate increases with the TG level(Fabbrini et al., 2008). VLDLs deliver TGs

from the liver to adipose tissue and contributes expansion of adipose tissue(Chiba, Nakazawa, Yui,

Kaneko, & Shimokado, 2003). Together, these reports suggest that adipose tissue is also remodeled

by the liver in HFD-fed mice and patients with NAFLD. Therefore, the increase of hepatic TGs by

AAV-*mir20b* is unlikely affected by AAV-*mir20b* on epididymal adipose tissue, and the increase in fat

content (Figure 5-figure supplement 3) may be a consequence of increased hepatic TG levels.

The expression of glycerolipid synthetic genes, including *AGPATs*, *GPATs*, and *DGATs*, was

increased by OA treatment, but *MIR20B* overexpression did not influence the expression of these

genes except for that of *DGAT1*. However, treatment with anti-MIR20B significantly reduced the

expression of glycerolipid synthetic genes, including *GPATs* and *DGATs*, under OA treatment (Figure

4C, N). These results suggested that MIR20B is necessary but not sufficient to induce the expression

of glycerolipid synthetic genes under OA treatment. We have shown that OA induces the expression

of *MIR20B* (Figure 1C, D), which can explain why *MIR20B* overexpression did not show an additional

enhancement under OA treatment. The increase in *DGAT1* expression induced by MIR20B might

contribute to the increase in TG formation and capacity to store OA. This could change the flux of

oleyl-CoA to TG synthesis, not β-oxidation with reduced expression of lipid oxidation-associated

genes (Figure 4B). Thus, we can expect that the decrease in OA uptake and increase in TG formation

induced by MIR20B resulted in reduced amounts of OA or oleyl-CoA inside the cell. However, as lipid

consumption through FA oxidation is decreased by MIR20B, free OA or oleyl-CoA might be

maintained at a stably increased level compared to that of OA-untreated MIR NC or MIR20B condition,

and the impact of the changes in OA or oleyl-CoA levels on the transcriptional phenotype might not be

significant as found in a constant elevated level of MIR20B by OA.

Recent reports suggest that some transcription factors regulate metabolic homeostasis by directly

mediating the expression of miRNAs(Yang & Wang, 2011). E2F1, which is a member of the E2F

transcription factor family, regulates myoblast differentiation and proliferation *via* the auto-regulatory

feedback loop between E2F1 and MiR20b in muscle(Luo, Li, Yi, Nie, & Zhang, 2016). Both the

hepatic expression and activity of E2F1 are increased during obesity. E2F1 deficiency protects

against obesity- and diabetes-induced liver steatosis in mouse models(Zhang et al., 2014).

Additionally, E2F1 induced chronic inflammation and hepatic lipid metabolism during NAFLD

development(Denechaud et al., 2016). Consistent with these results, we observed that the expression

of E2F1 was significantly increased in the fatty liver of both mice and humans, and its expression was

positively correlated with that of MIR20B (Figure 10). These results suggest that E2F1 may be an

upstream regulator of MIR20B in the liver, and this upregulation of MIR20B regulates lipid metabolism

in the pathogenesis of NAFLD.

NAFLD patients can develop NASH, which is characterized by hepatic steatosis complicated by

chronic hepatocellular damage and severe inflammation with fibrosis, potentially developing into

cirrhosis and HCC(Corey & Kaplan, 2014). This indicates that suppression of NAFLD progression is

the primary option for preventing the development of HCC. Recent reports suggest that *Ppara* KO

mice fed an MCD developed much more severe NASH than wild-type mice, and the expression of

Ppara in HCC tissue was significantly lower than that in normal liver tissue(Montagner et al., 2016).

PPARα activation contributes to the inhibition of HCC cell proliferation Thus, hepatic PPARα plays a

crucial role in tumorigenesis in the liver(Lefebvre, Chinetti, Fruchart, & Staels, 2006). Interestingly, it

has been reported that upregulated MiR20b highly regulates cancer cell proliferation and promotes

proliferation of H22 hepatocellular carcinoma cells (Peng et al., 2019; Xia et al., 2020). Thus, plasma

MIR20B can be a promising target in liver cancer development. Indeed, we observed that the level of

MIR20B was increased in NAFLD patients, but even robustly increased in the NASH stage.

Furthermore, we observed that the hepatic function of MiR20b dramatically regulates the genes

involved in inflammation and fibrosis by directly repressing *Ppara* in MCD-fed mice. Thus, our study

strongly suggested that MIR20B regulates the pathogenesis of NAFLD, but might also be relevant in

the development of severe stages of liver fibrosis and even in HCC.

Fibrates are FDA-approved medications for the treatment of patients with dyslipidemia

(hypercholesterolemia, mixed dyslipidemia, hypertriglyceridemia)(Katsiki et al., 2013). Fibrates are

well-known PPARα agonists, and PPARα is a widely accepted promising target for the treatment of

NAFLD and NASH. In animal studies, fenofibrate improved hypertriglyceridemia and hepatic steatosis.

In a small pilot trail, fenofibrate improved ALT and AST levels and hepatocellular ballooning

degeneration, but did not show significant improvements in steatosis, lobular inflammation, fibrosis, or

NAS score(Fernandez-Miranda et al., 2008). Clofibrate, another PPARα agonist, did not show clinical

benefit in the treatment of NASH(Laurin et al., 1996). Recently, the effect of omega-3 poly-

unsaturated FAs (n-3 PUFA), which are ligands for PPARα, on patients with NAFLD was evaluated.

Omacor, which consists of purified docosahexaenoic and eicosapentaenoic acids as ethyl esters,

showed a linear decrease in liver fat percentage without improvements in fibrosis score(Scorletti et al.,

2014). Omacor has undergone a phase 3 clinical trial but the results have not been disclosed

(NCT01277237). These results show that the effect of PPARα agonists on NAFLD is limited. Our

results suggest one of the possible reasons for this limited effect. Here, we show that PPARα

expression is suppressed by increased MIR20B expression, which may weaken the efficacy of

fibrates. Indeed, our results showed that the effects of fenofibrate were limited in AAV-*Mir20b*-treated

mice, and that the combination of fenofibrate and AAV-anti-*Mir20b* further improved the NAFLD

phenotype in MCD-fed mice. Nowadays, it is generally accepted that treatment with PPARα agonists

alone is limited and that a combination strategy is necessary for NAFLD. The inhibition of MIR20B

presents an opportunity for the development of novel therapeutics for NAFLD.

Acknowledgments

This research was funded by Korea Mouse Phenotyping Project (2016M3A9D5A01952411), the

National Research Foundation of Korea (NRF) grant funded by the Korea government

(2020R1F1A1061267, 2018R1A5A1024340, NRF-2021R1I1A2041463), the Future-leading Project

Research Fund (1.210034.01) of UNIST to J.H.C. and the National Research Foundation of Korea

(NRF) grant funded by the Korea government (2020R1I1A1A01074940) to H-J.J.

Competing Interest Statement

No conflicts of interest relevant to this manuscript.

References

Agarwal, V., Bell, G. W., Nam, J. W., & Bartel, D. P. (2015). Predicting effective microRNA target sites
in mammalian mRNAs. *Elife, 4*. doi:10.7554/eLife.05005

Aran, D., Camarda, R., Odegaard, J., Paik, H., Oskotsky, B., Krings, G., . . . Butte, A. J. (2017).
Comprehensive analysis of normal adjacent to tumor transcriptomes. *Nat Commun, 8*(1),
1077. doi:10.1038/s41467-017-01027-z

Bartel, D. P. (2004). MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell, 116*(2), 281-
297. doi:10.1016/s0092-8674(04)00045-5

Brunt, E. M., Janney, C. G., Di Bisceglie, A. M., Neuschwander-Tetri, B. A., & Bacon, B. R. (1999).
Nonalcoholic steatohepatitis: a proposal for grading and staging the histological lesions. *Am J
Gastroenterol, 94*(9), 2467-2474. doi:10.1111/j.1572-0241.1999.01377.x

Brunt, E. M., Kleiner, D. E., Wilson, L. A., Belt, P., Neuschwander-Tetri, B. A., & Network, N. C. R.
(2011). Nonalcoholic fatty liver disease (NAFLD) activity score and the histopathologic
diagnosis in NAFLD: distinct clinicopathologic meanings. *Hepatology, 53*(3), 810-820.
doi:10.1002/hep.24127

Chakravarthy, M. V., Pan, Z., Zhu, Y., Tordjman, K., Schneider, J. G., Coleman, T., . . . Semenkovich,
C. F. (2005). "New" hepatic fat activates PPARalpha to maintain glucose, lipid, and
cholesterol homeostasis. *Cell Metab, 1*(5), 309-322. doi:10.1016/j.cmet.2005.04.002

Corey, K. E., & Kaplan, L. M. (2014). Obesity and liver disease: the epidemic of the twenty-first
century. *Clin Liver Dis, 18*(1), 1-18. doi:10.1016/j.cld.2013.09.019

de Alwis, N. M., & Day, C. P. (2008). Non-alcoholic fatty liver disease: the mist gradually clears. *J
Hepatol, 48 Suppl 1*, S104-112. doi:10.1016/j.jhep.2008.01.009

Denechaud, P. D., Lopez-Mejia, I. C., Giralt, A., Lai, Q., Blanchet, E., Delacuisine, B., . . . Fajas, L.
(2016). E2F1 mediates sustained lipogenesis and contributes to hepatic steatosis. *J Clin
Invest, 126*(1), 137-150. doi:10.1172/JCI81542

Ding, J., Li, M., Wan, X., Jin, X., Chen, S., Yu, C., & Li, Y. (2015). Effect of miR-34a in regulating
steatosis by targeting PPARalpha expression in nonalcoholic fatty liver disease. *Sci Rep, 5*,
13729. doi:10.1038/srep13729

Dufour, J. F., Caussy, C., & Loomba, R. (2020). Combination therapy for non-alcoholic
steatohepatitis: rationale, opportunities and challenges. *Gut, 69*(10), 1877-1884.
doi:10.1136/gutjnl-2019-319104

Fabbrini, E., Sullivan, S., & Klein, S. (2010). Obesity and nonalcoholic fatty liver disease: biochemical,
metabolic, and clinical implications. *Hepatology, 51*(2), 679-689. doi:10.1002/hep.23280

Fernandez-Miranda, C., Perez-Carreras, M., Colina, F., Lopez-Alonso, G., Vargas, C., & Solis-
Herruzo, J. A. (2008). A pilot trial of fenofibrate for the treatment of non-alcoholic fatty liver
disease. *Dig Liver Dis, 40*(3), 200-205. doi:10.1016/j.dld.2007.10.002

Francque, S., Verrijken, A., Caron, S., Prawitt, J., Paumelle, R., Derudas, B., . . . Staels, B. (2015).
PPARalpha gene expression correlates with severity and histological treatment response in
patients with non-alcoholic steatohepatitis. *J Hepatol, 63*(1), 164-173.
doi:10.1016/j.jhep.2015.02.019

Hoang, S. A., Oseini, A., Feaver, R. E., Cole, B. K., Asgharpour, A., Vincent, R., . . . Sanyal, A. J.
(2019). Gene Expression Predicts Histological Severity and Reveals Distinct Molecular
Profiles of Nonalcoholic Fatty Liver Disease. *Sci Rep, 9*(1), 12541. doi:10.1038/s41598-019-
48746-5

Hosui, A., Tatsumi, T., Hikita, H., Saito, Y., Hiramatsu, N., Tsujii, M., . . . Takehara, T. (2017). Signal
transducer and activator of transcription 5 plays a crucial role in hepatic lipid metabolism
through regulation of CD36 expression. *Hepatol Res, 47*(8), 813-825. doi:10.1111/hepr.12816

Jang, H. J., Lim, S., Kim, J. M., Yoon, S., Lee, C. Y., Hwang, H. J., . . . Choi, J. H. (2020).
Glucosylceramide synthase regulates adipo-osteogenic differentiation through synergistic
activation of PPARgamma with GlcCer. *FASEB J, 34*(1), 1270-1287.
doi:10.1096/fj.201901437R

Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., . . . D'Eustachio, P. (2020).
The reactome pathway knowledgebase. *Nucleic Acids Res, 48*(D1), D498-d503.
doi:10.1093/nar/gkz1031

Jin, X., Chen, Y. P., Kong, M., Zheng, L., Yang, Y. D., & Li, Y. M. (2012). Transition from hepatic
steatosis to steatohepatitis: unique microRNA patterns and potential downstream functions
and pathways. *J Gastroenterol Hepatol, 27*(2), 331-340. doi:10.1111/j.1440-
1746.2011.06864.x

Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y., & Morishima, K. (2017). KEGG: new perspectives
on genomes, pathways, diseases and drugs. *Nucleic Acids Res, 45*(D1), D353-D361.
doi:10.1093/nar/gkw1092

Katayama, M., Wiklander, O. P. B., Fritz, T., Caidahl, K., El-Andaloussi, S., Zierath, J. R., & Krook, A.
(2019). Circulating Exosomal miR-20b-5p Is Elevated in Type 2 Diabetes and Could Impair
Insulin Action in Human Skeletal Muscle. *Diabetes, 68*(3), 515-526. doi:10.2337/db18-0470

Katsiki, N., Nikolic, D., Montalto, G., Banach, M., Mikhailidis, D. P., & Rizzo, M. (2013). The role of
fibrate treatment in dyslipidemia: an overview. *Curr Pharm Des, 19*(17), 3124-3131.
doi:10.2174/13816128113199170020

Laurin, J., Lindor, K. D., Crippin, J. S., Gossard, A., Gores, G. J., Ludwig, J., . . . McGill, D. B. (1996).
Ursodeoxycholic acid or clofibrate in the treatment of non-alcohol-induced steatohepatitis: a
pilot study. *Hepatology, 23*(6), 1464-1467. doi:10.1002/hep.510230624

Lefebvre, P., Chinetti, G., Fruchart, J. C., & Staels, B. (2006). Sorting out the roles of PPAR alpha in
energy metabolism and vascular homeostasis. *J Clin Invest, 116*(3), 571-580.
doi:10.1172/JCI27989

Liberzon, A., Birger, C., Thorvaldsdottir, H., Ghandi, M., Mesirov, J. P., & Tamayo, P. (2015). The
Molecular Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst, 1*(6), 417-
425. doi:10.1016/j.cels.2015.12.004

Lim, S., Jang, H. J., Park, E. H., Kim, J. K., Kim, J. M., Kim, E. K., . . . Suh, P. G. (2012).
Wedelolactone inhibits adipogenesis through the ERK pathway in human adipose tissue-
derived mesenchymal stem cells. *J Cell Biochem, 113*(11), 3436-3445. doi:10.1002/jcb.24220

Lopez-Sanchez, G. N., Dominguez-Perez, M., Uribe, M., Chavez-Tapia, N. C., & Nuno-Lambarri, N.
(2021). Non-alcoholic fatty liver disease and microRNAs expression, how it affects the
development and progression of the disease. *Ann Hepatol, 21*, 100212.
doi:10.1016/j.aohep.2020.04.012

Lopez-Velazquez, J. A., Carrillo-Cordova, L. D., Chavez-Tapia, N. C., Uribe, M., & Mendez-Sanchez,
N. (2012). Nuclear receptors in nonalcoholic Fatty liver disease. *J Lipids, 2012*, 139875.
doi:10.1155/2012/139875

Loyer, X., Paradis, V., Henique, C., Vion, A. C., Colnot, N., Guerin, C. L., . . . Rautou, P. E. (2016).
Liver microRNA-21 is overexpressed in non-alcoholic steatohepatitis and contributes to the
disease in experimental models by inhibiting PPARalpha expression. *Gut, 65*(11), 1882-1894.
doi:10.1136/gutjnl-2014-308883

Luo, W., Li, G., Yi, Z., Nie, Q., & Zhang, X. (2016). E2F1-miR-20a-5p/20b-5p auto-regulatory
feedback loop involved in myoblast proliferation and differentiation. *Sci Rep, 6*, 27904.
doi:10.1038/srep27904

Martens, M., Ammar, A., Riutta, A., Waagmeester, A., Slenter, D. N., Hanspers, K., . . . Kutmon, M.
(2021). WikiPathways: connecting communities. *Nucleic Acids Res, 49*(D1), D613-D621.
doi:10.1093/nar/gkaa1024

McKenna, N. J., Cooney, A. J., DeMayo, F. J., Downes, M., Glass, C. K., Lanz, R. B., . . . O'Malley, B.
W. (2009). Minireview: Evolution of NURSA, the Nuclear Receptor Signaling Atlas. *Mol
Endocrinol, 23*(6), 740-746. doi:10.1210/me.2009-0135

Montagner, A., Polizzi, A., Fouche, E., Ducheix, S., Lippi, Y., Lasserre, F., . . . Guillou, H. (2016). Liver
PPARalpha is crucial for whole-body fatty acid homeostasis and is protective against NAFLD.
Gut, 65(7), 1202-1214. doi:10.1136/gutjnl-2015-310798

Nelson, B. S., Fulenwider, H. D., Nennig, S. E., Smith, B. M., Sequeira, M. K., Chimberoff, S. H., . . .
Schank, J. R. (2019). Escalated Alcohol Self-Administration and Sensitivity to Yohimbine-
Induced Reinstatement in Alcohol Preferring Rats: Potential Role of Neurokinin-1 Receptors
in the Amygdala. *Neuroscience, 413*, 77-85. doi:10.1016/j.neuroscience.2019.06.023

Pawlak, M., Lefebvre, P., & Staels, B. (2015). Molecular mechanism of PPARalpha action and its
impact on lipid metabolism, inflammation and fibrosis in non-alcoholic fatty liver disease. *J
Hepatol, 62*(3), 720-733. doi:10.1016/j.jhep.2014.10.039

792 Peng, L., Li, S., Li, Y., Wan, M., Fang, X., Zhao, Y., . . . Xuan, Y. (2019). Regulation of BTG3 by
793    microRNA-20b-5p in non-small cell lung cancer. *Oncol Lett, 18*(1), 137-144.
794    doi:10.3892/ol.2019.10333

795 Rakhshandehroo, M., Knoch, B., Muller, M., & Kersten, S. (2010). Peroxisome proliferator-activated
796    receptor alpha target genes. *PPAR Res, 2010*. doi:10.1155/2010/612089

797 Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). limma powers
798    differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
799    *Res, 43*(7), e47. doi:10.1093/nar/gkv007

800 Scorletti, E., Bhatia, L., McCormick, K. G., Clough, G. F., Nash, K., Hodson, L., . . . Study, W. (2014).
801    Effects of purified eicosapentaenoic and docosahexaenoic acids in nonalcoholic fatty liver
802    disease: results from the Welcome* study. *Hepatology, 60*(4), 1211-1221.
803    doi:10.1002/hep.27289

804 Streleckiene, G., Inciuraite, R., Juzenas, S., Salteniene, V., Steponaitiene, R., Gyvyte, U., . . .
805    Skieceviciene, J. (2020). miR-20b and miR-451a Are Involved in Gastric Carcinogenesis
806    through the PI3K/AKT/mTOR Signaling Pathway: Data from Gastric Cancer Patients, Cell
807    Lines and Ins-Gas Mouse Model. *Int J Mol Sci, 21*(3). doi:10.3390/ijms21030877

808 Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., . . . Mesirov,
809    J. P. (2005). Gene set enrichment analysis: a knowledge-based approach for interpreting
810    genome-wide expression profiles. *Proc Natl Acad Sci U S A, 102*(43), 15545-15550.
811    doi:10.1073/pnas.0506580102

812 Wilson, C. G., Tran, J. L., Erion, D. M., Vera, N. B., Febbraio, M., & Weiss, E. J. (2016). Hepatocyte-
813    Specific Disruption of CD36 Attenuates Fatty Liver and Improves Insulin Sensitivity in HFD-
814    Fed Mice. *Endocrinology, 157*(2), 570-585. doi:10.1210/en.2015-1866

815 Xia, L., Li, F., Qiu, J., Feng, Z., Xu, Z., Chen, Z., & Sun, J. (2020). Oncogenic miR-20b-5p contributes
816    to malignant behaviors of breast cancer stem cells by bidirectionally regulating CCND1 and
817    E2F1. *BMC Cancer, 20*(1), 949. doi:10.1186/s12885-020-07395-y

818 Yang, Z., & Wang, L. (2011). Regulation of microRNA expression and function by nuclear receptor
819    signaling. *Cell Biosci, 1*(1), 31. doi:10.1186/2045-3701-1-31

820 Ye, D., Zhang, T., Lou, G., Xu, W., Dong, F., Chen, G., & Liu, Y. (2018). Plasma miR-17, miR-20a,
821    miR-20b and miR-122 as potential biomarkers for diagnosis of NAFLD in type 2 diabetes
822    mellitus patients. *Life Sci, 208*, 201-207. doi:10.1016/j.lfs.2018.07.029

823 Zhang, Y., Xu, N., Xu, J., Kong, B., Copple, B., Guo, G. L., & Wang, L. (2014). E2F1 is a novel
824    fibrogenic gene that regulates cholestatic liver fibrosis through the Egr-1/SHP/EID1 network.
825    *Hepatology, 60*(3), 919-930. doi:10.1002/hep.27121

Figures legend

Figure 1. ***MIR20B*** **expression is significantly increased in the livers of dietary obese mice and**

humans.

The miRNA regulatory networks for NR genes downregulated in the transcriptome of NAFLD

patients. The adjusted p-values in the table represent the enrichment of miRNA targets in the

downregulated NR genes (hypergeometric distribution) (A). The expression of miRNAs was measured

in indicated condition by quantitative RT-PCR (B-G). Hepatic miRNA levels of patients with steatosis

(n = 4) or NASH (n = 6) were normalized to those of normal individuals (n = 11). $*P < 0.05$, $**P < 0.01$

and $***P < 0.001$ *vs* normal individuals. $^{\#}P < 0.05$ and $^{\#\#}P < 0.01$ *vs* patients with steatosis (B).

MIR20B levels from HepG2 cells (C) and Huh-7 cells (D) treated with OA for 24 hours were

normalized to no treatment (NT). Hepatic *Mir20b* levels from C57BL/6J mice fed high fat diet (HFD)

(E), or methionine-deficient diet (MCD) (F) were normalized to normal chow diet (NCD). Hepatic

Mir20b levels from leptin-deficient *Lep$^{ob/ob}$* mice were normalized to lean wild mice (G). Values

represent means ± SEM (n = 3 - 5). $*P < 0.05$, $**P < 0.01$ *vs* NT in cells or NCD-fed mice.

Figure 1-figure supplement 1. Clinical characteristics of control individuals (N=11), steatosis

(steatosis > 50%, N=4) and NASH patients (N=6).

Individual clinical characteristics (A) and summary statistics (B) of patients. Data were presented as

mean ± SEM. $*P < 0.05$, $**P < 0.01$ *vs* Normal. BMI, Body Mass Index; AST, aspartate

aminotransferase; ALT, alanine aminotransferase; GGT, γ-glutamyl transpeptidase; ALP, alkaline

phosphatase; INR, international normalized ratio.

Figure 1-figure supplement 2. Pearson correlation between clinical characteristics of patients

and ***MIR20*B** **expression.**

Pearson correlation between metabolic parameters and *MIR20B* mRNA levels in individual clinical

patients. Correlation analysis of BMI (kg/m^2) (A), AST (IU/L) (B), ALT (IU/L) (C), and fasting glucose

(mg/dL) (D) with *MIR20B* expression ($2^{-\Delta Ct}$) in all population. BMI, Body Mass Index; AST, aspartate

aminotransferase; ALT, alanine aminotransferase.

Figure 1-figure supplement 3. Hepatic gene expression associated with steatosis,

inflammation and fibrosis is increased in dietary or genetic obese mice.

Genes involved in steatosis, inflammation, and fibrosis were measured by quantitative RT-PCR in

C57BL/6J mice fed high fat diet (HFD) (A) or methionine-deficient diet (MCD) (B), and were

normalized to normal chow diet (NCD). Leptin-deficient *Lep$^{ob/ob}$* mice were normalized to lean wild

mice (C). Values represent means ± SEM (n = 3 - 5). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ *vs* NCD-fed

mice.

Figure 1-figure supplement 4. *MIR20B* expression significantly increases in OA-treated

hepatocytes and the livers of dietary obese mice.

The expression of miRNA candidates from regulatory network analysis was measured by

quantitative RT-PCR in OA-treated HepG2 cells (A) or Huh-7 cells (B) for 24 hours, and liver of

C57BL/6J mice fed a high fat diet (HFD) (C). It was normalized to miRNA expression in no treatment

(NT) or normal chow diet (NCD). Values represent means ± SEM (n = 3-5). ns, not significant.

*$P < 0.05$, **$P < 0.01$ *vs* NT in cells or NCD-fed mice.

Figure 2. *PPARA* is a direct target of MIR20B.

Overexpressed *MIR20B* induces hepatic lipid accumulation in HepG2 cells treated with OA (1 mM).

Oil Red O staining showed intracellular lipid accumulation in HepG2 cells. Scale bar is 400 µm (A).

TG (B) and Cholesterol levels (C) were examined in OA-treated HepG2 cells transfected with MIR NC

or MIR20B. RNA-seq was performed on sample from HepG2 cell with or without overexpression of

MIR20B.Volcano plot of the gene expressions (log2 fold change) compared to the negative control

from RNA-seq analysis (D). Top ranked GSEA in overexpressed *MIR20B* compared to MIR NC in

HepG2 cells (E). The primary ranked enrichment plot of nuclear receptors pathway (F). Heatmap of

the genes in NR pathway upon *MIR20B* overexpression compared to control (G). Expression of

primary ranked nuclear receptors pathway genes from RNA-seq analysis in HepG2 cells, Huh-7 cells,

and primary hepatocytes transfected with MIR20B were normalized to each cells transfected with MIR

NC (H). Western blot analysis of PPARα on MIR NC or MIR20B transfected cells. The intensity of

PPARα blot was normalized to that of HSP90 (I). Venn diagram of predicted targets for MIR20B in

four major database system (J). Graphic image of the conserved binding motifs of *MIR20B* within

3'UTR mRNA of *PPARA* (K). Luciferase activities of MIR20B-transfected HepG2 cells and Huh-7 cells

containing the luciferase reporter DNA constructs for wild-type or mutated 3'UTR of *PPARA* were

normalized to those or MIR NC-transfected cells (L, M). Values represent means ± SEM (n = 3). *$P <$

0.05, **$P < 0.01$, ***$P < 0.001$ *vs* MIR NC. ##$P < 0.01$ *vs* MIR NC + OA. \$\$\$$P < 0.001$ *vs* MIR20B (30

pmol/ml).

Figure 2-figure supplement 1. Analysis of PPARα related pathway in RNA-seq of *MIR20B*

overexpressed HepG2 cells.

HepG2 cells were transfected either with MIR NC or MIR20B and analyzed by RNA-seq. Analysis of

the MIR20B compared to the negative control (n = 3) *via* PCA plot from RNA-seq data (A). Gene set

enrichment pathway analysis of PPARα-related pathways, including PPARA pathway and PPAR

signaling pathway in overexpressed *MIR20B* compared to MIR NC (B, C).

Figure 2-figure supplement 2. *PPARA* is the primary target of the overlapped candidates.

The seven predicted MIR20B targets (*PPARA, NAGK, USP46, ZNFX1, ITGB8, EPHA5, BMPR2*)

were overlapped in miRNA target prediction programs. HepG2 cells were transfected with MIR NC or

MIR20B and the expression of seven predicted MIR20B targets were measured by quantitative RT-

PCR. Relative values are normalized to MIR NC. Values represent means ± SEM (n = 3). *$P < 0.05$,

***$P < 0.001$ *vs* MIR NC.

Figure 2-figure supplement 3. Correlation between the expression of nuclear receptors and

MIR20B **during NAFLD progression.**

The expression of nuclear receptors (*PPARA, RORA, RORC, THRB*), potential target genes of

MIR20B, was measured in the liver of patient with indicated condition by quantitative RT-PCR. The

level of nuclear receptors in patients with steatosis or NASH was normalized to that of normal

individuals (A). Pearson correlation was analyzed between the expression ($2^{-\Delta Ct}$) of *MIR20B* and

nuclear receptors in all study population (B-E). Data were presented as mean ± SEM. ns, not

significant. *$P < 0.05$ and **$P < 0.01$ *vs* normal individuals. ##$P < 0.01$ *vs* patients with steatosis.

Figure 2-figure supplement 4. The expression of *PPARA* is regulated both in human and mice.

The expression of *PPARA* was measured in HepG2, Huh-7 cells, and mice liver tissues by

quantitative RT-PCR. *PPARA* expression levels from HepG2 cells (A) and Huh-7 cells (B) treated with

OA for 24 hours were normalized to no treatment (NT). Hepatic *Ppara* expression levels from

C57BL/6J mice fed a HFD (C) were normalized to NCD. Values represent means ± SEM (n = 3-5).

*$P < 0.05$, **$P < 0.01$ *vs* NT in cells or NCD-fed mice.

Figure 2-figure supplement 5. *PPARA*-3'UTR mutant blocks the effects of MIR20B on lipid

metabolism. Schematic image of expressing plasmids for *PPARA*-3'UTR wild-type (WT) or mutated

(Mut) (A). Lipid accumulation in 1 mM OA-treated HepG2 cells transfected with plasmids and miRNA

was stained by Oil Red O. Scale bar is 400 μm (B, Left). The staining was normalized to that of

HepG2 cells transfected with vector and MIR NC without OA (B, Right). ns, not significant. *$P < 0.05$,

***$P < 0.001$ *vs* MIR NC. ##$P < 0.01$, ###$P < 0.001$ *vs* MIR NC + OA. The expression of genes related

to FA β-oxidation (C) and FA uptake (D) was measured by quantitative RT-PCR. Relative values are

normalized to that of HepG2 cells transfected with vector and MIR NC without OA. Values represent

means ± SEM (n = 3). ns, not significant. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$.

FIGURE 3. The effects of MIR20B are mediated by PPARα.

HepG2 cells transfected with indicated miRNA, siRNA, or *PPARA* expression vector. Western blot

analysis of PPARα. The intensity of PPARα blot was normalized to that of HSP90 (A, D). Luciferase

activity using the luciferase reporter DNA constructs containing PPRE (*PPAR* response element) was

transfected in HepG2 cells. Luciferase activity was normalized to renilla activity (B, E). mRNA level of

genes involved in FA β-oxidation, lipogenesis, and FA uptake was analyzed by real-time qPCR (C

and F). After transfected with MIR NC or MIR20B, HepG2 cells were treated with fenofibrate (100 µM).

Protein level of PPARα was analyzed by western blot and normalized to that of HSP90 (G). The

transcriptional activity of PPARα was measured (H). Genes involved in of FA β-oxidation, lipogenesis

and FA uptake were determined by real-time qPCR (I). Relative values are normalized to MIR NC.

Values represent means ± SEM (n = 3). ns, not significant. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ *vs* MIR

NC. [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.001$ *vs* MIR20B or anti-MIR20B. [\$]$P < 0.05$, [\$\$]$P < 0.01$, [\$\$\$]$P < 0.001$

vs MIR NC + fenofibrate.

FIGURE 4. MIR20B regulates fatty acid metabolism.

HepG2 cells were transfected with MIR20B or anti-MIR20B and treated with OA for 24 hours. The

expression of *MIR20B* (A, L) and genes related to FA β-oxidation, lipogenesis, FA uptake (B, M) and

glycerolipid synthesis (C, N) were measured by quantitative RT-PCR. Representative mitochondrial

metabolites were measured in HepG2 cells (D, O). The expression of genes related to mitochondrial

biogenesis (E, P) were measured by quantitative RT-PCR. The mitochondrial copy of VIPR1 and MT-

ATP6 were determined (F, Q). OCR (G, R), basal and maximal OCR (H, S), and ATP levels (I, T)

were measured in HepG2 cells. FA uptake (J, U) and β-oxidation (K, V) activity were measured using

[9,10-[3]H(N)]-Palmitic Acid and normalized to the total protein content. Relative values are normalized

to MIR NC. Values represent means ± SEM (n = 3-5). ns, not significant. *$P < 0.05$, **$P < 0.01$,

***$P < 0.001$ *vs* MIR NC. [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.001$ *vs* MIR NC + OA. [\$]$P < 0.05$, [\$\$]$P < 0.01$,

[\$\$\$]$P < 0.001$ *vs* MIR20B or anti-MIR20B, respectively.

Figure 4-figure supplement 1. The expression of PPARα target genes is regulated by MIR20B.

HepG2 cells were transfected with MIR20B (A) or anti-MIR20B (B) and treated with OA for 24 hours.

The expression of PPARα target genes was measured by quantitative RT-PCR. Relative values are

normalized to MIR NC. Values represent means ± SEM (n = 3). ns, not significant. *$P < 0.05$,

**$P < 0.01$, ***$P < 0.001$ *vs* MIR NC. #$P < 0.05$, ##$P < 0.01$, ###$P < 0.001$ *vs* MIR NC + OA

Figure 4-figure supplement 2. MIR20B enhances lipid accumulation in primary hepatocytes

under OA-treatment.

Primary hepatocytes were transfected with MIR NC or MIR20B, subsequently treated with OA (1

mM). Representative Oil Red O staining revealed intracellular lipid accumulation (A). The level of

triglyceride (B) and cholesterol (C) were analyzed. The expression of *MIR20B* (D) and genes related

to FA β-oxidation, lipogenesis and FA uptake (E) was measured by quantitative RT-PCR. Values

represent means ± SEM (n = 3). ns, not significant. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ *vs* MIR NC.

#$P < 0.05$, ##$P < 0.01$, *vs* MIR NC + OA. $$$P < 0.01$, $$$$P < 0.001$, *vs* MIR20B.

Figure 4-figure supplement 3. Ketogenesis is not regulated by MIR20B in HepG2 cells.

HepG2 cells were transfected with MIR20B, anti-MIR20B, or MIR NC as control. The expression of

genes involved in ketogenesis (*HMGCL, ACAT1 and ACAT2*) were measured by quantitative RT-PCR.

Relative values are normalized to MIR NC (A, C). The concentration of β-hydroxybutyrate which

generated by ketogenesis was examined in HepG2 cells (B, D). Values represent means ± SEM

(n = 3). ns, not significant. *$P < 0.05$ *vs* MIR NC.

Figure 4-figure supplement 4. Inhibition of MIR20B alleviates lipid accumulation in HepG2 cells.

HepG2 cells and primary hepatocytes were transfected with MIR NC or anti-MIR20B, subsequently

treated with OA (1 mM). Representative Oil Red O staining revealed intracellular lipid accumulation (A,

D). The level of triglyceride (B, E) and cholesterol (C, F) were analyzed. The expression of *MIR20B* (G)

and genes related to FA β-oxidation, lipogenesis and FA uptake (H) were measured by quantitative

RT-PCR. HepG2 cells were infected with lentivirus containing shRNA non-targeting control (shNC) or

shRNA targeting *PPARA* (*shPPARA*) and transfected with either MIR NC or anti-MIR20B,

subsequently treated with OA (1 mM). Expression of *PPARA* and *FASN* was measured by

quantitative RT-PCR (I). Values represent means ± SEM (n = 3). ns, not significant. *P < 0.05,

**P < 0.01, ***P < 0.001 *vs* MIR NC. #P < 0.05, ##P < 0.01 *vs* MIR NC+OA. $$P$ < 0.01, $$$P$ < 0.001 *vs*

anti-MIR20B or anti-MIR20B + OA, respectively.

Figure 4-figure supplement 5. The expression of genes associated with FA uptake by MIR20B

in HepG2 cells.

HepG2 cells were transfected with MIR20B (A) or anti-MIR20B (B), subsequently treated with OA (1

mM) for 24 hours. The expression of genes related to FA uptake were measured by quantitative RT-

PCR. Relative values are normalized to MIR NC. Values represent means ± SEM (n = 3). ns, not

significant.

1001

1002　**Figure 4-figure supplement 6. The expression of *MIR20B* by OA in HepG2 cells.**

1003　 HepG2 cells were incubated with OA (1 mM) for indicated time. The expression of *MIR20B* was

1004　measured by quantitative RT-PCR. Relative values are normalized to NT. Values represent

1005　means ± SEM (n = 3). **P < 0.01, ***P < 0.001 *vs* NT. #P < 0.05 *vs* OA 24 h.

1006

Figure 5. MiR20b promotes hepatic steatosis in HFD-fed mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n = 10

per group) for 12 weeks and administered with indicated AAVs. Hepatic expression of *Mir20b* (A) and

PPARα (B), body weight (C), the ratio of fat mass to body weight (D), the ratio of lean mass to body

weight (E), representative images and weight of liver (F), representative images of H&E staining and

Oil Red O staining of liver slides (G), NAFLD activity score (NAS) (H), hepatic TG (I), serum AST (J),

serum ALT (K), glucose tolerance (L), insulin tolerance (M), fasting glucose (N), fasting insulin (O),

and HOMA-IR (P) were analyzed in indicated mice. Genes related to FA β-oxidation, lipogenesis and

FA uptake (Q) and PPARα target genes (R) were determined by quantitative RT-PCR. Relative

values are normalized to AAV-Control NCD. The intensity of PPARα blot was normalized to that of

HSP90. Values represent means ± SEM (n = 10). ns, not significant. $*P < 0.05$, $**P < 0.01$,

$***P < 0.001$ *vs* AAV-Control NCD. [#]$P < 0.05$, [##]$P < 0.01$, [###]$P < 0.001$ *vs* AAV-Control HFD. [$]$P < 0.05$,

[$$$]$P < 0.001$ *vs* AAV-*Mir20b* NCD. Scale bar is 100µm

1020

Figure 5-figure supplement 1. The expression of *Mir20b* in adipose tissues of AAV injected

mice. C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n

= 10 per group) for 12 weeks and administered with indicated AAVs. All tissues were harvested at the

same time. Quantitative RT-PCR analysis of *Mir20b* expression in epididymal white adipose tissue (A),

inguinal white adipose tissue (B), brown adipose tissue (C) in AAV-Control or AAV-*Mir20b* injected

mice. Values represent means ± SEM (n = 10). ns, not significant. $*P < 0.05$ *vs* AAV-Control NCD.

1027

Figure 5-figure supplement 2. MiR20b significantly regulates *Ppara* in HFD-fed mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n =

10 per group) for 12 weeks and administered with indicated AAVs. The expression of nuclear

receptors, MiR20B target candidates from regulatory network analysis and RNA-seq, was measured

by quantitative RT-PCR in liver of AAV-Control or AAV-*Mir20b* injected mice. Values represent

means ± SEM (n = 10). ns, not significant. $*P < 0.05$, $**P < 0.01$ *vs* AAV-Control NCD. [##]$P < 0.01$ *vs*

AAV-Control HFD.

1035

Figure 5-figure supplement 3. The weight of adipose tissues of AAV injected mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n =

10 per group) for 12 weeks and administered with indicated AAVs. All tissues were harvested at the

same time. Wet weight of epididymal white adipose tissue (A), inguinal white adipose tissue (B), and

brown adipose tissue (C) in AAV-Control or AAV-*Mir20b* injected mice. Values represent

means ± SEM (n = 10). ns, not significant. $***P < 0.01$ *vs* AAV-Control NCD. [#]$P < 0.05$ *vs* AAV-Control

HFD.

1043

Figure 5-figure supplement 4. The expression of genes involved in lipolysis in adipose tissue

and serum FFA in AAV injected mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n = 10

per group) for 12 weeks and administered with indicated AAVs. The expression of genes involved in

lipolysis was analyzed by quantitative RT-PCR in epididymal white adipose tissue of AAV-Control or

AAV-*Mir20b* injected mice (A). Serum FFA (B) was also examined. Relative values are normalized to

AAV-Control. Values represent means ± SEM (n = 10). ns, not significant. $**P < 0.01$, $***P < 0.001$ *vs*

AAV-Control NCD.

1052

Figure 6. Inhibition of MiR20b alleviates hepatic steatosis in HFD-fed mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n = 10

per group) for 12 weeks and administered with indicated AAVs. Hepatic expression of *Mir20b* (A) and

PPARα (B) and body weight (C), the ratio of fat mass to body weight (D), the ratio of lean mass to

body weight (D), representative images and weight of liver (F), representative images of H&E staining

and Oil Red O staining of liver slides (G), NAFLD activity score (NAS) (H), hepatic TG (I), serum AST

(J), and serum ALT (K), glucose tolerance (L), insulin tolerance (M), fasting glucose (N), fasting insulin

(O), and HOMA-IR (P) were analyzed in indicated mice. Genes related to FA β-oxidation, lipogenesis

and FA uptake (Q) and PPARα target genes (R) were determined by quantitative RT-PCR. Relative

values are normalized to AAV-Control NCD. The intensity of PPARα blot was normalized to that of

HSP90. Values represent means ± SEM (n = 10). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ *vs* AAV-Control

NCD. $^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.001$ *vs* AAV-Control HFD. $^{\$}P < 0.05$, $^{\$\$}P < 0.01$, $^{\$\$\$}P < 0.001$ *vs*

AAV-anti-*Mir20b* NCD. Scale bar is 100μm.

1066

Figure 6-figure supplement 1. The expression of *Mir20b* in adipose tissues of AAV injected

mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n =

10 per group) for 12 weeks and administered with indicated AAVs. All tissues were harvested at the

same time. Quantitative RT-PCR analysis of *Mir20b* expression in epididymal white adipose tissue (A),

inguinal white adipose tissue (B), brown adipose tissue (C) in AAV-Control or AAV-anti-*Mir20b*

injected mice. Relative values are normalized to AAV-Control. Values represent means ± SEM

(n = 10). ns, not significant.

1075

Figure 6-figure supplement 2. Inhibition of MiR20b significantly increases hepatic *Ppara* in

HFD-fed mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n =

10 per group) for 12 weeks and administered with indicated AAVs. The expression of nuclear

receptors, MIR20B target candidates from regulatory network analysis and RNA-seq, was measured

by quantitative RT-PCR in liver of AAV-Control or AAV-anti-MIR20B injected mice. Values represent

means ± SEM (n = 10). ns, not significant. *$P < 0.05$, ***$P < 0.001$ *vs* AAV-Control NCD. [###] $P < 0.001$

vs AAV-Control HFD.

1084

Figure 6-figure supplement 3. The weight of adipose tissues of AAV injected mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n = 10

per group) for 12 weeks and administered with indicated AAVs. All tissues were harvested at the

same time. Wet weight of epididymal white adipose tissue (A), inguinal white adipose tissue (B), and

brown adipose tissue (C) in AAV-Control or AAV-anti-*Mir20b* injected mice. Values represent

means ± SEM (n = 10). ns, not significant. ***$P < 0.01$ *vs* AAV-Control NCD.

1091

Figure 6-figure supplement 4. The expression of genes involved in lipolysis in adipose tissue

and serum FFA in AAV injected mice.

C57BL/6J mice were fed on normal chow diet (NCD, n = 10 per group) or high fat diet (HFD, n =

10 per group) for 12 weeks and administered with indicated AAVs. The expression of genes involved

in lipolysis was analyzed by quantitative RT-PCR in epididymal white adipose tissue of AAV-Control

or AAV-anti-*Mir20b* injected mice (A). Serum FFA (B) was also examined. Relative values are

normalized to AAV-Control. Values represent means ± SEM (n = 10). ns, not significant. ***$P < 0.001$

vs AAV-Control NCD.

1100

Figure 7. The effects of fenofibrate are limited in *Mir20b*-introduced mice.

C57BL/6J mice were fed on high fat diet (HFD) for 12 weeks and administered with indicated AAVs

(n = 3 - 5 per group). Then, mice injected with vehicle or fenofibrate (100 mg/kg) for 4 weeks. Hepatic

Mir20b (A) and PPARα expression (B), body weight (C), representative images and weight of liver (D),

the ratio of fat mass to body weight (E), the ratio of lean mass to body weight (F), representative

images of H&E staining and Oil Red O staining of liver slides (G), NAFLD activity score (NAS) (H),

hepatic TG (I), serum AST (J), and serum ALT (K), glucose tolerance (L), insulin tolerance (M), fasting

glucose (N), fasting insulin (O), and HOMA-IR (P) were analyzed in indicated mice. Genes related to

FA β-oxidation, lipogenesis and FA uptake were determined by quantitative RT-PCR (Q). Relative

values are normalized to AAV-Control HFD. The intensity of PPARα blot was normalized to that of

HSP90. The intensity of PPARα blot was normalized to that of HSP90. Values represent

means ± SEM (n = 3 - 5). ns, not significant, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$ *vs* AAV-Control HFD.

#$P < 0.05$ *vs* AAV-*Mir20b* HFD. $$P < 0.05$, $$$P < 0.01$, $$$$P < 0.001$ *vs* AAV-Control HFD + fenofibrate.

Scale bar is 100μm.

1115

Figure 8. MiR20b promotes liver inflammation and fibrosis in MCD-Fed Mice.

Heatmap (A) and correlation (B) of hepatic nuclear receptor gene expression in RNA-seq (Figure 2)

with public databases of liver fibrosis or NASH patients. The values are fold change (FC) compared to

each control samples. C57BL/6J mice were fed on normal chow diet (NCD, n = 5 per group) or

methionine-deficient diet (MCD, n = 5 per group) for 4 weeks and administered with indicated AAVs.

Before 1 week of MCD challenge, mice were injected with AAV-Control or AAV-anti-*Mir20b*. Hepatic

Mir20b (C) and PPARα expression (D, E, F), representative images of liver (G), H&E staining, Oil Red

O, staining and Sirius Red staining of liver slides (H), NAFLD activity score (NAS) (I), fibrosis score (J),

hepatic TG (K), serum AST (L), and serum ALT (M) were analyzed in indicated mice. Genes related to

inflammation (N) and fibrosis (O) were determined by quantitative RT-PCR. Relative values are

normalized to AAV-Control NCD. The intensity of PPARα blot was normalized to that of HSP90.

Values represent means ± SEM (n = 5). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ vs AAV-Control NCD. $^{\#}P <$

0.05, $^{\#\#}P < 0.01$ vs AAV-Control MCD. Scale bar is 100μm.

1129

Figure 9. Inhibition of MiR20b with fenofibrate ameliorates liver inflammation and fibrosis in

MCD-fed Mice.

C57BL/6J were fed a methionine-deficient diet (MCD) for 4 weeks with administration of indicated

AAVs (n = 5 - 6 per group). Then, mice injected with vehicle or fenofibrate (100 mg/kg) for 4 weeks.

Hepatic *Mir20b* (A) and PPARA expression (B, C). Representative images of liver (D), hepatic TG (E),

H&E staining, Oil Red O staining, Sirius Red staining of liver slides (F), NAFLD activity score (NAS)

(G), fibrosis score (H), serum AST (I), and serum ALT (J) were analyzed in indicated mice. Genes

related to inflammation (K) and fibrosis (L) were determined by quantitative RT-PCR. Relative values

are normalized to AAV-Control MCD. The intensity of PPARα blot was normalized to that of HSP90.

Values represent means ± SEM (n = 5 - 6). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ vs AAV-Control MCD.

$^{\#}P < 0.05$, $^{\#\#}P < 0.01$, $^{\#\#\#}P < 0.001$ vs AAV-Control MCD + fenofibrate. $^{\$}P < 0.05$, $^{\$\$}P < 0.01$ vs AAV-

anti-*Mir20b* MCD. Scale bar is 100μm.

1142

Figure 10. E2F1 is upregulated in both NAFLD patients and mice model.

The expression of *E2F1* was analyzed by quantitative RT-PCR. Hepatic *E2F1* expression levels of

steatosis or NASH patients were normalized to those of normal patients. $*P < 0.05$ and $**P < 0.01$ vs

normal patients (A). *E2F1* expression levels from HepG2 cells (B) and Huh-7 cells (C) treated with OA

for 24 hours were normalized to no treatment (NT). Hepatic *E2f1* expression levels from C57BL/6J

mice fed a HFD (D) and a MCD (E) were normalized to NCD. Values represent means ± SEM (n = 3 -

11). $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ vs NT in cells or NCD-fed mice, respectively.

Figure 1. *MIR20B* expression is significantly increased in the livers of dietary obese mice and humans.

Figure 2. *PPARA* is a direct target of MIR20B.

Figure 3. The effects of MIR20B are mediated by PPARα.

Figure 4. MIR20B regulates fatty acid metabolism.

Figure 5. MiR20b promotes hepatic steatosis in HFD-fed mice.

Figure 6. Inhibition of MiR20b alleviates hepatic steatosis in HFD-fed mice.

Figure 7. The effects of fenofibrate are limited in *Mir20b*-introduced mice.

Figure 8. MiR20b promotes liver inflammation and fibrosis in MCD-Fed Mice.

Figure 9. Inhibition of MiR20b with fenofibrate ameliorates liver inflammation and fibrosis in MCD-fed Mice.

Figure 10. E2F1 is upregulated in both NAFLD patients and mice model.