1
2
3

Reducing susceptibility distortion related image blurring in diffusion MRI EPI data

6
7

Ian A. Clark[1], Martina F. Callaghan[1], Nikolaus Weiskopf[1,2,3], Eleanor A. Maguire[1*], Siawoosh Mohammadi[4*]

[1]Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University College London, London, UK

[2]Department of Neurophysics, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[3]Felix Bloch Institute for Solid State Physics, Faculty of Physics and Earth Sciences, Leipzig University, Leipzig, Germany

[4]Institute of Systems Neuroscience, University Medical Centre Hamburg-Eppendorf, Hamburg, Germany

*Correspondence

Eleanor Maguire, Wellcome Centre for Human Neuroimaging, UCL Queen Square Institute of Neurology, University College London, 12 Queen Square, London, WC1N 3AR, UK**;** e.maguire@ucl.ac.uk

Siawoosh Mohammadi, Institute of Systems Neuroscience, University Medical Centre Hamburg-Eppendorf, Hamburg, Germany; s.mohammadi@uke.de

Keywords: diffusion MRI; distortion correction; arithmetic mean; weighted average; multi-parameter mapping; g-ratio mapping; myelin; SPM

Word count: 5982

Number of figures: 8

Number of tables: 1

ABSTRACT

Diffusion magnetic resonance imaging (MRI) is an increasingly popular technique in basic and clinical neuroscience. One promising application is to combine diffusion MRI with myelin maps from complementary MRI techniques such as multi-parameter mapping (MPM) to produce g-ratio maps that represent the relative myelination of axons and predict their conduction velocity. Statistical Parametric Mapping (SPM) can process both diffusion data and MPMs, making SPM the only widely accessible software that contains all the processing steps required to perform group analyses of g-ratio data in a common space. However, limitations have been identified in its method for reducing susceptibility-related distortion in diffusion data. More generally, susceptibility-related image distortion is often corrected by combining oppositely phase-encoded images (blip-up and blip-down) using the arithmetic mean (AM), however, this can lead to blurred images. In this study we sought to (1) improve the susceptibility-related distortion correction for diffusion MRI data in SPM; (2) deploy an alternative approach to the AM to reduce image blurring in diffusion MRI data when combining blip-up and blip-down EPI data after susceptibility-related distortion correction; and (3) assess the benefits of these changes for g-ratio mapping. We found that the new processing pipeline, called consecutive HySCO, improved distortion correction when compared to the standard approach in the ACID toolbox for SPM. Moreover, using a weighted average (WA) method to combine the distortion corrected data from each phase-encoding polarity achieved greater overlap of diffusion and more anatomically faithful structural white matter probability maps derived from minimally distorted multi-parameter maps as compared to the AM. Third, we showed that the consecutive HySCO WA performed better than the AM method when combined with multi-parameter maps to perform g-ratio mapping. These improvements mean that researchers can conveniently access a wide range of diffusion-related analysis methods within one framework because they are now available within the open-source ACID toolbox as part of SPM, which can be easily combined with other SPM toolboxes, such as the hMRI toolbox, to facilitate computation of myelin biomarkers that are necessary for g-ratio mapping.

INTRODUCTION

Diffusion magnetic resonance imaging (MRI) is an increasingly popular method in neuroscience and clinical research. Diffusion MRI permits visualisation of the mobility of water molecules within brain tissue, and provides measures of microstructural integrity and anatomical connectivity (Conturo et al., 1999; Shimony et al., 1999; Le Bihan et al., 2001; Kaden et al., 2007). Knowledge of how different brain regions communicate is essential for understanding brain anatomy and its relationship with cognitive processes (Jbabdi et al., 2015; Hodgetts et al., 2017; Assaf et al., 2019; Movahedian Attar et al., 2020), and diffusion imaging in the clinic can be used for various purposes including the characterisation of strokes and tumours (Neumann-Haefelin et al., 1999; Bergui et al., 2001).

However, diffusion MRI data are generally acquired with echo planar imaging (EPI; Turner and Le Bihan, 1990) – a fast MRI acquisition technique that is prone to spatial distortions related to off-resonance effects (Jezzard and Balaban, 1995). It is, therefore, important to correct spatial distortions in diffusion data to provide as accurate a representation as possible of the true anatomy. Fortunately, various methods to address spatial distortions in diffusion MRI are available. Field inhomogeneities due to magnetic susceptibility variations can be measured with additional MRI sequences for field mapping (Jezzard and Balaban, 1995; Reber et al., 1998; Hutton et al., 2002) or estimated from the distorted images themselves using physical models (Chang and Fitzpatrick, 1992). A popular and efficient approach uses two EPI images with reversed phase-encoding directions – also known as "blip-up" and "blip-down" images (Figure 1A) – to estimate the field map and correct the susceptibility-related distortions (Andersson et al., 2003; Weiskopf et al., 2005; Holland et al., 2010; Ruthotto et al., 2012; Breman et al., 2020). The issue remains, however, that even after distortion correction is applied, the corrected diffusion MRI images can suffer from additional image blurring artefacts because the compression of voxels leads to a loss of information. As illustrated in Figure 1, blip-up and blip-down images are often combined by calculating the arithmetic mean (AM) of the two images to provide a single diffusion image (Figure 1B, top).

Insert Figure 1 about here

Avoiding residual spatial distortions and image blurring is particularly important when combining diffusion MRI with other quantitative MRI contrasts that can reveal additional information that would otherwise not be available (Mohammadi and Callaghan, 2018). For example, multi-parameter mapping (MPM; Weiskopf et al., 2013) is an emerging neuroimaging technique that provides a comprehensive approach to acquiring multiple quantitative biomarkers, namely, the longitudinal relaxation rate R_1, proton density PD, effective transverse relaxation rate R_2^*, and magnetization transfer saturation MT_{sat} (Tabelow et al., 2019). These quantitative parameters contain specific and complementary information regarding the microstructural properties of brain tissue, such as the extent of myelination, iron and water concentration (e.g. Weiskopf et al., 2015; Kirilina et al., 2020). Combining diffusion MRI with MPM thus allows for the examination of microstructural information related to the neuronal conduction velocity in the form of the axonal g-ratio (Mohammadi et al., 2015a; Mohammadi and Callaghan, 2020). Furthermore, by combining diffusion MRI and MPM structural maps, it becomes possible to perform multivariate analyses with biomarkers sensitive to complementary tissue features (Draganski et al., 2011; Chowdhury et al., 2013). While various methods using reverse phase-encoding are available to efficiently reduce susceptibility-related distortions (for example, see Gu and Eklund, 2019), these implementations often focus

only on diffusion MRI data. As such, to combine diffusion MRI with other quantitative MRI
data, such as MPM for g-ratio mapping, the use of multiple toolboxes implemented in different
software packages, is currently required.

FSL, together with TBSS for group analyses, is probably the most widely used package
for diffusion MRI (Smith et al., 2004), with TOPUP for distortion correction (Andersson et al.,
2003) and eddy for motion and eddy current correction (Andersson and Sotiropoulos, 2016).
To examine the g-ratio, however, the diffusion data have to be combined with a quantitative
MRI biomarker for the myelin compartment, requiring laboratory-specific in-house software
that is not widely available (Stikov et al., 2015; Cercignani et al., 2017; Mancini et al., 2018).
In another analysis package, SPM (www.fil.ion.ucl.ac.uk/spm/), the ACID toolbox
(www.diffusiontools.com) can be used for distortion correction using HySCO2 (Ruthotto et
al., 2012; Ruthotto et al., 2013; Macdonald and Ruthotto, 2018) and motion and eddy current
correction using ECMOCO (Mohammadi et al., 2010; Mohammadi et al., 2015b). Moreover,
the hMRI toolbox, also in SPM (www.hmri.info; Tabelow et al., 2019), provides an easy-to-
use framework to generate quantitative structural maps from MPM data. SPM, therefore, has
the advantage of being able to process both diffusion and MPM data, making SPM the only
widely accessible software that contains all the processing steps required to perform group
analyses of g-ratio data in a common space. However, limitations have been identified in the
ability of the HySCO2 methodology to reduce susceptibility-related distortion (Gu and Eklund,
2019), highlighting a need for improvement. Moreover, the HySCO2 method uses the AM to
combine the oppositely phase-encoded data after correction, meaning that blurring artefacts are
typically present.

Here, we sought to improve the pipeline for the HySCO2-based distortion correction,
an approach we call consecutive HySCO, and to employ a different method to combine the
corrected blip-up and blip-down images to reduce image blurring by using a weighted average
(WA). We achieved this by down-weighting regions that were squeezed (e.g. in the blip-up
images) while up-weighting the same (stretched) regions in the opposite (blip-down) image.
To determine regions that were stretched and squeezed, we used the Jacobian of the HySCO2
estimated field map. We compared the performance of the proposed WA approach to the
standard AM combination used by HySCO2 in SPM. Where appropriate, we also benchmarked
these changes against the widely available default pipeline in FSL (as used in the human
connectome project; Glasser et al., 2013). Finally, given that the overall goal was to have all
the processing steps necessary to perform g-ratio analysis within one (SPM) package, we also
assessed whether there was any benefit of using the WA corrected diffusion data over the AM
corrected data for g-ratio mapping.

MATERIALS AND METHODS

Participants

Eighteen participants took part in the study. They were aged between 20 and 40 years old,
reported no psychological, psychiatric, neurological or behavioural health conditions. The
mean age of the sample was 30.0 years (SD = 6.07) and included 9 females and 9 males.
Participants were reimbursed £10 per hour for taking part which was paid at study completion.
All participants gave written informed consent and the study was approved by the University
College London Research Ethics Committee.

Diffusion MRI data acquisition

Diffusion-weighted images were collected on Siemens Magnetom 3T TIM Trio systems using 32 channel head coils. The protocol used was the Multi-Band Accelerated EPI Pulse Sequence developed by the Centre for Magnetic Resonance Research at the University of Minnesota (R012a-c, R013a on VB17, https://www.cmrr.umn.edu/multiband/; Feinberg et al., 2010; Xu et al., 2013). Acquisition parameters were: resolution = 1.7 mm isotropic; FOV = 220 mm × 220 mm × 138 mm; 60 directions with 6 interleaved b0 images, echo time (TE) = 112 ms, repetition time (TR) = 4.84s, with a multiband acceleration factor of 3. The sequence was performed 4 times – twice with b-values of 1000 and twice with b-values of 2500. The first acquisition of each set of b-values was performed with phase-encoding in the anterior to posterior direction (blip-up), the second in the posterior to anterior direction (blip-down).

Structural MRI data acquisition

Whole brain structural maps of magnetisation transfer (MT) saturation, at an isotropic resolution of 800μm, were derived from an MPM quantitative imaging protocol (Weiskopf et al., 2013; Callaghan et al., 2015; Callaghan et al., 2019). This protocol consisted of the acquisition of three multi-echo gradient-echo acquisitions with either PD, T1 or MT weighting. Each acquisition had a TR of 25 ms. PD weighting was achieved with an excitation flip angle of 6^0, which was increased to 21^0 to achieve T1 weighting. MT weighting was achieved through the application of a Gaussian RF pulse 2 kHz off resonance with 4ms duration and a nominal flip angle of 220^0. This acquisition had an excitation flip angle of 6^0. The field of view was 256mm head-foot, 224mm AP, and 179mm right-left. The multiple gradient echoes per contrast were acquired with alternating readout gradient polarity at eight equidistant echo times ranging from 2.34 to 18.44ms in steps of 2.30ms using a readout bandwidth of 488 Hz/pixel. Only six echoes were acquired for the MT weighted volume to facilitate the off-resonance pre-saturation pulse within the TR. To accelerate the data acquisition, partially parallel imaging using the GRAPPA algorithm was employed in each phase-encoded direction (anterior-posterior and right-left) with forty integrated reference lines and a speed up factor of two. Calibration data were also acquired at the outset of each session to correct for inhomogeneities in the RF transmit field (Lutti et al., 2010; Lutti et al., 2012).

Diffusion MRI distortion correction pipelines

Standard HySCO pipeline. The diffusion MRI data were first processed using the standard pre-processing pipeline available in the ACID toolbox (www.diffusiontools.com) within SPM12 (www.fil.ion.ucl.ac.uk/spm). The blip-up and blip-down data were separately corrected first for motion and eddy current artefacts (Mohammadi et al., 2010), and then for susceptibility-related distortion artefacts using the HySCO2 module (Ruthotto et al., 2012; Ruthotto et al., 2013; Macdonald and Ruthotto, 2018).

The new consecutive HySCO pipeline. As with the standard HySCO pipeline, the blip-up and blip-down data were first separately corrected for motion and eddy current artefacts and then for susceptibility-related distortion artefacts using the HySCO2 module. Tensor fitting (Mohammadi et al., 2013) was then performed separately on each of the distortion corrected blip-up and blip-down datasets to estimate Fractional Anisotropy (FA) maps. HySCO2 was then repeated but now using the distortion corrected and brain-masked FA maps as input instead of b0 images; the second HySCO2 field map being consecutively applied to the "pre-corrected" diffusion MRI data (Figure 2).

Insert Figure 2 about here

FSL pipeline. While our main interest was in comparing the two HySCO approaches, the
diffusion MRI data were also processed based on the widely-used human connectome project
pipeline (Glasser et al., 2013) in FSL 6.0.1 (Smith et al., 2004) to provide external
benchmarking. The blip-up and blip-down data were separately corrected for distortions using
TOPUP (Andersson et al., 2003), which was applied to the data along with correction for
motion and eddy current artefacts using eddy (Andersson and Sotiropoulos, 2016) with the
replace outliers option enabled (Andersson et al., 2016).

Diffusion MRI phase-encoding combination methods

For both the consecutive HySCO and FSL methods, the distortion-corrected data sets with
opposite phase-encoding polarity were combined using the ACID toolbox within SPM12.

First, the consecutive HySCO blip-up and blip-down distortion corrected data were
combined by calculating the AM of each blip-up and blip-down image pair.

Second, the consecutive HySCO blip-up and blip-down distortion corrected data were
combined using a WA. This aimed to minimise information loss due to susceptibility distortion
blurring induced by local spatial compression. To maximise the effective spatial resolution
when combining the two images acquired with opposite phase-encoding directions, a region
that was stretched in one of the images, increasing effective resolution, was up-weighted
whereas the opposite image was down-weighted using a WA:

218
$$S^{wa} = \frac{S^\uparrow f(Jac_\uparrow) + S^\downarrow f(Jac_\downarrow)}{f(Jac_\uparrow) + f(Jac_\downarrow)} \quad (1)$$

219   with $f(x) = 2 - \frac{2}{exp((abs(x) - x_0)\,k_0) + 1}$ being a Fermi function with two parameters (x_0, k_0),
220   tuning the sensitivity of the weighting of the Jacobian in a given voxel. Both parameters were
221   heuristically optimized. Note that the Jacobian of the field map estimated in the initial HySCO
222   step (Step 1, Figure 2) was used for weighting. The Jacobians were defined as follows:

223
$$Jac_\downarrow = 1 - \frac{db_H}{dp} = 2 - Jac_\uparrow \quad (2)$$

224   with b_H being the HySCO field in units of displacement in the phase-encoding direction in
225   mm and p being the phase-encoding direction – here the anterior-posterior direction.

226   Finally, to provide a benchmark, the FSL blip-up and blip-down distortion corrected
227   data were combined by calculating the AM of each blip-up and blip-down image pair.

Structural MRI data pre-processing

The structural MRI data were processed for each participant using the hMRI toolbox (Tabelow
et al., 2019) within SPM12. The default toolbox configuration settings were used, with the
exception that correction for imperfect spoiling was additionally enabled (Corbin and
Callaghan, 2021). The output MT saturation map, used as the anatomically faithful structural
image, quantified the degree of saturation of the steady state signal induced by the application
of the off-resonance pre-pulse, having accounted for spatially varying T_1 times and RF field
inhomogeneity (Helms et al., 2008; Weiskopf et al., 2013).

Each participant's MT saturation map was segmented into white matter probability
maps using the unified segmentation approach (Ashburner and Friston, 2005), but with no bias

field correction (since the MT saturation map does not suffer from any bias field modulation)
and using the tissue probability maps developed by Lorio et al. (2016).

Comparison of the diffusion MRI distortion correction methods

Analyses to investigate the differences between the diffusion MRI distortion correction
methods were performed in native space. For each participant, two FA maps were estimated
for each distortion correction method (standard HySCO, consecutive HySCO, FSL); one FA
map from the distortion corrected blip-up data and the other from the distortion corrected blip-
down data. FA maps were estimated using the Diffusion Kurtosis Fit (DKI) in the ACID
toolbox. Each FA map was then co-registered to an individual's MT saturation white matter
probability map. The difference between the individual blip-up and blip-down FA maps was
calculated, restricted to a white-matter mask. The white-matter mask was generated by
thresholding the MT saturation white matter probability maps above 0.9. From the resulting
difference image, the root mean square over all voxels in the white matter mask was calculated
for each individual for each distortion correction method.

Two-tailed paired t-tests, calculated in SPSS v25 and thresholded at $p < 0.05$, were then
performed to test for significant differences between the root mean square for each of the
distortion correction methods across participants. Effect sizes are reported as Hedge's g_{av},
which is Cohen's d calculated for repeated measures and corrected for the positive bias caused
by using sample estimates (Lakens, 2013). Paired t-tests were used instead of a repeated
measures ANOVA to treat each distortion correction method independently.

Comparison of the diffusion MRI phase-encoding combination methods

Analyses to compare the effects of the phase-encoding combination methods were also
performed in native space. For each participant, three pairs of FA and b0 maps were estimated,
one pair for each phase-encoding combination method (consecutive HySCO AM, consecutive
HySCO WA, FSL). The FA and b0 maps were estimated using DKI fit in the ACID toolbox.

Multichannel segmentation, using the unified segmentation approach (Ashburner and
Friston, 2005), but with the tissue probability maps developed by Lorio et al. (2016), was
performed on each FA and b0 map pair to define the white matter tissue segments. These
diffusion-derived white matter probability maps were then co-registered to each individual's
MT saturation white matter probability map. This allowed for comparison between the white
matter probability map defined from the distortion corrected diffusion images, and the white
matter probability map from the MT saturation structural map. The percentage overlap between
the diffusion white matter probability map and the structural white matter probability map was
then calculated for each phase-encoding combination method.

As well as investigating the overlap of the diffusion and structural white matter
probability maps as a whole, we also focused our analyses on regions that suffered from greater
distortion. To identify these areas, for each participant, a map of the Jacobians was calculated
from the first HySCO iteration. The co-registration transformations determined above were
applied to each participant's Jacobian map to co-register the Jacobian and MT saturation
structural images. The percentage overlap of the diffusion and structural white matter
probability maps was calculated in regions of higher distortions. The masks for the
aforementioned regions were determined by thresholding the Jacobian map at values higher
than 10%, 20% and 30% deviation from unity.

Two-tailed paired t-tests, calculated in SPSS v25 and thresholded at $p < 0.05$, were
performed to test for significant differences between the percentage overlaps for each of the

phase-encoding methods (consecutive HySCO AM, consecutive HySCO WA, FSL), at each
level of distortion (whole white matter map, 10%, distortion, 20% distortion, 30% distortion).
As with the distortion correction methods analyses, effect sizes are reported as Hedge's $g_{av,}$
(Lakens, 2013) and paired t-tests were used to treat each phase-encoding method
independently.

Application to g-ratio mapping

Our final set of analyses aimed to assess any benefits of the consecutive HySCO WA approach
on g-ratio mapping by comparing it to the use of consecutive HySCO AM on the generation
of g-ratio maps. As neuroscientific studies utilising the g-ratio are typically performed at the
group level, analyses were performed in group space. The g-ratio was calculated from axonal-
water fraction (AWF) and MT saturation maps according to Ellerbrock and Mohammadi
(2018):

$$g_{MR} = \sqrt{1 - \frac{MVF_{MR}}{MVF_{MR} + AVF_{MR}}} \qquad (3)$$

with MVF_{MR} being the myelin-volume fraction estimated from the MT saturation map and
AVF_{MR} being the axonal-volume fraction. The AVF_{MR} was estimated as $AVF_{MR} =$
$(1 - MVF_{MR})$ AWF according to Stikov et al. (2015), where AWF was obtained by combining
the intra-cellular fraction (ν_{icvf}) and isotropic fraction (ν_{iso}) maps from the neurite orientation
dispersion and density imaging (NODDI) toolbox
(http://mig.cs.ucl.ac.uk/index.php?n=Tutorial.NODDImatlab; Zhang et al., 2012) as AWF =
$(1 - \nu_{iso}) \nu_{icvf}$. The MT saturation map was obtained from the hMRI toolbox as described
above. For calibration of the MT saturation map to a myelin-volume fraction map ($MVF_{MR} =$
αMT_{sat}), we used the g-ratio based calibration method as reported in Ellerbrock and
Mohammadi (2018) and Mohammadi and Callaghan (2020), with a reference g-ratio value of
0.7 in the splenium (Mohammadi et al., 2015a), yielding $\alpha = 0.217$.

Transformation from native to group space was implemented using the voxel-based
quantification (VBQ) approach as implemented in the hMRI toolbox (Draganski et al., 2011),
which allows for preservation of the quantitative values of the g-ratio maps during smoothing
(Tabelow et al., 2019). Briefly, inter-subject registration using DARTEL (Ashburner, 2007)
was performed on the segmented MT saturation grey and white matter probability maps. The
resulting DARTEL template and deformations were then used to normalize the MT saturation,
g-ratio and Jacobian maps to Montreal Neurological Institute (MNI) space at 1.5 x 1.5 x 1.5mm
resolution with a tissue weighted smoothing kernel of 6mm full width at half maximum.

Differences in g-ratio mapping were determined by comparing the calculated g-ratio
values. As the g-ratio can only be defined when the white matter of the diffusion and structural
(MT saturation) maps overlap, a g-ratio of zero is recorded for non-overlapping voxels
(Mohammadi and Callaghan, 2020). We predicted, therefore, that the poorer overlap between
the AM diffusion data and structural white matter map (due to the higher blurring in the AM
diffusion data) would increase the number of voxels with a g-ratio of zero per participant,
decreasing the averaged g-ratio values at the group-level in regions with high susceptibility-
related distortions.

To quantify the reduction of group-averaged g-ratio values within regions suffering
from high levels of distortion, for each participant we extracted the mean g-ratio from the AM
and WA g-ratio maps from regions previously identified as suffering from distortion. This was
performed using a mask created from thresholding the Jacobian map at 20% (see also
Comparison of the diffusion MRI phase-encoding combination methods section above). We

then compared these to the averaged g-ratio value in white matter using two tailed paired t-
tests, thresholded at p < 0.05, calculated in SPSS v25 with effect sizes reported as Hedge's $g_{av,}$
(Lakens, 2013). We hypothesised that smaller g-ratio values would be apparent in the AM-
based g-ratio map because smoothing with zero-valued g-ratio map voxels, indicative of
misalignment, would systematically lower the g-ratio values.

Finally, to localise the voxels where the AM-based g-ratio values were smaller than the
WA-based g-ratios, we performed a voxel-wise comparison of the AM and WA g-ratio maps
within the same 20% Jacobian distortion mask. We did this using two-tailed paired t-tests to
identify significant differences in g-ratio values at statistical thresholds of p < 0.05 family wise
error (FWE) with a minimum cluster size of 10 voxels.

RESULTS

Comparison of diffusion MRI distortion correction pipelines

The average root mean square of the difference in FA map alignment between the blip-up and
blip-down data was significantly smaller for the consecutive HySCO approach in comparison
to the standard HySCO approach for all participants (Figure. 3, mean difference = 0.017, t(17)
= 8.65, p < 0.001, Hedge's g_{av} = 0.91), indicating better distortion correction by the consecutive
HySCO pipeline. Of note, smaller deviations (Figure 3) were observed between the blip-up and
blip-down data after FSL distortion correction in comparison to the standard HySCO approach
(mean difference = 0.040, t(17) = 10.38, p < 0.001, Hedge's g_{av} = 2.49) and the consecutive
HySCO approach (mean difference = 0.023, t(17) = 8.63, p < 0.001, Hedge's g_{av} = 1.72).

348
349 *Insert Figure 3 about here*
350

Comparison of diffusion MRI phase-encoding combination methods

Having established that consecutive HySCO gave better distortion correction than standard
HySCO, we then turned to examining the consecutive HySCO AM and consecutive HySCO
WA phase encoding combination methods. The improvement to the spatial accuracy gained by
using the WA to combine the opposite phase-encoding data is shown in Figure 4. In panels A
and B, additional white matter, in line with the anatomically accurate white matter tissue
probability derived from the MT saturation map, can be observed in the ventromedial prefrontal
cortex when using the WA compared to using the AM. In panels C and D, additional and better
definition of the white matter, again in line with the anatomically accurate MT saturation white
matter tissue probability map, is evident in the inferior temporal gyrus when using the WA
compared to using the AM. Using the WA seems, therefore, to provide better alignment
between the diffusion and quantitative structural maps.

363
364 *Insert Figure 4 about here*
365

To provide a quantitative examination, we also calculated the mean percentage overlap
between the diffusion and MT saturation white matter tissue probability maps for the
consecutive HySCO AM, consecutive HySCO WA and, as an additional benchmark, FSL (see
Figure 5). This quantitative comparison supported the qualitative visual inspection.

Insert Figure 5 about here

Using a WA to combine the consecutive HySCO distortion corrected blip-up and blip-
down data resulted in greater percentage overlap between the diffusion and MT saturation
white matter tissue probability maps when compared to the consecutive HySCO AM. This was
particularly evident in regions suffering from high levels of distortion (whole white matter:
mean difference = 0.89, $t(17) = 3.10$, $p = 0.007$, Hedge's $g_{av} = 0.40$; 10% distortion: mean
difference = 7.81, $t(17) = 11.92$, $p < 0.001$, Hedge's $g_{av} = 2.27$; 20% distortion: mean difference
= 17.66, $t(17) = 20.73$, $p < 0.001$, Hedge's $g_{av} = 4.38$; 30% distortion: mean difference = 26.23,
$t(17) = 25.94$, $p < 0.001$, Hedge's $g_{av} = 5.64$). Of note, using the AM to combine the FSL
distortion corrected blip-up and blip-down data also resulted in greater percentage overlap
between the diffusion and MT saturation white matter tissue probability maps compared to
using the AM to combine the consecutive HySCO distortion corrected data (whole white
matter: mean difference = 2.53, $t(17) = 8.65$, $p < 0.001$, Hedge's $g_{av} = 1.22$; 10% distortion:
mean difference = 8.15, $t(17) = 13.50$, $p < 0.001$, Hedge's $g_{av} = 2.40$; 20% distortion: mean
difference = 14.09, $t(17) = 16.88$, $p < 0.001$, Hedge's $g_{av} = 3.46$; 30% distortion: mean
difference = 18.97, $t(17) = 17.42$, $p < 0.001$, Hedge's $g_{av} = 3.92$).

Interestingly, across the whole of the white matter map, a larger percentage overlap
between the diffusion and MT saturation white matter tissue probability maps was observed
when using the AM to combine the FSL distortion corrected blip-up and blip-down data
compared to the consecutive HySCO WA (mean difference = 1.64, $t(17) = 6.47$, $p < 0.001$,
Hedge's $g_{av} = 0.88$), although there was no difference between the methodologies at 10%
distortion (mean difference = 0.34, $t(17) = 0.68$, $p = 0.51$, Hedge's $g_{av} = 0.13$). However, we
found that using the WA to combine the consecutive HySCO distortion corrected blip-up and
blip-down data resulted in greater percentage overlap of the diffusion and MT saturation white
matter tissue probability maps compared to FSL in regions suffering from greater levels of
distortion (20% distortion: mean difference = 3.58, $t(17) = 6.13$, $p < 0.001$, Hedge's $g_{av} = 1.19$;
30% distortion: mean difference = 7.26, $t(17) = 9.62$, $p < 0.001$, Hedge's $g_{av} = 1.93$) (Figure
5).

The consecutive HySCO WA method seems, therefore, to be particularly advantageous
in regions suffering from high distortion, for example, in the ventromedial prefrontal cortex (as
shown in Figure 4A and B), inferior temporal gyri (Figure 4C and D), as well as the middle
temporal gyri, medial temporal lobe (including the anterior hippocampi), and the brain stem
(Figure 6).

Insert Figure 6 about here

Application to g-ratio mapping

Finally, we set out to assess whether there was any benefit of using the consecutive HySCO
WA diffusion data over the consecutive HySCO AM data for g-ratio mapping. Figure 7 shows
that the coverage of the g-ratio map across white matter is enhanced when using the WA
combination compared to the AM combination. The focus is the ventromedial prefrontal cortex
and the brain stem, two regions that suffer from high levels of distortion. As can be observed,
greater accuracy was obtained in the white matter probability maps and more information was
available in the AWF map. Consequently, this resulted in a larger area where the g-ratio could

be defined when using the diffusion data corrected using the WA compared to performing the
same calculations on the AM combination.

418
419   *Insert Figure 7 about here*

421   We next sought to examine the g-ratio mapping data at the group level, predicting
higher mean g-ratio values when there was better overlap between the diffusion data and the
MT saturation white matter tissue probability map. As our previous analyses identified that the
WA combination resulted in greater overlap between the diffusion and MT saturation white
matter tissue probability maps in areas suffering from high levels of distortion, we predicted
that improvements in g-ratio mapping would be evident when using the WA methodology in
such regions.

428   To investigate this, we first compared the mean g-ratio values from across the whole of
white matter, finding significantly greater WA-based g-ratios in comparison to the AM-based
g-ratios (WA-based mean = 0.55; AM-based mean = 0.54; mean difference = 0.0071, $t(17)$ =
4.58, $p < 0.001$, Hedge's g_{av} = 0.45). To create a non-biased baseline mean g-ratio value, for
each participant we calculated a combined mean g-ratio for the AM and WA white matter
maps. We then extracted the mean g-ratio from the regions suffering from high levels of
distortion (identified by thresholding the Jacobian map at 20%), and subtracted this from the
combined mean g-ratio in order to provide a measure of the difference in g-ratio between highly
distorted regions and the whole of white matter. We predicted that the difference in mean g-
ratio values between the highly distorted regions and the combined mean g-ratio across the
whole of the AM and WA white matter maps would be smaller for the WA-based g-ratios. This
is because we expected the g-ratio values in the AM-based g-ratio map to be lower in highly
distorted regions, and thus further away from the combined mean g-ratio. This would be due
to the poorer overlap between the diffusion and MT saturation white matter tissue probability
maps increasing the number of voxels with a g-ratio of zero.

443   Indeed, a smaller difference between the g-ratio values in highly distorted regions and
the combined mean g-ratio was apparent for the WA-based g-ratios compared to the AM-based
g-ratios (mean difference = 0.092, $t(17)$ = 12.76, $p < 0.001$, Hedge's g_{av} = 1.98; Figure 8A).
This suggests, therefore, that in regions suffering from high levels of distortion, the WA-based
g-ratio values were closer to the white-matter average than AM-based g-ratio values, thus
providing a better representation of g-ratio values in regions with high susceptibility-related
distortions.

450   Finally, to further localise where, within the regions with high susceptibility-related
distortions, the AM-based g-ratios were significantly smaller than the WA-based g-ratios,
voxel-wise statistics were performed. As can be seen in Figure 8B, higher g-ratio values were
identified (top row) when using the WA combination in regions suffering from the greatest
levels of distortion (bottom row), including the ventromedial prefrontal medial frontal cortex,
brain stem, and inferior and middle temporal gyri (see also Table 1). No significant voxels were
identified for the reverse contrast.

457
458   *Insert Figure 8 and Table 1 about here*

DISCUSSION

Our aims in this study were to improve the susceptibility-related distortion correction for diffusion MRI data in SPM, to deploy an alternative approach to reducing image blurring in diffusion MRI data when combining blip-up and blip-down EPI data after susceptibility-related distortion correction, and to assess any benefits of these changes for g-ratio mapping. We found that the new processing pipeline, consecutive HySCO, improved distortion correction. Moreover, the WA compared to the commonly used AM combination method showed a distinct benefit following distortion correction in SPM, and this performed well when benchmarked against FSL, particularly in brain areas with strong susceptibility-related distortions. Finally, we showed that the consecutive HySCO WA performed better than the AM method when combined with other quantitative MRI parameters – MPM – to perform g-ratio mapping. These improvements mean that researchers can conveniently access a wide range of diffusion-related analysis methods within one framework because they are now available within the open-source ACID toolbox as part of SPM, which can be easily combined with other SPM toolboxes, such as the hMRI toolbox, to facilitate computation of myelin biomarkers that are necessary for g-ratio mapping.

The new diffusion MRI distortion correction pipeline in ACID

Recent work has highlighted that while the distortion correction module in SPM-ACID (HySCO2) performed well in correcting distortion in b0 images, there were limitations in its ability to reduce susceptibility-related distortion (Gu and Eklund, 2019), highlighting a need for improvement. To enhance the performance of the HySCO2 module, we performed two consecutive applications of the HySCO2 module instead of the standard single application, with ECMOCO for motion and eddy current correction performed in both the standard and consecutive HySCO pipelines. The second iteration of HySCO2 in the consecutive HySCO pipeline used FA maps computed after the initial distortion correction as input. We found that the consecutive HySCO pipeline improved distortion correction as compared to the standard HySCO pipeline, while noting that it was less efficient than the FSL TOPUP and eddy pipeline.

WA combination – a method to reduce susceptibility-distortion related blurring artefacts

The new consecutive HySCO pipeline was used to test the performance of an alternative phase-encoding combination approach. Areas that are strongly squeezed by susceptibility-related distortions suffer from blurring artefacts when taking the AM of the susceptibility-related distortion corrected blip-up and blip-down images. We proposed the WA combination to reduce this blurring artefact. The WA combination uses the Jacobian of the distortion field (estimated by the HySCO2 module in ACID during susceptibility-related distortion correction) to down-weight areas that are squeezed in, for example, the blip-up image, while up-weighting the same (stretched) regions in the opposite blip-down image. As illustrated in Figure 4, compared to the consecutive HySCO AM, using the consecutive HySCO WA combination resulted in the identification of additional and better spatially defined white matter, in line with the more anatomically accurate MT saturation white matter tissue probability map that was used here as a reference.

The benefits of the WA phase-encoding combination method were also apparent when using quantitative comparisons. Comparing the use of the WA and AM combination methodologies (following distortion, eddy current and motion correction in SPM) identified an increase in the percentage overlap of the diffusion and MT saturation white matter tissue

probability maps when using the WA. This was apparent across the whole white matter map,
but particularly when focusing on regions suffering from high levels of distortion (e.g.
ventromedial pre-frontal cortex, inferior and middle temporal gyri, medial temporal lobe, brain
stem). The WA also compared well with FSL, which was used for benchmarking, and indeed
showed a greater percentage overlap of the diffusion and MT saturation white matter tissue
probability maps for regions suffering from high levels of distortion.

Application to g-ratio mapping

The beneficial effect of using the consecutive HySCO WA combined diffusion data was
illustrated for g-ratio mapping, where blurring effects can reduce the overlap between the
diffusion-MRI-based axonal biomarker and the myelin marker maps. The blurring artefacts in
diffusion MRI (which are decreased when using a WA combination) lead to a worse
delimitation between grey and white matter, which in turn can propagate towards the resulting
g-ratio map (Mohammadi and Callaghan, 2020). As the MR g-ratio is defined only in white
matter (Stikov et al., 2015), it is typically created by combining an axonal water fraction
estimated in the white matter probability maps from diffusion MRI (here, the AWF) and a
white matter myelin biomarker map (here, the MT saturation). In current implementations,
whenever one of the two constituents is not defined, the g-ratio value cannot be calculated,
resulting in undefined areas for the g-ratio map which are set to zero (see Figure 7 in
Mohammadi and Callaghan, 2020). By using the consecutive HySCO WA method to combine
the diffusion data, higher quality data are available to calculate the AWF, enabling the g-ratio
map to cover a larger area of the white matter than when using the consecutive HySCO AM
method (illustrated in our Figure 7).

As these undefined areas vary between individuals, we additionally assessed the
beneficial effects of the WA over the AM combination for g-ratio mapping at the group level.
We predicted that less overlap between the diffusion and MT saturation white matter tissue
probability maps would increase the number of voxels with a g-ratio of zero, decreasing the
local g-ratio values in the group average. Indeed, this was what we found. In regions suffering
from high levels of distortion, the WA-based g-ratio values were closer to the white-matter
average than the AM-based g-ratio values. This prediction was further supported by the
observation that the AM-based g-ratio values were significantly smaller than WA-based g-ratio
values in regions with high susceptibility-related distortions. The improved definition of white
matter in regions suffering from high levels of distortion by the WA technique seems, therefore,
to also result in better representation of g-ratio values in regions with high susceptibility-related
distortions.

Limitations and future directions

We developed and tested our consecutive HySCO WA methodology within the SPM
framework to facilitate the combination of diffusion and quantitative MT saturation maps, the
latter of which has been developed uniquely within SPM (Tabelow et al., 2019). As such, we
aimed to improve the tools available in ACID and SPM. It may be that applying the WA
methodology developed here to other susceptibility-related distortion techniques would also
provide further benefits, however, examining this was beyond the scope of the current study.

We note that, even with the improvements to the initial SPM distortion correction
pipeline, FSL TOPUP and eddy provided better initial distortion, motion and eddy correction
than the SPM equivalents. This is potentially due to the consecutive HySCO pipeline using
multiple interpolation steps, introducing additional blurring effects. Future work will aim to
develop a methodology to combine these interpolation steps, further increasing the

effectiveness of HySCO distortion correction. Of note, however, despite the poorer
performance of HySCO susceptibility-related distortion correction as compared to FSL, the
WA combination in ACID overcame these initial differences in distortion correction between
the two toolboxes. Further development of the distortion correction methods in HySCO may
result in even greater anatomical accuracy in the future.

Conclusions

We present a new WA approach to combining blip-up and blip-down EPI images (after
susceptibility-related distortion correction) that reduces image blurring in diffusion MRI data.
In addition, use of the WA method in combination with other quantitative structural MRI maps
that can be processed within SPM improved g-ratio mapping at the group level in regions with
high susceptibility-related distortions. This WA approach is now available in the open-source
SPM-ACID toolbox together with a new processing pipeline for improved susceptibly-related
distortion correction.

DATA AVAILABILITY

On publication the tools developed as part of this article will be made freely available to
download from (www.diffusiontools.com). The raw diffusion and structural MRI data are part
of a larger project and will be made available once the construction of a dedicated data-sharing
portal has been finalised. In the meantime, requests for the raw data can be sent to
e.maguire@ucl.ac.uk.

ETHICS STATEMENT

This study involving human participants was approved by the University College London
Research Ethics Committee (project ID: 6743/001). All participants gave written informed
consent.

AUTHOR CONTRIBUTIONS

IC: Methodology, Investigation, Formal analysis, Writing - original draft, Writing - review &
editing. **MC:** Formal analysis, Writing - review & editing. **NW:** Resources, Writing - review
& editing. **EM:** Conceptualization, Methodology, Formal analysis, Writing - original draft,
Writing - review & editing, Supervision, Funding acquisition. **SM:** Conceptualization,
Methodology, Formal analysis, Writing - original draft, Writing - review & editing.

FUNDING INFORMATION

IC and EM were supported by a Wellcome Principal Research Fellowship to EM
(101759/Z/13/Z). IC, MC and EM were also supported by a Wellcome Strategic Award to the
Wellcome Centre for Human Neuroimaging (203147/Z/16/Z). SM was supported by the ERA-
NET NEURON (hMRI- ofSCI), the Federal Ministry of Education and Research (BMBF;
01EW1711A and B), and the German Research Foundation (DFG Priority Program 2041
"Computational Connectomics", [MO 2397/5-1; MO 2249/3–1], DFG Emmy Noether Stipend:
MO 2397/4-1), and the Forschungszentrums Medizintechnik Hamburg (fmthh; grant
01fmthh2017). MC is supported by the MRC and Spinal Research Charity through the ERA-

NET Neuron joint call (MR/R000050/1). NW was supported by the European Research
Council under the European Union's Seventh Framework Programme (FP7/2007-2013) / ERC
grant agreement n° 616905; the European Union's Horizon 2020 research and innovation
programme under the grant agreement No 681094; the BMBF (01EW1711A & B) in the
framework of ERA-NET NEURON.

ACKNOWLEDGEMENTS

598 Thanks to Anna Monk, Victoria Hotchin and Gloria Pizzamiglio for assistance with data
599 collection. Thanks also to Lars Rutthotto for helpful discussions.

CONFLICTS OF INTEREST

601 The Wellcome Centre for Human Neuroimaging (London, UK) and the Max Planck Institute
602 for Human Cognitive and Brain Sciences (Leipzig, Germany) have institutional research
603 agreements with Siemens Healthcare. NW was a speaker at an event organized by Siemens
604 Healthcare and was reimbursed for the travel expenses.

REFERENCES

Andersson, J. L. R., Graham, M. S., Zsoldos, E. and Sotiropoulos, S. N. (2016). Incorporating
outlier detection and replacement into a non-parametric framework for movement and
distortion correction of diffusion MR images. *NeuroImage,* 141, 556-572. doi:
10.1016/j.neuroimage.2016.06.058
Andersson, J. L. R., Skare, S. and Ashburner, J. (2003). How to correct susceptibility
distortions in spin-echo echo-planar images: application to diffusion tensor imaging.
NeuroImage, 20(2), 870-888. doi: 10.1016/S1053-8119(03)00336-7
Andersson, J. L. R. and Sotiropoulos, S. N. (2016). An integrated approach to correction for
off-resonance effects and subject movement in diffusion MR imaging. *NeuroImage,*
125, 1063-1078. doi: 10.1016/j.neuroimage.2015.10.019
Ashburner, J. (2007). A fast diffeomorphic image registration algorithm. *NeuroImage,* 38(1),
95-113. doi: 10.1016/j.neuroimage.2007.07.007
Ashburner, J. and Friston, K. J. (2005). Unified segmentation. *NeuroImage,* 26(3), 839-851.
doi: 10.1016/j.neuroimage.2005.02.018
Assaf, Y., Johansen-Berg, H. and Thiebaut de Schotten, M. (2019). The role of diffusion MRI
in neuroscience. *NMR Biomed.,* 32(4), e3762. doi: 10.1002/nbm.3762
Bergui, M., Zhong, J., Bradac, G. B. and Sales, S. (2001). Diffusion-weighted images of
intracranial cyst-like lesions. *Neuroradiology,* 43(10), 824-829. doi:
10.1007/s002340100595
Breman, H., Mulders, J., Fritz, L., Peters, J., Pyles, J., Eck, J., et al. (2020). An image
registration-based method for EPI distortion correction based on opposite phase
encoding (COPE), In: Špiclin Ž., McClelland J., Kybic J., Goksel O. (eds),
International Workshop on Biomedical Image Registration, Cham.doi: 10.1007/978-
3-030-50120-4_12
Callaghan, M. F., Josephs, O., Herbst, M., Zaitsev, M., Todd, N. and Weiskopf, N. (2015).
An evaluation of prospective motion correction (PMC) for high resolution
quantitative MRI. *Front. Neurosci.,* 9(97). doi: 10.3389/fnins.2015.00097
Callaghan, M. F., Lutti, A., Ashburner, J., Balteau, E., Corbin, N., Draganski, B., et al.
(2019). Example dataset for the hMRI toolbox. *Data in Brief,* 25, 104132. doi:
10.1016/j.dib.2019.104132

Cercignani, M., Giulietti, G., Dowell, N. G., Gabel, M., Broad, R., Leigh, P. N., et al. (2017).
637   Characterizing axonal myelination within the healthy population: a tract-by-tract
638   mapping of effects of age and gender on the fiber g-ratio. *Neurobiol. Aging,* 49, 109-
639   118. doi: 10.1016/j.neurobiolaging.2016.09.016
640 Chang, H. and Fitzpatrick, J. M. (1992). A technique for accurate magnetic resonance
641   imaging in the presence of field inhomogeneities. *IEEE Trans. Med. Imaging,* 11(3),
642   319-329. doi: 10.1109/42.158935
643 Chowdhury, R., Lambert, C., Dolan, R. J. and Düzel, E. (2013). Parcellation of the human
644   substantia nigra based on anatomical connectivity to the striatum. *NeuroImage,* 81,
645   191-198. doi: 10.1016/j.neuroimage.2013.05.043
646 Conturo, T. E., Lori, N. F., Cull, T. S., Akbudak, E., Snyder, A. Z., Shimony, J. S., et al.
647   (1999). Tracking neuronal fiber pathways in the living human brain. *Proc. Natl. Acad.*
648   *Sci. USA,* 96(18), 10422. doi: 10.1073/pnas.96.18.10422
649 Corbin, N. and Callaghan, M. F. (2021). Imperfect spoiling in variable flip angle T1 mapping
650   at 7T: quantifying and minimising impact *Magn. Reson. Med.* doi:
651   10.1002/mrm.28720
652 Draganski, B., Ashburner, J., Hutton, C., Kherif, F., Frackowiak, R. S. J., Helms, G., et al.
653   (2011). Regional specificity of MRI contrast parameter changes in normal ageing
654   revealed by voxel-based quantification (VBQ). *NeuroImage,* 55(4), 1423-1434. doi:
655   10.1016/j.neuroimage.2011.01.052
656 Ellerbrock, I. and Mohammadi, S. (2018). Four in vivo g-ratio-weighted imaging methods:
657   Comparability and repeatability at the group level. *Front. Neurosci.,* 39(1), 24-41.
658   doi: 10.1002/hbm.23858
659 Feinberg, D. A., Moeller, S., Smith, S. M., Auerbach, E., Ramanna, S., Glasser, M. F., et al.
660   (2010). Multiplexed echo planar imaging for sub-second whole brain fMRI and fast
661   diffusion imaging. *PLOS ONE,* 5(12), e15710. doi: 10.1371/journal.pone.0015710
662 Glasser, M. F., Sotiropoulos, S. N., Wilson, J. A., Coalson, T. S., Fischl, B., Andersson, J. L.,
663   et al. (2013). The minimal preprocessing pipelines for the Human Connectome
664   Project. *NeuroImage,* 80, 105-124. doi: 10.1016/j.neuroimage.2013.04.127
665 Gu, X. and Eklund, A. (2019). Evaluation of six phase encoding based susceptibility
666   distortion correction methods for diffusion MRI. *Front. Neuroinform.,* 13, 76. doi:
667   10.3389/fninf.2019.00076
668 Helms, G., Dathe, H. and Dechent, P. (2008). Quantitative FLASH MRI at 3T using a
669   rational approximation of the Ernst equation. *Magn. Reson. Med.,* 59(3), 667-672. doi:
670   10.1002/mrm.21542
671 Hodgetts, C. J., Postans, M., Warne, N., Varnava, A., Lawrence, A. D. and Graham, K. S.
672   (2017). Distinct contributions of the fornix and inferior longitudinal fasciculus to
673   episodic and semantic autobiographical memory. *Cortex,* 94, 1-14. doi:
674   10.1016/j.cortex.2017.05.010
675 Holland, D., Kuperman, J. M. and Dale, A. M. (2010). Efficient correction of inhomogeneous
676   static magnetic field-induced distortion in Echo Planar Imaging. *NeuroImage,* 50(1),
677   175-183. doi: 10.1016/j.neuroimage.2009.11.044
678 Hutton, C., Bork, A., Josephs, O., Deichmann, R., Ashburner, J. and Turner, R. (2002).
679   Image distortion correction in fMRI: A quantitative evaluation. *NeuroImage,* 16(1),
680   217-240. doi: 10.1006/nimg.2001.1054
681 Jbabdi, S., Sotiropoulos, S. N., Haber, S. N., Van Essen, D. C. and Behrens, T. E. (2015).
682   Measuring macroscopic brain connections in vivo. *Nat. Neurosci.,* 18(11), 1546-1555.
683   doi: 10.1038/nn.4134

Jezzard, P. and Balaban, R. S. (1995). Correction for geometric distortion in echo planar
images from B0 field variations. *Magn. Reson. Med.,* 34(1), 65-73. doi:
10.1002/mrm.1910340111
Jones, G. W. and Mahadevan, L. (2013). Planar morphometry, shear and optimal quasi-
conformal mappings. *Proc. R. Soc. Ser. A: Math. Phy.,* 469(2153), 20120653. doi:
10.1098/rspa.2012.0653
Kaden, E., Knösche, T. R. and Anwander, A. (2007). Parametric spherical deconvolution:
Inferring anatomical connectivity using diffusion MR imaging. *NeuroImage,* 37(2),
474-488. doi: 10.1016/j.neuroimage.2007.05.012
Kirilina, E., Helbling, S., Morawski, M., Pine, K., Reimann, K., Jankuhn, S., et al. (2020).
Superficial white matter imaging: Contrast mechanisms and whole-brain in vivo
mapping. *Science Advances,* 6(41), eaaz9281. doi: 10.1126/sciadv.aaz9281
Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a
practical primer for t-tests and ANOVAs. *Front. Psychol.,* 4(863). doi:
10.3389/fpsyg.2013.00863
Le Bihan, D., Mangin, J.-F., Poupon, C., Clark, C. A., Pappata, S., Molko, N., et al. (2001).
Diffusion tensor imaging: Concepts and applications. *J. Magn. Reson. Imaging,* 13(4),
534-546. doi: 10.1002/jmri.1076
Lorio, S., Fresard, S., Adaszewski, S., Kherif, F., Chowdhury, R., Frackowiak, R. S., et al.
(2016). New tissue priors for improved automated classification of subcortical brain
structures on MRI. *NeuroImage,* 130, 157-166. doi:
10.1016/j.neuroimage.2016.01.062
Lutti, A., Hutton, C., Finsterbusch, J., Helms, G. and Weiskopf, N. (2010). Optimization and
validation of methods for mapping of the radiofrequency transmit field at 3T. *Magn.*
Reson. Med., 64(1), 229-238. doi: 10.1002/mrm.22421
Lutti, A., Stadler, J., Josephs, O., Windischberger, C., Speck, O., Bernarding, J., et al. (2012).
Robust and fast whole brain mapping of the RF transmit field B1 at 7T. *PLOS ONE,*
7(3), e32379. doi: 10.1371/journal.pone.0032379
Macdonald, J. and Ruthotto, L. (2018). Improved susceptibility artifact correction of echo-
planar MRI using the alternating direction method of multipliers. *Journal of*
Mathematical Imaging and Vision, 60(2), 268-282. doi: 10.1007/s10851-017-0757-x
Mancini, M., Giulietti, G., Dowell, N., Spanò, B., Harrison, N., Bozzali, M., et al. (2018).
Introducing axonal myelination in connectomics: A preliminary analysis of g-ratio
distribution in healthy subjects. *NeuroImage,* 182, 351-359. doi:
10.1016/j.neuroimage.2017.09.018
Mohammadi, S. and Callaghan, M. F. (2018). Image Analysis 1. In M. Cercignani, N. G.
Dowell & P. S. Tofts (Eds.), *Quantitative MRI of the Brain: Principles of Physical*
Measurement (2 ed., pp. 303-324): CRC Press. doi: 10.1201/b21837
Mohammadi, S. and Callaghan, M. F. (2020). Towards in vivo g-ratio mapping using MRI:
unifying myelin and diffusion imaging. *J. Neurosci. Methods*, 108990. doi:
10.1016/j.jneumeth.2020.108990
Mohammadi, S., Carey, D., Dick, F., Diedrichsen, J., Sereno, M. I., Reisert, M., et al.
(2015a). Whole-brain in-vivo measurements of the axonal g-ratio in a group of 37
healthy volunteers. *Front. Neurosci.,* 9(441). doi: 10.3389/fnins.2015.00441
Mohammadi, S., Freund, P., Feiweier, T., Curt, A. and Weiskopf, N. (2013). The impact of
post-processing on spinal cord diffusion tensor imaging. *NeuroImage,* 70, 377-385.
doi: 10.1016/j.neuroimage.2012.12.058
Mohammadi, S., Möller, H. E., Kugel, H., Müller, D. K. and Deppe, M. (2010). Correcting
eddy current and motion effects by affine whole-brain registrations: Evaluation of

three-dimensional distortions and comparison with slicewise correction. *Magn. Reson.*
Med., 64(4), 1047-1056. doi: 10.1002/mrm.22501
Mohammadi, S., Tabelow, K., Ruthotto, L., Feiweier, T., Polzehl, J. and Weiskopf, N.
(2015b). High-resolution diffusion kurtosis imaging at 3T enabled by advanced post-
processing. *Front. Neurosci.,* 8(427). doi: 10.3389/fnins.2014.00427
Movahedian Attar, F., Kirilina, E., Haenelt, D., Pine, K. J., Trampel, R., Edwards, L. J., et al.
(2020). Mapping short association fibers in the early cortical visual processing stream
using in vivo diffusion tractography. *Cerebral Cortex,* 30(8), 4496-4514. doi:
10.1093/cercor/bhaa049
Neumann-Haefelin, T., Wittsack, H.-J., Wenserski, F., Siebler, M., Seitz, R. J., Mödder, U.,
et al. (1999). Diffusion- and perfusion-weighted MRI. *Stroke,* 30(8), 1591-1597. doi:
10.1161/01.STR.30.8.1591
Reber, P. J., Wong, E. C., Buxton, R. B. and Frank, L. R. (1998). Correction of off
resonance-related distortion in echo-planar imaging using EPI-based field maps.
Magn. Reson. Med., 39(2), 328-330. doi: 10.1002/mrm.1910390223
Ruthotto, L., Kugel, H., Olesch, J., Fischer, B., Modersitzki, J., Burger, M., et al. (2012).
Diffeomorphic susceptibility artifact correction of diffusion-weighted magnetic
resonance images. *Phys. Med. Biol.,* 57(18), 5715. doi: 10.1088/0031-
9155/57/18/5715
Ruthotto, L., Mohammadi, S., Heck, C., Modersitzki, J. and Weiskopf, N. (2013).
Hyperelastic susceptibility artifact correction of DTI in SPM *Bildverarbeitung für die*
Medizin 2013 (pp. 344-349): Springer. doi: 10.1007/978-3-642-36480-8_60
Shimony, J. S., McKinstry, R. C., Akbudak, E., Aronovitz, J. A., Snyder, A. Z., Lori, N. F., et
al. (1999). Quantitative diffusion-tensor anisotropy brain MR imaging: Normative
human data and anatomic analysis. *Radiology,* 212(3), 770-784. doi:
10.1148/radiology.212.3.r99au51770
Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-
Berg, H., et al. (2004). Advances in functional and structural MR image analysis and
implementation as FSL. *NeuroImage,* 23, Supplement 1(0), S208-S219. doi:
10.1016/j.neuroimage.2004.07.051
Stikov, N., Campbell, J. S. W., Stroh, T., Lavelée, M., Frey, S., Novek, J., et al. (2015). In
vivo histology of the myelin g-ratio with magnetic resonance imaging. *NeuroImage,*
118, 397-405. doi: 10.1016/j.neuroimage.2015.05.023
Tabelow, K., Balteau, E., Ashburner, J., Callaghan, M. F., Draganski, B., Helms, G., et al.
(2019). hMRI – A toolbox for quantitative MRI in neuroscience and clinical research.
NeuroImage, 194, 191-210. doi: 10.1016/j.neuroimage.2019.01.029
Turner, R. and Le Bihan, D. (1990). Single-shot diffusion imaging at 2.0 tesla. *J. Magn.*
Reson., 86(3), 445-452. doi: 10.1016/0022-2364(90)90023-3
Weiskopf, N., Klose, U., Birbaumer, N. and Mathiak, K. (2005). Single-shot compensation of
image distortions and BOLD contrast optimization using multi-echo EPI for real-time
fMRI. *NeuroImage,* 24(4), 1068-1079. doi: 10.1016/j.neuroimage.2004.10.012
Weiskopf, N., Mohammadi, S., Lutti, A. and Callaghan, M. F. (2015). Advances in MRI-
based computational neuroanatomy: from morphometry to in-vivo histology. *Curr.*
Opin. Neurol., 28(4). doi: 10.1097/WCO.0000000000000222
Weiskopf, N., Suckling, J., Williams, G., Correia, M., Inkster, B., Tait, R., et al. (2013).
Quantitative multi-parameter mapping of R1, PD*, MT, and R2* at 3T: a multi-center
validation. *Front. Neurosci.,* 7(95). doi: 10.3389/fnins.2013.00095
Xu, J., Moeller, S., Auerbach, E. J., Strupp, J., Smith, S. M., Feinberg, D. A., et al. (2013).
Evaluation of slice accelerations using multiband echo planar imaging at 3T.
NeuroImage, 83, 991-1001. doi: 10.1016/j.neuroimage.2013.07.055

Zhang, H., Schneider, T., Wheeler-Kingshott, C. A. and Alexander, D. C. (2012). NODDI:
Practical in vivo neurite orientation dispersion and density imaging of the human
brain. *NeuroImage,* 61(4), 1000-1016. doi: 10.1016/j.neuroimage.2012.03.072

TABLE 1 Regions and peak voxel coordinates with significantly ($p < 0.05$ FWE corrected)
higher g-ratio values within the 20% distortion mask when using the weighted average
combination compared to the arithmetic mean.

Region	Peak voxel coordinates	t
Left middle temporal gyrus	-60 -11 -18	10.81
Right middle temporal gyrus	56 -5 -27	9.13
Corpus callosum	5 32 3	8.97
Right brain stem	15 -20 -30	8.42
Left brain stem	-14 -18 -29	8.34
Right medial orbital gyrus	18 29 -14	7.93
Right frontal pole	12 63 -3	7.80
Left ventromedial frontal cortex	-8 57 -17	7.55
Left inferior temporal gyrus	-53 -44 -20	6.98
Right ventromedial frontal cortex	8 59 -14	6.88
Left frontal pole	-11 60 -2	6.41
Right inferior temporal gyrus	56 -44 -17	6.03
Left ventral diencephalon	-11 -3 -15	5.98

FIGURE 1 Illustration (modified from Jones and Mahadevan, 2013) of the origin of blurring artefacts after combining distortion corrected diffusion data, and how blurring can be avoided. **(A)** Susceptibility-related distortions in two EPI images acquired with reverse phase-encoding directions (i.e. "blip-up" and blip-down" images) can lead to localized squeezing and stretching. Importantly, where one image is squeezed (red box, top) the reverse phase-encoded image (green box, bottom) is stretched. **(B)** After distortion correction and resampling, the two images are typically combined into an arithmetic mean (AM, red-green-dashed box, top). Here, we propose an alternative combination, the weighted average (WA, green box, bottom), where only the stretched boxes are combined, which retains the spatial information. Therefore, spatial information is lost in the AM combination leading to blurring artefacts whereas it is retained in the WA combination.

FIGURE 2 The new consecutive HySCO pipeline. In step 1, as per the standard HySCO pipeline, the HySCO module is applied to the original b0 images, estimating the HySCO field (HySCO$_{b0}$). In step 2, tensor fitting is performed on the distortion corrected blip-up and blip-down data (b0$_{b0}$) to estimate Fractional Anisotropy maps (FA$_{b0FA}$), which are then used in a second round of HySCO for estimating the residual susceptibility related distortions HySCO$_{FA}$ (b0FA). The HySCO field map is the displacement in the phase-encoding direction in mm.

FIGURE 3 The root mean square of the difference in FA map calculated from the blip-up and blip-down data, depicted for each distortion correction methodology. **(A)** Each individual data point. **(B)** The mean and standard deviation of the sample for each distortion correction methodology. *** $p < 0.001$

FIGURE 4 Examples of the improvements in alignment between the white matter tissue
probability maps derived from the diffusion data and MT saturation map when using the WA
phase-encoding combination method compared to the AM. **(A)** In one example participant
(native space) additional white matter is evident in the ventromedial prefrontal cortex when
using the WA. **(B)** A close-up of the white matter shown in (A). **(C)** Additional and better
definition of the white matter in the inferior temporal gyrus of a different example participant
(native space) when using the WA. **(D)** A close-up of the white matter shown in (C). The
structural white matter map is the white matter tissue probability map derived from the MT
saturation map. HySCO = consecutive HySCO, AM = arithmetic mean, WA = weighted
average.

FIGURE 5 Mean (± standard deviation across participants) percentage overlap of the white matter probability maps created from the different phase-encoding combination diffusion data with the MT saturation white matter tissue probability map. The 10%, 20% and 30% distortion graphs focus on regions suffering from greater distortion, determined by thresholding the Jacobian map at distortion values higher than 10%, 20% and 30% respectively. HySCO = consecutive HySCO, AM = arithmetic mean, WA = weighted average. ** $p < 0.01$, *** $p < 0.001$.

FIGURE 6 Examples of regions where greater distortion was observed across the participants.
Images are the average deviation from unity of the Jacobian maps (top row) and average MT
saturation map (bottom row) across participants for the whole sample in Montreal Neurological
Institute (MNI) space. In the deviation from unity of the Jacobian maps, values away from zero
indicate areas suffering from greater distortion (see equation 2). Inter-subject registration was
performed using DARTEL as implemented in SPM12 (Ashburner, 2007). The DARTEL
template and deformations were used to normalize each participant's Jacobians and MT
saturation map to the stereotactic space defined by the MNI template (at 1.5 x 1.5 x 1.5 mm
resolution). VMPFC = ventromedial prefrontal cortex.

FIGURE 7 The impact of reduced blurring artefacts on g-ratio mapping when using the
weighted average (WA) combination compared to the arithmetic mean (AM), illustrated for a
single participant, in their native space. Depicted in the 1st column are the constituents of the
g-ratio: myelin-volume fraction (MVF) based on the MTsat (top), axonal-water fraction (AWF)
using AM (middle), and AWF using WA (bottom). In the 2nd column are the associated white
matter probability maps (c2): MT c2 (top), AM c2 (middle), and WA c2 (bottom). In the 3rd
column is the deviation from unity of the Jacobian map (top, where values away from zero
indicate areas suffering from greater distortion, see equation 2) and the resulting g-ratio maps
AM g-ratio (middle), and WA g-ratio (bottom). The red boxes and crosshairs highlight the
ventromedial prefrontal cortex, and the orange boxes the brain stem, both of which suffer from
high levels of distortion. Greater accuracy was obtained in the white matter probability maps
when using the WA phase-encoding combination method, resulting in a larger area where the
g-ratio was defined.

FIGURE 8 Comparison of g-ratio values when using the weighted average (WA) and arithmetic mean (AM) combinations. **(A)** Mean (± standard deviation) difference in g-ratios between highly distorted regions (defined by the 20% Jacobian distortion mask) and the whole of white matter. The smaller difference evident after the WA combination suggests the g-ratio values were closer to the white matter average, providing a better representation of g-ratio values in regions with high susceptibility-related distortions. **(B)** Voxel-wise comparison within the 20% Jacobian distortion mask showing regions with significantly higher g-ratio values when using the WA in comparison to the AM combination (top row), and their overlap with regions suffering from high levels of distortion. This is observable via the deviation from unity of the Jacobian map (bottom row, where values away from zero indicate areas suffering from greater distortion, see equation 2). Comparison images are thresholded at $p < 0.05$ FWE corrected and displayed on the average MT saturation maps for the whole sample. The left images show the ventromedial prefrontal cortex and brain stem, the right images show the inferior and middle temporal gyri.