# Human cortical areas are sparsely connected: Combining histology with diffusion MRI to estimate the absolute number of axons

**Burke Q. Rosen[1]\*, Eric Halgren[1,2]**

[1]Neurosciences Graduate Program, University of California, San Diego, La Jolla CA 92093
[2]Depts. of Neurosciences & Radiology, University of California, San Diego, La Jolla CA 92093

\*bqrosen@health.ucsd.edu
  3510 Dunhill St.
  San Diego, CA 92101
  858-405-9660

**Short Title**:
Human cortical areas are sparsely connected

**Author Roles**:
E.H. : conceptualization, supervision, funding acquisition, writing – original draft preparation, writing review & editing
B.Q.R. : investigation, formal analysis, data curation, methodology, software, visualization, writing – original draft preparation, writing review & editing

**Competing Interest Statement**:
The authors declare no competing interests.

**Keywords**:
diffusion MRI | structural connectivity | white matter | neuroanatomy | Human Connectome Project

**Preprint Servers**:
A version of this article was deposited in bioRxiv, https://doi.org/10.1101/2021.06.07.447453.

**Abstract** The tracts between cortical areas are conceived as playing a central role in cortical

information processing, but their actual numbers have never been determined in humans. Here we

estimate the absolute number of axons linking cortical areas from a whole-cortex diffusion-MRI (dMRI)

connectome, calibrated using the histologically-measured callosal fiber density. Median connectivity is

estimated as ~6200 axons between cortical areas within-hemisphere and ~1300 axons inter-

hemispherically, with axons connecting functionally-related areas surprisingly sparse. For example, we

estimate that <5% of the axons in the trunk of the arcuate and superior longitudinal fasciculi connect

Wernicke's and Broca's areas. These results suggest that detailed information is transmitted between

cortical areas either via linkage of the dense local connections or via rare, extraordinarily privileged

long-range connections.

## Introduction

The major tracts connecting cortical areas have long been central to models of information-processing

in the human brain [1]. Such models have been refined and applied to development and disease with

the advent of diffusion-MRI or dMRI [2]. However, dMRI only provides *relative* connectivity, not the

absolute number of axons. Relative connectivity is very useful in many circumstances but more

constraints would be possible if the absolute number of axons connecting different cortical areas

could be estimated. Here we describe and apply a novel method for translating from dMRI-derived

streamlines to axon counts.

The ratio between these two measures is obtained by comparing the number of streamlines pass

through the corpus callosum to the number of axons, as measured with histology. The corpus callosum

is uniquely well-suited for this purpose. Electron microscopy (EM) can be used to unambiguously

count the total number of callosal axons because the limits of the callosum are well-defined, axons are

aligned, and sections can be cut perpendicular to their axis. Likewise, dMRI-derived streamlines can be

unambiguously and exhaustively assigned to the callosum because the source and destination of all

axons passing through the callosum as a whole are well-defined. In contrast to the ipsilateral fiber

tracts, in which the majority of axons enter or leave the tract at point between the fascicular terminals,

commissural axons must all be connecting the two hemispheres. Fortunately, despite these

differences, the mean and variability of cross-sectional axon density of the corpus callosum is quite

similar to that of telencephalic white matter in general [3–5], permitting the streamline:axon ratio

calculated from callosal fiber to be applied to intra-hemispheric connections.

It has long been recognized that as the number of cortical neurons increases, maintaining the same

probability of connectivity between neurons would require that axon number increase approximately

with the square of neuron number, and this would require too much volume, impose an unsustainable

metabolic load [6], and actually decrease computational power due to conduction delays [7]. The

consequent imperative to minimize long distance cortico-cortical fibers has been posited to be

reflected in exponential decline in cortical connectivity with distance [8], and to be partially

compensated for with a small-world graph architecture [9], granting special properties to rare long-

distance fibers in a log-normal neural physiology and anatomy [10]. However, this organizing principle

is rarely explicitly addressed in terms of individual axon counts, nor in a manner both granular and

exhaustive with respect to cortical areas. In our histologically calibrated dMRI-derived estimation of this

intercortical axon counts, we find that the widespread cortical integration implied by behavioral and

mental coherence, and routinely observed in widespread physiological synchronization, belies a

surprising small absolute number of long-range axons connecting cortical areas.

**Materials and Methods**

The basic principle of our method is very simple. Given a dMRI-based measure of total inter-

hemispheric connectivity in arbitrary units and the physical number of axons traversing the corpus

callosum, the conversion factor between the two can be obtained by dividing the first by the second.

Specifically, we started with the total inter-hemispheric tractography strength reported in our dMRI

connectome of the Human Connectome Project (HCP) cohort [11]. For each individual, the cross-

sectional area of the corpus callosum was obtained using the standard FreeSurfer structural MRI

pipeline [12]. Multiplying this number by a histologically ascertained callosal fiber density [3] yields an

estimate of the number of axons traversing an individual's corpus callosum. Dividing this count by

individuals' inter-hemispheric dMRI streamline value yields the conversion ratio from the arbitrarily-

scaled dMRI metric to the absolute number of axons. Note that this procedure is independent from the

scale of the dMRI metric, requiring only that it be proportional to the absolute number of fibers. A

moderate proportionality has been observed in a comparison of dMRI and retrograde tracing in

macaques [13,14]. Therefore, while the ratio itself is study-specific, the procedure can be applied to

any dMRI tractography method or parameter set, provided that the dMRI method returns a continuous

distribution of connectivity values and has reasonably similar sensitivity to callosal and ipsilateral fiber

tracts. To demonstrate this, we repeated the analysis using data from an alternate tractography of the

HCP cohort [15]. For detailed methodology, see S1 Appendix: Extended Methods and Results.

**Results**

Our previous dMRI study [11] includes estimated tractography streamlines between all 360 parcels of

the HCP-MMP1.0 atlas [16] for each of the 1,065 individuals in the HCP dMRI cohort. The sum

connectivity between the 180 left hemisphere cortical parcels and the 180 right ($180^2$ parcel pairs)

constitutes the total callosal connectivity, on average $1.25 \times 10^8$ streamlines. Based on our assumed

fiber density of $3.7 \times 10^5$ axons/mm$^2$ [3] and measured callosal cross-sectional area (mean =

689.45mm$^2$), we estimate this cohort to have $2.6 \times 10^8$ callosal fibers on average. The mean quotient

between these two quantities is 2.00 axons-per-streamline, a ratio specific to the dMRI methodology

and parameters used.

Applying the conversion factor to the inter-parcel connectivity from our prior study yields an

estimate of the absolute number of axons connecting different cortical areas (Fig 1A), $2.43 \times 10^9$ axons

in total. This implies that <22% of the ~$11.5 \times 10^9$ cortical pyramidal cells [17,18] project outside their

parcel with the remainder being short-range horizontal or U-fibers. Furthermore, because 51% of inter-

areal axons are to adjacent areas, <11% of pyramidal cells project beyond the adjacent parcel. Axons

are approximately log-normally distributed among non-adjacent parcel pairs, with adjacent pairs

having disproportionally high axon counts (Fig 1B). Median connectivity is ~6,200 axons between

cortical areas in the same hemisphere, and ~1,300 inter-hemispherically. The number of axons in the

median *inter*-hemispheric connection is ~20% of that in the median *intra*-hemispheric connection,

similar to that found with histological tracing in macaques [19]. The number of callosal axons is not

significantly affected by subject sex, though subjects ages 26-30 have slightly more inter-hemispheric

axons than subjects ages 22-15 (S3 Fig). This sparse long-range connectivity is consistent with its

exponential fall-off measured with dMRI in humans and histologically in other mammals [8,11] and with

previous statistical estimates based on traditional neuroanatomy [20]. In comparison to the number of

axons that would be necessary for complete interconnectivity of neurons in different cortical areas with

each other, the estimated number is ~$10^{11}$ times smaller (see S1 Appendix for calculation).

Our method requires that dMRI-tractography performs in a roughly similar manner when applied to

intra- versus inter-hemispheric connections. We evaluated this similarity by comparing linear

regressions of distance-matched log-transformed streamlines, and found no difference between the

slope or intercept of inter- and intra-hemispheric connections (Fig 1C); within the common distance

domain a paired r-test showed no difference in correlation, $p = 0.58$. The total count of a parcel's

axons to or from all other areas is much less variable when normalized by the parcel's area (Fig 1D,E).

Multiplying the total number of fibers by the effective fiber cross-sectional area (the inverse callosal

packing density, $2.7 \times 10^{-6}$ mm$^2$/axon) yields $6.6 \times 10^3$ mm$^2$ of cortex or 3.7% of the total white-gray

interface. Note that the effective axonal area includes myelin, supporting cells and intercellular space

in addition to the axon proper. The cross-sectional packing density of human prefrontal white matter is

quite similar to callosal values with an average of $3.5 \times 10^5$ myelinated axons/mm$^2$ [4] after correction for

tissue shrinkage, and this varies among prefrontal regions by a less than factor of 2 [5]. The percent of

fiber-allocated cortical area is quite similar to the ~4% of total cortical fibers Schüz and Braitenberg

estimated [20] are contained in the corpus callosum and long fascicles. The remaining ~95% of the

cortical gray-white interface area is likely occupied by the short range U-fiber system which is difficult

to assess with dMRI.

We estimated the fraction of axons traversing the entire length of the arcuate / superior lateral

fasciculus (AF/SLF) between termination fields centered on Broca's and Wernicke's areas. The total

number of AF/SLF axons was derived by multiplying the tracts' cross-sectional areas by mean axon

density of ipsilateral tracts. Using published estimates of the cross-sectional areas of 160.6 mm$^2$ and

51.5 mm$^2$ for the left and right AF and 213.8 mm$^2$ and 174.4 for the left and right SLF [21], and a

shrinkage-corrected axon density of $3.5 \times 10^5$ axons/mm$^2$ [4], yields a total of $1.3 \times 10^8$ and $0.8 \times 10^8$ axons

in the left and right AF/SLF. These values were compared to the number of tractography-derived axons

connecting AF/SLF termination fields [16] according to consensus definitions from reference [22], see

S4 Fig. When the anterior termination field, centered on Broca's area is defined as HCP-MMP1.0

parcels 44, 45, 6r, IFSa, IFSp, and FOP4 and the posterior termination field, centered on Wernicke's

area as parcels PSL, RI, STV, and PFcm, trans-terminal axons account for only 0.6% and 0.8% of tract

axons in the left and right hemispheres. If the termination fields are liberally expanded to also include

parcels 47l and p47r in Broca's and parcels PF, PFm, and PGi in Wernicke's then these percentages

increase to a still modest 1.9% and 2.9% of total tract axons. Please note that in this calculation all

axons between Broca's and Wernicke's areas are assumed to pass through the AF/SLF; to the degree

that some pass outside the AF/SLF, our estimates should be decreased.

The volume of hemispheric white matter (WM) occupied by axons between cortical parcels is equal

to the sum of all axons' lengths multiplied by their cross-sectional areas. The mean fiber tract lengths of

connections were taken from our prior dMRI analysis [11]. For the effective cross-sectional area of

axons we again assumed an effective cross-sectional area equivalent to the inverse of the callosal fiber

density [3]. So calculated, the volume occupied by cortico-cortical and hippocampo-cortical fibers is

$4.3 \times 10^5$ mm$^3$, or about 96% of the total MRI-assessed WM volume. This implies that the number of

long-range fibers cannot be larger than our estimate unless the axon density calculated from the

corpus callosum histology is mistakenly low, but this is a linear effect that would need to be

unrealistically inaccurate for it to change our main conclusions. Another possibility is that axon density

was correctly measured for the corpus callosum but is higher for intra-hemispheric fibers *and* the

number of axons per streamline is higher intra- than inter-hemispherically. These both need to be the

case because there is no space available in the hemispheric white matter to contain more axons unless

they are smaller. However, the histological data indicates that axonal density is approximately the

same intra- and inter-hemispherically [3,4] and the comparison of distance-matched streamlines (Fig.

1*C*). suggests that the number of axons per streamline intra- and inter-hemispherically is also

approximately equal. The remaining WM volume may seem insufficient for connections with

subcortical structures. However, the major subcortical structures known to communicate with the

cortex (thalamus, amygdala, striatum, nucleus basalis, locus coeruleus, etc.) contain, in sum, less than

1% of the number of cortical neurons [23–25], see S1 Appendix: Extended Methods and Results. Even

under the unlikely assumption that all excitatory thalamic neurons project to the cortex with a density

across all cortical parcels proportionally to their area, their axons would only comprise ~4% of the total

WM volume.

We conducted a series of simulations examining the consequences of possible errors in axonal

packing density and axon-to-streamline ratio. As noted above, these parameters are co-constrained by

the physical volume available for white matter. At our derived ratio and assumed packing density, the

volume of inter-areal axons is just under the observed total white matter volume. Large net errors in

the assumed packing density would result in total inter-areal axon volumes that are inconsistent with

the hemispheric white matter volume, a value that is well established. Consequently, in our

simulations, axonal packing density and axon-to-streamline ratio were reciprocally changed in order to

maintain a constant total cerebral white matter volume. Since our parameter estimates are based

primarily on fibers passing through the callosum, which are longer in general than intra-hemispheric

fibers, parameter inaccuracies are more likely for short fibers. In order to evaluate the effect of such

inaccuracies, we first simulated the effect of assuming that dMRI sensitivity is systematically reduced

(and thus axon count underestimated) for shorter fibers. Uniformly doubling the number of axons with

lengths < 40 mm in this way yields only a 37.5% increase in the total number of inter-areal axons, and

because rank-order of observations is not altered the median axon counts are unchanged.

Conceivably, our parameter estimates are accurate for the longest fibers (which tend to pass through

the callosum) but are progressively less accurate for shorter inter-areal distances. Consequently, we

simulated assuming that our packing density and dMRI sensitivity estimates at the longest fiber lengths

were as in our base model, and then increased the dMRI streamline-to-axon ratio linearly as the fibers

got shorter. Axon density was adjusted to maintain white matter volume constant. Even with

unrealistically large systematic and matched errors in the two estimates, median inter-areal axon count

are only modestly increased, see S5 Fig. For example, adjustments resulting in twice the axon density

and half the sensitivity of dMRI to axons at short inter-parcel distances increase the median number of

inter-parcel axons by only ~36%. As reviewed below, errors in inter-areal axon count due to miss-

estimation of axonal packing density are likely to be relatively unbiased with respect to inter-areal fiber

length, and reported measures vary less than a factor of two. Our simulations indicate that such errors

would have only marginal effects on the median number of inter-areal axons.

In order to demonstrate that the key principles and findings of this report are robust to the details

of tractography procedure, we repeated our analysis on the tractography data of Arnatkeviciute and

colleagues [15]. These data consist of a 972 subject subset of the HCP cohort and use the same

parcellation but a different method to reconstruct the fiber streamlines. These data contain fewer

streamlines and the estimated numbers of axons per connection are broadly comparable, being

somewhat fewer but within an order of magnitude of our primary estimate. We find $8.4 \times 10^8$ total axons

and $1.1 \times 10^8$ callosal axons per subject on average, with medians of ~1100 and ~130 axons for pairwise

intra- and inter-hemispheric connections between cortical areas (S2 Fig), i.e., even more sparse than

those calculated from our tractography data. These data confirm that, independent of tractography

procedure, cortical areas are sparsely connected.

**Discussion**

In this study we estimated the absolute number of axons interconnecting cortical areas by calibrating

dMRI-based tractography using the histologically ascertained cross-sectional fiber density of the

corpus callosum and found that long-range cortico-cortical connections are quite sparse. Our method

depends on histological estimates of callosal axon packing density and leverages the unique

properties of the callosum. It assumes proportionality between the number of axons connecting two

areas and the number of dMRI tractography streamlines for the areal pair, and it assumes approximate

parity between dMRI sensitivity to inter- and intra-hemispheric connections. In order to estimate the

volume occupied by inter-parcel cortico-cortical connections, we make the further assumption that the

effective axonal packing density is reasonably uniform across the hemispheric white matter.

Although there are few reports of axonal density in the corpus callosum [3,26], they are consistent

with each other and with reports of axonal density of intra-hemispheric tracts [4]. Our study relies on

the data reported by Aboitiz et al. [3], because it is the most systematic count of which we are aware.

There is a later study, [26], which provides a value only slightly lower, $2.83 \times 10^5$ vs $3.7 \times 10^5$ axons/mm$^2$

after correction for tissue shrinkage [27]. However, since this later study was primarily a survey of axon

diameter with packing density only incidentally recorded, it is conceivable that density values derived

from could be small underestimates. Mis-estimated tissue shrinkage is possible but likely to be <10%

and callosal areas were measured with a well-validated and widely-used in vivo method [12]. The axon

count estimate does not require that the packing densities of the corpus callosum and ipsilateral white

matter be the same, but rather the lesser assumption that the axon to dMRI streamline ratio be

reasonably uniform across the various instances of long distance cortico-cortical connectivity.

However, if the packing densities of ipsilateral and callosal long distance connections are also similar

then the total hemispheric white matter volume provides an absolute constraint on the number of long

distance connections, and this provides a powerful validation of our estimate.

The literature provides converging evidence that the fiber packing density of white matter varies by

at most a factor of 2 across the cortex, including the corpus callosum. A histological study of human

cortex found only a 7% difference in the axon densities of the callosum versus the superior longitudinal

and inferior occipitofrontal fasciculi [26]. Zikopolous and Barbas [4] found that the cross-sectional

packing density of human prefrontal white matter is remarkably similar to callosal values with an

average of 3.5x10$^5$ myelinated axons/mm$^2$ after correction for tissue shrinkage and that this varies

among prefrontal regions by a less than factor of 2 [5]. In addition, while dMRI-based estimates of axon

density and caliber are imperfect, they suggest that axon density varies by less than 2-fold both

between major ipsilateral tracts and within each tract along their length [28]. While packing density is

not directly equivalent to axon caliber, the two are likely inversely related. Axon diameter, as estimated

with dMRI, varies by 20% among ipsilateral tracts and by at most a factor of 2.2 among white matter

voxels, including those of the corpus callosum [29,30]. Overall, distributions of dMRI-derived axon

diameter for the callosum [31] and whole cerebrum [30] are similar with the bulk of values between 2.5

and 5 μm. Histological measurements in macaques concur that axon diameters are very similar within

the callosal and non-callosal segments of major fasciculi and vary by less than 2-fold across the cortex

[32]. Furthermore, even this limited variation in diameter is not systematically dependent on the length

of axons but rather on their regions of origin and termination [33].

Concerning dMRI- to-axon-count proportionality, it has been shown in macaques that there is a

moderate linear correlation between dMRI-traced streamlines and the number of fibers identified with

histological tracers connecting cortical regions [13,14]. Based on these data, Donahue and colleagues

[13] concluded that dMRI tractography was capable of quantitively describing cortico-cortical white

matter tracts with approximately order-of-magnitude precision using a high-quality dataset such as the

HCP. In line with previous statistical estimates of whole-cortex axon counts based on traditional

neuroanatomy [20], the numbers of axons in pairwise connections are probably correct within an order

of magnitude. Estimates of this precision are useful as we find that that inter-areal connectivity derived

from dMRI varies over more than seven orders of magnitude.

While we derive a single dMRI-to-axon count factor, it is likely that the true conversion ratio varies

somewhat among connections due to local microstructural differences other than axon count such as

axon caliber, packing density, or myelination. However, the major fasciculi (including the corpus

callosum) have similar axon calibers and packing densities, varying by only a factor of ~2 across the

cortex when examined in humans histologically [4,5,26] or with dMRI [28–30] and histologically in

macaques [32]. More generally, packing density is a function of local cellular interactions, especially

with oligodendrocytes, and because these are the same in the corpus callosum and intra-hemispheric

white matter, the *a priori* expectation is that packing density would be similar, as histological data

supports. Myelination has only a modulatory effect on dMRI-detected anisotropy, with most of the

effect derived from axonal membranes [34]. Nevertheless, microstructural variation in dMRI-to-axon

count ratio may be a source of noise in our estimates. Our simulations of miss-estimation of dMRI-to-

axon ratio and packing density show that these errors only marginally affect estimated axon counts and

do not alter our conclusions. Our findings were also similar when we repeated our analysis on an

alternative, somewhat sparser dMRI tractography dataset [15], demonstrating that the details of the

tractography algorithm do not affect the overall tenor of our results. While the scope of this study is

limited to long-range fibers, we note that shorter, more superficial U-fibers are systematically less

myelinated and of lesser caliber than their inter-areal counterparts [5] and therefore the procedure

outlined here may require modification before being applied to them.

As previously stated, this methodology assumes a reasonable degree of parity in the sensitivity of

the dMRI tractography to intra- and inter-hemispheric fiber tracts. Consistent with this assumption, we

found little difference between distance-matched dMRI connectivity for callosal versus ipsilateral

connections. If callosal axons were more easily detectable, this would be reflected as an upward

displacement of the inter-hemispheric trace (blue) above the intra-hemispheric trace (green) across the

entire distance domain in Fig 1C, which is not evident. This parity is perhaps unsurprising because

over most of their trajectories, inter-hemispheric fibers are subjected to the same crossing fiber issues

as intra-hemispheric. Specifically, the corpus callosum is a distinct tract for only about 15-35 mm, but its fibers range in length up to about 300 mm. Thus, the fraction of an inter-hemispheric tract that resides within the callosum is inversely proportional to its total length. Consequently, if there were enhanced detection of callosal fibers as streamlines by dMRI, one would expect the inter-hemispheric (blue) trace in Fig 1C to be elevated primarily at short fiber lengths and depressed at long fiber lengths, resulting in a noticeably steeper slope for the inter-hemispheric than the intra-hemispheric trace, which is not observed. These data support the applicability of the scaling factor derived from inter-hemispheric fibers to intra-hemispheric fibers.

The corpus callosum was used to calibrate the estimate because of its unique properties: it has a well-defined cross-sectional area, more than ~99% of inter-hemispheric cortico-cortical axons are routed through it [3,35], and essentially no fibers leave or enter the tract between the two hemispheres. This is in sharp contrast to non-commissural fasciculi. While it may be commonly assumed that within major cortical fasciculi the majority of axons terminate or originate at the ends of the tract and thus carry information along its entire length, an alternative conception is that these large bundles are composed mostly of axons shorter than the total fascicular length which enter and exit the tract at various points. By analogy, the former assumption likens a tract to a tunnel, where all traffic is trans-terminal, whereas the latter conceives of tracts as like interstate highways, where very few vehicles travel the entire route. In a supplementary analysis we compared these models for the arcuate / superior lateral fascicular (AF/SLF) system. The total number of AF/SLF axons was estimated using the tract diameters [21] and packing densities [4] taken from the literature. The number of trans-terminal axons was determined by defining termination field parcels centered on restricted and inclusive definitions of Broca's and Wernicke's areas [22]. Depending on the assumptions, only about 1-5% of the axons in a middle section of these fasciculi are trans-terminal. These percentages may be over-

estimates since they assume that all fibers between the posterior and anterior areas travel through the

AF/SLF. However, even if these values are a 2-fold underestimate, it suggests that only a small fraction

of the axons in the ipsilateral fasciculi are trans-terminal. The evidence indicates that the 'highway'

model is more apt for the ipsilateral fiber tracts and this conception is consistent with neural wiring

being driven, in large part, by exponential distance rules [8]. This of course, does not apply to the

corpus callosum, as there is no inter-terminal cortex to project into.

Importantly, the reconceptualization of major intra-hemispheric tracts as containing few fibers

connecting their distant terminals is still consistent with the long- and well-established impression from

blunt dissection [36] and dMRI orientation maps [37] that the hemispheric white matter is largely

composed of well-defined long-distance tracts. Indeed, we estimate that ~96% of the white matter is

composed of inter-areal axons. What these observations suggest is that the major tracts arise from the

tendency of axons to grow in alignment with axons which are already present using established

mechanisms of adhesion and fasciculation [38]. Thus axons are free to join tracts at various points, and

tend to proceed together (fasciculate), but again are free to leave whenever they approach their own

target. In other words, axons are joined in a given tract because they share a direction rather than an

origin and destination.

The inter-areal axon counts we derive here permit other interesting quantitative estimates which

may inform models of cortical neurophysiology. For example, the connections between Wernicke's

and Broca's areas are thought to integrate receptive and expressive aspects of language, but we

estimate that there are only ~58,000 axons between the core cortical parcels in these regions (parcels

44 and PSL), fewer than two for each word in an average university student's vocabulary [39]. Another

example where quantitative appreciation of direct axonal connections may influence neuro-cognitive

models are the hippocampo-cortical interactions subserving recent memory, which are commonly

posited to carry information regarding the contents of the memory trace during memory formation,

consolidation and retrieval. We estimated that areas distant from the hippocampus, notably the

dorsolateral prefrontal cortex, may be connected to it by <10 axons/mm$^2$ (Fig 1F) including both

efferent and afferent axons, yet hippocampo-prefrontal interactions are considered crucial for

contextual recall [40]. We estimated the average neural density in the cortex as ~92,300 neurons/mm$^2$

by dividing the $16.34 \times 10^9$ cortical neurons (including interneurons) from [17] by the $1.77 \times 10^5$ mm$^2$

mean white–gray surface area of the HCP cohort used. If the sparse connectivity suggested by our

calculations is correct, it implies that hippocampo-dorsolateral prefrontal interactions in memory are

likely mediated by polysynaptic pathways.

These constraints encourage consideration of models of cortical function where connections are

dense but mainly local, i.e. a small world network with intense interconnections within modules and

sparse projections between them. While this general principle is widely accepted, the scale of the vast

gulf in absolute connectivity between local and long-range connections is startling. This network

architecture provides for the wide and efficient distribution of information created by local processing

within modules [8,9]. A more uniformly connected cortex would require more white matter,

necessitating a more voluminous cerebrum and the human cortex is near the limit after which an

increase in size reduces computational power [6,7]. Functionally, a deep reservoir of weak connections

enables a large number of states and eases state transitions [10]. Modeling suggests that long range

covariance and even synchrony can be achieved through activation of multi-synaptic pathways rather

than direct connections [41,42], and possible signs of these have been observed experimentally in

humans [43,44].

While the long-range direct cortico-cortical axons are few in number, we note that axons are

heterogeneous and that these counts are a limited proxy for true inter-areal connectivity. Axons,

especially those connecting architectonically similar regions, may have a disproportionate impact on

the flow of information despite their rarity [45–47], by virtue of their morphology (e.g. greater

diameter, larger termination fields, greater axonal arborization, or more numerous *en passant*

varicosities) or by molecular synaptic specializations. For example, while only ~5% of the synapses to

V1 layer 4 come from the lateral geniculate body in the macaque [48], they have an outsized effect on

their firing [49]. The importance of rare inter-module connections might also be enhanced if they are

focused on a small location within cortical parcels (i.e. the rich club [50]), but this has not been

convincingly demonstrated. Lastly, it is useful to note that these quantitative considerations are

radically different in other species, where the smaller number of cortical neurons and shorter inter-

areal distances allow greater connectivity between cortical areas, as well as a larger proportion of

subcortical connectivity [7].

**Fig 1.** The number of axons estimated to interconnect the 360 cortical parcels of the HCP-MMP1.0

atlas. **(A)** Connectivity matrix of intra-hemispheric axon counts, averaged across the two hemispheres.

Parcels are ordered into 10 functional networks. **(B)** Histograms showing the distribution of inter- and

intra-hemispheric pairwise axon counts. Physically adjacent and non-adjacent parcel pair intra-

hemispheric histograms are stacked. Median connectivity, indicated, is ~6,200 axons between cortical

areas in the same hemisphere, and ~1,300 inter-hemispherically. **(C)** Comparison of intra- and inter-

hemispheric dMRI connectivity as a function fiber tract distance. Pairwise values averaged within 15

fiber-length bins. Shading shows bootstrapped 95% confidence intervals. **(D-F)** axon counts and

densities averaged across the two hemispheres and visualized on the left fsaverage template cortex

[12]. **(D)** total axons connecting each parcel to all others. **(E)** axons connecting each parcel to all

others, normalized by the reference parcel's area. **(F)** axons connecting the hippocampus to the rest of

the cortex, normalized by the area of the cortical parcel, shown in log scale.

## Acknowledgements

This work was supported by National Institutes of Health grants RF1MH117155 and R01NS109553 (EH). Data were
provided, in part, by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van
Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 National Institutes of Health (NIH) Institutes and
Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems
Neuroscience at Washington University. The funders had no role in study design, data collection and analysis,
decision to publish, or preparation of the manuscript.

## Data availability

410  The processed data used in this study are available at  https://doi.org/10.5281/zenodo.5912108. Datafiles are in
411  Matlab v7.3 format. The Human Connectome Project's raw imaging data and FreeSurfer outputs may be
412  downloaded from https://db.humanconnectome.org.

## Code availability

415  The main standalone analysis Matlab code used in this study is available at
416  https://doi.org/10.5281/zenodo.5912108. Additional non-standalone supporting code will be made available
417  upon request.

## References

1.     Geschwind N. Disconnexion syndromes in animals and man. Brain. 1965;88: 237–294.
2.     Assaf Y, Johansen-Berg H, Thiebaut de Schotten M. The role of diffusion MRI in neuroscience. NMR
Biomed. 2019;32: e3762. doi:https://doi.org/10.1002/nbm.3762
3.     Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus callosum. Brain Res.
1992;598: 143–153. doi:10.1016/0006-8993(92)90178-C
4.     Zikopoulos B, Barbas H. Changes in prefrontal axons may disrupt the network in autism. J Neurosci.
2010;30: 14595–14609.
5.     Zikopoulos B, García-Cabezas MÁ, Barbas H. Parallel trends in cortical gray and white matter architecture
and connections in primates allow fine study of pathways in humans and reveal network disruptions in
autism. PLoS Biol. 2018;16: e2004559.
6.     Bullmore E, Sporns O. The economy of brain network organization. Nat Rev Neurosci. 2012;13: 336–349.
doi:10.1038/nrn3214
7.     Hofman MA. Evolution of the human brain: when bigger is better. Front Neuroanat. 2014;8: 15.
8.     Markov NT, Ercsey-Ravasz M, Van Essen DC, Knoblauch K, Toroczkai Z, Kennedy H. Cortical high-density
counterstream architectures. Science. 2013. pp. 1238406–1238406. doi:10.1126/science.1238406
9.     Bassett DS, Bullmore ET. Small-world brain networks revisited. Neurosci. 2017;23: 499–516.
10.    Buzsáki G, Mizuseki K. The log-dynamic brain: how skewed distributions affect network operations. Nat
Rev Neurosci. 2014;15: 264–278. doi:10.1038/nrn3687

11.    Rosen BQ, Halgren E. A Whole-Cortex Probabilistic Diffusion Tractography Connectome. eNeuro. 2021;8:
439           ENEURO.0416-20.2020. doi:10.1523/ENEURO.0416-20.2020
440    12.    Fischl B. FreeSurfer. Neuroimage. 2012;62: 774–781. doi:10.1016/j.neuroimage.2012.01.021
441    13.    Donahue CJ, Sotiropoulos SN, Jbabdi S, Hernandez-Fernandez M, Behrens TE, Dyrby TB, et al. Using
442           Diffusion Tractography to Predict Cortical Connection Strength and Distance: A Quantitative Comparison
443           with Tracers in the Monkey. J Neurosci. 2016;36: 6758–6770. doi:10.1523/JNEUROSCI.0493-16.2016
444    14.    van den Heuvel MP, de Reus MA, Feldman Barrett L, Scholtens LH, Coopmans FMT, Schmidt R, et al.
445           Comparison of diffusion tractography and tract-tracing measures of connectivity strength in rhesus
446           macaque connectome. Hum Brain Mapp. 2015;36: 3064–3075. doi:10.1002/hbm.22828
447    15.    Arnatkeviciute A, Fulcher BD, Oldham S, Tiego J, Paquola C, Gerring Z, et al. Genetic influences on hub
448           connectivity of the human connectome. Nat Commun. 2021;12: 4237. doi:10.1038/s41467-021-24306-2
449    16.    Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, et al. A multi-modal parcellation of
450           human cerebral cortex. Nature. 2016;536: 171–8. doi:10.1038/nature18933
451    17.    Azevedo FAC, Carvalho LRB, Grinberg LT, Farfel JM, Ferretti REL, Leite REP, et al. Equal numbers of
452           neuronal and nonneuronal cells make the human brain an isometrically scaled-up primate brain. J Comp
453           Neurol. 2009;513: 532–541. doi:10.1002/cne.21974
454    18.    DeFelipe J, Fariñas I. The pyramidal neuron of the cerebral cortex: Morphological and chemical
455           characteristics of the synaptic inputs. Prog Neurobiol. 1992;39: 563–607. doi:10.1016/0301-
456           0082(92)90015-7
457    19.    Barbas H, Hilgetag CC, Saha S, Dermon CR, Suski JL. Parallel organization of contralateral and ipsilateral
458           prefrontal cortical projections in the rhesus monkey. BMC Neurosci. 2005;6: 32. doi:10.1186/1471-2202-
459           6-32
460    20.    Schüz A, Braitenberg V. The human cortical white matter: quantitative aspects of cortico-cortical long-
461           range connectivity. Cortical areas Unity Divers. 2002; 377–385.
462    21.    Yeh F-C. Shape analysis of the human association pathways. Neuroimage. 2020;223: 117329.
463           doi:https://doi.org/10.1016/j.neuroimage.2020.117329
464    22.    Tremblay P, Dick AS. Broca and Wernicke are dead, or moving past the classic model of language
465           neurobiology. Brain Lang. 2016;162: 60–71. doi:https://doi.org/10.1016/j.bandl.2016.08.004
466    23.    Xuereb JH, Perry RH, Candy JM, Perry EK, Bonham JR. Nerve cell Loss in the thalamus in Alzheimer's
467           disease and Parkinson's disease. Brain. 1991;114: 1363–1379.
468    24.    Arcelli P, Frassoni C, Regondi MC, De Biasi S, Spreafico R. GABAergic neurons in mammalian thalamus: a
469           marker of thalamic complexity? Brain Res Bull. 1997;42: 27–37.
470    25.    Avino TA, Barger N, Vargas M V, Carlson EL, Amaral DG, Bauman MD, et al. Neuron numbers increase in
471           the human amygdala from birth to adulthood, but not in autism. Proc Natl Acad Sci. 2018;115: 3710–3715.
472    26.    Liewald D, Miller R, Logothetis N, Wagner HJ, Schüz A. Distribution of axon diameters in cortical white
473           matter: an electron-microscopic study on three human brains and a macaque. Biol Cybern. 2014;108:
474           541–557. doi:10.1007/s00422-014-0626-2
475    27.    Schüz A, Palm G. Density of neurons and synapses in the cerebral cortex of the mouse. J Comp Neurol.
476           1989;286: 442–455.
477    28.    De Santis S, Drakesmith M, Bells S, Assaf Y, Jones DK. Why diffusion tensor MRI does well only some of the
478           time: variance and covariance of white matter tissue microstructure attributes in the living human brain.
479           Neuroimage. 2014;89: 35-44.
480    29.    Huang SY, Tian Q, Fan Q, Witzel T, Wichtmann B, McNab JA, et al. High-gradient diffusion MRI reveals
481           distinct estimates of axon diameter index within different white matter tracts in the in vivo human brain.
482           Brain Struct Funct. 2020;225: 1277–1291. doi:10.1007/s00429-019-01961-2
483    30.    Fan Q, Nummenmaa A, Witzel T, Ohringer N, Tian Q, Setsompop K, et al. Axon diameter index estimation
484           independent of fiber orientation distribution using high-gradient diffusion MRI. Neuroimage. 2020;222:
485           117197.
486    31.    Horowitz A, Barazany D, Tavor I, Bernstein M, Yovel G, Assaf Y. In vivo correlation between axon diameter
487           and conduction velocity in the human brain. Brain Struct Funct. 2015;220: 1777–1788.
488    32.    Tomasi S, Caminiti R, Innocenti GM. Areal differences in diameter and length of corticofugal projections.
489           Cereb Cortex. 2012;22: 1463–1472.

33.   Innocenti GM, Vercelli A, Caminiti R. The Diameter of Cortical Axons Depends Both on the Area of Origin
and Target. Cereb Cortex. 2014;24: 2178–2188. doi:10.1093/cercor/bht070
34.   Beaulieu C. The basis of anisotropic water diffusion in the nervous system – a technical review. NMR
Biomed. 2002;15: 435–455. doi:https://doi.org/10.1002/nbm.782
35.   Highley JR, Esiri MM, McDonald B, Roberts HC, Walker MA, Crow TJ. The size and fiber composition of the
anterior commissure with respect to gender and schizophrenia. Biol Psychiatry. 1999;45: 1120–1127.
36.   Schmahmann JD, Pandya DN. Cerebral White Matter – Historical Evolution of Facts and Notions
Concerning the Organization of the Fiber Pathways of the Brain. J Hist Neurosci. 2007;16: 237–267.
doi:10.1080/09647040500495896
37.   Hagler DJ, Ahmadi ME, Kuperman J, Holland D, McDonald CR, Halgren E, et al. Automated white-matter
tractography using a probabilistic diffusion tensor atlas: Application to temporal lobe epilepsy. Hum Brain
Mapp. 2009;30: 1535–1547. doi:10.1002/hbm.20619
38.   Spead O, Poulain FE. Trans-axonal signaling in neural circuit wiring. Int J Mol Sci. 2020;21: 5170.
39.   Brysbaert M, Stevens M, Mandera P, Keuleers E. How Many Words Do We Know? Practical Estimates of
Vocabulary Size Dependent on Word Definition, the Degree of Language Input and the Participant's Age
. Frontiers in Psychology . 2016. p. 1116. Available:
https://www.frontiersin.org/article/10.3389/fpsyg.2016.01116
40.   Jin J, Maren S. Prefrontal-Hippocampal Interactions in Memory and Emotion   . Frontiers in Systems
Neuroscience . 2015. Available: https://www.frontiersin.org/article/10.3389/fnsys.2015.00170
41.   Bazhenov M, Rulkov NF, Timofeev I. Effect of synaptic connectivity on long-range synchronization of fast
cortical oscillations. J Neurophysiol. 2008;100: 1562–1575. doi:10.1152/jn.90613.2008
42.   Vicente R, Gollo LL, Mirasso CR, Fischer I, Pipa G. Dynamical relaying can yield zero time lag neuronal
synchrony despite long conduction delays. Proc Natl Acad Sci. 2008;105: 17157 LP – 17162.
doi:10.1073/pnas.0809353105
43.   Dickey CW, Verzhbinsky IA, Jiang X, Rosen BQ, Kajfez S, Stedelin B, et al. Widespread ripples synchronize
human cortical activity during sleep, waking, and memory recall. bioRxiv. 2021; 2021.02.23.432582.
doi:10.1101/2021.02.23.432582
44.   Mak-McCully RA, Deiss SR, Rosen BQ, Jung KY, Sejnowski TJ, Bastuji H, et al. Synchronization of Isolated
Downstates (K-Complexes) May Be Caused by Cortically-Induced Disruption of Thalamic Spindling. PLoS
Comput Biol. 2014;10. doi:10.1371/journal.pcbi.1003855
45.   Beul SF, Barbas H, Hilgetag CC. A Predictive Structural Model of the Primate Connectome. Sci Rep.
2017;7: 1–12. doi:10.1038/srep43176
46.   Deco G, Sanz Perl Y, Vuust P, Tagliazucchi E, Kennedy H, Kringelbach ML. Rare long-range cortical
connections enhance human information processing. Curr Biol. 2021;31: 4436-4448.e5.
doi:https://doi.org/10.1016/j.cub.2021.07.064
47.   Aboitiz F, López J, Montiel J. Long distance communication in the human brain: timing constraints for
inter-hemispheric synchrony and the origin of brain lateralization. Biol Res. 2003;36: 89–99.
48.   Peters A, Payne BR, Budd J. A numerical analysis of the geniculocortical input to striate cortex in the
monkey. Cereb Cortex. 1994;4: 215–229.
49.   Chung S, Ferster D. Strength and orientation tuning of the thalamic input to simple cells revealed by
electrically evoked cortical suppression. Neuron. 1998;20: 1177–1189.
50.   Van Den Heuvel MP, Sporns O. Rich-club organization of the human connectome. J Neurosci. 2011;31:
15775–15786.
51.   Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K. The WU-Minn Human
Connectome Project: An overview. Neuroimage. 2013;80: 62–79. doi:10.1016/j.neuroimage.2013.05.041
52.   Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. Probabilistic diffusion tractography with
multiple fibre orientations: What can we gain? Neuroimage. 2007;34: 144–155.
doi:10.1016/j.neuroimage.2006.09.018
53.   Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. Neuroimage. 2012;62: 782–790.
doi:10.1016/j.neuroimage.2011.09.015
54.   Tournier JD, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: A fast, flexible and open
software framework for medical image processing and visualisation. Neuroimage. 2019;202: 116137.

doi:10.1016/j.neuroimage.2019.116137

55.    Tournier JD, Calamante F, Connelly A. Improved probabilistic streamlines tractography by 2nd order
integration over fibre orientation distributions. Proceedings of the international society for magnetic
resonance in medicine. John Wiley & Sons, Inc. New Jersey, USA; 2010.

56.    Meyer M, Desbrun M, Schröder P, Barr AH. Discrete Differential-Geometry Operators for Triangulated 2-
Manifolds. 2003. pp. 35–57. doi:10.1007/978-3-662-05105-4_2

57.    Wegiel J, Flory M, Kaczmarski W, Brown WT, Chadman K, Wisniewski T, et al. Partial Agenesis and
Hypoplasia of the Corpus Callosum in Idiopathic Autism. J Neuropathol Exp Neurol. 2017;76: 225–237.
doi:10.1093/jnen/nlx003

58.    Williams EJ. The comparison of regression variables. J R Stat Soc Ser B. 1959;21: 396–399.

59.    Ji JL, Spronk M, Kulkarni K, Repovš G, Anticevic A, Cole MW. Mapping the human brain's cortical-
subcortical functional network organization. Neuroimage. 2019;185: 35–57.
doi:10.1016/j.neuroimage.2018.10.006

60.    Blinkov SM, Glezer II. The human brain in figures and tables: a quantitative handbook. Basic Books; 1968.

61.    von Bartheld CS, Bahney J, Herculano-Houzel S. The search for true numbers of neurons and glial cells in
the human brain: A review of 150 years of cell counting. J Comp Neurol. 2016;524: 3865–3895.
doi:10.1002/cne.24040

62.    Beckmann H, Lauer M. The human striatum in schizophrenia. II. Increased number of striatal neurons in
schizophrenics. Psychiatry Res Neuroimaging. 1997;68: 99–109.

63.    Baker KG, Halliday GM, Hornung J-P, Geffen LB, Cotton RGH. Distribution, morphology and number of
monoamine-synthesizing and substance P-containing neurons in the human dorsal raphe nucleus.
Neuroscience. 1991;42: 757–775.

64.    Mouton PR, Pakkenberg B, Gundersen HJG, Price DL. Absolute number and size of pigmented locus
coeruleus neurons in young and aged individuals. J Chem Neuroanat. 1994;7: 185–190.

65.    Eriksen N, Stark AK, Pakkenberg B. Age and Parkinson's disease-related neuronal death in the substantia
nigra pars compacta. Birth, life death dopaminergic neurons Subst Nigra. 2009; 203–213.

66.    Mufson EJ, Ginsberg SD, Ikonomovic MD, DeKosky ST. Human cholinergic basal forebrain:
chemoanatomy and neurologic dysfunction. J Chem Neuroanat. 2003;26: 233–242.

67.    Rivara C, Sherwood CC, Bouras C, Hof PR. Stereologic characterization and spatial distribution patterns of
Betz cells in the human primary motor cortex. Anat Rec Part A Discov Mol Cell Evol Biol An Off Publ Am
Assoc Anat. 2003;270: 137–151.

68.    Wada A, Goto J, Goto N, Kawamura N, Matsumoto K. Are There One Million Nerve Fibres in the Human
Medullary Pyramid? Okajimas Folia Anat Jpn. 2001;77: 221–224. doi:10.2535/ofaj1936.77.6_221

## S1 Appendix: Extended Methods and Results

### Diffusion MRI

The diffusion-weighted MRI (dMRI) connectivity data used to estimate inter-parcel axon totals was

described in our prior report [1] whose data are available in full at

https://doi.org/10.5281/zenodo.4060485. Preprocessed imaging data for 1,065 healthy young adults

were retrieved from the Human Connectome Project WU-Minn consortium 1200 release [2]

(https://db.humanconnectome.org). We segmented the cortex into 180 parcels per hemisphere

following the HCP-MMP1.0 atlas [3] and performed probabilistic tractography using the probtrackX

tool [4] from FSL [5] using tractography parameters detailed in our prior report [1]. This analysis

yielded a 360 x 360 dMRI connectivity matrix for each subject which provided the example data for

applying the method presented here. For probtrackX-derived connectivity, the raw unit of dMRI

connectivity strength is the number of streamlines connecting seed and target regions. The absolute

magnitude of this metric is arbitrary as it scales with the number of samples drawn from the probability

distribution of diffusion parameters in each voxel, in our analysis 5,000.

The individual pairwise streamline counts and fiber tract length matrices for the replication analysis

were described in [6] whose data are available at https://doi.org/10.5281/zenodo.4733297. These

data consist of a 972 subject subset of the 1,065 WU-Minn HCP subjects included in our primary

analysis. Arnatkeviciute and colleagues similarly segmented the cortex following the HCP-MMP1.0

atlas [3]. In contrast to our primary dataset, tractography was performed with MRtrix3 [7] using second-

order integration over fiber orientation distributions (iFOD2) [8], see [6] for details. When confined to

the same subject subset and non-null connections, the pairwise tract lengths and log-transformed

streamline counts are highly correlated ($r^2$ = 0.78 and $r^2$ = 0.69, respectively) between Arnatkeviciute

et al. [6] and our study [1]. The rate of exponential decay in the number of streamlines vs fiber tract

distance is also similar, $\lambda$ = 26.1 mm in their study vs 23.4 mm in ours, though less connectivity variance

is explained by distance in their results ($r^2$ = 0.12 vs. $r^2$ = 0.52). Results for these tractography data are

shown in S2 Fig.

Cross-sectional area of the corpus callosum, white matter volume, and parcel areas

The HCP WU-Minn consortium 1200 release [2] includes the standard outputs from the FreeSurfer

`recon-all` pipeline [9]. With default parameters, this pipeline labels the 1 $mm^3$ voxels of the `aseg.mgz`

volume which contain the corpus callosum with a fixed 5mm lateral extent. Dividing the number of

callosum-label voxels by 5 yields an estimate of its cross-sectional area, in $mm^2$. The total white matter

volume is reported in the `aseg.stats` output of `mri_segstats`. Parcel areas and adjacency were

determined [10] on triangular mesh of each subject's reconstruct white matter surface, or white matter

– gray matter interface (`?h.white`). These data were used for both the primary and replication analyses.

Cross-sectional fiber packing density in corpus callosum and other hemispheric white
matter

Our estimate of callosal fiber density was based on the electron microscopy (EM) study by Aboitiz and

colleagues [11] who reported a fiber density of 38 per 100 $\mu m^2$ or 3.7x$10^5$ axons/$mm^2$ after accounting

for tissue shrinkage to 96% volume after Epon fixation [12]. This is consistent with their finding of

2.65x$10^5$ axons/$mm^2$ as measured with light-microscopy in twenty brains as well, assuming shrinkage

to 43% volume for paraffin fixation [12] and an estimated ~20% of fibers not detectable with light

microscopy [11], as well as the 2.83x$10^5$ non-exhaustively counted axons per $mm^2$ reported in a more

recent EM study of callosal axon diameter in two brains [13]. This value was derived by dividing the

reported mean count of axons per callosal region by the callosal area surveyed, and correcting for EM

preparation tissue shrinkage. An additional light microscopy study of callosal axon density in eleven

control cortices reports an average of 1.12x$10^5$ myelinated axons per $mm^2$ [14], assuming shrinkage to

shrinkage to 43% volume. This somewhat lower figure may be explained by the inclusion of four

children's cortices in their cohort, or by mis-assumed shrinkage, but all reports are within half an order-

of-magnitude of each other.

<u>Distance matched intra- vs. inter-hemispheric dMRI connectivity</u>
One assumption of the method described in this report is that the dMRI tractography methodology

has approximately similar sensitivity to callosal and ipsilateral connections. This is difficult to determine

without ground truth connectivity, as intra- and inter-hemispheric connections may have true

anatomical differences. Nevertheless, we examined distance-matched intra- vs inter-hemispheric

connections (Fig 1*C*) and found no statistical evidence that they are different. Pairwise connections

were averaged in 15 distance bins, log-transformed and linear fits obtained for intra- and inter-

hemispheric connections. For the overlapping distance domain, the intra-hemispheric slope is -0.30,

95% confidence interval = [-0.33 -0.26] with an intercept of 6.18 [5.91 6.45] and the inter-hemispheric

slope is -0.26 [-0.29 -0.24] and intercept of 5.93 [5.72 6.14]; the 95% confidence intervals of both

parameters are overlapping. In addition, a paired r-test of correlation differences [15] failed to reject

the null hypothesis, $p$ = 0.58. The parity between distance-matched intra- and inter-hemispheric

connections also holds when the linear fit is obtained for over the entire distance domain, with an intra-

hemispheric slope and intercept -.31 [-0.33 -0.28] and 6.24 [6.01 6.46], respectively; -.28 [-0.30 -0.25]

and 6.00 [5.78 6.23] for inter-hemispheric connections. In the replication analysis with alternative

tractography data [6] (S2 Fig C) the paired r-test likewise failed to reject the null hypothesis, $p$ = 0.34.

The intra-hemispheric slope and intercept are -.37 [-0.37 -0.78] and 4.19 [3.25 5.13], respectively; -.25

[-0.30 -0.20] and 3.30 [2.82 3.79] for inter-hemispheric connections.

<u>Estimating axon counts and axon volume from dMRI</u>
From the histological estimate of callosal fiber density, MRI-derived measures of callosal cross-

sectional area and white matter connectivity, the conversion factor between dMRI connectivity and

physical axons can be estimated as follows:

Eq. 1.1 $N_{callosum} = A_{callosum} * D_{callosum}$

Eq. 1.2 $C_{axon} = \frac{Ncallosum}{M_{dMRI}}$

Where $A_{callosum}$ is the cross-sectional area (mm$^2$), $D_{callosum}$ is the assumed fiber density (3.7x10$^5$

axons/mm$^2$) of the corpus callosum, $N_{callosum}$ is the estimated total number of callosal fibers, $M_{dMRI}$ is the

magnitude of dMRI callosal connectivity, in our case, the number of tractography streamlines, and $C_{axon}$

is the conversion factor from dMRI to physical fibers (axons/streamline).

In addition to connectivity strength, or number of streamlines, the dMRI tractography analysis also

yields a mean streamline length for every parcel-pair connection. After converting dMRI connectivity

strength to physical axon count as described above, the total volume of the estimated axons can be

calculated as follows:

Eq. 2.1 $WM_{volume} = \sum_{i=1}^{N_{pair}} l_i * n_i * C_{axon} * \frac{1}{D_{callosum}}$

Eq. 2.2 $WM_{volume} = \sum_{i=1}^{N_{pair}} l_i * n_i * \frac{A_{callosum}}{M_{dMRI}}$

Where $WM_{volume}$ is the estimated axonal white matter volume (mm$^3$) and $N_{pair}$ is the total number of

parcel pairs. $l_i$ and $n_i$ are the length (mm) and number of dMRI streamlines in the $i^{th}$ parcel pair. Eq. 2.1

can be simplified into Eq. 2.2 by substituting Eqs. 1.1 and 1.2 into 2.1.

667
668    <u>Parcel grouping into functional networks</u>
669    The parcel ordering and the ten functional networks shown in Fig 1*A* were defined in our previous

670    report [1] and were modified from twelve networks identified by Ji and colleagues in a resting-state

671    fMRI study on the HCP cohort [16] by merging the primary and secondary visual networks as well as

672    the ventral and posterior multimodal networks.

673
674
675

## Effects of sex and age on inter-hemispheric connectivity

In their histological study of 20 cortices , Aboitiz and colleagues [11] found no significant effects of sex

or age on the total number of fibers, fiber density, and cross-sectional area of the corpus callosum. As

we assume a constant fiber density, our estimate of the total number of inter-hemispheric fibers is a

linear function of the callosal cross-sectional area. As the ages of the HCP 1200 cohort are grouped

into four broad ranges in the open access data, we investigated the effects of sex and age on the

estimated number inter-hemispheric axons with 2x4 fixed effects ANOVA, treating the age group

factor as categorical. Mean and individual values are shown in S3 Fig. Note that while the number of

pairwise axons is approximately log-normally distributed across areal pairs, it is approximately

normally distributed across individuals.

We found that, on average, males have $2.61 \times 10^8$ inter-hemispheric axons, 95% confidence interval

$= [2.58 \times 10^8 \ 2.64 \times 10^8]$ and females have $2.50 \times 10^8 \ [2.47 \times 10^8 \ 2.52 \times 10^8]$ axons. Subjects ages 22-25, 26-

30, 31-35, and 36+ year old have $2.50 \times 10^8 \ [2.46 \times 10^8 \ 2.55 \times 10^8]$, $2.58 \times 10^8 \ [2.55 \times 10^8 \ 2.61 \times 10^8]$, $2.55 \times 10^8$

$[2.51 \times 10^8 \ 2.59 \times 10^8]$, and $2.35 \times 10^8 \ [2.15 \times 10^8 \ 2.54 \times 10^8]$ axons, respectively. Intervals show

bootstrapped 95% confidence. We find no effect of sex on the number of inter-hemispheric axons,

$F_{1,1064} = 3.471$, $p = 0.0627$. Removing the one female outlier does not bring the mean difference into

statistical significance, $F_{1,1063} = 3.653$, $p = 0.0562$. The interaction of sex and age on the number of

inter-hemispheric axons is likewise not significant $F_{3,1064} = 2.282$, $p = 0.0776$. Age does significantly

affect the number of inter-hemispheric neurons, $F_{3,1064} = 5.100$, $p = 0.0017$. Post-hoc pairwise tests

show that only the difference between the 22-25 and 26-30 year old age groups is significant, $F_{1,1064} =$

7.646, $p = 0.0058$. These results are broadly consistent with those of Aboitiz et al. [11]. A more in-

depth examination of the effects of age on corpus callosum requires a cohort with which includes

more older subjects as well as more precise age values.

## Thalamo-cortical fiber volume estimation

The scope of our previous dMRI tractography study [1] was confined to cortico-cortical (including

cortico-hippocampal) connectivity. In order to estimate the total volume of thalamo-cortical axons, we

first calculated the number of excitatory thalamic neurons by summing the total number of neurons in

the thalamus [17], excluding the reticular nucleus, zona incerta, limitans/suprageniculate, and

subthalamus, and including from the remainder the ~62% proportion of thalamic neurons that are

excitatory [18]. The number of such neurons is $22.6 \times 10^6$. We assumed that each such thalamic neuron

projects a single axon to the cortex, and we allocated these axons to cortical parcels in proportion to

their area. The inverse callosal packing density was used for effective cross-sectional area, as with

cortico-cortical connections. To estimate the length of thalamo-cortical fiber tracts, for each parcel $i$,

we first identified the thalamic voxel nearest to parcel's centroid. We then used the inter-parcel fiber

tract length [1] from the centroid of parcel $i$ to the identified parathalamic parcel centroid. To this

figure we added the Euclidean distance from the parathalamic centroid to the nearest voxel of the

thalamus, ensuring that estimated thalamo-cortical fiber tract length is always non-zero. As with

cortico-cortical connections, the total white matter volume of each connection was estimated by taking

the product of the number of axons, the effective cross-sectional area, and the estimated fiber tract

length (Eq. 2).

## Estimation of proportion of actual inter-areal axons to the number that would be
## needed for complete inter-connectivity

We estimated that the total number of inter-areal axons is $\sim 2.43 \times 10^9$. Assuming n = $16.34 \times 10^9$ total

cortical neurons (including interneurons) [19] , the total number of connections needed for complete

whole-cortex connectivity is n(n-1) = $2.67 \times 10^{20}$. We estimated the neural density of the cortex as

$\sim 92,300$ neurons/mm$^2$ by dividing $16.34 \times 10^9$ by $1.77 \times 10^5$ mm$^2$ mean white–gray surface area of the

HCP cohort used. The total number of connections needed for complete inter-connectivity within each

area is approximately 1.18x10$^{18}$, given by

Eq. 3.1 $N_{completeIntra-areal} = \Sigma_{i=1}^{Nparcel}\left(A_{parcel} * D_{neuron}\right)^2$

Where A$_{parcel}$ is the area of each parcel and D$_{neuron}$ the neural density. Subtracting the connections

needed for within-area inter-connectivity from whole cortex interconnectivity yields 2.66x10$^{20}$, which is

1.10x10$^{11}$ times greater than the number of inter-areal contacts we calculate with our method.

<u>Number of neurons in non-cortical regions communicating with the cortex</u>

The number of neurons in different human brain regions has been reviewed by Blinkov and Glexer [21]

and von Bartheld et al. [22]. The number of neurons in the amygdala have been estimated as ~13x10$^6$

[23], striatum as ~55x10$^6$ [24], and thalamus as ~23x10$^6$ (thalamocortical, see above). These are the

only recipients or targets of cortical connections with significant numbers of neurons. Other locations,

such those which are the origins of serotonin [25], norepinephrine [26], dopamine [27], or

acetylcholine [28] afferents to the cortex all have ~50,000 axons each. Summed together, the number

of cells in all of the above structures comes to ~0.5% of the number of cortical cells. Direct projections

outside of the brain are exceedingly rare, with the number of Betz cells in the primary motor cortex

[29] or medullary pyramid [30] both estimated as ~100,000. It is difficult to escape the conclusion that,

at least in humans, the cortex communicates mainly with itself.

744
745

<u>References</u>

1.    Rosen BQ, Halgren E. A Whole-Cortex Probabilistic Diffusion Tractography Connectome. eNeuro. 2021;8:
ENEURO.0416-20.2020. doi:10.1523/ENEURO.0416-20.2020
2.    Van Essen DC, Smith SM, Barch DM, Behrens TEJ, Yacoub E, Ugurbil K. The WU-Minn Human
Connectome Project: An overview. Neuroimage. 2013;80: 62–79. doi:10.1016/j.neuroimage.2013.05.041
3.    Glasser MF, Coalson TS, Robinson EC, Hacker CD, Harwell J, Yacoub E, et al. A multi-modal parcellation of
human cerebral cortex. Nature. 2016;536: 171–8. doi:10.1038/nature18933

4.    Behrens TEJ, Berg HJ, Jbabdi S, Rushworth MFS, Woolrich MW. Probabilistic diffusion tractography with
754           multiple fibre orientations: What can we gain? Neuroimage. 2007;34: 144–155.
755           doi:10.1016/j.neuroimage.2006.09.018

756    5.    Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. Neuroimage. 2012;62: 782–790.
757           doi:10.1016/j.neuroimage.2011.09.015

758    6.    Arnatkeviciute A, Fulcher BD, Oldham S, Tiego J, Paquola C, Gerring Z, et al. Genetic influences on hub
759           connectivity of the human connectome. Nat Commun. 2021;12: 4237. doi:10.1038/s41467-021-24306-2

760    7.    Tournier JD, Smith R, Raffelt D, Tabbara R, Dhollander T, Pietsch M, et al. MRtrix3: A fast, flexible and open
761           software framework for medical image processing and visualisation. Neuroimage. 2019;202: 116137.
762           doi:10.1016/j.neuroimage.2019.116137

763    8.    Tournier JD, Calamante F, Connelly A. Improved probabilistic streamlines tractography by 2nd order
764           integration over fibre orientation distributions. Proceedings of the international society for magnetic
765           resonance in medicine. John Wiley & Sons, Inc. New Jersey, USA; 2010.

766    9.    Fischl B. FreeSurfer. Neuroimage. 2012;62: 774–781. doi:10.1016/j.neuroimage.2012.01.021

767    10.    Meyer M, Desbrun M, Schröder P, Barr AH. Discrete Differential-Geometry Operators for Triangulated 2-
768           Manifolds. 2003. pp. 35–57. doi:10.1007/978-3-662-05105-4_2

769    11.    Aboitiz F, Scheibel AB, Fisher RS, Zaidel E. Fiber composition of the human corpus callosum. Brain Res.
770           1992;598: 143–153. doi:10.1016/0006-8993(92)90178-C

771    12.    Schüz A, Palm G. Density of neurons and synapses in the cerebral cortex of the mouse. J Comp Neurol.
772           1989;286: 442–455.

773    13.    Liewald D, Miller R, Logothetis N, Wagner HJ, Schüz A. Distribution of axon diameters in cortical white
774           matter: an electron-microscopic study on three human brains and a macaque. Biol Cybern. 2014;108:
775           541–557. doi:10.1007/s00422-014-0626-2

776    14.    Wegiel J, Flory M, Kaczmarski W, Brown WT, Chadman K, Wisniewski T, et al. Partial Agenesis and
777           Hypoplasia of the Corpus Callosum in Idiopathic Autism. J Neuropathol Exp Neurol. 2017;76: 225–237.
778           doi:10.1093/jnen/nlx003

779    15.    Williams EJ. The comparison of regression variables. J R Stat Soc Ser B. 1959;21: 396–399.

780    16.    Ji JL, Spronk M, Kulkarni K, Repovš G, Anticevic A, Cole MW. Mapping the human brain's cortical-
781           subcortical functional network organization. Neuroimage. 2019;185: 35–57.
782           doi:10.1016/j.neuroimage.2018.10.006

783    17.    Xuereb JH, Perry RH, Candy JM, Perry EK, Bonham JR. Nerve cell Loss in the thalamus in Alzheimer's
784           disease and Parkinson's disease. Brain. 1991;114: 1363–1379.

785    18.    Arcelli P, Frassoni C, Regondi MC, De Biasi S, Spreafico R. GABAergic neurons in mammalian thalamus: a
786           marker of thalamic complexity? Brain Res Bull. 1997;42: 27–37.

787    19.    Azevedo FAC, Carvalho LRB, Grinberg LT, Farfel JM, Ferretti REL, Leite REP, et al. Equal numbers of
788           neuronal and nonneuronal cells make the human brain an isometrically scaled-up primate brain. J Comp
789           Neurol. 2009;513: 532–541. doi:10.1002/cne.21974

790    20.    DeFelipe J, Fariñas I. The pyramidal neuron of the cerebral cortex: Morphological and chemical
791           characteristics of the synaptic inputs. Prog Neurobiol. 1992;39: 563–607. doi:10.1016/0301-
792           0082(92)90015-7

793    21.    Blinkov SM, Glezer II. The human brain in figures and tables: a quantitative handbook. Basic Books; 1968.

794    22.    von Bartheld CS, Bahney J, Herculano-Houzel S. The search for true numbers of neurons and glial cells in
795           the human brain: A review of 150 years of cell counting. J Comp Neurol. 2016;524: 3865–3895.
796           doi:10.1002/cne.24040

797    23.    Avino TA, Barger N, Vargas M V, Carlson EL, Amaral DG, Bauman MD, et al. Neuron numbers increase in
798           the human amygdala from birth to adulthood, but not in autism. Proc Natl Acad Sci. 2018;115: 3710–3715.

799    24.    Beckmann H, Lauer M. The human striatum in schizophrenia. II. Increased number of striatal neurons in
800           schizophrenics. Psychiatry Res Neuroimaging. 1997;68: 99–109.

801    25.    Baker KG, Halliday GM, Hornung J-P, Geffen LB, Cotton RGH. Distribution, morphology and number of
802           monoamine-synthesizing and substance P-containing neurons in the human dorsal raphe nucleus.
803           Neuroscience. 1991;42: 757–775.

26.    Mouton PR, Pakkenberg B, Gundersen HJG, Price DL. Absolute number and size of pigmented locus

coeruleus neurons in young and aged individuals. J Chem Neuroanat. 1994;7: 185–190.

27.   Eriksen N, Stark AK, Pakkenberg B. Age and Parkinson's disease-related neuronal death in the substantia

nigra pars compacta. Birth, life death dopaminergic neurons Subst Nigra. 2009; 203–213.

28.   Mufson EJ, Ginsberg SD, Ikonomovic MD, DeKosky ST. Human cholinergic basal forebrain:

chemoanatomy and neurologic dysfunction. J Chem Neuroanat. 2003;26: 233–242.

29.   Rivara C, Sherwood CC, Bouras C, Hof PR. Stereologic characterization and spatial distribution patterns of

Betz cells in the human primary motor cortex. Anat Rec Part A Discov Mol Cell Evol Biol An Off Publ Am

Assoc Anat. 2003;270: 137–151.

30.   Wada A, Goto J, Goto N, Kawamura N, Matsumoto K. Are There One Million Nerve Fibres in the Human

Medullary Pyramid? Okajimas Folia Anat Jpn. 2001;77: 221–224. doi:10.2535/ofaj1936.77.6_221

816
**S2 Fig: Replication results.** Results obtained by repeating the analysis using an alternative

tractography dataset [1] (A) Connectivity matrix of intra-hemispheric axon counts, averaged across the

two hemispheres. Parcels are ordered into 10 functional networks. (B) Histograms showing the

distribution of inter- and intra-hemispheric pairwise axon counts. Physically adjacent and non-adjacent

parcel pair intra-hemispheric histograms are stacked. Median connectivity, shown in gray, is ~2,500

axons between cortical areas in the same hemisphere, and ~300 inter-hemispherically. Parcel pairs

with zero axons connecting them are represented by the bars left of the y-axis. (C) dMRI connectivity as

a function fiber tract distance. Pairwise values averaged within 15 fiber-length bins. Shading shows

bootstrapped 95% confidence intervals. (D-F) axons counts and densities averaged across the two

hemispheres and visualized on the left fsaverage template cortex [2]. (D) total axons connecting each

parcel to all others. (E) axons connecting each parcel to all others, normalized by the reference parcel's

area. (F) axons connecting the hippocampus to the rest of the cortex, normalized by the area of the

cortical parcel, shown in log scale.

830
831    1. Arnatkeviciute A, Fulcher BD, Oldham S, Tiego J, Paquola C, Gerring Z, et al. Genetic influences on hub
832         connectivity of the human connectome. Nat Commun. 2021;12: 4237. doi:10.1038/s41467-021-24306-2
833    2. Fischl B. FreeSurfer. Neuroimage. 2012;62: 774–781. doi:10.1016/j.neuroimage.2012.01.02

839
**840    S3 Fig: Effects of sex and age on the inter-hemispheric connectivity.** Each individual's total

number of estimated inter-hemispheric axons is shown with a marker. The black horizontal bars show

the group means and vertical bars the bootstrapped 95% confidence intervals these means. Shading

shows a kernel density estimate of the group distributions. Note that while the number of pairwise

axons is approximately log-normally distributed across areal pairs, it is approximately normally

distributed across individuals. The only significant group difference is between the 22-15 and 25-30

age groups, $F_{1,1064}$ = 7.646, $p$ = 0.0058. Interactions between sex and age effect were not significant.

As we assume a constant fiber density, our estimate the total number of inter-hemispheric fibers is a

linear multiple of the callosal cross-sectional area.

849
**S4 Fig: Arcuate / superior lateral fasciculus termination fields.** In order to estimate the fraction of

tract axons that travel the entire length of the tract, it's termination fields, centered on Broca's and

Wernicke's areas, were manually defined in terms of HCP-MMP1.0 parcels according to consensus

definitions [1]. (A) Conservative definitions where the anterior termination field, in teal, is composed of

parcels 44, 45, 6r, IFSa, IFSp, and FOP4 and the posterior termination field, in yellow, is composed of

parcels PSL, RI, STV, and PFcm. For this definition trans-terminal axons account for 0.6% and 0.8% of

tract axons in the left and right hemispheres. (B) Liberal definitions in which parcels 47l and p47r were

added anteriorly and parcels PF, PFm, and PGi were added posteriorly, resulting in trans-terminal

axons accounting for 1.9% and 2.9% of tract axons in the left and right hemispheres. Gray lines indicate

the approximate locations at which the tract diameter, used to estimate the total number of tract axons

were ascertained [2].

861
862    1. Tremblay P, Dick AS. Broca and Wernicke are dead, or moving past the classic model of language
863          neurobiology. Brain Lang. 2016;162: 60–71. doi:https://doi.org/10.1016/j.bandl.2016.08.004
864    2. Yeh F-C. Shape analysis of the human association pathways. Neuroimage. 2020;223: 117329.
865        doi:https://doi.org/10.1016/j.neuroimage.2020.117329
866
867

**S5 Fig: Simulated effect of short axon underestimation on pairwise axon counts.** In order to

explore the possibility that dMRI tractography is less sensitive to shorter connections the streamline-to-

axon ratio was increased linearly with inverse fiber length using a range of slope parameters. Packing

density was assumed to be reciprocally decreased in order to fix a constant total cerebral white matter

volume. (A) Distributions of Intra- and inter-hemispheric inter-areal axons counts. Gray histograms

show the effect of uniformly halfling the ratio and doubling the assumed packing density. (B) intra-

hemispheric and (C) inter-hemispheric adjustments to the # of pairwise axons as a function of inverse

fiber tract length and the resultant increase in median axon count as a function of the adjustments'

slope parameter. At slope = 0, values are unadjusted from the primary analysis. Large adjustments

only increase median counts modestly in the context of the log-normal distribution. For example, for

intra-hemispheric connections (B), at a slope of 0.004, the number of the shortest axons is about

doubled (i.e., corresponding to doubling the axon density and halving the sensitivity of dMRI to

axons), but the median number of inter-parcel axons only increases by ~36%.