Asexual freshwater snails make poor mate choice decisions

Sydney Stork[1], Joseph Jalinsky[1], Maurine Neiman[1,2]

[1]Department of Biology, University of Iowa, Iowa City, IA, 52242, USA

[2]Department of Gender, Women's, and Sexuality Studies, University of Iowa, Iowa City, IA, 52242, USA.

Abstract

Once-useful traits that no longer contribute to fitness tend to decay over time. We address

whether the expression of mating-related traits that increase the fitness of sexually reproducing

individuals but are likely less useful or even costly to asexual counterparts seems to exhibit

decay in the latter. *Potamopyrgus antipodarum* is a New Zealand freshwater snail characterized

by repeated transitions from sexual to asexual reproduction. The frequent coexistence of sexual

and asexual lineages makes *P. antipodarum* an excellent model for the study of mating-related

trait loss. We used a mating choice assay including sexual and asexual *P. antipodarum* females

and conspecific (presumed better choice) vs. heterospecific (presumed worse choice) males to

evaluate the loss of behavioural traits related to sexual reproduction. We found that sexual

females engaged in mating behaviours with conspecific mating partners more frequently and for

a greater duration than with heterospecific mating partners, while asexual females seemed to lack

the ability to make a choice. These results suggest that selection acting to maintain mate choice

in asexual *P. antipodarum* is weak or ineffective relative to sexual females and that asexual

reproduction likely contributes to the evolutionary decay of behavioural traits in this system.

Keywords: Asexuality, sexual reproduction, *Potamopyrgus antipodarum*, vestigial traits,

parthenogenesis, trait decay

Introduction

Adaptations that were once important for survival that subsequently become useless or impose new fitness costs tend to decay over time in a process known as vestigialisation [1-3]. Many examples of this phenomenon can be seen throughout evolutionary history, including the loss of limbs in snakes [4], eye structures in cave-dwelling fish [5], and wings in many insect species [6].

While the genetic mechanisms underlying vestigialisation depend on evolutionary context, traits are most commonly lost by the accumulation of selectively neutral or deleterious mutations [7, 8]. The decay of presently useless characteristics exemplifies evolutionary tradeoffs: the decay of one trait frees up energy and resources to be used elsewhere by the organism [7]. For example, Roff [6] suggests that the evolution of flightlessness from flighted ancestors allows for allocation of resources for reproduction. This framework implies that trait loss could translate into increased fitness for organisms for which the lost trait is no longer useful.

The decay of traits specific to mating and/or sexual reproduction is one potential consequence of transitions from sexual to asexual reproduction. These traits are typically required for reproductive success in sexual organisms. By definition, asexual organisms do not need external genetic contribution to reproduce, with the exception of sperm-dependent forms of parthenogenesis (e.g., gynogenesis). Mating behaviours are themselves associated with numerous costs [9, 10]. Accordingly, vestigialisation of mating-related traits in asexual lineages – including those involved in mate choice – is expected, particularly in a scenario where such behaviours and their maintenance are costly [3, 11, 12]. Alternatively, these traits may be selectively neutral or maintained by selection in asexuals, but only in a context where the costs

do not outweigh the benefits of trait maintenance [13, 14]. Several studies have supported these

ideas, finding abnormal or fully decayed traits involved in mating and reproduction

(spermathecae in asexual *Timema* stick insects [13]; sperm in asexual *P. antipodarum* males

[15]) in asexual organisms. These results suggest that these traits are not being maintained by

selection, at least to the extent of the same traits in sexual counterparts. Our focus is on

behaviours: assuming that mate quality contributes to fitness of sexual but not asexual females,

do mating-related behaviours related to the likelihood that mate preferences are associated with

mate quality [16-19] decay in asexual females relative to sexual counterparts?

Potamopyrgus antipodarum is a New Zealand freshwater snail characterized by the

frequent coexistence of phenotypically and ecologically similar sexual and asexual lineages [20,

21], making this species an excellent model to study the vestigialization of traits involved in

sexual reproduction. Previous studies have shown that mating behaviour in sexually reproducing

male *P. antipodarum* does not differ from that of the presumed asexual males produced

occasionally by asexual females [22]. Sexual males also show poor ability to discriminate

between favourable and unfavourable mates, mating with equal frequency with sexual and

asexual females and with parasitically castrated vs. healthy females [23]. Prior emphasis on

investigation of male mating behaviour is largely a result of the assumption that males are

physically in control of mating outcomes [24], despite a lack of studies assessing the female role.

These previous results set the stage for a direct comparison between sexual and asexual *P.*

antipodarum females. In a mating paradigm like that demonstrated for *P. antipodarum* where

males seem to lack adequate mate discrimination, differences in mating between sexual and

asexual females can be attributed to mating differences between these two female groups.

Therefore, we hypothesize that sexual female *P. antipodarum* are better able to choose a more

favourable mate when presented with a choice than asexual counterparts.

We evaluated this hypothesis by comparing the mating behaviour of sexual and asexual

P. antipodarum females with sexual male *P. antipodarum* and sexual male *Potamopyrgus*

estuarinus. *Potamopyrgus estuarinus* is a closely related outgroup species with minimal habitat

overlap with *P. antipodarum* [25]. Female *P. antipodarum* mating efforts with *P. antipodarum*

males are presumed to be favourable relative to mating with *P. estuarinus* because of the

possibility to generate offspring only with the former. Mating behavior (but not offspring

production) has been shown to occur between male and female *P. antipodarum* and *P. estuarinus*

[26], thereby providing *P. antipodarum* females with an adequate alternative choice.

Materials and Methods

Snail Selection and Care

We haphazardly selected three sets of 10 sexually mature (>3 mm long [27, 28]) sexual (N=30)

and asexual (N=30) female *P. antipodarum* descended from one sexual and one asexual female,

respectively, collected from Lake Mapourika, New Zealand, in 2018. These six sets of 10 female

snails will hereafter be referred to as 'populations.' We also haphazardly selected three sets of 20

each sexually mature (possessing a visible penis [29]) sexual *P. antipodarum* males (N=60) and

Potamopyrgus estuarinus males (N=60). All *P. antipodarum* males were chosen from sexual

lineages established by females sampled from New Zealand Lakes Mapourika and Selfe in 2018.

These lineages were separately derived from the lineages from which we drew females. We

isolated each population for approximately 14 days prior to behavioural trials to control for

previous exposure to males and possible recent mating. We painted female shells with light-

coloured nail polish to aid in differentiation between females and males when analysing video

footage.

We housed each population in a 0.95 L plastic cup containing ~700 mL of water. Snails

were fed *Spirulina* (a common *P. antipodarum* lab food) three times weekly and were given

powdered chalk as a source of dietary calcium once weekly. Cups were maintained at 16°C on a

12:12 hour light:dark cycle (feeding and maintenance following [24]).

Mating trials occurred in standard four-inch petri dishes. We covered the exterior of each

dish in electrical tape to control for external visual stimuli. To begin each trial for the sexual

females, we placed one sexual *P. antipodarum* female, one sexual *P. antipodarum* male, and one

P. estuarinus male equidistant from one another around the edges of the inside of the petri dish

(Fig. 1). After the three snails were placed, we filled the petri dish to about 75% total volume

with carbon-filtered tap water. We used this same setup for asexual *P. antipodarum* females,

with the asexual female replacing the sexual female. No snail was used in more than one trial.

Observation of Mating Behavior

We conducted mating trials three times weekly at approximately 1500 hours, for three hours. As

the maximum copulation duration for *P. antipodarum* is about two hours [23], we established

this time frame to allow for multiple mating attempts to occur. We recorded trials using a USB

webcam and time-lapse video software VideoVelocity (v3.7.2090 [30]). The three snails per trial

were allowed to move and interact freely without physical barriers inside the petri dish.

We loosely followed previous studies of mating in *P. antipodarum* (e.g., [24]) to define

attempted mating behaviour as any physical interaction resulting in one snail mounting the shell

of another. It is relevant to note that it is otherwise impossible to otherwise visually confirm

sexual contact (joining of genitalia) between *P. antipodarum* because this phenomenon is

obscured by the shells. Once this mounting behaviour was initiated, we arbitrarily established

that any such interaction lasting 15 seconds or longer would be counted as an instance of mate

choice; interactions shorter than 15 seconds were potentially too short to be discernible with our

time-lapse software and were excluded. Fifteen seconds is also more than enough time for snails

to reorient and move on following an incidental – versus mating – encounter. Both the

frequency (number of mating interactions females had with either *P. antipodarum* or *P.*

estuarinus males) and duration (length of time (seconds) that females and males were in physical

contact) of mating interactions between females and either *P. antipodarum* or *P. estuarinus*

males were recorded. Hereafter, mating pairs will be referred to with the female listed first as

either SexF or AsexF, respectively, and the male referred to by species [*P. antipodarum* male =

PAM; *P. estuarinus* male = PEM (e.g., SexF/PAM refers to a sexual female and male *P.*

antipodarum pair)].

We also used this experimental framework to establish expected outcomes and determine

how these outcomes would inform our hypotheses. First, a successful experiment would require

that sexual females mate more frequently and for longer with *P. antipodarum* males than with *P.*

estuarinus males. This outcome is critical in indicating that sexual females possess the ability to

discriminate between preferential and poor mating partners. Absence of a difference in mating

behaviour across male type for sexual females would indicate that these females are unable to

differentiate between mates or that observed contact is reflective of behaviour outside of mating.

Mating-related trait decay in asexual females should be evident in relatively weak or absent mate

choice for *P. antipodarum* vs. *P. estuarinus* males relative to choices in sexual female *P.*

antipodarum.

Statistical Analyses

We used the Shapiro-Wilk test to determine that both the mating frequency (sexual females: $W =$

0.769, $p = 8.18\text{x}10^{-8}$; asexual females: $W = 0.810$, $p = 1.12\text{x}10^{-5}$) and the mating duration (sexual

females: $W = 0.642$, $p = 1.05\text{x}10^{-9}$; asexual females: $W = 0.493$, $p = 3.17\text{x}10^{-8}$) datasets violated

normality assumptions of parametric analysis. Accordingly, we used nonparametric approaches

to analyse the data. First, we used Wilcoxon rank-sum analyses to determine whether mating

durations and frequencies of sexual and asexual females, respectively, with *P. antipodarum*

males differed from mating durations and frequencies with *P. estuarinus* males. As another

means of comparing mating behaviour across sexual and asexual females, we calculated

discrimination scores - defined as the difference between mating frequency and mating duration,

respectively, with male *P. antipodarum* versus male *P. estuarinus* – for each female. We then

used Kruskal-Wallis tests to determine whether there was a significant difference in the mating

frequency and mating duration discrimination scores, respectively, between sexual and asexual

females. Finally, we used Kruskal-Wallis tests to compare the difference in mating frequency

and duration between individual sexual and asexual females. All analyses were executed with R

3.6.2 [31], the ggplot2 (v3.3.3; [32]), ggpubr (v0.4.0; [33]) and dplyr (v1.0.4; [34]) packages.

Results

Sexual females mated significantly more often with *P. antipodarum* males (mean 1.41 ± 1.39 SD

matings) than with *P. estuarinus* males (mean 0.67 ± 0.78 SD matings) ($W = 483$, $p = 0.0303$)

(Fig. 2A). By contrast, asexual females mated with *P. antipodarum* males (mean 1.20 ± 1.50 SD

matings) with the same frequency as with *P. estuarinus* males (1.45 ± 1.43 SD matings) ($W =$

172, $p = 0.427$) (Fig. 2B).

Sexual females also mated for significantly longer with *P. antipodarum* males (405.56 ±

520.95 SD seconds) than with *P. estuarinus* males (129.92 ± 349.74 SD seconds ($W = 520$, $p =$

0.00608) (Fig. 2C). Asexual females did not differ in the duration of mating with *P. antipodarum*

(138.73 ± 269.67 SD seconds) and *P. estuarinus* males (52.82 ± 54.88 SD seconds) ($W = 202$, p

$= 0.967$) (Fig. 2D).

The discrimination score analysis revealed that sexual but not asexual females with a

choice between *P. antipodarum* and *P. estuarinus* males tend to make mating behaviour decision

with respect to mating frequency that favour the former (Kruskal-Wallis $X^2 = 4.47$, df = 1, $p =$

0.0344). By contrast, there was no significant difference in difference scores for mating duration

between sexual and asexual females (Kruskal-Wallis $X^2 = 1.79$, df = 1, $p = 0.181$).

Discussion

Our experiment revealed that sexual female *P. antipodarum* exhibited significantly higher

mating frequency and longer mating duration with conspecific vs. heterospecific males. These

results are as predicted if sexual female *P. antipodarum* are able to discriminate between mates

of higher vs. lower quality. By contrast, asexual *P. antipodarum* females showed no evidence for

mate discrimination or preference. Together, these findings suggest that mate choice is exhibited

by sexual but not asexual *P. antipodarum* females.

Evolutionary decay of mating-related traits in asexual contexts is a plausible explanation

for these results and is consistent with a previous study demonstrating evidence for decay of

sperm traits in the male offspring occasionally produced by asexual female *P. antipodarum* [15].

Despite absence of evidence for discriminatory mating behaviour, asexual females mated

at a nearly identical frequency with *P. antipodarum* males as sexual females (Fig. 2). The high

prevalence of mating behaviour combined with the lack of observable discriminatory behaviour

in asexual females suggests that either copulatory behaviour is still necessary and/or beneficial,

or simply that this behaviour has not fully decayed. That asexual females nevertheless engage in

mating behaviour at all could be a function of the relatively recent derivation of most asexual

female *P. antipodarum* from sexual counterparts [35]. With this in mind, it would be interesting

to repeat this experiment with females from older asexual lineages (*sensu* [24] but departing

from this prior study so that females vs. males have mate choice capabilities).

Another, non-mutually exclusive explanation is that copulatory behaviour could be

required for maximal reproductive output in *P. antipodarum* ([36]; reviewed in [37]). Neiman

[38] speculated that asexual females that have no direct use for sperm contributed by males

might still benefit from mating if copulatory stimuli are required for maximization of

reproduction, though Neiman [39], showed that this scenario is unlikely for *P. antipodarum*.

Alternatively, male ejaculate may provide nutritive benefits to asexual females. This possibility

is supported by the fact that sperm storage structures are maintained in asexual *P. antipodarum*

females [40]; though evidence that these structures might also function for the digestion of waste

in asexual *P. antipodarum* suggests the potential for a non-mating-specific or even unique

function for this organ in asexuals [41]. Follow-up studies of whether and how sperm/ejaculate

are used in asexual *P. antipodarum* and other asexual taxa will help clarify the prevalence and

persistence of copulatory behaviour in asexual females.

Author Contributions

SS, JJ, and MN conceived of and designed the experiment. SS executed the experiment and collected the data. SS and MN analysed the data. SS created figures and drafted the manuscript. JJ and MN revised the manuscript. All authors gave final approval for publication and agree to be held accountable for the work performed therein.

Acknowledgments

We thank the Iowa Center for Research by Undergraduates for contributing to mentorship and training for SS, Dr. Lori Adams for her input and guidance through the Honors Program for SS; John Logsdon, Mike Winterbourn, and Mary Morgan-Richards for help in snail collection; and Marissa Roseman for setting up, characterizing, and maintaining the snail lineages that we used. Josephine Bliss also contributed to snail care.

Funding

SS was supported by the Iowa Center for Research by Undergraduates and by Linda and Rick Maxson. NSF grant DEB-1753851 provided support for snail maintenance and care.

Figure Legends

Figure 1. One *P. estuarinus* male (A), one *P. antipodarum* male (B), and one sexual or asexual *P. antipodarum* female (C) were placed approximately equidistant from one another around the edges of the petri dish. Each female (C) was painted a light colour with nail polish.

Figure 2. Violin plots of mating frequency (A, B) and duration (C, D) of sexual (A, C; "SexF") and asexual (B, D; "AsexF") females with *P. antipodarum* ("PAM") and *P. estuarinus* ("PEM") males. Median values are indicated with black dots.

Literature Cited

1. Darwin C. 1859 *The Origin of Species by Means of Natural Selection or, the Preservation of Favoured Races in the Struggle for Life*. Oxford, UK: Oxford University Press.

2. Wiedersheim R. 1895 *The Structure of Man: an Index to His Past History*. London, UK: Macmillan and Company.

3. Van der Kooi CJ, Schwander T. 2014 On the fate of sexual traits under asexuality. *Biol. Rev.* **89**, 805-819. (doi:10.1111/brv.12078)

4. Lande R. 1978 Evolutionary mechanisms of limb loss in tetrapods. *Evolution.* **32**, 73-92. (doi:10.2307/2407411)

5. Jeffery WR. 2005 Adaptive evolution of eye degeneration in the Mexican blind cavefish. *J. Hered.* **96**, 185-196. (doi:101093/jhered/esi028)

6. Roff DA. 1990 The evolution of flightlessness in insects. *Ecol. Monog.* **60**, 389-421. (doi:10.2307/1943013)

7. Fong DW, Kane TC, Culver DC. 1995 Vestigialization and loss of nonfunctional characters. *Annu. Rev. Ecol. Syst.* **26**, 249-268. (doi:10.1146/annurev.es.26.110195.001341)

8. Lahti DC, Johnson NA, Ajie BC, Otto SP, Hendry AP, Blumstein DT, Coss RG, Donohue K, Foster SA. 2009 Relaxed selection in the wild. *Trends Ecol. Evol.* **24**, 487-496. (doi:10.1016/j.tree.2009.03.010)

9. Williams GC. 1966 Natural selection, the costs of reproduction, and a refinement of Lack's principle. *Am. Nat.* **100**, 687-690. (doi:10.1086/282461)

10. Magnhagen C. 1991 Predation risk as a cost of reproduction. *Trends Ecol. Evol.* **6**, 183-186. (doi: 10.1016/0169-5347(91)90210-O)

11. Carson HL, Chang LS, Lyttle TW. 1982 Decay of female sexual behavior under parthenogenesis. *Science*. **218**, 68-70. (doi:10.1126/science.218.4567.68)

12. Kampfraath AA, Dudink TP, Kraaijeveld K, Ellers J, Zizzari ZV. 2020 Male sexual trait decay in two asexual springtail populations follows neutral mutation accumulation Theory. *Evol. Biol.* **47**, 285-292. (doi:10.1007/s11692-020-09511-z)

13. Schwander T, Crespi BJ, Gries R, Gries G. 2013 Neutral and selection-driven decay of sexual traits in asexual stick insects. *Proc. R. Soc. Lond. B* **280**, 20130823. (doi:10.1098/rspb.2013.0823)

14. Kraaijeveld K, Anvar SY, Frank J, Schmitz A, Bast J, Wilbrandt J, Petersen M, Ziesmann T, Niehuis O, De Knijff P, *et al*. 2016 Decay of sexual trait genes in an asexual parasitoid wasp. *Genome Biol. Evol.* **8**, 3685-3695. (doi:10.1093/gbe/evw273)

15. Jalinsky J, Logsdon Jr JM, Neiman M. 2020 Male phenotypes in a female framework: Evidence for degeneration in sperm produced by male snails from asexual lineages. *J. Evol. Biol.* **33**, 1050-1059. (doi:10.1111/jeb.13632)

16. Zahavi A. 1975 Mate selection—a selection for a handicap. *J. Theor. Biol.* **53**, 205-214. (doi: 10.1016/0022-5193(75)90111-3)

17. Kirkpatrick M. 1982 Sexual selection and the evolution of female choice. *Evolution*. **36**, 1-12. (doi:10.2307/2407961)

18. Heywood JS. 1989 Sexual selection by the handicap mechanism. *Evolution*. **43**, 1387-1397. (doi:10.1111/j.1558-5646.1989.tb02590.x)

19. Kokko H, Brooks R, Jennions MD, Morley J. 2003 The evolution of mate choice and mating biases. *Proc. R. Soc. Lond. B* **270**, 653-664. (doi:10.1098/rspb.2002.2235)

20. Wallace C. 1992 Parthenogenesis, sex and chromosomes in *Potamopyrgus*. *J. Molluscan Stud*. **58**, 93-107. (doi:10.1093/mollus/58.2.93)

21. Dybdahl MF, Lively CM. 1995 Diverse, endemic and polyphyletic clones in mixed populations of a freshwater snail (*Potamopyrgus antipodarum*). *J. Evol. Biol*. **8**, 385-398. (doi:10.1046/j.1420-9101.1995.8030385.x)

22. Soper DM, Hatcher KM, Neiman M. 2016 Documentation of copulatory behaviour in triploid male freshwater snails. *Ethol. Ecol. Evol*. **28**, 110-116. (doi:10.1080/03949370.2015.1030781)

23. Neiman M, Lively CM. 2005 Male New Zealand mud snails (*Potamopyrgus antipodarum)* persist in copulating with asexual and parasitically castrated females. *Am. Midl. Nat*. **154**, 88-96. (doi:10.1674/0003-0031(2005)154[0088:MNZMSP]2.0.CO;2)

24. Nelson AE, Neiman M. 2011 Persistent copulation in asexual female *Potamopyrgus antipodarum*: evidence for male control with size-based preferences. *Int. J. Evol. Biol*. **2011**, 439046. (doi:10.4061/20110439046)

25. Haase M. 2003 Clinal variation in shell morphology of the freshwater gastropod *Potamopyrgus antipodarum* along two hill☐country streams in New Zealand. *J. R. Soc. NZ* **33**, 549-560. (doi: 10.1080/03014223.2003.9517743)

26. Stork, SE. 2019 Unpublished data.

27. Richards DC, Shinn DC. 2004 Intraspecific competition and development of size structure in the invasive snail *Potamopyrgus antipodarum* (Gray, 1853). *Am. Malacol. Bull*. **19**, 33-37.

28. McKenzie VJ, Hall WE, Guralnick RP. 2012 New Zealand mudsnails (*Potamopyrgus antipodarum*) in Boulder Creek, Colorado: environmental factors associated with fecundity of a parthenogenic invader. *Can. J. Zool*. **91**, 30-36. (doi:10.1139/cjz-2012-0183)

29. Haase M. 2008 The radiation of hydrobiid gastropods in New Zealand: a revision including the description of new species based on morphology and mtDNA sequence information. *System. Biodivers*. **6**, 99-159. (doi:10.1017/S1477200007002630)

30. CandyLabs. (2019). VideoVelocity Time-Lapse Capture Studio [Computer software]. Vancouver, Canada.

31. R Core Team. 2017 R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/

32. Villanueva RA, Chen ZJ, Wickham H. 2016 ggplot2: Elegant graphics for data analysis using the grammar of graphics

33. Kassambara A (2020). ggpubr: 'ggplot2' Based Publication Ready Plots. R package version 0.4.0. https://CRAN.R-project.org/package=ggpubr

34. Wickham H, François R, Henry L, Müller K (2021). dplyr: A Grammar of Data Manipulation. R package version 1.0.4. https://CRAN.R-project.org/package=dplyr

35. Neiman M, Jokela J, Lively CM. 2005 Variation in asexual lineage age in *Potamopyrgus antipodarum*, a New Zealand snail. *Evolution*. **59**, 1945-1952. (doi:10.1111/j.0014-3820.2005.tb01064.x)

36. Barnes H, Barnes M. 1977 Studies on the reproduction of cirripedes. I. Introduction: copulation, release of oocytes, and formation of the egg lamellae. *J. Exp. Mar. Biol. Ecol*. **27**, 195-218. (doi:10.1016/0022-0981(77)90124-1)

37. Eberhard WG, Cordero C. 1995 Sexual selection by cryptic female choice on male seminal products-a new bridge between sexual selection and reproductive physiology. *Trends Ecol. Evol*. **10**, 493-496. (doi:10.1016/S0169-5347(00)89205-8)

38. Neiman M. 2004 Physiological dependence on copulation in parthenogenetic females can reduce the cost of sex. *Anim. Behav*. **67**, 811-822. (doi:10.1016/j.anbehav.2003.05.014)

39. Neiman M. 2006 Embryo production in a parthenogenetic snail (*Potamopyrgus antipodarum*) is negatively affected by the presence of other parthenogenetic females. *Invertebr. Biol*. **125**, 45-50. (doi:10.1111/j.1744-7410.2006.00038.x)

40. Dillon RT. 2000 *The Ecology of Freshwater Molluscs*. Cambridge, UK: Cambridge University Press.

41. Fretter V, Graham A. 1962 *British Prosobranch Molluscs. Their Functional Anatomy and Ecology*. London, UK: Ray Society.