Asexual freshwater snails make poor mate choice decisions

Sydney Stork[1], Joseph Jalinsky[1], Maurine Neiman[1,2,*]

[1]Department of Biology, University of Iowa, Iowa City, IA, 52242, USA

[2]Department of Gender, Women's, and Sexuality Studies, University of Iowa, Iowa City, IA,

52242, USA.

*Corresponding author: maurine-neiman@uiowa.edu, 324 BB, Iowa City, IA, 52242, USA.

Abstract

Once-useful traits that no longer contribute to fitness tend to decay over time. Here, we address

whether the expression of mating-related traits that increase the fitness of sexually reproducing

individuals but are likely less useful or even costly to asexual counterparts seems to exhibit

decay in the latter. *Potamopyrgus antipodarum* is a New Zealand freshwater snail characterized

by repeated transitions from sexual to asexual reproduction. The frequent coexistence of sexual

and asexual lineages makes *P. antipodarum* an excellent model for the study of mating-related

trait loss. We used a mating choice assay including sexual and asexual *P. antipodarum* females

and conspecific (presumed better choice) vs. heterospecific (presumed worse choice) males to

evaluate the loss of behavioral traits related to sexual reproduction. We found that sexual females

engaged in mating behaviors with conspecific mating partners more frequently and for a greater

duration than with heterospecific mating partners. By contrast, asexual females mated at similar

frequency and duration as sexual females, but did not mate more often or for longer duration

with conspecific vs. heterospecific males. These results are consistent with a scenario where

selection acting to maintain mate choice in asexual *P. antipodarum* is weak or ineffective

relative to sexual females and, thus, where asexual reproduction contributes to the evolutionary

decay of mating-related traits in this system.

Introduction

Adaptations that were once important for survival but subsequently become useless or impose

new fitness costs tend to decay over time in a process known as vestigialization (Darwin 1859;

Wiedersheim 1895; Van der Kooi and Schwander 2014). Many examples of this phenomenon

can be seen throughout evolutionary history, including the loss of limbs in snakes (Lande 1978),

eye structures in cave-dwelling fish (Jeffery 2005), and wings in many insect species (Roff

1990).

While the genetic mechanisms underlying vestigialization depend on evolutionary

context, traits are most commonly lost by the accumulation of selectively neutral mutations

(Fong et al 1995; Lahti et al 2009). The decay of presently useless characteristics exemplifies

evolutionary tradeoffs: the decay of one trait frees up energy and resources to be used elsewhere

by the organism (Fong et al 1995). For example, Roff (1990) suggests that the evolution of

flightlessness from flighted ancestors allows for allocation of resources for reproduction. This

framework implies that trait loss could translate into increased fitness for organisms for which

the lost trait is no longer useful.

The decay of traits specific to mating and/or sexual reproduction (and above and beyond

the loss of fitness expected under the accumulation of harmful mutations expected under asexual

reproduction (Muller 1964; Hill and Robertson 1966)) is one potential consequence of transitions

from sexual to asexual reproduction. These traits are typically required for reproductive success

in sexual organisms. By definition, asexual organisms do not need external genetic contribution

to reproduce, with the exception of sperm-dependent forms of parthenogenesis (e.g.,

gynogenesis). Mating behaviors are themselves associated with numerous costs (Williams 1966;

Magnhagen 1991). Accordingly, vestigialization of mating-related traits in asexual lineages –

including those involved in mate choice – is expected, particularly in a scenario where such

behaviors and their maintenance are costly (Van der Kooi and Schwander 2014; Carson et al

1982; Kampfraath et al 2020). Alternatively, these traits may be selectively neutral or maintained

by selection in asexuals, but only in a context where the costs do not outweigh the benefits of

trait maintenance (Schwander et al 2013; Kraaijeveld et al 2016). Several studies have supported

these ideas, finding abnormal or fully decayed traits involved in mating and reproduction

(spermathecae in asexual *Timema* stick insects (Schwander et al 2013); sperm in asexual *P.*

antipodarum males (Jalinsky et al 2020)) in asexual organisms. These results suggest that these

traits are not being maintained by selection, at least to the extent of the same traits in sexual

counterparts. Our focus is on behaviors: under the assumptions that mate preferences are

associated with mate quality (Zahavi 1975; Kirkpatrick 1982; Heywood 1989), that mate quality

contributes to fitness of sexual but not asexual females, that the genes underlying mating

behaviors are not pleiotropic with respect to other traits also important to asexual females, and

that there is at most limited gene flow between sexual and asexual conspecifics, we ask whether

mating-related behaviors associated with mate preference decay in asexual females relative to

sexual counterparts.

Potamopyrgus antipodarum is a New Zealand freshwater snail characterized by the

frequent coexistence of phenotypically and ecologically similar obligately sexual and obligately

asexual lineages (Lively 1987; Wallace 1992), making this species an excellent model to study

the vestigialization of traits involved in sexual reproduction. Previous studies have shown that

mating behavior in sexually reproducing male *P. antipodarum* does not differ from that of the

presumed asexual males produced occasionally by asexual females (Soper et al 2016). While

gene flow mediated by these males is formally possible (Neiman et al 2011), separate lines of

evidence for distinct genomic consequences of asexuality in *P. antipodarum* (Sharbrough et al

2018; McElroy et al 2021) suggests that such gene flow is rare at most. Sexual males also show

poor ability to discriminate between favorable and unfavorable mates, mating with equal

frequency with sexual and asexual females and with parasitically castrated vs. healthy females

(Neiman and Lively 2005). Prior emphasis on investigation of male mating behavior is largely a

result of the assumption that males are physically in control of mating outcomes (Neiman and

Neiman 2011), despite a lack of studies assessing the female role. These previous results set the

stage for a direct comparison between sexual and asexual *P. antipodarum* females. In a mating

paradigm like that demonstrated for *P. antipodarum* where males seem to lack adequate mate

discrimination, differences in mating between sexual and asexual females can be attributed to

mating differences between these two female groups. Therefore, we hypothesize that sexual

female *P. antipodarum* are better able to choose a more favorable mate when presented with a

choice than asexual counterparts.

We evaluated this hypothesis by comparing the mating behavior of sexual and asexual *P.*

antipodarum females with sexual male *P. antipodarum* and sexual male *Potamopyrgus*

estuarinus. *Potamopyrgus estuarinus* is a closely related and phenotypically similar species that

nevertheless has minimal habitat overlap with *P. antipodarum* (Haase 2003). Sexual female *P.*

antipodarum mating efforts with conspecific males are presumed to be favorable relative to

mating with *P. estuarinus* because of the possibility to generate offspring only with the former.

Preference for *P. antipodarum* vs. *P. estuarinus* should be selectively neutral for asexual female

P. antipodarum, for which eggs develop in the absence of males and fertilization. We chose to

use *P. estuarinus* males as a heterospecific (poor) alternative choice in this experiment because

the two species are phenotypically similar and because mating behavior (but not offspring

production) has been shown to occur between male and female *P. antipodarum* and *P. estuarinus*

(Stork, unpublished), thereby providing *P. antipodarum* females with an adequate alternative

choice.

<u>Materials and Methods</u>

Snail Selection and Care

We haphazardly selected three sets of 10 sexually mature (>3 mm long; Richards and Shinn

2004; McKenzie et al 2012; Larkin et al 2016) sexual (N=30) and asexual (N=30) female *P.*

antipodarum descended from one sexual and one asexual female, respectively, collected from

Lake Mapourika, New Zealand, in 2018. These six sets of 10 female snails will hereafter be

referred to as 'populations.' We also haphazardly selected three sets of 20 each sexually mature

(possessing a visible penis, Haase 2008) sexual *P. antipodarum* males (N=60) and

Potamopyrgus estuarinus males (N=60). All *P. antipodarum* males were chosen from sexual

lineages established by females sampled from New Zealand Lakes Mapourika and Selfe in 2018.

These lineages were separately derived from the lineages from which we drew females. All *P.*

estuarinus males were collected from the Ashley River estuary in New Zealand in February,

2020.

All snails used in the experiment had been housed in 10-L plastic tanks filled about 75%

full with carbon-filtered tap water and fed *Spirulina* algae, a common *Potamopyrgus* lab food,

three times weekly. These tanks were maintained in our constant-temperature snail room, which

is held at 16°C and on a 12:12 hour light:dark cycle. Snails were given powdered chalk as a

source of dietary calcium once weekly (feeding and maintenance following Nelson and Neiman

(2011)). We isolated each population used for the experiment in a 0.95 L plastic cup containing

~700 mL of water for approximately 14 days prior to behavioral trials to control for previous

exposure to males and possible recent mating. We painted female shells with light-colored nail

polish to aid in differentiation between females and males when analyzing video footage.

Feeding and maintenance for the populations in cups followed our standard snail room protocol.

Mating trials occurred in standard four-inch petri dishes. We covered the exterior of each

dish in electrical tape to control for external visual stimuli. To begin each trial for the sexual

females, we placed one sexual *P. antipodarum* female, one sexual *P. antipodarum* male, and one

P. estuarinus male equidistant from one another around the edges of the inside of the petri dish

(Fig. 1). After the three snails were placed, we filled the petri dish to about 75% total volume

with carbon-filtered tap water. We used this same setup for asexual *P. antipodarum* females,

with the asexual female replacing the sexual female. No snail was used in more than one trial.

Observation of Mating Behavior

We conducted each mating trial at approximately 1500 hours, for three hours. As the maximum

copulation duration for *P. antipodarum* is about two hours (Neiman and Lively 2005), we

established this three-hour time frame to allow for multiple mating attempts to occur. We

recorded trials using a USB webcam and time-lapse video software VideoVelocity (v3.7.2090,

CandyLabs 2019). The three snails per trial were allowed to move and interact freely without

physical barriers inside the petri dish.

We loosely followed previous studies of mating in *P. antipodarum* (e.g., Nelson and

Neiman 2011) to define attempted mating behavior as any physical interaction resulting in one

snail mounting the shell of another. It is otherwise impossible to visually confirm sexual contact

(joining of genitalia) in *Potamopyrgus* because this phenomenon is internal relative to the

operculum and is thus obscured by the shells. Once this mounting behavior was initiated, we

arbitrarily established that any such interaction lasting 15 seconds or longer would be counted as

an instance of mate choice; interactions shorter than 15 seconds were potentially too short to be

discernible with our time-lapse software and were excluded. Fifteen seconds is also more than

enough time for snails to reorient and move on following an incidental – versus mating –

encounter. Both the frequency (number of mating interactions each female had with conspecific

vs. heterospecific males) and duration (length of time (seconds) that each female was in physical

contact with each male type) of mating interactions between females and conspecific vs.

heterospecific males were recorded.

We also used this experimental framework to establish expected outcomes and determine

how these outcomes would inform our hypotheses. First, a fair test of the trait decay hypothesis

would require that sexual females mate more frequently and for longer with conspecific males

than with *P. estuarinus* males. This outcome is critical in indicating that sexual females seem to

possess the ability to discriminate between preferential and poor mating partners in our

experimental setting. Absence of a difference in mating behavior across male type for sexual

females would not support the trait decay hypothesis, suggesting instead, for example, that these

females are unable to differentiate between mates or that observed contact is reflective of

behavior outside of mating. Mating-related trait decay in asexual females consistent with the trait

decay hypothesis would be reflected in relatively weak or absent mate choice for conspecific vs.

heterospecific males relative to choices in sexual female *P. antipodarum*.

Statistical Analyses

We used the Shapiro-Wilk test to determine that both the mating frequency (sexual females: $W =$

0.769, $p = 8.18 \times 10^{-8}$; asexual females: $W = 0.810$, $p = 1.12 \times 10^{-5}$) and the mating duration (sexual

females: $W = 0.642$, $p = 1.05 \times 10^{-9}$; asexual females: $W = 0.493$, $p = 3.17 \times 10^{-8}$) datasets violated

normality assumptions of parametric analysis. Accordingly, we used nonparametric approaches

to analyze the data. Analyses for mating duration were executed using the total mating duration

for each individual (Tables S1, S2). First, we used Mann-Whitney U tests to compare overall

mating frequency and duration between individual sexual and asexual females. Second, we used

Wilcoxon rank-sum analyses to determine whether mating durations and frequencies of sexual

and asexual females, respectively, with conspecific males differed from mating durations and

frequencies with heterospecific males. As another means of comparing mating behavior across

sexual and asexual females, we calculated discrimination scores - defined as the difference

between mating frequency and mating duration, respectively, with conspecific vs. heterospecific

males – for each female. We then used Mann-Whitney U tests to determine whether there was a

significant difference in the mating frequency and mating duration discrimination scores,

respectively, between sexual and asexual females. All analyses were executed with R 3.6.2 (R

Core Team 2017), the ggplot2 (v3.3.3; Villanueva et al 2016), ggpubr (v0.4.0; Kassambara

2020) and dplyr (v1.0.4; Wickham et al 2021) packages.

<u>Results</u>

Technical challenges with the recording software and camera hardware ultimately rendered data

from three sexual female and 10 asexual female trials impossible to extract, leaving us with trials

from 27 sexual females and from 20 asexual females. Comparison of the mating frequency and

duration of these sexual and asexual females revealed no differences between the two female

types for either trait (Frequency: Mann-Whitney $U = 288$, $z = -0.38$, $p = 0.704$; Duration: Mann-

Whitney $U = 187$, $z = 1.78$, $p = 0.075$). Wilcoxon rank-sum analyses demonstrated that sexual

females mated significantly more often with conspecific males (median = 1.0 matings, IQR =

2.0) than with heterospecific males (median = 1.0 matings, IQR = 1.0) ($W = 483$, $p = 0.030$) (Fig.

2a). By contrast, asexual females mated with conspecific males (median = 1.0 matings, IQR =

2.0) with the same frequency as with heterospecific males (median = 1.0, IQR = 2.5) ($W = 172$, p

= 0.427) (Fig. 2b).

Sexual females also mated for significantly longer with conspecific males (median =

242.72 seconds, IQR = 1447.82) than with heterospecific males (median = 22.76 seconds, IQR =

102.25) ($W = 520$, $p = 0.006$) (Fig. 2c). Asexual females did not differ in the duration of mating

with conspecific (median = 30.65 seconds, IQR = 377.43) and heterospecific males (median =

62.50 seconds, IQR = 163.84) ($W = 202$, $p = 0.967$) (Fig. 2d).

The discrimination score analysis revealed that sexual but not asexual females with a

choice between conspecific and heterospecific males tend to make mating behavior decisions

with respect to mating frequency that favor the former (Mann-Whitney $U = 175.5$, $z = 2.02$, $p =$

0.043). By contrast, there was no significant difference in discrimination scores for mating

duration between sexual and asexual females (Mann-Whitney $U = 208$, $z = 1.32$, $p = 0.187$).

Discussion

Our experiment revealed that sexual female *P. antipodarum* exhibited significantly higher

mating frequency and longer mating duration with conspecific vs. heterospecific males. These

results are as predicted if sexual female *P. antipodarum* are able to discriminate between mates

of higher vs. lower quality and are important in demonstrating that discriminatory behavior can

occur in the context of our experiment. By contrast, although asexual *P. antipodarum* females

mated at the same frequency and for the same duration as their sexual counterparts, the asexuals

showed no evidence for mate discrimination or preference. Together, these findings are

consistent with a scenario where sexual but not asexual female *P. antipodarum* exhibit mate

choice. Evolutionary decay of mating-related traits in an asexual context is a plausible

explanation for these results and is consistent with a previous study demonstrating evidence for

decay of sperm traits in the male offspring occasionally produced by asexual female *P.*

antipodarum (Jalinsky et al 2020). Another non-mutually exclusive explanation for our results is

provided by the possibility that asexual female *P. antipodarum* are experiencing more general

degradation of phenotype associated with the accumulation of harmful mutations that is expected

to accompany obligately asexual reproduction (Muller 1964; Hill and Robertson 1966). While a

relatively high rate of accumulation of potentially harmful mutations has been observed in

asexual *P. antipodarum* relative to sexual conspecifics (e.g., Sharbrough et al 2018), that asexual

P. antipodarum females are similar to or even outperform sexual females when it comes to key

determinants of fitness like fecundity (Paczesniak et al 2019) and growth rate and age at

reproductive maturity (Larkin et al 2016) suggests that these mutations might not be translating

into major phenotypic consequences. We also cannot formally exclude the possibility that

asexual female snails might demonstrate mating preferences in a different context, though the

fact that we observed such preferences in closely related sexual counterparts in the same

experiment suggests that this explanation is not likely to entirely explain the different outcomes

between sexual and asexual females. Finally, it is important to acknowledge that our study only

included female *P. antipodarum* from one lake and that we only compared one sexual lineage to

one asexual lineage. This Covid19-imposed limitation on our experimental design means that

when possible, it will be important to repeat the experiment using additional lineages to

determine whether our results are indeed more broadly generalizable.

The exhibition of similar mating behavior to their sexual counterparts combined with the

lack of observable discriminatory behavior in asexual females could be explained by a situation

where copulatory behavior – even in the absence of a connection between egg fertilization and

embryo development – is still necessary and/or beneficial although *choice* of the mating partner

is no longer important. For example, Neiman (2004) speculated that asexual females that have no

direct use for sperm contributed by males might still benefit from mating if copulatory stimuli

are required for maximization of reproduction, though Neiman (2006), showed that this scenario

is unlikely for *P. antipodarum*. Alternatively, male ejaculate may provide nutritive benefits to

asexual females. This possibility is supported by the fact that sperm storage structures are

maintained in asexual *P. antipodarum* females (Dillon 2000), though evidence that these

structures might also function for the digestion of waste in asexual *P. antipodarum* suggests the

potential for a non-mating-specific or even unique function for this organ in asexuals (Fretter and

Graham 1962). Follow-up studies of whether and how sperm/ejaculate are used in asexual *P.*

antipodarum and other asexual taxa will help clarify the prevalence and persistence of

copulatory behavior in asexual females.

Another and perhaps simpler explanation is that mating behavior has not fully decayed in

P. antipodarum, and that the ability to discriminate is lost before the behavior as a whole

disappears. That asexual females nevertheless engage in mating behavior at all could be a

function of the relatively recent derivation of most asexual female *P. antipodarum* from sexual

counterparts (Neiman et al 2005). With this in mind, it would be valuable to repeat this

experiment with females from older asexual lineages (*sensu* Nelson and Neiman 2011) but

departing from this prior study so that females vs. males have mate choice capabilities.

Acknowledgments

We thank the Iowa Center for Research by Undergraduates for contributing to mentorship and training for SS, Dr. Lori Adams for her input and guidance through the Honors Program for SS; John Logsdon, Mike Winterbourn, and Mary Morgan-Richards for help in snail collection; and Marissa Roseman for setting up, characterizing, and maintaining the snail lineages that we used. Josephine Bliss also contributed to snail care. This work was supported by the Iowa Center for Research by Undergraduates, by Linda and Rick Maxson, and via NSF grant DEB-1753851. We also acknowledge very constructive and helpful critiques from Prof. Ken Kraajiveld, Prof. Ingo Schlupp, and an anonymous reviewer that greatly improved the paper.

Declarations

Funding: This work was supported by the Iowa Center for Research by Undergraduates, by Linda and Rick Maxson, and via NSF grant DEB-1753851.

Conflicts of interest: Not applicable.

Ethics approval: Not applicable.

Consent to participate: Not applicable.

Consent for publication: All authors gave final approval for publication and agree to be held accountable for the work performed therein.

Availability of data and material: All data are available in the appendices. Snail videos are too large to readily post online but are available upon request from the authors.

Code availability: Not applicable.

Authors' contributions: Sydney Stork, Joseph Jalinsky, and Maurine Neiman conceived of and designed the experiment. Sydney Stork executed the experiment and collected the data. Sydney

Stork and Maurine Neiman analyzed the data. Sydney Stork created figures and drafted the manuscript. Sydney Stork, Joseph Jalinsky, and Maurine Neiman revised the manuscript.

References

Barnes H, Barnes M (1977) Studies on the reproduction of cirripedes. I. Introduction: copulation, release of oocytes, and formation of the egg lamellae. J Exp Mar Biol Ecol 27:195-218.

CandyLabs. (2019) VideoVelocity Time-Lapse Capture Studio [Computer software]. Vancouver, Canada.

Carson HL, Chang LS, Lyttle TW (1982) Decay of female sexual behavior under parthenogenesis. Science 218:68-70.

Darwin C (1859) The origin of species by means of natural selection or, the preservation of favoured races in the struggle for life. Oxford University Press. Oxford.

Dillon RT (2000) The ecology of freshwater molluscs. Cambridge University Press, Cambridge.

Eberhard WG, Cordero C (1995) Sexual selection by cryptic female choice on male seminal products-a new bridge between sexual selection and reproductive physiology. Trends Ecol Evol 10:493-496.

Fretter V, Graham A (1962) British prosobranch molluscs. Their functional anatomy and ecology. Ray Society, London.

Fong DW, Kane TC, Culver DC (1995) Vestigialization and loss of nonfunctional characters. Annu Rev Ecol Syst 26:249-268.

Haase M (2003) Clinal variation in shell morphology of the freshwater gastropod *Potamopyrgus antipodarum* along two hill-country streams in New Zealand. J R Soc NZ 33:549-560.

Haase M (2008) The radiation of hydrobiid gastropods in New Zealand: a revision including the description of new species based on morphology and mtDNA sequence information. System Biodivers 6:99-159.

Heywood JS (1989) Sexual selection by the handicap mechanism. Evolution 43: 1387-1397.

Hill WG, Robertson A (1966) The effect of linkage on limits to artificial selection. Genet Res 8:269-294.

Jalinsky J, Logsdon Jr JM, Neiman M (2020) Male phenotypes in a female framework: Evidence for degeneration in sperm produced by male snails from asexual lineages. J Evol Biol 33:1050-1059.

Jeffery WR (2005) Adaptive evolution of eye degeneration in the Mexican blind cavefish. J Hered 96:185-196.

Kampfraath AA, Dudink TP, Kraaijeveld K, Ellers J, Zizzari ZV (2020) Male sexual trait decay in two asexual springtail populations follows neutral mutation accumulation Theory Evol Biol 47:285-292.

Kassambara A (2020) ggpubr: 'ggplot2' Based Publication Ready Plots. R package version 0.4.0. https://CRAN.R-project.org/package=ggpubr.

Kirkpatrick M (1982) Sexual selection and the evolution of female choice. Evolution 36:1-12.

Kokko H, Brooks R, Jennions MD, Morley J (2003) The evolution of mate choice and mating biases. Proc R Soc Lond B 270:653-664.

Kraaijeveld K, Anvar SY, Frank J, Schmitz A, Bast J, Wilbrandt J, Petersen M, Ziesmann T, Niehuis O, De Knijff P, den Dunnen JT, Ellers J (2016) Decay of sexual trait genes in an asexual parasitoid wasp. Genome Biol Evol 8:3685-3695.

Lahti DC, Johnson NA, Ajie BC, Otto SP, Hendry AP, Blumstein DT, Coss RG, Donohue K, Foster SA (2009) Relaxed selection in the wild. Trends Ecol Evol 24:487-496.

Larkin K, Tucci, C, Neiman M (2016) Effects of polyploidy and reproductive mode on life history trait expression. Ecology and Evolution 6:765-778.

Lande R (1978) Evolutionary mechanisms of limb loss in tetrapods. Evolution 32:73-92.

Lively CM (1987) Evidence from a New Zealand snail for the maintenance of sex by parasitism. Nature 328:519-521.

Magnhagen C (1991) Predation risk as a cost of reproduction. Trends Ecol Evol 6:183-186.

McElroy KE, Müller S, Lamatsch DK, Bankers L, Fields PD, Jalinsky JR, Sharbrough J, Boore JL, Logsdon JM Jr, Neiman M (2021) Asexuality associated with marked genetic expansion of tandemly repeated rDNA and histone genes. Mol Biol Evol msab121: https://doi.org/10.1093/molbev/msab1121.

McKenzie VJ, Hall WE, Guralnick RP (2012) New Zealand mudsnails (*Potamopyrgus antipodarum*) in Boulder Creek, Colorado: environmental factors associated with fecundity of a parthenogenic invader. Can J Zool 91:30-36.

Muller HJ (1964) The relation of recombination to mutational advance. Mut Res 1:2-9.

Neiman M (2004) Physiological dependence on copulation in parthenogenetic females can reduce the cost of sex. Anim Behav 67:811-822.

Neiman M (2006) Embryo production in a parthenogenetic snail (*Potamopyrgus antipodarum*) is negatively affected by the presence of other parthenogenetic females. Invertebr Biol 125:45-50.

Neiman M, Lively CM. 2005 Male New Zealand mud snails (*Potamopyrgus antipodarum)* persist in copulating with asexual and parasitically castrated females. Am Midl Nat 154:88-96.

Neiman M, Jokela J, Lively CM (2005) Variation in asexual lineage age in *Potamopyrgus antipodarum*, a New Zealand snail. Evolution 59:1945-1952.

Neiman M, Paczesniak D, Soper DM, Baldwin AT, Hehman G (2011) Wide variation in ploidy level and genome size in a New Zealand freshwater snail with coexisting sexual and asexual lineages. Evolution 65:3202-3216.

Nelson AE, Neiman M (2011) Persistent copulation in asexual female *Potamopyrgus antipodarum*: evidence for male control with size-based preferences. Int J Evol Biol 2011: 439046.

Paczesniak D, Klappert K, Kopp K, Neiman M, Seppälä K, Lively CM, Jokela J (2019) Parasite resistance predicts fitness better than fecundity in a natural population of the freshwater snail *Potamopyrgus antipodarum*. Evolution 73:1634-1646.

R Core Team (2017) R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing. https://www.R-project.org/.

Richards DC, Shinn DC (2004) Intraspecific competition and development of size structure in the invasive snail *Potamopyrgus antipodarum* (Gray, 1853). Am Malacol Bull 19:33-37.

Roff DA (1990) The evolution of flightlessness in insects. Eco Monogr 60:389-421.

Schwander T, Crespi BJ, Gries R, Gries G (2013) Neutral and selection-driven decay of sexual traits in asexual stick insects. Proc R Soc Lond B 280:20130823.

Sharbrough J, Luse M, Boore JL, Logsdon JM Jr, Neiman M (2018) Radical amino acid mutations persist longer in the absence of sex. Evolution 72:808-824.

Soper DM, Hatcher KM, Neiman M (2016) Documentation of copulatory behavior in triploid male freshwater snails. Ethol Ecol Evol 28:110-116.

Van der Kooi CJ, Schwander T (2014) On the fate of sexual traits under asexuality. Biol Rev 89: 805-819.

Villanueva RA, Chen ZJ, Wickham H (2016) ggplot2: Elegant graphics for data analysis using the grammar of graphics.

Wallace C (1992) Parthenogenesis, sex and chromosomes in *Potamopyrgus*. J Molluscan Stud 58:93-107.

Wickham H, François R, Henry L, Müller K (2021) dplyr: A Grammar of Data Manipulation. R package version 1.0.4. https://CRAN.R-project.org/package=dplyr.

Wiedersheim R (1895) *The Structure of Man: an Index to His Past History*. London, UK: Macmillan and Company.

Williams GC (1966) Natural selection, the costs of reproduction, and a refinement of Lack's principle. Am Nat 100:687-690.

Zahavi A (1975) Mate selection—a selection for a handicap. J Theor Biol 53:205-214.

Figure Legends

Figure 1. One *P. estuarinus* male (A), one *P. antipodarum* male (B), and one sexual or asexual *P. antipodarum* female (C) were placed approximately equidistant from one another around the edges of the petri dish. Each female (C) was painted a light colour with nail polish.

Figure 2. Violin plots of mating frequency (a, b) and total mating duration (c, d) of sexual (a, c; "SexF") and asexual (b, d; "AsexF") females with *P. antipodarum* ("PAM") and *P. estuarinus* ("PEM") males. Individual data points are indicated with black dots.