TITLE:

The rapid developmental rise of somatic inhibition disengages hippocampal dynamics from self-motion

AUTHORS:

Robin F. Dard[1], Erwan Leprince[1], Julien Denis[1], Shrisha Rao-Balappa[2], Dmitrii Suchkov[1], Richard Boyce[1], Catherine Lopez[1], Marie Giorgi-Kurz[1], Tom Szwagier[1,3], Theo Dumont[1,3], Hervé Rouault[2], Marat Minlebaev[1], Agnes Baude[1], Rosa Cossart[1*], Michel A. Picardo[1*]

[1]Aix Marseille Univ, INSERM, INMED (U1249), Turing Centre for Living systems, Marseille, France

[2]Aix-Marseille Univ, Université de Toulon, CNRS, CPT (UMR 7332), Turing Centre for Living systems, Marseille, France

[3] Mines ParisTech, PSL Research University

* Correspondence: michel.picardo@inserm.fr and rosa.cossart@inserm.fr

ABSTRACT:

Early electrophysiological brain oscillations recorded in preterm babies and newborn rodents are initially mostly ignited by bottom-up sensorimotor activity and only later can detach from external inputs. This is a hallmark of most developing brain areas including the hippocampus, which in the adult brain, functions in integrating external inputs onto internal dynamics. Such developmental disengagement from external inputs is likely a fundamental step for the proper development of cognitive internal models. Despite its importance, the developmental timeline and circuit basis for this disengagement remain unknown. To address this issue, we have investigated the daily evolution of CA1 dynamics and underlying circuits during the first two postnatal weeks of mouse development using two-photon calcium imaging in non-anesthetized pups. We show that the first postnatal week ends with an abrupt shift in the representation of self-motion in CA1. Indeed, most CA1 pyramidal cells switch from activated to inhibited by self-generated movements at the end of the first postnatal week whereas the majority of GABAergic neurons remain positively modulated throughout this period. This rapid switch occurs within two days and follows the rapid anatomical and functional surge of local somatic GABAergic innervation. The observed change in dynamics is consistent with a two-population model undergoing a strengthening of inhibition. We propose that this abrupt developmental transition inaugurates the emergence of internal cognitive models.

INTRODUCTION

The adult hippocampus serves multiple cognitive functions including navigation and memory. These functions rely on the ability of hippocampal circuits to integrate external inputs conveying multi-sensory, proprioceptive, contextual and emotional information onto internally-generated dynamics. Therefore, the capacity to produce internally coordinated neuronal activity detached from environmental inputs is central to the cognitive functions of the hippocampus such as planning and memory (Buzsáki, 2015; Buzsáki and Moser, 2013). In contrast to the adult situation, the developing hippocampus, like many developing cortical structures, is mainly driven by bottom-up external environmental and body-derived signals, including motor twitches generated in the spinal cord and/or the brainstem (Dooley et al., 2020; Inácio et al., 2016; Karlsson et al., 2006; Mohns and Blumberg, 2010; Rio-Bermudez et al., 2020; Valeeva et al., 2019a). These produce early sharp waves conveyed by inputs from the entorhinal cortex (Valeeva et al., 2019a). The emergence of self-organized sequences without reliance on external cues in the form of sharp wave ripples (SWRs) is only observed after the end of the second postnatal week and sequential reactivations even a week later (Farooq and Dragoi, 2019; Muessig et al., 2019). Therefore, early hippocampal activity as measured with electrophysiological recordings is first externally-driven while the emergence of internal dynamics is protracted. The timing and the circuit mechanisms of the switch between motion-guided and internally produced hippocampal dynamics remain unknown. They have been proposed to rely on the maturation of CA3 and extrinsic hippocampal inputs, however, a possible role of local connectivity, in particular recurrent somatic inhibition cannot be excluded (Cossart and Khazipov, 2021a).

Local GABAergic interneurons could be critically involved in this phenomenon for several reasons. First, both theoretical and experimental work suggest that self-organized internal neuronal network dynamics require feedback connections to produce an emergent state of activity independently from the incoming input (Hopfield and Tank, 2005; Hopfield, 1982; Yuste, 2015). Feedback circuits are mainly GABAergic in CA1 (but not necessarily inhibitory), given the scarcity of recurrent glutamatergic connections in that hippocampal sub-region (Bezaire and Soltesz, 2013). Second, GABAergic interneurons, in particular the perisomatic subtypes, are long known to shape the spatial and temporal organization of internal CA1 dynamics (Buzsáki, 2015; Lee et al., 2014; Soltesz and Losonczy, 2018; Valero et al., 2015). However, GABAergic perisomatic cells display a delayed maturation profile both at structural (Jiang et al., 2001; Marty et al., 2002; Morozov and Freund, 2003; Tyzio et al., 1999) and functional levels (Ben-Ari, 2002; Doischer et al., 2008; Jiang et al., 2001; Khazipov et al., 2004; Marty et al., 2002; Morozov and Freund, 2003; Murata and Colonnese, 2020; Tyzio et al., 1999), and the precise developmental timeline for their postnatal development remains unknown, partly due to the difficulty in labeling them (Donato et al., 2017) .

Here we investigate the evolution of CA1 dynamics during the first and second postnatal weeks of
mouse development with an eye on the specific patterning of activity of CA1 GABAergic neurons. To
this aim, we adapted two-photon calcium imaging of CA1 dynamics using virally-expressed GCaMP6
through a cranial window in non-anesthetized pups. We show that the first postnatal week ends with an
abrupt switch in the representation of self-motion in CA1: principal neurons were synchronized by
spontaneous movement before P9, whereas self-motion decreased their activity after that time point.
Consistent with a two-population neuronal model, this switch was locally paralleled by the rapid
anatomical and functional surge of somatic GABAergic interneurons and no significant change in
external inputs. Self-generated bottom-up inputs may thus directly contribute to the emergence of
somatic GABAergic inhibition and in this way calibrate local circuits to the magnitude of external inputs
prior to the opening of experience-dependent plasticity.

RESULTS

Progressive evolution of CA1 neuronal dynamics

In order to induce stable and early expression of the calcium indicator protein GCaMP6s, pups were
injected with the AAV1-hSyn-GCaMP6s.WPRE.SV40 virus in the brain lateral ventricle on the day of
birth (P0, **Figure 1A and Figure 1 - figure supplement 1A**). Five to twelve days after injection, the
hippocampal CA1 region of non-anesthetized pups was imaged through a window implant placed on
the same day (**Figure 1A, see methods**). This acute window implant did not alter hippocampal
dynamics as revealed by *in vivo* bilateral silicon probes electrophysiological recordings of early sharp
waves (eSW, **Figure 1 - figure supplement 1B**) in P6-8 and P11 pups (N=4 and N=2 respectively)
expressing GCaMP6s (with a frequency of 2.6 eSW/min (25 % 1.15 eSW/min and 75 % 4.16 eSW/min)
for the ipsilateral side and 3.49 eSW/min (25 % 1.96 eSW/min and 75 % 5.1 eSW/min, *p-value* = 0.39)
for the contralateral side (**Figure 1 - figure supplement 1B**). eSW synchronization between
hemispheres was also preserved (peak at 0.087 +/- 0.027, **Figure 1 - figure supplement 1B**) which is
in agreement with previous data (Graf et al., 2021; Valeeva et al., 2019b). Finally, we checked for the
presence of other types of oscillations in both hemispheres and observed a peak in the theta range in the
P11 mouse pups in both hemispheres (Ipsi: peak at 4.3 Hz of amplitude $3.6\text{x}10^{-3}$ +/- $1.2\text{x}10^{-3}$ mV^2/Hz;
Contra: peak at 4.1 Hz of amplitude $2\text{x}10^{-3}$ +/- $3\text{x}10^{-4}$ mV^2/Hz (jackknife std)). Therefore, the presence
of the window implant preserved the previously described network activity of the early postnatal period
in developing rodents. Thus, we pursued the description of early multineuron CA1 dynamics using
calcium imaging (62 imaging sessions, 35 mouse pups aged between 5 and 12 days, yielding a total of
33,412 cells, see **Table 1** for details on each session and their inclusion in the figures).

Age	SubjectID	SessionID	N_Cells	Main Figures												Supplementary Figures					
				1A	1B	1C	2A	2B	2C	2D	3A	3B	4C	4D	5C	1S1A	2S1A	2S1B	3S2A	3S2B	3S2C
5	190320_190325	190325_a000	633	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		190325_a001	919	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
5	191205_191210_0	191210_a000	316	Y	Y	Y	Y	Y	Y	Y	Y*	N	N	N	Y	Y	Y	N	Y	Y	Y
		191210_a001	128	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
5	191205_191210_1	191210_a000	512	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
		191210_a001	117	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
5	200306_200311	200311_a000	388	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	Y	N	Y	Y
		200311_a001	272	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	Y	Y	Y
6	180201_180207	180207_a001	566	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		180207_a002	405	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
6	190921_190927_1	190927_a000	240	N	Y	Y	Y	Y	Y	N	Y	N	N	N	N	Y	Y	N	N	Y	Y
6	191205_191211_1	191211_a000	566	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	Y
		191211_a001	395	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
7	180201_180208	180208_a000	515	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		180208_a001	846	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		180208_a002	742	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		180208_a003	622	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
7	190226_190305	190305_a000	410	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
7	190320_190327	190327_a000	824	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		190327_a001	678	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		190327_a002	655	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
7	200103_200110_0	200110_a000	453	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
		200110_a001	486	Y	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
7	200206_200213	200213_a000	611	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	Y
		200213_a001	322	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	N	N	Y	Y
8	181009_181017	181017_a000	627	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
		181017_a001	588	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
8	181016_181024	181024_a005	565	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	N	Y	N
8	190921_190929_1	190929_a000	698	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
8	191205_191213	191213_a000	712	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	N	N	Y	Y
		191213_a001	297	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	N	N	Y	Y
8	191212_191220	191220_a001	537	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
		191220_a003	620	N	Y	Y	Y	Y	Y	Y	Y	N	N	N	Y	Y	Y	N	Y	Y	Y
9	171211_171220	171220_a001	372	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
9	190211_190220_0	190220_a000	378	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
		190220_a001	942	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
		190220_a002	813	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
		190220_a003	797	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
9	190305_190314	190314_a000	443	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
		190314_a001	500	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
9	190313_190322	190322_a000	844	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
		190322_a001	696	N	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	N	Y	N
9	190921_190930	190930_a001	1050	N	Y	Y	Y	Y	Y	Y	N	N	N	N	N	Y	Y	N	N	Y	Y
		190930_a003	849	N	Y	Y	Y	Y	Y	Y	N	N	N	N	N	Y	Y	N	N	Y	Y
9	200108_200117_1	200117_a001	34	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N
9	210226_210307_1	210307_a000	36	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	N	N
10	190211_190221_0	190221_a003	607	N	Y	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N
		190221_a005	928	N	Y	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N
10	190226_190308	190308_a000	546	N	Y	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N
		190308_a001	596	N	Y	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N
10	190921_191001	191001_a000	486	Y	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	Y	Y	Y
10	210226_210308_1	210308_a000	34	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N
11	190211_190222	190222_a000	606	N	Y	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N
11	190921_191002	191002_a001	567	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	Y	Y	Y
12	171029_171110	171110_a000	404	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	N	Y	N
		171110_a002	654	N	Y	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N
12	190828_190909	190909_a000	396	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	N	Y	Y	N	N	Y	N

		190909_a001	430	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	N	Y	Y	N	N	Y	N
12	190911_190923	190923_a001	413	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	N	Y	N
12	190921_191003	191003_a001	294	N	Y	Y	Y	Y	Y	Y	N	Y	N	N	N	Y	Y	N	N	Y	Y
12	191122_191204	191204_a000	782	N	Y	Y	Y	Y	Y	Y	N	Y*	N	N	Y	Y	Y	N	Y	Y	Y
		191204_a001	650	Y	Y	Y	Y	Y	Y	Y	N	Y	N	N	Y	Y	Y	N	Y	Y	Y

Table 1: Details on the 62 imaging sessions from the 35 mouse pups showing the number of cells recorded in each session and how they were used in the figures. In main figure 3A, * shows mouse pups that are used for illustration. Y: included in the panel, N: not included.

The contours of the imaged neurons and their calcium fluorescence events were extracted using Suite2P (Pachitariu et al., 2017) and DeepCINAC (Denis et al., 2020), respectively. Representative examples of fields of view, contour maps and activity raster plots from recordings in P5, P7, P10 and P12 mouse pups are shown in Figure 1A. Consistent with previous electrophysiological studies (Leinekugel et al., 2002; Mohns and Blumberg, 2008; Valeeva et al., 2019a), spontaneous neuronal activity in the CA1 region of P5-6 pups alternated between recurring population bursts (Synchronous Calcium Events, SCEs), and periods of low activity (**Figure 1A**). In P5-6 mouse pups more than half of the detected calcium transients occurred within SCEs (P5: median value 55 % N=4, n=8; P6: median value 55 % N=3, n=5; N: mice, n: imaging sessions, **Figure 1B**). Activity then became progressively continuous as evidenced by the linear decrease in the proportion of calcium transients occurring during SCEs (r^2=0.78, p<0.0001) to finally reach 33 % in P12 mouse pups (P12: N=5, n=8, **Figure 1B**). These changes in CA1 dynamics cannot be explained by a change in the activation rate of individual neurons as this parameter did not significatively change during the developmental period studied here (**Figure 1C**). Even though there was globally no significant change in the rate of neuronal activation as measured with calcium imaging from P5 to P12, we observed a small, but not significant decrease at P9-10 consistent with the description of a transient period of 'neural quiescence' at the beginning of the second postnatal week (Dominguez et al., 2021). We conclude that CA1 dynamics progressively evolve from discontinuous to continuous during the first two postnatal weeks, in agreement with previous electrophysiological studies (Cossart and Khazipov, 2021b; Mohns and Blumberg, 2008; Valeeva et al., 2019a).

Figure 1: Evolution of CA1 dynamics during the first two postnatal weeks

(A). Schematic of the experimental timeline. On postnatal day 0 (P0), 2µL of a non-diluted viral solution were injected in the left lateral ventricle of mouse pups. From five to twelve days after injection (P5-P12), acute surgery for window implantation above the corpus callosum was performed and followed by 2-photon calcium imaging recordings. Top panel: Four example recordings are shown to illustrate the imaging fields of view in the stratum pyramidale of the CA1 region of the hippocampus (scale bar: 100 µm). Middle panel: Contour maps showing the cells detected using Suite2p in the corresponding fields of view. Bottom panel: Raster plots inferred by DeepCINAC activity classifier, showing 300

randomly selected cells over the first 5 minutes of recording obtained for these imaging sessions (P5,
P7, P10, P12, see full raster plots for these imaging sessions in Figure 1 - figure supplement 1D). In the
raster plot from the P5 mouse the blue rectangle illustrates one synchronous calcium event (SCE). Scale
bar for time is 1 min. CICADA configuration files to reproduce example rater plots and cell contours
are available in Figure 1 - Source Data 1. **(B).** Evolution of the ratio of calcium transients within SCEs
over the total number of transients across age. Each dot represents a mouse pup and is color coded from
light gray (P5) to black (P12), the open blue circles represent the median of the age group. The red line
represents the linear fit of the data with r^2=0.78 and p<0.0001. Results to build the distribution as well
as CICADA configuration file to reproduce the analysis are available in Figure 1 - Source Data 1. **(C).**
Evolution of the number of transients per minute across the first two postnatal weeks. Each dot
represents the mean transient frequency from all cells imaged in one animal and is color coded from
light gray (P5) to black (P12). The open blue circles represent the median of the age group. No
significant influence of the age was found (Kruskal-Wallis test, 8 groups, KW stat =12.63, p-value =
0.0816). Results to build the distribution as well as CICADA configuration file to reproduce the analysis
are available in Figure 1 - Source Data 1.

Early synchronous calcium events correlate with spontaneous motor activity

Previous extracellular electrophysiological recordings indicated that, in developing rodents, CA1

dynamics followed spontaneous motor activity during the first postnatal week (Karlsson et al., 2006;

Rio-Bermudez et al., 2020; Valeeva et al., 2019a). Hence, we next examined the relationship between

population activity and movement as monitored using either piezo recordings or infrared cameras (see

methods). We first computed peri-movement time histograms (PMTHs) plotting the fraction of active

neurons centered on the onset of all movements. In mouse pups younger than P9, movements were

followed by a significant increase in the percentage of active cells (**Figure 2A**, P5-8 median above

chance level, **Supplementary movies 1-3**). In contrast, after P9, movements were followed by a

significant decrease of activity (**Figure 2A,** P10-12 median below chance level, **Supplementary**

Movies 4-6). Short myoclonic movements such as twitches/startles, happening during periods of active

sleep (Gramsbergen et al., 1970; Jouvet-Mounier and Astic, 1968; Karlsson et al., 2006) as opposed to

longer movements, happening mostly during wakefulness, may induce different activity patterns in the

hippocampus (Mohns and Blumberg, 2008). However, we found that both movement types did not

significatively differ in their impact on CA1 activity (**Figure 2 - figure supplement 1A**). Furthermore

calcium imaging was combined with nuchal EMG recordings in one P5 mouse pup, and an increase in

the percentage of active cells was observed following movements occurring both during REM sleep

and wakefulness (**Figure 2 - figure supplement 1B**). All movement types were thus combined in

further analysis. Post-movement activity was next computed, as defined by the number of active cells

in the two seconds following movement onset divided by the number of active cells within a four

seconds-long time window centered on movement onset (see methods, **Figure 2B and Figure 2 - figure**

supplement 1C). The median post-movement activity progressively decreased from P5 to P9 until it

suddenly dropped at P10 and stabilized until P12. P9 marked the transition in the relationship between

movement and CA1 activity. Indeed, the median post-movement activity exceeded 50 % from P5 to P8

(P5: 71 %, P6: 65 %, P7: 60 %, P8: 56 %). This is consistent with the evolution of PMTHs (**Figure**
2A). After P9, the median post-movement activity was lower than 50 % (P10: 39 %, P11: 35 %, P12:
40 %), thus revealing the inhibitory action of movement on activity. We next defined as 'inhibiting
movements' all the movements with a post-movement activity lower than 40 % and computed their
proportion in each mouse (**Figure 2C**). The proportion of 'inhibiting' movements was stable before P9
(P5: 11 %, P6: 16 %, P7: 10 %, P8: 15 %). Again, P9 marks a transition since we observed that
approximately half of the movements were followed by an inhibition of CA1 activity in P10-12 mice
(P10: 55 %, P11: 58 %, P12: 48 %). The proportion of 'inhibiting' movements varies with age as a
sigmoïd function with P9 being the transition time point (V50=9.015, r^2=0.75). In line with the
emergence of movement-induced inhibition, the fraction of neurons significatively associated with
immobility also increased with age (**Figure 2D**) suggesting an "internalization" of neuronal activity.
Altogether, these results indicate that the end of the first postnatal week marks a transition in the
evolution of CA1 dynamics, with both a decorrelation and an internalization of neuronal activity. We
next investigated the circuit mechanisms supporting these changes.

Figure 2: Linking CA1 dynamics to movement during the first two postnatal weeks.

(A). Peri-Movement-Time-Histograms (PMTH) representing the percentage of active cells centered on the onset of the mouse movements. The dark line indicates the median value, the two thick grey lines represent the 25th and 75th percentiles from the distribution made of all median PMTHs from the sessions included in the group. Overall are included: P5: N=4, n=8, P6: N=3, n=5; P7: N=5, n=12; P8: N=5, n=8; P9: N=5, n=11; P10: N=1, n=1; P11: N=1, n=1; P12: N=5, n=7 (N: number of mice, n: number of imaging sessions). In all panels, the thin straight gray lines represent the 5th percentile, the median and the 95th percentile of the distribution made of all median PMTHs resulting from surrogate raster plots from the sessions included in the group. Results to build the PMTH as well as CICADA configuration file to reproduce the analysis are available in Figure 2 - Source Data 1. **(B).** Distribution of post-movement activity across age. Each boxplot is built from all detected movements for the given age group. Whiskers represent the 5th and 95th percentiles with post-movement activity falling above or below represented as small dots. The average post-movement activity observed for each mouse pup is represented by the large dots color coded from light gray (P5) to black (P12). The red area illustrates the movement falling in the category of 'inhibiting' movements. Results to build the distributions as well as CICADA configuration file to reproduce the analysis are available in Figure 2 - Source Data 1. **(C).** Distribution of the proportion of 'inhibiting' movements across age. Each dot represents a mouse pup and is color coded from light gray (P5) to black (P12). The open blue circles represent the median of the age group. The red line shows a sigmoïdal fit with V50=9.015, r^2=0.75. Results to build the distribution as well as CICADA configuration file to reproduce the analysis are available in Figure 2 - Source Data 1. **(D).** Distribution of the proportion of significatively immobility-associated cells as a function of age. Each dot represents a mouse and is color coded from light gray (P5) to black (P12). The open blue circles represent the median of the age group. Results to build the distribution as well as CICADA configuration file to reproduce the analysis are available in Figure 2 - Source Data 1.

GABAergic neurons remain activated by spontaneous movement throughout the two first postnatal weeks

As a first step to identify the circuit mechanisms for this switch, we focused on local circuits and disentangled the respective contribution of local GABAergic neurons and principal cells to CA1 dynamics as well as their relation to movement. To this aim, we identified GABAergic neurons with the expression of a red reporter (tdTomato) in *GAD67Cre* pups virally infected with AAV9-FLEX-CAG-tdTomato and AAV1.hSyn.GCaMP6s (**Figure 3A, B**). In addition, we used these imaging experiments to train a cell classifier inferring interneurons in the absence of any reporter with 91 % reliability (Denis et al., 2020). "Labeled" and "inferred" GABAergic neurons were combined into a single group referred to as "interneurons" in the following (**Figure 3 - figure supplement 2B, C**). As illustrated in a representative raster plot from a P5 mouse, both pyramidal cells (black) and interneurons (red) were activated during movement (vertical grey lines**, Figure 3A**). This was confirmed when computing the PMTH for pups aged less than P9, with the activation of the two neuronal populations after movement exceeding chance level (P5-8: N=17, n=33, pyramidal cells: baseline value = 0.51 %, peak value = 2.1 %, interneurons: baseline value = 2.1 %, peak value 7.9 %, N: number of mice, n: number of imaging sessions, **Figure 3A**). In line with above results (**Figure 2A**), pups older than P9 showed a significant reduction in the proportion of active pyramidal cells following movement (**Figure 3B**, P10-12: N=7, n=9, baseline value = 1.3 %, trough value = 0.4 %). In contrast, interneurons remained significatively activated following movement even past P9 (**Figure 3B,** P10-12: N=7, n=9,

baseline value = 3.9 %, peak value = 10 %). We conclude that the link between movement and activity
evolves differentially towards the start of the second postnatal week when comparing pyramidal neurons
and GABAergic interneurons, the former being inhibited or detached from movements while the latter
remaining activated. This suggests that pyramidal neurons could be directly inhibited by local
interneurons after the first postnatal week, following a functional maturation of GABAergic outputs
onto principal cells. Alternatively, this could result from differential changes in the synaptic inputs
driving both cell types. In the following, we have addressed both, non-mutually exclusive, hypotheses.

Figure 3: Differential recruitment of CA1 glutamatergic and GABAergic neurons

(**A).** Top panel: Imaged field of view and associated raster plot from an example imaging session in the
stratum pyramidale from one P5 *Gad67Cre* mouse pup (scale bar = 100µm). Imaged neurons expressed
GCaMP6s. Interneurons were identified by the Cre-dependent expression of the red reporter tdTomato.
In the raster plot neurons are sorted according to their identification as pyramidal cells (black) or
interneurons (red), vertical gray lines indicate movements of the mouse. Scale bar: 60 seconds. Bottom
panel: PMTHs for pyramidal cells and interneurons combining all imaging sessions from mice aged
between P5 and P8. The dark line indicates the median value and the thick grey lines represent the 25th
and 75th percentiles from the distribution made of all median PMTH obtained from the sessions included
in the group. Thin gray lines represent the 5th, median and 95th from the distribution made of all median
PMTH obtained from surrogate raster plots from the sessions included in the group (**B).** Same as (A).
But illustration is made with one P12 *Gad67Cre* mouse pup and PMTHs are built with all imaging

sessions from pups aged between P10 and P12. Note the presence of red labeled processes in the neuropil of the *stratum pyramidale* of P12 in contrast to P5. Results to build the PMTHs as well as CICADA configuration files to reproduce the analysis are available in Figure 3 - Source Data 1.

We first compared the developmental time course of extra-hippocampal synaptic afferences onto CA1 GABAergic neurons and pyramidal cells using a rabies retrograde tracing method (Wickersham et al., 2007). We focused on changes that may occur around the end of the first postnatal week. To do so, two groups were compared, an *early* (AAV1-hSyn-FLEX-nGToG-WPRE3 -Helper virus- injected at P0, SAD-B19-RVdG-mCherry -pseudotyped defective rabies virus- at P5 and immunohistochemistry (IHC) at P9, **Figure 3 - figure supplement 1A-C**) and a *late* one (AAV1-hSyn-FLEX-nGToG-WPRE3 -Helper virus- injected at P0, SAD-B19-RVdG-mCherry -pseudotyped defective rabies virus- at P9, IHC at P13, **Figure 3 - figure supplement 1D-F**). Injections were performed in either *GAD67Cre* or *Emx1Cre* pups in order to specifically target GABAergic or glutamatergic cells, respectively. Four *GAD67Cre* pups (2 early and 2 late injections) and three *Emx1Cre* pups (1 early and 2 late) were analyzed with injection sites restricted to the hippocampus. Starter and retrogradely labeled cells were found all over the ipsilateral hippocampus. For both *GAD67Cre* and *Emx1Cre* pups, we found no striking difference in the retrogradely labeled extra-hippocampal regions between the *early* and *late* groups. In agreement with previous studies (Supèr and Soriano, 1994), we found that GABAergic and glutamatergic neurons in the dorsal hippocampus received mainly external inputs from the entorhinal cortex, the medial septum and the contralateral CA3 area (retrogradely labeled cells in these regions were found in 4 out of 4 *GAD67Cre* pups and 3 out of 3 *Emx1Cre* pups, **Figure 3 - figure supplement 1**). Thus, we could not reveal any major switch in the nature of the extra-hippocampal inputs impinging onto local CA1 neurons. Thus, we next explored the maturation of local somatic GABAergic innervation given its significant evolution throughout that period (Jiang et al., 2001; Marty et al., 2002; Morozov and Freund, 2003) as well as our observation of a dense tdTomato signal in the pyramidal layer from *GAD67Cre* mouse pups at P12 (**Figure 3B**), not visible at P5 (**Figure 3A**).

Abrupt emergence of a functional somatic GABAergic innervation at the beginning of the second postnatal week

We first analyzed the anatomical development of somatic GABAergic innervation within the CA1 pyramidal layer from P3 to P11, focusing on the innervation from putative parvalbumin-expressing Basket Cells (PVBCs), its main contributor. To this aim, we performed immunohistochemistry against Synaptotagmin2 (Syt2) which has been described as a reliable marker for parvalbumin positive inhibitory boutons in cortical areas (**Figure 4A**, (Sommeijer and Levelt, 2012)). Using a custom-made

Fiji-plugin (RINGO, see methods), we quantified the surface of the pyramidal cell layer covered by

Syt2 labeling at different stages and found that between P3 and P7, PV innervation remained stable

(median values: P3: 0.34 %, P5: 0.57 %, P7: 0.49 %, 3 mice per group, **Figure 4B**). However, around

P9, a sudden increase in the density of positive labelling was observed (P9: 1.03 %, P11: 1.48 %, 3 mice

per group, **Figure 4B**). These results are consistent with previous work (Jiang et al., 2001; Marty et al.,

2002), as well as with our tdTomato labeling (**Figure 3A-B**). They also match the transition observed

in CA1 dynamics (**Figure 3**). We next tested whether GABAergic axons in the pyramidal layer were

active during periods of movement. In order to increase success rate for imaging axons in the pyramidal

layer, we restricted these experiments to P9-10, i.e., when axons start densely innervating the CA1 layer.

To do so, we restrained the expression of the calcium indicator GCaMP6s to the axon (Broussard et al.,

2018) of interneurons using *GAD67Cre* mouse pups and specifically imaged axonal arborisation in the

pyramidal cell layer (**Figure 4C, left panel**). Fluorescence signals were extracted from axonal branches

using PyAmnesia (a method to segment axons, see methods, **Figure 4C, right panel**), and then

normalized using z-score (see methods). As expected, (see **Figure 3B**), an increase in the fluorescent

signal from GABAergic axonal branches was observed following movement (P9-10: n=3 mice, **Figure**

4D). As a result, we reasoned that an increase in perisomatic GABAergic inhibition could contribute to

the reduction of activity observed after movement during the second postnatal week in pyramidal

neurons.

Figure 4: Emergence of perisomatic GABAergic innervation

(**A**). Representative example confocal images of the CA1 region in a P7 (left) and a P11 (right) mouse
pup. DAPI staining was used to delineate the *stratum pyramidale* (sp) from the *stratum radiatum* (sr)
and *stratum oriens* (so, top row). Synaptotagmin-2 labeling (Syt2) is shown in the top and bottom rows.
Illustrated examples are indicated by red dots in the associated quantification in (B). Scale bar= 20 µm.
(**B**). Fraction of the pyramidal cell layer covered by Syt2 positive labeling as a function of age. Each
gray dot represents the average percentage of coverage from two images taken in the CA1 region of a
hippocampal slice. Open black dots are the average values across brain slices from one mouse pup. Blue
arrows indicate the slices used for illustration in (A). A significant effect of age was detected (One-Way
Anova, F=13.11, p=0.0005). Multiple comparison test shows a significant difference between age
groups (Bonferroni's test, *: p<0.05, **: p<0.01, ***: p <0.001). (**C**). Averaged image of a field of view
in the pyramidal cell layer of a P9 *GAD67Cre* mouse pup injected with a Cre-dependent Axon-
GCaMP6s indicator (left) and the segmented image resulting from PyAmnesia (right). (**D**). PMTH
showing the df/f signal centered on the onsets of animal movement (N=3, n=3). The dark gray line
indicates the median value, and the light gray lines represent 25[th] and 75[th] percentile. Results obtained
from surrogates are represented by light gray lines.

Increasing feedback inhibition in two-population models reproduces the developmental transition

To test whether an increase in perisomatic inhibition alone can explain the switch in network dynamics between the first and second postnatal weeks, we simulated a two-population network mimicking the development of perisomatic innervation (**Figure 5A**, see methods). Using a rate model and a Leaky Integrate and Fire (LIF) model, we show that increasing the strength of perisomatic inhibition can account for the experimentally observed decrease of responses to movement-like feedforward inputs (**Figure 5B**). Durations of feed-forward inputs were chosen similar to experimental movement durations (see **Figure 5 - figure supplementary 1A** for a log-normal of the movement durations). When inhibition is weak, the average activity of the pyramidal neurons increases at the onset of a given twitch. Then, it quickly relaxes to the baseline with a timescale that follows the synaptic time constant (**Figure 5B**, left panel). In the presence of strong inhibition, there is a reduction in response to movement inputs. In addition, due to strong feedback inhibition following the movement responses, network activity relaxes to the baseline with an undershoot (**Figure 5B**, right panel), recapitulating the experimental findings (see **Figure 2A**). Similar PMTHs were obtained for interneurons (see **Figure 5 - figure supplementary 1 B**).

To further show the importance of perisomatic inhibition on network dynamics, we computed the cross-correlogram between interneurons activity and pyramidal cell activity in the absence of movement (P5-8 light gray curve, P9-12 dark curve, **Figure 5C, left panel**). For P9-12, we observe a rapid drop in the correlation at positive time which means that elevated inhibitory activity is followed by a strong decrease in excitatory activity. Notice that this drop is absent for P5-8, where the cross-correlogram is mostly symmetric around t=0. Both the rate and LIF models displayed similar activity correlograms as the experimental data, with an undershoot in the presence of strong inhibitory feedback for positive time. We showed through numerical simulations that similar changes could be observed in a more realistic spiking network model (**Figure 5C, right panel**). Auto-correlograms of the excitatory and inhibitory activity were also measured and compared to our model predictions (see **Figure 5 - figure supplementary 1C**).

The consistency of our models with the experimental cross-correlograms, which were computed from the activity recorded during periods of immobility, further shows that the observed network dynamics and, in particular, the correlation undershoots most likely result from recurrent perisomatic inhibition rather than a feedforward drive from upstream areas. Therefore, in our model, the maturation of perisomatic inhibition alone was sufficient to support a switch in network dynamics.

Figure 5: Modeling the effects of perisomatic inhibition on pyramidal cell response

(A). The model consists of two populations, Excitatory (E) and Inhibitory (I) receiving feedforward input I_{ext}. The interaction strengths, Jab represent the effect of the activity of population b on a. We study the effects of perisomatic inhibition on the activity of pyramidal cells by varying the parameter J_{EI}. **(B).** PMTH, response of excitatory neurons to pulse input in the rate model and LIF network. **(C).** Cross-correlations during periods of immobility, in experimental data (left), rate model (middle), and LIF network (right). In the rate model, dotted lines are the predicted correlation from the analytic expressions and solid lines are the results from numerical integration.

DISCUSSION:

Using for the first time *in vivo* two-photon calcium imaging in the hippocampus of non-anesthetized mouse pups and a deep-learning based approach to infer the activity of principal cells and interneurons, we show that the end of the first postnatal week marks a salient step in the anatomical and functional development of the CA1 region. Indeed, within two days (P8-10), the link between CA1 principal cells activity and self-triggered movements is inverted and neurons are preferentially active during immobility periods. This is likely due to the time-locked anatomical and functional rise of somatic GABAergic activity given that interneurons remain highly active throughout this period, including in response to spontaneous movements. In this way, CA1 circuits start detaching from external inputs. Given the importance of internal dynamics for hippocampal function and cortical circuits operation in general, this is likely to be a critical general step in the proper maturation of cognitive circuits.

Early postnatal calcium activity in CA1 is driven by sensorimotor inputs

We found that, until P7-9, spontaneous movements are followed by a significant peak in calcium events in the CA1 principal cell layer and that most neuronal activity occurs during synchronous calcium events. This early link between sensorimotor inputs and early cortical dynamics has been previously reported using electrophysiological recordings in various areas and species, including humans (Milh et al., 2006). Here we extend that observation to calcium transients, which not only indirectly report action-potential firing as well as other modes of cell activation during development but also critically regulate activity-dependent genetic processes. In addition, we could describe the response to these movements with single-cell resolution. It is possible that other movements, like whisker movements, that we have not detected also contribute to the patterning of CA1 activity. It is also possible that self-generated activity from other sensory organs but independent from movement, like the retina or the olfactory bulb, also contribute to hippocampal dynamics, the same way they are relayed to sensory cortices. Interestingly, in contrast to previous reports (Tiriac et al., 2014), we did not observe any significant difference between twitches (occurring mainly during active sleep) and longer, more complex movements. This may reveal a difference between calcium imaging and electrophysiology, the former sampling from a larger population but at a lower temporal and spike signal resolution. The patterning of CA1 dynamics in the large imaged population did not reveal any obvious spatial distribution for movement-activated cells but we cannot exclude that these would vary along the radial and transverse directions, which are the two main axes of principal cell development (Caviness, 1973), and are differentially targeted by perisomatic PV basket cells (Lee et al., 2014; Valero et al., 2015).

Passed the end of the first postnatal week, between P8 and P10, a significant decrease in the fraction of coactive principal cells following movement was observed (while interneurons remained mostly activated by movement). We cannot exclude that some spikes fell below the threshold for detection of calcium events. In this case, rather than full inhibition, it may be that a strong shortening of the time window for neuronal integration occurred (due to feedback inhibition), which would limit the number of spikes produced by principal cells and thus keep them below detection levels. Yet, a novel machine-learning based algorithm (Denis et al., 2020) was used since it was especially designed to infer activity in the dense CA1 pyramidal cell layer. This change in the polarity of principal cells' response to movements is quite abrupt as it happens within less than two days (between P8 and 10). This contrasts with the progressive evolution of single cell firing frequencies but matches the fast redistribution of neuronal firing towards immobility periods. In this way, hippocampal neuronal dynamics "internalize" as they stop being driven by movements and preferentially occur within rest.

This "internalization" of hippocampal dynamics is reminiscent of similar phenomena observed in other cortical areas, such as the barrel cortex where whisker stimulation induces a reduction in the size of cell assemblies following P9 while the same stimulation widens cell assembly size a few days

before (Mòdol et al., 2019). It is also reminiscent of the recently described transient quiescent period observed in the somatosensory cortex using extracellular electrophysiological recordings (Dominguez et al., 2021). Last, it goes in hand with a sparsification of activity, which is a general developmental process supported by the emergence of inhibition (Golshani et al., 2009; Rochefort et al., 2009; Wolfe et al., 2010).

Circuit basis for the movement-triggered inhibition of CA1 dynamics

Our results demonstrate that the "internalization" of CA1 dynamics occurring at the end of the first postnatal week most likely relies on structural changes in local CA1 circuits rather than rewiring of the long-range extrahippocampal connectivity.

The long-range circuits mediating the bottom-up flow of self-triggered or externally generated sensory information to the hippocampus are starting to be elucidated. The two main structures directly transmitting sensorimotor information to the dorsal CA1 are the entorhinal cortex and septum. The former processes multisensory information from all sensory cortices (visual, auditory, olfactory, somatosensory), including movement-related sensory feedback (Rio-Bermudez and Blumberg, 2021)and was shown to be activated by spontaneous twitches prior to CA1 (Mohns and Blumberg, 2010; Rio-Bermudez et al., 2020; Valeeva et al., 2019a) while the latter is more likely to be involved in transmitting internal information (Fuhrmann et al., 2015; Wang et al., 2015) as well as unexpected environmental stimuli (Zhang et al., 2018). In addition to these two canonical pathways, one cannot exclude the involvement of a direct connection from the brainstem, given their existence in the adult and their role in promoting sleep as well as motor twitches (Liu et al., 2017; Szőnyi et al., 2019). However, our retrograde tracing experiments did not reveal any direct connection between the CA1 cells and the brainstem at the early ages analyzed here. In addition, we found that both CA1 interneurons and principal cells receive inputs from the septum and entorhinal cortex before the time of the switch (i.e. P9) and that there was no major qualitative change of inputs after, as expected from previous work (Supèr and Soriano, 1994). Still, these experiments do not allow a quantitative assessment of the number of inputs nor the type of inputs (GABAergic, cholinergic, etc.) and we cannot fully exclude that a stronger, or different source of excitatory drive would be impinging onto interneurons after the switch. Therefore, future optogenetic and slice physiology work is needed to characterize the bottom-up information flow onto specific components of the local CA1 circuits. Similarly, one cannot exclude a change in the CA3 to CA1 connectivity. Indeed, Schaffer collaterals are known to reach CA1 roughly around the end of the first postnatal week (Durand et al., 1996). In addition, roughly at the time of the switch, do we see the emergence of SWRs (Buhl and Buzsaki, 2005), a pattern strongly relying on CA3 inputs and perisomatic GABAergic transmission. However, we could not restrict the pool of starter cells to the CA1 region in our retrograde viral tracing experiments, which precluded analysis of the development of CA3-CA1 connectivity. Interestingly, among the external inputs onto CA1 described

above, the entorhinal cortex and CA3 were both shown to exert a mild influence on the organization of
intrinsic CA1 dynamics, possibly pointing at a critical role of local interneurons in this process (Zutshi
et al., 2021).

As indicated by our computational model, the disengagement from movement of CA1 dynamics
can be fully explained by the observed rise in anatomical (Syt2 labeling) and functional (axonal GCaMP
imaging) connectivity from perisomatic GABAergic cells onto pyramidal cells at the onset of the second
postnatal week. This increased connectivity could not be easily captured with our retrograde viral
labeling since the absence of early PV expression precludes the identification of PV basket cells, the
most prominent subtype of perisomatic GABAergic cells, among retrogradely labelled cells in *Emx1Cre*
pups. Early anatomical studies had already indicated that an increase of somatic GABAergic inhibition,
including from CCK-basket cells, occurred in CA1 during the first postnatal week (Danglot et al., 2006;
Jiang et al., 2001; Marty et al., 2002; Morozov and Freund, 2003). However, this rise was expected to
be more progressive and not as abrupt as observed here, as it happened within two days. If the axonal
coverage of the stratum pyramidale by PV-basket cells axons increases, we cannot exclude that this is
a general phenomenon, concerning all perisomatic subtypes, including soma-targeting CCK-expressing
basket cells which develop anatomically at around the same time (Morozov and Freund, 2003) or
chandelier cells. In addition, our computational model indicates that the emergence of feed-back
inhibition is sufficient to reproduce the developmental shift observed here, which could also involve
other types of CA1 interneurons, including dendrite-targeting ones.

Interestingly, a similar rise of somatic GABAergic axonal coverage occurs in the barrel cortex
at the same time. Indeed, recent connectomic mapping using 3D-electron microscopy in that region
revealed that the preferential targeting of cell bodies by GABAergic synapses increased almost threefold
between postnatal days 7 and 9 (Gour et al., 2020), whereas two-photon imaging of putative GABAergic
somatic axons in the same region revealed broader domains of co-activation (Mòdol et al., 2019). This
time period for the shift may be synchronous within brain regions involved in sensorimotor integration
such as the hippocampus and somatosensory cortex. Otherwise, PV expression was shown to develop
sequentially in a region-specific manner (Reh et al., 2020) following their intrinsic developmental age
(Donato et al., 2017).

We found that many principal cells are inhibited by movement while most imaged GABAergic
cells remained activated during the second postnatal week. This therefore indirectly suggests a net
inhibitory effect of GABAergic transmission after the first postnatal week. This is expected since the
shift from excitatory to inhibitory synaptic transmission was reported to occur earlier in the
hippocampus *in vivo* (Murata and Colonnese, 2020). On a side note, the lack of somatic GABAergic
inputs before P7 indicates that the early excitatory GABAergic drive in CA1 circuits likely originates
from non-somatic GABAergic interneurons, which include long-range, dendrite-targeting or

interneuron-specific interneurons. The circuit role of excitatory GABAergic transmission should be
revisited taking into account this new finding.

The movement-associated inhibition can result equally from feedforward (direct activation
from movement-transmitting inputs such as the entorhinal cortex) or feedback (from local CA1 cells)
inhibition. Our experiments do not allow disentangling both circuits. A progressive strengthening of
feedback inhibition circuits through the strengthening of local CA1 principal cells inputs onto
perisomatic GABAergic neurons is also possible, as it occurs in the developing somatosensory cortex
(Anastasiades and Butt, 2012). The inhibition of activity following movement is likely to be occurring
during a transient developmental period. Indeed, in the adult, both interneurons and principal cells
usually increase their activity as the animal moves (Fuhrmann et al., 2015). Therefore the switch
observed here opens another developmental time-window that probably closes with the emergence of
perineuronal nets and cell activation sequences at the end of the third developmental week (Farooq and
Dragoi, 2019; Horii-Hayashi et al., 2015; Muessig et al., 2019). We would like to propose this
developmental window, to be the critical period for CA1 development, a period during which
experience-dependent plasticity can be observed.

Conclusion

Cognitive hippocampal maps rely on two forms of representation, one that is map-based or allocentric
and the other that is self-referenced, or egocentric and requires body movement. We would like to
propose that the early postnatal period described here, where the hippocampus learns the statistics of
the body, and which terminates with the rise of a recurrent inhibitory network is a key step for the
emergence of an internal model onto which exploration of the external world can be grafted. An
imbalance between internal and environmental hippocampal representations due to a miswiring of local
somatic inhibition could have major outcomes. It could be at the basis of several neurodevelopmental
disorders, including autism spectrum disorders (ASD) and schizophrenia. Interestingly, both disorders
have been associated with an aberrant maturation of PV-expressing interneurons (Gogolla et al., 2014;
Jurgensen and Castillo, 2015; Lewis et al., 2005). In addition, the proper development of the peripheral
sensory system, which is partly initiating the early CA1 dynamics reported in our study, is also critically
involved in ASD (Orefice et al., 2016). The period described here corresponds to the third trimester of
gestation and likely extends postnatally given the protracted integration of GABAergic interneurons
into functional circuits in the human brain (Murphy et al., 2005; Paredes et al., 2016). Future work
should determine when a similar rise in somatic inhibition occurs in human infants and test whether it
could constitute a valuable biomarker for cognitive neurodevelopmental disorders.

Figure 1 – figure supplement 1

512 **Figure 1 - figure supplement 1**

513 **(A).** Distribution of the number of cells imaged in each field of view as a function of age. Each dot
514 represents a field view and is color coded from light gray (P5) to black (P12). The open blue circles
515 represent the median of each group. No effect of age was observed on the number of imaged cells. **(B).**
516 Top panels: Example of the electrophysiological recording done in the ipsi (hippocampus with window

implant) and contralateral (intact hippocampus) hippocampi. Use of multisite silicon probe allowed simultaneous recordings in different hippocampal layers (strati oriens (so), pyramidale (sp), radiatum (sr) and lacunosum moleculare (slm)). The eSW is marked by an asterisk. Bottom panels: box plot displays the eSW occurrence. Each dot represents the eSW rate per minute in a mouse pup (N=6). Box plots show median, the bottom and top edges of the box indicate the 25th and 75th percentiles, the whiskers extend to the most extreme data points not considered outliers. Bottom right panel: Cooccurrence of eSWs detected in both hippocampi. **(C)**. Developmental changes in power spectral density of the recorded LFP in the ipsi (hippocampus with window implant) and contralateral (intact hippocampus) hippocampi. Gray and black lines are the averages between the animals of P6-8 and P11 age ranges, respectively. Shading shows jackknife standard deviation. Note, presence of peak in theta frequency band (4-7 Hz) in P11 animals. **(D).** Raster plots from the imaging sessions used for illustration in Figure 1A, showing all cells over the full recording, scale bar is 2 minutes.

Figure 2 - figure supplement 1

(A). PMTHs for each age group built on twitches only (top row) or on complexe movements only (bottom row). Procedure to group sessions is similar to the one described in Figure 2 **(B).** PMTHs from one P5 mouse pup (including 2 imaging sessions) combining all detected movements during REM sleep ('P5 -REM') or all detected movement during Wakefulness ('P5 - AWAKE'). **(C).** Definition of the post-movement activity. **(D).** Distribution of the proportion of cells significatively associated with movement as a function of age. Each dot represents a mouse pup and is color coded from light gray (P5) to black (P12). The open blue circles represent the median for each group.

Figure 3 - figure supplement 1

(**A**). Experimental timeline used to investigate the inputs received by pyramidal cells (*Emx1Cre* mice) and interneurons (*GAD67Cre* mice) during the first postnatal week (early group). At P0 and P5 we injected the helper virus AAV1-hSyn-FLEX-nGToG-WPRE3 and the rabies SAD-B19-RVdG-mCherry into the hippocampus of *GAD67Cre* (Figure 3B) mice and *Emx1Cre* mice (Figure 3C) respectively. Brains are next collected for histological/anatomical analysis at P9. (**B**). Confocal images from early group *GAD67Cre* mice showing the injection sites in the dorsal hippocampus on sagittal slices at lower magnification (top left, scale bar= 1 mm), and higher magnification (top right, scale

bar=100 µm). Primo infected cells are labelled in green, the starter cell indicated by the arrow shows co-expression of the helper virus (green) and rabies virus (red). Presynaptic cells, labelled only in red, are localized in the entorhinal cortex (bottom left, scale bar= 1 mm) and in the medial septum (bottom right, scale bar= 1 mm). Hpc= Hippocampus; Ent. Ctx= Entorhinal Cortex; MS= Medial Septum **(C).** same as in (B) but for *Emx1Cre* mice. Scale bars from top left to bottom right: 200 µm, 100 µm, 500µm, 100 µm and 500 µm. Middle bottom image is a higher magnification of the bottom left panel. **(D).** Same as in (A) but for the late group (P9-P13). **(E).** Same as in (B) but for the late group. Scale bars from top left to bottom right: 1 mm, 100 µm, 1 mm and 1mm. **(F).** Same as in (C) but for the late group. Scale bars from top left to bottom right: 200 µm, 100 µm, 500µm, and 500 µm.

Figure 3 - figure supplement 2

(**A**). Top row: confocal images taken from a 69 µm z-stack (z-step: 1.5 µm) of a brain slice from a
Gad67Cre mouse pup injected with both AAV9-FLEX-tdTomato and AAV1.hSyn.GCaMP6s showing
the co-infection between tdTomato and GCaMP6s in hippocampal interneurons. Bottom row: maximum
intensity projections of the example z-stack showed in (A). On the maximal intensity projection from
tdTomato signal white arrows indicate interneurons not expressing GCaMP6s, gray arrows indicate
interneurons expressing GCaMP6s. The proportion of tdTomato cells expressing GCaMP6s is shown
on the plot for each layer of the region CA1 of the hippocampus. The proportion of tdTomato cells
expressing GCaMP6s *in vivo* not only take into account the co-expression of the two but represent the
proportion of tdTomato cells having a GCaMP6s signal that was not classified as 'noise' explaining the
relatively low proportion in comparison with the observed co-infection rate. Scale bars= 50 µm (**B**).
Evolution of the number of transient per minute observed in 'labelled' interneurons and 'inferred'
interneurons from P5 to P12. Each dot represents the average obtained from one mouse and is color
coded from light gray (P5) to black (P12). Filled dots represent 'inferred' interneurons, open dots

577 represent 'labelled' interneurons. **(C).** PMTHs combined all imaging sessions between P5 and P12
578 showing the activation after movement of both 'labelled' and 'inferred' interneurons.

Figure 5 – figure supplement 1

(A). Movement duration histograms. These histograms were fit by log-normal distributions. **(B).** PMTHs of the interneurons predicted by the rate and LIF models. **(C).** Auto-correlograms of the activity for the pyramidal cells and interneurons.

Parameter	Symbol	Value
Rate model		
Excitatory time constant	τ_E	0.7s
Inhibitory time constant	τ_I	3.0s
Connectivity matrix		
	J_{EE}	0.8
	J_{IE}	0.5
	J_{II}	−0.2
	J_{EI}^{weak}	−0.0
	J_{EI}^{strong}	−0.4
Noise amplitude	η	$0.1s^{-1/2}$
External input		
	H_E	0.5
	H_I	0.1
External drive increase	α	3.0
Twitch amplitude		
	J_E	7.0
	J_I	25.0
LIF model		
Exc. synapse time constant	τ_E	1.0s
Inh. synapse time constant	τ_I	3.0s
Membrane time constant	τ_m	20.0 ms
Threshold potential	V_{thr}	20.0 mV
Reset potential	V_{reset}	0.0mV
Connectivity		
Nb. of excitatory neurons	N_E	512
Nb. of inhibitory neurons	N_I	128
Connections from exc. neurons	K_E	32
Connections from inh. neurons	K_I	16
Connectivity matrix		
	J_{EE}	0.5
	J_{IE}	2.0
	J_{II}	−0.3
	J_{EI}^{weak}	−0.05
	J_{EI}^{strong}	−0.8
Noise amplitude	η	$1.0s^{-1/2}$
External input		
	H_E	1.0
	H_I	0.1
Twitch amplitude		
	J_E	2.0
	J_I	10.0
Twitch duration (lognormal)		
Mean parameter	μ	−0.2
Std parameter	σ	0.8

Figure 5 – Supplementary table 1:

592 Details of the parameters used in the rate and LIF models

 SUPPLEMENTARY MOVIES

Supplementary Movies 1-3: Three examples of calcium imaging movies from P5 mouse pups centered
on the onset of a twitch. The twitch is indicated by T on the upper left corner of the movie. Imaging 2
times speed up.

Supplementary Movies 4-6: Three examples of calcium imaging movies from P12 mouse pups centered
on the onset of a complex movement. The complex movement is indicated by M on the upper left corner
of the movie. Imaging 2 times speed up.

SOURCE DATA

Figure 1 - Source Data 1. Contains 3 folders. **1A**: contains the 4 raw '.tiff' files illustrating the field of
views and the CICADA configuration files necessary to plot the contours map and raster plots used for
the illustration. **1B**: contain (in an '.xlsx' file) the numerical data used to plot the evolution of the
transient in SCE and the CICADA configuration file necessary to reproduce the analysis. **1C**: contain
(in an '.xlsx' file) the numerical data used to plot the evolution of the transient per minute and the
CICADA configuration file necessary to reproduce the analysis.

Figure 2 - Source Data 1. Contains 4 folders. **2A**: contain (in an '.xlsx' file) the numerical data used to
plot all the PMTHs (in Figure 2A) and the CICADA configuration file necessary to reproduce the
analysis. **2B**: contain (in an '.xlsx' file) the numerical data used to plot Figure 2B and the CICADA
configuration file necessary to reproduce the analysis. **2C**: contain (in an '.xlsx' file) the numerical data
used to plot Figure 2C and the CICADA configuration file necessary to reproduce the analysis. **2D**:
contain (in an '.xlsx' file) the numerical data used to plot Figure 2D and the CICADA configuration
file necessary to reproduce the analysis.

Figure 3 - Source Data 1. Contains 3 folders. **'example_FoVs':** contains the '.tiff' images used for
illustration in Figure 3. **'example_raster_plots'**: contains the 2 CICADA configuration files necessary
to reproduce the raster plots used for illustration in Figure 3. **'psths'**: contains (in an '.xlsx' file) the
numerical data used to plot Figure 3 PMTHs and the CICADA configuration file necessary to reproduce
the analysis.

Figure 4 - Source Data 1. Contains 4 folders. **4A**: contains '.tiff' images used for illustration in Figure
4A. **4B**: contains the numerical data of the plot in Figure 4B. **4C**: contains the CICADA configuration
file necessary to plot the contour map shown in Figure 4C. **4D**: contains (in an '.xlsx' file) the numerical
data used to plot Figure 4C PMTH and the CICADA configuration file necessary to reproduce the
analysis.

MATERIALS AND METHODS

MICE

All experiments were performed under the guidelines of the French National Ethics Committee for Sciences and Health report on "Ethical Principles for Animal Experimentation" in agreement with the European Community Directive 86/609/EEC (Apafis#18-185 and #30-959).

EXPERIMENTAL PROCEDURES AND DATA ACQUISITION

Viruses. *In vivo* calcium imaging experiments were performed using AAV1-hSyn-GCaMP6s.WPRE.SV40 (pAAV.Syn.GCaMP6s.WPRE.SV40 was a gift from Douglas Kim & GENIE Project (Addgene viral prep #100843-AAV1; http://n2t.net/addgene:100843; RRID:Addgene_100843)), AAV9-FLEX-CAG-tdTomato (pAAV-FLEX-tdTomato was a gift from Edward Boyden (Addgene viral prep # 28306-AAV9 ; http://n2t.net/addgene:28306 ; RRID:Addgene_28306)), AAV9-hSyn-FLEX-axon-GCaMP6s (pAAV-hSynapsin1-FLEx-axon-GCaMP6s was a gift from Lin Tian (Addgene viral prep # 112010-AAV9 ; http://n2t.net/addgene:112010 ; RRID:Addgene_112010)). Retrograde tracing experiments were performed using AAV1-hSyn-FLEX-nGToG-WPRE3 (Charité #BA-096) and SAD-B19-RVdG-mCherry (Gift from the Conzelmann laboratory).

Intracerebroventricular injection. This injection protocol was adapted from already published methods (Kim et al., 2014). Mouse pups were anesthetized on ice for 3 – 4 min, and 2 µL of viral solution (titration at least 1 x 10^{13} vg / mL) were injected in the left lateral ventricle whose coordinates were estimated at the 2/5 of the imaginary line between the lambda and the eye at a depth of 0.4 mm. Correct injection was visualized by the spreading of the viral-dye mixture (1/20 of fast blue). In SWISS mouse pups we injected 2 µL of AAV2.1-hSyn-GCAMP6s.WPRE.SV40, in *GAD67Cre* mouse pups we injected either a mix of 1.3µL of AAV2.1-hSyn-GCAMP6s.WPRE.SV40 with 0.7µL of AAV9-FLEX-CAG-tdTomato or 2µL of AAV9-hSyn-FLEX-axon-GCaMP6s.

Intra-hippocampal injection. When hippocampal viral injections were performed at P0 (AAV1-hSyn-FLEX-nGToG-WPRE3), mouse pups were anesthetized by inducing hypothermia on ice and maintained on a dry ice-cooled stereotaxic adaptor (Stoelting, # 51615) with a digital display console (Kopf, Model 940). Dorsal hippocampus was targeted by empirically determined coordinates, based on the Atlas of the Developing Mouse Brain (Paxinos et al., 2020), using transverse sinus and superior sagittal sinus as reference: 0.8 mm anterior from the sinus intersection; 1.5 mm lateral from the sagittal sinus; 1.1 mm depth from the skull surface. Under aseptic conditions, an incision was made in the skin, the skull was exposed and gently drilled (Ball Mill, Carbide, #¼ .019" -0.500 mm diameter-, CircuitMedic). Ten nL of undiluted viral solution were injected using an oil-based pressure injection

system (Nanoject III, Drummond Scientific, rate of 5 nL/min). The tip of the pipette was broken to achieve an opening with an internal diameter of 30-40 μm. When hippocampal viral injections were performed at P5 or P9 (SAD-B19-RVdG-mCherry), pups were anesthetized using 3 % isoflurane in a mix of 90 % O_2-10 % air and maintained during the whole surgery (~0:30 h) between 1 % and 2.5 % isoflurane. Body temperature was monitored and maintained at 36°C. Analgesia was controlled using Buprenorphine (0.05 mg/ kg). Under aseptic conditions, an incision was made in the skin, the skull was exposed, antero-posterior and medio-lateral coordinates of the dorsal hippocampus were estimated by eye looking at the skull sutures. The skull was gently drilled and 10 nL of a viral solution were injected (Nanoject III, Drummond Scientific, rate of 5 nL/min) at a depth of 1.25 mm below the dura.

Window implant surgery. The surgery to implant a 3-mm-large cranial window above corpus callosum was adapted from previous methods (Villette et al., 2015). Anesthesia was induced using 3 % isoflurane in a mix of 90 % O_2-10 % air and maintained during the whole surgery (~1:30 h) between 1 % and 2.5 % isoflurane. Body temperature was controlled and maintained at 36°C. Analgesia was controlled using Buprenorphine (0.05 mg/ kg). Coordinates of the window implant were visually estimated. Then a small custom-made headplate was affixed with cyanoacrylate and dental acrylic cement. The skull was removed and the cortex was gently aspirated until the appearance of the external capsule/alveus. At the end of the cortectomy, we sealed a 3 mm glass window diameter circular cover glass (#1 thickness, Warner Instrument) attached to a 3 mm diameter large and 1.2 mm height cannula (Microgroup INC) with Kwik-Sil adhesive (WPI) and fixed the edge of the glass with cyanoacrylate. We let the animal recover on a heated pad for at least one hour before the imaging experiment.

Imaging. Two-photon calcium imaging experiments were performed on the day of the window implant using a single beam multiphoton pulsed laser scanning system coupled to a microscope (TriM Scope II, LaVision Biotech). The Ti:sapphire excitation laser (Chameleon Ultra II, Coherent) was operated at 920 nm. GCaMP fluorescence was isolated using a bandpass filter (510/25). Images were acquired through a GaAsP PMT (H7422-40, Hamamatsu) using a 16X immersion objective (NIKON, NA 0.8). Using Imspector software (LaVision Biotech), the fluorescence activity from a 400 μm x 400 μm field of view was acquired at approximately 9 Hz with a 1.85 μs dwell time per pixel (2 μm/pixel). Imaging fields were selected to sample the dorsal CA1 area and maximize the number of imaged neurons in the *stratum pyramidale*. Piezo signal, camera exposure time, and image triggers were synchronously acquired and digitized using a 1440A Digidata (Axon instrument, 50 kHz sampling) and the Axoscope 10 software (Axon instrument). During the imaging session, body temperature is continuously controlled.

Behavioral recordings. Simultaneously with imaging experiments, mouse motor behavior was monitored. In a first group of animals, motor behavior was monitored using two or three piezos attached to the paws of the animal. The signal from the piezo was acquired and digitized using a 1440A Digidata

and the Axoscope 10 software. In a second group of animals, pups were placed and secured on an
elevated platform (with the limbs hanging down on each side without touching the ground nor the
support, as described here (Blumberg et al., 2015)). Motor behavior was monitored using two infrared
cameras (Basler, acA1920-155um) positioned on each side of the animal. For each camera, a square
signal corresponding to the exposure time of each frame from the camera was acquired and digitized
using a 1440A Digidata and the Axoscope 10 software.

Recording of electromyogram activity in neonatal mice. The vigilance state of neonatal mice was
assessed through analysis of electromyogram (EMG) signals obtained from a single insulated Tungsten
wire (A-M Systems 795500) implanted in the nuchal muscle. A stainless-steel wire (A-M Systems
786000) wire inserted on the skull surface above the cerebellum and secured in place with dental cement
served as the reference electrode. Signals from the electrodes were first passed through a headstage pre-
amplifier before being digitized at 16000 Hz (Digital Lynx SX, Neuralynx (the pre-amplifier and
digitizer were both from Neuralynx, as was the acquisition software (Cheetah))) and saved to a hard
disk. TTL signals from the imaging and camera acquisition systems were simultaneously recorded as
well to enable precise synchronization of EMG recordings with the camera and imaging data.

In vivo extracellular electrophysiological recordings. Multisite probes (16 channels silicon probes with
50 um separation distance, NeuroNexus, USA) were used to record electrophysiological activity below
the window implant and in the intact hippocampus. To do so, we positioned the mouse pup (that had
previously undergone a window implant) on the experimental setup. To head fix the animal the skull
surface was covered with a layer of dental acrylic except the area above the intact hippocampus. In the
intact (contralateral) hippocampus the electrodes were positioned using the stereotaxic coordinates of
approximately 1.5 mm anterior to lambda and 1.5 mm lateral from the midline. Hippocampus under the
window was recorded through the hole drilled in the window implant. Both multisite silicon probes
were positioned at the depth to record *strata oriens* (SO), *pyramidale* (SP), *radiatum* (SR) and
lacunosum moleculare (SLM). After the positioning of the electrodes, the animal was left in the setup
for one hour to recover followed by 2 hours recordings of the neuronal activity in both hippocampi
simultaneously.

Histological processing. Pups were deeply anaesthetized with a mix of Domitor and Zoletil (0.9 and 60
mg/kg, respectively), then transcardially perfused with 4 % PFA in 0.1M PBS (PBS tablets, 18912-014,
Life technologies). For perisomatic innervation analysis, brains were post-fixed overnight at 4°C in 4
% PFA in 0.1M PBS, washed in PBS, cryo-protected in 30 % sucrose in PBS, before liquid nitrogen
freezing. Brains were then sectioned using a cryostat (CM 3050S, Leica) into 50 μm thick slices
collected on slides. Sections were stored at -20°C until further usage. For tracing experiments, brains
from *GAD67Cre* and *Emx1Cre* pups, were post-fixed overnight at 4 °C in 4 % PFA in 0.1M PBS,
washed in PBS and sectioned using a vibratome (VT 1200 s, Leica) into sagittal 70–80 μm thick slices.

Sections were stored in 0.1M PBS containing 0.05 % sodium azide until further usage. Immunocytochemistry was processed as described previously (Bocchio et al., 2020) . Briefly, sections were blocked with PBS-Triton (PBST) 0.3 % and 10 % normal donkey serum (NDS), and incubated with a mix of up to three primary antibodies simultaneously diluted in PBST with 1 % NDS overnight at room temperature with the following primary antibodies: rabbit anti-dsRed (1:1000; Clontech, AB_10013483), chicken anti-GFP (1:1000, Aves Labs, GFP-1020, AB_10000240), mouse anti-synaptotagmin-2 (1:100; Developmental Studies Hybridoma Bank, AB_2315626). After several washes, according to the mixture of primary antibodies, the following secondary antibodies were used: donkey anti-chicken Alexa 488 (1:500, SA1-72000), donkey anti-rabbit Alexa 555 (1:500, ThermoFisher, A31570), donkey anti-mouse Alexa 488 (1:500, ThermoFisher, A21202), donkey anti-mouse Alexa 647 (1:500, ThermoFisher, A31571). After Hoechst counterstaining, slices were mounted in Fluoromount. Epifluorescence images were obtained with a Zeiss AxioImager Z2 microscope coupled to a camera (Zeiss AxioCam MR3) with an HBO lamp associated with 470/40, 525/50, 545/25, and 605/70 filter cubes. Confocal images were acquired with a Zeiss LSM-800 system equipped with a tunable laser providing excitation range from 405 to 670 nm. For quantifying synaptotagmin-2, 11µm thick stacks were taken (z=1µm, pixel size=0,156µm) with the confocal microscope using a plan-Achromat 40x/1,4 Oil DIC objective.

DATA PREPROCESSING

Motion correction. Image series were motion corrected either by finding the center of mass of the correlations across frames relative to a set of reference frames (Miri et al., 2011) or using the NoRMCorre algorithm available in the CaImAn toolbox (Pnevmatikakis and Giovannucci, 2017) or both.

Cell segmentation. Cell segmentation was achieved using Suite2p (Pachitariu et al., 2017). Neurons with pixel masks including processes (often the case for interneurons located in the stratum oriens) were replaced by soma ROI manually drawn in ImageJ and matched onto Suite2p contours map using CICADA (**C**alcium **I**maging **C**omplete **A**utomated **D**ata **A**nalysis, source code available on Cossart lab GitLab group ID: 5948056). In experiments performed on *GAD67Cre* animals, tdTomato labelled interneurons were manually selected in ImageJ and either matched onto Suite2p contours map or added to the mask list using CICADA.

Axon segmentation. Axon segmentation was performed using pyAMNESIA (a **Py**thon pipeline for analysing the **A**ctivity and **M**orphology of **NE**urons using **S**keletonization and other **I**mage **A**nalysis techniques, source code available on Cossart lab GitLab group ID: 5948056). pyAMNESIA proposes a novel image processing method based on 3 consecutive steps: 1) extracting the axonal morphology of the image (skeletonization), 2) discarding the detected morphological entities that are not functional

ones (branch validation), and 3) grouping together branches with highly correlated activity (branch
clustering). To extract the skeleton, we first perform 3D gaussian smoothing and averaging of the
recording, producing an image that summarizes it; on this image are then successively applied a
histogram equalization, a gaussian smoothing, an adaptive thresholding, and finally a Lee
skeletonization (Suen et al., 1994), allowing for the extraction of the skeleton mask and the
morphological branches. To ensure the functional unity of the segmented branches, we only kept those
that illuminated uniformly, where uniformity was quantified by the skewness of the pixel distribution
of the branch during a calcium transient (branch validation). To cluster the valid branches based on their
activity, we first extracted their average trace -- being the average image intensity along the branch for
each frame -- and then clustered the branches traces using t-SNE and HDBSCAN algorithms with
Spearman correlation metric (branch clustering).

Cell type prediction. Cell type prediction was done using the DeepCINAC cell type classifier (Denis et
al., 2020). Briefly, a neuronal network composed of a convolutional neural network (CNN) and Long
Short-Term Memory (LSTM) was trained using labeled interneurons, pyramidal cells and noisy cells
to predict the cell type using 100 frames long movie patches centered on the cell of interest. Each cell
was classified as interneuron, pyramidal cell or noise. Cells classified as 'noisy cells' were removed
from further analysis. 'Labelled interneurons' were first kept in a separate cell type category and added
to the interneurons list.

Activity inference. Activity inference was done using DeepCINAC classifiers (Denis et al., 2020).
Briefly, a classifier composed of CNN and LSTM was trained using manually-labelled movie patches
to predict neuronal activation based on movie visual inspection. Depending on the inferred cell type,
activity inference was done using either a general classifier or an interneuron-specific classifier.
Activity inference resulted in a (cells x frames) matrix giving the probability for a cell to be active at
any single frame. We used a 0.5 threshold in this probability matrix to obtain a binary activity matrix
considering a neuron as active from the onset to the peak of a calcium transient.

Behavior. Piezo signals were manually analyzed in a custom-made graphical user interface (Python
Tkinter) to label the onset and offset of 'twitches', 'complex movements' and 'unclassified movements'.
Twitches were defined as brief movements (a few hundred milliseconds-long) occurring within periods
of rest and detected as rapid deflections of the piezo signal. 'Complex movements' were defined as
periods of movement lasting at least 2 seconds; 'unclassified movements' were short movements that
differed from twitches and complex movements. Analysis of video tracking was done using CICADA
and behavior was manually annotated in the BADASS (Behavioral Analysis Data And Some Surprises)
GUI.

Neurodata without border (NWB: N) embedding. For each imaging session, imaging data, behavioral
data, cell contours, cell type prediction, calcium traces and neuronal activity inference were combined
into a single NWB: N file (following the guidelines provided on https://www.nwb.org/). Our NWB: N
data set is accessible on DANDI archive (https://gui.dandiarchive.org/#/) - ref DANDI:000219.

MODELING

Network implementation. We constructed a simple rate model and subsequently a more realistic spiking
network in order to test our hypothesis that an increase in perisomatic inhibition could explain the switch
in network dynamics between the first and second postnatal week. Both models consisted of one
excitatory and one inhibitory population with recurrent interactions (**Figure 5A**, supplementary
methods). The development of perisomatic innervation was simulated by increasing the strength from
inhibitory to excitatory cells (J_{EI}). The external input to the model was composed of a constant and a
white noise term. To estimate the responses to twitch-like inputs, an additional feedforward input
composed of short pulses was fed to the network. In the rate model, the rates represented the population
averaged activities. The spiking network was constructed with Leaky Integrate and Fire (LIF) neurons.
The network connectivity was sparse and each neuron received inputs from randomly selected neurons.
Presynaptic spikes resulted in exponentially decaying postsynaptic currents. All codes used for the
modelling are available at https://gitlab.com/rouault-team-public/somatic-inhibition/

DATA ANALYSIS

Sample-size estimation. This study being mainly exploratory in the sense that the evolution of
population activity in the CA1 region of the hippocampus using large scale imaging has not been
described before, we have not been able to use explicit power calculation based on an expected size
effect.

Histological quantifications. Confocal images of synaptotagmin-2 immunostaining were analyzed
using RINGO (RINGs Observation), a custom-made macro in Fiji. We first performed a max-intensity
projection of the Z-stack images of the top 6 μm from the slice surface, then images were cropped to
restrict the analysis to the pyramidal cells layer. Obtained images were denoised using Fiji "remove
background" option and then by subtracting the mean intensity of the pixels within a manually drawn
ROI in the background area (typically the cell body of a pyramidal neuron). Denoised images were then
binarized using a max-entropy thresholding (Fiji option). Finally, particles with size between 0.4 μm²
and 4 μm² were automatically detected using the Fiji "Analyse particle" option. We then computed the
proportion of the pyramidal cell layer (i.e., surface of the cropped region) covered by positive
synaptotagmin-2 labelling.

In vivo electrophysiology. The neuronal activity recorded from both hippocampi in vivo using a 64-channel amplifier (DIPSI, France) was analyzed post-hoc. Firstly, data was downsampled to 1 kHz to save disc space. The local field potential (LFP) was band-passed (2-100 Hz) using the wavelet filter (Morlet, mother wavelet of order 6) and the common reference was subtracted to exclude the bias produced by volume conducted fluctuations of LFP. Sharp Wave events (SWs) were detected using a threshold approach. Firstly, LFP was bandpassed (2-45 Hz) and the difference between lfp recorded in the strati oriens,pyramidale and radiatum was calculated. Events were considered as SWs if: i) LFP reversion was observed in the stratum pyramidale, ii) their peak amplitude in the resulting trace exceeded the threshold of 4 standard deviations calculated over the entire trace (the threshold corresponds to p values below 0.01). The occurrence rate of SW was calculated over the entire recording and normalized to 1 min. SW co-occurrence was also calculated by cross correlating the SW timestamps from ipsilateral and contralateral hippocampi using a bin size of 10 milliseconds. Spectral analysis was carried out using the Chronux toolbox (Chronux, toolbox, 2010). Spectral power was estimated using direct multi-taper estimators (3 time-bandwidth product and 5 tapers).

Statistics for in vivo electrophysiology. Group comparisons were done using non parametric Wilcoxon rank sum test for equal medians, p-value of 0.05 was considered significant. Variability of the estimates was visualized as shaded bands of standard deviation computed using jackknife.

Vigilance state determination in neonatal mice. All analysis of EMG data was completed using custom scripts in MATLAB. For each experiment, the raw EMG data was first downsampled to 1000 Hz and subsequently high-pass filtered at 300 Hz and rectified. The processed data was then plotted to allow for manual inspection. Consistent with prior reports (Mohns and Blumberg, 2010), the data was primarily composed of alternating periods of high EMG tone (referred to as wakefulness) associated with 'complex' movements as well as periods of low EMG tone associated with a general behavioral quiescence and the presence of periodic brief myoclonic twitches (referred to as 'active sleep' due to the frequent observation of muscle twitches (Mohns and Blumberg, 2010). For vigilance state determination, we therefore utilized a protocol similar to that described previously (Rio-Bermudez et al., 2015). For both the 'high' and 'low' EMG tone conditions, 5 periods, each 1 s in duration, were first sampled from locations spread out over the entire recording length. Data from the samples were then pooled for each condition and the average value of the rectified signal was determined. Next, the midpoint between the average rectified signal values calculated for the 'high' and 'low' EMG tone conditions was determined for subsequent use as a threshold to separate periods of non-wakefulness (below the midpoint threshold value) from periods of wakefulness (above the midpoint threshold value), while the quarter point between these two values was calculated to further separate periods of non-wakefulness into active sleep (below the quarter point threshold value) or a sleep-wake transitory state (above the quarter point threshold value but below the midpoint threshold value). Once these thresholds

were determined, the entire length of data was divided into 1 s non-overlapping bins and the average filtered rectified EMG signal was determined for each. A hypnogram was then created by automatically applying the threshold-derived criteria to the binned averaged data. Data bins scored as being active sleep were further analyzed to determine the presence of muscle twitches; this was accomplished by automatically identifying those bins whose filtered rectified average value exceeded 5 x the mean value determined from the low EMG tone representative samples. As a final step, the hypnogram and filtered rectified EMG signal data were plotted and manually inspected to ensure the accuracy of results. The hypnogram was then incorporated in the final NWB: N file to serve in the definition of the epochs of wakefulness and active sleep.

Analysis of calcium imaging data. Analysis was performed using CICADA, a custom-made python toolbox allowing for the automatic analysis of calcium imaging data in the NWB format (Cossart lab Gitlab, group ID 5948056, Project ID: 14048984). CICADA can be installed following the installation guidelines presented at https://gitlab.com/cossartlab/cicada. Each figure panel resulting from an analysis performed in CICADA is associated with a configuration file provided as supplemental data. These configuration files can be loaded in CICADA with the option 'Load a set of parameters' allowing for the replication of the analysis.

Calcium transient frequency analysis. Analysis launched from CICADA 'Transient's frequency' analysis. The transient frequency for each cell was computed using the count of calcium transient onsets divided by the duration of the recording and was then averaged across all cells imaged in one given mouse pup across one or more imaging sessions.

Synchronous Calcium Event (SCE) detection. Analysis launched from CICADA 'SCE description' analysis. SCEs were defined as the imaging frames within which the number of co-active cells exceeded the chance level as estimated using a reshuffling method. Briefly, an independent circular shift was applied to each cell to obtain 300 surrogate raster plots. We computed the 99th percentile of the distribution of the number of co-active cells from these surrogates and used this value as a threshold to define the minimal number of co-active cells in a SCE. Peak of synchrony above this threshold separated by at least 5 imaging frames (500 ms) were defined as SCE frames. To compute the percentage of transients within SCEs we counted, for each cell, the number of its calcium transients (from onset to peak) crossing SCE frames and divided it by its total number of calcium transients. We averaged the obtained values over all the cells imaged per animal.

Peri-Movement-Time-Histograms (PMTH). Analysis launched from CICADA 'Population level PSTH' analysis. A 20-second-long time-window centered on movement onset was used. For each movement within an imaging session, the number of cells activated or the sum of all cell's fluorescence was calculated for each time bin in that 20-second-long window. We obtain as many values as

movements per time bin; for each individual imaging session the 25th, the median and 75th percentiles of the distributions of these values per time bin are computed and divided by the number of imaged cells. To display the percentage of active cells at a given time bin, these values were multiplied by 100. To combine imaging sessions in an age group (*i.e.,* P5, 6, 7, 8, 9, 10, 11, 12) all the median PMTHs from individual imaging sessions belonging to the given group were stacked and we represented at each time bin the 25th percentile, the median and the 75th percentile value of these median PMTHs. To evaluate chance level around movement onsets, 500 surrogate raster plots per imaging session were computed, and the above procedure was used to obtain chance level in each imaging session and then grouped. PMTH obtained from fluorescence signals were built from DF/F calcium traces.

Movement-related inhibition. Analysis launched from CICADA 'Activity ratio around epochs' analysis. A 4 seconds long window centered on the onset of movements was used. The total number of cells activated during this time period was calculated. If less than 40 % of these cells were activated within 2 seconds following movement onset, the movement was classified as an 'inhibiting' movement. This procedure was applied to all detected movements to obtain for each mouse pup the proportion of 'inhibiting' movements.

Movement- and immobility-associated cells. Analysis launched from CICADA 'Epoch associated cells' analysis. The number of transients per cell occurring during movement or immobility was calculated. These transient onsets were then circularly shifted 100 times and the same calculation was performed on each roll. We used the 99[th] percentile of this distribution as a threshold above which the cell was considered as associated with movement or immobility. Finally, the proportion of cells associated with rest or immobility was calculated for each imaged mouse.

Statistics. Statistical tests were performed using GraphPad (Prism).

Perisomatic inhibition onto pyramidal neurons in CA1 is established during early stages of development. Firstly, we investigated the effect of perisomatic inhibition on response to twitch-like inputs in a rate model. It allowed us to analytically characterise and compare the changes in the dynamics of the model under the conditions of weak and strong inhibition. Secondly, we show through numerical simulations that similar changes can be observed in a more realistic spiking network model. We show that increasing the strength of perisomatic inhibition in our models can account for the experimentally observed decrease of responses to twitch like feedforward inputs.

Rate model

The rate model consists two populations (Excitatory and Inhibitory) with interaction strengths J_{ab}, $\left(a, b \in \{E, I\}, J_{ab} > 0\right)$. They receive feedforward input of strengths $J_{a0} > 0$, from an external excitatory population with average firing rate r_0. The rates $\left(r_E, r_I\right)$ represent the population averaged activities. The time scale of their evolution is determined by τ_E and τ_I and follow,

$$\begin{bmatrix} \dot{r}_E \\ \dot{r}_I \end{bmatrix} = \begin{bmatrix} -\frac{r_E}{\tau_E} \\ -\frac{r_I}{\tau_I} \end{bmatrix} + f\left(\begin{bmatrix} J_{EE} & -J_{EI} \\ J_{IE} & -J_{II} \end{bmatrix} \begin{bmatrix} r_E \\ r_I \end{bmatrix} + \begin{bmatrix} J_{E0} \\ J_{I0} \end{bmatrix} r_0 \right)$$

Where, $f(x)$ is the neuronal transfer function. We choose it to be threshold linear. i.e., $f(x) = [x]_+$. For positive x, the dynamics in matrix notation can be written as,

$$\dot{r} = (J - I/\tau)r + r_0 = Mr + r_0$$

The fixed point of this system is given by,

$$r^* = -M^{-1}r_0$$

Where,

$$M^{-1} = \frac{1}{\det(M)} \begin{bmatrix} -J_{II} - \frac{1}{\tau_I} & J_{EI} \\ -J_{IE} & J_{EE} - \frac{1}{\tau_E} \end{bmatrix}$$

$$\det(M) = J_{EI}J_{IE} - J_{EE}J_{II} + \frac{J_{II}}{\tau_E} - \frac{J_{EE}}{\tau_I} + \frac{1}{\tau_E\tau_I}$$

Linear stability

The fixed point r^* is stable to small perturbations if the real part of all the eigenvalues of the Jacobian matrix $M(r^*)$ are negative. The eigenvalues $\lambda_{\pm}$ can be expressed as,

$$\lambda_{\pm} = \frac{1}{2}\left(Tr(M) \pm \sqrt{Tr(M)^2 - 4\,det(M)}\right)$$

where,

$$Tr(M) = J_{EE} - J_{II} - \frac{1}{\tau_E} - \frac{1}{\tau_I}$$

Equivalently, the system is always stable if $det(M) > 0$ and $Tr(M) < 0$.

Requiring, $r_e^* > 0$ and $r_i^* > 0$ gives the conditions,

$$(J_{II} + \frac{1}{\tau_I})J_{E0} - J_{EI}J_{I0} > 0 \implies \frac{J_{E0}}{J_{I0}} > \frac{J_{EI}}{J_{II} + 1/\tau_I}$$

$$J_{IE}J_{E0} - (J_{EE} - \frac{1}{\tau_E})J_{I0} > 0 \implies \frac{J_{E0}}{J_{I0}} > \frac{J_{EE} - 1/\tau_E}{J_{IE}}$$

$det(M) > 0$ gives,

$$-(J_{EE} - \frac{1}{\tau_E})(J_{II} + \frac{1}{\tau_I}) + J_{EI}J_{IE} > 0 \implies \frac{J_{EI}}{J_{II} + 1/\tau_I} > \frac{J_{EE} - 1/\tau_E}{J_{IE}}$$

Combining the inequalities above gives the constraints for stable non-zero rates,

$$\frac{J_{E0}}{J_{I0}} > \frac{J_{EI}}{J_{II} + 1/\tau_I} > \frac{J_{EE} - 1/\tau_E}{J_{IE}}$$

When the solutions are stable, small perturbations will decay to zero. Twitches can be considered as perturbations around the fixed point. The transient response to such short impulses can be expressed as $r(t) = C_1 exp(\lambda_+) + C_2 exp(\lambda_2)$.

For a system with external white noise , $\xi(t)$ as input we have,

$$\dot{r} = Mr + r_0 + \sqrt{\Sigma}\,\xi(t)$$

$$\langle \xi(t) \rangle_t = 0, \quad \langle \xi(t)\xi(s) \rangle_t = \delta(t - s)$$

we set the off diagonal elements to zero, i.e, $\Sigma_{ij} = 0$ if $i \neq j$.

Let $\delta r(t) = r(t) - r^*$, so the linearized system is,

$$\dot{\delta r} = M \delta r + \sqrt{\Sigma}\, \xi(t) dt$$

This can be seen as a 2D Ornstein - Ulenbeck process defined as:
$dx(t) = - A\, x(t)\, dt + B\, dW(t)$, which is well documented (Gardiner, 1985, p 109-111) and we can immediately write down the expression for the covariance matrix $C(\tau)$. With $(A = - M)$ we have,

$$C(\tau) = \langle \delta r(t) \delta r^T(t + \tau) \rangle = exp[-A\tau]\, \sigma$$

$$\sigma = \frac{\det \mathbf{A}\, \mathbf{\Sigma} + [\mathbf{A} - \mathbf{Tr(A)1}]\, \mathbf{\Sigma}\, [\mathbf{A} - \mathbf{Tr(A)1}]^{\mathbf{T}}}{2\mathbf{Tr(A)}\, \det(\mathbf{A})}$$

Spiking model: LIF network

Excitatory and inhibitory neurons are modelled as Leaky Integrate and Fire (LIF) neurons. The LIF network consists N_E excitatory neurons and N_I inhibitory neurons with exponentially decaying postsynaptic currents. Each neuron receives exactly K_E excitatory and K_I inhibitory inputs from randomly selected neurons in the network. And we assume that the network is sparse i.e. $K << N$.

The evolution of sub-threshold membrane voltage V_i^a of neuron i in population $a \in \{E, I\}$, is given by,

$$\frac{dV_i^a}{dt} = -\frac{V_i^a}{\tau_m} + I_{syn} + I_{ext}, \qquad I_{syn} = \sum_{b,j} J_{ij}^{ab} S_j^{ab}$$

$$\frac{dS_j^{ab}}{dt} = -\frac{S_j^{ab}}{\tau_{syn}} + \sum_{t_j^b} \delta\left(t - t_j^b\right)$$

When the membrane voltage reaches the threshold, $V_{thresold}$, it is reset to V_{reset}. τ_{syn} is the synaptic time constant, t_j^b is the spike time of neuron (i, b). The coupling strengths are $J_{ij}^{ab} = J_{ab}$, if there is a connection from neuron (j, b) to (i, a) and zero otherwise. The contribution of external inputs is represented by I_{ext}. I_{syn} represents the total synaptic currents due to spikes. Spikes are modelled as delta functions.

If a neuron (j, b) emits a spike at time t_j^b and projects to a post-synaptic neuron (i, a), this will result in a change of the membrane voltage V_i^a of the post-synaptic neuron by an

amount J_{ab} in a time τ_{syn}. The membrane voltage decays exponentially to its resting potential in a time τ_m.

Simulations and data analysis

The network simulations were conducted using custom code written in Python and C++ and all the analysis was done in Python. We use the forward Euler method to solve the set of coupled ode's with a time step of 0.1ms.

Covariance

Given a stationary stochastic process X_t with mean $\mu_X = E[X_t]$, the autocovariance is given by,

$$C_{XX}(\tau) = E\left[\left(X_t - \mu_X\right)\left(X_{t-\tau} - \mu_X\right)\right]$$

The covariance of X_t with another process Y_t is defined as,

$$C_{XY}(\tau) = E\left[\left(X_t - \mu_X\right)\left(Y_{t-\tau} - \mu_Y\right)\right]$$

Bibliography

Gardiner, C. W. (1985). *Handbook of stochastic methods* (Vol. 3). Berlin: springer.

https://doi.org/10.1080/713821616

ACKNOWLEDGMENTS:

This work was supported by the European Research Council under the European Union's, Horizon 2020
research and innovation program Grant#646925, the Fondation Bettencourt Schueller, and a NWB seed
grant # R20046AA. The project leading to this publication has received funding from the «
Investissements d'Avenir » French Government program managed by the French National Research
Agency (ANR-16-CONV-0001) and from Excellence Initiative of Aix-Marseille University -
A*MIDEX. R.F.D. was funded by the "Ministere de l'Enseignement Supérieur, de la Recherche et de
l'Innovation" and by the Fondation pour la Recherche Médicale Grant FDT202106012824. E.L. was
funded by the "Ministere de l'Enseignement Supérieur, de la Recherche et de l'Innovation" and by
A*Midex foundation and the French National Research Agency funded by the French Government «
Investissements d'Avenir » program (NeuroSchool, nEURo*AMU, ANR-17-EURE-0029 grant). J.D.
was supported by the Fondation pour la Recherche Médicale Grant FDM20170638339. M.P was
supported by the Fondation pour la Recherche Médicale Grant ARF20160936186. T. D and T. S. were
funded by « Investissements d'Avenir » French Government program managed by the French National
Research Agency (ANR-16-CONV-0001) and from Excellence Initiative of Aix-Marseille University
- A*MIDEX. We would like to thank Dr. Pierre-Pascal Lenck-Santini for providing valuable feedback
on our research project. We also would like to thank Marion Sicre for her help with *GAD67Cre*
experiments. We are grateful to Pr. Hannah Monyer for providing the *GAD67Cre* mouse lines. We
thank the Centre de Calcul Intensif d'Aix-Marseille for granting access to its high-performance
computing resources. The rabies virus was a gift from Conzelman laboratory.

AUTHORS CONTRIBUTION:

RFD, MAP, and R.C designed research. RFD and EL performed stereotaxic injections in viral
injections, RFD performed the surgery for window implant, RFD and MAP performed two photon
calcium imaging experiments. CL, MK, AB, RFD and MAP performed histology experiments. MM,
DS, RFD and MAP performed electrophysiological recordings, MM and DS analyzed these recordings.
RFD and RB performed electromyogram recordings coupled with in vivo imaging and RB analyzed the
EMG data. RFD, JD, MAP, and R.C designed the analysis pipeline. RFD and JD, wrote DeepCINAC,
CICADA and RINGO algorithms. TS and TD wrote PyAmnesia. SRB and HR designed and simulated
the network models. RFD, MAP and RC wrote the paper.

REFERENCE LIST:

Anastasiades PG, Butt SJB. 2012. A Role for Silent Synapses in the Development of the
Pathway from Layer 2/3 to 5 Pyramidal *Cell*s in the Neocortex. *Journal of Neuroscience*
32:13085–13099. doi:10.1523/jneurosci.1262-12.2012
Ben-Ari Y. 2002. Excitatory actions of gaba during development: the nature of the nurture.

Nature Reviews Neuroscience **3**:728–739. doi:10.1038/nrn920

Bezaire MJ, Soltesz I. 2013. Quantitative assessment of CA1 local circuits: Knowledge base for interneuron-pyramidal cell connectivity. *Hippocampus* **23**:751–785. doi:10.1002/hipo.22141

Blumberg MS, Sokoloff G, Tiriac A, Rio-Bermudez CD. 2015. A valuable and promising method for recording brain activity in behaving newborn rodents. *Dev Psychobiol* **57**:506–517. doi:10.1002/dev.21305

Bocchio M, Gouny C, Angulo-Garcia D, Toulat T, Tressard T, Quiroli E, Baude A, Cossart R. 2020. Hippocampal hub neurons maintain distinct connectivity throughout their lifetime. *Nature Communications* **11**:1–19. doi:10.1038/s41467-020-18432-6

Buhl DL, Buzsaki G. 2005. Developmental emergence of hippocampal fast-field "ripple" oscillations in the behaving rat pups. Neuroscience **134**:1423–1430. doi:10.1016/j.neuroscience.2005.05.030

Buzsáki G. 2015. Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. Hippocampus **25**:1073–1188. doi:10.1002/hipo.22488

Buzsáki G, Moser EI. 2013. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience* **16**:130–138. doi:10.1038/nn.3304

Caviness VS. 1973. Time of neuron origin in the hippocampus and dentate gyrus of normal and reeler mutant mice: an autoradiographic analysis. *The Journal of Comparative Neurology* **151**:113–120. doi:10.1002/cne.901510203

Cossart R, Khazipov R. 2021a. How development sculpts hippocampal circuits and function. *Physiol Rev*. doi:10.1152/physrev.00044.2020

Cossart R, Khazipov R. 2021b. How development sculpts hippocampal circuits and function. *Physiological Reviews (accepted)*.

Danglot L, Triller A, Marty S. 2006. The development of hippocampal interneurons in rodents. Hippocampus 16:1032–1060. doi:10.1002/hipo.20225

Denis J, Dard RF, Quiroli E, Cossart R, Picardo MA. 2020. DeepCINAC: A Deep-Learning-Based Python Toolbox for Inferring Calcium Imaging *Neuron*al Activity Based on Movie Visualization. *Eneuro* **7**:ENEURO.0038-20.2020. doi:10.1523/eneuro.0038-20.2020

Doischer D, Hosp JA, Yanagawa Y, Obata K, Jonas P, Vida I, Bartos M. 2008. Postnatal Differentiation of Basket Cells from Slow to Fast Signaling Devices. Journal of Neuroscience 28:12956–12968. doi:10.1523/jneurosci.2890-08.2008

Dominguez S, Ma L, Yu H, Pouchelon G, Mayer C, Spyropoulos GD, Cea C, Buzsáki G, Fishell G, Khodagholy D, Gelinas JN. 2021. A transient postnatal quiescent period precedes emergence of mature cortical dynamics 1–34. doi:10.1101/2021.02.17.430487

Donato F, Jacobsen RI, Moser M-B, Moser EI. 2017. Stellate cells drive maturation of the entorhinal-hippocampal circuit. *Science* eaai8178-17. doi:10.1126/science.aai8178

Dooley JC, Glanz RM, Sokoloff G, Blumberg MS. 2020. Self-Generated Whisker Movements Drive State-Dependent Sensory Input to Developing Barrel Cortex. *Curr Biol* **30**:2404-2410.e4. doi:10.1016/j.cub.2020.04.045

Durand GM, Kovalchuk Y, Konnerth A. 1996. Long-term potentiation and functional synapse induction in developing hippocampus. Nature **381**:71–75. doi:10.1038/381071a0

Farooq U, Dragoi G. 2019. Emergence of preconfigured and plastic time-compressed sequences in early postnatal development. Science **363**:168–173. doi:10.1126/science.aav0502

Fuhrmann F, Justus D, Sosulina L, Kaneko H, Beutel T, Friedrichs D, Schoch S, Schwarz MK, Fuhrmann M, Remy S. 2015. Locomotion, Theta Oscillations, and the Speed-Correlated Firing of Hippocampal Neurons Are Controlled by a Medial Septal Glutamatergic Circuit.

Neuron **86**:1253–1264. doi:10.1016/j.neuron.2015.05.001

Gogolla N, Takesian AE, Feng G, Fagiolini M, Hensch TK. 2014. Sensory Integration in Mouse Insular Cortex Reflects GABA Circuit Maturation. Neuron **83**:894–905. doi:10.1016/j.neuron.2014.06.033

Golshani P, Goncalves JT, Khoshkhoo S, Mostany R, Smirnakis S, Portera-Cailliau C. 2009. Internally Mediated Developmental Desynchronization of Neocortical Network Activity. Journal of Neuroscience 29:10890–10899. doi:10.1523/jneurosci.2012-09.2009

Graf J, Zhang C, Marguet SL, Herrmann T, Flossmann T, Hinsch R, Rahmati V, Guenther M, Frahm C, Urbach A, Neves RM, Witte OW, Kiebel SJ, Isbrandt D, Hübner CA, Holthoff K, Kirmse K. 2021. A limited role of NKCC1 in telencephalic glutamatergic neurons for developing hippocampal network dynamics and behavior. *Proc National Acad Sci* **118**:e2014784118. doi:10.1073/pnas.2014784118

Gramsbergen A, Schwartze P, Prechtl HFR. 1970. The postnatal development of behavioral states in the rat. Dev Psychobiol 3:267–280. doi:10.1002/dev.420030407

Hopfield J, Tank D. 2005. Computing with neural circuits: a model. Science **233**:625–633. doi:10.1126/science.3755256

Hopfield JJ. 1982. Neural networks and physical systems with emergent collective computational abilities. Proc National Acad Sci **79**:2554–2558. doi:10.1073/pnas.79.8.2554

Horii-Hayashi N, Sasagawa T, Matsunaga W, Nishi M. 2015. Development and Structural Variety of the Chondroitin Sulfate Proteoglycans-Contained Extracellular Matrix in the Mouse Brain. *Neural Plast* **2015**:1–12. doi:10.1155/2015/256389

Inácio AR, Nasretdinov A, Lebedeva J, Khazipov R. 2016. Sensory feedback synchronizes motor and sensory neuronal networks in the neonatal rat spinal cord. Nature Communications 7:13060. doi:10.1038/ncomms13060

Jiang M, Oliva AA, Lam T, Swann JW. 2001. GABAergic neurons that pioneer hippocampal area CA1 of the mouse: morphologic features and multiple fates. The Journal of Comparative Neurology **439**:176–192. doi:10.1002/cne.1341

Jouvet-Mounier D, Astic L. 1968. [Study of the course of sleep in the young rat during the 1st postnatal month]. *Comptes Rendus Des Séances De La Société De Biologie Et De Ses Fil* **162**:119–23.

Jurgensen S, Castillo PE. 2015. Selective Dysregulation of Hippocampal Inhibition in the Mouse Lacking Autism Candidate Gene CNTNAP2. *J Neurosci* **35**:14681–14687. doi:10.1523/jneurosci.**166**6-15.2015

Karlsson KAE, Mohns EJ, Prisco GV di, Blumberg MS. 2006. On the co-occurrence of startles and hippocampal sharp waves in newborn rats. Hippocampus 16:959–965. doi:10.1002/hipo.20224

Khazipov R, Khalilov I, Tyzio R, Morozova E, Ben-Ari Y, Holmes GL. 2004. Developmental changes in GABAergic actions and seizure susceptibility in the rat hippocampus. *European Journal of Neuroscience* **19**:590–600. doi:10.1111/j.0953-816x.2003.03**152**.x

Lee S-H, Marchionni I, Bezaire M, Varga C, Danielson N, Lovett-Barron M, Losonczy A, Soltesz I. 2014. Parvalbumin-Positive Basket Cells Differentiate among Hippocampal Pyramidal Cells. Neuron 82:1129–1144. doi:10.1016/j.neuron.2014.03.034

Leinekugel X, Khazipov R, Cannon R, Hirase H, Ben-Ari Y, Buzsáki G. 2002. Correlated bursts of activity in the neonatal hippocampus in vivo. Science **296**:2049–2052. doi:10.1126/science.1071111

Lewis DA, Hashimoto T, Volk DW. 2005. Cortical inhibitory neurons and schizophrenia. Nature Reviews Neuroscience 6:312–324. doi:10.1038/nrn1648

Liu K, Kim J, Kim DW, Zhang YS, Bao H, Denaxa M, Lim S-A, Kim E, Liu C, Wickersham IR,

Pachinis V, Hattar S, Song J, Brown SP, Blackshaw S. 2017. Lhx6-positive GABA-releasing neurons of the zona incerta promote sleep. *Nature Publishing Group* **548**:1–24. doi:10.1038/nature23663

Marty S, Wehrlé R, Alvarez-Leefmans FJ, Gasnier B, Sotelo C. 2002. Postnatal maturation of Na+, K+, 2Cl- cotransporter expression and inhibitory synaptogenesis in the rat hippocampus: an immunocytochemical analysis. European Journal of Neuroscience 15:233–245. doi:10.1046/j.0953-816x.2001.01854.x

Milh M, Kaminska A, Huon C, Lapillonne A, Ben-Ari Y, Khazipov R. 2006. Rapid Cortical Oscillations and Early Motor Activity in Premature Human Neonate. *Cerebral Cortex* **17**:1582–1594. doi:10.1093/cercor/bhl069

Miri A, Daie K, Arrenberg AB, Baier H, Aksay E, Tank DW. 2011. Spatial gradients and multidimensional dynamics in a neural integrator circuit. *Nat Neurosci* **14**:1150–1159. doi:10.1038/nn.2**888**

Mòdol L, Bollmann Y, Tressard T, Baude A, Che A, Duan ZRS, Babij R, García NVDM, Cossart R. 2019. Assemblies of Perisomatic GABAergic Neurons in the Developing Barrel Cortex. Neuron 1–18. doi:10.1016/j.neuron.2019.10.007

Mohns EJ, Blumberg MS. 2010. Neocortical Activation of the Hippocampus during Sleep in Infant Rats. Journal of Neuroscience 30:3438–**344**9. doi:10.1523/jneurosci.4832-09.2010

Mohns EJ, Blumberg MS. 2008. Synchronous Bursts of Neuronal Activity in the Developing Hippocampus: Modulation by Active Sleep and Association with Emerging Gamma and Theta Rhythms. Journal of Neuroscience 28:10134–10144. doi:10.1523/jneurosci.1967-08.2008

Morozov YM, Freund TF. 2003. Postnatal development and migration of cholecystokinin-immunoreactive interneurons in rat hippocampus. Neuroscience 120:923–939. doi:10.1016/s0306-4522(03)00409-3

Muessig L, Lasek M, Varsavsky I, Cacucci F, Wills TJ. 2019. Coordinated Emergence of Hippocampal Replay and Theta Sequences during Post-natal Development. *Current Biology* **29**:834-840.e4. doi:10.1016/j.cub.2019.01.005

Murata Y, Colonnese MT. 2020. GABAergic interneurons excite neonatal hippocampus in vivo. *Science advances* **6**:eaba1430. doi:10.1126/sciadv.aba1430

Murphy KM, Beston BR, Boley PM, Jones DG. 2005. Development of human visual cortex: A balance between excitatory and inhibitory plasticity mechanisms. Dev Psychobiol 46:209–221. doi:10.1002/dev.20053

Orefice LL, Zimmerman AL, Chirila AM, Sleboda SJ, Head JP, Ginty DD. 2016. Peripheral Mechanosensory Neuron Dysfunction Underlies Tactile and Behavioral Deficits in Mouse Models of ASDs. Cell 166:299–313. doi:10.1016/j.cell.2016.05.033

Pachitariu M, Stringer C, Dipoppa M, Schröder S, Rossi LF, Dalgleish H, Carandini M, Harris KD. 2017. Suite2p: beyond 10,000 neurons with standard two-photon microscopy. *Biorxiv* 061507. doi:10.1101/061507

Paredes MF, James D, Gil-Perotin S, Kim H, Cotter JA, Ng C, Sandoval K, Rowitch DH, Xu D, McQuillen PS, Garcia-Verdugo J-M, Huang EJ, Alvarez-Buylla A. 2016. Extensive migration of young neurons into the infant human frontal lobe. Science **354**:aaf7073. doi:10.1126/science.aaf7073

Paxinos, Halliday, Watson, Kassem. 2020. Atlas of the Developing Mouse Brain.

Pnevmatikakis EA, Giovannucci A. 2017. NoRMCorre: An online algorithm for piecewise rigid motion correction of calcium imaging data. *Journal of Neuroscience Methods* **291**:83–94. doi:10.1016/j.jneumeth.2017.07.031

Reh RK, Dias BG, Nelson CA, Kaufer D, Werker JF, Kolb B, Levine JD, Hensch TK. 2020.

Critical period regulation across multiple timescales. Proc National Acad Sci **117**:23242–
23251. doi:10.1073/pnas.1820836117

Rio-Bermudez CD, Blumberg MS. 2021. Sleep as a window on the sensorimotor foundations
of the developing hippocampus. Hippocampus. doi:10.1002/hipo.23334

Rio-Bermudez CD, Kim J, Sokoloff G, Blumberg MS. 2020. Active Sleep Promotes Coherent
Oscillatory Activity in the Cortico-Hippocampal System of Infant Rats. *Cerebral cortex (New*
York, NY : 1991) **30**:2070–2082. doi:10.1093/cercor/bhz223

Rio-Bermudez CD, Sokoloff G, Blumberg MS. 2015. Sensorimotor Processing in the
Newborn Rat Red Nucleus during Active Sleep. J Neurosci 35:8322–8332.
doi:10.1523/jneurosci.0564-15.2015

Rochefort NL, Garaschuk O, Milos R-I, Narushima M, Marandi N, Pichler B, Kovalchuk Y,
Konnerth A. 2009. Sparsification of neuronal activity in the visual cortex at eye-opening.
Proceedings of the National Academy of Sciences of the United States of America
106:15049–15054. doi:10.1073/pnas.0907660106

Soltesz I, Losonczy A. 2018. CA1 pyramidal cell diversity enabling parallel information
processing in the hippocampus. Nature Neuroscience 21:1–10. doi:10.1038/s41593-018-
0118-0

Sommeijer J-P, Levelt CN. 2012. Synaptotagmin-2 is a reliable marker for parvalbumin
positive inhibitory boutons in the mouse visual cortex. *PLoS ONE* **7**:e35323.
doi:10.1371/journal.pone.0035323

Suen CY, Wang PSP, LEE S-W, LAM L, SUEN CY. 1994. Thinning Methodologies for
Pattern Recognition. *S Mach Perc* 239–261. doi:10.1142/9789812797858_0012

Supèr H, Soriano E. 1994. The organization of the embryonic and early postnatal murine
hippocampus. II. Development of entorhinal, commissural, and septal connections studied
with the lipophilic tracer DiI. The Journal of Comparative Neurology 344:101–120.
doi:10.1002/cne.903440108

Szőnyi A, Sos KE, Nyilas R, Schlingloff D, Domonkos A, Takács VT, Pósfai B, Hegedüs P,
Priestley JB, Gundlach AL, Gulyás AI, Varga V, Losonczy A, Freund TF, Nyiri G. 2019.
Brainstem nucleus incertus controls contextual memory formation. Science **364**:eaaw0445-
15. doi:10.1126/science.aaw0445

Tiriac A, Del Rio-Bermudez C, Blumberg MS. 2014. Self-Generated Movements with
"Unexpected" Sensory Consequences. Curr Biol 24:2136–2141.
doi:10.1016/j.cub.2014.07.053

Tyzio R, Represa A, Jorquera I, Ben-Ari Y, Gozlan H, Aniksztejn L. 1999. The Establishment
of GABAergic and Glutamatergic Synapses on CA1 Pyramidal Neurons is Sequential and
Correlates with the Development of the Apical Dendrite 1–11.

Valeeva G, Janackova S, Nasretdinov A, Rychkova V, Makarov R, Holmes GL, Khazipov R,
Lenck-Santini P-P. 2019a. Emergence of Coordinated Activity in the Developing Entorhinal-
Hippocampal Network. Cerebral cortex (New York, NY : 1991) **29**:906–920.
doi:10.1093/cercor/bhy309

Valeeva G, Nasretdinov A, Rychkova V, Khazipov R. 2019b. Bilateral Synchronization of
Hippocampal Early Sharp Waves in Neonatal Rats. *Frontiers in Cellular Neuroscience* **13**:29.
doi:10.3389/fncel.2019.00029

Valero M, Cid E, Averkin RG, Aguilar J, Sanchez-Aguilera A, Viney TJ, Gomez-Dominguez
D, Bellistri E, Prida LM de la. 2015. Determinants of different deep and superficial CA1
pyramidal cell dynamics during sharp-wave ripples. Nature Neuroscience 18:1281–1290.
doi:10.1038/nn.4074

Villette V, Malvache A, Tressard T, Dupuy N, Cossart R. 2015. Internally Recurring

Hippocampal Sequences as a Population Template of Spatiotemporal Information. Neuron
88:357–366. doi:10.1016/j.neuron.2015.09.052

Wang DV, Yau H-J, Broker CJ, Tsou J-H, Bonci A, Ikemoto S. 2015. Mesopontine median
raphe regulates hippocampal ripple oscillation and memory consolidation. Nature
Neuroscience 18:728–735. doi:10.1038/nn.3998

Wickersham IR, Lyon DC, Barnard RJO, Mori T, Finke S, Conzelmann K-K, Young JAT,
Callaway EM. 2007. Monosynaptic Restriction of Transsynaptic Tracing from Single,
Genetically Targeted Neurons. Neuron 53:639–647. doi:10.1016/j.neuron.2007.01.033

Wolfe J, Houweling AR, Brecht M. 2010. Sparse and powerful cortical spikes. *Current*
Opinion in Neurobiology **20**:306–312. doi:10.1016/j.conb.2010.03.006

Yuste R. 2015. From the neuron doctrine to neural networks. *Nat Rev Neurosci* **16**:487–497.
doi:10.1038/nrn3962

Zhang G-W, Sun W-J, Zingg B, Shen L, He J, Xiong Y, Tao HW, Zhang LI. 2018. A Non-
canonical Reticular-Limbic Central Auditory Pathway via Medial Septum Contributes to Fear
Conditioning. Neuron 97:406-417.e4. doi:10.1016/j.neuron.2017.12.010

Zutshi I, Valero M, Fernández-Ruiz A, Buzsáki G. 2021. Extrinsic control and intrinsic
computation in the hippocampal CA1 circuit. Neuron. doi:10.1016/j.neuron.2021.11.015