The impact of housing conditions on porcine adult stem cell populations differ between adipose tissue and skeletal muscle

Audrey Quéméner, Frédéric Dessauge, Marie-Hélène Perruchot, Nathalie Le Floc'h, Isabelle Louveau

PEGASE, INRAE, Institut Agro, Saint Gilles, France

Corresponding author:

Isabelle Louveau

16 Le Clos, 35590 Saint Gilles, France

Email address: isabelle.louveau@inrae.fr

Abstract

Background. In pigs, the ratio between lean mass and fat mass determines production efficiency and is strongly influenced by the number and size of cells in tissues. During growth, the increase in the number of cells results from the recruitment of different populations of multipotent adult stem cells residing in the tissues. We hypothesized that the impact of a hygiene challenge on the proportions of adult stem cells in adipose tissue and skeletal muscle may differ between pigs with different residual feed intake (RFI), a measure of feed efficiency.

Methods. At the age of 11 weeks, Large White pigs from two lines divergently selected for low and high RFI were housed in two contrasting hygiene conditions (good *vs* poor). After six weeks of challenge, pigs were slaughtered (n = 5-9/group). Samples of subcutaneous adipose tissue and *longissimus* muscle were collected, and cells from the stromal vascular fraction (FSV), which includes adult stem cell populations, were isolated from each tissue. Adipose and muscle cell populations from the FSV were phenotyped by flow cytometry using antibodies that targeted different cell surface markers (CD45 to identify hematopoietic cells; CD34, CD38, CD56 and CD140a to identify mesenchymal stem cells (MSC) with adipogenic and/or myogenic potential).

Results. Adipose tissue and muscle shared some common MSC populations although MSC diversity was higher in muscle than in adipose tissue. In muscle, the $CD45^-CD56^+CD34^-CD140a^+$ and $CD45^-CD56^+CD34^+CD140a^+$ cell populations were abundant. Of these two cell populations, only the proportions of $CD45^-CD56^+CD34^+CD140a^+$ cells increased ($P \leq 0.05$) in pigs housed in poor hygiene compared with good hygiene conditions. For the $CD45^-CD56^-CD34^-$ cell population, present in low proportion, there was an interaction between hygiene condition and genetic line ($P \leq 0.05$) with a decrease in low RFI pigs housed in poor hygiene conditions. In adipose tissue, the two abundant MSC populations were $CD45^-CD56^-CD34^-$ and $CD45^-CD56^+CD34^-$. The proportion of $CD45^-CD56^-CD34^-$ cells increased ($P \leq 0.05$) whereas the proportion of $CD45^-CD56^+CD34^-$ tended to decrease ($P \leq 0.1$) in pigs housed in poor conditions. This study shows that the proportions of some MSC populations were affected by hygiene of housing conditions in a tissue-dependent manner in pigs of both RFI lines. Therefore, these cell populations could be targeted to modulate growth and body composition in growing animals.

Introduction

During postnatal growth and development, farm animals like pigs are exposed to a diversity of environmental stimuli (Colditz et al., 2016). Stressful situations like transport and housing in poor hygiene conditions can result in immune system hyper-activation. The animal ability to cope with these different challenges will influence the growth of their productive tissues and subsequently their growth efficiency and robustness. Fat animals are generally less efficient in the conversion of feed into meat (Sillence, 2004). Moreover, as excess body fat triggers the production of immune and inflammatory cells, the lean-to-fat ratio in the body may influence the ability to fight against infection, and thus the robustness of animals (Patience, Rossoni-Serão & Gutiérrez, 2015). Hence, a better control of body composition in growing pigs requires a better understanding of the dynamics and flexibility of muscle and adipose tissue growth when the environment surrounding pigs is modified. The mass of skeletal muscle and adipose tissue is largely determined by the number and size of muscle fibers and adipocytes (Lefaucheur, 2010; Louveau et al., 2016). This increase in cell number results from recruitment of multipotent mesenchymal stromal/stem cells (MSCs) resident in tissues (Dominici et al., 2006). These cells comprise heterogeneous populations of undifferentiated cells able of self-renewal with the capacity to differentiate into cells of the mesenchymal lineage (adipocytes, myocytes or osteocytes for instance). These MSCs have now emerged as important targets to control growth and final body composition in livestock (Johnson, 2013; Dodson et al., 2015).

The identification of MSC populations is based on the use of a combination of cell surface markers and flow-cytometry (Bourin et al., 2013; Relaix et al., 2021). Thus, MSCs are positive for CD105, CD90, CD73 and negative for hematopoietic markers CD11b, CD14, CD34 and CD45. More specifically, both skeletal muscle and adipose tissue cells include resident MSCs that share common cell surface markers. For example, the CD56 marker, also known as a neural cell adhesion molecule (NCAM) and a myogenic marker in muscle (Pisani et al., 2010a; Wilschut et al., 2011; Uezumi et al., 2014), has been also found in MSCs from porcine adipose tissue. Recently, the phenotypic marker CD140a, or platelet derived growth factor receptor alpha (PDGFRα) emerged as an important MSC marker. It has been shown that CD140a$^+$ cells derived from muscle or adipose tissue have adipogenic potential (Lee & Granneman, 2012, Uezumi et al., 2010, 2016; Sun, Berry & Olson, 2017). Finally, CD38, one of the main NAD-degrading

enzymes in mammalian tissues, has been suggested as a novel marker of committed

preadipocytes in mice (Carrière et al., 2017). Other studies have shown that the combination of

CD34 and CD56 markers allow the discrimination between myogenic and adipogenic

progenitors in human and pig muscle (Pisani et al., 2010a,b; Perruchot et al., 2013). Taken

together, these cell surface markers could be used to decipher the functional signatures of muscle

and adipose tissue.

Thus, despite the well-known relationship between adipose tissue and skeletal muscle throughout

life (Crisan et al., 2008; De Carvalho et al., 2019), there is a notable lack of studies investigating

deeply these two tissues. In this context, we propose to characterize and compare MSCs

populations from adipose and skeletal muscle tissues in response to an environmental challenge

in two pig lines divergently selected for feed efficiency (Gilbert et al., 2007). With a genetic

selection mainly based on production criteria for more than 50 years, pigs are potentially more

susceptible to infectious diseases and have a weaker immune response (Knap & Doeschl-Wilson,

2020). In order to determine if genetic selection has an impact on the immune response of the

best performing pigs, a sanitary challenge (good *versus* poor hygiene of housing conditions) was

recently carried out on two lines of pigs divergent for their residual feed intake (RFI), a measure

of feed efficiency (Chatelet et al., 2018). It has been shown that this sanitary challenge reduced

the growth performance of pigs and stimulated their immune system (Le Floc'h et al., 2014). We

hypothesized that hygiene of housing conditions will impact the proportions of hematopoietic

and/or MSC cells in subcutaneous adipose tissue (SCAT) and skeletal muscle and that the

response may differ between pigs with different RFI.

Materials & Methods

Ethical approval/declaration

The experiment was performed in the INRAE UE3P experimental facility (Saint-Gilles, France; (doi.org/10.15454/1.5573932732039927E12) in compliance with the ethical standards of the European Community (Directive 2010/63/EU) and was approved by the regional ethical committee (Comité Rennais d'Ethique en matière d'expérimentation animale or CREEA Rennes). The experiment approval number is APAFIS#494–2015082717314985.

Animals

For this experiment, a subset of pigs was selected from the period 1 of a larger study previously described by Chatelet et al. (2018). Growing Large-White pigs (males and females) originated from the 8th generation of two lines divergently selected for residual feed intake (RFI), a measure of feed efficiency (Gilbert et al., 2007). Briefly, animals from the high RFI (HRFI) and the low RFI (LRFI) lines were weaned at 4 weeks of age. At 11 weeks ($\approx$ 25 kg), half of the LRFI and HRFI pigs were then assigned to either a room with good hygiene conditions and the other half to a room with poor hygiene conditions. Good housing conditions included room cleaning, disinfection, and adoption of strict biosecurity precautions. In contrast, poor hygiene conditions consisted of no cleaning nor sanitation of the room after the previous occupation by non-experimental pigs. All pigs were housed in individual pens (85 x 265 cm) in concrete floor throughout the experimental period and were fed ad libitum a standard diet formulated to meet or exceed the nutritional requirement of growing pigs (9.47 MJ of net energy/kg, starch 44.2%; fat 3.1%; crude protein 15.3%. They had free access to water. After six weeks of challenge and after an overnight fast, animals (n = 5-9/experimental group) were euthanized by electrical stunning and exsanguination. Immediately after slaughter, samples of subcutaneous dorsal adipose tissue (SCAT) and of longissimus muscle were excised and placed in 37°C Krebs-Ringer-bicarbonate-HEPES buffer for cell isolation (Perruchot et al., 2013).

Adipose cell isolation

Cells were isolated from a sample of fresh dorsal subcutaneous adipose tissue by collagenase digestion as previously described (De clercq et al., 1997; Perruchot et al., 2013). Briefly, fresh

adipose tissue was cut into small pieces and dissociated by enzymatic digestion with collagenase

II and XI (800 U/mg; Sigma, St. Quentin Fallavier, France) under shaking in a dry bath for 45

min at 37°C. Then, a centrifugation at 400 g for 10 min was performed to separate floating

adipocytes from the pellet of stromal vascular fraction (SVF) cells. After resuspension, SVF cells

were successively filtered through a 200-µm and 25-µm nylon membranes (Dutscher, Brumath,

France). After isolation, SVF cells were placed in 90% Fetal Calf Serum (FCS) and 10%

dimethyl sulfoxide (1–2 million cells per mL) and frozen at − 150 °C.

Muscle cell isolation

Fresh muscle tissue devoid of visible connective tissue was finely chopped and digested for 3 x

20 min in a Ca^{2+}-free medium containing 0.25% trypsin (Invitrogen, Cergy-Pontoise, France),

1.5 mg/mL collagenase II (PAA Laboratoires, Les Mureaux, France), and 0.1% DNAse (Sigma,

St. Quentin Fallavier, France) in a shaking water bath at 37°C. After centrifugation at 800 g for

10 min at 4°C, cell pellet was re-suspended in ice-cold proliferation growth medium (PGM)

containing DMEM supplemented with 10% FCS (Invitrogen), 10% horse serum (Invitrogen),

and Penicillin 6,25 UI/ml -Streptomycin 6,25 µg/ml, (Fisher Scientific, Illkirch, France). Next,

cell suspension was successively filtered through a 200-µm and 50-µm Nylon mesh (Dutscher,

Brumath, France). After their collection in ice-cold PGM, isolated cells were placed in 90% Fetal

Calf Serum (FCS) and 10% dimethyl sulfoxide (1–2 million cells per mL) and frozen at

− 150 °C.

Flow cytometry analysis

Individual vials of cryopreserved cells were thawed in a 37°C water bath (2–3 min). After

thawing, cells were suspended in 10 mL of FCS (Sigma). After centrifugation at 500 g for 5 min

at room temperature (RT), cells were suspended in 10 mL of DPBS (Fisher Scientific, Illkirch,

France). Cells were then counted with the TC20™ Automated Cell Counter (Bio-Rad, Marnes-

la-Coquette, France). Next, cells were centrifuged at 500 g for 5 min at RT and suspended in 10

mL of FACS buffer. This FACS buffer contained MACSQuant Running Buffer (Miltenyi Biotec,

Paris, France) and 0.5% of MACS® BSA Stock Solution (Miltenyi Biotec). Cells were

dispensed at 50,000 cells for SCAT and 500,000 cells for muscle in each round bottom tube of 5

mL (Sarstedt, Nümbrecht, Germany). After centrifugation at 500 g for 5 min at RT, cells were

suspended in 100 µL of FACS buffer. Cells were incubated in the dark on ice for 15 min with
labeled monoclonal or polyclonal antibodies coupled to different fluorochromes:
allophycocyanine (APC), brilliant violet 421 (BV421), fluorescein isothiocyanate (FITC),
phycoerythrin (PE), phycoerythrin-vio770 (PE-Vio770) (Table 1). After incubation with
antibodies, cells were washed in FACS buffer and centrifuged at 500 g for 5 min at 4°C. Labeled
cells were then suspended in 400 µL of FACS buffer. Labeled cells were analyzed using a
MACSQuant Analyzer 10 flow cytometer (Miltenyi Biotec), which was calibrated daily. A
minimum of 50,000 events was acquired per sample for SCAT and 200,000 events for muscle.
Flow cytometry analysis were performed with appropriate isotype matching negative and Flow
Minus One (FMO) controls. Data were analyzed using the FlowLogic software (Inivai, Mentone,
Victoria, Australia).

Table 1: List of antibodies used for flow cytometry analysis.

Antigen	Antibody	Reference	Supplier
CD34	CD34-PE (clone EP373Y)	ab223930	Abcam
Isotype control	IgG-PE (clone EPR25A)	ab209478	Abcam
CD38	CD38-PE-Vio770 (clone REA 671)	130-119-159	Miltenyi Biotec
Isotype control	IgG1-PE-vio770 (REA 293)	130-113-440	Miltenyi Biotec
CD45	CD45-FITC (clone K252.1E4)	MCA1222F	BIO RAD
Isotype control	IgG1-FITC (clone MCA928F)	MCA928F	BIO RAD
CD56	CD56-BV421 (clone NCAM16.2)	562751	BD Biosciences
Isotype control	(BALB/c)-BV421 IgG2b, κ	562748	BD Biosciences
CD140a	CD140a-APC (clone REA 637)	130-122-942	Miltenyi Biotec
Isotype control	IgG1-APC (REA 293)	130-113-434	Miltenyi Biotec

Statistical analyzes

Statistical analyzes were performed using the GraphPad Prism software (version 8.4.3 (686), San Diego, CA, USA). The data were subjected to the two-way analysis of variance (ANOVA). Hygiene of housing conditions, RFI line and their interactions were considered as fixed effects. Results are presented as mean ± standard error of mean (SEM). Differences were considered statistically significant at P ≤ 0.05 and were discussed as a trend for 0.05 < P < 0.1.

Results

Growth performance and body composition of the studied groups

Growth performance (Chatelet et al., 2018) and body composition (Sierżant et al., 2019) of pigs have been previously described for a greater number of experimental pigs. Data for the subset of pigs investigated in the current study are shown in Table 2. First, there was no significant interaction between housing conditions and RFI lines for pig performance and their body composition. Final body weights, average daily gain and backfat weight did not differ between the experimental groups (P > 0.1). Interestingly, irrespective of housing conditions, the relative loin weight was lower in HRFI pigs than in LRFI pigs (P = 0.005). Finally, irrespective of the RFI line, the loin/backfat ratio was higher (P = 0.045) in pigs raised in poor conditions than in pigs raised in good conditions.

	Good		Poor		*P-values*[1]		
	LRFI	**HRFI**	**LRFI**	**HRFI**	**Hyg**	**Line**	**Hyg x Line**
Initial body weight (kg)	27.8 ± 0.7	25.7 ± 1.7	28.2 ± 0.7	25 ± 1.4	0.402	0.495	0.289
Final body weight (kg)	54.6 ± 2.6	51.1 ± 1.9	54.0 ± 3.4	49.8 ± 3.3	0.783	0.174	0.958
Average daily gain (g/day)	626 ± 67	590 ± 59	603 ± 84	564 ± 48	0.750	0.614	0.919
Relative loin weight (%)	27.6 ± 0.3	27.0 ± 0.3	28.4 ± 0.2	27.2 ± 0.5	0.137	**0.005**	0.274
Relative backfat weight (%)	4.9 ± 0.2	5.0 ± 0.1	4.6 ± 0.3	4.6 ± 0.3	0.125	0.903	0.962
Loin/backfat ratio	5.6 ± 0.2	5.5 ± 0.2	6.5 ± 0.5	6.0 ± 0.3	**0.045**	0.322	0.638

Table 2: Growth performance of low (LRFI) and high (HRFI) residual feed intake pigs housed in good or poor hygiene conditions for six weeks.

Values are means ± SEM (n = 5-9 pigs/experimental group). Boldface highlights significant differences (P ≤ 0.05). [1]Probability values for the effect of hygiene conditions (Hyg), genetic lines (Line), and their interaction.

Description of the phenotyping strategy

To identify the cell populations specific and common to SCAT and muscle, we focused our

analysis on a panel of five cell surface markers: CD45, CD34, CD38, CD56 and CD140a (Figure

1).

Figure 1: Diagram of the strategy used for cell immunophenotyping by flow cytometry.

Adult stem cells were isolated from adipose tissue and skeletal of growing pigs with low and

high residual feed intake and housed in good or poor hygiene conditions for six weeks. A

common analytical strategy was used for both tissues. However, adipose tissue was phenotyped

at the level of triple lineage because of its low quantities of $CD45^-CD56^-CD34^+$ and $CD45^-$

$CD56^+CD34^+$ cells (white squares).

First, gating was performed on living cells that were negative for the CD45 in order to remove

hematopoietic cells positive for the CD45 marker. Next, we analyzed the following combinations

in SCAT and muscle: $CD45^-CD34^+$, $CD45^-CD38^+$, $CD45^-CD56^-$, $CD45^-CD56^+$, $CD45^-CD140a^+$.

Then, we performed a deeper phenotyping for cell populations predominant in both SCAT and

muscle, respectively $CD45^-CD56^-$ and $CD45^-CD56^+$ (Figures 2 and 3). Based on the literature,

we evaluated within these two cell populations, the expression of the CD34 marker as named:

$CD45^-CD56^-CD34^-$, $CD45^-CD56^-CD34^+$, $CD45^-CD56^+CD34^-$ and $CD45^-CD56^+CD34^+$

populations. Given the low proportion (<0.5%) of some cell populations in SCAT, we only

analyzed the expression of the CD38 and CD140a cell surface markers in muscle in order to

highlight the proportion of the following populations: CD45⁻CD56⁻CD34⁻CD38⁺, CD45⁻CD56⁻

CD34⁻CD140a⁺, CD45⁻CD56⁻CD34⁺CD38⁺, CD45⁻CD56⁻CD34⁺CD140a ⁺, CD45⁻CD56⁺CD34⁻

CD38⁺, CD45⁻CD56⁺CD34⁻CD140a⁺, CD45⁻CD56⁺CD34⁺CD38⁺ and CD45⁻

CD56⁺CD34⁺CD140a⁺. Altogether, this gating strategy allowed us to highlight specific and

common cell populations in both at SCAT and muscle.

277
278 **Figure 2: Representative flow cytometry plots illustrating the immunophenotyping strategy**

279 **used for adult stem cells isolated from adipose subcutaneous tissue of growing pigs.** The

280 captions correspond to the cell populations highlighted in red. In addition, the red boxes delimit

281 the positivity threshold of studied cell populations established according to the appropriate

282 isotype and Fluorescence Minus One controls.

Figure 3: Representative flow cytometry plots illustrating the immunophenotyping strategy used for adult stem cells isolated from skeletal muscle tissue of growing pigs. The captions correspond to the cell populations highlighted in red. In addition, the red boxes delimit the positivity threshold of studied cell populations established according to the appropriate isotype and Fluorescence Minus One controls.

Analysis of cell populations identified in subcutaneous adipose tissue

There was a significant proportion of CD45$^+$ cells (23.1%) in SCAT with no difference between

the four experimental groups (Table 3). We next analyzed the occurrence of co-staining for the

four cell surface markers CD56, CD34, CD38 and CD140a in order to further delineate the

different populations within CD45$^-$ cells dissociated from SCAT. With triple staining, the

proportions of the CD45$^-$CD56$^-$CD34$^-$ population was higher (+11.7%, P = 0.034) in poor than in

good hygiene of housing conditions whatever the RFI line (Figure 4A) whereas the proportion of

the CD45$^-$CD56$^+$CD34$^-$ population tended to decrease (-17.2%, P = 0.099) in poor hygiene

conditions compared with good conditions (Figure 4C). Of note, the other cell populations with

double or triple staining, such as CD45$^-$CD56$^+$CD34$^+$ cells (Figure 4E) were present in very low

proportions (<1%) in SCAT (Table 3).

Table 3: Percentage of adult stem cell populations in subcutaneous adipose tissue of low (LRFI) and high (HRFI) residual feed intake pigs housed in good or poor hygiene conditions for six weeks.

Cell populations	Good		Poor		*P-values[1]*		
	LRFI	HRFI	LRFI	HRFI	Hyg	Line	Hyg x Line
CD45$^+$	18.2 ± 7.1	22.7 ± 7.1	26.7 ± 5.1	24.1 ± 5.4	0.434	0.880	0.571
CD45$^-$	81.5 ± 7.1	77.0 ± 7.1	72.7 ± 5.2	75.7 ± 5.4	0.429	0.905	0.558
CD45$^-$CD34$^+$	0.5 ± 0.2	0.3 ± 0.1	0.4 ± 0.1	0.6 ± 0.1	0.530	0.844	*0.071*
CD45$^-$CD38$^+$	0.3 ± 0.1	0.2 ± 0.1	0.4 ± 0.1	0.5 ± 0.1	*0.082*	0.779	0.212
CD45$^-$CD140a$^+$	0.5 ± 0.1	0.1 ± 0.02	0.3 ± 0.1	0.3 ± 0.1	0.607	*0.059*	**0.036**
CD45$^-$CD56$^-$	27.4 ± 6.0	29.5 ± 5.2	42.4 ± 3.7	36.8 ± 5.1	**0.035**	0.724	0.448
CD45$^-$CD56$^-$CD34$^-$	26.8 ± 6.0	29.2 ± 5.2	41.9 ± 3.7	36.4 ± 5.0	**0.034**	0.757	0.442
CD45$^-$CD56$^-$CD34$^+$	0.6 ± 0.2	0.2 ± 0.04	0.5 ± 0.2	0.3 ± 0.1	0.746	0.106	0.591
CD45$^-$CD56$^+$	53.9 ± 12.2	47.3 ± 9.9	30.1 ± 6.2	38.6 ± 10.0	0.101	0.920	0.437
CD45$^-$CD56$^+$CD34$^-$	53.7 ± 12.3	47.1 ± 10.0	29.8 ± 6.2	38.3 ± 10.0	0.*099*	0.922	0.439
CD45$^-$CD56$^+$CD34$^+$	0.2 ± 0.1	0.2 ± 0.1	0.2 ± 0.1	0.3 ± 0.1	0.230	0.653	0.693

Values are means ± SEM (n = 6-9 pigs/experimental group). Boldface highlights significant differences (P ≤ 0.05) and italicized characters show a trend (0.05 < P < 0.10).

[1]Probability values for the effect of hygiene conditions (Hyg), genetic lines (Line), and their interaction.

Figure 4: Percentage of adult stem cell populations in subcutaneous adipose tissue (SCAT)

and muscle from low (LRFI, white bar) and high (HRFI, black bar) residual feed intake

pigs housed in good or poor hygiene conditions. Percentage of cell populations are expressed

as means ± SEM (n= 5-9/experimental group).

Analysis of cell populations identified in skeletal muscle

The proportion of CD45$^+$ cells in muscle was in the range of 20-23% (Table 4) and increased (P = 0.020) in pigs housed in poor conditions compared with pigs housed in good conditions for both RFI lines. Next, we analyzed the expression of CD34, CD38, CD140a and CD56 within the CD45$^-$ cell population (Figures 1 and 3). Despite their high proportions (>50%) in muscle, the CD45$^-$CD34$^+$ and CD45$^-$CD140a$^+$ cell populations did not differ significantly between the four groups. Furthermore, there was an interaction for the CD45$^-$CD56$^-$ cell population (P = 0.011). Interestingly, within this CD45$^-$CD56$^-$ cell population, we analyzed the expressions of CD34 and CD140a. We also observed a significant interaction for the CD45$^-$CD56$^-$CD34$^-$ cell population (P = 0.042) (Figure 4B). For the CD45$^-$CD56$^-$CD34$^-$CD140a$^+$ cells, their proportion tended to decrease (P = 0.072) in poor compared with good hygiene conditions. Moreover, within the CD45$^-$CD56$^+$ cell population, the proportion of CD45$^-$CD56$^+$CD34$^-$ cells tended to be higher in HRFI pigs (+10%, P = 0.101) than in LRFI pigs (Figure 4D). Concerning the CD45$^-$CD56$^+$CD34$^+$ cells, their proportions do not vary whatever the genetic line or the housing conditions (Figure 4F). Finally, by adding the CD140a marker in the analysis, we highlighted that, irrespective of the genetic line, the CD45$^-$CD56$^+$CD34$^+$CD140a$^+$ cells increased (+ 11.3%) in poor conditions (P = 0.052) compared with good conditions (Table 4).

Table 4: Percentage of cell populations in muscle of low (LRFI) and high (HRFI) residual feed intake pigs housed in good or poor hygiene conditions for six weeks.

Cell populations	Good		Poor		*P-values*[1]		
	LRFI	HRFI	LRFI	HRFI	Hyg	Line	Hyg x Line
CD45$^+$	20.1 ± 1.5	18.6 ± 1.4	23.0 ± 0.9	23.1 ± 1.9	**0.020**	0.620	0.560
CD45$^-$	79.0 ± 1.5	80.6 ± 1.4	76.4 ± 1.0	76.1 ± 2.1	**0.030**	0.678	0.552
CD45$^-$CD34$^+$	53.3 ± 7.3	47.9 ± 5.8	58.2 ± 3.7	49.5 ± 4.6	0.583	0.242	0.790
CD45$^-$CD38$^+$	0.1 ± 0.02	0.1 ± 0.02	0.1 ± 0.03	0.3 ± 0.1	0.121	0.447	*0.091*
CD45$^-$CD140a$^+$	55.4 ± 9.7	67.2 ± 6.4	69.1 ± 3.5	67.2 ± 0.8	0.327	0.478	0.327
CD45$^-$CD56$^-$	5.6 ± 0.9	2.6 ± 0.5	2.3 ± 0.3	2.8 ± 0.5	**0.024**	*0.073*	**0.011**
CD45$^-$CD56$^-$CD34$^-$	4.4 ± 0.9	2.0 ± 0.4	1.7 ± 0.3	1.8 ± 0.2	**0.020**	*0.057*	**0.042**
CD45$^-$CD56$^-$CD34$^-$CD140a$^+$	1.8 ± 0.7	1.0 ± 0.3	0.5 ± 0.1	0.5 ± 0.2	*0.072*	0.362	0.351
CD45$^-$CD56$^-$CD34$^+$	1.2 ± 0.5	0.6 ± 0.2	0.5 ± 0.1	1.0 ± 0.4	0.735	0.855	0.115
CD45$^-$CD56$^-$CD34$^+$CD140a$^+$	0.6 ± 0.4	0.3 ± 0.2	0.2 ± 0.04	0.7 ± 0.4	0.943	0.904	0.210
CD45$^-$CD56$^+$	73.3 ± 1.4	77.9 ± 1.5	74.1 ± 1.1	73.2 ± 2.4	0.224	0.245	*0.099*
CD45$^-$CD56$^+$CD34$^-$	19.0 ± 5.9	30.8 ± 6.5	16.2 ± 3.0	24.4 ± 7.1	0.436	0.101	0.764
CD45$^-$CD56$^+$CD34$^-$CD140a$^+$	17.9 ± 5.9	29.3 ± 6.4	15.6 ± 2.9	23.6 ± 7.0	0.499	0.108	0.771
CD45$^-$CD56$^+$CD34$^+$	54.0 ± 6.8	46.8 ± 6.1	57.7 ± 3.6	48.4 ± 5.2	0.652	0.169	0.856
CD45$^-$CD56$^+$CD34$^+$CD140a$^+$	36.3 ± 5.5	37.7 ± 5.4	52.7 ± 3.3	43.0 ± 6.2	**0.052**	0.433	0.298

Values are means ± SEM (n = 5-7 pigs/experimental group). Boldface highlights significant differences (P ≤ 0.05) and italicized character shows a trend (0.05 < P < 0.10). [1]Probability values for the effect of hygiene conditions (Hyg), genetic lines (Line), and their interaction. The proportions of CD45$^-$CD56$^-$CD34$^-$CD38$^+$, CD45$^-$CD56$^-$CD34$^+$CD38$^+$, CD45$^-$CD56$^+$CD34$^-$CD38$^+$ and CD45$^-$CD56$^+$CD34$^+$CD38$^+$ populations were below 0.2%.

Discussion

The current study was conducted to determine whether the exposure of growing HRFI and LRFI pigs to poor hygiene of housing conditions impact the proportions of adult stem cell populations resident in muscle and adipose tissue, two tissues of importance for body composition and meat quality traits. Whatever the species considered, the characterization of adult stem cells in muscle and adipose tissue is fairly recent and is conventionally carried out using flow cytometry which is based on the use of antibodies directed against cell surface proteins (Bourin et al., 2013; Relaix et al., 2021). For the current cell immunophenotyping, we have used five well-known cell surface markers (CD45, CD56, CD34, CD38 and CD140a) to identify specific and common $CD45^-$ stem cell populations in both tissues. Moreover, we analyzed for the first time $CD45^+$ cells, a population of hematopoietic cells (Han et al., 2021) known to be increased after injury in muscle (Wosczyna et al., 2019). Our major finding is that the proportions of several cell populations were affected by hygiene of housing conditions in a tissue-dependent manner in pigs of both RFI lines.

The lower growth performance of pigs housed in poor hygiene conditions compared with pigs housed in good hygiene conditions described with a greater number of animals than in the current subset of animals (Chatelet et al., 2018) and the increase in their loin/backfat ratio was associated with changes in the relative proportions of some cell populations in adipose tissue and skeletal muscle. Even though health and growth performance of the most feed efficient LRFI pigs was less impaired by poor hygiene conditions (Chatelet et al., 2018), our findings at the cellular levels do not support a significant relationship between MSC and the better coping ability of LRFI compared with HRFI pigs. In the current study, there was only a significant interaction for the $CD45^-CD56^-CD34^-$ population in muscle, a relatively small population. For the other cell populations, there was no significant difference between the two lines. This differs slightly from a previous study showing some differences in some cell populations between the two RFI lines (Perruchot et al., 2020).

The current study clearly shows the presence of a relative high proportion of $CD45^+$ cells suggesting the presence of hematopoietic cells in both muscle and adipose tissue as shown in previous studies (Cousin et al., 2016; Wosczyna et al., 2019). Our study also indicates that the

proportion of CD45$^+$ cells in muscle increased with the exposure of pigs to poor hygiene of

housing conditions. This increase may be associated with the inflammation induced by the poor

hygiene of housing conditions. The stimulation of the immune system of pigs housed in these

conditions was assessed by an increase in plasma haptoglobin concentrations, an acute phase

protein (APP), and in white blood cell and granulocyte counts three weeks after the beginning of

the challenge (Chatelet et al., 2018). The origin of this increase in CD45$^+$ cells remains

unknown. It could result from an increase in CD45$^+$ resident cells in skeletal muscle.

Alternatively, it could come from other tissues as reported for MSCs in a recent review (Girousse

et al., 2021). A study based on the investigation of MSCs from adipose tissue and peripheral

blood of Iberian pigs supports an in vitro migration ability of these cells (Calle et al., 2018).

Resident MSCs could circulate toward distant inflamed tissues to allow their repair/regeneration.

Hence, the observed increase in CD45$^+$ cells may be associated with changes in local secretory

environment and may affect properties of other muscle cells and affect subsequent growth.

As described previously (Bourin et al., 2013; Girousse et al., 2021; Relaix et al., 2021) including

in our previous work (Perruchot et al., 2013, 2020), there was a diversity of adult stem cell

populations in skeletal muscle and adipose tissue of the different groups of animals. We

confirmed that CD45$^-$CD56$^+$ cells are predominant in both muscle and adipose tissue as observed

before (Perruchot et al., 2013). In porcine muscle, these cells are mainly myogenic (Perruchot et

al., 2013). In human and pig, CD45$^-$CD56$^+$CD34$^-$ were identified as satellite cells (Wilschut et

al., 2008; Pisani et al., 2010b; Lewis et al., 2014). In adipose tissue, the presence of CD45$^-$CD56$^+$

is much less documented. We further highlighted a higher cell diversity in muscle than in

adipose tissue. There was especially a high proportion of CD34$^+$ and CD140a$^+$ cells within the

CD45$^-$ cells. In pigs, CD45$^-$CD34$^+$ cells have been previously identified in muscle as PW1$^+$

interstitial cells (PICs) with an ability to differentiate into skeletal myoblast/myotubes, smooth

muscle, and endothelial cells (Lewis et al., 2014). The CD45$^-$CD140a$^+$ cells were also located in

the interstitial space of muscle. In normal skeletal muscle and damaged skeletal muscle,

CD140a$^+$ cells exhibit fibrogenic and adipogenic potentials and were named fibro/adipogenic

progenitors (FAPs; Uezumi et al., 2010, 2014; Uezumi, Ikemoto-Uezumi & Tsuchida, 2014). In

muscle of growing pigs, we have shown that CD56$^+$CD140a$^+$ and CD56$^+$CD140a$^-$ were mainly

myogenic (Perruchot et al., 2020). Muscle stem cell heterogeneity has been described at several

other levels, including transcription factor, clonal capacity, metabolism, age dependent myogenic

potential, and recently functional response to environmental stress (Cho & Doles, 2019; Relaix et

al., 2021). This multitude of resident stem cells contributes likely together to the remarkable

regeneration capacities of muscle after injury (Mackey et al., 2017; Relaix et al., 2021). We also

included the CD38 marker in our analysis to determine whether it could be used to identify

committed preadipocytes as described in mice (Carrière et al., 2017). However, the very low

number of cells expressing this marker does not support these findings. The discrepancy between

the two studies may be related to the fact that the CD38 marker was detected in cells from

abdominal adipose tissue, an adipose depot exhibiting a greater expression of CD38 than in

SCAT.

We highlighted for the first time that the environment influence MSCs in growing pigs. The

CD45$^-$CD56$^+$CD34$^+$CD140a$^+$ cell population, one of the abundant populations in muscle,

increased clearly after an exposure of pigs to poor hygiene of housing conditions. We previously

found that CD56$^+$CD140a$^+$cells were myogenic (Perruchot et al., 2020). With the use of the

CD34 and CD56 markers, CD34$^+$ cells derived from human muscle were described to be myo-

adipogenic bipotent progenitors (Pisani et al., 2010a,b). Thus, we can hypothesize that these cells

exhibit myo/adipogenic potentials. It is interesting to note that CD56$^+$CD140a$^+$ cells were also

affected by a high-fat/high-fiber diet in porcine muscle (Perruchot et al., 2020). This suggests

that these cells may respond to changes of different factors. The current study further indicates

that there was an interaction between hygiene conditions and RFI lines for CD45$^-$CD56$^-$ cells

including CD45$^-$CD56$^-$CD34$^-$ cells representing a low proportion of cells in muscle. There was a

decrease of cells in poor hygiene condition only in LRFI pigs. These cells may be perivascular

cells as described for CD146$^+$CD34$^-$CD45$^-$CD56$^-$ cells in multiple human tissues including

muscle (Crisan et al., 2008; Shenoy & Bose, 2018). These cells may have a myo-adipogenic

potential. Indeed, we have shown that CD56$^-$CD140a$^+$ cells in porcine muscle exhibited both

adipogenic and myogenic potentials (Perruchot et al., 2020). Moreover, Shenoy & Bose (2018)

showed that pericytes from mouse liver were able to regenerate muscle fibers in muscle of

dystrophic (Mdx) mice. In addition, it has been shown in skeletal muscle of obese humans that

the CD56$^-$ cell fraction gives rise to white adipocytes (Laurens et al., 2016). Other studies have

reported the existence of adipogenic progenitors, not expressing the CD56 marker, in skeletal

muscle MSCs derived from lean individuals in humans (Vauchez et al., 2009; Lecourt et al.,

2010). Altogether, available data obtained in pigs and humans are consistent with a myo-

adipogenic potential of the CD45$^-$CD56$^-$CD34$^-$ cells.

Adipose tissue cell composition is less complex than skeletal muscle with two abundant CD45$^-$

cell populations, the CD45$^-$CD56$^-$CD34$^-$ and the CD45$^-$CD56$^+$CD34$^-$ populations. The proportion

of CD45$^-$CD56$^-$CD34$^-$ cells increased whereas the proportion of CD45$^-$CD56$^+$CD34$^-$ tended to

decrease in degraded hygiene conditions. The presence of CD45$^-$CD56$^+$CD34$^-$ cells in adipose

tissue is poorly documented in the literature. In two previous studies, we provided evidence that

CD56$^+$ cells from subcutaneous adipose tissue were less adipogenic than CD56$^-$ cells (Perruchot

et al., 2013, 2020). Some groups did not find any CD56$^+$ cells in adipose tissue, while others

attribute the expression of CD56 to natural killer cells (CD45$^+$). Overall, there is no consensus on

the expression of the CD56 marker in adipose tissue within the CD45$^-$ cells. Moreover, the

increase in CD45$^-$CD56$^-$CD34$^-$ cells in poor hygiene condition may increase later adipose tissue

growth in this group of pigs. Another possible hypothesis would be that the poor hygiene

condition of pig housing may lead to a delay in the development of adipose tissue, and therefore

the pool of CD45$^-$CD56$^-$CD34$^-$ cells, potential adipocyte precursors, would be less used in these

animals compared with those housed in good condition. Altogether, these findings are consistent

with studies demonstrating that the composition of the MSC compartment of adipose tissue and

skeletal muscle can be affected by several factors including nutritional factors (Perruchot et al.,

2020) during growth.

Conclusions

The findings gathered in this study clearly show that the relative proportions of hematopoietic

and of some MSC populations were affected by hygiene of housing conditions in a tissue-

dependent manner in pigs of both RFI lines. It further indicates that these cell populations may

play a significant role in adipose tissue and skeletal muscle homeostasis and can be targeted to

modulate growth and body composition in growing animals. Further investigations will be

needed to get a better understanding of cell interactions in both tissues.

Acknowledgements

The authors are very grateful to the staff of the experimental pig facilities from INRAE at UE3P (Saint-Gilles, France) and GenESI (Le Magneraud, Surgères, France) for animal care and slaughtering. They are also grateful to Hélène Gilbert (INRAE, UMR GenePhySe, Toulouse, France) for selection of the pig RFI lines. The authors also wish to thank Frédérique Mayeur and Christine Tréfeu (PEGASE, Saint-Gilles, France) for their contribution to the isolation of adipose and muscle cells.

Funding

The research leading to these results has received funding from the European Union's Seventh Framework Programme for Research, Technological Development and Demonstration (grant number 613574, PROHEALTH project) to support animal costs and from INRAE to support analytical measurements costs. Audrey Quéméner was supported by a PhD scholarship from INRAE (Phase division) and the research fund of Région Bretagne (France). Funders approved the general objectives of the study but have no roles in its design and data collection nor interfered with data interpretation and conclusions

Data accessibility

Data are available online: https://doi.org/10.15454/IXXVBB

Supplementary material

There is no supplementary material.

Conflict of interest disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article

References

Bourin P, Bunnell BA, Casteilla L, Dominici M, Katz AJ, March KL, Redl H, Rubin JP,

Yoshimura K, Gimble JM. 2013. Stromal cells from the adipose tissue-derived stromal

vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a

joint statement of the International Federation for Adipose Therapeutics and Science

(IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy* 15:641–

648. DOI: 10.1016/j.jcyt.2013.02.006.

Calle A, Barrajón-Masa C, Gómez-Fidalgo E, Martín-Lluch M, Cruz-Vigo P, Sánchez-Sánchez

R, Ramírez MÁ. 2018. Iberian pig mesenchymal stem/stromal cells from dermal skin,

abdominal and subcutaneous adipose tissues, and peripheral blood: in vitro

characterization and migratory properties in inflammation. *Stem Cell Research &*

Therapy 9:178. DOI: 10.1186/s13287-018-0933-y.

Carrière A, Jeanson Y, Côté J, Dromard C, Galinier A, Menzel S, Barreau C, Dupuis-Coronas S,

Arnaud E, Girousse A, Cuminetti V, Paupert J, Cousin B, Sengenes C, Koch-Nolte F,

Tchernof A, Casteilla L. 2017. Identification of the ectoenzyme CD38 as a marker of

committed preadipocytes. *International Journal of Obesity* 41:1539–1546. DOI:

10.1038/ijo.2017.140.

Chatelet A, Gondret F, Merlot E, Gilbert H, Friggens N, Floc'h NL. 2018. Impact of hygiene of

housing conditions on performance and health of two pig genetic lines divergent for

residual feed intake. *animal* 12:350–358. DOI: 10.1017/S1751731117001379.

Cho DS, Doles JD. 2019. Skeletal muscle progenitor cell heterogeneity. In: Birbrair A ed. *Stem*

Cells Heterogeneity in Different Organs. Advances in Experimental Medicine and

Biology. Cham: Springer International Publishing, 179–193. DOI: 10.1007/978-3-030-

24108-7_9.

Colditz IG, Hine BC, Colditz IG, Hine BC. 2016. Resilience in farm animals: biology,

management, breeding and implications for animal welfare. *Animal Production Science*

56:1961–1983. DOI: 10.1071/AN15297.

Cousin B, Casteilla L, Laharrague P, Luche E, Lorsignol A, Cuminetti V, Paupert J. 2016.

Immuno-metabolism and adipose tissue: The key role of hematopoietic stem cells.

Biochimie 124:21–26. DOI: 10.1016/j.biochi.2015.06.012.

Crisan M, Yap S, Casteilla L, Chen C-W, Corselli M, Park TS, Andriolo G, Sun B, Zheng B,

Zhang L, Norotte C, Teng P-N, Traas J, Schugar R, Deasy BM, Badylak S, Bűhring H-J,

Giacobino J-P, Lazzari L, Huard J, Péault B. 2008. A perivascular origin for

mesenchymal stem cells in multiple human organs. *Cell Stem Cell* 3:301–313. DOI:

10.1016/j.stem.2008.07.003.

De Carvalho FG, Justice JN, Freitas EC de, Kershaw EE, Sparks LM. 2019. Adipose tissue

quality in aging: how structural and functional aspects of adipose tissue impact skeletal

muscle quality. *Nutrients* 11:2553. DOI: 10.3390/nu11112553.

De clercq L, Mourot J, Genart C, Davidts V, Boone C, Remacle C. 1997. An anti-adipocyte

monoclonal antibody is cytotoxic to porcine preadipocytes in vitro and depresses the

development of pig adipose tissue. *Journal of Animal Science* 75:1791–1797. DOI:

10.2527/1997.7571791x.

Dodson MV, Allen RE, Du M, Bergen WG, Velleman SG, Poulos SP, Fernyhough-Culver M,

Wheeler MB, Duckett SK, Young MRI, Voy BH, Jiang Z, Hausman GJ. 2015. Invited

review: Evolution of meat animal growth research during the past 50 years: Adipose and

muscle stem cells. *Journal of Animal Science* 93:457–481. DOI: 10.2527/jas.2014-8221.

Dominici M, Le Blanc K, Mueller I, Slaper-Cortenbach I, Marini FC, Krause DS, Deans RJ,

Keating A, Prockop DJ, Horwitz EM. 2006. Minimal criteria for defining multipotent

mesenchymal stromal cells. The International Society for Cellular Therapy position

statement. *Cytotherapy* 8:315–317. DOI: 10.1080/14653240600855905.

Gilbert H, Bidanel J-P, Gruand J, Caritez J-C, Billon Y, Guillouet P, Lagant H, Noblet J, Sellier

P. 2007. Genetic parameters for residual feed intake in growing pigs, with emphasis on

genetic relationships with carcass and meat quality traits. *Journal of Animal Science*

85:3182–3188. DOI: 10.2527/jas.2006-590.

Girousse A, Mathieu M, Sastourné-Arrey Q, Monferran S, Casteilla L, Sengenès C. 2021.

Endogenous mobilization of mesenchymal stromal cells: a pathway for interorgan

communication? *Frontiers in Cell and Developmental Biology* 8. DOI:

10.3389/fcell.2020.598520.

Han TTY, Walker JT, Grant A, Dekaban GA, Flynn LE. 2021. Preconditioning human adipose-

derived stromal cells on decellularized adipose tissue scaffolds within a perfusion

bioreactor modulates cell phenotype and promotes a pro-regenerative host response.

Frontiers in Bioengineering and Biotechnology 9. DOI: 10.3389/fbioe.2021.642465.

Johnson SE. 2013. Growth and development symposium: Participation of adult tissue-restricted

stem cells to livestock growth and development1. *Journal of Animal Science* 91:57–58.

DOI: 10.2527/jas.2012-6103.

Knap PW, Doeschl-Wilson A. 2020. Why breed disease-resilient livestock, and how? *Genetics*

Selection Evolution 52:60. DOI: 10.1186/s12711-020-00580-4.

Laurens C, Louche K, Sengenes C, Coué M, Langin D, Moro C, Bourlier V. 2016. Adipogenic

progenitors from obese human skeletal muscle give rise to functional white adipocytes

that contribute to insulin resistance. *International Journal of Obesity* 40:497–506. DOI:

10.1038/ijo.2015.193.

Le Floc'h N, Knudsen C, Gidenne T, Montagne L, Merlot E, Zemb O. 2014. Impact of feed

restriction on health, digestion and faecal microbiota of growing pigs housed in good or

poor hygiene conditions. *Animal* 8:1632–1642. DOI: 10.1017/S1751731114001608.

Lecourt S, Marolleau J-P, Fromigué O, Vauchez K, Andriamanalijaona R, Ternaux B,

Lacassagne M-N, Robert I, Boumédiene K, Chéreau F, Marie P, Larghéro J, Fiszman M,

Vilquin J-T. 2010. Characterization of distinct mesenchymal-like cell populations from

human skeletal muscle in situ and in vitro. *Experimental Cell Research* 316:2513–2526.

DOI: 10.1016/j.yexcr.2010.04.020.

Lee Y, Granneman J. 2012. Seeking the source of adipocytes in adult white adipose tissues.

Adipocyte 1:230–236. DOI: 10.4161/adip.20804.

Lefaucheur L. 2010. A second look into fibre typing – Relation to meat quality. *Meat Science*

84:257–270. DOI: 10.1016/j.meatsci.2009.05.004.

Lewis FC, Henning BJ, Marazzi G, Sassoon D, Ellison GM, Nadal-Ginard B. 2014. Porcine

skeletal muscle-derived multipotent PW1pos/Pax7neg interstitial cells: isolation,

characterization, and long-term culture. *STEM CELLS Translational Medicine* 3:702–

712. DOI: https://doi.org/10.5966/sctm.2013-0174.

Louveau I, Perruchot M, Bonnet M, Gondret F. 2016. Invited review: Pre- and postnatal adipose

tissue development in farm animals: from stem cells to adipocyte physiology. *Animal*

10:1839–1847. DOI: 10.1017/S1751731116000872.

Mackey AL, Magnan M, Chazaud B, Kjaer M. 2017. Human skeletal muscle fibroblasts

stimulate in vitro myogenesis and in vivo muscle regeneration. *The Journal of Physiology*

595:5115–5127. DOI: https://doi.org/10.1113/JP273997.

Patience JF, Rossoni-Serão MC, Gutiérrez NA. 2015. A review of feed efficiency in swine:

biology and application. *Journal of Animal Science and Biotechnology* 6:33. DOI:

10.1186/s40104-015-0031-2.

Perruchot M-H, Dessauge F, Gondret F, Louveau I. 2020. Response of adult stem cell

populations to a high-fat/high-fiber diet in skeletal muscle and adipose tissue of growing

pigs divergently selected for feed efficiency. *European Journal of Nutrition*. DOI:

10.1007/s00394-020-02418-7.

Perruchot M, Lefaucheur L, Barreau C, Casteilla L, Louveau I. 2013. Age-related changes in the

features of porcine adult stem cells isolated from adipose tissue and skeletal muscle.

American Journal of Physiology-Cell Physiology 305:C728–C738. DOI:

10.1152/ajpcell.00151.2013.

Pisani DF, Clement N, Loubat A, Plaisant M, Sacconi S, Kurzenne J-Y, Desnuelle C, Dani C,

Dechesne CA. 2010a. Hierarchization of myogenic and adipogenic progenitors within

human skeletal muscle. *Stem Cells* 28:2182–2194. DOI:

https://doi.org/10.1002/stem.537.

Pisani DF, Dechesne CA, Sacconi S, Delplace S, Belmonte N, Cochet O, Clement N,

Wdziekonski B, Villageois AP, Butori C, Bagnis C, Santo JPD, Kurzenne J-Y, Desnuelle

C, Dani C. 2010b. Isolation of a highly myogenic CD34-negative subset of human

skeletal muscle cells free of adipogenic potential. *STEM CELLS* 28:753–764. DOI:

https://doi.org/10.1002/stem.317.

Relaix F, Bencze M, Borok MJ, Der Vartanian A, Gattazzo F, Mademtzoglou D, Perez-Diaz S,

Prola A, Reyes-Fernandez PC, Rotini A, Taglietti. 2021. Perspectives on skeletal muscle

stem cells. *Nature Communications* 12:692. DOI: 10.1038/s41467-020-20760-6.

Shenoy P, Bose B. 2018. Hepatic perivascular mesenchymal stem cells with myogenic

properties. *Journal of Tissue Engineering and Regenerative Medicine* 12. DOI:

10.1002/term.2503.

Sierżant K, Perruchot M, Merlot E, Le Floc'h N, Gondret F. 2019. Tissue-specific responses of

antioxidant pathways to poor hygiene conditions in growing pigs divergently selected for

feed efficiency. *BMC Veterinary Research* 15:341. DOI: 10.1186/s12917-019-2107-2.

Sillence MN. 2004. Technologies for the control of fat and lean deposition in livestock. *The*

Veterinary Journal 167:242–257. DOI: 10.1016/j.tvjl.2003.10.020.

Uezumi A, Fukada S, Yamamoto N, Ikemoto-Uezumi M, Nakatani M, Morita M, Yamaguchi A,

Yamada H, Nishino I, Hamada Y, Tsuchida K. 2014. Identification and characterization

of PDGFR α + mesenchymal progenitors in human skeletal muscle. *Cell Death &*

Disease 5:e1186–e1186. DOI: 10.1038/cddis.2014.161.

Uezumi A, Fukada S, Yamamoto N, Takeda S, Tsuchida K. 2010. Mesenchymal progenitors

distinct from satellite cells contribute to ectopic fat cell formation in skeletal muscle.

Nature Cell Biology 12:143–152. DOI: 10.1038/ncb2014.

Uezumi A, Ikemoto-Uezumi M, Tsuchida K. 2014. Roles of nonmyogenic mesenchymal

progenitors in pathogenesis and regeneration of skeletal muscle. *Frontiers in Physiology*

5. DOI: 10.3389/fphys.2014.00068.

Vauchez K, Marolleau J-P, Schmid M, Khattar P, Chapel A, Catelain C, Lecourt S, Larghéro J,

Fiszman M, Vilquin J-T. 2009. Aldehyde dehydrogenase activity identifies a population

of human skeletal muscle cells with high myogenic capacities. *Molecular Therapy*

17:1948–1958. DOI: 10.1038/mt.2009.204.

Wilschut KJ, Jaksani S, Dolder JVD, Haagsman HP, Roelen BAJ. 2008. Isolation and

characterization of porcine adult muscle-derived progenitor cells. *Journal of Cellular*

Biochemistry 105:1228–1239. DOI: https://doi.org/10.1002/jcb.21921.

Wilschut KJ, van Tol HTA, Arkesteijn GJA, Haagsman HP, Roelen BAJ. 2011. Alpha 6 integrin

is important for myogenic stem cell differentiation. *Stem Cell Research* 7:112–123. DOI:

10.1016/j.scr.2011.05.001.

Wosczyna MN, Konishi CT, Perez Carbajal EE, Wang TT, Walsh RA, Gan Q, Wagner MW,

Rando TA. 2019. Mesenchymal stromal cells are required for regeneration and

homeostatic maintenance of skeletal muscle. *Cell Reports* 27:2029-2035.e5. DOI:

10.1016/j.celrep.2019.04.074.