1
2
3
4
5   The Ebola virus matrix protein clusters phosphatidylserine, a critical step in viral budding
6
7
8

9   Monica L. Husby[1,2], Souad Amiar[1,2¶], Laura I. Prugar[3¶], Emily A. David[1&], Caroline B.
10 Plescia[1&], Kathleen E. Huie[3&], Jennifer M. Brannan[3], John M. Dye[3], Elsje Pienaar[2,4],
11 Robert V. Stahelin[1,2*]
12
13
14
15 [1] Department of Medicinal Chemistry & Molecular Pharmacology, Purdue University,
16 West Lafayette, Indiana, United States of America
17
18 [2] Purdue Institute of Inflammation, Immunology and Infectious Disease (PI4D), Purdue
19 University, West Lafayette, Indiana, United States of America
20
21 [3] United States Army Medical Research Institute of Infectious Diseases USAMRIID, Fort
22 Detrick, Frederick, Maryland, United States of America
23
24 [4] Weldon School of Biomedical Engineering, Purdue University, West Lafayette, Indiana,
25 United States of America
26
27
28 *Corresponding author
29 E-mail:rstaheli@purdue.edu (RVS)
30
31 [¶] These authors contributed equally to this work
32 [&] These authors also contributed equally to this work

33
34
35 Running title: Targeting PS levels to inhibit EBOV
36
37 **Keywords:** Ebola virus, fendiline, filovirus, matrix protein, phosphatidylserine, plasma
38 membrane, viral budding, VP40

Abstract

Phosphatidylserine (PS) has been shown to be a critical lipid factor in the assembly and spread of numerous lipid enveloped viruses. Here, we describe the ability of the Ebola virus (EBOV) matrix protein eVP40 to induce clustering of PS and promote viral budding *in vitro*, as well as the ability of an FDA approved drug, fendiline, to reduce PS clustering subsequently reducing virus budding and entry. To gain mechanistic insight into fendiline inhibition of EBOV replication, multiple *in vitro* assays were employed including imaging, viral budding and viral entry assays. Fendiline reduced the PS content in mammalian cells and PS in the plasma membrane, reducing the ability of VP40 to form new virus particles. Further, particles that do form from fendiline treated cells have altered particle morphology and decreased infectivity capacity. These complementary studies reveal the mechanism by which filovirus matrix proteins cluster PS to enhance viral assembly, budding, and spread from the host cell while also laying the groundwork for fundamental drug targeting strategies.

Introduction

Ebola virus (EBOV), which was first discovered in 1976, has been of much concern since the unprecedented 2014-16 outbreak in Western Africa (Breman *et al*, 2016). These fears were exacerbated with the 2018-2020 EBOV outbreak in the Democratic Republic of Congo (>2200 fatalities) as well as a small outbreak there in 2021. The FDA approved an EBOV vaccine (efficacy when administered prior to virus exposure) in 2019 and a monoclonal antibody cocktail in 2020 (Graul *et al*, 2020; Mullard 2020); however, the duration and breadth of these outbreaks underscore the dangers of reoccurring outbreaks and the imminent need to develop small molecule counter measures to treat patients who test positive for EBOV. Further, there is still a large gap in knowledge in how EBOV hijacks host cell components to replicate and spread from cell-to-cell, elucidation of which may identify new drug targets.

In the Filoviridae family, EBOV and Marburg virus (MARV) are lipid enveloped negative-sense single stranded RNA viruses (Banadyga *et al*, 2016; Mühlberger 2007). One commonly overlooked characteristic of many pathogenic viruses, including EBOV and MARV, is their lipid envelope, which is acquired from the host cell they infect. Furthermore, lipid enveloped negative-strand RNA viruses possess limited viral machinery, often encoding for just a handful of viral proteins. Amongst these viral proteins is the multi-functional matrix protein. These matrix proteins, including the VP40 protein of EBOV (eVP40) and MARV (mVP40), are essential to viral assembly and egress (Stahelin 2014; Madara *et al,* 2015). In fact, independent expression of eVP40 or mVP40 leads to the production of virus-like particles (VLPs), nearly indistinguishable from infectious virions (Jasenosky *et al*, 2001; Noda *et al*, 2002; Licata *et al*, 2003). Although these matrix

proteins travel through different trafficking pathways within cells, they coalesce at the

plasma membrane (PM) to form the viral matrix, which directs viral assembly, budding

and the acquisition of their characteristic lipid envelope (Panchal *et al*, 2003; Jasenosky

& Kawoaka 2004; Wang *et al*, 2010; Bornholdt *et al*, 2013; Oda *et al*, 2015; Amiar &

Stahelin 2020; Wan *et al*, 2020). Importantly, phosphatidylserine (PS) has been

implicated in recruiting matrix proteins to the PM and coordinating the assembly of

progeny virions (Adu-Gyamfi *et al*, 2015; Wijesinghe & Stahelin 2015; Bobone *et al*, 2017;

Del Vecchio *et al*, 2018).

While lipids play a critical role in assembly of new viral particles, lipids are also

actively involved in viral entry in a phenomenon known as "apoptotic mimicry". Apoptotic

mimicry is central to the efficient entry of numerous lipid-enveloped viruses (Jemiielity *et*

al, 2013; Moller-Tank *et al*, 2013; Carnec *et al*, 2015). During apoptotic mimicry, PS is

transferred from the inner to the outer leaflet of the PM; this causes PS to become a

component of the outer viral envelope during infection (Adu-Gyamfi *et al*, 2015; Amara &

Mercer 2015; Nanbo & Kawoaka 2010). Subsequently, the exposed PS in the viral

envelope is recognized by target cell receptors for viral uptake, continuing the viral

lifecycle (Kondratowicz *et al*, 2011; Moller-Tank *et al*, 2013; Brunton *et al*, 2019).

The two bilayers of the PM have varying compositions of four main phospholipid

classes asymmetrically distributed across the two bilayers (Bell *et al*, 1981; Van Meer *et*

al, 2008). However, the most abundant anionic lipid within the inner leaflet of the PM is

PS, a frequent participant in peripheral protein recruitment (Cho & Stahelin 2005; Stahelin

et al, 2014; Kerr *et al*, 2018). Extensive work has looked at the dynamic nature of lipids

within the PM, including PS, and their tendency to cluster into domains several hundred

nanometers in size (Fairn *et al*, 2011; Hirama *et al*, 2017; Bobone *et al*, 2017). Clustering

of anionic lipids into domains enriches regions of the PM with anionic charge, creating a

platform for electrostatic interactions at the PM interface for peripheral protein

recruitment. This phenomenon has been reported between PS and the matrix protein of

influenza A virus (Bobone *et al*, 2017). Although significant work has underscored the

importance of PS in filovirus budding and entry (Moller-Tank *et al*, 2013; Moller-Tank &

Maury 2014; Soni & Stahelin 2014; Stahelin 2014; Adu-Gyamfi *et al*, 2015; Del Vecchio

et al, 2018; Brunton *et al*, 2019), the molecular details of the interaction has not been

explored in the context of the lateral organization of PS, matrix assembly or implications

on viral spread.

The FDA approved drug, fendiline, has been reported to reduce PS levels within

the PM inner leaflet (Cho *et al*, 2016; van der Hoeven *et al*, 2018), which was sufficient

to inhibit the oncogenic protein K-Ras PM localization and signaling (van der Hoeven *et*

al, 2013; van der Hoeven *et al*, 2018). Fendiline was initially approved by the FDA in the

1970s as a non-selective calcium channel blocker to treat coronary heart disease (Bayer

et al, 1987); however, these recently identified off target properties were found to be

calcium independent and associated with the indirect inhibition of acid sphingomyelinase

(ASM) (Cho *et al*, 2016; van der Hoeven *et al*, 2018). eVP40 has been shown to utilize

PS for PM localization, assembly, and production of progeny virions (Adu-Gyamfi *et al*,

2015; Del Vecchio *et al*, 2018; Johnson *et al*, 2021); however, detailed molecular insight

into this relationship is lacking. To delineate the molecular architecture and requirements

of PS concentration on VP40 assembly, oligomerization and budding, we employed

biochemical and biophysical assays *in vitro* and in cells. We also tested the ability and

mechanism by which eVP40 clusters PS *in vitro* and in cells. We hypothesized that

reduction of PS from the PM with fendiline treatment would perturb EBOV assembly and

inhibit viral budding. Lastly, fendiline treatment was tested as a potential therapy for

inhibition of EBOV budding and spread in biosafety level (BSL)-2 and BSL-4 models of

infection.

Results

EBOV VP40 localizes to PS enriched regions in synthetic membranes and in living

cells

eVP40 has been reported to have selectivity and high affinity for PS (Ruigrok *et al*,

2000; Scianimanico *et al*, 2000; Adu-Gyamfi *et al*, 2015; Del Vecchio *et al*, 2018) and

PI(4,5)P$_2$ (Johnson *et al*, 2016; Johnson *et al*, 2018), both *in vitro* and in cells. To establish

an *in vitro* system for eVP40-PS colocalization, we employed a giant unilamellar vesicle

(GUV) system with fluorescently labelled His$_6$-eVP40 and a fluorescent PS molecule,

TopFluor®   TMR-PS   (tetramethylrhodamine-PS   (TMR-PS)).

Dipalmitoylphosphatidylserine (DPPS) was chosen for this assay as eVP40 associates

with saturated PS containing membranes *in vitro* more significantly than PS mono- or

diunsaturated containing membranes (data not shown).   In GUVs supplemented with

40% DPPS, a homogeneous ring structure of eVP40-Alexa488 surrounding the GUV

membrane (Fig EV1A, *middle panel*) was observed and a random plot profile analysis

indicated a small overlap of the two fluorescent signals at the membrane (Fig EV1C,

indicated by the asterisk). To investigate further the ability of eVP40-Alexa488 to

associate with membranes similar to the PM, GUVs containing both 40% DPPS and 2.5%

PI(4,5)P$_2$ were used (Fig EV1A *right panel*)(See Material and Methods section for

important temperature preparation guidelines). The plot profile analysis of the image in

Expanded View Fig 1A revealed a strong overlap between the two fluorescence signals

(eVP40 and TMR-PS) (Fig EV1D). The protein enrichment analysis also supported this

previous observation with an enrichment index of ~5.4 $\pm$ 2, four times more than

membranes devoid of PI(4,5)P$_2$ (Fig EV1E). These data are in agreement with the

previously published findings that eVP40 requires both PS and PI(4,5)P$_2$ for efficient

membrane binding and oligomerization and suggest eVP40 is able to enrich PS at sites

of VP40 oligomerization.

To expand upon our findings, we investigated if eVP40 localized to PS enriched

regions within the plasma membrane of live cells. To visualize PS and protein localization

simultaneously, we transiently expressed EGFP-fused proteins in HEK293 cells and

supplemented the cells with TMR-PS immediately prior to imaging (Fig EV2A). To test

the hypothesis that eVP40 localizes to PS enriched regions of the PM, we examined the

plot profile analysis of TMR-PS and eVP40 by expressing functionally unique EGFP fused

proteins: WT-eVP40, Lact-C2 (a PS binding reporter), K224A-eVP40 (a PS-binding

residue mutant (Del Vecchio *et al*, 2018), and WE/A-eVP40 (oligomerization deficient

mutant (Adu-Gyamfi *et al*, 2012; Bornholdt *et al*, 2013). The fluorescence profile of EGFP-

WT-eVP40 vs. TMR-PS revealed a strong overlap between the two fluorophores (Fig

EV2A,D). This cellular data corroborates our *in vitro* data, demonstrating that EGFP-

eVP40 localizes to PS enriched regions of both model membranes and in the PM of cells.

Additionally, there was no significant fluorescence signal overlap between the EGFP-

K224A-eVP40 mutant and TMR-PS (Fig EV2A,E), which supports the requirement of PS

binding for PM localization of eVP40 (Del Vecchio *et al*, 2018). Importantly, random plot

profile analysis revealed a moderate overlap in the fluorescence signals of the

oligomerization deficient mutant WE/A-eVP40 and TMR-PS (Fig EV2A,F). This is

important to note as this protein is still able to interact with PS at the PM, however, is

unable to properly oligomerize (Hoenen *et al*, 2010; Adu-Gyamfi *et al*, 2012; Adu-Gyamfi

et al, 2013). These results suggest VP40 interacts with PS at the PM inner leaflet as a

dimer without significant oligomerization, in line with VP40 *in vitro* lipid-binding (Del

Vecchio *et al*, 2018).

EBOV-VP40 enhances clustering of PS in synthetic membranes and in living cells

Biophysical and molecular studies into PS dynamics in model membranes and

living cells revealed that PS basally distributes into clustered domains enriched with PS

(Fairn *et al*, 2011; Hirama *et al*; 2017; Bobone *et al*, 2017). Interestingly, cellular proteins

such as Annexins are known to significantly enhance the clustering of PS (Menke *et al*,

2005) and viral proteins such as M1 of Influenza A virus have a selectivity for these PS

clusters (Bobone *et al*, 2017). However, detailed examination of PS clustering and

whether filovirus matrix proteins such as eVP40 alter the organization of PS has not yet

been explored.

Confocal 3D reconstruction of GUVs (with DPPS and PI(4,5)P$_2$) indicated different

structures of TMR-PS clusters observed where eVP40-Alexa488 fluorescence was

enriched (Fig EV1A, *three right columns*). Furthermore, the Mander's coefficient index of

correlation was quantified (between the TMR-PS and eVP40-Alexa488) with varying lipid

compositions (Fig 1B) demonstrating a statistically significant increase in PS clustering

when eVP40-Alexa488 was incubated with GUVs containing both DPPS and PI(4,5)P$_2$

(Fig. 1b; *p<0.0001). These results suggest that eVP40 induced PS clustering *in vitro,*

which was significantly enhanced in the presence of both PS and PI(4,5)P$_2$, akin to the

lipid composition typically found in the PM inner leaflet.

Next, we were interested if eVP40 was able to induce PS clustering in synthetic

membranes in the absence of PI(4,5)P$_2$. Therefore, we performed a TMR self-quenching

experiment as described previously (Zhao *et al*, 2010; Wen *et al*, 2018). The TMR

fluorescent group has potent self-quenching properties when two molecules or more are

brought in close distance from each other. We tested the ability of eVP40 to undergo TMR

self-quenching when fluorescent PS was incorporated in lipid vesicles, as a secondary

effect to eVP40-induced PS-clustering. We also tested different eVP40 concentrations

and different PS ratios to investigate if TMR self-quenching was concentration dependent.

The strongest TMR self-quenching in all membranes was observed at 1.5 μM eVP40 (Fig

1C). As expected, higher PS molar ratios resulted in stronger eVP40 induction of TMR

self-quenching except for a high molar ratio of DPPS (60% molar ratio). However, the

addition of PI(4,5)P$_2$ at 2.5% molar ratio to 60% DPPS-containing membranes rescued

TMR self-quenching similarly to 40% DPPS-containing membranes. Further, DPPS-

containing membranes at 60% molar ratio with or without PI(4,5)P$_2$ displayed a maximum

of TMR-PS self-quenching at 2 μM eVP40. These observations may indicate a saturation

of the liposome membranes with eVP40 at high PS concentrations. Altogether, this assay

demonstrated that eVP40 is able to cluster PS, which is enhanced in the presence of

PI(4,5)P$_2$.

183   We next examined if eVP40 enhanced PS clustering in the PM of cells. As

184 previously mentioned, PS selectively localizes into clustered regions (Fairn *et al*, 2011;

185 Hirama *et al*; 2017; Bobone *et al*, 2017); and we were able to detect a basal level of PS

186 enriched clusters in our control GFP expressing cells, with PS clusters accounting for

187 approximately ~8% of the PM (Fig 1D *top panel* and Fig 1E). Our method was further

188 validated by expressing an additional control protein with a glycosylphosphatidylinositol

189 membrane anchor conjugated to GFP (GFP-GPI), which revealed PS clusters in ~8% of

190 the PM area (Fig 1E; Representative image in Fig EV3B *top panel*).

191   We next sought to determine if our technique accurately captured enhanced PS

192 clustering using the established EGFP-Annexin A2 PS reporter (Menke *et al*, 2005). As

193 shown in Fig 1D-E*,* expression of EGFP-Annexin A2 significantly enhanced PS clustering

194 roughly 2-fold, compared to EGFP expressing cells (***p=0.0001). Taken together, these

195 findings corroborate the previously reported effect of Annexin A2 on PS organization

196 (Menke *et al*, 2005), as well as validate the method developed for our assay. In contrast,

197 expression of EGFP-PLCδ-PH (PI(4,5)P$_2$) or EGFP-LactC2 did not significantly alter the

198 extent of PS clustering (Fig 1E). This confirmed that transient expression of fluorescently

199 conjugated lipid-binding proteins is not sufficient to enhance PS clustering at the PM.

200   Next, we evaluated the effect of eVP40 expression on PS organization across the

201 PM. EGFP-WT-eVP40 increased PS clustering by ~2 fold (*p=0.004), similar to the PS

202 clustering observed with Annexin A2 (Fig 1D,E). However, expression of the PS-binding

203 deficient mutant EGFP-K224A-eVP40 showed no significant change in PS clustering (Fig

204 1D,E), supporting the hypothesis that eVP40 must interact with PS to promote its

205 clustering at the PM. Additionally, to investigate if eVP40 matrix oligomerization was

important for PS clustering, we expressed EGFP-WE/A-eVP40 in HEK293 cells. It is

important to note that this mutant still colocalizes with PS at the PM (Fig EV2A,F) albeit

to a lesser extent than WT (Adu-Gyamfi *et al*, 2012). Although the WE/A-eVP40 and PS

interaction is maintained in cells, no significant increase in PS clustering was observed

(Fig 1D,E). To the best of our knowledge, this is the first account of a filovirus matrix

protein modulating the organization of PS within the PM. Moreover, these results

demonstrate that both membrane binding and oligomerization of eVP40 is central to

eVP40-mediated PS clustering.

eVP40 membrane binding and oligomerization are dependent on

phosphatidylserine content in lipid membranes

Next, we hypothesized eVP40 may require PS clustering for productive

interactions at the PM during assembly. Enrichment of PS within regions of the PM would

provide additional PS molecules available to recruit eVP40 to platforms of viral budding.

To investigate how increasing the amount of PS within membranes dictates eVP40

membrane affinity, surface plasmon resonance (SPR) was performed with His_6-eVP40

and large unilamellar vesicles (LUVs) with increasing concentrations of PS (POPC matrix

with PS added from 1% to 22 mol% PS; Fig 2A-C). eVP40 displayed moderate binding to

LUVs with 1% PS, with an apparent affinity of 2.5 µM (Fig 2A). However, increasing the

concentration of PS to 11% increased the apparent affinity of eVP40 to 0.65 µM (Fig 2B).

eVP40 displayed even stronger affinity to vesicles with 22% PS, with an apparent affinity

(K_d) of ~0.18 µM (Fig 2C). These results indicate that by increasing the amount of PS in

membranes, the affinity of eVP40 to lipid membranes can be modulated. This finding

supports the hypothesis that PS clustering may be a mechanism for the virus to provide

the necessary electrostatic contacts needed for matrix assembly during viral production.

Once at the PM, VP40 oligomerizes into the extensive matrix that gives rise to the

stability and structure of the virion. Previously, Adu-Gyamfi et al (Adu-Gyamfi *et al*, 2015)

highlighted the importance of PS in this process, where a cell line deficient in PS synthesis

showed a significant reduction in eVP40 oligomerization. Moreover, our confocal

clustering data (Fig 1D,E) revealed that eVP40 oligomerization is crucial for modulating

PS organization into clustered domains. To investigate how increasing PS concentration

alters eVP40 oligomerization, we utilized chemical crosslinking of His$_6$-eVP40 which had

been incubated with LUVs of increasing PS concentration (Fig 2D,E). Introduction of 15%

PS into LUVs led to extensive eVP40 oligomerization beyond dimeric eVP40 compared

to when 0% PS LUVs were used (Fig 2D *lane 2*, Fig 1E). We next tested LUVs containing

30% and 60% PS and found that eVP40 oligomerization was even more significantly

detected than when just 15% PS was used (Fig 2D *lane 3 and lane 4, respectively*, Fig

1E). Compared to LUVs with 0% PS, both 30% and 60% PS led to a significant increase

in eVP40 oligomerization (*p=0.021 and *p=0.017, respectively). Further, eVP40

oligomerization appeared to saturate when 30% PS was included, as increasing PS

content to 60% did not increase eVP40 oligomerization (compared to 30% PS). Taken

together, these studies suggest a dynamic relationship between PS clustering and eVP40

affinity and oligomerization as a critical step in eVP40 assembly.

Total cellular and plasma membrane levels of phosphatidylserine are reduced by

fendiline treatment

A recent study reported an FDA-approved drug, fendiline, inhibited K-Ras PM

localization and signaling (Cho *et al*, 2016) and reduced PM PS content in MDCK cells

(Cho *et al*, 2016). Therefore, it was our goal to determine if fendiline could also reduce

PS levels in the human cell line HEK293, a cell line commonly used in BSL-2 filovirus

studies. The initial finding that fendiline reduced PS levels within the PM (40% reduction,

IC_{50} ~3uM) was conducted in MDCK cells using thin-layer chromatography (Cho *et al*,

2016), therefore it had not been established if this effect was cell-type specific. To address

this, we first established fendiline's toxicity in HEK293 cells after 24 and 48 hours of

treatment. No significant toxicity was observed in treatments up to 5 µM fendiline (Fig

EV4A). To evaluate the effect of fendiline on PS in HEK293 cells, cells were treated with

fendiline for 48 hours and lipids were extracted and quantified by liquid chromatography-

tandem mass spectrometry (LC-MS/MS). We observed a significant reduction in cellular

PS levels compared to DMSO treated cells, after 48 hour treatment with 1 µM fendiline

(~18% reduction; *p=0.012) and 5 µM fendiline (~30% reduction; ***p=0.0003) (Fig 3A).

Fendiline exhibited no selectivity in reducing different PS species, as 5 µM fendiline

reduced long chain (C>38) and saturated PS species nearly equally (Fig EV4B). It is

important to note that the effect of fendiline on PS was specific in that fendiline treatment

had no significant effect on another anionic phospholipid, phosphatidic acid (Fig EV4C).

Therefore, our data supports the reported finding that fendiline reduced total cellular

levels of PS, and that the effect is not cell dependent.

As PS is an integral anionic component of the PM inner leaflet, we sought to

confirm that fendiline treatment also reduced PS levels within the PM in HEK293 cells.

PS localization within the PM has been readily studied by expressing EGFP-LactC2 in

mammalian cells (Yeung *et al*, 2008; Kay *et al*, 2012). Therefore, HEK293 cells

expressing EGFP-LactC2 were imaged at 24 hours (Fig EV4D) and 48 hours (Fig 3B)

post-treatment with increasing concentrations of fendiline. Single doses of 500 nM

fendiline had no effect on EGFP-LactC2 PM localization at 24 or 48 hours post treatment

(Fig EV4E and Fig 3C, respectively). However, we found a ~30% reduction in PM EGFP-

LactC2 localization after 24 hours of treatment for both 1 µM (**p=0.0003) and 5 µM

fendiline (**p=0.0045) (Fig EV4E). However, a single dose of 1 µM fendiline treatment did

not significantly affect Lact-C2 PM localization after 48 hours of treatment (Fig 3C).

Conversely, a single dose of 5 µM fendiline significantly reduced Lact-C2 PM localization

even at 48 hours post treatment (~30% reduction; **p=0.0031; Fig. 3c), a reduction similar

to that observed at 24 hours post treatment.

Fendiline has been known to target L-type calcium channels in the T-tubules

(invaginations of the plasma membrane) of muscle cells thereby inhibiting calcium

movement and acting as a vasodilator (Bayer & Mannhold 1987). However, in a number

of non-muscle cell lines, fendiline has been shown to increase calcium levels, including

hepatoma cells (Cheng *et al*, 2001), oral cancer cells (Huang *et al*, 2009), PC prostate

cancer cells (Jan *et al*, 2001), and canine kidney cells (MDCK) (Jan *et al*, 2000). Fendiline

was also specific in the K-Ras inhibitory affect among a number of L-type calcium channel

blockers (van der Hoeven *et al*, 2013) strongly suggesting the fendiline mechanism of

action is directly linked to PS depletion and not changes in calcium. Fendiline was shown

to inhibit ASM (van der Hoeven *et al*, 2015) thereby reducing plasma membrane

sphingomyelin, ceramide, PS and cholesterol. Restoration of ASM activity lead to the

relocalization of the PS reporter LactC2 to the PM (van der Hoeven et al, 2015). Lastly,

studies in HEK293 cells, the major cell line used herein, demonstrated endogenous

calcium channel currents could be distinguished from heterologous expression of L-type

calcium channels (Berjukow *et al*, 1996).

Thus, based upon rigor of prior research the most likely hypothesis is that fendiline

inhibits ASM to alter sphingomyelin levels, which are necessary to maintain proper PS

levels in the plasma membrane. Recently, fendiline analogs have been developed with

nanomolar potency and have been shown to displace ASM, which is membrane bound in

acidic lysosomes leading to ASM degradation (Wang *et al*, 2021) While the direct

mechanism of PS depletion in the PM is not fully understood, transbilayer communication

may take place across the plasma membrane between sphingomyelin levels on the outer

leaflet and phosphatidylserine and cholesterol on the inner leaflet (Llorente *et al*, 2013;

Skotland & Sandvig 2019).

Fendiline reduces PS clustering

Next, we hypothesized that reduced levels of PS within the PM would therefore

reduce the degree of PS clustering. To determine if fendiline treatment reduced the

degree of PS clustering, we utilized the Number & Brightness technique (N&B). N&B is a

quantitative fluorescence microscopy technique that allows one to detect the aggregation

state of proteins with pixel resolution in real time (Digman *et al*, 2008). Previously, N&B

was used to quantify PS clustering by analyzing the N&B profile of EGFP-LactC2 (Bobone

et al, 2017). To accurately capture PS clustering at the PM, imaging was performed at a

focal plane near the cell surface (Fig EV4F).

To evaluate PS clustering, HEK293 cells expressing EGFP-LactC2 were treated

with control or fendiline for 48 hours and the EGFP-LactC2 N&B profile was examined

(Fig 3D,E). Three different cluster bin sizes were examined, 1—5, 5-10 and >10. The

average percentage of pixels in each bin was calculated and plotted (Fig 3E). Within

control-treated cells, significant aggregation of EGFP-LactC2 was observed, with 25%

present in complexes of 5-10 LactC2 molecules and ~10% in complexes of >10 LactC2

molecules (Fig 3D *top panel* and Fig 3E). This corroborates previous work investigating

PS clustering, which found EGFP-LactC2 clusters up to 15 molecules in size (Bobone *et*

al, 2017). Treatment of cells with 5 µM fendiline abolished the presence of EGFP-LactC2

complexes (>10 molecules) and significantly reduced the number of complexes of 5-10

LactC2 molecules large (from ~8% in DMSO to ~0% in 5 µM fendiline; **p=0.0043) (Fig

3D *bottom panel* and Fig 3E). Moreover, there was a significant (~23%) increase in

EGFP-LactC2 complexes of ~1-5 molecules in size in cells treated with 5 µM fendiline

compared to control treated cells (****p<0.0001) (Fig 3D *bottom panel* and Fig 3E). Taken

together, this data suggests that fendiline treatment disrupted large PS-dependent

LactC2 complexes which was compensated by an increase in smaller PS-dependent

complexes. Therefore, fendiline may possess antiviral properties by disassembling PS

enriched regions that would otherwise have been used as platforms for viral assembly.

While the mechanism by which fendiline reduces PS clustering is not known, it has

been hypothesized that sphingomyelin changes on the outer plasma membrane can alter

the distribution of PS in the inner leaflet (Skotland & Sandvig 2019). This is thought to be

attributed to the long fatty acid chains of sphingomyelin interacting with the hydrophobic

chains of PS, especially that of PS 18:0/18:1 (Skotland & Sandvig 2019), the most

predominant PS species in the plasma membrane. The changes in cholesterol in the

plasma membrane could also be partially responsible as optimal levels of sphingomyelin

are required for maintenance of cholesterol and PS levels in the plasma membrane inner

leaflet (Maekawa *et al*, 2016 Sci Rep) and PS is necessary for proper cholesterol

distribution across the plasma membrane (Maekawa & Fairn 2015).

Fendiline significantly inhibits authentic EBOV replication

To determine if the FDA-approved drug fendiline was able to inhibit authentic

filovirus replication and spread, we first established the toxicity of fendiline in Vero E6

cells (Fig EV5A) an established model for BSL-4 filovirus studies (Moe *et al*, 1981), and

then monitored the efficacy of fendiline at inhibiting EBOV replication in a BSL-4 setting.

Vero E6 cells were used to examine filovirus replication 48, 72 and 96-hours post-

infection at different multiplicity of infection (MOI) with EBOV (Kikwit) (Fig 4). Following

infection, plates were separated into three post-infection treatment groups (day 0, every

day dosing (e.d.), or every other day dosing (e.o.d.)).

Fendiline was most effective at reducing EBOV infection in vitro at the highest 20

µM concentrations in each treatment group with statistically significant inhibition observed

for EBOV at each time point and each treatment group (excluding EBOV 48 hours,

e.d.)(****$p<0.0001$, **$p<0.0066$) (Fig 4). Percent inhibition was directly affected by timing

of treatments following infection. Furthermore, EBOV infection treatment with fendiline

e.d. had the highest inhibition on viral spread at each time point for 20 µM fendiline

treatments. Cells of the e.o.d. treatments group, which did not receive treatment

immediately following infection with virus, had a dramatically reduced degree of inhibition

as compared to the day 0 and e.d. treatment groups, both of which received fendiline

immediately following viral infection of one hour.

Fendiline reduced EBOV-VP40 localization to the plasma membrane

As both EBOV and MARV-VP40 assembly at the PM is in part governed by PS,

we first analyzed both EGFP-eVP40 and EGFP-mVP40 PM localization in cells treated

with fendiline for 24 or 48 hours. Treatment with 1 µM and 5µM fendiline had no significant

effect on eVP40 PM at 24 hours post treatment (Fig EV5B,C); therefore, EGFP-mVP40

PM localization was not assessed further at 24 hours. Since no significant change in

EGFP-mVP40 PM localization was observed after 48 hours with either 1 µM or 5 µM

fendiline treatment (Fig 5A *bottom panel* and Fig 5B), mVP40 was excluded from further

experiments. In agreement with our results thus far, 1 µM fendiline did not significantly

inhibit EGFP-eVP40 PM localization after 48 hours of treatment (Fig 5A *top panel* and Fig

5C). However, treatment with 5 µM fendiline for 48 hours led to a modest reduction in

EGFP-eVP40 PM localization (~6% reduction compared to control treated cells; not

significant as p=0.08) (Fig 5A *top panel* and Fig 5C). However, the reduction of eVP40

PM localization was not robust enough to lead to the observed inhibition of EBOV by

fendiline treatment in our BSL-4 studies (Fig 4). One possible explanation is that a

limitation of this technique is the inability to differentiate the extent of VP40

oligomerization occurring using basic confocal microscopy. Therefore, it is possible that

fendiline reduced PS levels within the PM, but not significantly enough to block VP40's

ability to bind to the PM.

VP40 oligomerization is significantly reduced by fendiline treatment

PS is also a key factor promoting the self-assembly of VP40 into the matrix layer

of the budding virion (Adu-Gyamfi *et al*, 2013; Adu-Gyamfi *et al*, 2015). This self-assembly

process has been highlighted in our *in vitro* crosslinking data (Fig 2D,E) as well as

previously reported in live cells utilizing the N&B technique (Adu-Gyamfi *et al*, 2012). To

assess how fendiline impacted eVP40 oligomerization in cells, we examined the

oligomerization profile of EGFP-eVP40 using the previously described N&B (Adu-Gyamfi

et al, 2012; Adu-Gyamfi *et al*, 2015; Johnson *et al*, 2016) (Fig 5D,E). The crystal structure,

biochemical analysis, and recent Cryo-ET of eVP40 at the PM strongly suggests eVP40

binds to the PM as a dimer (Bornholdt *et al*, 2013; Del Vecchio *et al*, 2018; Wan *et al*,

2020). eVP40 dimers subsequently oligomerize into larger oligomers via CTD-CTD

interactions between dimers, leading to the eVP40 matrix layer formation (Wan *et al*,

2020). Therefore, for our data to coincide with the current models of eVP40

oligomerization, EGFP-eVP40 oligomers were grouped into bins based on multiples of

the dimer. Since analyzing eVP40 oligomerization by an increase in every dimer leads

to an abundance of data as well as partial overlap in fluorescence signal for each dimer

addition, we simplified the data presentation and analysis to different bins of eVP40

oligomers   (i.e. dimer-hexamer, hexamer-12mer, 12mer-18mer, and >18mer). The

average percentage of pixels in each bin was calculated and plotted for HEK293 cells

expressing EGFP-eVP40 and treated with either the control or indicated concentration of

fendiline for 48 hrs (Fig 5D,E).

Large eVP40 oligomeric structures corresponding to each bin size were readily

detectable at the PM in control treated cells (Fig 5D *top panel* and Fig 5E), with ~72% of

eVP40 found as a dimer-hexamer, ~16% as a hexamer-12mer, ~8% as a 12mer-18mer,

and 3% in complexes >18mer. Treatment with 1 µM fendiline led to a ~8% increase in

dimeric-hexameric eVP40, and small decreases in the larger oligomeric structures,

although no changes were statistically significant (Fig 5D *middle panel* and Fig 5E).

However, the oligomeric profile of eVP40 was statistically different when cells were

treated with 5 µM fendiline. Following 5 µM fendiline treatment there was a significant

increase in eVP40 found in the dimer-hexameric state (~13% increase; **p=0.0035),

which was counterbalanced by an equal reduction in the larger oligomeric states (~6%

reduction for hexamer-12mer, 5% reduction for 12mer-18mer, and ~3% reduction for

eVP40 structures >18mer) (Fig 5D *bottom panel* and Fig 5E). These results support our

hypothesis that by reducing PS concentration and therefore the pool of PS available for

clustering, eVP40 is unable to properly oligomerize once it traffics and binds to the PM.

This, in combination with the modest reduction in eVP40 PM binding following fendiline

treatment, may therefore impact the production of viral particles as suggested from our

BSL-4 studies.

Fendiline reduced VLP production at the plasma membrane

As fendiline reduced VP40 oligomerization, we sought to determine the effect of

fendiline treatment on VLP production using functional budding assays. No significant

effect on VLP production was observed for cells treated with 0.5 µM or 1 µM fendiline at

either 24 (Fig 6A *lane 3,4* and Fig 6B) or 48 hours post-treatment (Fig 6C *lanes 3,4* and

Fig 6D). However, treatment with one dose of 5 µM fendiline for 24 hours led to a ~25%

reduction in VLP production (Fig 6A *lane 5* and Fig 6B) compared to DMSO treated cells

(Fig 6A *lane 2* and Fig 6B). More importantly, this reduction in VLP production was even

more robust when monitored at 48 hours post-treatment, with a statistically significant

~60% reduction in the relative budding efficiency of 5 µM fendiline treated cells (*p-

0.0260) (Fig 6C *lane 5* and Fig 6D). The reduction in VLPs is supported by our previous

findings that a single dose of 5 µM fendiline reduced PS levels, PS clustering and the

extent of eVP40 oligomerization at the PM. Therefore, we hypothesize that reduced virus

budding is at least partially responsible for fendiline efficacy in authentic EBOV studies

(Fig 4).

To further investigate the reduction of VLP production in fendiline treated cells, and

to determine if there were any observable morphological changes in VLPs, scanning

electron microscopy (SEM) experiments were performed (Fig 6E-F). Micrographs of cells

expressing FLAG-eVP40 and treated with 5 µM fendiline revealed minimal VLP

production at the PM compared to control treated cells (Fig 6F). These findings support

the hypothesis that fendiline treatment considerably reduces the production of EBOV

VLPs.

VLP morphology is altered by fendiline treatment

The structure and stability of filoviruses is derived from the VP40 matrix underlying

the lipid envelope of virions. Therefore, we utilized transmission electron microscopy

(TEM) of purified VLPs to determine if disturbing matrix assembly and altering the lipid

components of the PM with fendiline treatment changed VLP morphology and possibly

infectivity (Fig 7A-C). During filoviral entry, surface exposed GP and viral envelope PS

interact with the receptor T-cell immunoglobulin receptor-1 (TIM-1) (Moller-Tank *et al*,

2013; Brunton *et al*, 2019). To recapitulate entry-competent VLPs (eVLPs), we co-

expressed eVP40 with the Ebola virus glycoprotein (eGP). We performed TEM of eVLPs

purified from control and 5 µM fendiline treated cells and used ImageJ software to analyze

VLP length and diameter (Fig 7A-C). Control eVLPs were heterogenous in length with a

mean length of 4.1 µm ± 2.9 (Fig 7A *left panel* and Fig 7B). Control eVLPs diameter also

exhibited a level of heterogeneity but had a fairly consistent diameter of 75 nm ± 12.9,

which is similar to previous studies of both virions and VLPs (Noda *et al*, 2002; Liu 2014)

(Fig 7A *left panel* and Fig 7C). The length and diameter of eVLPs derived from 5 µM

fendiline treated cells were significantly less than control eVLPs. Strikingly, fendiline

treatment reduced eVLP length by ~35%, from 4.1 µm to 2.7 µm (*p=0.0139) (Fig 7A *right*

panel and Fig. 7B) and modestly reduced eVLP diameter (*p=0.043) (Fig 7A *right panel*

and Fig 7C). To ensure that eVLPs derived from fendiline treated cells were not more

susceptible to damage during the purification, circular dichroism thermal melting was

performed and no difference in eVLP stability was observed (Appendix Fig S1). Reduced

eVLP length and diameter could translate into reduced infectivity (e.g., less PS and less

surface area and membrane available to bind TIM-1), therefore we next sought to

determine the effect of fendiline on eVLP entry.

Fendiline blocks EBOV eVLP entry

A common characteristic of viral infectivity is the relationship between virion

associated PS and the TIM-1 receptor on target cells (Kondratowicz *et al*, 2011; Moller-

Tank *et al*, 2013; Wang *et al*, 2017; Dejarnac *et al*, 2018; Brunton *et al*, 2019). Moreover,

it has been previously reported that other ASM inhibitors blocked EBOV infectivity (Miller

et al, 2012). To determine if fendiline treatment reduced the entry of eVLPs, we performed

a fluorescent based entry assay using 1,1'-dioctadecyl-3,3,3',3'-

tetramethylindocarbocyanine perchlorate (DiI) labelled eVLPs (Nanbo *et al*, 2010; Kuroda

et al, 2015; Nanbo *et al*, 2019). By testing entry of eVLPs derived from fendiline treated

cells rather than the entry of eVLPs on fendiline treated cells, we were able to determine

how fendiline treatment affected eVLP entry rather than how inhibition of ASM in target

cells affected eVLP entry (as previously described (Miller *et al*, 2012). If entry of the eVLPs

was not altered by fendiline treatment, one would expect a dose-dependent increase in

infectivity with increasing TIM-1 expression. Conversely, if eVLP entry was inhibited by

fendiline treatment, a dose-dependent increase in eVLP entry would not be observed with

increasing TIM-1 expression.

TIM-1 overexpression was increased in target cells and a detectable and

significant dose-dependent increase in control eVLP entry was observed, by more than

200% in the highest TIM-1 overexpressing cells (compared to no TIM-1 overexpression;

Fig 7D *top panel* and Fig 7E). Remarkably, no measurable increase in eVLP entry was

observed for fendiline derived-VLPs across any of the TIM-1 overexpressing target cell

conditions (Fig 7D *bottom panel* and 7E). From this comparison, these results suggest

that the impaired entry of fendiline eVLPs is a result of reduced PS in the viral envelope,

either from smaller VLPs or a lower % of PS content. These findings in combination with

the observed reduction in VLP formation further substantiate the significant reduction of

EBOV infection observed in our live virus studies following fendiline treatment.

Mathematical model of in vitro experiments

We next used a mathematical model to predict how the effects of fendiline on both

viral budding and entry combine to produce the observed effects in the BSL-4 assays.

We calibrated our mathematical model (equations 1-3) to experimental data from the

budding, entry and cellular infection assays using approaches and parameter settings

outlined in the methods and Table 1. Results from our two-phase calibration procedure

are shown in Fig 8A-F and Appendix Fig S2. The model captures key features of the data

including a progressive increase in percentage of infected cells over time, differences

between MOI as well as limited cell death in the first 48 hours of the experiment (Appendix

Fig S2).

The dynamics behind these calibrated figures suggested that fendiline treatment

significantly delayed the infection process (Fig 8G,H), resulting in the observed decrease

in percent infected cells with treatment over 4 days. The effects of fendiline on budding

and entry are estimated to have similar pharmacodynamics (PD), with entry effects

estimated to have a slightly stronger response (lower C_{50} and higher E_{max}) compared to

budding (Table 1). Based on PD parameters, the response to fendiline was estimated to

be weaker in the BSL-4 assays, as is evident by higher C_{50} values and lower E_{max} values

compared to the budding and entry assays (Table 1, Fig 8I). These PD parameter

differences between budding and entry assays vs BSL-4 results could suggest that other

parts of the viral life cycle not affected by fendiline (not quantified explicitly in these

experiments) become rate limiting in the BSL-4 assays, thereby reducing the overall effect

of fendiline on infection progression. In summary, a mathematical model consistent with

three independent experimental systems, predicts a combination of budding and entry

effects resulting in the observed BSL-4 effects, and estimates PD parameters for each

mechanism.

Discussion

The host cell PM is exploited by filoviruses for their assembly and budding, where

they can egress the host cell to form a new virion. The matrix protein, VP40, is the main

driver of this process as it harbors a high affinity for lipids in the PM inner leaflet. Lipid

binding by VP40, which includes selectivity for PS (Ruigrok *et al*, 2000; Scianimanico *et*

al, 2000; Adu-Gyamfi *et al*, 2015; Del Vecchio *et al*, 2018) and PI(4,5)P$_2$ (Johnson *et al*,

2016; Johnson *et al*, 2018) drives and stabilizes, respectively, VP40 oligomers that are

necessary for viral budding. In fact, VP40 has been shown to be sufficient (in the absence

of other filovirus proteins) to form VLPs from the host cell PM that are nearly

indistinguishable from virions (Jasenosky *et al*, 2001; Noda *et al*, 2002; Licata *et al*, 2004).

Again, this underscores the unique properties of VP40 structure and sequence, which

provides a template for host lipid binding, oligomerization, and sufficient information to

encode cues for scission to complete the viral budding process. Further, VP40 derived

VLPs enter cells in a PS-dependent manner despite the absence of the EBOV

glycoprotein (Jemielity *et al*, 2013). While some of the basic principles between VP40 and

PM interactions have been previously revealed, the mechanistic consequences of the

interactions and their potential pharmacological targeting have remained unknown.

In this study, we demonstrated that VP40 was able to cluster PS *in vitro* and in the

PM of cells. VP40-dependent PS clustering required PS binding and efficient PM

localization of VP40, as the K224A-VP40 mutant, previously defined as a PS-binding

residue did not significantly increase PS clustering. VP40 oligomerization was also crucial

to PS clustering efficiency as a VP40 oligomerization deficient mutant, which can still

exhibit PM localization, significantly reduced PS clustering. This is an important point as

the PM of host cells is generally thought to harbor 20-30 mol% PS in the inner leaflet. In

fact, VP40 effectively binds and oligomerizes on PS-containing membranes with

compositions close to the PS-content of the PM inner leaflet. In contrast, PS

concentrations below 15 mol% didn't provide robust affinity and oligomerization compared

to those with 22 mol% PS and greater. PS clustering and PS content may also play a

critical role in the loss of asymmetry that occurs during the EBOV budding process. For

instance, pooling of PS in distinct regions of VP40 assembly may provide a cue for

scramblases shown to distribute PS to the outer leaflet of the PM during the viral budding

process (Nanbo *et al*, 2010; Younan *et al*, 2018).

A recent study demonstrated that silencing PSS1, an enzyme responsible for PS

synthesis in mammalian cells was sufficient to inhibit EBOV replication (Younan *et al*,

2018). However, to the best of our knowledge, small molecules aimed at targeting host

cell lipid distribution have not previously been tested against EBOV. Fendiline was a

logical choice to form an initial hypothesis of an FDA-approved drug that could inhibit

EBOV budding as it was recently shown to lower PM PS (Cho *et al*, 2016) and inhibit K-

Ras signaling (van der Hoeven *et al*, 2013; van der Hoeven *et al*, 2018) suggesting this

FDA-approved drug may be sufficient to inhibit EBOV budding. Indeed, in BSL-4

experiments, fendiline was able to inhibit EBOV replication >75% at 20 μM when given

every day post-infection. Follow up mechanistic studies demonstrated that fendiline

efficacy was due to inhibition of VP40 oligomerization, viral budding and viral entry. Thus,

fendiline, which reduced the PS content of HEK293 cells by ~30%, subsequently reduced

PS clustering and VP40 oligomerization necessary for efficient viral budding. VLPs that

did form from fendiline-treated cells had an overall reduced length and surface area,

which likely combined with the reduced PS-content of the virus or VLPs to limit

subsequent viral entry. Thus, disruption of PM PS content by one small molecule was

sufficient to effect at least three important steps in the filovirus life cycle. Importantly, an

increase in intracellular calcium due to fendiline can be ruled out in reducing and/or

inhibiting EBOV and VP40 budding as calcium has been shown to be a critical component

of VLP formation and virus replication (Han Z *et al*, 2015). Thus, an increase of calcium

in HEK293 cells would be expected to increase and not decrease budding as we've

observed.

Overall, this study lends credence to the hypothesis that host processes may be

targeted to inhibit viral replication and spread. While the potency of fendiline may be low,

the combination of fendiline with other FDA-approved drugs that have shown efficacy

against EBOV (Johansen *et al*, 2013; Johansen *et al*, 2015; Nelson *et al*, 2016; Dyall *et*

al, 2018; Jasenosky *et al*, 2019) hold further promise. Fendiline was effective at 5 μM for

reducing PS content in cells and the relationship to fendiline concentrations used *in vivo*

is an important point to address in the future. Fendiline has been used in both single

(200-1200 mg) and multiple doses (400 mg, twice) per day, which was followed with

pharmacokinetic evaluation yielding maximal plasma levels in patients of 9-170 ng/mL

(Weyhenmeyer *et al*, 1987). At the upper range of fendiline detected in the plasma, 170

ng/mL = 540 nM fendiline. Thus, at the upper value of fendiline detected from oral dosing,

the concentration was lower than the 5 μM used to lower cellular PS. Part of the issue in

fendiline dosing is rapid metabolism with a bioavailability of ~20% depending on the mg

dose given (Kurovetz *et al*, 1982). Thus, the issue isn't necessarily fendiline toxicity and

efficacy but drug metabolism of fendiline in the plasma and through the hepatic 1st pass

effect that limit plasma concentration available (Kurovetz *et al*, 1982). Thus, future

studies geared towards more stable drugs that can alter PS content through a similar

mechanism of action (via ASM inhibition) may be beneficial.

Recently, fendiline analogs have been generated that have nanomolar potencies

against K-Ras PS-dependent plasma membrane localization and were effective at

inhibiting cancer cell proliferation in the low micromolar range (Wang *et al*, 2021).

Fendiline has also been shown to block SARS-CoV-2 S-mediated fusion (Xiao *et al*, 2020)

and cause *Leishmania infantum* death via mitochondrial depolarization (Reimão *et al*,

2016). Thus, our work and these studies lay a framework to improved pharmacological

targeting strategies against either VP40 matrix assembly or PS clustering that would not

only reduce viral budding and spread, but lower subsequent viral entry, which partially

relies on PS in the viral envelope (Moller-Tank *et al*, 2013; Jemielity *et al*, 2013; Moller-

Tank *et al*, 2014; Brunton *et al*, 2019). Notwithstanding pharmacological principles

learned from this study, a critical balance between VP40 and PS has been resolved

demonstrating a critical need for VP40 clustering in the assembly and budding process.

VP40 oligomers are needed for enhanced PS clustering where PS clustering seems to

ensure optimal VP40 oligomerization.

Material and Methods

Reagents & solutions

PBS, DMEM, ionomycin and Lipofectamine^{TM} LTX with Plus^{TM} Reagent, and

Invitrogen^{TM} Molecular Probes^{TM} Hoechst 3342 were purchased from Fisher Scientific,

heat-inactivated fetal bovine serum (FBS) was purchased from Hyclone, and Minimum

Essential Medium (MEM) was purchased from Corning. Invitrogen Live Cell Imaging

Solution, DiI Stain, Halt^{TM} Protease Inhibitor Cocktail, Pierce BCA Assay kit, and BS3

were purchased from ThermoFisher Scientific. Non-essential amino acids (NEAA) were

purchased from Sigma Aldrich and L-glutamine was purchased from Gibco. Alexa Fluor^{TM}

488 C_5 Maleimide (Alexa-488) for protein conjugation was purchased from Invitrogen.

Fendiline was purchased from Cayman Chemical Company, prepared in DMSO and

stored at -20 °C. Ultra-Pure Grade DMSO was purchased from VWR and the Ni-NTA

slurry was purchased from Qiagen. L1 chips for SPR experiments were purchased from

GE Healthcare. For cell viability assays, CellTiter-Glo® was purchased from Promega.

Antibody information for immunoblotting and immunofluorescence can be found in Table

2. Ten percent neutral buffered formalin was purchased from Val Tech Diagnostics

(Brackenridge, PA). Cell staining buffer was purchased from BioLegend. stain was

purchased from Fisher Scientific.

Plasmids

EGFP, EGFP-eVP40 and EGFP-WE/A-eVP40 were prepared as described

previously (Adu-Gyamfi *et al*, 2012; Johnson *et al*, 2016). GFP-K224A-eVP40 was

prepared by site directed mutagenesis (Johnson *et al*, 2018) GFP-mVP40 was used as

described previously (Wijesinghe & Stahelin 2015; Amiar *et al*, 2021). The GFP-LactC2

plasmid was a kind gift from Sergio Grinstein (University of Toronto). GFP-PLCδPH was

a kind gift from Tamas Balla (NIH). pCAG-GPI-GFP was a gift from Anna Katerina

Hadjantonakis (Addgene #32601). pEGFP-N3-Annexin A2 was a gift from Volker Gerke

& Ursula Rescher (Addgene #10796). pCAGGS-FLAG-eVP40 (NR49337) and

pcDNA3.1-eGP (NR-19814) were obtained from BEI Services. pCAGGS-TIM-1 was from

Heinz Feldmann (Watt *et al*, 2014).

Lipids and LUV preparation

POPC (#850457), DPPC (#850355), cholesterol (#700000), DPPS (#840037),

POPE (#850757), POPS (#840034), Brain PI(4,5)P$_2$ (#840046), and TopFluor® TMR-PS

(#810242) were purchased from Avanti Polar Lipids, Inc. (Alabaster, AL) and stored in

chloroform and/or methanol at -20°C until use.. For large unilamellar vesicle (LUV)

preparation used in SPR and chemical crosslinking experiments, lipid mixtures were

prepared at the indicated compositions (using a POPC matrix as the background and

control and the indicated additions of PS at the expense of POPC), dried down to lipid

films under a continuous stream of N$_2$, and stored at -20°C until further use. On each day

of experiments, LUVs were brought to room temperature, hydrated in either SPR buffer

(10 mM HEPES, 150 mM NaCl, pH 7.4) or chemical crosslinking buffer (260 µM Raffinose

pentahydrate in PBS, pH 7.4), vortexed vigorously, and extruded through a 100 nm (SPR

experiments) or 200 nm (chemical crosslinking experiments) filter. Vesicle size was

confirmed by dynamic light scattering using a DelsaNano S Particle Analyzer (Beckman

Coulter, Brea, CA).

Preparation and imaging of GUVs

GUVs were prepared by a gentle hydration method (Reeves *et al*, 1969; Darszon

et al, 1980; Yamashita *et al*, 2002). Briefly, 1 mM of lipid control mixture was made and

contained 1,2-dipalmitoyl-sn-glycero-3-phosphocholine (DPPC), cholesterol (Chol) and

fluorescent phosphatidylserine (TopFluor® TMR-PS) at 90:9.8:0.2% molar ratio. For PS

clustering and eVP40 binding analysis, 1,2-dipalmitoyl-sn-glycero-3-phosphoserine

(DPPS) alone or with brain phosphatidylinositol 4,5-bisphosphate $PI(4,5)P_2$ were added

at 40 and 2.5% molar ratios, respectively, and the ratios of DPPC were adjusted

accordingly. The lipid mixtures were prepared in 5 mL round-bottom glass flasks and the

chloroform was removed with rotary movements under a continuous stream of N_2 and

even heat at 55°C until complete evaporation of the solvent. The lipid films were then

hydrated overnight at 50°C (DPPC:Chol:TopFluor TMR-PS) and 55°C

$(DPPC:Chol:DPPS:PI(4.5)P_2:TopFluor$ TMR-PS) in an appropriate volume of GUV

hydration buffer (150 mM NaCl, 10 mM HEPES, 0.5 M sucrose, pH 7.4).

For imaging, the freshly hydrated GUVs were diluted 10 times in GUV dilution

buffer (150 mM NaCl, 10 mM HEPES, 0.5 M glucose, pH 7.4) and placed on a 6-mm

diameter chamber made from a silicon sheet using a core sampling tool (EMS # 69039-

60). The silicon chamber was mounted on a 1.5-mm clean coverglass (EMS # 72200-31)

pre-coated with 1 mg/mL BSA. The set up was then assembled in an Attofluor chamber

(Invitrogen # A7816) and 1.25 µM eVP40-Alexa488 was added. GUVs imaged for eVP40

and PS colocalization and clustering analysis was performed at 37°C on a Nikon Eclipse

Ti Confocal inverted microscope (Nikon Instruments, Japan), using a Plan Apochromat

60x 1.4 numerical aperture oil objective and a 100x 1.45 numerical aperture oil objective,

respectively. A 488 nm argon laser was used to excite GFP and a 561 nm argon laser

was used to excite TopFluor®-TMR-PS. The 3D reconstruction was performed using

ImageJ. Mander's correlational analysis was performed using the plugin JACoP (Bolte &

Cordelières 2006).

Cell culture, transfections, pharmacological treatments

All BSL-2 studies were performed using HEK293 cells obtained from the American

Type Culture Collection and cultured in DMEM supplemented with 10% FBS and 1% PS.

Transient transfections were performed using Lipofectamine^{TM} LTX + PLUS^{TM} Reagent,

according to the manufacturer's protocol. All transfections were performed in DMEM

supplemented with 10% FBS. Treatment with fendiline (in DMSO) occurred at 5-hours

post-transfection in DMEM supplemented with 10% FBS. BSL-4 assays were also

performed using Vero E6 cells cultured in MEM, 5% heat-inactivated fetal bovine serum,

1% L-glutamine, and 1% NEAA. Both HEK293 and Vero E6 cells were cultured and

incubated at 37°C, 5% CO_2, 80% humidity.

Immunoblotting

Samples prepared for western blotting analysis were first separated using SDS-

PAGE (8% for chemical crosslinking and 12% for cell lysates and VLPs). Following

transfer onto a nitrocellulose membrane, membranes were blocked with 5% MILK-TBST

and analyzed with their respective antibodies (See Table 2). Antibodies were detected

using an ECL detection reagent and imaged on the ImageQuant LAS 4000 or Amersham

Imager 600 (GE Healthcare Life Sciences). All quantitative analysis derived from western

blotting was performed using densitometry analysis in ImageJ.

Protein purification and Alexa-488 C$_5$ Maleimide labelling

The His$_6$-eVP40-pET46 expression vector was a kind gift from Erica Ollmann

Saphire (La Jolla Institute for Immunology) and was expressed and grown in Rosetta^{TM} 2

BL21 DE3 competent cells (Merck Millipore, Billerica MA). The pet28a-His$_6$-Lact C2

bacterial expression plasmid was a kind gift from Dr. Sergio Grinstein. His$_6$-eVP40 and

His$_6$-LactC2 were grown and purified as described previously (Johnson *et al*, 2016).

Following elution from a Ni-NTA slurry (Qiagen), the protein samples were then further

purified using size exclusion chromatography on a HiLoad 16/600 Superdex 200 pg

column (ÄKTA pure, GE Healthcare). The desired fractions containing dimeric VP40 or

monomeric LactC2 were collected, concentrated and stored in 10 mM Tris, 300 mM NaCl,

pH 8.0. Protein concentration was calculated using the Pierce BCA assay and the protein

was stored at 4°C for no longer than 14 days.

Labeling of eVP40 cysteine residues was carried out using Alexa-488- C$_5$-

maleimide. eVP40 dimer was treated with a 1.5-fold molar excess of Alexa-488-C5-

maleimide dissolved in DMSO for 2 hours at room temperature in maleimide labelling

buffer (20 mM NaPi solution pH 7.4, 150 mM NaCl, 4 M Guanidine HCl). The labeling

reaction was quenched by adding diothiothreitol (final concentration of 50 mM) and the

labeled protein was separated from the non-conjugated dye on a HiLoad 16/600

Superdex 200 pg column. Fractions containing eVP40 dimer were collected and

concentrated. The labelling efficiency and protein concentration were estimated using a

NanoDrop according to Invitrogen's instructions.

Plasma membrane localization confocal microscopy

Live cell imaging experiments were performed at 24 hours and 48 hours post

treatment. Experiments to quantify fluorescent protein PM localization were performed on

a Zeiss LSM 710 inverted microscope using a Plan Apochromat 63x 1.4 numerical

aperture oil objective. A 488 nm argon laser was used to excite GFP/EGFP. GFP-LactC2

PM localization was quantified ratiometrically by comparing the PM signal vs. the cytosolic

signal. GFP-EBOV-VP40 PM localization was quantified ratiometrically by comparing the

PM signal vs. the total fluorescence signal within the cell.

TMR-PS quenching experiment

The Top Fluor TMR quenching analysis was performed as described previously

(Zhao *et al*, 2010). Briefly, LUVs with 0.5% TopFluor® TMR-PS and increasing amounts

of DPPS (0, 1, 5, 10, 20, 40, 60%) with or without $PI(4.5)P_2$ (2.5% with 60% DPPS) were

made as described above. LUVs were then mixed at a final concentration of 40 μM with

corresponding concentrations of eVP40 dimer in 10 mM HEPES, 160 mM NaCl pH 7.4 in

a black/clear bottom 96-well plate and incubated at 37$°$C for 30 min. The Top Fluor TMR

was excited at 547 nm and the fluorescence was recorded from 555 to 600 nm with no

cutoff wavelength using the plate reader SpectraMax M5e (Molecular Devices, St Jose,

CA). The fluorescence quenching ratio, ΔF, was calculated according to the ratio (F_0/F),

where F_0 is the fluorescence in absence of protein and F is the fluorescence at a given

eVP40 concentration.

Cellular Top Fluor TMR-PS clustering confocal microscopy

Each experimental day, a 100 µM working stock of TopFluor® TMR-PS in

methanol was prepared. Immediately prior to imaging, cells were placed in 4°C for 5 min.

The working stock was diluted to a final 500 nM TopFluor® TMR-PS solution in 3 mg/mL

BSA/PBS. The 500 nM TopFluor® TMR-PS/BSA/PBS solution was incubated with cells

at 4°C for 10 min, rinsed three times with cold PBS, and immediately imaged in fresh cold

PBS. Top Fluor TMR was excited at 560 nm and GFP was excited at 488 nm. For PS

clustering analysis, a custom macro in ImageJ was used. Prior to the macro analysis,

background was subtracted and the contrast was enhanced. To isolate the PM area, a

default threshold was applied. To isolate PS clusters the Moments analysis thresholding

(Tsai 1985) was applied. Following the moments analysis thresholding, the custom

ImageJ macro was applied: despeckle, close-, fill holes, and remove outliers (radius=5,

threshold=50). The sum of the remaining particles area was calculated, as well as the PM

area. %PS clustering was calculated according to the ratio ($Area_{clusters}/Area_{plasma\ membrane}$).

Number & Brightness

Number & Brightness experiments were performed as described previously (Adu-

Gyamfi *et al*, 2017; Bobone *et al*, 2017; Johnson *et al*, 2018) on a Zeiss LSM 880 upright

microscope using a LD "C-Apochromat" 40x/1.1 W Corr M27 objective. HEK293 cells

expressing either GFP, GFP-LactC2 or GFP-eVP40 were treated with fendiline (1 or 5

µM) for 48-hours prior to N&B analysis. Cells were imaged in phenol-free live cell imaging

solution. For each experimental day, the brightness value of a monomer was determined

in cells expressing monomeric GFP. Each image was acquired using the same laser

power (0.01), resolution (256x256), pixel dwell time (16 us), frames (50), and zoom (pixel

size of 50 nm). SimFCS Globals Software (Laboratory for Fluorescence Dynamics,

University of California, Irvine, CA) was used for analysis.

Chemical crosslinking

His$_6$-eVP40 and His$_6$-LactC2 were purified as previously described in *Protein*

Purification. LUVs containing POPC and Brain PI(4,5)P$_2$ (2.5%) with varying PS mol%

composition (0, 15,30, 60%) were prepared as previously described in *Lipids & Vesicle*

Preparation. Experimental protocol was adapted from (Johnson *et al*, 2016), and the

manufacturers protocol for BS^3 (ThermoFisher) with each step performed at RT. In brief,

protein (final concentration of 0.3 μM in PBS pH 7.4) was mixed with LUVs (final

concentration = 660 μM) at a 1:1 volumetric ratio for 30 min. Protein bound LUVs were

separated from unbound protein through centrifugation (75,000 x *g*, 30 min, 22°C), and

resuspended in PBS (pH 7.4) buffer containing BS^3 (final concentration of 200 μM).

Samples were incubated for 45 min, quenched with glycine for 15 min and then analyzed

through western blotting. Following immunoblotting, oligomerization of VP40 was

quantified ratiometrically by comparing VP40$_o$ vs. VP40$_{m+d}$ (where VP40$_o$ is the oligomeric

VP40 band density (>75 kDa) and VP40$_{m+d}$ is the sum of monomeric and dimeric eVP40

(~37 and 74 kDa) band density).

Surface plasmon resonance

To determine the affinity of His$_6$-eVP40 to LUVs with increasing PS concentrations,

SPR was performed. SPR experiments were performed at 25°C using a Biacore X100 as

described previously (Adu-Gyamfi *et al*, 2015). In brief, an L1 chip was coated at 5 μL/min

with LUVs containing 0% PS (100% POPC, PS added at the expense of POPC) on flow

cell 1 and either 1%, 11% or 22 mol% POPS on flow cell 2 (LUV preparation described in

previous section, *Lipids and Vesicle Preparation)*. The LUV conjugated chip was

stabilized by washing with 50 mM NaOH and blocked with 0.1 mg/mL BSA (in SPR buffer)

at a flow rate of 10 µL/min until the response on each flow cell was <100 response units

(RU). For quantitative affinity analysis, each concentration of eVP40 was injected for 540

s at a flow rate of 10 µL/min with a 180 s delay, and the difference in response between

flow cell 1 and flow cell 2 was recorded (ΔRU). The apparent K_d of vesicle binding was

determined   using   the   non-linear   least   squares   analysis: $Req = Rmax/(1 +$

$\frac{Kd}{C}$)where R_{eq} (measured in RU) is plotted against protein concentration (C). R_{max} is the

theoretical maximum RU response and K_d is the apparent membrane affinity. Data were

fit using the Kaleidagraph fit parameter of (m0*m1)/(m0+m2);m1=1100;m2=1. ΔRU data

was normalized in GraphPad Prism 8 for windows (La Jolla, CA) and plotted in

Kaleidagraph (Reading, PA).

Lipidomics

HEK293 cells were treated with the indicated concentration of fendiline for 48

hours, collected through centrifugation, rinsed with PBS and protein concentration was

determined. Cells were pelleted, flash frozen in liquid N_2 and stored at -80°C until

subsequent LC/MS/MS processing by Avanti Polar Lipids, Inc.. Prior to LC-MS/MS

analysis, lipids were extracted using the Folch method (Folch *et al*, 1957). The bottom

chloroform layer taken after centrifugation was diluted with internal standards for LPA,

LPS, PA and PS for quantization by injection on LC-MS/MS. Samples were injected on a

LC-MS/MS method using a Waters Acquity UPLC / AB Sciex 5500 MS system performing

reversed phase separation of LPA and LPS and PA and PS components with MS/MS

detection. Each molecular species identified by the [M-H] m/z of its acyl carbon:double

bond (CC:DB i.e. 34:2 PA) was quantified against the response of the internal standards

of known concentration. Content of individual and total LPA/PA and LPS/PS was

reported. Values were corrected to 1×10^6 cells for all samples.

BSL-4 immunofluorescence assay

Vero E6 cells were seeded at 2e4 cells/well in black 96-well poly-D-lysine treated

plates (Greiner Bio-One Cellcoat®). Twenty-four hours prior to infection, fendiline was

diluted in 0.5% DMSO and Vero E6 cell culture media at indicated concentrations and

added to cells. An equivalent percentage of DMSO in culture media served as the vehicle

control. Following pretreatment, compound was removed, and cells were incubated with

Ebola virus (Kikwit) at a multiplicity of infection (MOI) of 0.1 or 1.0 in a BSL-4 located at

USAMRIID. Following absorption for 1 hour, virus inoculum was removed and cells were

washed. Plates were divided into three post-infection treatment groups (day 0, every day-

e.d., every other day-e.o.d.), and received either culture media or freshly prepared

fendiline or vehicle control. Cells were then treated daily with freshly prepared compound

or left to incubate based on their designated treatment group. At 48 hours (MOI=1.0), 72

or 96 hours (MOI=0.1) post infection, cells were washed with PBS and submerged in 10%

neutral buffered formalin for 24 hours prior to removal from the BSL-4 laboratory. Formalin

was removed and cells were washed with PBS. Cells were blocked with 3% BSA/PBS

cell staining buffer (BioLegend) and incubated at 37°C for 2 hours. Ebola virus GP-

specific mAb KZ52, diluted in 3% BSA/PBS, was added to appropriate wells containing

infected cells and incubated at room temperature for 2 hours. Cells were washed three

times with PBS prior to addition of goat anti-human or goat anti-mouse IgG-Alexa-488

secondary antibody. Following 1-hour incubation with secondary antibody, cells were

washed 3 times prior to counterstaining with Hoechst's stain diluted in PBS. Cells were

imaged and percent of virus infected cells calculated using the Operetta High Content

Imaging System and Harmony® High Content Imaging and Analysis Software

(PerkinElmer).

VLP collections & functional budding assays

HEK293 cells were transfected and treated with fendiline as described in the

previous section, *Cell Culture, Transfection & Pharmacological Treatments*. Budding

assays were performed as described previously (Johnson *et al*, 2016; Harty 2018). In

brief, VLP containing supernatants were harvested from cells and clarified through low

speed centrifugation. Clarified VLPs were loaded onto a 20% sucrose cushion in STE

buffer (10 mM TRIS, 100 mM NaCl, 1 mM EDTA, pH 7.6), isolated through

ultracentrifugation, and resuspended in either 150 mM ammonium bicarbonate (functional

budding assays), 2.5% glutaraldehyde in 0.1 M cacodylate buffer (TEM experiments),

STE buffer (entry assays) or 0.1 M phosphate buffer for CD and thermal melting (PB; 0.02

M sodium phosphate monobasic, 0.08 M sodium phosphate dibasic, pH 7.4). VLP

samples were stored at -80°C for functional budding assays, -20°C for entry assays or

4°C for TEM and CD analysis.

For functional budding assays, cell lysate samples were harvested and lysed on

ice with RIPA buffer (150mM NaCl, 5mM EDTA pH=8, 50mM Tris pH 7.4, 1% Triton-X,

0.1% SDS, 0.5% deoxycholic acid) supplemented with Halt^{TM} Protease Inhibitor Cocktail.

Prior to separation on a 12% SDS-PAGE gel, cell lysate and VLP sample volume loading

were normalized to sample protein content, determined by a BCA assay. Gels were

transferred to a nitrocellulose membrane and immunoblotted was performed as described

previously in the section, *Immunoblotting*. Following ECL detection, VP40 cell lysate

(VP40$_{CL}$) expression was normalized to the respective GAPDH band density. The relative

budding index was calculated according to the ratio of density$_{VLP}$/density$_{C+VLP}$ (where

density$_{VLP}$ is the eVP40 VLP band density and density$_{C+VLP}$ is the eVP40 cell lysate + VLP

band density).

Scanning electron microscopy

HEK293 cells were transfected with FLAG-eVP40 and treated as described in the

previous section, *Cell Culture, Transfections & Pharmacological Treatments.* Cells were

scraped and collected through low-speed centrifugation at 48 hours post-transfection, and

stored in primary fixative (2% glutaraldehyde, 2% paraformaldehyde in 0.1 M cacodylate

buffer, pH 7.35) at 4°C until processing. During processing, samples were fixed to

coverslips and post-stained with 1% osmium tetroxide in 0.1 M cacodylate buffer.

Samples were extensively rinsed with water and dehydrated with a graded series of

ethanol followed by drying in a Tousimis 931 Supercritical Autosamdri® device. Prior to

imaging, samples were coated with 3 nm Iridium. A Field Emission Scanning Electron

Microscope Magellan 400 (FEI) (Hillsboro, OR) was used to collect images, with

assistance from Tatyana Orlova at the Notre Dame Integrated Imaging Facility.

Transmission electron microscopy imaging

VLPs were purified as previously described in *VLP Collections & Functional*

Budding Assays. Following ultracentrifugation, VLPs were resuspended in fixative (2.5%

glutaraldehyde in 0.1 M cacodylate buffer). Purified VLPs were applied onto glow

discharged carbon formvar grids and negatively stained using 4% uranyl acetate.

Samples were imaged with a FEI Tecnai T12 electron microscope equipped with a

tungsten source and operating at 80 kV. VLP length and diameter measurements were

quantified using ImageJ software. For diameter analysis, eight different diameters were

measured across random areas on each VLP, and the mean diameter was reported.

Circular dichroism

VLPs were produced in the presence of DMSO or fendiline and purified as

previously described in the *VLP collections* section. PB buffer was added to a 10-mm

path-length Spectrosil Far UV Quartz cuvette (Starna Cells CatID: 21-Q-10) and a

background spectra was collected and autosubtracted from VLP samples. VLP samples

were loaded into the 10-mm cuvette diluted in PB buffer to an approximate final protein

concentration of 30 µg/mL. Circular dichroism spectra of VLP samples were collected

between 200-280 nm at a 0.2 nm step size with 0.5 s time-per-point (with adaptive

sampling) using a Chirascan spectrometer (Applied Photophysics, Leatherhead, UK).

Absorbance spectra (Abs) and detector signal (hv) were collected simultaneously as

controls. After collecting the CD spectra, a microstir bar was added to the cuvette and

thermal melting was run from 20°C-93°C using a 0.5°C step ramping at 1.00°C/minute

with a tolerance of 0.20°C; simultaneously, spectra were collected at a single wavelength

of 220 nm with a time-per-point of 24 seconds. Absorbance spectra (Abs) and detector

signal (hv) were collected simultaneously as controls. At the end of thermal melting

measurement collection, temperature set-points were replaced with temperatures

measured by the sample handling unit. Data was converted from Chirascan filetype to

CSV and then extracted into GraphPad PRISM 7. Using the first derivative of the circular

dichroism spectral signal in respect to temperature, the maxima was taken as the melting

point of the sample. Melting temperatures of the three replicates were averaged for the

reported T_m.

DiI entry assay

VLPs produced from HEK293 cells expressing FLAG-eVP40 and eGP were

purified as previously described in the *Functional Budding Assays* section. DiI entry

assays were performed as described previously (Kuroda *et al*, 2015; Nanbo *et al*, 2018).

In brief, following ultra-centrifugation VLPs were resuspended in STE buffer and further

purified by filtering through a 0.22 μm filter. Protein content of VLP samples were

normalized to 0.1 μg/mL using STE buffer. VLPs were labeled with DiI for 1 hr at RT with

gentle agitation (final DiI = 0.06 μM). Following incubation, labeled VLP samples were

concentrated down to equal volumes, and brought up to volume in phenol-free MEM with

2% FBS and 4% BSA.

HEK293 cells were transfected with TIM-1 for 24 hours prior to incubation with DiI-

labeled VLPs and briefly rinsed with phenol-free MEM with 2% FBS and 4%BSA. DiI-

VLPs were added to TIM-1 expressing HEK293 cells, spinoculated for 45 min at 4°C, and

allowed to incubate for 1 hr at 37°C. Plates were then rinsed with PBS, fixed with 4%

paraformaldehyde in PBS, their nuclei stained with Hoechst 3342, and stored at 4°C until

imaging. During image acquisition, z-stacks were acquired of 10-15 frames (1 μm steps).

Toxicity analysis

HEK293 and Vero E6 cell toxicity following fendiline treatment was tested at the

indicated time points using the Cell Titer Glo Viability Assay (Promega, Madison WI)

according to the manufacturer's protocol. In brief, HEK293 cells were treated with the

indicated concentration of fendiline or control for 24 or 48 hours. Vero E6 cells were

treated for 24 hours, the drug was removed and the cells were replenished with Vero E6

culture media, to mirror the corresponding ebolavirus infections at BSL-4. Following the

one-hour mock infection, cells were washed with PBS and plates were divided into three

treatment groups (day 0, e.d., e.o.d.), and cells received either culture media or freshly

prepared fendiline or vehicle control and were then treated daily with freshly prepared

compound or left to incubate based on their designated treatment group. At 48, 72, and

96 hours following mock infection, and mirroring the post infection fixation time points,

CellTiter-Glo® reagent was added to each well in accordance with the manufacturer's

instructions. Both HEK293 and Vero E6 toxicity assays luminescence readings were

recorded using a SpectraMax® M5 (Molecular Devices®) plate reader.

Mathematical model of *in vitro* experiments

We implemented a system of ordinary differential equations (ODEs) to describe

the dynamics of host target cells, infected cells and free virus in different combinations

reflecting the *in vitro* experimental systems used here. These equations are similar to

those used to simulate Ebola virus dynamics in earlier work (Nguyen *et al*, 2015;

Martyushev *et al*, 2016).

$\frac{dT}{dt} = -\beta TV$ (1)

$\frac{dI}{dt} = \beta TV - \delta I$ (2)

$\frac{dV}{dt} = pI - cV$ (3)

Where *T*, *I* and *V* represent numbers of susceptible target host cells, infected cells and

free virions respectively and parameters are described in Table 2.

We modified appropriate parameters in equations 1-3 to represent the following

experimental systems during calibration of the mathematical model:

• Viral budding assay (set β=0, $T(0) = 0$, $I(0) = 2.625 \times 10^6$, $V(0) = 0$, vary δ, c, p)

• Viral entry assay (set p=0, $T(0) = 6.3 \times 10^5$, $I(0) = 0$, $V(0) = 6.3 \times 10^3$, vary β, δ, c)

• BSL-4 Cellular infection assay (BSL4) ($T(0) = 5 \times 10^4$, $I(0) = 0$, $V(0) = 5 \times 10^4$ (MOI 1)

or $V(0) = 5 \times 10^3$ (MOI 0.1) , vary β, δ, c, p)

Fendiline treatment effects are simulated using E_{max} dose response curves:

977                    $$f_X = E_{max} \frac{C^H}{C_{50}^H + C^H}$$

Where *C*: concentration of fendiline, *E*$_{max}$: maximum effect of fendiline, *H*: hill constant for

the dose response curve, and *C*$_{50}$ concentration with 50% of *E*$_{max}$ efficacy. *E*$_{max}$, *C*$_{50}$ and

H are fitted separately for fendiline effects on budding (*f*$_{budding}$) or entry (*f*$_{entry}$). Fendiline

efficacy (f_X) is defined as a fraction where f_X=0 implies no effect and f_X=1 implies 100%

inhibition of X (X = budding or entry). Fendiline effects are integrated into equations (1-3)

by multiplying β by (1-f_{entry}) and multiplying p by (1-$f_{budding}$). Fendiline concentrations are

assumed to be constant over the observation periods based on low *in vitro* degradation

rates of the drug. Daily treatment in the BSL-4 assays (e.d.) are simulated by removing

all free virus particles from the equations at each dosing time.

We calibrated the model in two stages. First, we calibrated to the budding and

entry assays. The uncoupling of budding and entry in this data allows us to define

biologically feasible ranges for the effects of fendiline on budding and entry separately.

Using these feasible ranges, we proceed to calibrate the full model to the BSL-4 data

(day-1/0 and e.d.). Therefore, we allow the budding and entry assays to inform the BSL-

4 simulations without imposing strict assumptions about the equivalency between the two

systems. In this way we progressively build complexity into the model accounting for

fendiline effects on viral budding, entry, and infection progression.

Calibrating to budding and entry assays we restrict the value of p to be larger than

1 and c to be between 0 and 5. These assumptions are in line with previous estimates,

and are necessary to qualitatively reproduce viral production observed, but not quantified,

in the budding assays. Parameters are estimated using Matlab's non-linear least squares

optimization algorithm. Parameter bounds and final values are defined in Table 2.

1001

Statistical testing

All experiments were done in triplicate (unless otherwise noted and figure legends

indicate the number of replicates). For analysis of eVLP diameter and length from TEM

experiments, as well as total PA levels between control and 5 µM fendiline treated cells,

a two-tailed t-test was performed. For all experiments which contained >2 experimental

groups, a one-way ANOVA with Dunnett's multiple comparisons was performed on raw

data. Lastly, for N&B analysis, a two-way ANOVA with Dunnett's multiple comparisons

was performed.

1010
1011 **Data Availability**

1012   No data were deposited in a public database and any additional data from this

1013 paper can be requested from the authors.

1014
Acknowledgements

These studies were supported by the NIH (AI081077) to R.V.S and the Indiana CTSI to

E.P. and R.V.S. (UL1TR001108). M.L.H. was partially supported by a NIH T32 fellowship

(T32 GM075762), E.A.D. was supported by NIH T32 fellowship (GM132024), and C.B.P.

was supported by NIH T32 fellowship (AI148103). We are grateful for lipidomic analysis

by Avanti Polar Lipids, Inc., and for research support by Dr. Nathan Dissinger. The

authors acknowledge the use of the facilities of the Bindley Bioscience Center, a core

facility of the NIH-funded Indiana Clinical and Translational Sciences institute and the use

of the Purdue Life Science Electron Microscopy facility.

1024
Author Contributions

Conceptualization: M.L.H. and R.V.S. Investigation: M.H., S.A., L.I.P., E.A.D., C.B.P., and

K.E.H. Methodology: M.H., S.A., L.I.P., E.A.D., C.B.P., K.E.H., and L.I.P. designed the

BSL-4 experiments. Formal analysis: M.H., S.A., L.I.P., E.A.D., C.B.P., L.I.P. and K.E.H.

performed and analyzed the BSL-4 experiments. E.P. designed and performed the

mathematical modeling and analysis. Funding acquisition: M.L.H., E.A.D., C.B.P., E.P.,

and R.V.S. Project administration: R.V.S. Supervision: R.V.S. oversaw the in vitro and

BSL-2 work, J.M.B. and J.D. oversaw the BSL-4 work, E.P. oversaw the mathematical

modeling and analysis. Writing - original draft: M.L.H. and R.V.S. Writing – review and

editing: all authors.

1035

1036 **Conflict of Interest**

1037 The authors declare that they have no conflict of interest.

References

Adu-Gyamfi E, Digman MA, Gratton E, Stahelin R V (2012) Investigation of Ebola VP40 assembly and oligomerization in live cells using number and brightness analysis. *Biophys J* 102: 2517–2525

Adu-Gyamfi E, Soni SP, Xue Y, Digman MA, Gratton E, Stahelin RV (2013) The Ebola virus matrix protein penetrates into the plasma membrane: a key step in viral protein 40 (VP40) oligomerization and viral egress. *J Biol Chem* 288: 5779-5789

Adu-Gyamfi E, Johnson KA, Fraser ME, Scott JL, Soni SP, Jones KR, Digman MA, Gratton E, Tessier CR, Stahelin RV (2015) Host Cell Plasma Membrane Phosphatidylserine Regulates the Assembly and Budding of Ebola Virus. *J Virol* 89: 9440-9453

Amara A, Mercer J (2015) Viral apoptotic mimicry. *Nat Rev Microbiol* 13: 461–469

Amiar S, Stahelin RV (2020) The Ebola virus matrix protein VP40 hijacks the host plasma membrane to form virus envelope. *J Lipid Res* 61: 971

Amiar S, Husby ML, Wijesinghe KJ, Angel S, Bhattarai N, Gerstman BS, Chapagain PP, Li S, Stahelin RV (2021) Lipid-specific oligomerization of the Marburg virus protein VP40 is regulated by two distinct interfaces for virion assembly. *J Biol Chem* 296: 100796

Banadyga L, Dolan MA, Ebihara H (2016) Rodent-Adapted Filoviruses and the Molecular Basis of Pathogenesis. *J Mol Biol* 428: 3449–3466

Bayer R, Mannhold R. (1987) Fendiline: a review of its basic pharmacological and clinical properties. *Pharmatherapeutica* 5: 103–136

Bell RM, Ballas LM, Coleman RA (1981) Lipid topogenesis. *J Lipid Res* 22: 391–403

Berjukow S, Doring F, Froschmayr M, Grabner M, Glossman H, Hering S (1996) Endogenous calcium channels in human embryonic kidney (HEK293) cells. *Br J Pharmacol* 118: 748-753

Bobone S, Hilsch M, Storm J, Dunsing V, Herrmann A, Chiantia S (2017) Phosphatidylserine Lateral Organization Influences the Interaction of Influenza Virus Matrix Protein 1 with Lipid Membranes. *J Virol* 91: 1–15.

Bolte S, Cordelières FP (2006) A guided tour into subcellular colocalization analysis in light microscopy. *J Microsc* 224: 213–232

Bornholdt ZA, Noda T, Abelson DM, Halfmann P, Wood MR, Kawaoka Y, Sahpire EO (2013) Structural rearrangement of ebola virus vp40 begets multiple functions in the virus life cycle. *Cell* 154: 763–77

Breman JG, Heymann DL, Lloyd G, McCormick JB, Miatudila M, Murphy FA, Muyembé-Tamfun JJ, Piot P, Ruppol JF, Sureau P, *et al* (2016) Discovery and Description of Ebola Zaire Virus in 1976 and Relevance to the West African Epidemic during 2013-2016. *J Infect Dis* 214: S93–S101

Brunton B, Rogers K, Phillips EK, Brouillette RB, Bouls R, Butler NS, Maury W (2019) TIM-1 serves as a receptor for Ebola virus in vivo, enhancing viremia and pathogenesis. *PLoS Negl Trop Dis* 13: e0006983

Carnec X, Meertens L, Dejarnac O, Perera-Lecoin M, Hafirassou ML, Kitaura J, Ramdasi R, Schwartz O, Amara A (2015) The Phosphatidylserine and Phosphatidylethanolamine Receptor CD300a Binds Dengue Virus and Enhances Infection. *J Virol* 90: 92–102

Cheng JS, Wang JL, Lo YK, Chou KJ, Lee KC, Liu CP, Chang HT, Jan CR (2001) Effects

of the antianginal drug fendiline on Ca2+ movement in hepatoma cells. *Hum Exp*
Toxicol 20: 359-364

Cho KJ, van der Hoeven D, Zhou Y, Maekawa M, Ma X, Chen W, Fairan GD, Hancock
JF (2015) Inhibition of Acid Sphingomyelinase Depletes Cellular
Phosphatidylserine and Mislocalizes K-Ras from the Plasma Membrane. *Mol Cell*
Biol 36: 363–374

Cho W, Stahelin RV (2005) Membrane-protein interactions in cell signaling and
membrane trafficking. *Annu Rev Biophys Biomol Struct* 34: 119-151

Darszon A, Vandenberg CA, Schönfeld M, Ellisman MH, Spitzer NC, Montal M (1980)
Reassembly of protein-lipid complexes into large bilayer vesicles: perspectives for
membrane reconstitution. *Proc Natl Acad Sci U S A* 77: 239–243

Dejarnac O, Hafirassou ML, Chazal M, Versapuech M, Gaillard J, Perera-Lecoin M,
Umana-Diaz C, Bonnet-Madin L, Carnec X, Tinevez JY *et al* (2018) TIM-1
Ubiquitination Mediates Dengue Virus Entry. *Cell Rep* 23: 1779–1793

Del Vecchio K, Frick CT, Gc JB, Oda S, Gerstman BS, Saphire EO, Chapagain PP,
Stahelin RV (2018) A cationic, C-terminal patch and structural rearrangements in
Ebola virus matrix VP40 protein control its interactions with phosphatidylserine. *J*
Biol Chem 293: 3335–3349

Digman MA, Dalal R, Horwitz AF, Gratton E (2008) Mapping the number of molecules
and brightness in the laser scanning microscope. *Biophys J* 94: 2320–2332

Dyall J, Nelson EA, DeWald LE, Guha R, Hart BJ, Zhou H, Postnikova E, Logue J, Vargas
WM, Gross R, *et al* (2018) Identification of Combinations of Approved Drugs With
Synergistic Activity Against Ebola Virus in Cell Cultures. *J Infect Dis* 218: S672–
S678

Fairn GD, Schieber NL, Ariotti N, Murphy S, Kuerschner L, Webb RI, Grinstein S, Parton
RG (2011) High-resolution mapping reveals topologically distinct cellular pools of
phosphatidylserine. *J Cell Biol* 194: 257–227

Folch J, Lees M, Sloane Stanley GH (1957) A simple method for the isolation and
purification of total lipides from animal tissues. *J Biol Chem* 226: 497–509

Graul, AI, Pina, P, Tracy, M, Sorbera, L (2020) This year's new drugs and biologics. *Drugs*
Today 56: 47-103

Han Z, Madara JJ, Herbert A, Prugar LI, Ruthel G, Lu J, Liu Y, Liu W, Liu X, Wrobel JE
et al (2015) Calcium regulation of hemorrhagic fever virus budding: Mechanistic
implications for host-oriented therapeutic intervention. *PLoS Pathog* 11: e1005220

Harty RN (2018) Hemorrhagic Fever Virus Budding Studies. In: *Hemorrhagic Fever*
Viruses: Methods and Protocols, Salvato MS (ed) pp. 209–215. Springer New York,
New York, USA.

Hirama T, Das R, Yang Y, Ferguson C, Won A, Yip CM, Kay JG, Grinstein S, Parton RG,
Fairn GD (2017) Phosphatidylserine dictates the assembly and dynamics of
caveolae in the plasma membrane. *J Biol Chem* 292: 14292–14307

Hoenen T, Biedenkopf N, Zielecki F, Jung S, Groseth A, Feldmann H, Becker S (2010)
Oligomerization of Ebola virus VP40 is essential for particle morphogenesis and
regulation of viral transcription. *J Virol* 84: 7053–7063

Huang CC, Huang CJ, Cheng JS, Liu SI, Chen IS, Tsai JY, Chou CT, Tseng PL, Jan CR
(2009) Fendiline-evoked [Ca2+]i rises and non-Ca2+-triggered cell death in human
oral cancer cells. *Hum Exp Toxicol* 28: 41-48

Jan CR, Tseng CJ, Chen WC (2000) Fendiline increases [Ca2+]I in Madin Darby canine
kidney (MDCK) cells by releasing internal Ca2+ followed by capacitative Ca2+
entry. *Life Sci* 66: 1053-1062
Jan CR, Lee KC, Chou KJ, Cheng JS, Wang JL, Lo YK, Chang HT, Tang KY, Yu CC,
Huang JK (2001) Fendiline, an anti-anginal drug, increases intracellular Ca2+ in
PC3 human prostate cancer cells. *Cancer Chemother Pharmacol* 48: 37-41
Jasenosky LD, Cadena C, Mire CE, Borisevich V, Haridas V, Ranjbar S, Nambu A, Bavari
S, Soloveva V, Sadukhan S *et al* (2019) The FDA-Approved Oral Drug
Nitazoxanide Amplifies Host Antiviral Responses and Inhibits Ebola Virus. *iScience*
19: 1279–1290
Jasenosky LD, Neumann G, Lukashevich I, Kawaoka Y (2001) Ebola virus VP40-induced
particle formation and association with the lipid bilayer. *J Virol* 75: 5205–5214
Jasenosky LD, Kawaoka Y (2004) Filovirus budding. Virus Res. 106: 181–188.
Jemielity S, Wang JJ, Chan YK, Ahmed AA, Li W, Monahan S, Bu X, Farzan M, Freeman
GJ, Umetsu DT *et al* (2013) TIM-family Proteins Promote Infection of Multiple
Enveloped Viruses through Virion-associated Phosphatidylserine. *PLoS Pathog* 9:
e1003232.
Johansen LM, Brannan JM, Delos SE, Shoemaker CJ, Stossel A, Lear C, Hoffstrom BG,
Evans Dewald L Schornberg KL, Scully C *et al* (2013) FDA-approved selective
estrogen receptor modulators inhibit Ebola virus infection. Sci Transl Med 5:
190ra79
Johansen LM, DeWald LE, Shoemaker CJ, Hoffstrom BG, Lear-Rooney CM, Stossel A,
Nelson E, Delos SE, Simmons JA, Greiner JM *et al* (2015) A screen of approved
drugs and molecular probes identifies therapeutics with anti-Ebola virus activity. *Sci*
Transl Med 7: 290ra89
Johnson KA, Budicini MR, Urata S, Gerstman BS, Chapagain PP, Li S, Stahelin RV
(2018) PI(4,5)P2 Binding Sites in the Ebola Virus Matrix Protein Modulate Assembly
and Budding. *bioRxiv* doi:10.1101/341248
Johnson, KA, Bhattarai N, Budicini MR, LaBonia CM, Baker SCB, Gerstman BS,
Chapagain PP, Stahelin RV (2021) Cysteine mutations in the Ebolavirus matrix
protein VP40 promote Phosphatidylserine bniding by increasing the flexibility of a
lipid-binding loop. *Viruses* 13: 1375
Johnson KA, Taghon GJF, Scott JL, Stahelin R V (2016) The Ebola Virus matrix protein,
VP40, requires phosphatidylinositol 4,5-bisphosphate (PI(4,5)P2) for extensive
oligomerization at the plasma membrane and viral egress. *Sci Rep* 6: 19125
Kay JG, Koivusalo M, Ma X, Wohland T, Grinstein S (2012) Phosphatidylserine dynamics
in cellular membranes. *Mol Biol Cell* 23: 2198–2212
Kerr D, Tietjen GT, Gong Z, Tajkhorshid E, Adams EJ, Lee KYC (2018) Sensitivity of
peripheral membrane proteins to the membrane context: A case study of
phosphatidylserine and the TIM proteins. *Biochim Biophys acta Biomembr* 1860:
2126–2133
Kondratowicz AS, Lennemann NJ, Sinn PL, Davey RA, Hunt CL, Moller-Tank S,
Meyerholz DK, Rennert P, Mullins PF, Brindley M, *et al* (2011) T-cell
immunoglobulin and mucin domain 1 (TIM-1) is a receptor for Zaire Ebolavirus and
Lake Victoria Marburgvirus. *Proc Natl Acad Sci U S A* 108: 8426–8431
Kuroda M, Fujikura D, Nanbo A, Marzi A, Noyori O, Kajihara M, *et al* (2011) Interaction

Between TIM-1 and NPC1 Is Important for Cellular Entry of Ebola Virus. *J Virol* 89: 6481–6493

Kurovetz WR, Brunner F, Beubler E, Weyhenmeyer R, Lohaus R, Grob M, Mayer D (1982) Single dose pharmacokinetics of fendiline in humans. *Eur J Drug Metab Pharmaco*kinet 7: 105-110

Licata JM, Simpson-Holley M, Wright NT, Han Z, Paragas J, Harty RN (2003) Overlapping Motifs (PTAP and PPEY) within the Ebola Virus VP40 Protein Function Independently as Late Budding Domains: Involvement of Host Proteins TSG101 and VPS-4. *J Virol* 77: 1812–1819

Liu L (2014) Fields Virology, 6th Edition. *Clin Infect Dis* 59: 613–613

Llorente A, Skotland T, Sylvänne T, Kauhanen D, Rog T, Orlowski A, Vattulainen I, Ekroos K, Sandvig K (2013) Biochim Biophys Acta 1831: 1302-1309

Madara JJ, Han Z, Ruthel G, Freedman BD, Harty RN (2015) The multifunctional Ebola virus VP40 matrix protein is a promising therapeutic target. *Future Virol* 10: 537-546.

Maekawa M, Fairn GD (2015) Complementary probes reveal that phosphatidylserine is required for the proper transbilayer distribution of cholesterol. *J Cell Sci* 128: 1422-1433.

Maekawa M, Lee M, Wei K, Ridgway ND, Fairn GD (2016) Staurosporines decrease ORMDL proteins and enhance sphingomyelin synthesis resulting in depletion of plasmalemmal phosphatidylserine. *Sci Rep* 6: 35762

Martyushev A, Nakaoka S, Sato K, Noda T, Iwami S (2016) Modelling Ebola virus dynamics: Implications for therapy. *Antiviral Res* 135: 62–73.

Menke M, Gerke V, Steinem C (2005) Phosphatidylserine membrane domain clustering induced by annexin A2/S100A10 heterotetramer. *Biochemistry* 44: 15296–15303

Miller ME, Adhikary S, Kolokoltsov AA, Davey RA (2012) Ebolavirus Requires Acid Sphingomyelinase Activity and Plasma Membrane Sphingomyelin for Infection. *J Virol* 86: 7473–7483

Moe JB, Lambert RD, Lupton HW (1981) Plaque assay for Ebola virus. *J Clin Microbiol* 13: 791–793.

Moller-Tank S, Kondratowicz AS, Davey RA, Rennert PD, Maury W (2013) Role of the phosphatidylserine receptor TIM-1 in enveloped-virus entry. *J Virol* 87: 8327–8341.

Moller-Tank S, Maury W (2014) Phosphatidylserine receptors: Enhancers of enveloped virus entry and infection. *Virology* 468-470: 565–580.

Mühlberger E (2007) Filovirus replication and transcription. *Future Virol* 2: 205–215.

Mullard, A (2020) FDA approves antibody cocktail for Ebola virus. *Nat Rev Drug Discov* 19: 827

Nanbo A, Imai M, Watanabe S, Noda T, Takahashi K, Neumann G, Halfmann P, Kawoaka Y (2010) Ebolavirus is internalized into host cells via macropinocytosis in a viral glycoprotein-dependent manner. *PLoS Pathog* 6: e1001121

Nanbo A, Kawaoka Y (2019) Molecular Mechanism of Externalization of Phosphatidylserine on the Surface of Ebola Virus Particles. *DNA Cell Biol* 38: 115–120

Nanbo A, Maruyama J, Imai M, Ujie M, Fujioka Y, Nishide S, Takada A, Ohba Y, Kawoaka Y (2018) Ebola virus requires a host scramblase for externalization of phosphatidylserine on the surface of viral particles. *PLoS Pathog* 14: e1006848

Nelson EA, Barnes AB, Wiehle RD, Fontenot GK, Hoenen T, White JM (2016)
Clomiphene and Its Isomers Block Ebola Virus Particle Entry and Infection with
Similar Potency: Potential Therapeutic Implications. *Viruses* 8: 206.

Nguyen VK, Binder SC, Boianelli A, Meyer-Hermann M, Hernandez-Vargas EA (2015)
Ebola virus infection modeling and identifiability problems. *Front Microbiol.* 6: 257

Noda T, Sagara H, Suzuki E, Takada A, Kida H, Kawaoka Y (2002) Ebola Virus VP40
Drives the Formation of Virus-Like Filamentous Particles Along with GP. *J Virol* 76:
4855–4865

Oda S, Noda T, Wijesinghe KJ, Halfmann P, Bornholdt ZA, Abelson DM, Armbrust T,
Kawaoka Y, Stahelin RV, Saphire EO (2015) Crystal Structure of Marburg Virus
VP40 Reveals a Broad, Basic Patch for Matrix Assembly and a Requirement of the
N-Terminal Domain for Immunosuppression. *J Virol* 90: 1839–1848

Panchal RG, Ruthel G, Kenny TA, Kallstrom GH, Lane D, Badie SS, Li L, Bavari S, Aman
MJ. (2003) In vivo oligomerization and raft localization of Ebola virus protein VP40
during vesicular budding. *Proc Natl Acad Sci U S A* 100: 15936–15941

Reeves JP, Dowben RM (1969) Formation and properties of thin-walled phospholipid
vesicles. *J Cell Physiol* 73: 49–60

Reimão JQ, Mesquita JT, Ferreira DD, Tempone AG (2016) Investigation of calcium
channel blockers as antiprotozoal agents and their interference in the metabolism
of Leishmania (L.) infantum. *Evid Based Complement Alternat Med* 2016: 1523691

Ruigrok RW, Schoehn G, Dessen A, Forest E, Volchkov V, Dolnik O, Klenk HD,
Weissenhorn W (2000) Structural characterization and membrane binding
properties of the matrix protein VP40 of Ebola virus. *J Mol Biol* 300: 103–112

Scianimanico S, Schoehn G, Timmins J, Ruigrok RH, Klenk HD, Weissenhorn W (2000)
Membrane association induces a conformational change in the Ebola virus matrix
protein. *EMBO J* 19: 6732–6741

Skotland T, Sandvig K (2019) The role of PS 18:0/18:1 in membrane function. *Nat*
Commun 10: 2752

Soni SP, Stahelin RV (2014) The Ebola virus matrix protein VP40 selectively induces
vesiculation from phosphatidylserine-enriched membranes. *J Biol Chem* 289:
33590–33597

Stahelin RV (2014) Membrane binding and bending in Ebola VP40 assembly and egress.
Front Microbiol 5: 300

Stahelin RV, Scott JL, Frick CT (2014) Cellular and molecular interactions of
phosphoinositides and peripheral proteins. *Chem Phys Lipids* 182: 3–18.

Tsai WH (1985) Moment-preserving thresolding: A new approach. *Comput Vision, Graph*
Image Process 29: 377–393

van der Hoeven D, Cho K, Ma X, Chigurupati S, Parton RG, Hancock JF (2013) Fendiline
inhibits K-Ras plasma membrane localization and blocks K-Ras signal
transmission. *Mol Cell Biol* 33: 237–251

van der Hoeven D, Cho K, Zhou Y, Ma X, Chen W, Naji A, Montufar-Solis D, Zuo Y, Kovar
SE, Levental I et al (2018) Sphingomyelin Metabolism Is a Regulator of K-Ras
Function. *Mol Cell Biol* 38: e00373-17

Van Meer G, Voelker DR, Feigenson GW (2008) Membrane lipids: Where they are and
how they behave. *Nat Rev Mol Cell Biol* 9: 112–124.

Wan W, Clarke M, Norris MJ, Kolesnikova L, Koehler A, Bornholdt ZA, Becker S, Saphire

EO, Briggs JA (2020) Ebola and Marburg virus matrix layers are locally ordered
assemblies of VP40 dimers. *Elife* 9: e592225
Wang YE, Park A, Lake M, Pentecost M, Torres B, Yun TE, Wolf MC, Holbrook MR,
Freiberg AN, Lee B (2010) Ubiquitin-regulated nuclear-cytoplasmic trafficking of the
Nipah virus matrix protein is important for viral budding. *PLoS Pathog* 6: e1001186
Wang J, Qiao L, Hou Z, Luo G (2017) TIM-1 Promotes Hepatitis C Virus Cell Attachment
and Infection. *J Virol* 91: e01583-16
Wang P, van der Hoeven D, Ye N, Chen H, Liu Z, Ma X, Montufar-Solis D, Rehl KM,
Cho KJ, Thapa S *et al* (2021) Scaffold repurposing of fendiline: Identification of
potent KRAS plasma membrane localization inhibitors. *Eur J Med Chem* 217:
113381.
Watt A, Moukambi F, Banadyga L, Groseth A, Callison J, Herwig A, Ebihara H, Feldmann
H, Hoenen T (2014) A Novel Life Cycle Modeling System for Ebola Virus Shows a
Genome Length-Dependent Role of VP24 in Virus Infectivity. *J Virol* 88: 10511–
10524
Wen Y, Vogt VM, Feigenson GW (2018) Multivalent Cation-Bridged PI(4,5)P2 Clusters
Form at Very Low Concentrations. *Biophys J* 114: 2630–2639
Weyhenmeyer R, Fenzl E, Apecechea M, Rehm KD, Dyde CJ, Johnson KJ, Friedel R
(1987) Tolerance and pharmacokinetics of oral fendiline. *Arzneimittelforschung* 37:
58-62
Wijesinghe KJ, Stahelin V (2015) Investigation of the Lipid Binding Properties of the
Marburg Virus. *J Virol* 90: 3074–3085
Xiao X, Wang C, Chang D, Wang Y, Dong X, Jiao T, Zhao Z, Ren L, Dela Cruz CS,
Sharma L *et al* (2020) Identification of potent and safe antiviral therapeutic
candidates against SARS-CoV-2. *Front Immunol* 11: 586572
Yamashita Y, Oka M, Tanaka T, Yamazaki M (2002) A new method for the preparation
of giant liposomes in high salt concentrations and growth of protein microcrystals
in them. *Biochim Biophys Acta* 1561: 129–134
Yeung T, Gilbert GE, Shi J, Silvius J, Kapus A, Grinstein S (2008) Membrane
phosphatidylserine regulates surface charge and protein localization. *Science* 319:
210–213
Younan P, Iampietro M, Santos RI, Ramanathan P, Popov VL, Bukreyev A (2018)
Disruption of Phosphatidylserine Synthesis or Trafficking Reduces Infectivity of
Ebola Virus. *J Infect Dis* 218: S475–S485
Zhao H, Hakala M, Lappalainen P (2010) ADF/cofilin binds phosphoinositides in a
multivalent manner to act as a PIP(2)-density sensor. *Biophys J* 98: 2327–2336
1304
1305

Figure Legends

Figure 1

Clustering of PS by eVP40 *in vitro* and in HEK 293 cells. A Representative 3D reconstructed confocal images of immobilized GUVs (DPPC:Cholesterol:DPPS:PI(4,5)P$_2$:TopFluor® TMR-PS(red)). Left panel: GUVs incubated without eVP40-Alexa488. Right three panels: GUVs incubated with 1.25 μM eVP40-Alexa488 (green). **B** Index of correlation (Mander's coefficient) between TopFluor® TMR-PS and eVP40-Alexa488 of different GUVs compositions incubated with 1.25 μM eVP40-Alexa488. Values are reported as mean ± s.d. A one-way ANOVA with multiple comparisons was performed (n = 3); ****$p<0.0001$. **C** % TopFluor® TMR-PS quenching by eVP40 using GUVs (DPPC:Cholesterol:TopFluor®TMR-PS + increasing mol% of PS), 2.5% PI(4,5)P$_2$ was added to GUVs with 60% PS. Fluorescence spectra were recorded (Ex: 547 nm; Em: 550-600 nm); n=2. **D** Representative confocal images of HEK293 cells expressing various GFP-fused proteins (green) and supplemented with TopFluor® TMR-PS (red); scale bar= 10 μm. Yellow arrows indicate high intensity PS fluorescence regions **E** %PM with PS clusters = area of high intensity fluorescent PS clusters over total plasma membrane area from images in panel D. Black bars are control proteins and blue bars are eVP40 proteins. Values are reported as mean ± s.d.; N>18, n=3; A one-way ANOVA was performed with multiple comparisons compared to the control GFP %PS clustering (***$p=0.0007$, **$p=0.004$). DPPC: dipalmitoyl-phosphatidylcholine; DPPS: dipalmitoyl-phosphatidylserine; GUVs: giant unilamellar vesicles; PS: phosphatidylserine; PM: plasma membrane.

Figure 2

Effect of PS concentration on eVP40 binding affinity to and oligomerization on membranes. A-C SPR demonstrates that eVP40 affinity to LUVs increases in relation to PS concentration. **A** Representative normalized sensorgram of His_6-eVP40 binding to LUVs containing 1% PS indicating an apparent affinity of 2.5 µM. **B** Representative normalized sensorgram of His_6-eVP40 binding to LUVs containing 11% PS indicating an apparent affinity of 0.65 µM. **C** Representative normalized sensorgram of His_6-eVP40 binding to LUVs containing 22% PS indicating an apparent affinity of 0.18 µM. **D-E** PS concentration in LUVs enhances the ability of His_6-eVP40 to oligomerize on membranes. **D** Representative western blot of chemical crosslinking performed on His_6-WT-eVP40 following incubation with LUVs of varying PS content (detected by Mouse α-His antibody & HRP-Sheep α-Mouse). **E** Oligomerization capacity was determined from the western blot band density ratio of oligomers/(monomer + dimer) from chemical crosslinking experiments. A one-way ANOVA was performed with multiple comparisons compared to the control 0% PS LUVs control (30% PS *p= 0.021; 60% PS *p=0.017). n=3**.** Values are reported as mean ± s.d.; SPR: surface plasmon resonance; LUVs: large unilamellar vesicles; PS: phosphatidylserine; HRP: horseradish peroxidase.

Figure 3

PS concentration, localization, and dynamics in fendiline treated HEK293 cells. A Lipidomic analysis (LC/MS/MS) of total lipids extracted from HEK293 cells treated with the indicated concentration of fendiline (48 hours) demonstrated a significant reduction of total cellular PS levels. Values are normalized to DMSO control and are reported as mean

± s.d.; n=3; A one-way ANOVA was performed with multiple comparisons to the control DMSO (*p=0.0120, ***p=0.0003). **B-C** Analysis of PS plasma membrane localization in response to fendiline treatment in HEK293 cells. **B** Representative confocal images from live cell imaging of HEK293 cells expressing GFP-LactC2 and treated with fendiline for 48 hours; scale bars= 10 µm. **C** Effect of fendiline on PS plasma membrane localization was calculated by the ratio of GFP fluorescence at the plasma membrane intensity/intracellular intensity. Values are normalized to DMSO control and are reported as mean ± s.d.; N>15, n=3; A one-way ANOVA was performed with multiple comparisons compared to the DMSO control (**p=0.0031) **D-E** Analysis of PS clustering in HEK293 cells in response to fendiline treatment through N&B analysis. **D** *Left panel:* Representative images from time-lapse (30 frames) imaging of HEK293 expressing GFP-LactC2 and treated with fendiline for 48 hours; scale bar= 5 µm. *Middle panel:* Brightness and Intensity plots for each representative image. *Right panel:* Selection map correlating each pixel in the representative image to an oligomerization state (b value) (red: monomer-5mer, green: 5mer-10mer, blue: >10mer). **E** Average % pixels quantification from panel (d)= Percentage of GFP-LactC2 with brightness values corresponding to monomer-5mer (~1.-1.5), 5mer-10mer (~1.5-1.9) and >10mer (>1.9) over the total pixels within each image. Values are reported as mean ± s.d.; N≥9, n=3; A two-way ANOVA was performed with Dunnett's multiple comparisons comparted to the control DMSO % average pixels (****p<0.0001, **p=0.0043).GFP-LactC2: phosphatidylserine sensor; N&B: Number & Brightness analysis; PM: plasma membrane; PS: phosphatidylserine.

Figure 4

Evaluation of fendiline efficacy in the inhibition of authentic EBOV spread. A-D Effect of fendiline on EBOV infection. **A** Representative confocal images of Vero E6 cells infected with EBOV (Kikwit) at the indicated MOI and treated with the indicated concentration of fendiline. Cells were pretreated 24 hours prior to infection with the indicated concentration of fendiline. Post infection, cells were treated 1 hour later (d -1/0), treated every day (e.d), or treated every other day (e.o.d) and fixed at either 48 hours, 72 hours or 96 hours post infection. (green=EBOV; blue= nuclei). White numbering in top right corner indicates %infection. **B-D** Quantification of % inhibition of EBOV by fendiline. **B** 48 hours (MOI 1.0) **C** 72 hours (MOI 0.1) **D** 96 hours (MOI 0.1). Values are reported as mean ± s.d. A one-way ANOVA was performed with multiple comparisons was performed. n=3. EBOV: Ebola virus; MOI: multiplicity of infection; d. -1/0: treatment 1 hour after infection; e.d.: treatment every day; e.o.d.: treatment every other day.

Figure 5

Analysis of eVP40 and mVP40 cellular localization and oligomerization following fendiline treatment. A-C Effect of fendiline on eVP40 and mVP40 PM localization in HEK293 cells after 48 hours of treatment. **A** Representative confocal images from live cell imaging experiments of HEK293 cells expressing EGFP-WT-eVP40 (top panel) and EGFP-WT-mVP40 (bottom panel) after 48 hours of fendiline treatment. scale bars= 10 µm. Effect of fendiline on eVP40 **(B)** and mVP40 **(C)** PM localization was quantified by the ratio of EGFP fluorescence intensity at the PM / total EGFP fluorescence intensity (and normalized to DMSO control). N>15, n=3. Values are reported as mean ± s.d. A one-way ANOVA with multiple comparisons was performed compared to the DMSO control.

D-E Analysis of eVP40 oligomerization in HEK293 cells in response to 48 hour fendiline treatment using N&B analysis. **D** *Left panel:* Representative images from time-lapse (30 frames) of HEK293 expressing EGFP-WT-eVP40 and treated with fendiline for 48 hours. scale bar = 5 µm. *Middle panel:* Brightness and Intensity plots for each representative image. *Right panel:* Selection map correlating each pixel in the representative image to an oligomerization state (b value) (red: monomer-hexamer, green: hexamer-12mer, blue: 12mer-24mer, pink: >24mer). **E** Average % pixel quantification from panel (d)= % of GFP-WT-eVP40 with brightness values corresponding to monomer-hexamer (~1.-1.6), hexamer-12mer (~1.6-2.0), 12mer-24mer (2.0-3.2) and >24mer (>3.2) over the total pixels within each image. Values are reported as mean ± s.d.; N≥9, n=3; A two-way ANOVA was performed with Dunnett's multiple comparisons compared to the control DMSO % average pixels (**p=0.0035). PM: plasma membrane; N&B: Number & Brightness analysis.

Figure 6

VLP production and morphology in HEK293 cells in the presence of fendiline. A-D Functional budding assays assessed at 24 hours (A-B) and 48 hours (C-D) post treatment. **A** Representative western blot of budding assays performed at 24 hours. VLP samples (top panel) and cell lysate samples (bottom panel) collected from HEK293 cells and immunoblotted for eVP40 expression; GAPDH served as a loading control. eVP40 detected by (Rabbit α-eVP40 and HRP-Goat α-Rabbit); GAPDH detected by mouse α-GAPDH and HRP-Sheep α-Mouse) **B** Quantification of relative budding index at 24 hours post fendiline treatment. Relative budding index was determined by the western blot band

density of eVP40 in the VLP fraction/(total eVP40 cell lysate + eVP40 VLP band density) and was normalized to the DMSO control. Cell lysate eVP40 band density was normalized to GAPDH band density prior to use in budding index quantification. n=3. Values are reported as mean ± s.d. A one-way ANOVA was performed with multiple comparisons compared to the DMSO control. **C** Representative western blot of budding assays performed at 48 hours. VLP samples (top panel) and cell lysate samples (bottom panel) collected from HEK293 cells and immunoblotted for eVP40 expression; GAPDH served as a loading control. eVP40 detected by (Rabbit α-eVP40 and HRP-Goat α-Rabbit); GAPDH detected by (Mouse α-GAPDH and HRP-Sheep α-Mouse) **D** Quantification of relative budding index at 48 hours post fendiline treatment. Relative budding index was determined by the western blot band density of eVP40 in the VLP fraction/(total eVP40 cell lysate + eVP40 VLP band density) and was normalized to the DMSO control. Cell lysate eVP40 band density was normalized to GAPDH band density prior to use in budding index quantification. n=3. Values are reported as mean ± s.d. A one-way ANOVA was performed with multiple comparisons compared to the DMSO control. (*p=0.0260) **E-F** SEM micrographs of HEK93 cells. **E** Representative micrographs of mock transfected HEK293 cells harvested after 48 hours of no treatment or DMSO treatment. **F** Representative micrographs of HEK293 cells expressing FLAG-eVP40 and harvested after 48 hours of no treatment, DMSO treatment, or the indicated concentration of fendiline. VLPs: virus like particles; SEM: scanning electron microscopy; GAPDH: glyceraldehyde 3-phosphate dehydrogenase; HRP: horseradish peroxidase.

Figure 7

Effect of fendiline on eVLP morphology and TIM-1 dependent eVLP entry. **A-C** TEM analysis of eVLP morphology. **A** Representative transmission electron micrographs of eVLPs purified from HEK293 cells expressing FLAG-eVP40 and eGP following 48 hours of DMSO (left panel) or 5 µM fendiline treatment (right panel). **B** Quantification of eVLP length (µm) of DMSO-derived eVLPs (black) and fendiline-derived eVLPs (blue). N>50, n=3. Values are reported as mean ± s.d. A two-tailed t-test was performed (**p=0.0139). **C** Quantification of eVLP diameter (nm) of DMSO-derived eVLPs (black) and fendiline-derived eVLPs (blue). N>50, n=3. Values are reported as mean ± s.d. A two-tailed t-test was performed (*p=0.0430). **D-E** Fluorescence based DiI TIM-1 dependent entry assay. **D** Representative confocal images from the DiI-entry assay comparing entry of eVLPs produced from DMSO (top panel) and fendiline-treated HEK293 cells (bottom panel) into target cells (HEK293 cells transiently expressing increasing amounts of TIM-1; 0.0 µg, 0.5 µg, 1.0 µg). A stack of 10 frames was acquired for each image. DiI (initially red) was recolored to yellow for easier observation in print; blue (Hoechst 3342 stain); scale bar = 10 µm. **E** Quantification of eVLP entry was performed by calculating the total number of DiI punctate / the total number of DiI-positive cells. Three images from each z-stack was quantified. N=9, n=3. Values are reported as mean ± s.d. A one-way ANOVA was performed with multiple comparisons against the 0.0 µg TIM-1 condition for both DMSO- and fendiline derived eVLPs.(****p<0.0001; **p=0.0093). eVLP: entry-competent viral like particles; TEM: transmission electron microscopy; TIM-1: t-cell immunoglobulin receptor-1; eVLPs: entry-competent VLPs; eGP: Ebola glycoprotein; DiI: 1,1'-Dioctadecyl-3,3,3',3'-Tetramethylindocarbocyanine Perchlorate.

Figure 8

Calibrated mathematical model reproduces key observations in multiple experimental datasets. Percentage infected cells is shown for various fendiline concentrations given prior to infection (d-1/0, **A-C**) or daily (e.d., **D-F**). (**A,D**) MOI 1; (**B,C,E,F**) MOI 0.1. (**G-H**) Model predicted cell and viral dynamics for MOI 0.1. (**i**) Model predicted dose response curves for fendiline effects on viral budding and entry in the BSL4 experiments.

Expanded View Figure Legends

Expanded View 1 (EV1)

Fluorescence profiles of PS and WT-eVP40 through confocal microscopy and plot profile analysis of fluorescently labeled GUVs and Alexa488-eVP40. **A** Representative confocal images of fluorescently labeled (TopFluor® TMR-PS) GUVs with varying lipid compositions (red) following incubation with eVP40-Alexa88 (green). Colocalization between eVP40 and PS within GUVs was indicated by plot profile analyses of fluorescence signals between TopFluor® TMR-PS (red dotted line) and eVP40-Alexa488 (green solid line) performed at indicated open yellow lines in **a** and shown in **B-D**. **B** Plot profile analysis of eVP40-Alexa488 and control GUVs (DPPC:Cholesterol:0.2mol% TopFluor® TMR-PS), **C** Plot profile analysis of eVP40-Alexa488 and PS GUVs (DPPC:Cholesterol:0.2mol% TopFluor® TMR-PS:DPPS). * indicates overlap in fluorescence signals. **D** Plot profile analysis of eVP40-Alexa488 and PS+PI(4,5)P$_2$ GUVs (DPPC:Cholesterol:0.2mol% TopFluor® TMR-PS:DPPS:PI(4,5)P$_2$). **E** Plot of WT eVP40 Alexa-488 membrane fluorescence/total fluorescence for lipid

compositions shown in a-d, n = 3, ****p<0.0001. PS: phosphatidylserine; DPPC: dipalmitoyl-phosphatidylcholine; DPPS: dipalmitoyl-phosphatidylserine.

Expanded View 2 (EV2)

Fluorescence profiles of PS and GFP-WT-eVP40 in HEK 293 cells through confocal microscopy. A Representative confocal images from live cell imaging of HEK293 expressing various GFP-fused proteins (GFP; green) following supplementation with TopFluor® TMR-PS (red). Solid white lines indicate where plot profile analysis was performed; scale bar= 10 μm. **B-C** Validation of ability to detect exogenously added fluorescently labelled PS within the inner leaflet of the plasma membrane of cells **B** Plot profile analysis of HEK293 cells expressing cytosolic GFP. **C** Plot profile analysis of HEK293 cells expressing the PS sensor GFP-LactC2 **D-E** Investigation of functionally distinct eVP40 proteins ability to bind to fluorescently labelled PS within the inner leaflet of the plasma membrane in living cells. **D** Plot profile analysis of HEK293 cells expressing GFP-WT-eVP40. **E** Plot profile analysis of HEK293 cells expressing GFP-K224A-eVP40 (PS-binding residue mutant). **F** Plot profile analysis of HEK293 cells expressing GFP-WE/A-eVP40 (oligomerization deficient mutant). TopFluor TMR-PS fluorescence signal intensity (red dotted line) and GFP fluorescence signal intensity (green solid line).

Expanded View 3 (EV3)

Validation of in vitro and cellular PS clustering experiments. **A** GUVs of varying compositions were imaged prior to and following the addition of 1.25 μM eVP40-Alexa. eVP40-Alexa 488 enrichment ratios at the membrane of the GUVs were calculated by the

ratio the Alexa488 fluorescence intensity at the GUV membrane / Alexa488 total fluorescence. Values are reported as mean ± s.d.; of n=3. A two-way ANOVA with multiple comparisons was performed. *p<0.0001. **B** Representative images of the step-wise image analysis workflow of quantifying PS clustering in living HEK293 cells expressing GFP-fused proteins using a custom ImageJ macro. scale bar= 10 μm. **C** Representative images from live cell imaging experiments of HEK293 cells expressing control GFP-fused proteins specific for the plasma membrane (GPI), and specific lipids, PS (LactC2) and PI(4,5)P$_2$ (PLCδ-PH). scale bar = 10 μm.

Expanded View 4 (EV4)

Profile of fendiline toxicity and efficacy in BSL-2 models. A CellTiter-Glo® viability results of HEK293 cells. HEK293 cells were treated with fendiline for 24 hours (black line) and 48 hours (blue line) and viability was assessed as a % viability of control (n = 3). **B** PS saturation analysis from lipidomic analysis (LC/MS/MS) of total lipids extracted from HEK293 cells treated with the indicated concentration of fendiline (48 hours). Values are normalized to DMSO control and are reported as mean ± s.d.; n=3; A one-way ANOVA was performed with multiple comparisons compared to the control DMSO. **C** PA level analysis from lipidomic analysis (LC/MS/MS) of total lipids extracted from HEK293 cells treated with 5 μM fendiline (48 hours). Values are normalized to DMSO control and are reported as mean ± s.d.; n=3; A two-tailed t-test was performed. **D-E** Analysis of PS plasma membrane localization in response to 24 hour fendiline treatment. **D** Representative confocal images from live cell imaging of HEK293 cells expressing GFP-LactC2 and treated with fendiline for 24 hours; scale bars = 10 μm. **E** Effect of fendiline

on PS plasma membrane localization was calculated by the ratio of GFP fluorescence at the (plasma membrane intensity/intracellular intensity). Values are normalized to DMSO control and are reported as mean ± s.d.; N>15, n=3; A one-way ANOVA was performed with multiple comparisons compared to the DMSO control. **p=0.0045; ***p=0.0003. **f** N&B analysis of HEK293 cells expressing the control GFP. Analysis was performed at 48 hours post treatment (DMSO) to align with N&B analysis performed on experiments with HEK293 cells expressing GFP-LactC2 or GFP-eVP40 and treated with the control or fendiline. *Left panel:* Representative images from time-lapse (30 frames) of HEK293 expressing EGFP and treated with fendiline for 48 hours. scale bar = 5 μm. *Middle panel:* Brightness and Intensity plots for representative image. *Right panel:* Selection map correlating each pixel in the representative image to an oligomerization state (b value) (red: monomer). PS: phosphatidylserine; PA: phosphatidic acid; N&B: number & brightness analysis.

Expanded View 5 (EV5)

Vero E6 toxicity and effect of fendiline on eVP40 plasma membrane localization at 24 hours post treatment. A CellTiter-Glo® viability results of Vero cells. Cells were treated with control or fendiline for 48 hours according to the BSL-4 infection model; d-1/0 (black line), e.d. (blue line) and e.o.d (gray line) and viability was assessed as a % viability of control (n = 3). **B-C** Effect of fendiline on eVP40 PM localization in HEK293 cells after 24 hours of treatment. **B** Representative confocal images from live cell imaging experiments of HEK293 cells expressing EGFP-WT-eVP40 after 48 hours of fendiline treatment. scale bars= 10 μm. **C** Effect of fendiline on eVP40 PM localization was

quantified by the ratio of EGFP fluorescence intensity at the PM / total GFP fluorescence

intensity (and normalized to DMSO control). N>15, n=3. Values are reported as mean ±

s.d. A one-way ANOVA with multiple comparisons was performed compared to the DMSO

control. PM: plasma membrane.

Tables

Table 1. Mathematical model parameter descriptions and values

Symbol	Description	Unit	Value from calibration to budding and entry assays	Ranges based on fits to budding and entry assays	Value from calibration to BSL4 data – using ranges defined by budding and entry assays
β	Infection rate constant of target cells by free virions	$(Virion.day)^{-1}$	N/A	0 - inf	6.6×10^{-6}
δ	Death rate constant of infected cells	Day^{-1}	N/A	0 - inf	4.6×10^{-6}
c	Decay rate constant of free virions	Day^{-1}	N/A	0 - 5	5
p	Viral production rate constant by infected cells	$Virions.(Infected\ cell.day)^{-1}$	N/A	1 - inf	49
H_b	Hill constant for Fendiline effects on VLP budding		2.7	0.5 - 7	3.4
H_e	Hill constant for Fendiline effects on VLP entry		11	0.5 - 20	4.7

C_{50}^{b}	Concentration with 50% of maximum Fendiline effect on VLP budding	µM	5.7	2 - 20	18
C_{50}^{e}	Concentration with 50% of maximum Fendiline effect on VLP entry	µM	4.8	0.5 - 20	15
E_{max}^{b}	Maximum Fendiline effect on VLP budding		0.98	0.3 - 1	0.3
E_{max}^{e}	Maximum Fendiline effect on VLP entry		0.76	0.3 - 1	0.33

Table 2. Antibodies used in immunoblotting

Target	Species	Tag	Dilution	Cat. #	Source
eVP40	Rabbit		1/40,000	0301-010	IBT Bioservices
Rabbit	Goat	HRP	1/5,000	205718	Abcam
GAPDH	Mouse		1/5,000	8245	Abcam
Mouse	Sheep	HRP	1/10,000	6808	Abcam
EBOV-GP (KZ52)	Human		1.0 µg/ml	0260-001	IBT Bioservices
Human	Goat	Alexa-488	0.7 µg/ml	A11013	Invitrogen
Mouse	Goat	Alexa-488	1.0 µg/ml	A11029	Invitrogen
6xHis	Mouse		1/1,000	A5588	Sigma

Figure 1

Figure 2

Figure 3

EBOV - Fendiline Treated

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Expanded View Figure 1

Expanded View Figure2

Expanded View Figure 3

Expanded View Figure 4

a.

Vero cells (48 hrs)

% viability of control

- Fendiline (d. -1/0)
- Fendiline (e.d.)
- Fendiline (e.o.d.)

[Fendiline]

c.

PM / Total fluorescence Intensity

DMSO   500 nM   1 μM   5 μM

[Fendiline]

b.

DMSO   500 nM   1 uM   5 uM

GFP-eVP40

Expanded View Figure 5