Cortical waves mediate the cellular response to electric fields

Authors: Qixin Yang[1,2], Yuchuan Miao[3], Leonard J. Campanello[1,2], Matt J. Hourwitz[4], Bedri

Abubaker-Sharif[3], Abby L. Bull[1,2], Peter N. Devreotes[3], John T. Fourkas[2,4], Wolfgang Losert[1,2]*.

Affiliations:

[1]Department of Physics, University of Maryland, College Park, MD 20742

[2]Institute for Physical Science and Technology, University of Maryland, College Park, MD

20742

[3]Department of Cell Biology, Johns Hopkins University, Baltimore, MD 21218

[4]Department of Chemistry & Biochemistry, University of Maryland, College Park, MD 20742

*Correspondence to: wlosert@umd.edu

Abstract:

Electrotaxis, the directional migration of cells in a constant electric field, is important in regeneration, development, and wound healing. Electrotaxis has a slower response and a smaller dynamic range than guidance by other cues, suggesting that the mechanism of electrotaxis share both similarities and differences with chemical-gradient-sensing pathways. We examined a mechanism centered on the excitable system consisting of cortical waves of biochemical signals coupled to cytoskeletal reorganization, which has been implicated in random cell motility. We use electro-fused giant *Dictyostelium discoideum* cells to decouple waves from cell motion and employ nanotopographic surfaces to limit wave dimensions and lifetimes. We demonstrate that wave propagation in these cells is guided by electric fields. The wave area and lifetime gradually increase in the first 10 minutes after an electric field is turned on, leading to more abundant and wider protrusions in the cell region nearest the cathode. The wave directions display "U-turn" behavior upon field reversal, and this switch occurs more quickly on nanotopography. Our results suggest that electric fields guide cells by controlling waves of signal transduction and cytoskeletal activity, which underlie cellular protrusions. Whereas surface receptor occupancy triggers both rapid activation and slower polarization of signaling pathways, electric fields appear to act primarily on polarization, explaining why cells respond to electric fields more slowly than to other guidance cues.

Introduction

Electrotaxis, which refers to the directed migration of cells under the guidance of an electric field (EF), is important in wound healing, development, and regeneration (Cortese et al., 2014; Lin et al., 2008; Zhao et al., 2006). EFs have been shown to cause several key signaling molecules to be

distributed asymmetrically across cells (Sato et al., 2009; Zhao et al., 2002, 2006), setting up cell

polarity. The one-order-of-magnitude range of EF strengths sensed by cells (Zhao et al., 2002) is

considerably smaller than the four-orders-of-magnitude concentration sensitivity in chemotaxis

(Harvath et al., 1991). Furthermore, whereas cells respond to chemical guidance cues on a time

scale of seconds and develop polarity over several minutes, the response to EFs can take up to 10

minutes or more after the EF is turned on (Wang et al., 2014; Zhao et al., 2006). These

differences raise the possibility that the rapid gradient sensing mechanisms do not serve as

primary mediators of EF sensing by cells. In this study, we examine whether, after turning on an

EF, the gradual polarization of the excitable biochemical networks that organize actin

polymerization comprises a slow-acting mediator of the cellular response to the EF.

Actin polymerization, coordinated with its associated signaling molecules, self-organizes into

microscale spatial regions that travel as waves across plasma membranes. These waves drive

various cell behaviors, such as migration and division (Bhattacharya et al., 2019; Bretschneider

et al., 2009; Flemming et al., 2020; Gerhardt et al., 2014; Gerisch, 2010). The wave system can

be described as a coupled signal transduction excitable network - cytoskeletal excitable network

(STEN-CEN) (Devreotes et al., 2017; Miao et al., 2019). STEN-CEN has the characteristics of

an excitable system, including exhibiting an activation threshold for wave initiation and

experiencing refractory periods. It has been shown that the STEN-CEN wave properties dictate

protrusion properties (Miao et al., 2019). Tuning the activity levels of key components in STEN-

CEN changes wave patterns, which leads to the transition of protrusion profiles. An

activator/inhibitor, reaction/diffusion system model successfully recapitulates the experimental

results (Bhattacharya et al., 2020; Bhattacharya & Iglesias, 2018). For simplicity, here we will

refer to STEN-CEN waves as cortical waves.

One challenge in investigating whether cortical waves can act as the mediators of EFs is that in

many of the cell types that show a strong response to EFs, the wave area is comparable to the cell

area. Furthermore, waves are generated at the leading edge of the cell during directed migration

(Xu et al., 2003), so that wave dynamics are tightly coupled with cell dynamics. For instance,

when a cell responds to an EF reversal, waves typically remain at the cell front as the cell turns.

It is not known whether the waves drive cells to turn or the cell polarity keeps the previous

leading edge more active so that this edge responds first.

To distinguish between wave response and cell motion, we produced electro-fused giant *D.*

discoideum (Neumann et al., 1980) with diameters up to ten times larger than that of an

individual cell. Multiple simultaneous waves can be generated across the surface contact area of

a giant cell (Gerhardt et al., 2014). These waves also generate actin-filled macropinosomes on

the dorsal membrane (Veltman et al., 2016). The giant cells provide an excellent opportunity to

study cortical wave dynamics in multiple cell regions simultaneously.

We further use nanotopography to alter the waves' spatial structures and characteristic timescales.

Upon contact with nanotopography, cells produce quasi-1D wave patches. The phenomenon of

guided actin polymerization by nanotopography is known as esotaxis (Driscoll et al., 2014),

which has been investigated in detail (Ketchum et al., 2018; Lee et al., 2020). There are several

advantages of incorporating nanotopography in our study. First, these waves persist for a shorter

time on nanotopography than on flat surfaces, enabling us to investigate whether wave systems

with different characteristic timescales respond to EFs differently. Second, waves on ridged

surfaces have shorter lifetimes than those on flat surfaces, and thus only propagate in local

regions of giant cells. Therefore, nanotopography allows us to distinguish between local and

global mediation of the EF response.

Results

Cortical waves and cell migration can be studied independently in giant cells

We imaged cells that simultaneously expressed both limE-RFP and PH_{Crac}-GFP. The former

allows us to monitor filamentous actin (F-actin), which represents CEN activities. The latter

enables us to monitor phosphatidylinositol-3,4,5-trisphosphate (PIP3), an indicator of STEN

activities. In single, differentiated cells, usually only one wave is generated at the leading edge

(Fig. 1a and Video 1), and the wave motion is coupled with cell motion. For instance, when the

cell in Fig. 1a changed its direction of motion, the wave remained at the leading edge (72 s - 120

s).

In giant cells, multiple waves were initiated randomly and propagated radially across the basal

membranes (Fig. 1b and Video 2). CEN is driven by STEN, but has a substantially shorter

characteristic timescale. Thus, PIP3 waves displayed band-like shapes, whereas F-actin appeared

across the bands with higher levels at the rims of PIP3 waves (Miao et al., 2019). As shown in

Fig. 1b, colliding waves did not cross, but instead rotated by 90° (Fig. 1b, 150 s - 200 s). This

behavior is suggestive of a refractory period following excitation, which is a hallmark of an

excitable system. On nanoridges, the giant cells generated multiple, quasi-1D patches of F-actin

and PIP3 with shorter lifetimes than on flat surfaces (Fig. 1c and Video 3). Some waves formed

and propagated for a short distance (Line 2 in Fig. 1c), whereas others formed and then quickly

dissipated (Line 3 in Fig. 1c). The wave dissipation can be explained in terms of an excitable

system with lateral inhibition, in which the dispersion of the inhibitor is faster than that of the

activator. Thus, the waves eventually dissipate due to the spatial accumulation of the inhibitor.

Prior studies have shown that in this situation, the excitable system threshold determines the

wave duration (Bhattacharya et al., 2020; Ermentrout et al., 1984). As was the case on flat

surfaces, 1D patches occurred throughout the basal surfaces on ridges, and thus were

independent of cell motion.

Waves were also generated on the dorsal planes. In contrast to basal waves, which propagated

across the surface contact (Video 2 and Video 3), dorsal waves were associated with membrane

deformations, and resembled macropinosomes (Video 4). Based on 3D lattice light-sheet images

of a cell plated on nanoridges (Fig. 1d and Video 5), activation of PIP3 and F-actin was

correlated in both basal waves and dorsal waves. However, the dorsal waves were primarily

generated in cuplike structures, whereas the stripe-like basal waves spanned the entire basal

plane. In all cases, PIP3 activity was coordinated with F-actin activity (Profiles in Fig. 1a, 1b,

and 1c), both in the absence (Fig. 1) and presence of an EF (Fig. 1 – figure supplement 1, Video

6). Therefore, in the experiments described below, we only monitored F-actin activity.

Fig. 1. STEN-CEN waves in single cells and giant cells. a. Snapshots of a differentiated, single

D. discoideum cell expressing limE-RFP and PHcrac-GFP, with cell boundaries denoted with

blue dashed lines. The right column shows the normalized intensity of limE and PHcrac from the

arrows in the merge images. The scale bars are 10 µm. **b**. Snapshots of an electrofused giant *D.*

discoideum cell on a flat surface, with scanning profiles in the right column. All scale bars are 10

µm. **c.** A snapshot of an electrofused giant cell on the ridged surface. The left kymographs are

from the line 2 and line 3 specified in the merged image. Line 2 shows a wave propagating along

nanoridges, and line 3 shows a wave that existed briefly and then dissipated. **d.** 3D

reconstruction (single time point) of a single *D. discoideum* cell plated on nanoridges, acquired

using a lattice light-sheet microscope. Here we show the top aspect view (top row) and the side

aspect view (bottom row). On the dorsal membrane of the cell, there are waves forming

microcytotic cups (triangle) on the curved membrane, and on the basal membrane, there are

streak-like waves (asterisk). The red channel represents limE-RFP, and the green channel

represents PHcrac-GFP. As both the side and top views show, the dorsal waves and basal waves

are independent structures, but both are composed of coordinated F-actin and PIP3.

EFs increase the area, duration, and speed of waves on nanoridges.

We found that giant cells respond to a narrow range (15 V/cm to 20 V/cm) of EF amplitudes

(Video 7 and Video 8), and that higher voltage (35 V/cm) damaged cells. The 1D waves

generated on nanoridges related to esotaxis enabled us to quantify the effects of a 20 V/cm EF

(all EFs used here are of this magnitude, see Fig. 2 – figure supplement 1and Methods for more

details about the electrotaxis experiments) on the areas, durations, and speeds of waves. Fig. 2a

shows snapshots of the dynamics of F-actin in a giant cell on parallel nanoridges with a 1.6 µm

spacing. In the absence of an EF (top row in Fig. 2a), individual actin polymerization events

were initiated in patches on the basal surfaces (Fig. 2a and Video 9).

In the first several minutes after turning on the EF, most patches propagated as a wave along a

single ridge (Fig. 2a, blue inset). After the EF was on for 10 min, some patches appeared to

undergo coordinated motion across several ridges (Fig. 2a, pink inset). We calculated the ratio of

F-actin occupancy to cell area, and found that an EF increased the overall level of actin

polymerization by a factor of two to three (Fig. 2b). Actin patches were larger in the presence of

an EF and organized into larger groups located preferentially at the cell front (bottom row in Fig.

2a, 20 min, and 25 min), leading to wider protrusions at cell fronts that drove directed cell

migration (Fig. 2 – figure supplement 2). To determine whether the groups comprised a single,

large wave growing across multiple ridges or multiple, small patches nucleated in close

proximity, we measured the dynamics of the patches using optical flow (Lee et al., 2020),

focusing on the patch edges (Fig. 2c, left image. See Methods for more details). If both edges of

a patch were moving in the same direction, the structure was classified as a single, large wave. If

the edges were not coordinated, the patch was classified as multiple, individual structures. This

method enabled us to capture accurately waves that span across multiple ridges and are moving

coordinately.

Once the large actin structures were classified, their instantaneous dimensions were measured

parallel and perpendicular to the ridges (Fig. 2c). Density scatter plots of both dimensions exhibit

elliptical contours (Fig. 2d), suggesting that nanotopography constrains wave growth. With an EF

parallel to the ridges, the waves broadened in both directions (Fig. 2d). The average increases in

wave dimension parallel and perpendicular to the ridges were 20% and 13%, respectively, and

the average increase in wave area was 44%. An increase in wave duration was also observed,

with the minimum wave area correlated to the duration (Fig 2e, black circles). The wave area

depends exponentially on the maximum wave duration (Fig. 2e, solid black lines), allowing us to

extract a characteristic wave time scale via

$$Area_{min} = C * e^{\frac{Duration}{T}} \quad .$$ (1)

Here C is a constant, and T is the characteristic time scale, which is 48 s with no EF and 61 s in

the presence of a 20 V/cm EF. This difference is consistent with the EF drawing the system

closer to the excitability threshold. An average increase of 9% in wave propagation speed was

also observed in the presence of an EF (Fig. 2f).

Fig. 2. EFs alter F-actin wave properties. a. limE images of a giant cell on nanoridges without

an EF (top) and in a 20 V/cm EF turned on at 0 min (bottom). **b**. The temporal change of the

percentage of the cell area occupied by limE without an EF (blue, $N_{cell} = 5$) and in a 20 V/cm EF

introduced at 0 min (red, $N_{cell} = 4$). The shaded areas represent the mean plus or minus one

standard deviation **c**. Division of groups of waves. The color represents the orientation of

optical-flow vectors according to the color wheel. The green arrows are the optical-flow vectors,

the length of which correspond to the magnitude of motion. The left image is an example of a

large structure composed of two independent substructures, where the vectors at the right edge

are not moving in the same direction. The wave scales in the directions perpendicular to (blue

arrows) and parallel to the ridges (orange arrow) were measured on the preprocessed waves. **d.**

Density scatter plots of wave scales parallel to ridges vs. perpendicular to ridges. **e.** Density

scatter plots of actin-wave dimension vs. actin-wave duration. For each wave duration, the 5

points with the smallest wave areas (black circles) were selected to fit the boundaries (solid black

lines). **f.** Distributions of wave propagation speeds before (blue, N_{wave} = 125) and after (red, N_{wave}

= 163) applying an EF. The analyses in d-f were based on N = 4 independent experiments. The

two distributions are different (Two-sample t-test, P = 0.017).

Figure 2 – source data 1. Related to Fig. 2b

Figure 2 – source data 2. Related to Fig. 2d

Figure 2 – source data 3. Related to Fig. 2e

Figure 2 – source data 4. Related to Fig. 2f

EFs guide the direction of actin waves.

Next, we consider the directional guidance of actin waves by EFs on nanoridges (Fig. 3a and

Video 9) and on flat surfaces (Fig. 3b and Video 10, see Methods for details). The EF was

introduced at 0 min (T_1), and in the first 2 min had little effect on the actin dynamics on any

surface. On nanoridges, actin waves continued to propagate preferentially along the ridges (Fig.

3a, T_1). On flat surfaces, the waves propagated radially in groups, as seen from the broad

distribution at T_1 in Fig. 3b. In the presence of an EF, the waves propagated preferentially

towards the cathode within ~15 min (Fig. 3a, 3b, T_2). The perpendicular spread was significantly

more limited on nanoridges (Fig. 3a, T_2).

The direction of the EF was reversed after the cell had commenced steady directional migration,

which took ~20 to 25 min. Following the field reversal, waves on ridged surfaces reoriented

toward the new cathode within 5 min (Fig. 3a, T_3). On flat surfaces, the wave propagation

direction was perpendicular to both the previous and the new EF directions at ~7 min after the

field reversal (Fig. 3b, T_3). Preferential propagation towards the new cathode occurred after ~13

min (Fig. 3b, T_4). The difference in response time between nanoridges and flat surfaces may be

related to the fact that waves persist longer on flat surfaces than on nanoridges (Fig. 3c).

Fig. 3d shows the continuous temporal changes of the orientation distributions. On nanoridges,

the preferred wave directions switched directly following EF reversal (left plot in Fig. 3d),

whereas on flat surfaces waves maintained a preferred direction that changed continuously in a

U-turn behavior (Right plot in Fig. 3d). In contrast, although single *D. discoideum* cells undergo

U-turns in response to EF reversal (Sato et al., 2007), giant cells did not (Video 10).

Wave turning may be related to differences in the patterns of wave expansion (Fig. 3e). On flat

surfaces, waves started from a small patch (Fig. 3e, S1) and eventually broke into band-shaped

waves (Fig. 3e, S4). During directed migration, the intermediate expansion of actin waves (Fig.

3e, S2, S3 in the top row) was biased by the EF, resulting in band-shaped waves propagating

preferentially towards the cathode (Fig. 3e, S4 in top row). After EF reversal, waves expanded in

all directions (Fig. 3e, S2, S3 in bottom row), such that optical-flow analysis captured turning

behavior more frequently.

We also simultaneously imaged limE-RFP at the basal plane (near the surface contact) and the

dorsal plane (6 µm higher). Dorsal waves (Video 11 and 12) are localized at cell fronts and

rearranged to the new fronts following EF reversal (Fig. 3 – figure supplement 1). Rather than

directly switching preferential direction, dorsal waves gradually turned toward the new cathode

(Fig. 3 – figure supplement 1), in a manner similar to that of basal waves on flat surfaces. Thus,

two different response times to EF reversal exist within the same cell, with a faster response for

basal waves guided by nanotopography and a slower response for the free dorsal waves. On flat

surfaces, the two responses are synchronized (Fig. 3 – figure supplement 2).

Fig. 3. EFs guide actin waves. a, b Optical-flow analysis of actin-wave dynamics in giant cells

on ridged and flat surfaces. The top row shows a time series of limE images for giant cells

overlaid with optical-flow vectors, the color of which is coded according to the color wheel. The

accompanying polar plots show the corresponding orientation displacements of optical-flow

vectors. For both a and b the EF was turned on at 0 min. The bottom time stamp indicates when

the EF was reversed from the cathode being on the right (red) to the cathode being on the left

(green). **c.** Distributions of wave duration from 3 independent days of experiments. The

distributions were weighted by wave area, because the number of long-lasting large waves on

flat surfaces (N_{wave} = 359) is smaller than the number of short-lived small patches on ridged

surfaces (N_{wave} = 658). Correspondingly, the absolute waves counts do not match the pixel-based,

optical-flow analysis in a and b. Based on a two-sample t-test on the wave areas on flat surfaces

vs. on ridges, the null hypothesis was rejected at the 5% significance level with $p = 2 \times 10^{-15}$. **d.**

Kymographs of orientation displacements of optical-flow vectors. The *x*-axes of the kymographs

represent time, and the *y*-axes represent orientation. The colors represent the proportions. The EF

was turned on at time T_1, and was reversed at the time denoted by the black arrow **e.** LimE

snapshots showing the patterns of actin-wave expansion during steady directed migration in a

constant EF (top) and after reversing the EF direction (bottom). The blue arrows point to specific

stages of wave expansion. W: Wave, S: Stage of wave expansion. The right panel is a cartoon

illustrating the patterns of actin-wave expansion during directed migration in EFs (top) and after

EFs were reversed (bottom).

Figure 3 – source data 1. Related to Fig. 3c

277   **Subcellular spatial inhomogeneity of the response to EFs on nanoridges.**

278   Although waves in migrating *D. discoideum* cells localize predominantly at the leading edge

279   (Weiner et al., 2007; Zhao et al., 2002), waves are observed across the basal layer in giant *D.*

280   *discoideum* cells. We analyzed the smaller, shorter-lived waves on nanoridges. Although the

281   wave locations were distributed essentially uniformly throughout cells in the absence of an EF,

282   more waves were generated at the cell fronts in the presence of an EF (Fig. 4a and Video 9). In

283   addition, the average area per wave was larger near the front of cells in an EF (Fig. 4b). We also

284   measured the wave properties in the single cells scattered throughout the field of view but did not

285   observe a corresponding gradient of wave properties among single cells closer to the cathode

286   versus the cells closer to the anode. This result indicates that the spatial inhomogeneity shown in

287   Figs. 4a, b was caused by the EF rather than by the absolute electrical potential relative to the

288   ground (Fig. 4 – figure supplement 1).

289

290   We explored the response of this inhomogeneity to EF reversal by tracking each wave location

291   relative to the cell centroid in the 12 min following EF reversal (Fig. 4c). New waves started to

292   appear near the side of the cell facing the new cathode within 3 min (Fig. 4c, left region of P_2),

293   whereas the complete inhibition of wave generation near the old cell front took longer (Fig. 4c,

294   right region of P_5). This observation suggests that the initiation at a new cell front and the

295   inhibition of waves at the old front are regulated by two distinct processes with different

296   timescales.

297

298   Next, we looked at the time required to switch propagation direction in different subcellular

299   regions following EF reversal. The basal membrane was segmented into an "old front" region

(facing the original cathode) and a "new front" region (facing the new cathode), as illustrated in

Fig. 4d. The distributions of wave propagation directions show that waves in the new front

region switched their preferential direction at ~4 min. In contrast, waves in the old front region

changed their preferential direction on a time scale of ~ 7 min (Figs. 4 e, f). Our analysis further

shows that larger waves in the old fronts are less sensitive to EF reversal than those in the new

fronts (Fig. 4 – figure supplement 2).

Fig. 4. Spatial inhomogeneity of the response to EFs on nanoridges. a. Density scatter plots

of the wave area vs. x position of the wave relative to the cell center. Nanoridges and EF are

orientated in the x-direction. The difference of x coordinates of cell center and wave location was

calculated, then the value was further normalized by the cell radius. Each point represents a wave,

and all the points were collected from 5 independent experiments. The left plot is for a period in

which there was no EF (N_{wave} = 296), and the right plot is for a period in which there was a 20

V/cm EF, during which the cells exhibited steady directional migration (N_{wave} = 224) . For each

experiment without an EF, the EF was always turned on several minutes later. Thus, we defined

the direction in which cathode was located in the presence of an EF as the positive direction in

the absence of an EF. The color code corresponds to the density of points. **b.** Average wave area

in sub-cellular regions. The points in a were sectioned, based on their x position relative to the

cell center (normalized by cell radius) at a bin size of 0.25 (8 sections in total from -1 to 1), and

calculated the average wave area in each section. **c.** Changes in actin waves' spatial distribution

in response to EF reversal; data from 6 independent experiments. The color of each plot is coded

according to the timeline displayed at the bottom of the panel. P_2-P_5: The EF was reversed, and

cells gradually developed polarization towards the new cathode. The number of waves in each

plot: N_{p2} = 272, N_{p3} = 277, N_{p4} = 193, N_{p5} = 246. **d.** A schematic illustrating the old and new

fronts of giant cells when the EF was reversed. **e.** Time stacks of orientation distributions of

optical-flow vectors at an old front and a new front. The EF was reversed from the cathode being

at the right (0) to the cathode being at the left (π) at 0 min. **f.** Comparisons of response time

between new fronts (green) and old fronts (orange) from multiple experiments (N_{cell} = 5). The *P*-

value was calculated using a pairwise t-test at the 5% significance level. **g.** Cartoon illustrating

different time scales of local wave propagation and global rearrangement of STEN-CEN

thresholds, in response to EF reversal.

<u>Figure 4 – source data 1. Related to Fig. 4a-c</u>

Discussion

By employing giant cells, in which the cortical waves are disentangled from cell motion, we demonstrate that EFs modulate cortical wave dynamics directly, providing a mechanism for cell guidance by EFs (Fig. 1 c, b, d). Our use of nanoridges to generate quasi-1D waves that are small, short-lived, and unable to turn (Fig. 1c) enabled detailed quantification of wave properties, demonstrating that EFs directly affect the abundance, locations, and directions of cortical waves.

EFs guide cortical wave dynamics

Previous studies have suggested that the basal cortical waves in *D. discoideum* are insensitive to external chemotactic gradients, whereas "pseudopods" at other regions in the same cells can be guided (Lange et al., 2016). This conclusion is surprising because the biochemical events traveling with the waves are the same as those occurring on pseudopods, and pseudopods with the dorsal cups on the same cells do respond to chemoattractants. Also, similar cortical waves in human mammary epithelial cells can be guided effectively by epidermal growth factors (Zhan et al., 2020). Additional input from the greater contact of giant *D. discoideum* cells with the surface may outweigh the effect of applied chemical gradients on the basal waves. Other studies have shown that single cells can integrate combinations of external chemical and mechanical stimuli.

Our work shows that in giant cells, waves of both F-actin polymerization (Fig. 3) and its upstream regulator PIP3 (Fig. 1 – figure supplement 1) are indeed guided by EFs. These biased biochemical and biomechanical events lead to more protrusions at the cell front than at the cell back, thus driving cell migration (Fig. 2 – figure supplement 2). The development of the biased

wave activities takes ~10 min following the introduction of an EF (Fig. 2a and Fig. 3), which is

much slower than the timescale of surface-receptor-regulated chemotaxis. The high resistance of

the cell membrane limits the effects of EFs on intracellular components, but EFs may act on the

charged lipids and molecular clusters. Thus, we suspect that the slow response results from the

electrophoresis of the charged membrane components involved in wave formation, which has a

characteristic time scale of 5 to 10 min (Allen et al., 2013; McLaughlin & Poo, 1981).

We further explored the dynamics in response to EF reversal at the subcellular level using

nanotopography (Fig 4). We observed that the new waves are induced to propagate towards the

current cathode within 2 to 3 min (Fig. 4e and Fig.4 – figure supplement 2), suggesting that

waves themselves can adapt quickly to the changing electrical environments. Because we only

observed the fast adaptation on ridged surfaces, this phenomenon may be related to the shorter

wave lifetimes on nanoridges than on flat surfaces. A short lifetime allows waves to be nucleated

at a higher rate on the nanoridges, leading to a rapid directional response. During this process,

the EF may regulate the wave nucleation through locally changing specific charged lipids, ion

fluxes, or local pH gradients (Crevenna et al., 2013; Frantz et al., 2008; Köhler et al., 2012;

Martin et al., 2011; Zhou & Pang, 2018).

EFs modulate the thresholds of the excitable wave system

Recent studies have shown that the cortical wave system can be described as a coupled signal

transduction and cytoskeletal excitable network. Based on both simulation and experimental

studies (Bhattacharya et al., 2020; Miao et al., 2017), it has been shown that the wave ranges,

durations, and speeds are determined by the local threshold of activation, which in turn are

regulated by the relative levels of activators and inhibitors (Miao et al., 2017, 2019).

Our quantification shows that guided waves become larger, faster, and more persistent in an EF

(Fig. 2), indicating that the excitable system is closer to its threshold for activation (Miao et al.,

2019). This effect may arise from enhanced positive feedback, reduced negative feedback, or

both. We further find that wave nucleation is enhanced at the cell front and suppressed at the

back (Fig. 4a, b). This subcellular inhomogeneity is consistent with a biased excitable network

framework (Iglesias & Devreotes, 2012; Meinhardt, 1999; Tang et al., 2014; Xiong et al., 2010),

which was added to the STEN-CEN model to introduce an internal spatial gradient in the local

threshold of wave initiation, akin to cell polarity.

Local excitation and global inhibition (Xiong et al., 2010), LEGI, schemes have effectively

recreated the features of both fast directional sensing and stable polarity in response to chemical

signals, which can lead to robust biased excitable network. Both directional sensing and stable

polarity can lead to a robust biased excitable network. For chemical signals, the directional

response from PIP3 occurs within seconds, whereas the establishment of stable polarity usually

requires many minutes. However, based on our analysis, establishing both directional response

(Fig. 3) and polarity (Fig. 4) in response to EFs requires 5 to 10 min. It is worth noting that PIP3

waves also sense EFs on a time scale of minutes (Fig. 1 – figure supplement 1). Our observation

suggests that EFs act on the polarity establishment rather than directional sensing. This

hypothesis is supported by a recent study showing that G-protein-coupled receptors (GPCRs),

which are the regulator in the LEGI model for *D. discoideum* that allows for sensing

chemoattractant on timescales of seconds, are not essential for electrotaxis (Zhao et al., 2002).

EFs act on waves, and waves determine cell behaviors

Our results raise the possibility that cortical wave dynamics are modulated directly by EFs and

that the waves in turn mediate cellular response. Waves travel across cell membranes to

coordinate the trailing edge with the front edge, and the cytoskeletal components in cortical

waves are involved in developing the stable polarity. On the other hand, the duration and turning

capacity of STEN-CEN waves directly impact the speed and characteristics of the cellular

response to EFs (Fig. 3) on a longer timescale than that of surface-receptor-regulated chemotaxis.

Our results shed light on how EFs modulate protrusions. Previous studies have shown that

various protrusions that drive cell motion, such as filopodia, lamellipodia (Miao et al., 2019), and

macropinocytotic cups (Video 4), are always associated with expanding waves near cell

perimeter. Our previous work has shown that changing wave properties by perturbing STEN-

CEN states leads to the transition of protrusion profiles, which indicates that wave properties

dictate the properties of the protrusions (Miao et al., 2019). Here we showed that EFs can alter

the waves differently on the two ends of the cell (Fig. 4a). As a result of these spatially

inhomogeneous wave properties, protrusions become more abundant and larger on one side of

the cell versus the other, which eventually leads to guidance of cell migration.

On flat surfaces, a slow U-turn is observed following EF reversal, whereas on nanoridges, faster

switching is observed. Thus, the response of migrating cells to a changing guidance cue can be

predicted from the characteristics of the waves driving the migration process. Indeed, the U-turn

behaviors of neutrophils and differentiated, single *D. discoideum* cells in response to EF reversal

(Hind et al., 2016; Sato et al., 2007; Srinivasan et al., 2003; Xu et al., 2003), which are usually

ascribed to stable cell polarity, may instead reflect the persistence and 2D turning behavior of

cortical waves in these environments (Fig. 3).

Nanoridges allow us to shed further light on the multiscale character of the system, because cells

include both short, 1D waves on the basal plane, and longer-lasting, 2D waves on the dorsal

plane. The different response times on the subcellular level due to different wave behaviors (Fig.

4 – figure supplement 2) provide strong evidence that cortical waves act as direct mediators of

EFs. Waves on different planes are similar in composition but are impacted differently by the EF.

We observed fast switching of wave directions in the basal plane near the ridged substrate and

slower turning of the waves in the dorsal plane within the same cell, indicating that the direction

of waves is controlled locally by external cues (Fig. 3 – figure supplement 1).

EFs provide a means to modulate cortical waves directly. On the other hand, biological

conditions that modulate wave characteristics may also speed up or suppress the cellular

response to directional cues. Longer-lasting waves offer persistence in the face of rapidly

changing gradients, whereas shorter waves yield faster adaptability to changing directional

signals. The durations of waves and their ability to turn together have a dominant effect on the

response of cells to an EF.

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Cell line *(D. discoideum)*	Aca null	https://doi.org/10.1016/S0092-8674(03)00081-3		The cell line was a gift from Carole A. Parent lab.
Cell line *(D. discoideum)*	PHcrac-GFP LimE-RFP	https://doi.org/10.1038/ncb3495		The cell line was a gift from Peter N. Devreotes lab.
software, algorithm	Optical flow analysis (run by MATLAB)	https://doi.org/10.1091/mbc.E19-11-0614		

Materials and Methods

<u>Cell line</u>

In the study, we used LimE-RFP aca null Dictyostelium discoideum (D.d.) and PHcrac-

GFP/LimE-RFP D.d cell lines. LimE-RFP aca null was a gift from Carole A. Parent lab

(https://doi.org/10.1016/S0092-8674(03)00081-3), and PHcrac-GFP-LimE-RFP was a gift from

Peter N. Devreotes lab (https://doi.org/10.1038/ncb3495). We have conducted the mycoplasma

contamination testing for both cell lines and did not detected contamination.

<u>Cell culture</u>

Dictyostelium discoideum cell lines were grown axenically in the HL5 medium. Aggregation

adenylyl cyclase null (ACA−) mutants, which do not produce cAMP and do not have chemotaxis

signal relay (Kriebel et al., 2003), were used in electrotaxis experiments to avoid chemotaxis.

The cells used also express limE-RFP as a reference for filamentous actin structures. G418 was

used as the selection medium during cell culture. For the experiments in Fig. 1/Fig.1 -

supplement figure 1, we used *Dictyostelium discoideum* co-expressing PHcrac-GFP and LimE-

RFP, and we used G418 as the selective medium. Note that an enhancement in LimE

concentration is associated with protrusions, as seen in fig. 2 -figure supplement 2, and that the

protrusions are biased to the side facing the cathode. Because protrusions are driven by F-actin

polymerization, we believe this observation rules out the possibility that LimE binding/unbinding

to/from F-actin itself is sensitive to EFs.

<u>Electrofusion</u>

Cells were washed twice with 17 mM Sorensen buffer (15 mM KH_2PO_4 and 2 mM Na_2HPO_4,

pH 6.0) and rolled for 30 min at a concentration of 1.5×10^7 mL^{-1}. Electrofusion was conducted

with a Gene Pulser Gen1 system. Three pulses of 1 kV at a 1 s interval were applied. After

electroporation, cells were relaxed for 5 min. Then cells were diluted to 5×10^5 mL^{-1} with normal

developing buffer (5 mM KH_2PO_4, 5 mM Na_2HPO_4, 2 mM $MgCl_2$ and 0.2 mM $CaCl_2$, PH 6.5)

and seeded into a customized electrotactic chamber, with dimensions 20 mm $\times$ 5 mm $\times$ 0.25 mm.

The aca null cell line that we used did not generate many waves in the vegetative stage, and

electro-fusion with 1 kV pulses stressed the cells. Thus cells were starved for 2 h before

experiments to generate more actin waves.

<u>Nanotopography</u> <u>fabrication</u>

The nanotopographic pattern used in these cell studies was fabricated through a technique known

as multiphoton absorption polymerization (MAP), as described elsewhere (C. M. LaFratta et al.,

2006; C. N. LaFratta et al., 2004). An ultrafast, pulsed laser beam (Coherent Mira 900-F, 76

MHz) was passed through a high-numerical-aperture microscope objective onto a

photopolymerizable resin, sandwiched between glass coverslips. A LabVIEW (National

Instruments) program allowed for control of the stage position and the shutter state, determining

where polymerization occurred (and did not) in the resin and allowing patterning. Once

fabrication was completed, the patterned sample was developed in ethanol twice for 3 min each

to remove unreacted monomer. The polymerized structure was baked at 110 °C for at least 1 h.

To produce the necessary number of replicate patterns with the same dimensions, an adapted

version of replica molding was performed (Sun et al., 2018). A hard polydimethylsiloxane (*h*-

PDMS) film containing hexanes to increase the resolution of feature replication was spin-coated

onto the functionalized structure made from MAP. The film was allowed to sit on the structure

for 2 h at room temperature and was then baked at 60 °C for 1 h. Regular PDMS (Sylgard 184)

was prepared at a 10:1 ratio of elastomer base to the curing agent by degassing and mixing. The

PDMS was poured onto the *h*-PDMS film, and molding was completed by baking at 60 °C for an

additional 70 min. The final mold was peeled from the glass slide supporting the MAP-patterned

structure.

The mold was used to produce replicas of the original pattern. A drop of the same acrylic resin

was placed on the patterned area of the PDMS mold, and then an acrylate-functionalized glass

coverslip was pressed firmly on top, spreading the sandwiched drop. Tape secured this system in

place. The resin was cured for a total of 5 min under a UV lamp (Blak-ray), producing a polymer

film. It should be noted that the PDMS mold is the negative relief pattern of the structure made

using MAP. Therefore, samples (or replicas) of the original pattern could be produced on a

relatively large scale with this method. The replicas were soaked in ethanol for at least 12 h

before use in the cell studies. We fabricated flat surface samples by using a PDMS mold with a

smooth surface.

Lattice light-sheet microscopy

The 3i lattice light-sheet microscope in the Johns Hopkins School of Medicine Microscope

Facility was used for 2-color, 3D imaging. Vegetative, single *Dictyostelium* cells were seeded on

a circular 5 mm coverslip patterned with nanoridges, which was immersed in a bath of standard

developing buffer throughout imaging.

Electrotaxis experiments

We 3D-printed electrotaxis chambers (Fig. 2 – figure supplement 1) with dimensions of 20 mm $\times$

5 mm $\times$ 0.25 mm and composed of a clear resin using a Formlabs Form2 3D-printer. Agar

bridges were used to isolate cell media from electrodes to minimize electrochemical products

and pH changes. 20 V/cm constant EFs were applied. Time-lapse images of the phase-contrast

channel and the RFP/GFP channel were recorded using PerkinElmer spinning-disk microscope at

a frame rate of 0.1 frames/s (Yokogawa CSU-X1 spinning-disk scan head (5000 rpm)) with

Hamamatsu EMCCD camera and Volocity analysis software.

<u>Optical-flow analysis and model fitting of actin polymerization dynamics</u>

We applied the Lukas-Kanade optical-flow method to quantify the direction of the intensity flow

in fluorescence videos. This algorithm produced pixel-basis vector fields of intensity motion.

Before applying the optical-flow algorithm, each image was smoothed by a 2D Gaussian filter (σ

$= 3$) to reduce noise. After the smoothing, we further removed the flow vectors created by noise

using optical-flow reliability as our criterion. The reliability is defined as the smallest

eigenvalues of the $A^T wA$ matrix, where w is a Gaussian weight matrix and A is the intensity

gradient matrix. The size of the weight matrix for *D. discoideum* was set at 19×19, with

standard deviation $\sigma = 2$ (0.42 µm).

We built a bimodal von Mises model to compare the actin and cellular responses accurately. A

von Mises distribution is given by

$$f_{VM}(\theta|\mu,\kappa) = \frac{e^{\kappa \cos(\theta-\mu)}}{2\pi I_0(\kappa)} \ . \tag{2}$$

534   where the peak location is μ and the concentration κ. The orientation distribution of optical-flow

535   vectors at each time point is fit with two von Mises distributions

$$f(\theta|\mu_1,\mu_2,\kappa,p_1,p_2) = p_1 f_{VM}(\theta|\mu_1,\kappa) + p_2 f_{VM}(\theta|\mu_1+\pi,\kappa) \tag{3}$$

537   where p is the proportion of each component. We use the constraints

$$\mu_1 - \mu_2 = \pi \tag{4}$$

and

$$p_1 + p_2 = 1 \, . \qquad\qquad (5)$$

Maximum likelihood estimation (MLE) is applied to estimate model parameters based on the

orientation of all the optical-flow vectors every 12 frames (2 min). With this model, we can

quantitatively study the temporal change of actin dynamics. The preferential direction is defined

as the μ with the largest proportion p at each time point.

<u>Quantification of actin wave properties</u>

The segmentation of actin waves was conducted based on the combined information from

fluorescence intensities and optical flow. We first applied the kmeans ($k = 3$) cluster (Kanungo et

al., 2002) to pick up the bright regions in the limE-RFP videos, then only kept the moving

objects by applying the reliability mask from the optical-flow analysis.

To classify the large actin structures composed of substructures moving independently, we

considered the optical flow at the edges of the large structure (Fig. 2c). If the optical-flow vectors

were moving in the same direction at both edges, the large structure was classified as a single

wave. Otherwise, the large structure was divided into multiple smaller patches. In the latter case,

a pronounced boundary was detectable between two substructures, then we used the detected

boundary to divide the large structure into multiple substructures.

After classification, we measured properties such as wave speed, wave duration, and wave area

to characterize STEN-CEN. Wave speed was measured by tracking the clusters of optical-flow

vectors oriented in similar directions. The detailed algorithm can be found in a prior publication

(Lee et al., 2020). To measure wave duration, we first tracked actin waves using a customized,

multi-object tracking tool based on the overlapping areas between frames. A unique

identification number was assigned to each wave, then wave duration, wave area (measured by

the Matlab function regionprops) were recorded for each wave.

Acknowledgments: We thank Q. Qing and M. Zhao for the discussion. **Competing interests:**

The authors declare that they have no competing interests.

References:

Allen, G. M., Mogilner, A., & Theriot, J. A. (2013). Electrophoresis of cellular membrane

components creates the directional cue guiding keratocyte galvanotaxis. *Current Biology*,

23(7), 560–568. https://doi.org/10.1016/j.cub.2013.02.047

Bhattacharya, S., Banerjee, T., Miao, Y., Zhan, H., Devreotes, P. N., & Iglesias, P. A. (2020).

Traveling and standing waves mediate pattern formation in cellular protrusions. *Science

Advances*, *6*(32), eaay7682. https://doi.org/10.1126/sciadv.aay7682

Bhattacharya, S., & Iglesias, P. A. (2018). Controlling excitable wave behaviors through the

tuning of three parameters. *Biological Cybernetics*, *113*(1), 61–70.

https://doi.org/10.1007/s00422-018-0771-0

Bhattacharya, S., Miao, Y., Devreotes, P. N., & Iglesias, P. A. (2019). A Coupled Excitable

Network Model Dictates Cortical Wave Patterns and Controls Cellular Protrusion

Morphology. *Biophysical Journal*, *116*(3), 121a. https://doi.org/10.1016/j.bpj.2018.11.675

Bretschneider, T., Anderson, K., Ecke, M., Müller-Taubenberger, A., Schroth-Diez, B.,

Ishikawa-Ankerhold, H. C., & Gerisch, G. (2009). The three-dimensional dynamics of actin

waves, a model of cytoskeletal self-organization. *Biophysical Journal*, *96*(7), 2888–2900.

585 https://doi.org/10.1016/j.bpj.2008.12.3942

586 Cortese, B., Palamà, I. E., D'Amone, S., & Gigli, G. (2014). Influence of electrotaxis on cell

587  behaviour. *Integrative Biology (United Kingdom)*, *6*(9), 817–830.

588  https://doi.org/10.1039/c4ib00142g

589 Crevenna, A. H., Naredi-Rainer, N., Schonichen, A., Dzubiella, J., Barber, D. L., Lamb, D. C., &

590  Wedlich-Söldner, R. (2013). Electrostatics control actin filament nucleation and elongation

591  kinetics. *Journal of Biological Chemistry*, *288*(17), 12102–12113.

592  https://doi.org/10.1074/jbc.M113.456327

593 Devreotes, P. N., Bhattacharya, S., Edwards, M., Iglesias, P. A., Lampert, T., & Miao, Y. (2017).

594  Excitable Signal Transduction Networks in Directed Cell Migration. *Annual Review of Cell*

595  *and Developmental Biology*, *33*(1), annurev-cellbio-100616-060739.

596  https://doi.org/10.1146/annurev-cellbio-100616-060739

597 Driscoll, M. K., Sun, X., Guven, C., Fourkas, J. T., & Losert, W. (2014). Cellular contact

598  guidance through dynamic sensing of nanotopography. *ACS Nano*, *8*(4), 3546–3555.

599  https://doi.org/10.1021/nn406637c

600 Ermentrout, G. B., Hastings, S. P., & Troy, W. C. (1984). Large Amplitude Stationary Waves in

601  an Excitable Lateral-Inhibitory Medium. *SIAM Journal on Applied Mathematics*, *44*(6),

602  1133–1149. https://doi.org/10.1137/0144081

603 Flemming, S., Font, F., Alonso, S., & Beta, C. (2020). How cortical waves drive fission of motile

604  cells. *Proceedings of the National Academy of Sciences of the United States of America*,

605  *117*(12), 6330–6338. https://doi.org/10.1073/pnas.1912428117

606 Frantz, C., Barreiro, G., Dominguez, L., Chen, X., Eddy, R., Condeelis, J., Kelly, M. J. S.,

607  Jacobson, M. P., & Barber, D. L. (2008). Cofilin is a pH sensor for actin free barbed end

formation: role of phosphoinositide binding. *Journal of Cell Biology*, *183*(5), 865–879.

https://doi.org/10.1083/jcb.200804161

Gerhardt, M., Ecke, M., Walz, M., Stengl, A., Beta, C., & Gerisch, G. (2014). Actin and PIP3

waves in giant cells reveal the inherent length scale of an excited state. *Journal of Cell*

Science, *127*(20), 4507–4517. https://doi.org/10.1242/jcs.156000

Gerisch, G. (2010). Self-organizing actin waves that simulate phagocytic cup structures. *PMC*

Biophysics, *3*(1), 1–9. https://doi.org/10.1186/1757-5036-3-7

Harvath, L., Robbins, D., Russell, A. A., & Seamon, K. B. (1991). Elevation of cAMP

differentially affects chemotactic responsiveness . *The Journal of Immunology*, *146 (1)*

22(Vol. 146, Issue 1).

Hind, L. E., Vincent, W. J. B., & Huttenlocher, A. (2016). Leading from the Back: The Role of

the Uropod in Neutrophil Polarization and Migration. *Developmental Cell*, *38*(2), 161–169.

https://doi.org/10.1016/j.devcel.2016.06.031

Iglesias, P. A., & Devreotes, P. N. (2012). Biased excitable networks: How cells direct motion in

response to gradients. *Current Opinion in Cell Biology*, *24*(2), 245–253.

https://doi.org/10.1016/j.ceb.2011.11.009

Kanungo, T., Mount, D. M., Netanyahu, N. S., Piatko, C. D., Silverman, R., & Wu, A. Y. (2002).

An efficient k-means clustering algorithms: Analysis and implementation. *IEEE*

Transactions on Pattern Analysis and Machine Intelligence, *24*(7), 881–892.

https://doi.org/10.1109/TPAMI.2002.1017616

Ketchum, C. M., Sun, X., Suberi, A., Fourkas, J. T., Song, W., & Upadhyaya, A. (2018).

Subcellular topography modulates actin dynamics and signaling in B-cells. *Molecular*

Biology of the Cell, *29*(14), 1732–1742. https://doi.org/10.1091/mbc.E17-06-0422

Köhler, S., Schmoller, K. M., Crevenna, A. H., & Bausch, A. R. (2012). Regulating Contractility of the Actomyosin Cytoskeleton by pH. *Cell Reports*, *2*(3), 433–439. https://doi.org/10.1016/j.celrep.2012.08.014

Kriebel, P. W., Barr, V. A., & Parent, C. A. (2003). Adenylyl cyclase localization regulates streaming during chemotaxis. *Cell*, *112*(4), 549–560. https://doi.org/10.1016/S0092-8674(03)00081-3

LaFratta, C. M., Li, L., & Fourkas, J. T. (2006). Soft-lithographic replication of 3D microstructures with closed loops. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(23), 8589–8594. https://doi.org/10.1073/pnas.0603247103

LaFratta, C. N., Baldacchini, T., Farrer, R. A., Fourkas, J. T., Teich, M. C., Saleh, B. E. A., & Naughton, M. J. (2004). Replication of two-photon-polymerized structures with extremely high aspect ratios and large overhangs. *Journal of Physical Chemistry B*, *108*(31), 11256–11258. https://doi.org/10.1021/jp048525r

Lange, M., Prassler, J., Ecke, M., Müller-Taubenberger, A., & Gerisch, G. (2016). Local Ras activation, PTEN pattern, and global actin flow in the chemotactic responses of oversized cells. *Journal of Cell Science*, *129*(18), 3462–3472. https://doi.org/10.1242/jcs.191148

Lee, R. M., Campanello, L., Hourwitz, M. J., Alvarez, P., Omidvar, A., Fourkas, J. T., & Losert, W. (2020). Quantifying topography-guided actin dynamics across scales using optical flow. *Molecular Biology of the Cell*, *31*(16), 1753–1764. https://doi.org/10.1091/mbc.e19-11-0614

Lin, F., Baldessari, F., Gyenge, C. C., Sato, T., Chambers, R. D., Santiago, J. G., & Butcher, E. C. (2008). Lymphocyte Electrotaxis In Vitro and In Vivo. *The Journal of Immunology*, *181*(4), 2465–2471. https://doi.org/10.4049/jimmunol.181.4.2465

Martin, C., Pedersen, S. F., Schwab, A., & Stock, C. (2011). Intracellular pH gradients in

migrating cells. *American Journal of Physiology - Cell Physiology*, *300*(3), 490–495.

https://doi.org/10.1152/ajpcell.00280.2010

McLaughlin, S., & Poo, M. M. (1981). The role of electro-osmosis in the electric-field-induced

movement of charged macromolecules on the surfaces of cells. *Biophysical Journal*, *34*(1),

85–93. https://doi.org/10.1016/S0006-3495(81)84838-2

Meinhardt, H. (1999). Orientation of chemotactic cells and growth cones: Models and

mechanisms. *Journal of Cell Science*, *112*(17), 2867–2874.

https://doi.org/https://doi.org/10.1242/jcs.112.17.2867

Miao, Y., Bhattacharya, S., Banerjee, T., Abubaker☐Sharif, B., Long, Y., Inoue, T., Iglesias, P.

A., & Devreotes, P. N. (2019). Wave patterns organize cellular protrusions and control

cortical dynamics. *Molecular Systems Biology*, *15*(3), 1–20.

https://doi.org/10.15252/msb.20188585

Miao, Y., Bhattacharya, S., Edwards, M., Cai, H., Inoue, T., Iglesias, P. A., & Devreotes, P. N.

(2017). Altering the threshold of an excitable signal transduction network changes cell

migratory modes. *Nature Cell Biology*, *19*(4), 329–340. https://doi.org/10.1038/ncb3495

Neumann, E., Gerisch, G., & Opatz, K. (1980). Cell fusion induced by high electric impulses

applied to Dictyostelium. *Naturwissenschaften*, *67*(8), 414–415.

https://doi.org/10.1007/BF00405493

Sato, M. J., Kuwayama, H., Van Egmond, W. N., Takayama, A. L. K., Takagi, H., Van Haastert,

P. J. M., Yanagida, T., & Ueda, M. (2009). Switching direction in electric-signal-induced

cell migration by cyclic guanosine monophosphate and phosphatidylinositol signaling.

Proceedings of the National Academy of Sciences of the United States of America, *106*(16),

6667–6672. https://doi.org/10.1073/pnas.0809974106

Sato, M. J., Ueda, M., Takagi, H., Watanabe, T. M., Yanagida, T., & Ueda, M. (2007). Input-

output relationship in galvanotactic response of Dictyostelium cells. *BioSystems*, *88*(3),

261–272. https://doi.org/10.1016/j.biosystems.2006.06.008

Srinivasan, S., Wang, F., Glavas, S., Ott, A., Hofmann, F., Aktories, K., Kalman, D., & Bourne,

H. R. (2003). Rac and Cdc42 play distinct roles in regulating PI(3,4,5)P3 and polarity

during neutrophil chemotaxis. *Journal of Cell Biology*, *160*(3), 375–385.

https://doi.org/10.1083/jcb.200208179

Sun, X., Hourwitz, M. J., Baker, E. M., Schmidt, B. U. S., Losert, W., & Fourkas, J. T. (2018).

Replication of biocompatible, nanotopographic surfaces. *Scientific Reports*, *8*(1), 1–9.

https://doi.org/10.1038/s41598-017-19008-z

Tang, M., Wang, M., Shi, C., Iglesias, P. A., Devreotes, P. N., & Huang, C. H. (2014).

Evolutionarily conserved coupling of adaptive and excitable networks mediates eukaryotic

chemotaxis. *Nature Communications*, *5*, 1–13. https://doi.org/10.1038/ncomms6175

Veltman, D. M., Williams, T. D., Bloomfield, G., Chen, B. C., Betzig, E., Insall, R. H., & Kay, R.

R. (2016). A plasma membrane template for macropinocytic cups. *ELife*,

5(DECEMBER2016), 24. https://doi.org/10.7554/eLife.20085

Wang, M. J., Artemenko, Y., Cai, W. J., Iglesias, P. A., & Devreotes, P. N. (2014). The

directional response of chemotactic cells depends on a balance between cytoskeletal

architecture and the external gradient. *Cell Reports*, *9*(3), 1110–1121.

https://doi.org/10.1016/j.celrep.2014.09.047

Weiner, O. D., Marganski, W. A., Wu, L. F., Altschuler, S. J., & Kirschner, M. W. (2007). An

actin-based wave generator organizes cell motility. *PLoS Biology*, *5*(9), 2053–2063.

https://doi.org/10.1371/journal.pbio.0050221

Xiong, Y., Huang, C. H., Iglesias, P. A., & Devreotes, P. N. (2010). Cells navigate with a local-

excitation, global-inhibition-biased excitable network. *Proceedings of the National*

Academy of Sciences of the United States of America, *107*(40), 17079–17086.

https://doi.org/10.1073/pnas.1011271107

Xu, J., Wang, F., Van Keymeulen, A., Herzmark, P., Straight, A., Kelly, K., Takuwa, Y.,

Sugimoto, N., Mitchison, T., & Bourne, H. R. (2003). Divergent signals and cytoskeletal

assemblies regulate self-organizing polarity in neutrophils. *Cell*, *114*(2), 201–214.

https://doi.org/10.1016/S0092-8674(03)00555-5

Zhan, H., Bhattacharya, S., Cai, H., Iglesias, P. A., Huang, C. H., & Devreotes, P. N. (2020). An

Excitable Ras/PI3K/ERK Signaling Network Controls Migration and Oncogenic

Transformation in Epithelial Cells. *Developmental Cell*, *54*(5), 608-623.e5.

https://doi.org/10.1016/j.devcel.2020.08.001

Zhao, M., Jin, T., McCaig, C. D., Forrester, J. V, & Devreotes, P. N. (2002). Genetic analysis of

the role of G protein-coupled receptor signaling in electrotaxis. *The Journal of Cell Biology*,

157(6), 921–927. https://doi.org/10.1083/jcb.200112070

Zhao, M., Song, B., Pu, J., Wada, T., Reid, B., Tai, G., Wang, F., Guo, A., Walczysko, P., Gu, Y.,

Sasaki, T., Suzuki, A., Forrester, J. V., Bourne, H. R., Devreotes, P. N., McCaig, C. D., &

Penninger, J. M. (2006). Electrical signals control wound healing through

phosphatidylinositol-3-OH kinase-γ and PTEN. *Nature*, *442*(7101), 457–460.

https://doi.org/10.1038/nature04925

Zhou, H. X., & Pang, X. (2018). Electrostatic Interactions in Protein Structure, Folding, Binding,

and Condensation. *Chemical Reviews*, *118*(4), 1691–1741.

https://doi.org/10.1021/acs.chemrev.7b00305

Funding

Air Force Office of Scientific Research grant (FA9550-16-1-0052)

- Qixin Yang
- Yuchuan Miao
- Leonard J. Campanello
- Matt J. Hourwitz
- Bedri Abubaker-Sharif
- Abby L. Bull
- Peter N. Devreotes
- John T. Fourkas
- Wolfgang Losert

National Insititute of Health (T32 GM136577)

- Bedri Abubaker-Sharif

The funders had no role in study design, data collection, and interpretation, or the decision to submit the work for publication.

a

b

c

T1   T2   T3   T4   T5

Fig. 1 – figure supplement 1. Colocalization of PIP3 and F-actin in an EF. a. Images of

PHcrac (top row) and LimE (middle row) in a 20 V/cm EF with the cathode located on the left.

The EF was turned on at 0 min and reversed at 34 min. From the combined images (bottom row),

it is clear that F-actin and PIP3 remained coordinated during electrotaxis. **b**. Kymograph of

angular distribution of F-actin (top) / PIP3 (bottom) wave motion. Optical-flow analysis was

applied to both the F-actin and PIP3 videos to measure wave motion. Then we stacked up the

angular distribution at each time point along the x-axis. The dynamics of F-actin/PIP3 in

response to EFs are similar. The color was coded according to the proportion of each bin in the

angular distribution. **c.** Polar plots of angular distributions from the time points specified in b.

Fig. 2 – figure supplement 1. Schematic of the 3D-printed chamber used for electrotaxis

experiments. a. Top view of the experimental setup. **b.** Back view of the setup. Agar bridges

isolate the cell media from electrodes to avoid changes in pH and the generation of

electrochemical products. The cells were injected using pipettes into the center channel (red box

highlighted in a). The height of the electrotaxis channel is ~0.25 mm. The channel was sealed

with a large substrate coverslip and a smaller top coverslip.

Fig. 2 – figure supplement 2. EFs induce keratocyte-mode migration, producing larger

protrusions at cell fronts. **a**. Shape dynamics of a giant *D. discoideum* cell at 10 s intervals. Cell

boundaries were outlined using an active contour algorithm. Boundaries are color-coded

according to time. Four stages were categorized: (S0) Random motion in the absence of an EF.

(S1) Random motion in the first 15 min in the presence of an EF. (S2) Transition state usually

with a retraction at the back. (S3) Steady migration state in the presence of an EF. **b**. A

schematic of local protrusion and retraction, where the solid line is the current frame, and the

dotted line represents the cell boundary 1 min later. A protrusive region (yellow) is defined as

one occupied in the new frame, not in the previous frame, and a retractive region (blue) is

defined as one occupied in the previous frame, not in the new frame. **c**. Kymograph of local

boundary motion. The x-axis represents time, the y-axis indicates boundary points, and the color

of each pixel corresponds to the speed. **d**. Correlation curves of boundary motion at different

time points. **e**. Correlation length (defined as the point at which curves in d reach a value of 1/2)

vs. time. Each curve represents an independent experiment conducted on a different day.

Fig. 3 – figure supplement 1. Basal actin waves reverse direction on nanoridges, whereas

dorsal waves turn. a. A schematic showing the two imaging planes used, with the morphology

of the substrate. **b.** LimE-RFP images recorded at the dorsal plane. Unlike the basal focal plane

images, which capture the complete basal wave dynamics, the dorsal plane images do not capture

the full dorsal wave motion. To avoid photobleaching and laser damage, we only imaged the

cross-section of the dorsal waves and tracked the cross-sections using optical-flow analysis.

c. A schematic introducing 4-parameter, bimodal von Mises (vM) fitting. The orientation

distribution (gray) was fitted to a bimodal vM distribution at each time point. The two vM

distributions were set to share the degree of concentration κ ($\kappa_1 = \kappa_2$), peak locations μ that were

set to be   apart ($\mu_1 = \mu_2 + 180°$). **d.** Temporal evolution of fitting parameters in response to

EF reversal. Among two fitted vM distributions, the one with a larger proportion (vM$_p$) was

tracked. For both plots, the curves represent the mean p of vM$_p$ with standard derivations from

multiple experiments (   ), and the colors represent the μ of vM$_p$.

Fig. 3 – figure supplement 2. Basal and dorsal waves on flat surfaces both turn in response

to EF reversal. Temporal evolution of cosine function of the fitted preferential direction of actin

wave propagation for giant cells on flat surfaces. The analysis was based on $N = 3$ experiments.

Fig. 4 – figure supplement 1. The spatial inhomogeneity of wave properties could be caused

either by the **EF or by the external electrical potential gradient relative to the ground.** To

explore these scenarios, we quantified the waves in single cells surrounding the giant cell in the

field of view, where the center of the field was defined as the origin. In contrast to the giant cells,

these single cells are scattered throughout the field of view but are not large enough for the

potential gradient to create significant intracellular polarization. Thus, if the spatial

inhomogeneity is caused by the external electrical potential gradient, we would observe a

gradient of wave properties from single cells located in the region between -50 μm and 50 μm. **a.**

A limE image. Single cells are highlighted with blue circles. **b, c.** Density scatter plots of wave

location vs. wave area for single cells (highlighted by blue circles in a) in the absence (b) and the

presence (c) of EF. Unlike in giant cells (Fig. 4a, 4b), the wave areas in single cells are spatially

homogeneous. The ratio of mean wave areas in the regions nearer the cathode (location > 0) to

those in regions farther from the cathodes (location < 0) was calculated for both single cells and

giant cells. In giant cells, this ratio increases from 0.85 to 2.00 with an EF (Fig. 4a), whereas for

single cells, the ratio is almost unchanged (1.08 without an EF and 0.98 with an EF). This

analysis was based on the experiments from 4 different days.

Figure 4 – figure supplement 1- source data 1. Related to Fig. 4 – figure supplement 1.

Fig. 4 – figure supplement 2. Analysis of the propagation of individual waves. a. Illustration

of analysis of the propagation of individual waves. We first segmented each wave. For all of the

optical-flow vectors within the wave region, we normalized the magnitudes and then calculated

the *x* projection of their resultant vector   (——   where   is the orientation of

optical-flow vector . For waves propagating unidirectionally,   is close to ±1, whereas for

waves extending bidirectionally,   is close to 0. **b.** Violin plots of   of waves near old and

new fronts, at different time windows illustrated by the bottom timeline. The violin plots show

the distribution of   weighted by wave area, and the centered boxplot (gray bars) show the

without any weighting. The analysis was based on $N = 7$ experiments performed over 4 different

days. As the plots show, individual waves in the old fronts reversed their directions towards new

cathodes (both violin plot and boxplot shift their centers towards <0 region) after 9 min (   ).

While for waves in the new fronts, larger waves shifted their directions to new cathode within 6

min (violin plot shifts to < 0 at   ), and smaller wave response time is comparable to waves in

the old fronts (boxplot shifts to < 0 at   ).

Figure 4 – figure supplement 2- source data 1. Related to Fig. 4 – figure supplement 2b.

837