Synthetic Cooling Agents in US-marketed E-cigarette Refill Liquids and Popular Disposable E-cigarettes: Chemical Analysis and Risk Assessment.

Sairam V. Jabba, DVM, PhD [* 1,2] ; Hanno C. Erythropel, PhD [* 2,3] ; Deyri Garcia Torres [4] ; Lauren

A. Delgado [4] ; Jackson G. Woodrow [4] ; Paul T. Anastas [5] ; Julie B. Zimmerman PhD [2,3] ; and

Sven-Eric Jordt, PhD [1,2,6]

1: Department of Anesthesiology, Duke University School of Medicine, Durham, NC

2: Yale Center for the Study of Tobacco Products (YCSTP), Department of Psychiatry, Yale

School of Medicine, New Haven, CT

3: Department of Chemical and Environmental Engineering, Yale University, New Haven, CT;

4 Yale College, New Haven, CT

5: School of Public Health, Yale University, New Haven, CT;

6: Corresponding author

* These authors contributed equally

Address for Corresponding Author

     Sven-Eric Jordt, PhD (sven.jordt@duke.edu)

     Department of Anesthesiology

     Duke University School of Medicine

     3 Genome Ct.

     Durham, NC 27710-3094

Word count: ~4550 words

Abstract word count: 250

Citations: 59

Abstract:

Background: Menthol, through its cooling and pleasant sensory effects, facilitates smoking and tobacco product initiation, resulting in the high popularity of mint/menthol-flavored E-cigarettes. More recently, E-cigarette vendors started marketing synthetic cooling agents as additives that impart a cooling effect but lack a characteristic minty odor. Knowledge about content of synthetic coolants in US-marketed E-cigarette products and associated health risks is limited.

Methods: E-liquid vendor sites were searched with the terms "koolada", "kool/cool", "ice", or WS-3/WS-23, denoting individual cooling agents, and relevant refill E-liquids were purchased. "Ice" flavor varieties of Puffbar, the most popular disposable E-cigarette brand, were compared with non-"Ice" varieties. E-liquids were characterized, and synthetic coolants quantified using GC/MS. Margin of exposure (MOE), a risk assessment parameter, was calculated to assess the risk associated with synthetic coolant exposure from E-cigarette use.

Results: WS-3 was detected in 24/25 refill E-liquids analyzed. All Puffbar flavor varieties contained either WS-23 (13/14) or WS-3 (5/14), in both "Ice"- and non-"Ice" flavors. Modeling consumption of WS-3 from vaped E-liquids, resulted in MOEs below the safe margin of 100 for most daily use scenarios. MOEs for WS-23 were <100 for 10/13 Puffbar flavors in all use scenarios. Puffbar power specifications are identical to Juul devices.

Conclusions: Synthetic cooling agents (WS-3/WS-23) were present in US-marketed E-cigarettes, at levels that may result in consumer exposures exceeding safety thresholds set by regulatory agencies. Synthetic coolants are not only found in mint-or menthol-flavored products, but also in fruit- and candy-flavored products, including popular disposable E-cigarette products such as Puffbar.

Implications: Synthetic cooling agents are widely used in "kool/cool"- and "ice"-flavored

E-liquids and in E-liquids without these labels, both as a potential replacement for menthol or to

add cooling 'notes' to non-menthol flavors. These agents may be used to bypass current and

future regulatory limits on menthol content in tobacco products, and not just E-cigarettes. Since

synthetic cooling agents are odorless, they may not fall under the category of "characterizing

flavor", potentially circumventing regulatory measures based on this concept. Regulators need

to consider the additional health risks associated with exposure to synthetic cooling agents.

Introduction

Additives and flavors are added to tobacco products to counteract the harshness, sensory irritation and bitter tastes of tobacco and nicotine. In the United States, the Family Smoking Prevention and Tobacco Control Act (FSPTCA) of 2009 authorized the Food and Drug Administration (FDA) to regulate constituents, additives, and flavors in tobacco products [1]. The FSPTCA banned the sales of flavored combustible cigarettes, with the exception of menthol cigarettes [1]. Menthol in tobacco products is well understood to reduce harshness, impart a pleasant minty and cooling flavor and increase palatability [2]. Menthol affects tobacco and nicotine use behaviors such as initiation, dependence, abuse liability and reduces the ability to quit, especially among the youth and young adults [2]. The use of mentholated tobacco products is associated with tobacco use disparities, with 85% of African American smokers using menthol cigarettes [2,3]. The US FDA announced in May 2021 that it will issue a rule to ban menthol cigarettes [4]. Menthol is also a popular flavor additive in E-cigarettes, especially after the FDA banned all flavored closed pod systems (eg. Juul) except for menthol and tobacco [5]. Sales of menthol Juul pods increased 6-fold from August 2019 to May 2020 (10.7% to 61.8%) when all other Juul flavors (besides tobacco) were removed from the marketplace [6,7].

Synthetic cooling agents share the sensory cooling effects of menthol, but lack menthol's strong minty odor, especially at the concentrations at which they are added to E-liquids [8,9]. These agents were developed beginning in the 1970s, to replace menthol in body care and food products. In the 1970s and 80s, the tobacco industry including R.J. Reynolds and Phillip Morris, carried out consumer tests with cigarettes containing synthetic cooling agents, to impart a "cooling without menthol" experience [8–11]. A 2018 analysis of combustible cigarettes (both menthol and non-menthol) marketed in Germany detected WS-3, indicating that coolants other than menthol are added to tobacco products [12,13]. In a recent study we demonstrated that European-marketed Juul menthol pods contained the synthetic cooling agent WS-3, while US and Canadian pods did not [14]. More recently Omaiye et al., demonstrated the presence of synthetic coolants in US-marketed disposables [15]. While synthetic cooling agents are sold to (Do-It-Yourself) DIY users in the US, often under the label "Koolada", the identity and concentrations of these agents in US-marketed E-cigarettes and E-cigarette refill liquids ("E-liquids") remains unknown [16].

While natural and synthetic mint- and cooling compound are widely used in food and other consumer products, there are rising concerns about their toxicity, especially in poorly regulated products such as E-cigarettes. For example, we detected concerning levels of

pulegone, a carcinogenic mint flavorant banned by FDA in food, in a variety of mint- and

menthol-flavored E-cigarette and smokeless tobacco products [17,18]. Synthetic cooling agents

such as WS-3 are classified by the Flavor Extracts Manufacturers Associaton (FEMA) as GRAS

(Generally Recognized As Safe). However, this applies only to their intended use in food

products, but not to E-cigarette use resulting in inhalational exposure [19]. The Joint Expert

Committee on Food Additives (JECFA) of the World Health Organization (WHO) established a

threshold of concern of 90 µg/day/person for intake of WS-3 and related compounds, a level

that is often exceeded by consumers [20]. Based on oral toxicity studies, an oral No-Observed-

Adverse-Effect-Level (NOAEL) of 8 mg/kg-bw for WS-3 [20–22] was established, and 5 mg/kg-bw

for WS-23, another widely used synthetic cooling agent. Chronic or higher doses were observed

to cause kidney and liver lesions [20,23]. *In vitro* genotoxicity studies in mammalian cells

demonstrated WS-23 to be clastogenic (i.e., inducing disruption or breakages of chromosomes),

leading to a call by JECFA for additional toxicological studies to further evaluate the safety of

WS-23 [20]. Exposure of airway-epithelial cells in-vitro to various concentrations of synthetic

coolants and WS-23 containing e-liquids also demonstrated a dose-dependent cellular toxicity

[15]. A recent inhalation exposure study in rodents by an E-liquid manufacturer for WS-23

determined a NOAEL of 29 mg/kg-bw (342 mg/m3) in rodents [24]. Due to their evident toxicity

and increased use in E-cigarettes, it is critical to evaluate the potential health risks associated

with synthetic coolant exposure from E-cigarette use.

In the present study, we determined the levels of synthetic cooling agents (WS-3, WS-

23) as well as important peppermint- (menthol and menthone) and spearmint- (carvone) flavor

chemicals, in US-marketed refill E-liquids and in disposable E-cigarettes of the brand Puffbar.

The rationale to include menthol, menthone and carvone in chemical analysis is that, they are

the most frequent mint flavors encountered in E-liquids. Several recent E-cigarette prevalence

studies indicated high use of disposables, especially 'ice'-flavored, among youth and young

adults [6,7,25–29]. Disposable E-cigarette brand, Puffbar, was selected because, according to the

2021 US National Youth Tobacco Survey it was determined as the most popular E-cigarette

device among middle and high school students [6,7,28]. In addition, the power specifications for

Puffbar devices were determined. We then assessed the health risk associated with synthetic

coolant exposure via E-cigarette use by calculating the Margin of Exposure (MOE), a risk

assessment parameter, for a range of product use scenarios modeling low, moderate and

frequent E-cigarette use.

Materials and Methods:

Analysis of synthetic cooling agents and other flavor content in US-marketed E-cigarette liquids and their aerosols:

E-liquid selection

E-cigarette vendor websites keyword searches for "koolada", "kool", "WS-3", and "WS-23" were carried out to identify E-liquids potentially containing synthetic coolants. Identified products, including a concentrate ("Agent Cool") for do-it-yourself (DIY) users, were purchased directly from vendors in 2019 and 2020. In addition, previously purchased menthol/mint/spearmint/wintergreen E-liquids (2019) were re-analyzed for synthetic coolant presence. In early 2020, all available flavors of "Puffbar" were purchased from puffecig.com due to the frequent occurrence of the words "cool/ice" in Puffbar flavor names (see Table 1,2 for an overview). Purchased E-liquids were stored in dark at room temperature.

Power specifications measurement

Puffbar devices were opened manually, and the heating coil was carefully pulled towards the front of the opened device by the connecting wires. Resistance was measured in three devices by clamping a digital multimeter (Fluke 116) to the connecting wires. Battery specifications were printed onto the battery pack and reported as such.

E-liquid chemical analysis

E-liquids were analyzed using an established GC-MS method and selected compounds (WS-3, WS-23, menthol, menthone, carvone, nicotine) were quantified in triplicate using an established GC-FID method [30]. To do so, commercially available standards of N-Ethyl-p-menthane-3-carboxamide (WS-3; 99%; CAS# 39711-79-0), (S)-nicotine (>99%; CAS# 54-11-5), (DL)-menthol (99%; CAS# 89-78-1), (D)-carvone (>96%; CAS# 2244-16-8; all Sigma-Aldrich, Saint Louis, MO), 2-isopropyl-N,2,3-trimethylbutyramide, (WS-23; >98%; CAS# 51115-67-4, TCI, Portland, OR), (L)-menthone (97%; CAS# 14073-97-3; Alfa Aesar, Haverhill, MA), propylene glycol (>99.5%; CAS# 57-55-6), and glycerol (99.6%; CAS 56-81-5; both Fisher Scientific, Hampton, NH) were used to construct calibration curves in the relevant concentration range. Samples were diluted in 1mL of methanol [JT Baker, Center Valley, PA] containing 1 g/L of 1,4-dioxane [Alfa Aesar] as internal standard (IS) and injected; the detailed analytical methods have been described previously [30]. PG/VG ratios for Puffbars were rounded to the nearest 5 for simplicity.

Coolant carryover analysis

To assess the carryover of the synthetic coolants WS-3 and WS-23 from refill E-liquids to E-cigarette-generated aerosol, the Suorin iShare refillable pod E-cigarette device was used as its power specifications are very similar to the Puffbar and Juul device (Suorin iShare: 3.7V, 1.8Ω, resulting max. power 7.6W; Puffbar and Juul: 3.7V, 1.7Ω, resulting max. power 8.1W). Moreover, Suorin remains highly popular among vapers, and especially high school age vapers, as a favorite refillable device [29]. The Sourin device was attached via a custom-3D printed connector to a previously described house-built vaping machine [31]. In brief, the device was controlled by a programmable Arduino board that activates a micro-diaphragm pump at an operator-defined puff length, inter-puff break, and number of puffs. A microneedle valve in combination with a flow meter (Omega, Norwalk, CT) was used to control the flow rate. Generated aerosol was trapped by a series of liquid nitrogen-chilled cold-finger traps. The carryover puffing regime was 20 puffs, 2L/min flow rate, 2.4s puff length, 80mL puff volume, 30s inter-puff break and carryover was determined in triplicate (three separate cartridges). The chosen puffing regime is at the lower spectrum of observed user behavior [31,32] and has been utilized previously to determine carryover in popular Juul E-cigarettes [33]. After 20 puffs were collected for each experiment, the traps were allowed to thaw and the captured material was taken up in 1mL of methanol containing IS and injected into the GC/FID. Percent carryover was calculated as coolant content in trapped aerosol over coolant content of consumed liquid, as determined by weighing the e-liquid-containing cartridge before and after the carryover experiment using an analytical balance (Sartorius Practum, Göttingen, Germany). E-liquids for carryover experiments were chosen based on type of synthetic coolant present (WS-3, WS-23), and based on high, average, or low synthetic coolant and/or nicotine content (Table1).

Margin of Exposure (MOE) calculation for synthetic cooling agents:

MOE, a preferred risk assessment parameter for toxic compounds by the JECFA [34,35], was calculated for WS-3 and WS-23 containing E-liquids using the following formula, as described previously:[36]

$$MOE = \frac{\textbf{NOAEL for synthetic coolant}}{\textbf{Estimated Exposure Dose (EED)}}$$

Where NOAEL is the No-Observed-Adverse-Effect-Level, a dose determined as the threshold producing a toxic effect in animal oral toxicity studies. Though benchmark dose (BMD) is the preferred toxicological threshold for MOE calculations, NOAEL-values are applicable when available toxicity data is not amenable for statistical modeling of BMD [37], as was the case

for WS-3 and WS-23. JECFA determined NOAELs for WS-3 and WS-23 are 8 and 5 mg/kg-

bw/day, respectively [21–23]. Daily Estimated Exposure Doses were determined from WS-3 and

WS-23 concentrations in the E-liquids and their estimated average daily amount consumed.

Nicotine concentrations in the majority of refill E-liquids ranged between 0.1-1.0%, and user

surveys found consumption volumes for such low nicotine E-cigarette liquids to be 9 mL/day,

ranging from 1.7-17.7 mL/day) [18,38,39]. To cover a wide range of user behaviours, MOEs for

these refill E-liquids were calculated for consumption volumes of 1-,3-,5-,10- and 15-mL/day.

For nicotine salt-containing refill liquids (nicotine content >3%), MOEs were calculated for 1-,3-

,5-mL/day to account for the higher nicotine content. Puffbar E-liquid MOEs were calculated for

consumption amounts of half-,1- and 2-Puffbar(s) to model volumes consumed by light,

moderate, and heavy users of 3-5% nicotine salt-containing disposable E-cigarettes (Puffbar

volume is 1.3 mL). For the calculations, a user body weight of 60 kg was assumed; to convert

E-cigarette liquid consumption from volume in mL/day to g/day, the densities of propylene glycol

and glycerol (1.04 and 1.26 g/cm^3, respectively) were taken into consideration.

Results

Chemical analysis of synthetic cooling agents, nicotine and other flavorants in US-marketed E-cigarette refill liquids, concentrates and their aerosols:

Using our search criteria, 25 E-cigarette refill liquids and 1 concentrate from 19 different brands were identified and their contents of synthetic coolants, mint-related compounds and nicotine identified by GC/MS and quantified by GC/FID. Of the analyzed liquids, 21 contained free-base nicotine and 4 nicotine salts. WS-3 was present in 24/25 E-liquids, with amounts varying from 150 µg/g-14,650 µg/g (Table1). One E-liquid contained WS-23 exclusively (Dillinger;1,860 µg/g)(Table1). The concentrate contained both WS-3 (8,840 µg/g) and WS-23 (124,060 µg/g)(Table1).

Carryover rates for WS-3 from 8-different E-liquids into aerosol ranged between 91-109% (vs. neat E-liquid content) (average: 105±6.4%; n=20; Table1). For the only refill E-liquid it was detected in, the carryover rate for WS-23 was 99±10% (vs. neat E-liquid content; n=3; Table1).

In addition to WS-3 and WS-23, menthol, menthone, carvone, and nicotine were also quantified, which were detected in 13/25 E-liquids for menthol (330-23,490 µg/g), 6/25 for menthone (30-25,450 µg/g) and carvone (10-6830 µg/g), and 4/25 for menthol, menthone, carvone, WS-3 combined (4,940-46,150 µg/g; see Table1 for details). Nicotine concentrations in 20/21 free-base nicotine containing E-liquids were below 1% (1.41-9.97 mg/g), except for Clove+Cool Hit (17.91 mg/g). Nicotine contents in nicotine-salts containing E-liquids ranged from 28.74-44.6 mg/g (Table1). Carryover rates for menthol, menthone and carvone were calculated from vaping of select refill E-liquids and ranged from 103-107%, 85-106% and 97-105%, respectively (Supplementary Table1). Taken together, these results suggest that synthetic coolants are added to US-marketed E-liquids of a wide range of flavors (Table1), including to those already containing the "natural" coolant menthol. Indicated PG/VG ratios for refill E-liquids in Table 1 represent label information.

Chemical analysis of synthetic cooling agents and menthol in US-marketed disposable Puffbar E-cigarettes:

Several Puffbar flavors are marketed as 'Ice' flavors, indicating the presence of a cooling agent. WS-23 was detected in 13/14 Puffbar flavors (1,610-36,660 µg/g), WS-3 in 5/14 flavors (1,880-10,770 µg/g) and menthol in 7/14 analyzed flavors (620-3,680 µg/g; Table2). 3/14 Puffbar flavors contained all three compounds, with a combined concentrations of 8,830-48,050

μg/g (Table2). The nicotine concentrations in the Puffbar E-liquids ranged from 29.4-40.7 mg/g. As no PG/VG ratios were indicated on the product label, ratios were measured by GC/FID (Table 2).

Puffbar power specifications:

Contrary to pod-based E-cigarettes, the disposable puffbar device does not contain a separate "pod" or cartridge that can be exchanged, but the e-liquid is housed within the metal frame of the device, where it is stored in an absorbent material much like in the first generation "cig-a-likes" [31]. The coil resistance was measured at 1.7Ω and the battery indicated a maximum power output of 3.7V, which means the device has the exact power specifications as the Juul device [14], in addition to its look that also seem to mimic the Juul device. Maximum power output is thus calculated as 8.05W.

Margin of Exposure calculations for synthetic coolants in US-marketed refill E-liquids and disposable E-cigarettes:

To assess the risk associated with daily intake of WS-3 and WS-23 from E-cigarette use, we calculated the MOEs for different use scenarios (see methods for details). For organ toxicity, a safety margin MOE of >100 is considered low risk, while MOEs <100 require prioritization by regulatory agencies for risk mitigation for a given additive [37,40]

For the E-liquid with the highest WS-3 concentration (Arctic Rush), MOEs ranged from 28 (1-mL/day) to 2 (15-mL/day;Table S2 and Figure1), and are thereby significantly lower than the recommended safety margin of 100. For consumption volumes of 3-mL/day and 5-mL/day, 13/24 and 18/24 respectively, of WS-3 containing refill E-liquids had a MOE of <100. For 10- and 15-mL/day, MOEs were calculated only for the 20 free-base nicotine containing E-liquids: For 10-mL/day and 15-mL/day consumption volumes, 16/20 and 19/20 WS-3 containing E-liquids respectively, yielded a MOE <100 (Table S2 and Figure1). MOEs for the one WS-23-containing refill E-liquid was <100 in all assessed consumption scenarios, except for 1-mL/day consumption (Table S2 and Figure1). The tested concentrate ("Agent Cool") contained both WS-3 and WS-23. Based on vendor recommendations to add 1-2 drops of concentrate per pod, we conservatively estimated MOEs for a resulting dilution range of 1-3%. Resulting MOEs for WS-23 when added at 1-3% to E-liquids were <100 at all consumption rates, except for the lowest consumption rate (1-mL/day) at 1% dilution. Taken together, these results indicate that users of synthetic coolants flavored E-liquids are exposed to WS-3 and WS-23 levels that can potentially pose a health risk upon long-term use.

Next, we calculated the MOEs for synthetic coolants for daily consumption of Puffbar

devices. For the Puffbar flavor with the highest content of both WS-3 and WS-23 (Cool Mint),

the MOEs for consumption of 0.5-2 Puffbars/day ranged from 11-2 for WS-23 and 57-14 for

WS-3, values that are significantly lower than the safety margin threshold of 100 (Table S3 and

Figure2). MOEs for WS-3 were lower than the safety threshold of 100 at the highest estimated

consumption rate (2-Puffbars/day) for all five WS-3 containing Puffbar flavors. MOEs for WS-3

were also <100 for Cucumber and Menthol flavors at 1-Puffbar/day consumption rate. For all

other flavors, WS-23 MOEs were <100 for all use scenarios, with the exception of Café Latte

(0.5 Puffbar/day), Cucumber (0.5 Puffbar/day) and Menthol (0.5 and 1-Puffbar/day). These

results indicate that users of Puffbar disposable E-cigarettes are potentially exposed to levels of

synthetic coolants that may pose organ toxicity risks, especially from WS-23 exposure.

Discussion:

We detected the presence of synthetic coolants (WS-3 and WS-23) in US-marketed E-cigarette refill liquids and almost all tested flavors of the popular disposable E-cigarette brand, Puffbar, which was found to be the most popular device among US middle- and high school students in 2021 [28]. Using a relatively low-power E-cigarette that resembles the power specifications of Puffbar and the popular Juul device, we demonstrate that these coolants can efficiently be aerosolized and carried over to reach user airways. Margins of Exposure (MOE) were calculated to assess the health risks associated with exposure to these coolants from E-cigarette use, a parameter used by the US FDA and other global regulatory agencies to regulate the safe use of food additives with carcinogenic and organotoxic effects. These calculations suggests that daily intake of WS-3 or WS-23 by E-cigarette users exceed levels considered acceptable for consumption by FDA and the European Flavour Association (EFFA).

Natural cooling agents such as menthol provide a cooling sensation by activating cold-sensitive trigeminal nerve fibers innervating the upper respiratory tract and oral cavity. These sensory neurons express TRPM8 ion channels, the receptor for menthol that is also activated by cooling [41,42]. Menthol also has counterirritant, antitussive and analgesic properties and inhibits respiratory irritation responses to tobacco smoke and its various irritant constituents through a TRPM8 dependent mechanism in rodent models [43,44]. Menthol, through its actions on TRPM8, attenuated the aversive oral effects of nicotine in mice and increased nicotine intake [45]. The half-maximal concentrations of menthol to activate human TRPM8 is 4µM [46,47]. This concentration is known to be exceeded after inhalation of menthol cigarette smoke [48]. A menthol concentration of 0.65µM (16 ppm), was sufficient to reduce sensory irritation by cigarette smoke exposure in mice, illustrating the strong counterirritant properties of menthol even at low concentrations [43]. In E-cigarette users, menthol is transferred to the airways with high efficiency, where menthol concentrations per puff are comparable to concentrations in puffs generated from menthol cigarettes [14,33]. Previous analytical studies demonstrated that menthol concentrations in marketed E-liquids range from 2-68,000 µg/mL (13 µM-435 mM) [49]. In the present study, menthol levels in E-cigarette refill liquids and Puffbar devices ranged from 330 – 23,490 µg/g (~2.4-173 mM) and 620-3,680 µg/g (~4.6-27 mM), respectively. These levels are sufficient to reduce sensory perception of irritation and harshness elicited by nicotine in humans [50]. Similar to menthol, synthetic cooling agents (including WS-3 and WS-23) also activate TRPM8 with increased efficacy (WS-3) and with comparable half-maximal concentrations (WS-3: 2-4µM; WS-23: 44µM) [46,51,52]. Several of the refill E-liquids analyzed in this study

contained WS-3 at levels comparable to menthol levels in mint- and menthol-flavored E-liquids

(150-14,650 µg/g; ~816 µM-80 mM), with some containing synthetic coolants in addition to

"natural" cooling agents. The concentrations of WS-3 in several Puffbar E-liquids, up to 1%

(1,880-10,770 µg/g; 10-59 mM), are similar to menthol concentrations in liquids of Juul, the

market-leading pod E-cigarette system in the US [14,33]. In addition, the majority of the Puffbar

E-liquids analyzed in this study contained WS-23 ranging between 1-3.7% (up to 250 mM).

Carryover of synthetic cooling agents from E-liquids into the vapor was highly efficient,

approximating 100%, as previously demonstrated for menthol (Table1, SI Table1) [33]. Based on

the puffing regimen in our study, the aerosol concentrations of WS-3 for majority of refill E-

liquids (18/24 E-liquids) was calculated to be higher than 5 ppm (range: 5.8-98 ppm). Similarly,

WS-3 and WS-23 concentrations for all of disposable Puffbars tested exceeded 10 ppm (WS-3:

12-72 ppm; WS-23: 13-303 ppm), which is sufficient to mediate TRPM8-dependent

counterirritant effects. Since these synthetic coolants are either more potent or comparable

TRPM8 agonists to menthol, it is likely they facilitate inhalation in vapers using the products

analyzed by us. One recent study also quantified menthol and the synthetic coolants (WS-3 and

WS-23) in Puffbar devices [15]. This study (Omaiye et al.,) found similar quantities of synthetic

coolants and a higher quantity of menthol in the Puffbar "Cool Mint" product, and higher

quantities of all 3 compounds in the "Menthol" product [15] . Some of these differences in amounts

could be that, the tested "Cool Mint" device in this study was smaller in size (1.3 mL) than what

was analyzed in Omaiye et al., study (3.2 mL). It is thus possible that new products were

introduced and that e-liquid contents might have been adjusted simultaneously. Another

observation to explain this difference could be the degree of e-liquid content variability (i.e.,

production inconsistency and manufacturing sites) of the Puffbar devices, such as evident in

WS-23 contents in the "Menthol" product in this study and menthol content in the "Cool Mint"

product reported earlier [15] . This clearly illustrates the variability and inconsistencies of the

manufacturing practices in the market place.

Our risk assessment analysis suggests that users of E-cigarettes containing synthetic

cooling agents are exposed to amounts of WS-3 and WS-23 that exceed the amounts

considered acceptable by regulators for intake from food. Detected amounts were much higher

than the thresholds set by JECFA for human intake of flavor additives of this particular class of

chemicals (aliphatic and aromatic amines and amides, FEMA Nos 1594-1601, 1606 and 1613)

at 90 µg/person/day, either added as individual flavoring or combined total [20]. Even consumption

of just 1-mL of WS-3 containing E-cigarette liquid (150-14650 µg/g) results in exposures

exceeding JECFA's safety threshold by ~2-180 fold. Vaping of 1 disposable Puffbar E-cigarette

containing synthetic cooling agents would expose a user to amounts of WS-3 (1,880-10770

µg/g) that exceed the threshold of concern by ~29-176 fold, or ~26-598 fold for WS-23 (3,940-

36660 µg/g). There were 5 Puffbar varieties that contained both WS-3 and WS-23, and when

added together, would expose a user vaping 1 Puffbar device to ~126-774 fold the

recommended maximal daily amount for amides, based on JECFA recommendations. These

results strongly suggest that E-cigarette users are exposed to synthetic coolant amounts which

would be considered unsafe when consumed in food, raising concerns for health risks upon

chronic exposure.

The available chemical toxicity margins for WS-3 and WS-23, such as provided by

JECFA, are derived from toxicological studies in animals in which coolants were administered

orally [20]. Calculating MOE for inhalation of a compound based on past oral toxicity studies

requires some caution. For human risk assessment and to calculate risk assessment

parameters from exposure to these chemicals through various routes, it is routine practice for

US and International regulatory agencies (US Environmental Protection Agency, EPA;

European Chemical Agency, ECHA; UK's Interdepartmental Group on Health Risks from

Chemicals, IGHRC) to employ route-to-route (R2R) extrapolation [53,54]. Regulatory agencies

agree that the absorption efficiency for chemicals by the respiratory system is higher than the

efficiency of the digestive tract and usually, an oral-to-inhalation ratio of 2 is applied for route-to-

route extrapolation, reflecting the higher absorption of toxicants following inhalation [53–55]. In this

study, R2R extrapolation was not used, however, if applied, the MOE values would be even

lower, reflecting an even higher risk associated with inhalation of these two synthetic cooling

agents at various daily consumption volumes. This would suggest even greater urgency for

regulatory measures mitigating the risk from WS-3 and WS-23 exposures in E-cigarette users.

Recently, an e-vapor product company (RLX technology, China) has conducted a nose-

only acute and sub-acute inhalation study for WS-23 (28-day exposure at doses of 19 and 29

mg/kg-bw/day) [24]. Whole-body plethysmography was conducted after 28 days of repeated

exposure to measure various parameters of pulmonary function. Compared to control and

solvent groups, mice exposed to WS-23 demonstrated significant changes in various

parameters, such as time of expiratory (Te), peak expiratory flow (PEF), relaxation time (RT),

minute volume (MV), respiratory rate (F), end-inspiration pause (EIP), end-expiratory pause

(EEP). These results indicate that WS-23 may alter pulmonary function of users of WS-23

containing E-cigarettes. However, the study reports no remarkable histopathological changes in

either the respiratory organs (nose, throat, trachea and lungs) or other visceral organs (liver,

kidney, heart etc.). It has to be noted that the histopathological analysis was qualitative, and not

based on a quantitative histopathologic scoring. The absence of histopathological evidence for

organ toxicity contradicts prior WS-23 oral toxicity studies in rats that reported kidney lesions

and hepatic toxicity starting at 10 mg/kg-bW/day (NOAEL of 5 mg/kg-bW/day) [20,21]. The

discrepancy between these observations could be due to the different routes of exposure (oral

vs. inhalation), or more importantly, may have resulted from the different exposure durations

(14-weeks oral dosing vs. 4-weeks inhalational exposure).

In vitro genotoxicity studies of WS-23 in mammalian cells showed evidence for

clastogenicity [20]. Clastogenicity was observed only in the presence of metabolic activation,

indicating that the effect is mediated by the formation of a reactive metabolite. Due to the

increased exposure to WS-23 from use of E-cigarettes, it is imperative that more studies are

conducted to understand the mechanisms of their metabolism and further identify WS-23's

reactive clastogenic metabolites.

Previous studies have examined the toxicological effects of synthetic cooling agents,

including WS-3 and WS-23, in isolation, and not in mixtures. As demonstrated in this study, refill

E-cigarette liquids and popular disposable E-cigarettes contain mixtures of natural and synthetic

cooling agents. Inhalation toxicity studies are required to examine the effects of acute, sub-

acute and chronic exposures to mixtures of these chemicals and marketed E-cigarettes that

contain such mixtures. Another limitation in assessing the risk associated from use of these E-

cigarettes is the rapidly evolving product variety and the dynamic nature of their composition.

The products analyzed in this study were purchased in the last 1-2 years. More recent marketed

products may contain different compositions. Hence, continuous monitoring of new E-cigarette

products by determining their chemical compositions for presence of potentially toxic flavorant

levels is critical for both regulatory and risk assessment purposes.

Following the withdrawal of flavored Juul cartridges from the US market, disposable

E-cigarette products of the brand "Puffbar" brand have become increasingly popular among US

youth and young adults [6,56]. In Puffbar products, synthetic cooling agents were not only found in

mint/menthol-flavored, or "cool"- or "ice"-labelled E-liquids, but in almost all products tested

(13/14), suggesting that a cooling effect is preferred by consumers. Cooling agents were added

to fruit-flavored E-liquids named "Orange Ice" and "Blueberry Ice", but also "Strawberry Banana"

and "Lychee", without the "Ice" label, and in "Cucumber" and "OMG" flavors. This illustrates the

utility of odorless synthetic cooling agents to add a cooling effect to flavors that likely would not

be favored by users when combined with menthol, due to the incompatible minty odor the

"natural" coolant menthol would add. Synthetic cooling agents may allow tolerance of more

intense flavor combinations that would otherwise be irritating.

The fact that synthetic coolants are odorless also raises another important question:

Legislation such as the FSPTCA or the European Tobacco Product Directive (TPD) [1,57] banned

"characterizing" flavorants from combustible tobacco products (with exemption for menthol in

the US), yet is unclear how synthetic coolants lacking a characterizing flavor would be treated. It

has to be noted that TPD does not allows additives in tobacco products that facilitates

inhalation, but no specific compounds that facilitate inhalation or method to measure facilitation

are currently listed in the TPD. So, a replacement of the cooling compound menthol (with a

minty characterizing flavor) with a synthetic coolants (lacking characterizing flavor) could be a

strategy to circumvent a menthol ban. Policy makers should consider this when implementing

menthol bans, such as both Canada and Germany have done in respective regulations that

either maintain a positive list of compounds permitted as additives in tobacco products

(Canada), or specifically ban a whole class of compounds that facilitates inhalation, including all

natural and synthetic derivatives of menthol and other cooling agents (Germany) [58,59].

Conclusions:

Synthetic cooling agents are added to US-marketed E-cigarettes in a wide range of amounts that are comparable and relative to menthol amounts in flavored E-cigarettes. Synthetic coolants were found not only in mint- and menthol-flavored products, but also in fruit-, dessert- and sweet-flavored E-cigarettes. These coolants are added to refill liquids, E-cigarette liquid diluents and popular disposable E-cigarettes such as Puffbar, at levels that pose potential health risks to users, thereby requiring for these products to be prioritized for risk mitigation measures.

Funding:

This work was supported by the National Institute on Drug Abuse (NIDA) of the National

Institutes of Health (NIH) and the Center for Tobacco Products of the US Food and Drug

Administration (FDA) (P50DA036151 and U54DA036151; Yale Tobacco Center of Regulatory

Science, TCORS); and National Institute of Environmental Health Sciences (NIEHS)

(R01ES029435 to Dr. Sven-Eric Jordt).

Declaration of Interests:

The funding organization had no role in the design and conduct of the study; the collection,

management, analysis, and interpretation of the data; the preparation, review, or approval of the

manuscript; nor in the decision to submit the manuscript for publication. The content is solely

the responsibility of the authors and does not necessarily represent the views of National

institutes of Health (NIH) or the Food and Drug Administration (FDA). Dr. Jordt reports receiving

personal fees from Hydra Biosciences LLC and Sanofi S.A. and nonfinancial support from

GlaxoSmithKline Pharmaceuticals outside the submitted work. No other financial disclosures

were reported by the authors of this paper.

Author contributions were as follows: SVJ, HCE and SEJ conceptualized and designed

the study; DGT, LD, and HCE acquired and analyzed the chemical analytical data; SVJ

analyzed and interpreted the data for risk analysis calculations; SVJ, SEJ and HCE drafted the

manuscript; SVJ, HCE, PTA, JBZ, and SEJ provided supervision; JBZ critically revised the

manuscript for important intellectual content.

Data Availability Statement:

The data underlying this article are available in the article and in its online supplementary

material.

1. FDA Center for Tobacco Products. Family Smoking Prevention and Tobacco Control Act -
An Overview. *FDA*. Published online June 3, 2020. Accessed June 7, 2021.
https://www.fda.gov/tobacco-products/rules-regulations-and-guidance/family-smoking-
prevention-and-tobacco-control-act-overview

2. FDA Center for Tobacco Products. Menthol and Other Flavors in Tobacco Products. *FDA*.
Published online April 28, 2021. Accessed May 14, 2021. https://www.fda.gov/tobacco-
products/products-ingredients-components/menthol-and-other-flavors-tobacco-products

3. Villanti AC, Mowery PD, Delnevo CD, Niaura RS, Abrams DB, Giovino GA. Changes in the
prevalence and correlates of menthol cigarette use in the USA, 2004–2014. *Tobacco
Control*. 2016;25(Suppl 2):ii14-ii20. doi:10.1136/tobaccocontrol-2016-053329

4. Food and Drug Administration. FDA Commits to Evidence-Based Actions Aimed at Saving
Lives and Preventing Future Generations of Smokers. FDA. Published April 30, 2021.
Accessed May 10, 2021. https://www.fda.gov/news-events/press-announcements/fda-
commits-evidence-based-actions-aimed-saving-lives-and-preventing-future-generations-
smokers

5. Office of the Commissioner. FDA finalizes enforcement policy on unauthorized flavored
cartridge-based e-cigarettes that appeal to children, including fruit and mint. FDA.
Published March 24, 2020. Accessed May 14, 2021. https://www.fda.gov/news-
events/press-announcements/fda-finalizes-enforcement-policy-unauthorized-flavored-
cartridge-based-e-cigarettes-appeal-children

6. Wang TW. E-cigarette Use Among Middle and High School Students — United States,
2020. *MMWR Morb Mortal Wkly Rep*. 2020;69. doi:10.15585/mmwr.mm6937e1

7. Ali FRM. E-cigarette Unit Sales, by Product and Flavor Type — United States, 2014–2020.
MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6937e2

8. Leffingwell JC. Cool without Menthol & Cooler than Menthol and Cooling Compounds as
Insect Repellents. Cool without Menthol & Cooler than Menthol. Published June 8, 2018.
Accessed May 14, 2021. http://www.leffingwell.com/cooler_than_menthol.htm

9. Leffingwell JC, Rowsell D. Wilkinson Sword Cooling Compounds: From the Beginnning to
Now. *Perfumer & Flavorist*. 2014;39(3):34-43.

10. Leffingwell JC. WILKINSON-SWORD "'CWM'". Published online March 5, 1975.
https://www.industrydocuments.ucsf.edu/docs/tkjn0089

11. Newman F. Memorandum From Philip Morris In-House Counsel (Newman) To Philip
Morris In-House Counsel (Holtzman) Providing Legal Advice And Analysis Regarding
Content Of Draft Press Release For New Product Introduction;   Northwind Comments On
3rd Tier Inquires. Published online March 14, 1981. Accessed May 24, 2021.
https://www.industrydocuments.ucsf.edu/docs/kgxy0101

12. Reger L; M J; Hahn, H; Hahn, J. Analysis of Menthol, Menthol-Like, and Other Tobacco
Flavoring Compounds in Cigarettes and in Electrically Heated Tobacco Products. *Beiträge
zur Tabakforschung International Contributions to Tobacco Research*. 2018;28(2):93-102.
doi:10.2478/cttr-2018-0010

13. Jabba SV, Jordt SE. Turbocharged Juul device challenges European tobacco regulators.
Eur Respir J. 2020;56(2). doi:10.1183/13993003.02430-2020

14. Erythropel HC, Anastas PT, Krishnan-Sarin S, O'Malley SS, Jordt SE, Zimmerman JB.
Differences in flavourant levels and synthetic coolant use between USA, EU and Canadian
Juul products. *Tob Control.* Published online April 27, 2020. doi:10.1136/tobaccocontrol-
2019-055500

15. Omaiye EE, Luo W, McWhirter KJ, Pankow JF, Talbot P. Flavour chemicals, synthetic
coolants and pulegone in popular mint-flavoured and menthol-flavoured e-cigarettes.
Tobacco Control. Published online June 30, 2021. doi:10.1136/tobaccocontrol-2021-
056582

16. Liquid Nicotine Wholesalers. Koolada 10% PG (TFA). Liquid Nicotine Wholesalers.
Accessed June 8, 2021. https://liquidnicotinewholesalers.com/flavor-apprentice-koolada-
10-pg.html

17. Jabba SV, Jordt SE. Risk Analysis for the Carcinogen Pulegone in Mint- and Menthol-
Flavored e-Cigarettes and Smokeless Tobacco Products. *JAMA Intern Med.* Published
online September 16, 2019. doi:10.1001/jamainternmed.2019.3649

18. Jabba SV, Jordt SE. Estimating Fluid Consumption Volumes in Electronic Cigarette Use-
Reply. *JAMA Intern Med.* 2020;180(3):468-469. doi:10.1001/jamainternmed.2019.6630

19. Hallagan J. The Safety Assessment and Regulatory Authority to Use Flavors – Focus on
E-Cigarettes. Published online March 3, 2015. Accessed June 8, 2021.
https://www.femaflavor.org/sites/default/files/FEMAGRAS_Ecig_March_3_2015.pdf

20. Joint Expert Committee on Food Additives, ed. *Evaluation of Certain Food Additives: Sixty-*
Ninth Report of the Joint FAO/WHO Expert Committee on Food Additives ; [Rome, 17 - 26
June 2008].; 2009. http://www.inchem.org/documents/jecfa/jecmono/v952je01.pdf

21. Cheng M. *A Subchronic Oral Toxicity Study of T0332.04 in Rats. Unpublished Report to*
the Flavor Manufacturers Association. Submitted to WHO by the Flavor and Extract
Manufacturers Association of the United States, Washington DC, USA. Hazleton
Laboratories, Madison, Wisconsin, USA.; 1982.

22. Miyata K. *Summary of 28-Day Repeated Dose Oral Toxicity Study of WS-3. Unpublished*
Report to the Flavor Manufacturers Association. Submitted to WHO by the Flavor and
Extract Manufacturers Association of the United States, Washington DC, USA.; 1995.

23. Joint FAO/WHO Expert Committee on Food Additives. Meeting (65th : 2005: Geneva
Switzerland), International Program on Chemical Safety. *Safety Evaluation of Certain Food*
Additives. World Health Organization, International Programme on Chemical Safety; 2006.

24. Wu ZH, Liu YS, Li XD, et al. Acute and subacute inhalation toxicity assessment of WS-23
in Sprague-Dawley rats. *J Appl Toxicol.* Published online March 23, 2021.
doi:10.1002/jat.4166

25.   Leventhal A, Dai H, Barrington-Trimis J, Sussman S. 'Ice' flavoured e-cigarette use among
young adults. *Tobacco Control.* Published online May 16, 2021.
doi:10.1136/tobaccocontrol-2020-056416

26.   Galimov A, Leventhal A, Meza L, et al. Prevalence of disposable pod use and consumer
preference for e-cigarette product characteristics among vape shop customers in Southern
California: a cross-sectional study. *BMJ Open.* 2021;11(10):e049604.
doi:10.1136/bmjopen-2021-049604

27.   Davis DR, Morean ME, Bold KW, et al. Cooling e-cigarette flavors and the association with
e-cigarette use among a sample of high school students. *PLOS ONE.*
2021;16(9):e0256844. doi:10.1371/journal.pone.0256844

28.   Park-Lee E. Notes from the Field: E-Cigarette Use Among Middle and High School
Students — National Youth Tobacco Survey, United States, 2021. *MMWR Morb Mortal*
Wkly Rep. 2021;70. doi:10.15585/mmwr.mm7039a4

29.   Gaiha SM, Lempert LK, McKelvey K, Halpern-Felsher B. E-cigarette devices, brands, and
flavors attract youth: Informing FDA's policies and priorities to close critical gaps. *Addictive*
Behaviors. 2022;126:107179. doi:10.1016/j.addbeh.2021.107179

30.   Erythropel HC, Jabba SV, DeWinter TM, et al. Formation of flavorant-propylene Glycol
Adducts With Novel Toxicological Properties in Chemically Unstable E-Cigarette Liquids.
Nicotine Tob Res. 2019;21(9):1248-1258. doi:10.1093/ntr/nty192

31.   Rosbrook K, Erythropel HC, DeWinter TM, et al. The effect of sucralose on flavor
sweetness in electronic cigarettes varies between delivery devices. *PLoS One.*
2017;12(10):e0185334. doi:10.1371/journal.pone.0185334

32.   Talih S, Balhas Z, Eissenberg T, et al. Effects of User Puff Topography, Device Voltage,
and Liquid Nicotine Concentration on Electronic Cigarette Nicotine Yield: Measurements
and Model Predictions. *Nicotine & Tobacco Research.* 2015;17(2):150.
doi:10.1093/ntr/ntu174

33.   Erythropel HC, Davis LM, de Winter TM, et al. Flavorant-Solvent Reaction Products and
Menthol in JUUL E-Cigarettes and Aerosol. *Am J Prev Med.* 2019;57(3):425-427.
doi:10.1016/j.amepre.2019.04.004

34.   WHO/IPC. *Environmental Health Criteria 239: Principles for Modellingdose–Response for*
the Risk Assessment of Chemicals. World Health Organization
https://apps.who.int/iris/bitstream/handle/10665/44065/WHO_EHC_240_8_eng_Chapter5.
pdf?sequence=8

35.   Benford D, Bolger PM, Carthew P, et al. Application of the Margin of Exposure (MOE)
approach to substances in food that are genotoxic and carcinogenic. *Food and Chemical*
Toxicology. 2010;48:S2-S24. doi:10.1016/j.fct.2009.11.003

36.   Jabba SV, Jordt SE. Risk Analysis for the Carcinogen Pulegone in Mint- and Menthol-
Flavored e-Cigarettes and Smokeless Tobacco Products. *JAMA Intern Med.* Published
online September 16, 2019. doi:10.1001/jamainternmed.2019.3649

37. Baumung C, Rehm J, Franke H, Lachenmeier DW. Comparative risk assessment of
tobacco smoke constituents using the margin of exposure approach: the neglected
contribution of nicotine. *Sci Rep*. 2016;6(1):35577. doi:10.1038/srep35577

38. Chaumont M, de Becker B, Zaher W, et al. Differential Effects of E-Cigarette on
Microvascular Endothelial Function, Arterial Stiffness and Oxidative Stress: A Randomized
Crossover Trial. *Sci Rep*. 2018;8(1):10378. doi:10.1038/s41598-018-28723-0

39. Smets J, Baeyens F, Chaumont M, Adriaens K, Van Gucht D. When Less is More: Vaping
Low-Nicotine vs. High-Nicotine E-Liquid is Compensated by Increased Wattage and Higher
Liquid Consumption. *Int J Environ Res Public Health*. 2019;16(5).
doi:10.3390/ijerph16050723

40. National Research Council (U.S.), ed. *Risk Assessment in the Federal Government:*
Managing the Process. National Academy Press; 1983.

41. McKemy DD, Neuhausser WM, Julius D. Identification of a cold receptor reveals a general
role for TRP channels in thermosensation. *Nature*. 2002;416(6876):52-58.
doi:10.1038/nature719

42. Peier AM, Moqrich A, Hergarden AC, et al. A TRP channel that senses cold stimuli and
menthol. *Cell*. 2002;108(5):705-715. doi:10.1016/s0092-8674(02)00652-9

43. Willis DN, Liu B, Ha MA, Jordt SE, Morris JB. Menthol attenuates respiratory irritation
responses to multiple cigarette smoke irritants. *FASEB J*. 2011;25(12):4434-4444.
doi:10.1096/fj.11-188383

44. Ha MA, Smith GJ, Cichocki JA, et al. Menthol attenuates respiratory irritation and elevates
blood cotinine in cigarette smoke exposed mice. *PLoS One*. 2015;10(2):e0117128.
doi:10.1371/journal.pone.0117128

45. Fan L, Balakrishna S, Jabba SV, et al. Menthol decreases oral nicotine aversion in
C57BL/6 mice through a TRPM8-dependent mechanism. *Tob Control*. 2016;25(Suppl
2):ii50-ii54. doi:10.1136/tobaccocontrol-2016-053209

46. Behrendt HJ, Germann T, Gillen C, Hatt H, Jostock R. Characterization of the mouse cold-
menthol receptor TRPM8 and vanilloid receptor type-1 VR1 using a fluorometric imaging
plate reader (FLIPR) assay. *Br J Pharmacol*. 2004;141(4):737-745.
doi:10.1038/sj.bjp.0705652

47. Klein AH, Iodi Carstens M, McCluskey TS, et al. Novel menthol-derived cooling
compounds activate primary and second-order trigeminal sensory neurons and modulate
lingual thermosensitivity. *Chem Senses*. 2011;36(7):649-658. doi:10.1093/chemse/bjr029

48. Tobacco Products Scientific Advisory Committee. *Menthol Cigarettes and Public Health:*
Review of the Scientific Evidence and Recommendations. US Department of Health and
Human Services, Food and Drug Administration; 2011:1-252. Accessed April 27, 2021.
https://wayback.archive-
it.org/7993/20170405201731/https://www.fda.gov/downloads/AdvisoryCommittees/Committ
eesMeetingMaterials/TobaccoProductsScientificAdvisoryCommittee/UCM269697.pdf

49. Omaiye EE, McWhirter KJ, Luo W, Tierney PA, Pankow JF, Talbot P. High concentrations
of flavor chemicals are present in electronic cigarette refill fluids. *Scientific Reports*.
2019;9(1):2468. doi:10.1038/s41598-019-39550-2

50. Rosbrook K, Green BG. Sensory Effects of Menthol and Nicotine in an E-Cigarette.
Nicotine & Tobacco Research. 2016;18(7):1588-1595. doi:10.1093/ntr/ntw019

51. Sherkheli MA, Vogt-Eisele AK, Bura D, Beltrán Márques LR, Gisselmann G, Hatt H.
Characterization of selective TRPM8 ligands and their structure activity response (S.A.R)
relationship. *J Pharm Pharm Sci*. 2010;13(2):242-253. doi:10.18433/j3n88n

52. Johnson S, Tian M, Sheldon G, Dowd E. Trigeminal Receptor Study of High-Intensity
Cooling Agents. *J Agric Food Chem*. 2018;66(10):2319-2323.
doi:10.1021/acs.jafc.6b04838

53. European Chemicals Agency (ECHA). *Guidance on Information Requirements and
Chemical Safety Assessment Chapter R.8: Characterisation of Dose [Concentration]-
Response for Human Health*. European Chemicals Agency; 2012.
https://echa.europa.eu/documents/10162/13632/information_requirements_r8_en.pdf

54. The Interdepartmental Group on Health Risks from Chemicals (IGHRC). *Guidelines on
Route-to-Route Extrapolation of Toxicity Data When Assessing Health Risks of Chemicals*.
Cranfield University; 2006.
http://www.iehconsulting.co.uk/IEH_Consulting/IEHCPubs/IGHRC/cr12.pdf

55. Schröder K, Escher SE, Hoffmann-Dörr S, Kühne R, Simetska N, Mangelsdorf I.
Evaluation of route-to-route extrapolation factors based on assessment of repeated dose
toxicity studies compiled in the database RepDose®. *Toxicol Lett*. 2016;261:32-40.
doi:10.1016/j.toxlet.2016.08.013

56. Ali FRM. E-cigarette Unit Sales, by Product and Flavor Type — United States, 2014–2020.
MMWR Morb Mortal Wkly Rep. 2020;69. doi:10.15585/mmwr.mm6937e2

57. European Union. Revision of the Tobacco Products Directive. Public Health - European
Commission. Published November 25, 2016. Accessed June 8, 2021.
https://ec.europa.eu/health/tobacco/products/revision_en

58. TabakerzV - Verordnung über Tabakerzeugnisse und verwandte Erzeugnisse. Accessed
June 8, 2021. https://www.gesetze-im-internet.de/tabakerzv/BJNR098010016.html

59. Health Canada. Order Amending the Schedule to the Tobacco Act (Menthol). Published
August 2, 2017. Accessed June 8, 2021. https://www.canada.ca/en/health-
canada/programs/consultation-changes-tobacco-act-address-menthol/order-amending-
schedule-tobacco-act-menthol.html

Figure Legends

Figure 1: Margins of Exposure (MOE) for synthetic cooling agents from E-Cigarette refill liquids and coolant concentrate. MOEs are plotted for representative WS-3 (left) and WS-23 (right) -containing E-liquids and diluents with high, intermediate and low coolant concentrations, for daily consumption of 1, 3, 5, 10 and 15 ml E-liquid. No-Observed-Adverse-Effect-Levels (NOAEL) determined by JECFA in animal studies of organ toxicity upon oral administration were used for MOE calculations. A MOE of <100 (dashed line) signals a safety concern and requires regulatory mitigation. For WS-23 concentrate (Agent Cool, TBD Liquids), MOEs were calculated for dilutions within the range of manufacturer-recommendations to 1 and 3%. MOEs of the oral average daily intake of US consumers (US-OADI) determined by JECFA are displayed for comparison.

Figure 2: Margins of Exposure (MOE) for synthetic cooling agents from Puffbar-branded disposable E-cigarettes. MOEs are plotted for exposures from Puffbar devices with flavored E-liquids containing both WS-3 and WS-23, at high (Cool mint), intermediate (Blueberry ice) or low (Menthol) concentrations, separated for each coolant (left: WS-3, right: WS-23), calculated for daily consumption of ½ Puffbar (0.65 mL), 1 Puffbar (1.3 mL) and 2 bars (2.6 mL). JECFA-determined NOAEL values were used for MOE calculations. MOEs of the oral average daily intake of US consumers (US-OADI) determined by JECFA are displayed for comparison.

Table 1: Concentrations of synthetic cooling agents (WS-3 and WS-23), menthol, menthone, carvone, nicotine and solvent chemicals in US-marketed E-cigarette refill liquids. Carryover rates for synthetic coolants are provided for select refill liquids. [a]single experiment. WS-3 and WS-23 concentrations are highlighted in yellow and green, respectively; Concentrate E-liquid highlighted in orange.

| | E-cigarette Flavor (Manufacturer) | Flavor Concentration ± s.d. (µg/g) | | | | | Nicotine concentration ± s.d. (mg/g) | PG/VG ratio (%/%) | Synthetic coolant carryover ± s.d. (%) |
		WS-3	WS-23	Menthol	Menthone	Carvone			
1	Arctic Rush (Central Vapors)	14650 ± 230					9.2 ± 0.09	50/50	106 ± 4
2	Frzn Mango (BLVK)	2450 ± 00		3640 ± 10			5.08 ± 0.03	30/70	
3	Nuclear White Menthol (Vapor HQ)	2440 ± 10		23490 ± 130	310 ± 20	190 ± 70	9.88 ± 0.02	50/50	104 ± 4
4	Rich Girl (Stunna)	2260 ± 30					3.31 ± 0.06	70/30	107 ± 11
5	Frozen Berries (Shinjin Vapor)	2220 ± 20		2320 ± 20			3.36 ± 0.05	30/70	
6	Frozen Limeade (Shinjin Vapor)	1540 ± 20		3470 ± 50			3.38 ± 0.04	30/70	
7	Cool Lime (Chill)	2110 ± 60					1.63 ± 0.05	30/70	
8	Tropicool (Chill)	2040 ± 40					3.02 ± 0.07	30/70	91[a]
9	Kool Cali (Chill)	1690 ± 20					4.20 ± 0.06	40/60	
10	Sakura 3 mg Nic (Sugoi)	1090 ± 20		910 ± 10			2.32 ± 0.03	30/70	105 ± 2
11	Coconut Ice (EC Blend)	2240 ± 10					1.41 ± 0.02	60/40	
12	Clove + Cool Hit (EC Blend)	1600 ± 20					17.91 ± 0.23	50/50	
13	Popsicle USA (EC Blend)	150 ± 10				10 ± 1		60/40	
14	Frozen (FJ's)	1550 ± 50						30/70	
15	Wintergreen Menthol (VaporHQ)	870 ± 20		4850 ± 30			9.72 ± 0.08	50/50	
16	Icy Menthol (Vista Vapors)	560 ± 10		9240 ± 90	1320± 50	2890 ± 10	9.80 ± 0.02	50/50	
17	Ice Dragon (ELD Flavors)	410 ± 30					2.75 ± 0.01	20/80	
18	Wintermint (Gothic Vapor)	330 ± 50		1110 ± 190		510 ± 10	9.03 ± 0.84	50/50	
19	Climate Change (What Juice)	290 ± 10		330 ± 10			9.48 ± 0.12	30/70	
20	Arctic (Vista Select)	280 ± 20		3470 ± 10	30 ± 10	930 ± 70	6.59 ± 0.06	30/70	
21	Dillinger (NastyJuice X Kilo)		1860 ± 40		280 ± 10		4.29 ± 0.07	30/70	99 ± 10
22	Kool Peach (Salty Man)	990 ± 20					46.4 ± 1.09	40/60	109 ± 1
23	Sakura Nic Salts 35 mg (Sugoi)	1170 ± 30		1170 ± 20			28.74 ± 0.43	50/50	
24	Sakura Nic Salts 50 mg (Sugoi)	1140 ± 60		1140 ± 90	14110 ± 670		39.19 ± 2.20	50/50	97[a]

25	Unicorn Nicotine Salt Spearmint (BLVK)	2950 ± 20		10920 ± 160	25450 ± 600	6830 ± 40	41.36 ± 0.33	40/60	107 ± 8
26	Agent Cool Super Concentrate (TBD Liquids)	8840 ± 130	124060 ± 720						

WS-3	
WS-23	Concentrate

Table 2: Chemical concentrations of synthetic cooling agents (WS-3 and WS-23), menthol and nicotine in flavors of US-marketed Puffbar disposable E-cigarettes. No menthone or carvone was found in any of the analyzed Puffbar e-liquids. PG/VG ratio measured in triplicate and rounded to next 5. Empty cells represent "not detected".

	Puffbar flavor	Flavor Concentration ± s.d. (µg/g)					Nicotine concentration ± s.d. (mg/g)	PG/VG ratio (%/%)
		WS-3	WS-23	Menthol	Menthone	Carvone		
1	Cool Mint	10770 ± 210	36660 ± 530	620 ± 50			32.79 ± 0.38	55/45
2	Café Latte		2300 ± 30	1500 ± 50			36.56 ± 0.30	55/45
3	Orange Ice	1750 ± 20	22950 ± 450				37.61 ± 0.76	30/70
4	Pomegranate						29.37 ± 0.47	50/50
5	Lush Ice		25830 ± 450	2250 ± 190			32.26 ± 0.53	55/45
6	Melon Ice		8810 ± 60				32.81 ± 0.29	65/35
7	Cucumber	3770 ± 20	3940 ± 10				31.67 ± 0.15	60/40
8	Lychee		26240 ± 240				33.22 ± 0.27	55/45
9	Menthol	3540 ± 30	1610 ± 1950	3680 ± 550			33.25 ± 0.22	45/55
10	Blueberry Ice	1880 ± 20	14120 ± 210	2300 ± 40			32.71 ± 0.45	50/50
11	Strawberry Banana		5420 ± 90				32.12 ± 0.52	50/50
12	OMG		7390 ± 330	1330 ± 40			34.01 ± 1.35	50/50
13	Peach Ice		11540 ± 270				33.11 ± 0.63	55/45
14	Banana Ice		13110 ± 50	2040 ± 110			40.66 ± 0.22	55/45

WS-3
WS-23
Menthol

Figure 1

Figure 2