Beyond tracking climate: niche evolution during native range expansion and its

implications for novel invasions

Nicky Lustenhouwer[1]* and Ingrid M. Parker[1]

[1] Department of Ecology and Evolutionary Biology, University of California, Santa Cruz, USA

* corresponding author: nlustenh@ucsc.edu

Abstract

Ecological niche models have been instrumental in understanding and forecasting the widespread

shifts in species distributions under global change. However, growing evidence of evolution in

spreading populations challenges their key assumption of niche conservatism, limiting model

transferability in space and time. Climate niche evolution has been studied extensively in

invasive species, but may also occur in native populations tracking climate change, when species

encounter novel abiotic conditions that vary with latitude. We compared niche shifts during

native range expansion and during invasion in *Dittrichia graveolens*, a Mediterranean annual

plant species that is currently undergoing both types of spread. We asked whether the species'

northward native range expansion in Eurasia matched climate change from 1901-1930 to 1990-

2019, or if further range expansion was promoted by niche evolution. In addition, we asked how

niche expansion in the native range affected forecasts of two ongoing invasions in Australia and

California. We quantified niche shifts in environmental space using the analytical framework of

niche centroid shift, overlap, unfilling, and expansion (COUE), and examined associated

distribution changes in geographic space using Maximum Entropy modeling. Our results show

that *D. graveolens* expanded its native range well beyond what would be sufficient to track

climate change, a shift associated with a 5.5% niche expansion to include more temperate

climates. In contrast, both invasions showed evidence of niche conservatism, with niche filling

depending on invader residence time. Including the expanded native niche in invasion

projections added new areas at risk of invasion, but none of these has been colonized at present.

We conclude that native populations may track climate change and adapt to novel local

conditions in parallel, causing an evolutionary expansion of the climate niche and more

widespread range expansion than expected based on climate change alone.

Keywords

Range shift; climate change; invasion; niche evolution; range expansion; habitat suitability

Introduction

Forecasting the widespread distribution shifts of both native and invasive species under global

change represents one of the major challenges in biodiversity conservation (Urban et al. 2016).

The climate niche has become a central concept in modeling efforts to understand species'

preferred climate conditions, where such suitable habitat will be present under future climates,

and which areas are at risk of invasion on other continents (Elith and Leathwick 2009).

Ecological niche models (ENMs; also known as habitat suitability models or species distribution

models) are widely used and applied to conservation planning (Araújo et al. 2011). However,

range-expanding populations have been demonstrated to rapidly adapt to novel environments in a

growing body of empirical studies (reviewed in Colautti and Lau 2015, Chuang and Peterson

2016), challenging the fundamental assumption of ENMs that niches are conserved in space and

time (Pearman et al. 2008). Newly developed mechanistic and hybrid species distribution models

that incorporate evolution of species' physiological limits (Hoffmann and Sgrò 2011, Catullo et

al. 2015) predict markedly different outcomes of climate-induced range shifts (Bush et al. 2016)

and invasions (Kearney et al. 2009) than traditional correlative ENMs. While there is thus a

compelling argument for incorporating evolution into range expansion forecasts (Nadeau and

Urban 2019), a better understanding of the degree to which climate niche evolution promotes

contemporary range expansions is necessary to make informed predictions.

Niche evolution has been studied extensively in invasive species, with studies comparing the

climate conditions occupied by populations in the native and invaded range. While there is

strong evidence that invading populations can rapidly evolve to reestablish adaptive clines along

environmental gradients similar to those in their native range (e.g., Boheemen et al. 2019),

whether species' ultimate niche limits are generally conserved during invasion (Petitpierre et al.

2012, Liu et al. 2020) or commonly shift (Early and Sax 2014, Atwater et al. 2018) is highly

debated. Niche stasis (*sensu* Pearman et al. 2008) is defined as the lack of change in either the

fundamental niche (the climate where a species can grow in the absence of biotic constraints and

geographic barriers) or realized niche (the actual climate conditions where a species is observed,

which is captured by ENMs). Niches may shift in the invaded range when not all areas with

similar climates are occupied (yet) due to dispersal limitation. Niche expansion occurs when

biotic interactions or dispersal barriers constraining the realized niche in the native range are

lifted in the new range, or when the fundamental niche itself evolves (Alexander and Edwards

2010). Examples of evolutionary changes that have been linked to climate niche expansion in

invasive populations include plant phenology responses to temperature or photoperiod (Dlugosch

and Parker 2008a, Colautti and Barrett 2013) and insect thermal and moisture tolerance (Kearney

et al. 2009, Hill et al. 2013).

Much less attention in the empirical literature has been paid to niche evolution during

contemporary native range shifts induced by climate change. Instead, species' responses to

global warming are commonly viewed as a "move, adapt, or die" conundrum, where populations

need to track suitable climates to higher latitudes and altitudes, or adapt *in situ*, or else they will

lag behind the pace of climate change and go extinct (Aitken et al. 2008, Corlett and Westcott

2013). However, poleward-spreading populations face a variety of novel abiotic (as well as

biotic) conditions, even if range expansion is initiated by climate change (Spence and Tingley

2020). For example, photoperiod and temperature seasonality increase non-linearly with latitude,

and plants experience a reduction in photosynthetically active radiation and light quality towards

the poles (Taulavuori et al. 2010, Saikkonen et al. 2012). These novel combinations of

temperature and photoperiod cues at higher latitudes require a plastic or evolutionary response

(Visser 2008). Thus, whether climate-mediated range shifts will involve simultaneous niche

evolution is now acknowledged as an important open question (Lee-Yaw et al. 2019, Nadeau and

Urban 2019). Recent empirical work has demonstrated rapid evolutionary responses to novel

abiotic conditions during contemporary native range shifts, including increased thermal niche

breadth in damselflies (Lancaster et al. 2015, Dudaniec et al. 2018) and earlier fall phenology in

plants (Lustenhouwer et al. 2018). Palaeoecological studies of the Pleistocene glaciations have

shown that dispersal, plasticity, evolution and niche conservatism have all played a role in past

range shifts with climate change (Martínez-Meyer et al. 2004, Nogués-Bravo et al. 2018).

Historic constraints (or the lack of them) on range expansion may inform how evolution will

affect recent range shifts with climate change (Diamond 2018).

Comparing native range shifts and invasions provides valuable insights into the drivers of niche

evolution during range expansion in both native and exotic ranges. Quantifying rapid adaptation

to novel climates during invasions can inform predictions about the degree of evolutionary

change that may be expected during native range expansions on similar time scales (Moran and

Alexander 2014). Similarly, the likelihood of niche shifts in the invaded range depends on

whether the native range is limited by biotic interactions or fundamental evolutionary constraints

such as genetic correlations between traits (Alexander and Edwards 2010). Yet there are also

differences between native and exotic range expansions in key processes driving evolutionary

change, most notably the level of gene flow. Disconnection from the historic native range could

lead to increased niche evolution in invading populations (Jakob et al. 2010), unless genetic

diversity and population sizes are too low (Hoffmann and Sgrò 2011). However, genetic

bottlenecks have little impact in invasions founded by multiple introductions (Dlugosch and

Parker 2008b, Smith et al. 2020). In the native range, the impact of ongoing gene flow will

depend on whether it has a maladaptive swamping effect or increases evolutionary potential, an

issue which is highly debated (Kottler et al. 2021). Theory suggests that adaptation during range

expansion under gene flow is governed by the balance between selection and genetic drift on the

one hand, and the steepness of the environmental gradient on the other hand (Polechová and

Barton 2015). Overall, authors have hypothesized that niche shifts are more common or rapid in

exotic than in native range expansions (Pearman et al. 2008, Wiens et al. 2019), but the empirical

data required to test this idea are largely absent.

In this study, we examine climate niche evolution during both native and exotic range expansion

by taking advantage of a species currently involved in both types of population spread. *Dittrichia*

graveolens (L.) Greuter is an annual plant in the Asteraceae with a native distribution around the

Mediterranean Basin (Brullo and De Marco 2000). The species has greatly expanded its native

range northward since the mid-20[th] century, coincident with rapid evolution of earlier flowering

time at the leading edge in response to shorter growing seasons (Lustenhouwer et al. 2018). *D.*

graveolens has a ruderal life history and produces large numbers of wind-dispersed seeds,

facilitating spread along roads where biotic interactions with other plant species play a minor

role in its ecological success. The species has invaded worldwide in most other regions with a

Mediterranean climate - first Australia (1860s; Parsons and Cuthbertson 2001) and South Africa

(GBIF.org 2020), then California (1980s; Preston 1997), and most recently Chile (Santilli et al.

2021). In Australia and California, *D. graveolens* is considered a noxious weed of high

management concern due to a combination of rapid spread and toxicity to livestock, impacts on

native plant communities, and human skin allergies (Parsons and Cuthbertson 2001, Brownsey et

al. 2013b, USDA 2013).

We ask: (a) Did *D. graveolens* simply track climate change during the native range expansion, or

was range expansion promoted by climate niche evolution? (b) Is there evidence of niche shifts

in the invaded ranges? (c) How does taking into account climate niche evolution during the

native range expansion change invasion predictions for California and Australia? To answer

these questions, we applied the COUE scheme of niche centroid shift, overlap, unfilling and

expansion (Guisan et al. 2014) to *D. graveolens*' native range expansion with climate change

(comparing the periods 1901-1930 and 1990-2019) and to the two exotic range expansions. This

method quantifies niches in environmental space and is widely used to study niche dynamics of

invasive species. In addition, to explore niche shifts during range expansion in geographic space,

we fit species distribution models using maximum entropy modeling (Maxent), which was

specifically designed for presence-only data (Phillips et al. 2006). Based on prior evidence for

rapid evolution of phenology at the northern range edge (Lustenhouwer et al. 2018), we

hypothesized that *D. graveolens*' climate niche expanded during native range expansion in

Eurasia. We expected greater niche filling and greater niche expansion in Australia than in

California, due to *D. graveolens*' much longer invasion history in the former region. Finally, we

hypothesized that forecasting the two invaded distributions based on the newly expanded native

range would increase our invasion risk assessment for a wider range of climates and geographic

areas.

Material and Methods

Data collection

Occurrence data

We compiled presence-only species occurrence data for *Dittrichia graveolens* (L.) Greuter and

its nomenclatural synonyms *Inula graveolens* (L.) Desf., *Cupularia graveolens* (L.) Godr. &

Gren., and *Erigeron graveolens* (L.). We used the Holarctic Floral Kingdom (Cox 2001) as our

study region (split between Eurasia/North Africa for the native range and North America for one

of the invaded ranges), to take into account the broadest range of environments available to *D.*

graveolens in its evolutionary history and facilitate projection of our models to other continents

(Merow et al. 2013). This study region also allows for comparison to other studies using the

same spatial extent (Petitpierre et al. 2012). Our primary data source was the Global Biodiversity

Information Facility (~75% of occurrences; GBIF.org 2020), supplemented by country-level

species occurrence databases (FloraWeb 2013, Nikolić 2015, Zając and Zając 2019, BSBI 2020,

Info Flora 2020), standard floras, articles in botanical journals, and herbarium records. All

citizen science records from iNaturalist in GBIF were verified manually (iNaturalist 2020). We

used QGIS v3.8.2 (QGIS Development Team 2019) to combine and convert all data to the

WGS84 coordinate system with decimal degrees latitude and longitude. Spatial grids (UTM,

MTB, etc.) were imported as cell centroids. To increase sampling density across the study

region, we also georeferenced localities without spatial coordinates (such as towns and other

geographic features described in sufficient detail) using GEOLocate (Rios 2020). Data quality

control included the removal of duplicate records and those located at (0,0) degrees or country

centroids. The final (expanded) native range dataset included 8157 occurrences. To study niche

shifts in the exotic range of *D. graveolens*, we selected the invasions in Australia and California

because they are both well-documented (using records from GBIF and Calflora 2021,

respectively).

Defining the historic native range limit

D. graveolens' historic native distribution is centered around the Mediterranean Basin, extending

eastward to the Middle East and NW-India, and northward into central France (Tutin et al. 1976,

Brullo and De Marco 2000). The first records of a large-scale northward range expansion of *D.*

graveolens appear for the 1950s in northern France, initially associated with salt mines and

followed by abundant spread along roads (Wagenitz 1966, Ciardo and Delarze 2005, Parent

2011). We chose 1901-1930 as the pre-expansion time window for our study, which matches the

earliest available information about *D. graveolens'* distribution in floras of France (Coste 1903,

Rouy 1903, Bonnier and Layens 1909) and the Balkan Peninsula (Hayek and Markgraf 1931), as

well as the earliest monthly climate data (see next section). We used the botanical literature,

dated species occurrences, and online databases to define a hypothesized *historic native range*

limit pre-expansion (Supporting Information). We considered administrative regions where *D.*

graveolens is reported as a native species to be part of the historic native range (e.g., Flora

Europaea, Tutin et al. 1976, and Euro+Med PlantBase, von Raab-Straube 2021), refining

country-level data using other data sources.

Environmental predictors

Monthly temperature and precipitation data were compiled from the Climatic Research Unit

(CRU TS4.04, Harris et al. 2020) for 1901-1930 (past) and 1990-2019 (present), and used to

calculate all 19 WORLDCLIM variables (Fick and Hijmans 2017) for each time period (dismo

package; Hijmans et al. 2017). In addition, we calculated the average total number of frost days

for September-December (hereafter "frost") for the same data sets. We selected 6 predictors

(Table 1) based on the biology of *D. graveolens*, the Mediterranean and temperate climates

characteristic of the expanded native range, and criteria limiting multicollinearity among

variables: a pairwise Pearson correlation of 0.75 or less and a Variance Inflation Factor (usdm

package, Naimi et al. 2014) below 5 (as recommended by Guisan et al. 2017). *D. graveolens* is a

late-season annual plant, establishing a rosette in late spring and fruiting in fall (Brownsey et al.

2013b), with earlier phenology occurring at higher latitudes (Lustenhouwer et al. 2018). In early

life stages, precipitation is key to the growth of a tap root (Brownsey et al. 2013a), whereas

plants are sensitive to cold and especially frost later in the year (Parsons and Cuthbertson 2001),

when the end of the growing season constrains plant fitness (Lustenhouwer et al. 2018). We

considered climate variables representing temperature, precipitation, and their variability.

Collinearity was evaluated over the entire Eurasian Holarctic. Based on *D. graveolens*' biology,

we discarded annual temperature and precipitation in favor of frost days during the reproductive

period and precipitation in the driest and warmest quarters. Temperature of these quarters was

highly correlated with the other selected variables and excluded.

<u>Modeling approach</u>

To combine climate and occurrence data, we scaled the latter down to the same resolution with

one record per grid cell (0.5° latitude and longitude, corresponding to ca. 55 by 55 km). Cells

containing occurrence records but no climate data (primarily covering sea rather than land, at

coastlines or islands) were excluded. We set up two primary datasets, past and present, for our

native range analyses covering the same Holarctic study region. The past dataset (representing

the historic native range) consisted of climate data for the period 1901-1930 and all occurrence

records located within the historic native range limit (n=399), assuming that occurrence locations

represent suitable habitat for *D. graveolens* regardless of the date of observation. The present

dataset (representing the expanded native range) consisted of climate data for the period 1990-

2019 and the complete set of occurrence records (n=746). We used present-day climate data for

the invaded range datasets. The Australia dataset contained all GBIF occurrences on the

continent. The California dataset covered all of North America in spatial extent but occurrence

data for California only, to focus on the west coast invasion. All analyses were conducted in R

v4.0.0 (R Core Team 2020), using code adapted from Di Cola et al. (2017), Guisan et al. (2017),

and Smith (2020a).

Niches in environmental space

To quantify niche dynamics during range expansion following the COUE framework (Guisan et

al. 2014), we applied the workflow developed by Broennimann et al. (2012), available in the

ecospat R package (Di Cola et al. 2017, Broennimann et al. 2020). This approach evaluates

niches in environmental space, which is defined by conducting a principal component analysis

(PCA) for all environmental variables in both study regions under comparison. Niches are

estimated by applying a kernel smoother function to the density of species occurrences from each

range in gridded environmental space along the first two PCA axes. Indices of niche change are

calculated exclusively for environments that are available in both study regions (using the 90^{th}

percentile to exclude marginal environments), because shifts to and from nonanalog climates do

not represent shifts in the fundamental niche (Guisan et al. 2014). *Niche stability* is defined as the

proportion of occurrences in the new niche that overlap in environmental space with the native

niche, while *niche expansion* (1-stability) is the proportion of occurrences in the new niche that

are located in novel environments. Finally, *niche unfilling* quantifies environmental space that is

occupied in the native range but that has not been filled in the new range (yet). It is calculated for

the native occurrences as the proportion located in climate conditions that are unoccupied (but

available) in the new range (Di Cola et al. 2017). Overall *niche overlap* is measured by

Schoener's *D* (Schoener 1970) and used to test for niche conservatism during range expansion

with two different tests (Warren et al. 2008, Broennimann et al. 2012). The niche equivalency

test randomly reallocates occurrences between the two niches, creating a null distribution of *D*

values based on 100 permutations to test whether the niches are identical (conducting one-sided

tests asking if the observed *D* is greater or lower than expected by chance). The niche similarity

test evaluates whether the two niches are more or less similar to each other than to other niches

selected at random from the study area (shifting niches across both time periods for the native

study area, but in the invaded study area only for Australia and California). We applied the above

workflow to niche changes between (i) the past and present dataset (native range expansion), (ii)

the past dataset and each invaded range, and (iii) the present dataset and each invaded range.

Projections in geographic space

To project niche changes in geographic space, we fit Maxent species distribution models

(Phillips et al. 2006) to the past dataset (Past Model) and present dataset (Present Model).

Maxent performs well for presence-only data under a range of sampling scenarios (Grimmett et

al. 2020). Because the geographic availability of our native range occurrence data was highly

concentrated in Western Europe, we used 3468 target background sites of taxonomically related

species to correct for bias in sampling effort (Phillips et al. 2009, Supporting Information).

Models were fit using the maxnet package in R (Phillips 2017, Phillips et al. 2017). We first

tuned the Past Model, using the trainMaxnet function (enmSdm package, Smith 2020b) to select

the optimal combination of features (starting with all classes) and regularization parameters (β of

0.5 and 1 to 10). The best model (lowest AIC) contained linear, quadratic and product features

with $\beta=0.5$. We fit the Present Model using the same features and regularization. To ask whether

D. graveolens simply tracked optimal climate conditions or expanded its native range beyond

that, we created three model projections: the Past Model on the past and present climate, and the

Present Model on the present climate. To evaluate model performance, we partitioned the past

and present datasets randomly into training and test data using 5 k-folds. We computed AUC

(dismo, Hijmans et al. 2017) and the Continuous Boyce Index (CBI; enmSdm, Smith 2020b) for

each k-fold and calculated the mean and standard deviation across models. To project areas at

potential risk of invasion in California and Australia, we applied a Minimum Presence Threshold

(the lowest habitat suitability at which *D. graveolens* is already present in the invaded range).

We then compared projections generated by the Past and Present model for each invaded range.

Results

Over the course of less than a century, *D. graveolens* has shifted its native range limit northward

by nearly 7 degrees latitude. During this range expansion, the climate niche expanded by 5% to

include more temperate environments with lower diurnal range and precipitation seasonality

(bio2, bio15), increased precipitation in the driest and warmest quarters (bio17, bio18), and

increased temperature seasonality (bio4) and fall frost (Fig. 1a,b). Niche overlap (Schoener's D =

0.71) of the historic and expanded native niche was significantly lower than expected by chance

(niche equivalency test; Fig. 1c, Table 2), indicating a niche shift during range expansion.

Nonetheless, the two niches were more similar to each other than to randomly selected niches in

the study area (niche similarity test; Fig. 1d, Table 2). Niche expansion was not driven by

climate change between the past and present period, which happened in a different direction in

environmental space (reduced frost and temperature seasonality; Fig. 1a).

In line with this climate niche expansion, we found that *D. graveolens* in Eurasia has expanded

its geographic range well beyond the extent sufficient to track climate change (Fig. 2; Supporting

Information). The Past Model, fit to the historic climate and occurrences (Supporting

Information), predicts suitable habitat for *D. graveolens* around the Mediterranean Basin and into

central France as expected (Fig. 2a). Model AUC was 0.89 ± 0.01, indicating good to excellent

performance (Araújo et al. 2005). Projecting this model onto the present climate (Fig. 2b), we

found a minor northward shift in favorable conditions, particularly adjacent to the original

northern range limit in France. However, the observed range expansion of *D. graveolens*

extended much further northward and eastward (Fig. 2c). The Present Model, fit to the present

climate and occurrences (Supporting Information), predicts a much wider distribution in Europe

(Fig. 2d), outperforming the Past Model especially when predicting the actual probability of

occurrence in the present (CBI of 0.93 and 0.61, respectively; Table 3).

The invasions in California and Australia exhibited contrasting niche dynamics, consistent with

their difference in invader residence time. In California, only a small subset of the climate

conditions in the native niche are already occupied (niche unfilling was 93% at the scale of North

America; Fig. 3d, Table 2). In contrast, the Australian invasion has already spread into most

areas that show similar climatic conditions to the native range (unfilling 6%; Fig. 3a, Table 2).

These contrasting results were reflected in the niche equivalency test, which showed that the

California niche was less similar to the historic native range than expected by chance, while the

Australian niche was more similar than expected (Table 2). We found near-complete niche

stability in both invasions, meaning that invasive populations occur almost exclusively in climate

conditions that also existed in the historic native range. Accordingly, niche expansion was low

(Table 2). Niche conservatism could not be rejected in the niche similarity test for either

invasion, indicating that it cannot be ruled out that any niche differentiation between native and

exotic ranges was driven by the availability of environments in the exotic ranges.

Niche change indices were similar for both invasions when including the native range expansion

(Table 2). Even though the peak of native occurrences shifted toward more temperate climates

(shading in Fig. 3e), California niche unfilling remained stable because the majority of native

occurrences already occurred outside the invaded climate space. Of the newly invadable climate

conditions, most are absent in Australia (Fig. 3b), but present in North America (Fig. 3e).

Combining projections of the Past and Present Maxent models onto Australia and California

revealed additional areas that would be at risk of invasion if niche shifts were to happen as in the

native range. However, neither invasion has yet advanced into areas solely included by the

Present Model (Fig. 3c, 3f).

Discussion

As many native species are shifting their ranges to track climate change, the ecological and

evolutionary drivers of range expansion have become a key focus of global change research

(Nadeau and Urban 2019). Ecological studies of native range shifts have found varying degrees

to which species keep up with shifting climate isotherms (Chen et al. 2011, Lenoir et al. 2020),

including some species that outpace climate warming. Two studies using a similar approach to

ours, comparing historic and current ranges using ENMs, found incomplete climate tracking (*Ilex*

aquifolium, Walther et al. 2005) or range contractions rather than expansions (montane rodents,

Pardi et al. 2020). In contrast, our results show that *D. graveolens* has undergone an extensive

northward expansion of its native range, beyond what would be expected based on climate

tracking alone.

At the start of the native range expansion, the first records of *D. graveolens* north of the historic

range limit in France (GBIF.org 2020) match the availability of new suitable habitat there with

climate change (Fig. 2b). Theory predicts that rapid evolution of both local adaptation and

increased dispersal during range expansion could cause populations to spread beyond shifting

climate isotherms, while expanding their fundamental niche to include colder climates (Kubisch

et al. 2013). Our results are consistent with such a scenario, where adaptation to climate

conditions at northern latitudes (Fig. 1) facilitated further spread of *D. graveolens* (Fig. 2c).

Niche expansion (5.5%, following Broennimann et al. 2012) was high relative to comparative

studies of niche shifts during invasion using the same metrics, falling in the 78th percentile of

expansion estimates by Petitpierre et al. (2012, n = 50 plant species), and the 60th percentile of

Liu et al. (2020, n = 211 plant species with COUE estimates). Niche expansion was low

compared to results in Early & Sax (2014). These authors found much higher niche expansion

overall than Petitpierre et al., which they attributed life history differences between the study

species; Early & Sax considered endemic species with small native ranges while Petitpierre et al.

studied weedy species that were widely distributed in their native range, more similar to *D.*

graveolens.

We found little evidence for niche expansion in the invaded range. Australia and California

differed strongly in niche unfilling, illustrating the importance of invader residence time in

invasion risk assessment (Wilson et al. 2007). Although the California invasion appears to be in

much earlier stages with further dispersal expected, both invasive niches almost exclusively

cover climatic conditions that are also present in the native niche (high stability), a pattern

known as climate matching, which has been found for many successful invasive species (Hayes

and Barry 2008). Interestingly, despite its longer history, the Australian niche barely expands

beyond the native niche, suggesting niche conservatism. Incorporating the native range

expansion into projections of niche change and invasion risk adds newly invadable

climates/habitat especially in North America (Fig. 3). However, given that neither invasion has

yet spread into these areas, it remains an open question whether any similar niche evolution can

be expected in the invaded ranges in the future. The strong overlap in climate conditions

occupied in the historic native range and both exotic ranges does suggest that invasive

populations originated from a Mediterranean part of the native range.

<u>Limitations</u>

Any niche modeling study is constrained by the available data. Although the native and invaded

distributions of *D. graveolens* are relatively well-documented, occurrence records do contain

temporal and spatial biases that may have affected our results. Because the majority of our

occurrence data was reported in recent decades, we included all occurrences south of the original

range limit in the historic native range, regardless of the date they were reported. It is possible

that past climate conditions at these locations were less favorable for *D. graveolens*. However,

because our study is focused on a northward range expansion tracking climate change, and the

historic native range now represents the trailing edge of the distribution, our assumption to treat

all occurrences as historical presences should underestimate the niche shift during range

expansion and therefore represents a conservative approach. To address the spatial bias in our

data, we employed the target background record approach to reflect sampling effort across the

study area. This solution was a great improvement over randomly chosen background points

(which resulted in overfitting to climate areas with high sampling effort), but was still not

optimal given the low number of target background points available compared to Maxent's

standard of 10,000. However, at our spatial resolution (0.5°) even near 215,000 target records

from 296 species covered only 3468 cells across the entire study area (Supporting Information).

Higher-resolution climate data would allow for more background points (Merow et al. 2013) but

was not available for the past time period of interest.

Niche expansion as quantified in this study can represent either a change in the realized niche

due to lifted biotic or dispersal constraints, or be caused by evolution of the fundamental niche.

D. graveolens has very high spread potential due to a combination of an annual life history,

wind-dispersed seeds and the production of tens of thousands of seeds per plant. No historic

barriers to dispersal are apparent in the native range. Moreover, *D. graveolens* has a typical

ruderal life history and has expanded its native range primarily along roadsides, with limited

biotic interactions with other species. Finally, previous work has demonstrated that rapid

evolution of earlier phenology increased plant fitness in northern leading-edge populations

(Lustenhouwer et al. 2018). We therefore argue that evolution of the fundamental niche is the

most plausible explanation for the observed climate niche expansion.

Broader implications

Our finding of greater niche expansion during native range expansion than during invasion

contradicts hypotheses in the literature that niche stasis should be more pronounced in the native

range (Pearman et al. 2008, Wiens et al. 2019). Rather than having a maladaptive swamping

effect, gene flow in the native range brought in genetic variation from the northern end of

existing latitudinal clines that was adaptive in northern climates (Lustenhouwer et al. 2018). In

order to definitively assess the evolutionary potential of *D. graveolens* in California and predict

whether this invasion will proceed into areas forecasted by the Present Model (Fig. 3c), empirical

studies comparing plants of native and invasive origin will be necessary. Nonetheless, our study

supports calls to limit multiple introductions of invasive species, even if they are widespread

already (Smith et al. 2020), because the introduction of genotypes from northern Europe to

California would put new areas at risk of invasion (Fig. 3e,f).

In conclusion, our results suggest that climate change may act as a catalyst for range expansion

and subsequent climate niche evolution in plants. The generality of this phenomenon for other

species will depend on their evolutionary potential (generation time, heritable genetic diversity),

range-limiting factors, and dispersal ability (Catullo et al. 2015). Recent work on invertebrates

suggests that range expansion can promote increased thermal and diet niche breadth at the range

edge in many species (Lancaster 2016, 2020). We encourage future studies validating native

range shift projections with observed spread (e.g., Araújo et al. 2005), to examine whether

climate niche evolution during native range expansion is common and causes populations to

spread further than expected under climate tracking. Ultimately, the study of both biological

invasions and native species threatened by climate change will benefit from a better

understanding of the drivers of niche evolution during range expansion.

Acknowledgements

We are grateful to Zarina Pringle for her assistance in the occurrence data collection. We also

thank A. Pliszko, M. Kaligarič, G. Király, P. Eliáš, and D. Schmidt for pointing us to occurrence

records in Poland, Slovenia, Slovakia, and Hungary. Finally, we thank the Botanical Society of

Britain and Ireland for providing data from the UK. Members of the Parker and Gilbert labs

provided helpful comments at several points in the study.

References

Aitken, S. N. et al. 2008. Adaptation, migration or extirpation: climate change outcomes for tree
populations. - Evol. Appl. 1: 95–111.

Alexander, J. M. and Edwards, P. J. 2010. Limits to the niche and range margins of alien species.
- Oikos 119: 1377–1386.

Araújo, M. B. et al. 2005. Validation of species–climate impact models under climate change. -
Glob. Change Biol. 11: 1504–1513.

Araújo, M. B. et al. 2011. Climate change threatens European conservation areas. - Ecol. Lett.
14: 484–492.

Atwater, D. Z. et al. 2018. Climatic niche shifts are common in introduced plants. - Nat. Ecol.
Evol. 2: 34–43.

Boheemen, L. A. van et al. 2019. Rapid and repeated local adaptation to climate in an invasive
plant. - New Phytol. 222: 614–627.

Bonnier, G. and Layens, G. 1909. Flore complète de la France et de la Suisse (comprenant aussi
toutes les espèces de Belgique, d'Alsace et de Lorraine). - Libraire générale de
l'enseignement.

Broennimann, O. et al. 2012. Measuring ecological niche overlap from occurrence and spatial
environmental data. - Glob. Ecol. Biogeogr. 21: 481–497.

Broennimann, O. et al. 2020. ecospat: Spatial Ecology Miscellaneous Methods. R package
version 3.1.

Brownsey, R. et al. 2013a. Seed and Germination Biology of Dittrichia graveolens (Stinkwort). -
Invasive Plant Sci. Manag. 6: 371–380.

Brownsey, R. et al. 2013b. Stinkwort is rapidly expanding its range in California. - Calif. Agric.
67: 110–115.

Brullo, S. and De Marco, G. 2000. Taxonomical revision of the genus *Dittrichia* (Asteraceae). -
Port. Acta Biol. 19: 341–354.

BSBI 2020. Botanical Society of Britain and Ireland Distribution Database. Records for
Dittrichia graveolens accessed June 5, 2020.

Bush, A. et al. 2016. Incorporating evolutionary adaptation in species distribution modelling
reduces projected vulnerability to climate change. - Ecol. Lett. 19: 1468–1478.

Calflora 2021. Calflora: Information on California plants for education, research and
conservation [web application]. - The Calflora Database [a non-profit organization].

Catullo, R. A. et al. 2015. Extending spatial modelling of climate change responses beyond the
realized niche: estimating, and accommodating, physiological limits and adaptive
evolution. - Glob. Ecol. Biogeogr. 24: 1192–1202.

Chen, I.-C. et al. 2011. Rapid range shifts of species associated with high levels of climate
warming. - Science 333: 1024–1026.

Chuang, A. and Peterson, C. R. 2016. Expanding population edges: theories, traits, and trade-
offs. - Glob. Change Biol. 22: 494–512.

Ciardo, F. and Delarze, R. 2005. Observations floristiques sur l'autoroute lémanique: espèces
nouvelles, halophiles et rares. - Saussurea Traveaux Société Bot. Genève 35: 83–89.

Colautti, R. I. and Barrett, S. C. H. 2013. Rapid adaptation to climate facilitates range expansion
of an invasive plant. - Science 342: 364–366.

Colautti, R. I. and Lau, J. A. 2015. Contemporary evolution during invasion: evidence for
differentiation, natural selection, and local adaptation. - Mol. Ecol. 24: 1999–2017.

Corlett, R. T. and Westcott, D. A. 2013. Will plant movements keep up with climate change? -
Trends Ecol. Evol. 28: 482–488.

Coste, H. 1903. Flore descriptive et illustrée de la France, de la Corse, et des contrées
limitrophes. - P. Klincksieck.

Cox, B. 2001. The biogeographic regions reconsidered. - J. Biogeogr. 28: 511–523.

Di Cola, V. et al. 2017. ecospat: an R package to support spatial analyses and modeling of
species niches and distributions. - Ecography 40: 774–787.

Diamond, S. E. 2018. Contemporary climate-driven range shifts: Putting evolution back on the
table. - Funct. Ecol. 32: 1652–1665.

Dlugosch, K. M. and Parker, I. M. 2008a. Invading populations of an ornamental shrub show
rapid life history evolution despite genetic bottlenecks. - Ecol. Lett. 11: 701–709.

Dlugosch, K. M. and Parker, I. M. 2008b. Founding events in species invasions: genetic
variation, adaptive evolution, and the role of multiple introductions. - Mol. Ecol. 17: 431–
449.

Dudaniec, R. Y. et al. 2018. Signatures of local adaptation along environmental gradients in a
range-expanding damselfly (*Ischnura elegans*). - Mol. Ecol. 27: 2576–2593.

Early, R. and Sax, D. F. 2014. Climatic niche shifts between species' native and naturalized
ranges raise concern for ecological forecasts during invasions and climate change. - Glob.
Ecol. Biogeogr. 23: 1356–1365.

Elith, J. and Leathwick, J. R. 2009. Species Distribution Models: Ecological Explanation and
Prediction Across Space and Time. - Annu. Rev. Ecol. Evol. Syst. 40: 677–697.

Fick, S. E. and Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces
for global land areas. - Int. J. Climatol. 37: 4302–4315.

FloraWeb 2013. *Dittrichia graveolens* in FloraWeb: Daten und Informationen zu Wildpflanzen
und zur Vegetation Deutschlands.

GBIF.org 2020. GBIF Occurrence Download for *Dittrichia graveolens* (L.) Greuter,
https://doi.org/10.15468/dl.s0qcs0.

Grimmett, L. et al. 2020. Presence-only species distribution models are sensitive to sample
prevalence: Evaluating models using spatial prediction stability and accuracy metrics. -
Ecol. Model. 431: 109194.

2017. 6. Environmental Predictors: Issues of Processing and Selection. - In: Guisan, A. et al.
(eds), Habitat Suitability and Distribution Models: With Applications in R. Ecology,
Biodiversity and Conservation. Cambridge University Press, pp. 61–109.

Guisan, A. 2017. Habitat suitability and distribution models, with applications in R. - Cambridge
University Press.

Guisan, A. et al. 2014. Unifying niche shift studies: insights from biological invasions. - Trends
Ecol. Evol. 29: 260–269.

Harris, I. et al. 2020. Version 4 of the CRU TS monthly high-resolution gridded multivariate
climate dataset. - Sci. Data 7: 109.

Hayek, A. and Markgraf, F. 1931. Prodromus florae peninsulae balcanica / A. Hayek; F.
Markgraf ; 2: Dicotyledoneae sympetalae.

Hayes, K. R. and Barry, S. C. 2008. Are there any consistent predictors of invasion success? -
Biol. Invasions 10: 483–506.

Hijmans, R. J. et al. 2017. dismo: Species Distribution Modeling.

Hill, M. P. et al. 2013. A predicted niche shift corresponds with increased thermal resistance in
an invasive mite, Halotydeus destructor. - Glob. Ecol. Biogeogr. 22: 942–951.

Hoffmann, A. A. and Sgrò, C. M. 2011. Climate change and evolutionary adaptation. - Nature
470: 479–485.

iNaturalist 2020. Available from https://www.inaturalist.org.

Info Flora 2020. *Dittrichia graveolens* in: Info Flora, the National Data and Information Center
on the Swiss Flora. www.infoflora.ch.

Jakob, S. S. et al. 2010. Population demography influences climatic niche evolution: evidence
from diploid American Hordeum species (Poaceae). - Mol. Ecol. 19: 1423–1438.

Kearney, M. et al. 2009. Integrating biophysical models and evolutionary theory to predict
climatic impacts on species' ranges: the dengue mosquito *Aedes aegypti* in Australia. -
Funct. Ecol. 23: 528–538.

Kottler, E. J. et al. 2021. Draining the swamping hypothesis: little evidence that gene flow
reduces fitness at range edges. - Trends Ecol. Evol. 36: 533–544.

Kubisch, A. et al. 2013. Predicting range shifts under global change: the balance between local
adaptation and dispersal. - Ecography 36: 873–882.

Lancaster, L. T. 2016. Widespread range expansions shape latitudinal variation in insect thermal
limits. - Nat. Clim. Change 6: 618–621.

Lancaster, L. T. 2020. Host use diversification during range shifts shapes global variation in
Lepidopteran dietary breadth. - Nat. Ecol. Evol.: 1–7.

Lancaster, L. T. et al. 2015. Latitudinal shift in thermal niche breadth results from thermal
release during a climate-mediated range expansion. - J. Biogeogr. 42: 1953–1963.

Lee-Yaw, J. A. et al. 2019. Range shifts and local adaptation: integrating data and theory towards
a new understanding of species' distributions in the Anthropocene. - New Phytol. 221:
644–647.

Lenoir, J. et al. 2020. Species better track climate warming in the oceans than on land. - Nat.
Ecol. Evol.: 1–16.

Liu, C. et al. 2020. Most invasive species largely conserve their climatic niche. - Proc. Natl.
Acad. Sci. 117: 23643–23651.

Lustenhouwer, N. et al. 2018. Rapid evolution of phenology during range expansion with recent
climate change. - Glob. Change Biol. 24: e534–e544.

Martínez-Meyer, E. et al. 2004. Ecological niches as stable distributional constraints on mammal
species, with implications for Pleistocene extinctions and climate change projections for
biodiversity. - Glob. Ecol. Biogeogr. 13: 305–314.

Merow, C. et al. 2013. A practical guide to MaxEnt for modeling species' distributions: what it
does, and why inputs and settings matter. - Ecography 36: 1058–1069.

Moran, E. V. and Alexander, J. M. 2014. Evolutionary responses to global change: lessons from
invasive species. - Ecol. Lett. 17: 637–649.

Nadeau, C. P. and Urban, M. C. 2019. Eco-evolution on the edge during climate change. -
Ecography in press.

Naimi, B. et al. 2014. Where is positional uncertainty a problem for species distribution
modelling? - Ecography 37: 191–203.

Nikolić, T. 2015. Flora Croatica Database, http://hirc.botanic.hr/fcd.

Nogués-Bravo, D. et al. 2018. Cracking the Code of Biodiversity Responses to Past Climate
Change. - Trends Ecol. Evol. 33: 765–776.

Pardi, M. I. et al. 2020. Testing climate tracking of montane rodent distributions over the past
century within the Great Basin ecoregion. - Glob. Ecol. Conserv. 24: e01238.

Parent, G. H. 2011. Etudes ecologiques et chorologiques sur la flore Lorraine. Note 26: quelques
taxons nouveaux ou meconnus de Lorraine française. Donnees rassemblees de 1993 a
2005. - Inst. Grand-ducal Luxemb. Sect. Sci. Nat. Phys. Mathématiques Nouv. Sér. 46:
5–40.

Parsons, W. T. and Cuthbertson, E. G. 2001. Stinkwort. - In: Noxious Weeds of Australia.
Second Edition. Csiro Publishing, pp. 281–283.

Pearman, P. B. et al. 2008. Niche dynamics in space and time. - Trends Ecol. Evol. 23: 149–158.

Petitpierre, B. et al. 2012. Climatic niche shifts are rare among terrestrial plant invaders. -
Science 335: 1344–1348.

Phillips, S. 2017. Maxnet: fitting "maxent" species distribution models with "glmnet". R package
version 0.1.2.

Phillips, S. J. et al. 2006. Maximum entropy modeling of species geographic distributions. -
Ecol. Model. 190: 231–259.

Phillips, S. J. et al. 2009. Sample selection bias and presence-only distribution models:
implications for background and pseudo-absence data. - Ecol. Appl. 19: 181–197.

Phillips, S. J. et al. 2017. Opening the black box: an open-source release of Maxent. - Ecography
40: 887–893.

Polechová, J. and Barton, N. H. 2015. Limits to adaptation along environmental gradients. -
Proc. Natl. Acad. Sci. 112: 6401–6406.

Preston, R. E. 1997. *Dittrichia graveolens* (Asteraceae), new to the California weed flora. -
Madroño 44: 200–203.

QGIS Development Team 2019. QGIS Geographic Information System. - Open Source
Geospatial Foundation.

R Core Team 2020. R: A language and environment for statistical computing. - R Foundation for
Statistical Computing.

Rios, N. E. 2020. GEOLocate. A platform for georeferencing natural history collections data. https://www.geo-locate.org/.

Rouy, G. 1903. XLIV. Rubiacées - XLVIII. Composées. - In: Flore de France, ou description des plantes qui croissent spontanément en France, en Corse et en Alsace-Lorraine. Deslis, pp. 205–206.

Saikkonen, K. et al. 2012. Climate change-driven species' range shifts filtered by photoperiodism. - Nat. Clim. Change 2: 239–242.

Santilli, L. et al. 2021. First record of *Dittrichia graveolens* (Asteraceae, Inuleae) in Chile. - Darwiniana Nueva Ser.: 31–38.

Schoener, T. W. 1970. Nonsynchronous Spatial Overlap of Lizards in Patchy Habitats. - Ecology 51: 408–418.

Smith, A. B. 2020a. Best practices in species distribution modeling: a workshop in R. Available at http://www.earthskysea.org/.

Smith, A. B. 2020b. enmSdm: Tools for modeling niches and distributions of species. R package version 0.5.2.9.

Smith, A. L. et al. 2020. Global gene flow releases invasive plants from environmental constraints on genetic diversity. - Proc. Natl. Acad. Sci. 117: 4218–4227.

Spence, A. R. and Tingley, M. W. 2020. The challenge of novel abiotic conditions for species undergoing climate-induced range shifts. - Ecography in press.

Taulavuori, K. et al. 2010. Growth responses of trees to arctic light environment. - In: Lüttge, U. et al. (eds), Progress in Botany 71. Progress in Botany. Springer, pp. 157–168.

Tutin, T. G. et al. 1976. Flora Europaea. Volume 4. Plantaginaceae to Compositae (and Rubiaceae). - Cambridge University Press.

Urban, M. C. et al. 2016. Improving the forecast for biodiversity under climate change. - Science 353: aad8466.

USDA 2013. Weed Risk Assessment for *Dittrichia graveolens* (L.) Greuter (Asteraceae) – Stinkwort.

Visser, M. E. 2008. Keeping up with a warming world; assessing the rate of adaptation to climate change. - Proc. R. Soc. B Biol. Sci. 275: 649–659.

2021. *Dittrichia graveolens* (E von Raab-Straube, Ed.). - EuroMed PlantBase

Wagenitz, G. 1966. Compositae I: Allgemeiner Teil, Eupatorium - Achillea. - In: Illustrierte Flora von Mitteleuropa. Parey, in press.

Walther, G.-R. et al. 2005. An ecological 'footprint' of climate change. - Proc. R. Soc. B Biol.
Sci. 272: 1427–1432.

Warren, D. L. et al. 2008. Environmental Niche Equivalency Versus Conservatism: Quantitative
Approaches to Niche Evolution. - Evolution 62: 2868–2883.

Wiens, J. J. et al. 2019. Rapid niche shifts in introduced species can be a million times faster than
changes among native species and ten times faster than climate change. - J. Biogeogr. 46:
2115–2125.

Wilson, J. R. U. et al. 2007. Residence time and potential range: crucial considerations in
modelling plant invasions. - Divers. Distrib. 13: 11–22.

Zając, A. and Zając, M. 2019. Distribution atlas of vascular plants in Poland: appendix. -
Institute of Botany, Jagiellonian University.

  Figures

Figure 1 Niche shift during native range expansion, comparing the past (1901-1930) and present

(1990-2019) datasets; (a) past and present niche in environmental space, with colors indicating

niche stability (purple), unfilling (orange) and expansion (blue), and the white arrow showing the

niche centroid shift. Dark shading shows the density of occurrences in the historic range. Lines

represent the climate of the Eurasian Holarctic in the past (orange) and present (blue), using

100% (solid) and 90% (dashed) of available climates. The black arrow indicates the climate

centroid shift between past and present. (b) principal component analysis of the entire

environmental space, with axes explaining 41.6% (x) and 29.0% (y) of variation. Arrows show

environmental predictors as defined in Table 1. (c, d) niche equivalency and niche similarity test,

with histograms showing null distributions and diamonds marking observed niche overlap.

Figure 2 Maxent habitat suitability results for (a) the Past Model (fit to the past climate and

historic native range occurrences) projected onto the past climate (1901-1930); (b) the Past

Model projected onto the present climate (1990-2019), indicating expected range expansion with

climate change; (c) the same projection with observed occurrences in the present, and (d) the

Present Model (fit to the present climate and all occurrences) projected onto the present climate.

Historic native range limit represented by the black line, and species occurrence records in the

historic (a,b) and expanded (c,d) native range by dots. Axes display degrees longitude (x) and

latitude (y). Projections for the entire study area available in the Supporting Information.

663

664   **Figure 3** Niche shift during invasion in Australia (a,b) and California (d,e), using the past native

665   (a,d) and present native (b,e) niche and climate as a reference. Colors indicate niche stability,

666   unfilling and expansion as in Fig. 1, with lines representing the climate of the native (orange) and

667   exotic (blue) study area. Shading shows the density of occurrences from the historic (a,d) and

668   expanded (b,e) native range in environmental space. Panels c and f show areas at risk of invasion

669   as projected by the Past and Present Model, with cells shaded from dark to light as follows: cell

670   included in the Past Model only, in both models, in the Present Model only, or in neither model.

671   Points indicate observation records for *D. graveolens* in Australia (c) and California (f).

672

673 **Tables**

674 **Table 1** Environmental predictors included in this study, with minimum (min), median (med)

675 and maximum (max) values across all pixels in each study area. Data from CRU TS4.04.

Predictor		Eurasian Holarctic (1901-1930)			Eurasian Holarctic (1990-2019)			North America			Australia		
		min	med	max	min	med	max	min	med	max	min	med	max
bio2	mean diurnal range (°C)	4.2	10.5	18.0	4.4	10.3	18.1	4.6	10.5	20.8	6.7	14.4	16.5
bio4	temperature seasonality (100xSD,°C)	289	1288	2448	295	1255	2365	108	1171	1705	113	545	679
bio15	precipitation seasonality (CV, mm)	21	61	173	23	63	176	19	63	183	21	94	162
bio17	precipitation driest quarter (mm)	0	34	470	0	35	503	0	44	403	0	18	236
bio18	precipitation warmest quarter (mm)	0	163	1383	0	169	1248	1	182	930	24	140	1032
frost	frost days Sep-Dec (count)	0	85	180	0	79	177	0	91	174	0	4	82

676 **Table 2** Niche shift metrics following the COUE framework, calculated using the intersection of

677 the 90^{th} percentile of environmental conditions in each range. Niche equivalency and similarity

678 tests were one-sided, with H_1 indicating the alternative hypothesis used.

Comparison		Niche shift metrics				Niche equivalency test		Niche similarity test	
niche 1	niche 2	D	expansion	stability	unfilling	H_1	P	H_1	P
past native	present native	0.71	0.05	0.95	0	lower	0.01**	greater	0.01**
past native	Australia	0.60	0	1.00	0.06	greater	0.01**	greater	0.12
present native	Australia	0.53	0	1.00	0.11	greater	0.01**	greater	0.07
past native	California	0.05	0.01	0.99	0.93	lower	0.01**	greater	0.31
present native	California	0.04	0.03	0.97	0.94	lower	0.01**	greater	0.24

680 **Table 3** Maxent model performance for projections of the Past Model in both time periods, and

681 the projection of the Present Model in the present. All presence and background points were

682 allocated randomly to training and test datasets using 5 k-folds for cross-validation (table shows

683 points per fold). Mean and standard deviation of AUC and CBI are given across the 5 k-folds.

Climate and occurrence data model	projection	n presences training	test	n background points training	test	AUC mean	SD	CBI mean	SD
Past Model	past dataset	319	80	2774	694	0.89	0.01	0.85	0.11
Past Model	present dataset	319	149	2774	694	0.76	0.01	0.61	0.14
Present Model	present dataset	597	149	2774	694	0.81	0.01	0.93	0.04