TRβ Agonism Induces Tumor Suppression and Enhances Drug Efficacy in Anaplastic Thyroid Cancer in Female Mice

Noelle E. Gillis[1,2,3], Lauren M. Cozzens*[1,4]*, Emily R. Wilson[1,5], Noah M. Smith[1], Jennifer A. Tomczak[1], Eric L. Bolf*[1,2]*, Frances E. Carr[1,2]**

[1] Department of Pharmacology, Larner College of Medicine, University of Vermont, Burlington, Vermont
[2] University of Vermont Cancer Center, University of Vermont, Burlington, Vermont
[3] Current address: Masonic Cancer Center, University of Minnesota, Minneapolis, Minnesota
[4] Current address: University of Colorado, Anschutz Medical Campus, Aurora, Colorado
[5] Current address: University of Utah, Salt Lake City, Utah

** *Corresponding author,* Email: frances.carr@med.uvm.edu

Short title: TRβ agonist blunts tumor progression

Keywords: GC-1, Sobetirome, thyroid hormone receptor, anaplastic thyroid cancer

Noelle E. Gillis, gilli431@umn.edu; University of Minnesota, Minneapolis MN, 55455 ORCID: 0000-0002-0005-4194

Lauren M. Cozzens, lauren.cozzens@cuanschutz.edu; University of Colorado at Anschutz, Aurora, CO 80045

Emily R. Wilson, emily.r.wilson@hci.utah.edu, University of Utah, Salt Lake City, UT 84116

Noah M. Smith, noah.smith@uvm.edu, University of Vermont, Burlington VT 05405

Jennifer A. Tomczak, jennifer.tomczak@uvm.edu; University of Vermont, Larner College of Medicine, Burlington, VT, 05405

Eric L. Bolf, eric.bolf@2seventybio.com, Cambridge MA 02142

Frances E. Carr, frances.carr@med.uvm.edu; University of Vermont, Larner College of Medicine, Burlington, VT, 05405   ORCID: 0000-0001-7266-8788

Word Count: 4990

ABSTRACT

Thyroid hormone receptor beta (TRβ) is a recognized tumor suppressor in numerous solid cancers. The molecular signaling of TRβ has been elucidated in several cancer types through re-expression models. Remarkably, the potential impact of selective activation of endogenous TRβ on tumor progression remains largely unexplored. We used cell-based and *in vivo* assays to evaluate the effects of the TRβ agonist Sobetirome (GC-1) on a particularly aggressive and dedifferentiated cancer, anaplastic thyroid cancer (ATC). Here we report that GC-1 reduced the tumorigenic phenotype, decreased cancer stem-like cell populations, and induced re-differentiation of the ATC cell lines with different mutational backgrounds. Of note, this selective activation of TRβ amplified the effects of therapeutic agents in blunting the aggressive cell phenotype and stem-cell growth. In xenograft assays, GC-1 alone inhibited tumor growth and was as effective as the kinase inhibitor, Sorafenib. These results indicate that selective activation of TRβ not only induces a tumor suppression program *de novo* but enhances the effectiveness of anti-cancer agents revealing potential novel combination therapies for ATC and other aggressive solid tumors.

INTRODUCTION

There is compelling evidence that the nuclear thyroid hormone receptor beta (TRβ) blunts tumor progression, invasion, and metastases in multiple solid tumors including thyroid, breast, colon and other cancers[1]. Whereas diminished expression and/or function of TRβ is often characteristic of advanced and aggressive cancers, cell-based and animal studies reveal that liganded re-expressed TRβ induces apoptosis, reduces an aggressive phenotype, decreases cancer stem cell populations, and slows tumor growth through modulation of a complex transcriptional network[2-9]. Transcriptional regulation by TRβ is critical for its function as a tumor suppressor because it also acts as both a signal transducer and facilitator of long-term epigenetic programming for maintenance of cell identity. Our recent studies revealed that TRβ in the presence of triiodothyronine (T_3) induced a tumor-suppressive transcriptomic program and cell redifferentiation in aggressive thyroid and breast cancer cells[2,7]. We identified several key mechanisms by which activation of TRβ could reduce tumor progression, implicating thyroid hormone signaling as a therapeutic opportunity in advanced cancers. Recent studies in breast [8], colon[10], and liver[5] cancers also emphasize this untapped therapeutic potential[11,12]. Despite the potential clinical benefits of TRβ tumor suppressor activity implicated by numerous cell-based and animal studies, TRβ selective agonists remain largely untested as cancer therapeutics as single agents or in combination with other treatments. Novel treatments are a critical need in the context of aggressive poorly differentiated tumors, such as anaplastic thyroid cancer (ATC), for which there are limited targeted therapies and thus far, no curative options.

As noted in ATC and other endocrine-related cancers, activation of

phosphatidylinositol 3-kinase (PI3K)–Akt–mammalian target of rapamycin (mTOR)

pathway (PI3K-Akt) and mitogen-activated protein kinase (MAPK) signaling are

recognized drivers of thyroid oncogenesis and blocking this activity has been a

therapeutic focus. Yet inhibitors of the major components of these and other signaling

cascades rarely provide a durable response as resistance quickly develops[13]. Current

molecular diagnostics provide a rationale for intervention with combination therapies but

thus far have not yielded significantly improved responses and potential use of

thyromimetics not systematically evaluated[14] [13,15,16]. The therapuetic use of non-isoform

selective thyromimetics is problematic due to the side-effects of hyperthyroidism and the

potential for adverse cardiovascular events that result from aberrant TRα activity.

To subvert these negative side effects, TRβ-selective thyromimetics have been

developed[17]. Several of these drugs have been shown to be effective for treatment of

metabolic and neurodegenerative disorders in clinical trials[17,18]. The TRβ agonist

Sobetirome (GC-1) has been extensively characterized and shown to preferentially bind

to and activate TRβ rather than TRα[19-22]. Thus, given our prior work demonstrating that

restoration of TRβ expression reduces the aggressive ATC phenotype[2,7,23], we here

tested the possibility that selective activation of endogenous TRβ with GC-1 might

reduce the tumor cell phenotype and inhibit tumor growth. In ATC cells with diverse

genetic backgrounds [24] and varying endogenous TRβ levels, GC-1 alone decreased the

aggressive phenotype, reduced cancer stem cell growth and increased the

effectiveness of MAPK, PI3K, and cell cycle inhibitors. GC-1 significantly increased

expression of thyroid-specific genes, inducing re-differentiation in ATC cells.

Importantly, we observed an increase in NIS protein and cellular iodide uptake,

implicating potential restoration of functionality. In a xenograft model, GC-1 reduced

tumor growth as effectively as a MAPK pathway inhibitor, Sorafenib. The combination of

GC-1 and Sorafenib maximally suppressed tumor growth. These observations establish

the foundation that activation of endogenous TRβ with isoform-selective agonists may

be an effective and practical adjuvant therapeutic strategy.

MATERIALS AND METHODS

Culture of thyroid cell lines. ATC cell lines were cultured in RPMI 1640 growth media

with L-glutamine (300 mg/L), sodium pyruvate and nonessential amino acids (1%)

(Corning), supplemented with 10% fetal bovine serum (Peak Serum) and penicillin-

streptomycin (200 IU/L) (Corning) at 37°C, 5% CO_2, and 100% humidity. Charcoal-

stripped fetal bovine serum (Sigma) was used for hormone induced gene expression

analysis. SW1736-EV and SW-TRβ cells were modified by lentiviral transduction as

recently described[2]. SW1736 and KTC-2 were authenticated by the Vermont Integrative

Genomics Resource at UVM using short tandem repeat profiles (SW1736, May 2019;

KTC-2, October 2019). 8505C and OCUT2 were authenticated by the U. Colorado using

short tandem repeat profiles (8505C, June 2013; OCUT-2, June 2018).

Pharmacological agents. T_3 (Sigma-Aldrich) was suspended in 1 M NaOH and GC-1

100% ethanol for stock concentrations of 1 mM before further dilution in culture

medium. Buparlisib, Sorafenib, Palbociclib, and Alpelisib (MedChemExpress) were

suspended in 100% DMSO for stock concentrations of 50-150 mM before further

dilution in culture medium.

Cell Growth and Viability Assays. Cell growth was measured by cell counting at

discrete time points. Cells were seeded in 12- well plates, then treated with GC-1,

Buparlisib, Alpelisib, Palbociclib, or Sorafenib to establish time and concentration effects

of therapeutics on cell growth. Cell viability was determined by a Sulforhodamine B

assay (Abcam) following the manufacturer's protocol. In brief, ATC cells were plated in

96-well clear flat-bottom plates at a density of 5,000 cells per well. Cells were fixed,

stained, and imaged using a plate reader according to manufacturer's instructions.

Migration Assay. Cell migration was determined by wound healing assay as previously

described[2,23]. Media was supplemented with GC-1 (10 nM) with and without 0.5 μM

Buparlisib, 0.5 μM Alpelisib, 1 nM Palbociclib, or 5 μM Sorafenib. Images were obtained

at 0, 16, 24, 48, and 72 hours or until 100% wound closure, depending on the cell line.

Percent closure was calculated relative to the area of the initial scratch.

Tumorsphere Assay. Tumorspheres formed from ATC cells were used to assess self-

renewal and sphere-forming efficiency as previously described[2]. Where indicated,

adherent cells were treated with 0.5 μM Buparlisib, 0.5 μM Alpelisib, 1 nM Palbociclib, 5

μM Sorafenib, or vehicle for 72 hours prior to tumorsphere-forming assay.

Tumorspheres were then cultured with or without GC-1 (10 nM) to evaluate the effects

of liganded TRβ on thyrosphere growth alone or after treatment with a therapeutic

agent.

Western Blot Analysis. Proteins were isolated from whole cells in lysis buffer and

visualized by Western blot as previously described[2,23,25]. Specific proteins were

detected with the indicated antibodies (Table 1); immunoreactive proteins were detected

by enhanced chemiluminescence (Thermo Fisher Scientific) on a ChemiDoc XRS+ (Bio-

Rad Laboratories).

RNA Extraction and Quantitative Real-Time PCR (qRT-PCR). Total RNA was

extracted using RNeasy Plus Kit (Qiagen) according to manufacturer's protocol. cDNA

was generated using 5X LunaScript RT SuperMix, and mRNA expression was

quantified by qRT-PCR using 2X Luna Universal qPCR Master Mix (New England

Biolabs) on a QuantStudio 3 real-time PCR system (Applied Biosystems). Fold change

in gene expression compared to housekeeping controls was calculated using the ddCT

method. Primer sequences are indicated in Table 2.

Iodide Uptake Assay. Cells were seeded at a density of 40,000 cells per well in a 96

well plate with or without 10 nM GC-1 for 48 hours. Media was discarded, and cells

were washed four times with warm PBS before incubation with 1 μM potassium iodide

(KI, Sigma-Aldrich) in PBS for 1 hour at 37°C. Wells were washed an additional four

times with warm PBS and 100 μL of ddH$_2$O were added to each well. 100 μl of 24 mM

ammonium cerium (IV) sulfate hydrate with 5.7% concentrated H$_2$SO$_4$ (Sigma-Aldrich)

and 100 μl of 24 mM sodium arsenite (III) with 205 mM sodium chloride (NaCl) and 50

mM sodium hydroxide (NaOH, Sigma-Aldrich) were added to each well. Plates were

incubated in the dark at room temperature for 30 minutes, and absorbance at 415 nm

was measured with a Synergy 2 Multi-Detection Microplate Reader (Agilent

Technologies).

***In Vivo* Evaluation of Sobetirome (GC-1) and Sorafenib.** The xenograft experiment

was approved by the Animal Care and Use Committee of the University of Vermont

(Protocol X0-018). Four-week-old athymic female nude mice (outbred homozygous

nude *Foxn1^{nu}/Foxn1^{nu}*) were purchased from Jackson Laboratory (Bar Harbor, ME,

USA) and were allowed to acclimatize for one week. Mice were given food and water *ad*

libitum. Mice were anesthetized using 3% isoflurane delivered at 1.5 L/min, and tumors

were established by subcutaneously injecting 1×10^6 8505C cells with a 26G needle in

100 μL of high concentration Matrigel (Corning Inc.) diluted 1:2 with base RPMI-1640

into each flank of 24 mice. We chose 8505C cells based on the rapid growth

characteristics *in vitro* observed in our own studies[26] and based on their ability to form

large tumors in *in vivo* studies by other groups[27-30]. One week later, mice were sorted

into four treatment groups to achieve approximately equal body weights within each

group. Mice were then administered 0.3 mg/kg GC-1, 10 mg/kg Sorafenib, both GC-1

and Sorafenib, or vehicle control every day via intraperitoneal injection using 27G

needles. Pharmacological agents were dissolved in 100% DMSO and diluted daily in

30% DMSO, 40% PEG300, and 30% PBS and incubated at 55°C for 10 minutes prior to

vortexing to encourage complete solubilization. Tumor dimensions were measured with

digital calipers, and the volumes were calculated by the following formula: $(\Pi \times a \times b^2) / $

6, where a represents the largest diameter and b is the perpendicular diameter. The

body weight of each animal was taken at least twice a week to monitor toxicity. After

treatment for 16 days, mice were euthanized with carbon dioxide, and the tumors were

harvested, fixed with formalin (Thermo Fisher Scientific), and stored at 4°C prior to slide

sectioning and immunohistochemistry.

Immunofluorescent Analysis of Tumor Tissues. Formalin fixed paraffin embedded

human thyroid xenograft tumor blocks were sectioned at 5 μm thickness onto glass

microscope slides, dried overnight, and then baked at 60°C for one hour. Sections were

deparaffinized in three changes of xylene for 15 minutes each, then dehydrated in a

series of graded ethanols (100% x 2, 95% x 2, 70%, 50%, water) for 5 minutes each.

Antigen retrieval utilizing 1X DAKO target Retrieval Solution (#S1699, lot# 11270951;

Agilent Technologies, Santa Clara, CA) in 50% glycerol was performed in a Decloaking

Chamber (SP1 - 100°C for 15 min, SP2 - 90°C for 1 min). Following antigen retrieval,

slides were cooled to room temperature in the antigen retrieval solution for 20 minutes.

Slides were then rinsed 2x 5 minutes in distilled water followed by 0.2% Tween-20 for

an additional 5 minutes. Tissue sections were blocked for non-specific antibody

reactivity by incubation with 10% normal goat serum diluted in phosphate buffered

saline (PBS)/ 3% bovine serum albumin (BSA)/ 0.3% Triton X-100 for 60 minutes at

room temperature. Immediately following blocking, slides were incubated with rabbit

polyclonal anti-Ki67 (Abcam, Cambridge, UK; #ab15580) at a concentration 1.4 µg/ml in

PBS/3.0% BSA/0.3% Triton X-100 overnight at 4°C. Slides were then rinsed 7x 5

minutes in PBS/3.0% BSA and incubated with goat anti-rabbit Alexa 488 conjugated

secondary antibody (ThermoFisher, #A11008) in PBS/3.0% BSA for 60 minutes at room

temperature. Staining was followed by 5 x 5 minute rinses with PBS/3.0% BSA.

Samples were counterstained with 4', 6-diamidino-2-phenylindole (DAPI)

(ThermoFisher, #D1306) at a concentration of 10 µg/ml in PBS / 3.0%BSA for 15

minutes then rinsed 3x 5 minutes in distilled water. Following rinses slides were

mounted with a #1.5 coverslip and DAKO IF mounting media (Agilent; #S3023) for

imaging.

Confocal Imaging of Human Thyroid Xenograft Tissue. Xenograft samples stained with anti-KI-67 and DAPI were imaged on a Nikon A1R HD point scanning confocal microscope with a 20x Plan Apo λ DIC objective lens (NA 0.75, WD 1000 μm) with 488 nm and 405 nm laser excitation respectively. Image acquisition was completed in galvanometic scanning mode (Nikon Instruments Inc., Melville NY). Images of the entire xenograft section were acquired at 12-bit resolution using the tile scan function. Images were saved in ND2 file format in NIS Elements (version 5.2.1, Nikon, Tokyo, Japan).

Indica Labs HALO Image Analysis. Images were analyzed using HALO image analysis (Indica Labs, Albuquerque NM; version 3.3.45) utilizing the HighPlex FL module (version 4.1.3). Regions of interest containing only tumor tissue were outlined on each tiled image. Nuclear detection was set to the DAPI channel and the nuclear detection settings were optimized to accurately segment nuclei in the images. Membrane and cytoplasmic threshold settings were optimized for Ki-67 staining in the Alexa 488 (green) channel. Cytoplasmic detection settings were set to 4095 (in a 12 bit image) so that no cytoplasmic background staining would be included in the analysis. Nuclear positive signal setting threshold was set per image, to most accurately reflect the antibody staining level observed in the image. Percent Ki-67 positive cells were summarized.

Ethics Statement and Animal Modeling. All animal procedures were approved by Institutional Animal Care and Use Committee (IACUC) at UVM. All animals were maintained in pathogen-free conditions and cared for in accordance with IACUC policies and certification. All surgeries were performed with isoflurane anesthesia. Temperature-controlled post-surgical monitoring was implemented to minimize suffering. Mice

carrying anaplastic thyroid tumors were euthanized at designated time points for tumor

collection. We used signs of ulceration or a maximum individual tumor size of 2000 mm^3

as a protocol-enforced endpoint.

Statistics. All statistical analyses were performed using GraphPad Prism 9.3.1 and

variance assumed similar between experimental groups. Paired comparisons were

analyzed by unpaired Student's t-test assuming two-tailed distribution. Group

comparisons were made by one-way ANOVA followed by Sidak's or Tukey's multiple

comparison test as appropriate. Multigroup analyses was by two-way ANOVA followed

by a Tukey's multiple comparison test. Data are represented as mean ± standard

deviation or standard error of the mean where indicated. Differences were considered

statistically significant at p ≤ 0.05. Area under the curve (AUC) at the 95th confidence

interval was used to evaluate statistical differences in growth and migration assays.

RESULTS

GC-1 blocks the tumorigenic phenotype of ATC cells transduced with TRβ.

We previously demonstrated that T_3 (10 nM) treatment of transduced SW1736

cells, in which TRβ is re-expressed (SW-TRβ), induced a tumor suppression

transcriptomic program and reduced cell growth when compared to cells transduced

with empty vector (SW-EV) [2]. We therefore tested the ability of GC-1 (10 nM) to reduce

ATC phenotypes. We found that GC-1 similarly decreased cell growth in four days (**Fig.**

1A). There was a 50% reduction in the overall growth rate of SW-EV cells and 75%

reduction in SW-TRβ cells (**Fig. 1B**), indicating potent activation of endogenous TRβ.

There was also a modest decrease in the cell migration rate upon treatment with GC-1

(**Fig. 1C**).

As cancer stem cells are thought to be responsible for tumor initiation and

recurrence, targeting these cells is key to successful treatment. Therefore, we tested

whether thyroid cancer stem cells are responsive to GC-1. Thyrosphere formation is

robust in SW-EV cells, and GC-1 induced a significant reduction in growth (**Fig. 1D**).

Re-expression of TRβ (SW-TRβ) reduces thyrosphere growth and addition of GC-1

reduces thyrosphere formation by approximately 80% (**Fig. 1D**). These results indicate

that GC-1 can activate TRβ to reduce the thyroid cancer stem cell population.

Apoptotic signaling is central to ligand-activated TRβ modulation of the ATC

aggressive phenotype [2]. Thus, we assessed, by immunoblot, the effects of GC-1 on

PARP and caspase 3 cleavage as markers of late-stage apoptosis (**Fig. 1E**). GC-1

significantly increased cleavage of PARP and caspase 3 in SW-TRβ consonant with

increased apoptotic signaling. Combined, these results indicate that GC-1 can activate

TRβ to reduce the aggressive phenotype in transduced ATC cells with TRβ re-

expressed and recapitulate effects seen with T_3 at comparable concentrations.

GC-1 slows growth of parental ATC cells.

Once we confirmed that GC-1 induced the same phenotypic changes as in our

transduced ATC cell line model, it was critical to assess whether selective activation of

endogenous TRβ could induce similar effects in unmodified ATC cell lines [24]. SW1736,

8505C, OCUT2, and KTC-2 cells all harbor *BRAF*[V600E] and *TERT* driver mutations, but

otherwise have diverse mutational backgrounds (summarized in **Fig. 2A**). Treatment of

each cell line with 10 nM GC-1 for four days decreased cell growth whereas treatment

with 10 nM T_3 did not induce a significant change (**Fig. 2B**). Our prior work confirmed

that endogenous TRβ expression is low but detectable in the ATC cell lines when

compared with normal thyroid cells [23]. In the present study, GC-1 but not T3 after 24 hr

treatment significantly increased TRβ gene expression in SW1736, 8505C, OCUT2 cells

(**Fig. 2C**). Neither GC-1 nor T_3 treatment of cells significantly altered TRβ protein levels

during that time period (**Fig. 2D,E).** Thus, GC-1 but not T_3 may selectively enhance

TRβ gene expression and contribute to stable protein levels in ATC cells.

GC-1 increases the efficacy of therapeutic drugs on cell viability.

Based on our previous work on activation of TRβ in ATC cells, we hypothesized that

GC-1 could enhance the efficacy of therapeutics that target selective pathways. Thus,

we chose representative drugs that targeted tumorigenic cell signaling including PI3K,

MAPK, and cell-cycle modulated by TRβ. Cell viability was calculated following

treatment for three days with increasing concentrations of the PI3K inhibitor Buparlisib

(pan-PI3K inhibitor) or Alpelisib (PI3K☐ mutant inhibitor) for OCUT2 cells which harbor a

gain-of-function *PIK3CA* mutation (**Fig. 3A**), the cell cycle inhibitor Palbociclib (**Fig. 3B**),

or the multi-tyrosine kinase inhibitor Sorafenib (**Fig. 3C**), with or without 10 nM GC-1.

Each therapeutic agent decreased cell viability in three days in a dose-dependent

manner. The addition of GC-1 significantly increased the efficacy of these agents at

lower concentrations. For example, 1 μM Buparlisib was required to reduce cell viability

below 50% in the absence of GC-1, but when combined with 10 nM GC-1 this could be

achieved with 0.1-0.5 μM Buparlisib (**Fig. 3A**). Similar trends were observed with all four

therapeutics and across the panel of cell lines used.

GC-1 enhances the effects of therapeutic agents on cell migration.

In addition to reducing cell viability, many therapeutics target pathways that are

involved in cell migration. To further evaluate the ability of GC-1 to enhance the efficacy

of select therapeutics, cells were treated with the indicated inhibitors at concentrations

determined to be effective based on cell viability assays (**Fig. 3**), and migration was

observed by assessing wound closure. Each inhibitor was able to successfully slow

migration, and this effect was increased when each of the inhibitors were combined with

GC-1, except for Buparlisib in KTC-2 cells (**Fig. 4B-E**). Notably, GC-1 alone was able

to prevent wound closure of all cell lines with similar success as each inhibitor alone

(**Fig. 4A-E**). These findings are similar to what we previously observed in experiments

using endogenous thyroid hormone, T_3, indicating GC-1 is a suitable alternative for TRβ

activation.

GC-1 blocks tumorsphere outgrowth and increases efficacy of therapeutic
agents.

Current treatment options for ATC primarily target hyperproliferative cells, but

aggressive cancers like ATC are enriched for long-lived cancer stem cells that are

thought to be a major mechanism for therapy resistance and tumor recurrence. We

previously demonstrated that ligand-activated TRβ regulates genes related to stemness

in ATC cells and that GC-1 could block thyrosphere outgrowth in our modified ATC cells

(**Fig. 1D**). We therefore tested the hypothesis that GC-1 could prevent thyrosphere

outgrowth after an initial treatment with an inhibitor that targets proliferation.

ATC cells were treated with the indicated inhibitor or vehicle for three days,

followed by re-plating in non-adherent conditions. We again observed that GC-1 alone

could reduce thyrosphere formation in unmodified ATC cell lines (**Fig. 5A-C**). None of

the inhibitors could completely prevent thyrosphere outgrowth when used alone;

however, Sorafenib was the most effective overall (**Fig. 5C**). The addition of GC-1 upon

removal of each inhibitor blocked nearly all thyrosphere outgrowth. This effect was

consistent across all four cell lines and all three inhibitors used. Similar results were

obtained with pre-treatment of 8505C cells with GC-1 followed by exposure to

therapeutics (**Sup. Fig. 1**). Treatment with GC-1 also decreases mRNA expression of a

subset of stemness markers (**Sup. Fig. 2**) This suggests that GC-1 may be particularly

effective at blocking cancer stem cell expansion.

GC-1 promotes re-differentiation of ATC cell lines.

In transduced cells, TRβ activation with T_3 reduces the aggressive phenotype

partially through re-differentiation. Therefore, we investigated the potential for GC-1 to

promote re-differentiation in parental SW1736 cells. Gene expression analysis using

RT-qPCR revealed that 24 hours of GC-1 exposure increased levels of thyroid-specific

gene transcripts (**Fig. 6A**). This transcriptomic program is summarized by the thyroid

differentiation score (TDS), which significantly increased in GC-1-treated cells (**Fig. 6B**).

Additionally, GC-1 alone decreased the overall number of CD24$^-$/CD44$^+$ cells present in

the monolayer population as evident by RT-qPCR (**Sup. Fig. 3A**). We also observed a

decrease in the ATC markers *ALCAM* (CD166) and *ALDH1A1* (**Sup. Fig. 3B**).

Intriguingly, *SLC5A5* expression was increased with 24 hours of GC-1 (**Fig. 6A**).

Therefore, we further investigated NIS protein levels after 48 hours of GC-1 treatment.

Both nascent and mature (glycosylated) NIS protein significantly increased following

GC-1 treatment (**Fig. 6C-D**). Iodide uptake into cells also increased (**Fig. 6E**), indicating

that the increased NIS protein was functional. Similar results were observed in all ATC

cell lines including in charcoal stripped serum in 8505C cells to confirm the agonist

selective effect (**Sup. Fig. 3**). These findings reveal a novel function of GC-1 to induce

re-differentiation in an ATC cell line model and offers a promising target to increase NIS

levels and iodide uptake.

GC-1 blunts tumor growth as effectively as Sorafenib *in vivo.*

To validate our cell-based findings *in vivo*, the effect of GC-1 alone or in

combination with Sorafenib was tested in a xenograft study. GC-1 blunted tumor growth

with no significant change in tumor volume over the treatment period similar to

Sorafenib and the combination of both treatments (**Fig. 7A**). At day 16 of treatment, GC-

1 induced a significant 60% reduction in tumor mass as did Sorafenib (**Fig. 7B, C**). The

potential for drug synergy was masked due to the equivalent efficacy with Sorafenib at

these doses (**Fig. 7A-C**). All mouse groups exhibited similar weight gains over the

course of treatment (**Fig. 7D**). Immunofluorescence image analysis (HALO) on excised

tumors revealed diminished Ki67 in all treatment groups (**Fig. 7E-F**). The combination of

GC-1 and Sorafenib resulted in a significant reduction in Ki67 expression (**Fig. 7F**).

Region of Interest (ROI) analysis revealed preferential distribution of Ki67 expression in

cells along the periphery of the tumor samples.

DISCUSSION

There is compelling evidence that loss of expression of TRβ, a member of the

thyroid hormone receptor (TR) family, through genomic modifications and epigenetic silencing is characteristic of thyroid tumors[3,31-33]. Numerous pre-clinical studies demonstrate the potent anti-tumorigenic effect of robust expression of the *THRB* gene. Re-expression of silenced TRβ using demethylating agents delays thyroid tumor progression *in vivo* [34], mice expressing a c-terminal frameshift of TRβ (*Thrb^{PV}*) spontaneously develop follicular thyroid cancer [35], and *Thrb$^{PV/PV}$ Kras^{G12D}* mice develop thyroid tumors with features of de-differentiated thyroid cancer [36]. Our studies in anaplastic thyroid cancer cells revealed that activation of TRβ induces a tumor suppression transcriptomic program, induces re-differentiation, and reduces the aggressive phenotype[2,25].

Clinical studies of altered thyroidal status have revealed the potential that activation of TR might be of therapeutic value. Patients who were rendered hypothyroxinemic and treated with T_3 had longer survival with end-stage solid tumors [37] and T_3 treatment enhanced chemosensitivity [38]. Following tyrosine kinase inhibitor therapy, patients demonstrated improved survival with treatment of hypothyroidism [39]. Given that TRβ agonists have been developed [40], it is remarkable that selective activation of TRβ has not yet been explored in cancer models. Selective agonists have been clinically successful for treatment of metabolic disorders, hyperlipidemia, hypercholesterolemia, and non-alcoholic steatohepatitis without the harmful cardiovascular effects mediated by TRα [41,42]. In particular Sobetirome (GC-1) and derivatives have been extensively characterized. GC-1 preferentially binds to TRβ and regulates gene expression comparable to T_3 [21,43] and thus a candidate tumor suppressor molecule notably for aggressive cancers such as ATC.

The global incidence of thyroid cancer is increasing faster than any other solid

tumor [44,45] and outcomes for patients with resistant or recurrent disease and poorly

differentiated thyroid cancers (PDTC) are extremely poor [46]. Patients with advanced or

metastatic anaplastic thyroid cancer (ATC) have a higher mortality rate than all other

endocrine cancers combined[47]. Thus, novel combinatorial therapies are of critical need.

In ATC tumors, current approaches aim to inhibit or disrupt multiple pathways

with combination therapies including blocking cell-cycle, cell survival and proliferation

[48,49]. Our prior work demonstrated that TRβ represses signaling pathways and cell-cycle

regulation through genomic mechanisms[2,23]. Therefore, in this study, we tested

representative drugs that target the MAPK (Sorafenib), PI3K (Buparlisib, Alpelisib), and

cell cycle (Pablociclib) to assess whether GC-1 could increase their effectiveness. GC-1

reduced the effective drug concentrations required to observe a reduction in cell

viability. One of our most intriguing observations is that GC-1 alone significantly reduced

thyrosphere formation in all ATC cell types. Remarkably, GC-1 in combination with any

of the drug treatments nearly eliminated thyrosphere formation.

GC-1 induced re-differentiation in the SW1736 cell line in 24 hours,

demonstrating potent transcriptional reprogramming. While we have previously

demonstrated the potential for TRβ to promote a more differentiated phenotype in ATC

cells, this is the first time we have observed the profound impact of GC-1 in a parental

cell line with low TRβ expression. Importantly, we observed a concomitant increase in

NIS that led to a higher intake of iodide. Since PDTC and ATC tumors often exhibit a

marked decrease in NIS expression and resistance to RAI (radioactive iodine) therapy,

the results presented here raise a provocative implication for the use of GC-1 in

combination with RAI. Critically, we found that GC-1 could blunt tumor growth as

effectively as Sorafenib in a xenograft study confirming the observations in cell-based

studies. The combination of GC-1 and Sorafenib did not further reduce the tumor

volume or mass over the study period. Future studies establishing the

pharmacodynamics of GC-1 in combination with different doses of therapeutic agents

will elucidate the most effective combinatorial paradigm.

Our findings demonstrate that selective activation of TRβ with GC-1 can reduce

the aggressive tumor phenotype, reduce the cancer stemness in ATC cells, increase

the sensitivity of these cells to therapeutic agents, and promote re-differentiation and

iodide uptake in transduced cells in which TRβ is re-expressed and in parental cell lines

with diverse genetic backgrounds. Moreover, GC-1 treatment is as effective as the

therapeutic Sorafenib in inhibiting tumor growth. These studies provide the foundation

for further investigation of the broad impact of GC-1 on mitigating tumor progression,

pharmaceutical toxicity, and development of resistance to treatment.

ACKNOWLEDGEMENTS:

SW1736 and KTC-2 cell lines were generously provided by Dr. John A. Copland III

(Mayo Clinic). 8505C and OCUT-2 cell lines were obtained and authenticated from the

U Colorado Cancer Center Tissue Culture Shared Resource supported by National

Cancer Institute P30CA046934. Human cell line authentication, NextGen sequencing,

automated DNA sequencing was performed in the Vermont Integrative Genomics

Resource (RRID# SCR_021775) at UVM. Immunofluorescence staining, imaging and

analysis was performed at the Microscopy Imaging Center at UVM (RRID#

SCR_018821). Confocal microscopy was performed on a Nikon A1R-HD point scanning

confocal microscope (NIH 1S10OD025030-01). We thank the Histology Research

Support Facility in the Department of Pathology and Laboratory Medicine at UVMMC.

We thank the Office of Animal Care Management for their expertise for *in vivo*

experiments.

Authors' Contribution Statement: NEG: Conceptualization, Formal analysis,

Investigation, Writing – Review & editing. **LMC:** Formal analysis, Investigation, Writing.

ERW: Formal analysis, Investigation. **NMS:** Investigation, Formal analysis. **JAT:**

Investigation, Writing – Review & editing. **ELB:** Conceptualization, – Review & editing.

FEC: Conceptualization, Resources, Writing-Review & editing, Supervision, Project

administration, Funding acquisition.

Funding Statement: The research reported was supported by grants from National

Institutes of Health U54 GM115516; National Cancer Institute 1F99CA245796-01; UVM

Cancer Center-Lake Champlain Cancer Research Organization (C3) 12577-21; and

UVM Larner College of Medicine.

Authors' Disclosure Statement: FC is a member of the Endocrinology Editorial Board.

Data Availability: Original data generated and analyzed during this study are included

in this published article or in the data repositories listed in References.

REFERENCES

1. Kim WG, Cheng SY. Thyroid hormone receptors and cancer. *Biochim Biophys Acta*. Jul 2013;1830(7):3928-36. doi:10.1016/j.bbagen.2012.04.002

2. Bolf EL, Gillis NE, Davidson CD, et al. Thyroid Hormone Receptor Beta Induces a Tumor-Suppressive Program in Anaplastic Thyroid Cancer. *Mol Cancer Res*. Oct 2020;18(10):1443-1452. doi:10.1158/1541-7786.MCR-20-0282

3. Aranda A, Martinez-Iglesias O, Ruiz-Llorente L, Garcia-Carpizo V, Zambrano A. Thyroid receptor: roles in cancer. *Trends Endocrinol Metab*. Sep 2009;20(7):318-24. doi:10.1016/j.tem.2009.03.011

4. Kim WG, Zhao L, Kim DW, Willingham MC, Cheng SY. Inhibition of tumorigenesis by the thyroid hormone receptor beta in xenograft models. *Thyroid*. Feb 2014;24(2):260-9. doi:10.1089/thy.2013.0054

5. Kowalik MA, Puliga E, Cabras L, et al. Thyroid hormone inhibits hepatocellular carcinoma progression via induction of differentiation and metabolic reprogramming. *J Hepatol*. Jun 2020;72(6):1159-1169. doi:10.1016/j.jhep.2019.12.018

6. Zhu L, Tian G, Yang Q, et al. Thyroid hormone receptor β1 suppresses proliferation and migration by inhibiting PI3K/Akt signaling in human colorectal cancer cells. *Oncol Rep*. Sep 2016;36(3):1419-26. doi:10.3892/or.2016.4931

7. Bolf EL, Gillis NE, Davidson CD, et al. Common tumor-suppressive signaling of thyroid hormone receptor beta in breast and thyroid cancer cells. *Mol Carcinog*. Dec 2021;60(12):874-885. doi:10.1002/mc.23352

8. López-Mateo I, Alonso-Merino E, Suarez-Cabrera C, et al. Thyroid Hormone Receptor β Inhibits Self-Renewal Capacity of Breast Cancer Stem Cells. *Thyroid*. Jan 2020;30(1):116-132. doi:10.1089/thy.2019.0175

9. Martínez-Iglesias O, Ruiz-Llorente L, Jurado CC, Aranda A. Chapter 11 - Thyroid Hormone Receptors and their Role in Cell Proliferation and Cancer. In: Ulloa-Aguirre A, Tao Y-X, eds. *Cellular Endocrinology in Health and Disease (Second Edition)*. Academic Press; 2021:229-246.

10. Pourvali K, Shimi G, Ghorbani A, Shakery A, Shirazi FH, Zand H. Selective thyroid hormone receptor beta agonist, GC-1, is capable to reduce growth of colorectal tumor in syngeneic mouse models. *J Recept Signal Transduct Res*. Apr 27 2022:1-8. doi:10.1080/10799893.2022.2032748

11. Davidson CD, Gillis NE, Carr FE. Thyroid Hormone Receptor Beta as Tumor Suppressor: Untapped Potential in Treatment and Diagnostics in Solid Tumors. *Cancers (Basel)*. Aug 24 2021;13(17)doi:10.3390/cancers13174254

12. Lee WK, Cheng SY. Targeting transcriptional regulators for treatment of anaplastic thyroid cancer. *J Cancer Metastasis Treat*. 2021;7doi:10.20517/2394-4722.2021.58

13. Pozdeyev N, Rose MM, Bowles DW, Schweppe RE. Molecular therapeutics for anaplastic thyroid cancer. *Semin Cancer Biol*. Apr 2020;61:23-29. doi:10.1016/j.semcancer.2020.01.005

14. Nabhan F, Dedhia PH, Ringel MD. Thyroid cancer, recent advances in diagnosis and therapy. *Int J Cancer*. Sep 1 2021;149(5):984-992. doi:10.1002/ijc.33690

15. McCrary HC, Aoki J, Huang Y, et al. Mutation based approaches to the treatment of anaplastic thyroid cancer. *Clin Endocrinol (Oxf)*. May 2022;96(5):734-742. doi:10.1111/cen.14679

16. Prete A, Matrone A, Gambale C, et al. Poorly Differentiated and Anaplastic Thyroid Cancer: Insights into Genomics, Microenvironment and New Drugs. *Cancers (Basel)*. Jun 26 2021;13(13)doi:10.3390/cancers13133200

17. Saponaro F, Sestito S, Runfola M, Rapposelli S, Chiellini G. Selective Thyroid Hormone Receptor-Beta (TRβ) Agonists: New Perspectives for the Treatment of Metabolic and Neurodegenerative Disorders. *Front Med (Lausanne)*. 2020;7:331. doi:10.3389/fmed.2020.00331

18. Li L, Song Y, Shi Y, Sun L. Thyroid Hormone Receptor-β Agonists in NAFLD Therapy: Possibilities and Challenges. *J Clin Endocrinol Metab*. Jun 16 2023;108(7):1602-1613. doi:10.1210/clinem/dgad072

19. Scanlan TS. Sobetirome: a case history of bench-to-clinic drug discovery and development. *Heart Fail Rev*. Mar 2010;15(2):177-82. doi:10.1007/s10741-008-9122-x

20. Wagner RL, Huber BR, Shiau AK, et al. Hormone selectivity in thyroid hormone receptors. *Mol Endocrinol*. Mar 2001;15(3):398-410. doi:10.1210/mend.15.3.0608

21. Luong XG, Stevens SK, Jekle A, et al. Regulation of gene transcription by thyroid hormone receptor β agonists in clinical development for the treatment of non-alcoholic steatohepatitis (NASH). *PLoS One*. 2020;15(12):e0240338. doi:10.1371/journal.pone.0240338

22. Bleicher L, Aparicio R, Nunes FM, et al. Structural basis of GC-1 selectivity for thyroid hormone receptor isoforms. *BMC Struct Biol*. Jan 31 2008;8:8. doi:10.1186/1472-6807-8-8

23. Davidson CD, Bolf EL, Gillis NE, Cozzens LM, Tomczak JA, Carr FE. Thyroid hormone receptor beta inhibits the PI3K-Akt-mTOR signaling axis in anaplastic thyroid cancer cell lines via genomic mechanisms. *Journal of the Endocrine Society*. 2021;doi:10.1210/jendso/bvab102

24. Landa I, Ibrahimpasic T, Boucai L, et al. Genomic and transcriptomic hallmarks of poorly differentiated and anaplastic thyroid cancers. *The Journal of Clinical Investigation*. 03/01/ 2016;126(3):1052-1066. doi:10.1172/JCI85271

25. Carr FE, Tai PW, Barnum MS, et al. Thyroid Hormone Receptor-beta (TRbeta) Mediates Runt-Related Transcription Factor 2 (Runx2) Expression in Thyroid Cancer Cells: A Novel Signaling Pathway in Thyroid Cancer. *Endocrinology*. Aug 2016;157(8):3278-92. doi:10.1210/en.2015-2046

26. Davidson CD, Tomczak JA, Amiel E, Carr FE. Inhibition of Glycogen Metabolism Induces Reactive Oxygen Species-Dependent Cytotoxicity in Anaplastic Thyroid Cancer in Female Mice. *Endocrinology*. Oct 23 2022;163(12)doi:10.1210/endocr/bqac169

27. Khan HY, Ge J, Nagasaka M, et al. Targeting XPO1 and PAK4 in 8505C Anaplastic Thyroid Cancer Cells: Putative Implications for Overcoming Lenvatinib Therapy Resistance. *Int J Mol Sci*. Dec 29 2019;21(1)doi:10.3390/ijms21010237

28. Lin SF, Lin JD, Hsueh C, Chou TC, Wong RJ. Effects of roniciclib in preclinical models of anaplastic thyroid cancer. *Oncotarget*. Sep 15 2017;8(40):67990-68000. doi:10.18632/oncotarget.19092

29. Yun HJ, Kim HJ, Kim J, et al. Synergistic Anticancer Activity of N-Hydroxy-7-(2-Naphthylthio) Heptanomide, Sorafenib, and Radiation Therapy in Patient-Derived Anaplastic Thyroid Cancer Models. *Int J Mol Sci*. Jan 7 2021;22(2)doi:10.3390/ijms22020536

30. Fang L, Liu YJ, Zhang YW, et al. Comparison of Proteomics Profiles Between Xenografts Derived from Cell Lines and Primary Tumors of Thyroid Carcinoma. *J Cancer*. 2021;12(7):1978-1989. doi:10.7150/jca.50897

31. Silva JM, Domínguez G, González-Sancho JM, et al. Expression of thyroid hormone receptor/erbA genes is altered in human breast cancer. Original Paper. *Oncogene*. 06/14/online 2002;21:4307. doi:10.1038/sj.onc.1205534

32. Wojcicka A, Piekielko–Witkowska A, Kedzierska H, et al. Epigenetic Regulation of Thyroid Hormone Receptor Beta in Renal Cancer. *PLOS ONE*. 2014;9(5):e97624. doi:10.1371/journal.pone.0097624

33. Park JW, Zhao L, Willingham M, Cheng S-y. Oncogenic mutations of thyroid hormone receptor β. *Oncotarget*. 02/28

11/20/received

01/20/accepted 2015;6(10):8115-8131.

34. Kim WG, Zhu X, Kim DW, Zhang L, Kebebew E, Cheng SY. Reactivation of the silenced thyroid hormone receptor beta gene expression delays thyroid tumor progression. *Endocrinology*. Jan 2013;154(1):25-35. doi:10.1210/en.2012-1728

35.   Suzuki H, Willingham MC, Cheng SY. Mice with a mutation in the thyroid hormone receptor beta
gene spontaneously develop thyroid carcinoma: a mouse model of thyroid carcinogenesis. *Thyroid*. Nov
2002;12(11):963-9. doi:10.1089/105072502320908295
36.   Zhu X, Zhao L, Park JW, Willingham MC, Cheng SY. Synergistic signaling of KRAS and thyroid
hormone receptor beta mutants promotes undifferentiated thyroid cancer through MYC up-regulation.
Neoplasia. Sep 2014;16(9):757-69. doi:10.1016/j.neo.2014.08.003
37.   Hercbergs A, Johnson RE, Ashur-Fabian O, Garfield DH, Davis PJ. Medically induced euthyroid
hypothyroxinemia may extend survival in compassionate need cancer patients: an observational study.
Oncologist. Jan 2015;20(1):72-6. doi:10.1634/theoncologist.2014-0308
38.   Huang J, Jin L, Ji G, et al. Implication from thyroid function decreasing during chemotherapy in
breast cancer patients: chemosensitization role of triiodothyronine. *BMC Cancer*. Jul 6 2013;13:334.
doi:10.1186/1471-2407-13-334
39.   Lechner MG, Vyas CM, Hamnvik OR, et al. Hypothyroidism During Tyrosine Kinase Inhibitor
Therapy Is Associated with Longer Survival in Patients with Advanced Nonthyroidal Cancers. *Thyroid*. Apr
2018;28(4):445-453. doi:10.1089/thy.2017.0587
40.   Davis PJ, Tang HY, Hercbergs A, Lin HY, Keating KA, Mousa SA. Bioactivity of Thyroid Hormone
Analogs at Cancer Cells. *Front Endocrinol (Lausanne)*. 2018;9:739. doi:10.3389/fendo.2018.00739
41.   Trost SU, Swanson E, Gloss B, et al. The thyroid hormone receptor-beta-selective agonist GC-1
differentially affects plasma lipids and cardiac activity. *Endocrinology*. Sep 2000;141(9):3057-64.
doi:10.1210/endo.141.9.7681
42.   Grover GJ, Mellstrom K, Ye L, et al. Selective thyroid hormone receptor-beta activation: a
strategy for reduction of weight, cholesterol, and lipoprotein (a) with reduced cardiovascular liability.
Proc Natl Acad Sci U S A. Aug 19 2003;100(17):10067-72. doi:10.1073/pnas.1633737100
43.   Yuan C, Lin JZ, Sieglaff DH, et al. Identical gene regulation patterns of T3 and selective thyroid
hormone receptor modulator GC-1. *Endocrinology*. Jan 2012;153(1):501-11. doi:10.1210/en.2011-1325
44.   Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. *CA Cancer J Clin*. Jan 2017;67(1):7-30.
doi:10.3322/caac.21387
45.   Pereira M, Williams VL, Hallanger Johnson J, Valderrabano P. Thyroid Cancer Incidence Trends in
the United States: Association with Changes in Professional Guideline Recommendations. *Thyroid*. Aug
2020;30(8):1132-1140. doi:10.1089/thy.2019.0415
46.   Glaser SM, Mandish SF, Gill BS, Balasubramani GK, Clump DA, Beriwal S. Anaplastic thyroid
cancer: Prognostic factors, patterns of care, and overall survival. *Head Neck*. Apr 2016;38 Suppl 1:E2083-
90. doi:10.1002/hed.24384
47.   Prasongsook N, Kumar A, Chintakuntlawar AV, et al. Survival in Response to Multimodal Therapy
in Anaplastic Thyroid Cancer. *J Clin Endocrinol Metab*. Dec 1 2017;102(12):4506-4514.
doi:10.1210/jc.2017-01180
48.   Alobuia W, Gillis A, Kebebew E. Contemporary Management of Anaplastic Thyroid Cancer. *Curr
Treat Options Oncol*. Aug 7 2020;21(10):78. doi:10.1007/s11864-020-00776-2
49.   Molinaro E, Romei C, Biagini A, et al. Anaplastic thyroid carcinoma: from clinicopathology to
genetics and advanced therapies. Review Article. *Nature Reviews Endocrinology*. 07/14/online
2017;13:644. doi:10.1038/nrendo.2017.76

Figure 1. GC-1 blocks tumorigenic phenotypes in ATC cells transduced with TRβ.

A. Growth curves demonstrate that SW-TRβ growth is significantly decreased after

treatment with GC-1 for 4 days (n=3). Significance (* p<0.05) was determined by a two-

way ANOVA and Tukey's multiple comparisons test. **B**. Area under the curve (AUC)

analysis of growth curve data. Significance (* p<0.05) was determined by t-test. **C.**

Treatment with GC-1 slows migration of SW-TRβ and SW-EV cells measured by

scratch assay. Significance (* p<0.05) was determined by t-test. **D.** GC-1 repressed

thyrosphere formation in SW-TRβ and SW-EV cells (n=4). Significance (* p<0.05) was

determined by t-test. **E.** Immunoblot demonstrates that 5 days of GC-1 treatment

induces apoptosis in SW-TRβ cells assessed by cleavage of PARP and caspase 3

(n=3).

Figure 2: GC-1 slows growth of parental ATC cells. **A**. Summary of mutated genes

in four ATC cell lines implemented in this study (17). **B**. Relative cell growth of

unmodified ATC cell lines was measured by cell counting assay after 4 days of

treatment with 10 nM GC-1 or 10 nM T_3. ignificance (* p<0.05) was determined by a

two-way ANOVA followed by Dunnett's multiple comparisons test; 3 independent

experiments were performed per treatment group. **C.** GC-1 increases TRβ mRNA

expression, measured by qPCR, in ATC cells relative to vehicle control at 24 hours. **D**.

TRβ protein is detectable by immunoblot in each cell line. **E.** GC-1 treatment induces a

modest but not significant increase in TRβ protein levels quantified by densitometry.

Figure 3. GC-1 increases efficacy of inhibitors. Cell viability of unmodified ATC cell

lines was measured by SRB assay after 3 days of treatment with increasing indicated

concentrations of buparlisib or alpelisib **A**, sorafenib **B**, or palbociclib **C** simultaneously

in combination with 10 nM GC-1. Significance (* $p < 0.05$) was determined by a two-way

ANOVA followed by Sidak's multiple comparisons test; 3 independent experiments were

performed per each treatment group. Area under the curve (AUC) analysis of each

dose-response experiment is shown in the inset.

Figure 4. GC-1 enhances the effects of inhibitors on cell migration. Cell migration

was measured by scratch assay. Representative images demonstrate reduced scratch

closure after two days when ATC cells are treated with GC-1 **A**. Area under the curve

(AUC) analysis shows SW1736 **B**, 8505C **C**, OCUT2 **D**, and KTC-2 **E** cells all exhibit

lower rates of scratch closer after treatment with GC-1 alone and further reduced

scratch closure rates after combined treatment with GC-1 and inhibitors. Significance (*

$p < 0.05$) was determined by t-test.

Figure 5. GC-1 blocks thyrosphere outgrowth and increases the efficacy of

therapeutic agents. Thyrosphere growth was determined for ATC cells after 3 days of

treatment with 0.5 µM buparlisib or alpelisib **A**, 10 µM sorafenib **B**, or 1 nM palbociclib **C**

under adherent culture conditions, followed by plating in conditions for spheroid growth

in the presence of 10 nM GC-1 for 7 days. GC-1 alone and each therapeutic agent

significantly blocked sphere formation in all cell types. GC-1 in addition to each

therapeutic agent further inhibited or completely blocked sphere formation. Significance

(* $p < 0.05$) was determined by two-way ANOVA followed by Sidak's multiple

comparisons test; 3 independent experiments were performed per each treatment

group; n=3 per analyses.

Figure 6. GC-1 induces expression of redifferentiation markers in SW1736 cells.

A. RT-qPCR for thyroid differentiation markers was performed on SW1736 cells treated

with or without 10 nM GC-1 for 24 hours. mRNA levels in cells treated with GC-1 are

relative to the untreated cells. Significance (* $p < 0.05$) was determined by Student's

unpaired t test; 3 independent experiments were performed per treatment group. **B.**

Thyroid differentiation score (TDS) was calculated using the expression level of the

genes in panel A, excluding *THRB*, as previously described (10, 17). Significance (*

$p < 0.0001$) was determined by Student's unpaired t test. **C.** SW1736 cells were treated

with or without 10 nM GC-1 in triplicate for 48 hours prior to protein extraction and

analysis via immunoblot. Normal-like thyroid Nthy-ORI cells were used as a positive

control. **D.** Relative NIS/GAPDH protein levels from C were quantified via ImageJ, and

significance (* $p < 0.01$) was determined by Student's unpaired t test. **E.** Iodide uptake

was measured after 48 hours of GC-1 treatment. Significance was determined by

Student's unpaired t test

(* $p < 0.001$).

Figure 7. GC-1 inhibits tumor growth as effectively as Sorafenib in an ATC

xenograft model. Nude mice were injected subcutaneously with 1 x 10^6 8505C cells.

Tumors were established after 1 week. Mice were assigned to 4 treatment groups by

weight and injected i.p. daily for 16 days with vehicle control, 10 mg/kg sorafenib, 0.3

mg/kg GC-1, or a combination of Sorafenib and GC-1. **A.** Percent change in tumor

volumes from day 1 of treatment; *** $p < 0.001$ **B.** Representative excised tumors from

each treatment. No evidence of metastases was noted in any group. Reduced

vascularization was observed with GC-1 treatment. **C.** Tumors were excised on Day 16

and weighed. Sorafenib (blue), GC-1 (red) induced a 60% reduction in final tumor mass

with no difference between the two treatments.　The combined treatments did not

further reduce the tumor mass over the time period. Significance indicated ** $p < 0.01$. **D.**

Mice were weighed three times a week following drug injections to monitor toxicity. No

significant change in animal weight was noted for any group. **E.** Representative images

of tumors stained for Ki-67 are illustrated. Regions of interest containing only tumor

tissue were outlined on each tiled image.　Nuclear detection was set to the DAPI

channel and the nuclear detection settings were optimized to accurately segment nuclei

in the images. **F.** Images were analyzed using HALO image analysis (Indica Labs,

Albuquerque NM; version 3.3.45) utilizing the HighPlex FL module (version 4.1.3).

Quantification of Ki-67 fluorescent intensity per cell is summarized. Sorafenib and GC-1

decreased Ki-67 but only significantly in combination. Significance (** $p < 0.01$) was

determined by one-way ANOVA followed by Sidak's multiple t-test.

Figure 1

Figure 2

A

Gene Function	Gene Name	SW1736	8505C	OCUT-2	KTC-2
		\multicolumn: Cell Line			
		\multicolumn: Mutation Type			
MAPK Signaling	BRAF	V600E	V600E	V600E	V600E
Telomerase	TERT (promoter)	124C-T	146C-T	146C-T	124C-T
DNA Homeostasis	TP53	Trunc.	Miss.		
PI3K Signaling	PIK3CA			Kinase Domain	
Cell-Cell Contact	NF2		Trunc.		
DNA Homeostasis	ATM		Trunc.		
TSH Signaling	TSHR	Miss.			
Histone Methyltransferase	KMT2C				Miss.
Histone Methyltransferase	KMT2D				Trunc.
Histone Acetyltransferase	EP300	Trunc.			
SWI/SNF Complex	ARID1A				In Frame
SWI/SNF Complex	SMARCA4		Miss.		

Figure 3

Figure 4

Figure 5

Figure 6

A

DIO1 | **DIO2** | **DUOX1** | **DUOX2** | **FOXE1** | **NKX2-1** | **SLC5A5**

SLC26A4 | **PAX8** | **TG** | **THRA** | **THRB** | **TPO** | **TSHR**

B

Thyroid Differentiation Score (TDS)

C

SW1736   Nthy-ORI

GC-1:   -   -   -   +   +   +   -

Glycosylated NIS — 68 kDa

Nascent NIS — 55 kDa

GAPDH — 37 kDa

D

NIS Protein Level

E

Iodide Uptake

Figure 7