A Thyroid Hormone Receptor Beta Specific Agonist Suppresses Anaplastic Thyroid

Cancer Cell Phenotype and Increases Efficacy of Therapeutic Agents

Noelle E. Gillis [1,2*], Cole D. Davidson [1,2*], Lauren M. Cozzens [1,2], Emily Wilson [1,2], Eric L. Bolf [1,2],

Jennifer A. Tomczak [1,2], Frances E. Carr [1,2**]

[1] Department of Pharmacology, Larner College of Medicine, Burlington, Vermont

[2] University of Vermont Cancer Center, Burlington, Vermont

[**] Corresponding author, Email: Frances.Carr@med.uvm.edu

[*] These authors contributed equally to this work

Short title: TRβ agonist inhibits ATC progression and increases therapeutic efficacy

Keywords: GC-1, Sobetirome, thyroid hormone receptor, anaplastic thyroid cancer, PI3K

inhibitors, MAPK inhibitor, cell cycle inhibitor

Abstract

Background: Anaplastic thyroid cancer (ATC) is one of the most lethal endocrine cancers, with

an average survival time of six months after diagnosis. These aggressive tumors have very

limited treatment options highlighting a need for a deeper understanding of its mechanisms for

development of more effective therapies. We have previously shown that the liganded thyroid

hormone receptor beta (TRβ) can function as a tumor suppressor and induce re-differentiation

in ATC cells. We therefore tested the hypothesis that selective activation of TRβ with

sobetirome (GC-1) could reduce the tumorigenic phenotypes of ATC cell lines and improve the

efficacy of clinically relevant therapeutics.

Methods: We used a panel of four ATC cell lines with variable genetic backgrounds to assess

the ability of GC-1 to reduce the aggressive phenotype. The effects of GC-1 alone or in

combination with buparlisib, alpelisib, sorafenib, and palbociclib on cell growth, viability, and

migration were determined and compared with the gene expression levels of selected markers.

The impact of these treatments on the cancer stem cell population was assessed by

tumorsphere assay. Thyroid differentiation markers were measured by gene analysis, and

sodium iodide symporter (NIS) protein level and function were determined.

Results: Our results show that GC-1 alone can decrease cell viability, growth, and slow cell

migration in all four ATC cell lines. In addition, GC-1 is able to further block each of these

phenotypes when combined with buparlisib, alpelisib, sorafenib, or palbociclib. GC-1 alone

blocks thyrosphere outgrowth in all cell lines and increases the efficacy of each of the

therapeutic agents tested. GC-1 increased NIS transcript and protein levels to allow for

increased iodide uptake in ATC cells.

Conclusion: Activation of TRβ with selective agonist sobetirome (GC-1) reduces the aggressive

phenotype and induced re-differentiation in ATC cells and increases the efficacy of therapeutic

agents that are currently used in the treatment of ATC. These results indicate that selective

activation of TRβ not only induces a tumor suppression program de novo but enhances the

effectiveness of anti-cancer agents.

Introduction

The global incidence of thyroid cancer, the most common endocrine malignancy, is increasing faster than any other solid tumor almost tripling over the past few decades (1,2). While the prognoses for patients with differentiated tumors receiving standard therapies are generally very good, outcomes for patients with resistant or recurrent disease and poorly differentiated thyroid cancers (PDTC) are extremely poor (3). Due to a lack of effective therapies, patients with advanced or metastatic anaplastic thyroid cancer (ATC) have a higher mortality rate than all other endocrine cancers combined with a mean survival time of 6 months (4). The pathogenesis of thyroid cancer is characterized by dysregulation of intracellular signaling pathways, abnormalities in expression of tumor suppressors, cell cycle regulators, and apoptotic signaling. ATC tumors are refractory to radioiodine therapy due to loss of sodium-iodide symporter (NIS), and traditional chemotherapy is of limited benefit. Activation of phosphatidylinositol 3-kinase (PI3K)–Akt–mammalian target of rapamycin (mTOR) pathway (PI3K-Akt) and mitogen-activated protein kinase (MAPK) signaling are recognized drivers of thyroid oncogenesis and blocking this activity is a therapeutic focus. Yet inhibitors of the major components of these signaling cascades rarely provide a durable response due to the development of resistance.

The aggressive behavior of ATC and poor outcome with conventional therapies has led to the continued evaluation of new drugs with variable levels of toxicity which limit their clinical use (5-7). Molecular diagnostics, notably identification of BRAF[V600E] mutations, provide rationale for intervention with combination therapies (8). Thus far, the use of combination treatments has not yielded significantly improved responses due to the development of resistance, and there are still no curative options. Therefore, there remains an urgent need for innovative and less-toxic therapeutic options for ATC.

There is compelling evidence that the ligand-activated thyroid hormone receptor beta

(TRβ) functions as a suppressor of tumor progression, invasion, and metastases in a number of

endocrine cancers, including differentiated thyroid tumors. Our recent studies revealed that TRβ

in the presence of triiodothyronine (T_3) induced a tumor-suppressive transcriptomic program in

anaplastic thyroid cancer (ATC) cells (9,10). We identified key signaling pathways by which

activation of TRβ could reduce tumor cell progression, by altered gene expression, implicating

thyroid hormone signaling as a potential therapeutic pathway in ATC. However, activation of

TRβ by non-isoform selective ligands such as T_3 is contraindicated as a therapeutic approach

due to potential for adverse side-effects. Isoform-selective T_3 analogs that preferentially bind to

and activate TRβ mitigate these deleterious effects and have been clinically successful for the

treatment of metabolic disorders (11-14). The analogue sobetirome, or GC-1, induces similar

transcriptomic changes to T_3 (15,16).

In our studies presented here, we establish the effects of GC-1 alone and in combination

with therapeutic agents in ATC cells with diverse genetic backgrounds (17). GC-1 alone

decreases the aggressive phenotype and increases the effectiveness of MAPK, PI3K, and cell

cycle inhibitors. GC-1 significantly increases expression of thyroid-specific genes, inducing re-

differentiation in ATC cells. Importantly, we observe an increase in NIS protein which leads to

enhanced iodide uptake. These observations indicate that selective activation with isomer-

specific agonists may be an effective and practical adjuvant therapeutic strategy.

MATERIALS AND METHODS:

Culture of thyroid cell lines. Anaplastic thyroid cancer cell lines (SW1736, 8505C, OCUT-2,

and KTC-2) and the normal-like thyroid cell line Nthy-ORI were cultured in RPMI 1640 growth

media with L-glutamine (300 mg/L), sodium pyruvate and nonessential amino acids (1%)

(Corning Inc, Corning, NY, USA), supplemented with 10% fetal bovine serum (Thermo Fisher

Scientific, Waltham, MA, USA) and penicillin-streptomycin (200 IU/L) (Corning) at 37°C, 5%

CO_2, and 100% humidity. For measuring *MYC* expression via RT-qPCR, cells were maintained

in media with 10% charcoal stripped fetal bovine serum (Life Technologies). SW1736 cells were

modified by lentiviral transduction as recently described (10). SW1736, KTC-2, and Nthy-ORI

were authenticated by the Vermont Integrative Genomics Resource at the University of Vermont

using short tandem repeat profiles and Promega GenePrint10 System (SW1736, May 2019;

KTC-2, October 2019; Nthy-ORI, August 2020). 8505C and OCUT2 were authenticated by the

University of Colorado using short tandem repeat profiles (8505C, June 2013; OCUT-2, June

2018).

Pharmacological agents. T_3 and GC-1 (Sigma, St. Louis, MO, USA) were suspended in 1 M

NaOH (T_3) and 100% ethanol (GC-1) for stock concentrations of 1 mM before further dilution in

culture medium. Buparlisib, sorafenib, palbociclib, and alpelisib (MedChemExpress, Monmouth

Junction, NJ, USA) were suspended in 100% DMSO for stock concentrations of 50-150 mM

before further dilution in culture medium.

Cell Growth and Viability Assays. Cell growth was measured by cell counting at discrete time

points. Cells were seeded in 12- well plates and growth optimized. Cells were treated with GC-

1, buparlisib, alpelisib, palbociclib, or sorafenib to establish time and concentration effects of the

select agents on cell growth. Cell viability was determined by a Sulforhodamine B assay

(Abcam, Cambridge, UK) following the manufacturer's protocol. In brief, ATC cells were plated

in 96-well clear flat-bottom plates at a density of 5,000 cells per well. Cells were fixed, stained,

and then imaged using a plate reader according to manufacturer's instructions.

Migration Assay. Cell migration was determined by wound healing assay. Cells were plated

and allowed to grow to 100% confluency. Four hours prior to scratching, cells were treated with

10 µg/ml mitomycin C (Sigma). A scratch was performed with a P1000 pipette tip and debris

was washed away with PBS. Media was supplemented with GC-1 (10 nM) with and without 0.5

µM buparlisib, 0.5 µM alpelisib, 1 nM palbociclib, or 5 µM sorafenib. Images were obtained at 0,

16, 24, 48, and 72 hours or until 100% wound closure, depending on the cell type. The wound

closure was measured using the ImageJ macro "Wound Healing Tool"

(https://github.com/MontpellierRessourcesImagerie/imagej_macros_and_scripts/wiki/Wound-

Healing-Tool). Percent closure was calculated relative to the area of the initial scratch.

Tumorsphere Assay. Cancer stem cell tumorspheres formed from ATC cells were used to

assess self-renewal and sphere-forming efficiency. For generating tumorspheres, adherent

monolayer cells were dissociated with Trypsin-EDTA, and single cells were moved to round-

bottom ultra-low attachment 24-well plates at a density of 1000 cells/well (Corning).

Tumorspheres were cultured in RPMI 1640 growth media supplemented with 20 ng/mL each of

epidermal growth factor (EGF) and fibroblastic growth factor (FGF) (Gold Bio). Where indicated,

adherent cells were treated with 0.5 µM buparlisib, 0.5 µM alpelisib, 1 nM palbociclib, 5 µM

sorafenib, or vehicle for 72 hours prior to thyrosphere-forming assay. Tumorspheres were then

cultured in the presence or absence of GC-1 (10 nM) to evaluate the effects of liganded TRβ on

thyrosphere growth alone or subsequent to a therapeutic agent treatment. Tumorspheres grew

for seven days and were then counted with an inverted microscope (Carl Zeiss, Oberkochen,

Germany).

Immunoblot Analysis. Proteins were isolated from whole cells in lysis buffer (20mM Tris-HCl

[pH 8], 137mM NaCl, 10% glycerol, 1% Triton X-100, and 2mM EDTA) containing Protease

Inhibitor Cocktail 78410 (Thermo Fisher Scientific). Proteins were resolved by polyacrylamide

gel electrophoresis on 10% sodium dodecyl sulfate gels EC60752 (Thermo Fisher Scientific)

and immobilized onto nitrocellulose membranes (GE Healthcare, Chicago, IL, USA) by

electroblot (Bio-Rad Laboratories, Hercules, CA, USA). Membranes were blocked with 5% w/v

BSA in TBS and 0.1% v/v Tween20 (Gold Bio) for one hour at room temperature. Specific

proteins were detected by immunoblotting with the indicated antibodies (Table 1);

immunoreactive proteins were detected by enhanced chemiluminescence (Thermo Fisher

Scientific) on a ChemiDoc XRS+ (Bio-Rad Laboratories).

RNA Extraction and Quantitative Real-Time PCR (qRT-PCR). Total RNA was extracted using

RNeasy Plus Kit (Qiagen, Hilden, DE) according to manufacturer's protocol. cDNA was then

generated using 5X LunaScript RT SuperMix, and mRNA expression was quantified by qRT-

PCR using 2X Luna Universal qPCR Master Mix (NEB, Ipswich, MA, USA) on a QuantStudio 3

real-time PCR system (Applied Biosystems, Foster City, CA, USA). Fold change in gene

expression compared to endogenous controls was calculated using the ddCT method. Primer

sequences are indicated in Table 2.

Iodide Uptake Assay. Cells were seeded at a density of 40,000 cells per well in a 96 well plate

with or without 10 nM GC-1 for 48 hours. Media were discarded, and cells were washed four

times with warm PBS before incubation with 1 μM potassium iodide (KI, Sigma) in PBS for 1

hour at 37°C. Wells were washed an additional four times with warm PBS and 100 μL of ddH$_2$O

were added to each well. A standard curve ranging from 100-700 nM KI was assayed in parallel.

100 μl of 24 mM ammonium cerium (IV) sulfate hydrate with 5.7% concentrated H$_2$SO$_4$ (Sigma)

and 100 μl of 24 mM sodium arsenite (III) with 205 mM sodium chloride (NaCl) and 50 mM

sodium hydroxide (NaOH, Sigma) were added to each well. The plate was incubated in the dark

at room temperature for 30 minutes, and absorbance at 415 nm was measured with a Synergy

2 Multi-Detection Microplate Reader (Agilent Technologies, Santa Clara, CA, USA).

Statistics. All statistical analyses were performed using GraphPad Prism software. Paired

comparisons were by T-test and group comparisons were made by a 2-way ANOVA followed by

a Tukey multiple comparisons test or Sidak's multiple analyses where indicated. Data are

represented as mean ± standard deviation. Significance is indicated * p<0.05. Area under the

curve (AUC) at the 95[th] confidence interval was used to evaluate statistical differences in growth

and migration assays.

RESULTS:

Sobetirome, GC-1, blocks the tumorigenic phenotype and cancer stem cell growth in ATC

cells transduced with TRβ.

168   We have previously demonstrated that T_3 (10 nM) treatment of transduced SW1736

cells, in which TRβ is re-expressed (SW-TRβ), induced a tumor suppression transcriptomic

program and reduced cell growth when compared to cells transduced with empty vector (SW-

EV) with low levels of endogenous TRβ (10). In the present studies, we tested the ability of GC-

1 (10 nM) to alter the ATC phenotype and cancer cell stemness. We found that GC-1

significantly decreased cell growth in four days (**Fig. 1A**). There was a 50% reduction in the

overall growth rate of SW-EV cells and 75% reduction in SW-TRβ cells, indicating potent

activation of endogenous TRβ (**Fig. 1B**).

176   As cancer stem cells are considered to be responsible for tumor initiation and tumor

recurrence after chemotherapy, targeting these cells is a key for successful treatment.

Therefore, a critical question is whether or not thyroid cancer stem cells are responsive to GC-1.

Thyrosphere formation is robust in SW-EV cells, and GC-1 induced a significant reduction in

growth (**Fig. 1C**). Re-expression of TRβ (SW-TRβ) reduces thyrosphere growth and the addition

of GC-1 induces a greater than 80% reduction in thyrosphere formation (**Fig. 1C**). These results

indicate that GC-1 alone can activate TRβ to reduce the thyroid cancer stem cell population.

183   Apoptotic signaling is a key factor in mediating ligand-activated TRβ modification of the

ATC aggressive phenotype (10). Thus, we assessed the effects of GC-1 on PARP and caspase

3 cleavage by immunoblot as markers of late-stage apoptosis (**Fig. 1D**). GC-1 significantly

increased the presence of cleaved PARP and cleaved caspase 3 in SW-TRβ consonant with

increased apoptotic signaling.

These results indicate that GC-1 can activate TRβ to reduce the aggressive phenotype

in transduced ATC cells with TRβ re-expressed and recapitulate the effects seen with T_3 at

comparable concentrations.

GC-1 slows growth of parental ATC cells.

Although it was confirmed that GC-1 could induce the same phenotypic changes in our

transduced ATC cell line model, it was critical to assess whether agonist-selective activation of

endogenous TRβ could induce similar effects in unmodified parental ATC cell lines with diverse

mutational backgrounds (17). SW1736, 8505C, OCUT2, and KTC-2 cells all harbor $BRAF^{V600E}$

and *TERT* driver mutations. SW1736 and 8505C have additional *TP53* mutations. SW1736 has

an additional *TSHR* mutation, and OCUT2 has a gain-of-function *PIK3CA* mutation

(summarized in **Fig. 2A**). Endogenous TRβ expression is low in all four ATC cell lines when

compared with normal thyroid cells (18). Treatment of each of these cell lines with 10 nM GC-1

for four days yielded a modest but significant decrease in cell growth in all four cell lines (**Fig.

2B-E**). Treatment with 10 nM T_3 did not significantly alter the growth of any of these four cell

lines. This result suggests that even low levels of endogenous TRβ can be selectively activated

with GC-1, and it is more effective at promoting tumor suppressor activity than T_3.

GC-1 increases the efficacy of selected therapeutic agents

As agonist activation of TRβ can induce a tumor suppression program and reduce the

aggressive tumor phenotype in ATC cells, a critical question is whether GC-1 can enhance the

efficacy of therapeutics that target selective pathways. Cell viability was calculated following

treatment for three days with increasing concentrations of the PI3K inhibitor buparlisib (pan-

PI3K inhibitor) or alpelisib (PI3K⬚ mutant inhibitor) for OCUT2 cells which harbor a gain-of-

function *PIK3CA* mutation (**Fig. 3A**), the cell cycle inhibitor palbociclib (**Fig. 3B**), or the multi-

tyrosine kinase inhibitor sorafenib (**Fig. 3C**), with or without 10 nM GC-1. Each of the

therapeutic agents decreased cell viability in three days in a dose-dependent manner. The

addition of GC-1 significantly increased the efficacy of these agents at lower concentrations. For

example, 1 μM buparlisib was required to reduce the cell viability below 50% in the absence of

GC-1, but when combined with 10 nM GC-1 this could be achieved with 0.1-0.5 μM buparlisib

(**Fig. 3A**). Similar trends were observed with all four therapeutics and across the panel of cell

lines used.

GC-1 enhances the effects of therapeutic agents on cell migration

In addition to reducing cell viability, many therapeutics target pathways that are involved

in cell migration. To further evaluate the ability of GC-1 to enhance the efficacy of select

therapeutics, cells were treated with the indicated inhibitors at concentrations determined to be

effective based on the cell viability assays (**Fig. 3**), and migration was observed by assessing

wound closure. Each inhibitor was able to successfully prevent wound closure, and this effect

was increased when each of the inhibitors were combined with GC-1, with the exception of

buparlisib in the KTC-2 cells (**Fig. 4B-E**). Notably, GC-1 alone was able to prevent wound

closure of all tumor cell lines with similar success as each inhibitor alone (**Fig. 4A-E**). These

findings are similar to what we previously observed in experiments using endogenous thyroid

hormone, T_3, indicating GC-1 is a suitable alternative for TRβ activation and mitigation of

malignant phenotype.

GC-1 blocks tumorsphere outgrowth and increases the efficacy of therapeutic agents.

Current treatment options for ATC primarily target hyperproliferative cells, but aggressive

cancers like ATC are enriched for long-lived cancer stem cells that are thought to be a major

mechanism for therapy resistance and tumor recurrence. The signaling pathways that govern

stemness and proliferation are separate and frequently oppose each other. We previously

demonstrated that ligand-activated TRβ regulates genes related to stemness in ATC cells and

that GC-1 could block thyrosphere outgrowth in our modified ATC cells (**Fig 1D**). We therefore

tested the hypothesis that GC-1 could prevent thyrosphere outgrowth after an initial treatment

with an inhibitor that targets proliferation.

ATC cells were treated with the indicated inhibitor or vehicle for three days, followed by

re-plating in non-adherent serum-free conditions. We again observed that GC-1 alone could

reduce thyrosphere formation in unmodified ATC cell lines (**Fig. 5A-C**). None of the inhibitors

could completely prevent thyrosphere outgrowth when used alone; however, sorafenib was the

most effective overall (**Fig. 5C**). Strikingly, the addition of GC-1 upon removal of each inhibitor

blocked nearly all thyrosphere outgrowth. This effect was consistent across all four cell lines and

all three inhibitors used in this experiment. This result suggests that GC-1 may be particularly

effective at blocking cancer stem cell expansion

GC-1 promotes re-differentiation of ATC cell lines.

In transduced cells, TRβ activation with T_3 reduces the aggressive phenotype partially

through induction of re-differentiation in ATC cell. Therefore, we investigated the potential for

GC-1 alone to promote re-differentiation in parental SW1736 cells. Gene expression analysis

using RT-qPCR revealed that 24 hours of GC-1 exposure increased levels of thyroid-specific

gene transcripts (**Fig. 6A**). This transcriptomic program is summarized by the thyroid

differentiation score (TDS), which significantly increased in the GC-1-treated cells (**Fig. 6B**).

Additionally, GC-1 alone decreased the overall number of CD24$^-$/CD44$^+$ cells present in the

monolayer population as evident by RT-qPCR (**Sup. Fig. 1**). We also observed a decrease in

the ATC markers *ALCAM* (CD166) and *ALDH1A1* (**Sup. Fig. 1**).

Intriguingly, *SLC5A5* expression was greatly increased with 24 hours of GC-1 (**Fig. 6A**).

Therefore, we further investigated NIS protein levels after 48 hours of GC-1 treatment. Not only

were nascent and mature, glycosylated NIS protein significantly increased following GC-1

treatment (**Fig. 6C-D**), iodide uptake increased as well (**Fig. 6E**). This indicated that the

increased NIS protein was functional. These findings reveal a novel function of GC-1 to induce

re-differentiation in an ATC cell line model and offers a promising target to increase NIS levels

and iodide uptake.

DISCUSSION:

Nuclear receptors (NRs) are master regulators of growth and differentiation and are

increasingly recognized as diagnostic and therapeutic targets in thyroid, breast, and other

hormone-dependent cancers. Yet, the actions of non-steroidal NRs in these cancers are not

well characterized (19).There is compelling evidence that loss of expression of TRβ, a member

of the thyroid hormone receptor (TR) family, through genomic modifications and epigenetic

silencing is characteristic of thyroid, breast and other endocrine-related tumors (20-26).

Multiple lines of evidence demonstrate the tumor suppressive activity of TRβ in thyroid

cancer. Numerous pre-clinical studies demonstrate the potent anti-tumorigenic effect of robust

expression of the TRβ gene. Re-expression of silenced TRβ using demethylating agents delays

thyroid tumor progression *in vivo* (27), mice expressing a c-terminal frameshift of TRβ (*Thrb^{PV}*)

spontaneously develop follicular thyroid cancer (28), and *Thrb$^{PV/PV}$ Kras^{G12D}* mice develop

thyroid tumors with features of de-differentiated thyroid cancer (29). Our studies in anaplastic

thyroid cancer cells revealed that activation of TRβ induces a tumor suppression transcriptomic

program and reduces the aggressive phenotype (9,10).

Agonists have been developed to target tissue specific expression of the most abundant

isomers (30).   Selective activation of TRβ in pre-clinical and clinical studies have yielded

exciting results with great promise for novel interventions. Patients who were rendered

hypothyroxinemic and treated with T_3 alone had longer survival with end-stage solid tumors (31)

and T_3 treatment enhanced chemosensitivity (32). Following tyrosine kinase inhibitor therapy,

patients demonstrated improved survival with treatment of hypothyroidism (33). Mimetics of T_3

that selectively act as TRβ agonists have been clinically successful for treatment of metabolic

disorders, hyperlipidemia, hypercholesterolemia and non-alcoholic steatohepatitis without the

cardiovascular effects mediated by TRα (11,12). Those in clinical trials include TRβ selective

sobetirome (GC-1) (14) with comparable transcriptomic effects of T_3 (16). Our findings

demonstrate that GC-1 alone decreases the ATC aggressive phenotype in transduced cells in

which TRβ is re-expressed and parental cell lines with diverse genetic backgrounds. These

results emphasize the generalizability of GC-1 to activate a TRβ tumor suppression program.

In anaplastic thyroid cancer tumors, several genetic alterations and disrupted signaling

pathways are usually found. Therefore, the treatment aim is to inhibit or disrupt several

pathways with combination therapies that block cell survival, proliferation typically by targeting

the MAPK, PI3K, and cell cycle pathways (7,34). There are many therapeutics that are in use in

combination or alone including multi-kinase inhibitors, epigenetic modulators, apoptosis-

inducing agents, and others. However, to date, none have been successful in long-term

treatment. Development of resistance and toxicity are limitations to their use.

In this study, we tested representative drugs that targeted the PI3K, MAPK, and cell

cycle pathways to assess whether GC-1 could alter the effectiveness of these drugs on their

targets.   Since we have demonstrated that activation of TRβ in ATC represses the PI3K

pathway through genomic mechanisms (10,18), GC-1 with either buparlisib or alpelisib would be

suppressing PI3K activity in concert. However, TRβ does not directly impact MAPK or cell cycle

signaling through transcriptomic modulation so GC-1 in combination with either sorafenib or

palbociclib demonstrated concordant effects.   In all studies, GC-1 reduced the effective

concentrations required to observe a reduction in cell viability. Not surprisingly, not all effects

were noted for cell migration as these events are mediated by distinct cell signaling pathways

modulated by TRβ. One of our most intriguing observations is the impact of GC-1 alone and in

combination with inhibitors on cancer stem cell growth in thyrosphere formation. In all ATC cell

types, GC-1 alone significantly reduced thyrosphere formation. Remarkably, GC-1 in

combination with any of the drug treatments nearly eliminated thyrosphere formation.

GC-1 induced re-differentiation in the SW1736 cell line in 24 hours, demonstrating

potent transcriptional reprogramming. While we have previously demonstrated the potential for

TRβ to promote a more differentiated phenotype in ATC cells, this is the first time we have been

able to demonstrate the profound impact of GC-1 in a parental cell line with low TRβ expression.

Importantly, we observed a concomitant increase in the sodium iodide symporter that led to a

higher intake of iodide. Since PDTC and ATC tumors often exhibit a marked decrease in NIS

expression and resistance to radioactive iodide (RAI) therapy, the results presented here raise a

provocative implication for the use of GC-1 in combination with RAI.

Our findings in this study demonstrate for the first time that selective activation of TRβ

with GC-1 (Sobetirome) can reduce the aggressive phenotype, reduce the cancer stemness in

anaplastic thyroid cancer cells, increase the sensitivity of these cells to therapeutic agents, and

promote re-differentiation and iodide uptake. These studies provide the foundation for further

establishing the broad impact of GC-1 alone and in combination with various therapeutics in

mitigating toxicity and the development of resistance to treatment.

ACKNOWLEDGEMENTS:

The research reported here was supported by grants from National Institutes of Health U54

GM115516 for the Northern New England Clinical and Translational Research Network;

National Cancer Institute 1F99CA245796-01; UVM Cancer Center-Lake Champlain Cancer

Research Organization (C3) 12577-21; and UVM Larner College of Medicine. SW1736 and

KTC-2 cell lines were generously provided by Dr. John Copland III (Mayo Clinic) and 8505C and

OCUT-2 cells were generously provided by Dr. Rebecca Schweppe (University of Colorado).

Human cell line authentication, NextGen sequencing, automated DNA sequencing and

molecular imaging was performed in the Vermont Integrative Genomics Resource supported by

the University of Vermont Cancer Center, Lake Champlain Cancer Research Organization, and

the UVM Larner College of Medicine. Additional human cell line authentication was performed

by the CU Cancer Center Tissue Culture Shared Resource supported by National Cancer

Institute P30CA046934.

Authors' Contribution Statement: Noelle E. Gillis: Conceptualization, Formal analysis,

Investigation, Writing – Review & editing. **Cole D. Davidson**: Conceptualization, Formal

analysis, Investigation, Writing – Review & editing, Visualization. **Lauren M. Cozzens:** Formal

analysis, Investigation. **Emily Wilson:** Investigation. **Eric L. Bolf:** Conceptualization, – Review

& editing. **Jennifer A. Tomczak:** Investigation, Writing – Review & editing. **Frances E. Carr:**

Conceptualization, Resources, Writing-Review & editing, Supervision, Project administration,

Funding acquisition.

Authors' Disclosure Statement: The authors declare they have no conflict of interest.

Funding Statement: The research reported here was supported by grants from National

Institutes of Health U54 GM115516 for the Northern New England Clinical and Translational

Research Network; National Cancer Institute 1F99CA245796-01; UVM Cancer Center-Lake

Champlain Cancer Research Organization (C3) 12577-21; and UVM Larner College of

Medicine.

REFERENCES

1. Siegel RL, Miller KD, Jemal A. Cancer Statistics, 2017. CA Cancer J Clin. 2017;67(1):7-30.

2. Pereira M, Williams VL, Hallanger Johnson J, Valderrabano P. Thyroid Cancer Incidence
Trends in the United States: Association with Changes in Professional Guideline
Recommendations. Thyroid. 2020;30(8):1132-1140.

3. Glaser SM, Mandish SF, Gill BS, Balasubramani GK, Clump DA, Beriwal S. Anaplastic
thyroid cancer: Prognostic factors, patterns of care, and overall survival. Head Neck. 2016;38
Suppl 1:E2083-2090.

4. Prasongsook N, Kumar A, Chintakuntlawar AV, Foote RL, Kasperbauer J, Molina J, Garces
Y, Ma D, Wittich MAN, Rubin J, Richardson R, Morris J, Hay I, Fatourechi V, McIver B, Ryder
M, Thompson G, Grant C, Richards M, Sebo TJ, Rivera M, Suman V, Jenkins SM, Smallridge
RC, Bible KC. Survival in Response to Multimodal Therapy in Anaplastic Thyroid Cancer. J Clin
Endocrinol Metab. 2017;102(12):4506-4514.

5. De Leo S, Trevisan M, Fugazzola L. Recent advances in the management of anaplastic
thyroid cancer. Thyroid Res. 2020;13(1):17.

6. Gentile D, Orlandi P, Banchi M, Bocci G. Preclinical and clinical combination therapies in the
treatment of anaplastic thyroid cancer. Med Oncol. 2020;37(3):19.

7. Molinaro E, Romei C, Biagini A, Sabini E, Agate L, Mazzeo S, Materazzi G, Sellari-
Franceschini S, Ribechini A, Torregrossa L, Basolo F, Vitti P, Elisei R. Anaplastic thyroid
carcinoma: from clinicopathology to genetics and advanced therapies. Nature Reviews
Endocrinology. 2017;13:644.

8. Subbiah V, Kreitman RJ, Wainberg ZA, Cho JY, Schellens JHM, Soria JC, Wen PY, Zielinski
C, Cabanillas ME, Urbanowitz G, Mookerjee B, Wang D, Rangwala F, Keam B. Dabrafenib and
Trametinib Treatment in Patients With Locally Advanced or Metastatic BRAF V600-Mutant
Anaplastic Thyroid Cancer. J Clin Oncol. 2018;36(1):7-13.

9. Carr FE, Tai PW, Barnum MS, Gillis NE, Evans KG, Taber TH, White JH, Tomczak JA,
Jaworski DM, Zaidi SK, Lian JB, Stein JL, Stein GS. Thyroid Hormone Receptor-beta (TRbeta)
Mediates Runt-Related Transcription Factor 2 (Runx2) Expression in Thyroid Cancer Cells: A
Novel Signaling Pathway in Thyroid Cancer. Endocrinology. 2016;157(8):3278-3292.

10. Bolf EL, Gillis NE, Davidson CD, Rodriguez PD, Cozzens L, Tomczak JA, Frietze S, Carr
FE. Thyroid Hormone Receptor Beta Induces a Tumor-Suppressive Program in Anaplastic
Thyroid Cancer. Mol Cancer Res. 2020;18(10):1443-1452.

11. Trost SU, Swanson E, Gloss B, Wang-Iverson DB, Zhang H, Volodarsky T, Grover GJ,
Baxter JD, Chiellini G, Scanlan TS, Dillmann WH. The thyroid hormone receptor-beta-selective
agonist GC-1 differentially affects plasma lipids and cardiac activity. Endocrinology.
2000;141(9):3057-3064.

12. Grover GJ, Mellstrom K, Ye L, Malm J, Li YL, Bladh LG, Sleph PG, Smith MA, George R,
Vennstrom B, Mookhtiar K, Horvath R, Speelman J, Egan D, Baxter JD. Selective thyroid
hormone receptor-beta activation: a strategy for reduction of weight, cholesterol, and lipoprotein
(a) with reduced cardiovascular liability. Proc Natl Acad Sci U S A. 2003;100(17):10067-10072.

13. Kelly MJ, Pietranico-Cole S, Larigan JD, Haynes NE, Reynolds CH, Scott N, Vermeulen J,
Dvorozniak M, Conde-Knape K, Huang KS, So SS, Thakkar K, Qian Y, Banner B, Mennona F,
Danzi S, Klein I, Taub R, Tilley J. Discovery of 2-[3,5-dichloro-4-(5-isopropyl-6-oxo-1,6-
dihydropyridazin-3-yloxy)phenyl]-3,5-dio xo-2,3,4,5-tetrahydro[1,2,4]triazine-6-carbonitrile (MGL-
3196), a Highly Selective Thyroid Hormone Receptor beta agonist in clinical trials for the
treatment of dyslipidemia. J Med Chem. 2014;57(10):3912-3923.

14. Elbers LP, Kastelein JJ, Sjouke B. Thyroid Hormone Mimetics: the Past, Current Status and
Future Challenges. Curr Atheroscler Rep. 2016;18(3):14.

15. Scanlan TS, Yoshihara HA, Nguyen NH, Chiellini G. Selective thyromimetics: tissue-
selective thyroid hormone analogs. Curr Opin Drug Discov Devel. 2001;4(5):614-622.

16. Yuan C, Lin JZ, Sieglaff DH, Ayers SD, Denoto-Reynolds F, Baxter JD, Webb P. Identical
gene regulation patterns of T3 and selective thyroid hormone receptor modulator GC-1.
Endocrinology. 2012;153(1):501-511.

17. Landa I, Ibrahimpasic T, Boucai L, Sinha R, Knauf JA, Shah RH, Dogan S, Ricarte-Filho JC,
Krishnamoorthy GP, Xu B, Schultz N, Berger MF, Sander C, Taylor BS, Ghossein R, Ganly I,
Fagin JA. Genomic and transcriptomic hallmarks of poorly differentiated and anaplastic thyroid
cancers. The Journal of Clinical Investigation. 2016;126(3):1052-1066.

18. Davidson CD, Bolf EL, Gillis NE, Cozzens LM, Tomczak JA, Carr FE. Thyroid hormone
receptor beta inhibits the PI3K-Akt-mTOR signaling axis in anaplastic thyroid cancer cell lines
via genomic mechanisms. Journal of the Endocrine Society. 2021.

19. Doan TB, Graham JD, Clarke CL. Emerging functional roles of nuclear receptors in breast
cancer. J Mol Endocrinol. 2017;58(3):R169-R190.

20. Silva JM, Domínguez G, González-Sancho JM, García JM, Silva J, García-Andrade C,
Navarro A, Muñoz A, Bonilla F. Expression of thyroid hormone receptor/erbA genes is altered in
human breast cancer. Oncogene. 2002;21:4307.

21. Kim W, Cheng S. Thyroid hormone receptors and cancer. Vol 1830. Biochim Biophys
Acta2013:3928-3936.

22. Aranda A, Martinez-Iglesias O, Ruiz-Llorente L, Garcia-Carpizo V, Zambrano A. Thyroid
receptor: roles in cancer. Trends Endocrinol Metab. 2009;20(7):318-324.

23. Jezequel P, Campone M, Gouraud W, Guerin-Charbonnel C, Leux C, Ricolleau G, Campion
L. bc-GenExMiner: an easy-to-use online platform for gene prognostic analyses in breast
cancer. Breast Cancer Res Treat. 2012;131(3):765-775.

24. Jezequel P, Frenel JS, Campion L, Guerin-Charbonnel C, Gouraud W, Ricolleau G,
Campone M. bc-GenExMiner 3.0: new mining module computes breast cancer gene expression
correlation analyses. Database (Oxford). 2013;2013:bas060.

25. Wojcicka A, Piekielko–Witkowska A, Kedzierska H, Rybicka B, Poplawski P, Boguslawska J,
Master A, Nauman A. Epigenetic Regulation of Thyroid Hormone Receptor Beta in Renal
Cancer. PLoS One. 2014;9(5):e97624.

26. Park JW, Zhao L, Willingham M, Cheng S-y. Oncogenic mutations of thyroid hormone
receptor β. Oncotarget. 2015;6(10):8115-8131.

27. Kim WG, Zhu X, Kim DW, Zhang L, Kebebew E, Cheng SY. Reactivation of the silenced
thyroid hormone receptor beta gene expression delays thyroid tumor progression.
Endocrinology. 2013;154(1):25-35.

28. Suzuki H, Willingham MC, Cheng SY. Mice with a mutation in the thyroid hormone receptor
beta gene spontaneously develop thyroid carcinoma: a mouse model of thyroid carcinogenesis.
Thyroid. 2002;12(11):963-969.

29. Zhu X, Zhao L, Park JW, Willingham MC, Cheng SY. Synergistic signaling of KRAS and
thyroid hormone receptor beta mutants promotes undifferentiated thyroid cancer through MYC
up-regulation. Neoplasia. 2014;16(9):757-769.

30. Davis PJ, Tang HY, Hercbergs A, Lin HY, Keating KA, Mousa SA. Bioactivity of Thyroid
Hormone Analogs at Cancer Cells. Front Endocrinol (Lausanne). 2018;9:739.

31. Hercbergs A, Johnson RE, Ashur-Fabian O, Garfield DH, Davis PJ. Medically induced
euthyroid hypothyroxinemia may extend survival in compassionate need cancer patients: an
observational study. Oncologist. 2015;20(1):72-76.

32. Huang J, Jin L, Ji G, Xing L, Xu C, Xiong X, Li H, Wu K, Ren G, Kong L. Implication from
thyroid function decreasing during chemotherapy in breast cancer patients: chemosensitization
role of triiodothyronine. BMC Cancer. 2013;13:334.

33. Lechner MG, Vyas CM, Hamnvik OR, Alexander EK, Larsen PR, Choueiri TK, Angell TE.
Hypothyroidism During Tyrosine Kinase Inhibitor Therapy Is Associated with Longer Survival in
Patients with Advanced Nonthyroidal Cancers. Thyroid : official journal of the American Thyroid
Association. 2018;28(4):445-453.

34. Alobuia W, Gillis A, Kebebew E. Contemporary Management of Anaplastic Thyroid Cancer.
Curr Treat Options Oncol. 2020;21(10):78.

Figure 1. GC-1 blocks tumorigenic phenotypes in ATC cells transduced with TRβ. A.

Growth curves demonstrate that SW-TRβ growth is significantly decreased after treatment with

GC-1 for 4 days (n=3). Significance (* $p<0.05$) was determined by a two-way ANOVA and

Tukey's multiple comparisons test. **B**. Area under the curve (AUC) analysis of growth curve

data. Significance (* $p<0.05$) was determined by t-test. **C.** Treatment with GC-1 slows migration

of SW-TRβ and SW-EV cells measured by scratch assay. Significance (* $p<0.05$) was

determined by t-test. D. GC-1 repressed thyrosphere formation in SW-TRβ and SW-EV cells

(n=4). Significance (* $p<0.05$) was determined by t-test. E. Immunoblot demonstrates that 5

days of GC-1 treatment induces apoptosis in SW-TRβ cells assessed by cleavage of PARP and

caspase 3 (n=3).

Figure 2: GC-1 slows growth of parental ATC cells. A. Summary of mutated genes in four

ATC cell lines implemented in this study (17). B-E. Relative cell growth of unmodified ATC cell

lines was measured by cell counting assay after 4 days of treatment with 10 nM GC-1 or 10 nM

T_3. Significance (* $p<0.05$) was determined by a two-way ANOVA followed by Dunnett's multiple

comparisons test; 3 independent experiments were performed per treatment group.

Figure 3. GC-1 increases efficacy of inhibitors. Cell viability of unmodified ATC cell lines was

measured by SRB assay after 3 days of treatment with increasing concentrations buparlisib or

alpelisib (A), sorafenib (B), or palbociclib (C) simultaneously in combination with 10 nM GC-1.

Significance (* $p<0.05$) was determined by a two-way ANOVA followed by Sidak's multiple

comparisons test; 3 independent experiments were performed per each treatment group.

Figure 4. GC-1 enhances the effects of inhibitors on cell migration. Cell migration was

measured by scratch assay. Representative images demonstrate reduced scratch closure after

two days when ATC cells are treated with GC-1 (A). Area under the curve (AUC) analysis

shows SW1736 (B), 8505C (C), OCUT2 (D), and KTC-2 (E) cells all exhibit lower rates of

scratch closer after treatment with GC-1 alone and further reduced scratch closure rates after

combined treatment with GC-1 and inhibitors. Significance (* p<0.05) was determined by t-test.

Figure 5. GC-1 blocks thyrosphere outgrowth and increases the efficacy of therapeutic

agents. Thyrosphere growth was determined for ATC cells after 3 days of treatment with 0.5 µM

buparlisib or alpelisib (A), 10 µM sorafenib (B), or 1 nM palbociclib (C) under adherent culture

conditions, followed by plating in conditions for spheroid growth in the presence of 10 nM GC-1

for 7 days. GC-1 alone and each therapeutic agent significantly blocked sphere formation in all

cell types. GC-1 in addition to each therapeutic agent further inhibited or completely blocked

sphere formation. Significance (* p<0.05) was determined by two-way ANOVA followed by

Sidak's multiple comparisons test; 3 independent experiments were performed per each

treatment group; n=3 per analyses.

Figure 6. GC-1 induces expression of redifferentiation markers in SW1736 cells. A. RT-

qPCR for thyroid differentiation markers was performed on SW1736 cells treated with or without

10 nM GC-1 for 24 hours. mRNA levels in cells treated with GC-1 are relative to the untreated

cells. Significance (* p<0.05) was determined by Student's unpaired t test; 3 independent

experiments were performed per treatment group. B. Thyroid differentiation score (TDS) was

calculated using the expression level of the genes in panel A, excluding *THRB*, as previously

described (10, 17). Significance (* p<0.0001) was determined by Student's unpaired t test. C.

SW1736 cells were treated with or without 10 nM GC-1 in triplicate for 48 hours prior to protein

extraction and analysis via immunoblot. Normal-like thyroid Nthy-ORI cells were used as a

positive control. D. Relative NIS/GAPDH protein levels from (C) were quantified via ImageJ, and

significance (* p<0.01) was determined by Student's unpaired t test. E. Iodide uptake was

measured after 48 hours of GC-1 treatment. Significance was determined by Student's unpaired

t test. (* p<0.001).

Supplemental Figure 1. RT-qPCR for markers of stemness and proliferation was performed on

SW1736 cells treated with or without 10 nM GC-1 for 24 hours. mRNA levels in cells treated

with GC-1 are relative to the untreated cells. Significance (* $p<0.05$) was determined by

Student's unpaired t test; 3 independent experiments were performed per treatment group.

Gene Function	Gene Name	Cell Line			
		SW1736	8505C	OCUT-2	KTC-2
		Mutation Type			
MAPK Signaling	BRAF	V600E	V600E	V600E	V600E
Telomerase	TERT (promoter)	124C-T	146C-T	146C-T	124C-T
DNA Homeostasis	TP53	Trunc.	Miss.		
PI3K Signaling	PIK3CA			Kinase Domain	
Cell-Cell Contact	NF2		Trunc.		
DNA Homeostasis	ATM		Trunc.		
TSH Signaling	TSHR	Miss.			
Histone Methyltransferase	KMT2C				Miss.
Histone Methyltransferase	KMT2D				Trunc.
Histone Acetyltransferase	EP300	Trunc.			
SWI/SNF Complex	ARID1A				In Frame
SWI/SNF Complex	SMARCA4		Miss.		

A

Day 0   Day 1   Day 2

vehicle

GC-1

B   **SW1736**

AUC Units

GC-1: - +   - +   - +   - +
   Vehicle Buparlisib Palbociclib Sorafenib

C   **8505C**

AUC Units

GC-1: - +   - +   - +   - +
   Vehicle Buparlisib Palbociclib Sorafenib

D   **OCUT2**

AUC Units

GC-1: - +   - +   - +   - +
   Vehicle Alpelisib Palbociclib Sorafenib

E   **KTC-2**

AUC Units

GC-1: - +   - +   - +   - +
   Vehicle Buparlisib Palbociclib Sorafenib

A

DIO1	DIO2	DUOX1	DUOX2	FOXE1	NKX2-1	SLC5A5
*	ns	*	*	ns	*	*

GC-1: - +

SLC26A4	PAX8	TG	THRA	THRB	TPO	TSHR
ns	*	*	ns	*	*	*

GC-1: - +

B

Thyroid Differentiation Score (TDS)

TDS

*

GC-1: - +

C

SW1736   Nthy-ORI

GC-1: - - - + + +   -

Glycosylated NIS   68 kDa

Nascent NIS   55 kDa

GAPDH   37 kDa

D

NIS Protein Level

Relative NIS/GAPDH

*

GC-1: - +

E

Iodide Uptake

pmol Iodide

*

GC-1: - +