Emerging single cell endothelial heterogeneity supports sprouting tumour angiogenesis and growth

Ken Matsumoto[1,2], Florian Rambow[3,4], Fabio Stanchi[1,2], Thomas Mathivet[5], Junbin Qian[6,7], Wolfgang Giese[8], Liqun He[9], Diether Lambrechts[6,7], Bin Zhou[10], Christer Betsholtz[9,11], Jean-Christophe Marine[3,4], Holger Gerhardt[1, 2, 8, 12, 13 *]

[1]Vascular Patterning laboratory, VIB Center for Cancer Biology, Leuven, Belgium

[2]Vascular Patterning laboratory, Department of Oncology, KU Leuven, Leuven, Belgium

[3]Laboratory for Molecular Cancer Biology, VIB Center for Cancer Biology, Leuven, Belgium

[4]Laboratory for Molecular Cancer Biology, Department of Oncology, KU Leuven, Leuven, Belgium

[5]Laboratoire de Neurosciences Expérimentales er Cliniques, Université de Poitiers, INSERM U1084, Poitiers, France

[6]Laboratory of Translational Genetics, VIB Center for Cancer Biology, Leuven, Belgium

[7]Laboratory of Translational Genetics, Center for Human Genetics, KU Leuven, Leuven, Belgium

[8]Integrative Vascular Biology, Max-Delbrück Center for Molecular Medicine, 13125 Berlin, Germany

[9]Department of Immunology, Genetics and Pathology, Rudbeck Laboratory, Uppsala University, Uppsala, Sweden

[10]State key Laboratory of Cell Biology, CAS center for Excellence in Molecular Cell Science, Institute of Biochemistry and Cell Biology, Shanghai Institutes for Biological Sciences, University of Chinese Academy of Sciences, Chinese Academic of Sciences,

Shanghai 200031, China.

[11]Integrated Cardio Metabolic Centre (ICMC), Karolinska Institute, Huddinge, Sweden

[12]DZHK (German Center for Cardiovascular Research) Partner site Berlin

[13]Berlin Institute of Health (BIH), Berlin, Germany

[*]Correspondence: H. Gerhardt, email: holger.gerhardt@mdc-berlin.de

Keywords: *Csf1r*, lineage tracing, tumour angiogenesis, intravital imaging, single cell

transcriptional profiling

Conflict of Interest: The authors have declared that no conflict of interest exists.

SUMMARY

Blood vessels supplying tumors are often dysfunctional and generally heterogeneous. The

mechanisms underlying this heterogeneity remain poorly understood. Here, using

multicolor lineage tracing, *in vivo* time-lapse imaging and single cell RNA sequencing in

a mouse glioma model, we identify tumour-specific blood endothelial cells that originate

from cells expressing the receptor for colony stimulating factor 1, *Csf1r*, a cytokine which

controls macrophage biology. These *Csf1r* lineage endothelial cells (CLECs) form up to

10% of the tumour vasculature and express, besides classical blood endothelial cell

markers, a gene signature that is distinct from brain endothelium but shares similarities

with lymphatic endothelial cell populations. *in silico* analysis of pan-cancer single cell

RNAseq datasets highlights the presence of a comparable subpopulation in the

endothelium of a wide spectrum of human tumours. We show that CLECs actively

contribute to sprouting and remodeling of tumour blood vessels and that selective

depletion of CLECs reduces vascular branching and tumour growth. Our findings indicate

that a non-tumour resident Csf1r-positive population is recruited to tumours,

differentiates into blood endothelial cells to contribute to vascularization and, thereby,

tumour growth.

INTRODUCTION

Endothelial cells form a single cell layer lining the inner walls of blood vessels and play

critical roles in organ homeostasis and disease progression. Once formed, following

embryonic and early post-natal development, blood vessels retain a high level of

adaptability to meet changing metabolic or hemodynamic requirements or enable further

tissue growth including in tumours. How local vascular networks respond to these

adaptive challenges, rapidly expand, remodel and reestablish homeostasis remains a

timely research topic, with many open questions still unresolved. In particular in

pathologies, knowledge of the precise nature and origin of new vessel formation,

endothelial activation and differentiation steps, and of the mechanisms driving vessel

dysmorphia, or preventing effective revascularization in ischemic diabetic complication

is critical for new therapeutic approaches. New blood vessel formation in the adult was

long believed to arise exclusively by sprouting and proliferation of endothelial cells from

local blood vessels without any de novo differentiation from progenitor cells. However,

Asahara and coworkers isolated mononuclear cells from human peripheral blood and

identified circulating endothelial progenitor cells (EPCs) [1]. These cells were shown to

derive from the bone marrow and contribute to endothelial cells in blood vessels in

hindlimb ischemia and tumour xenograft mouse model [2,3]. By fluorescent in situ

hybridization with sex chromosome-specific probes in patients with cancer after bone

marrow transplantation, Peters and coworkers demonstrated that bone marrow-derived

stem cells contribute to endothelial cells in human tumour endothelium [4]. The resulting

concept of vasculogenesis in the adult raised the prospect of novel therapeutic approaches

for ischemic vascular disease and for targeting tumour angiogenesis [5,6]. Interestingly,

EPCs also contributed to newly forming lymphatic vessels [7,8]. A vast number of

publications since have supported or challenged these initial discoveries, and the

occurrence, origin and significance of EPCs continues to be a matter of debate [9-12].

Whereas EPCs and other circulating progenitor cells were long thought to origin from the

bone marrow, recent studies challenged this idea and suggested the existence of a

different yet unknown vascular niche for endothelial progenitor cells that can contribute

to blood vessel formation[13].

RESULTS

A Csf1r cell lineage contributes to blood vascular endothelial cells during glioma

growth in mice.

In order to study dynamic macrophage recruitment and blood vessel patterning *in vivo*

during glioma growth, we adapted a surgical cranial window model for two-photon

microscopy allowing repeated visualization of the same animal over time. We generated

spheroids of syngeneic C57BL/6 mouse glioma cells (CT2A or GL261) [14,15] modified to

stably express blue fluorescent protein (BFP), and injected them into the mouse cortex,

followed by implantation of a glass coverslip [16]. By crossing transgenic *Csf1r (Colony*

stimulating factor 1 receptor)– specific tamoxifen-inducible Cre driver mice [17] with the

mTmG Cre reporter mouse line [18], we induced membrane-targeted GFP expression in

Csf1r-expressing cells and used this stable lineage trace to follow the cell population over

time (Fig. 1a, b). As expected, the vast majority of tumour associated macrophages were

labelled by GFP in line with their *Csf1r*-dependent recruitment and expansion[19].

Unexpectedly, however, through the combination of longitudinal live two-photon

imaging and lineage tracing, we discovered GFP-positive cells contributing to the tumour

vasculature during glioma growth (Fig. 1c). These *Csf1r* lineage endothelial cells

(CLECs) frequently emerged at tip cell positions of the growing vasculature, heading

blood vessel sprouts in the mouse glioma (Fig. 1d, Supplementary Movie1). Additionally,

CLECs were found bridging adjacent capillaries during anastomosis (Fig. 1e).

Fluorescent dextran injection confirmed that CLECs contributed to - and were

incorporated in the endothelial lining of - functionally perfused blood vessels including

lumenized sprouts headed by endothelial tip cells (Fig. 1f). Counterstaining on fixed

samples identified that all cells forming blood vessels including the lineage-traced CLECs

expressed typical endothelial markers (CDH5, CD31) and lacked macrophage and

myeloid cell markers (F4/80, CD45) (Fig. 1g, h, Extended Data Fig. 1). BFP was not

expressed in CLECs, demonstrating that they were not derived from glioma cells (Fig.

1g).

To gain more insight into the nature and significance of CLECs, we isolated and

quantified the cell population in the mouse glioma model. By 4 weeks of tumour

development (close to the specified humane endpoint of 5 weeks) up to 10% of the tumour

vasculature was comprised of cells carrying the *Csfr1* lineage trace. Similar frequencies

were found in tumours derived from either of the mouse glioma cell lines, CT2A (Fig. 2a,

b), or GL261 (Fig. 2c). Polyclonal antibody staining indicated that *Csf1r* may also be

expressed in some brain capillary endothelial cells [20,21]. Recent single cell sequencing of

adult mouse brain endothelial cells however did not confirm this notion[22], suggesting that

CLECs are unlikely to originate from brain endothelium. Indeed, tamoxifen mediated

activation of our transgenic *Csfr1* lineage trace did not label brain endothelial cells in

mice lacking tumours, sham operated as well as in the healthy brain parenchyma

surrounding implanted tumours (Fig. 2d, e, f). *Csf1r* is important for the embryonic

development of microglia and its expression is also detected in adult microglia by single

cell RNA sequencing. Surprisingly however, Csf1r-Mer-iCre-Mer mediated

recombination of the Cre-Reporter expression was also not observed in microglia of the

healthy adult mouse brain. Irrespective of this apparent discrepancy between reported

microglial *Csf1r* expression and the lack of Csf1r- Mer-iCre-Mer mediated recombination

in microglia, these data also exclude microglia as a potential source of CLECs.

Furthermore, we observed CLECs also in tumours originating from B16F1 melanoma

cells implanted under the skin; thus, this *Csfr1*-lineage endothelial cell contribution not

only occurs in brain tumours with microglia in their environment, but also in other tumour

types and locations (Fig. 2g, h). Notably, we also found *Csf1r*-lineage lymphatic

endothelial cells in B16F1 skin melanomas indicating that CLECs can contribute to both

blood and lymphatic vessels (Extended Data Fig. 2).

CLECs do not originate from hematopoietic niche in bone-marrow or spleen.

Given the abundant labelling of macrophages and the important role of *Csf1r* in the

recruitment of bone-marrow derived monocytes and differentiated macrophages, we

asked whether CLECs are also derived from the bone-marrow. To test this possibility, we

generated bone marrow chimeras. Flow cytometric analysis confirmed 97.3 ± 1.6 %

reconstitution of wild-type recipient bone marrow by transplanted *Csf1r-Mer2.Cre^{mTmG}*

bone marrow after 8 weeks post transplantation (Extended Data Fig. 3). Analysis of

glioma blood vessels in these bone marrow chimeras (BM$^{Csf1r-Mer2.Cre-mTmG}$) failed to

identify bone marrow derived CLECs, although the numbers of tumour macrophage were

similar to controls (Fig. 3a). Conversely, chimeras of wild-type donor bone marrow

transplanted into *Csf1r-Mer2.Cre^{mTmG}* recipients (*Csf1r-Mer2.Cre^{mTmG}*::BM^{WT}),

displayed similar CLECs numbers in glioma vessels as observed earlier in non-chimeric

Csf1r-Mer2.Cre^mTmG mice (Fig. 3b). These results were confirmed by intravital imaging

in the glioma of *Csf1r-Mer2.Cre^mTmG*::BM^WT (Fig. 3c, Supplemental Movie 2). Thus, the

bone marrow chimera and lineage tracing experiments strongly suggest that CLECs do

not originate from bone marrow-derived cells, nor do they represent previously suggested

transdifferentiation events from bone marrow-derived macrophages. Surgical removal of

the spleen also did not affect CLECs numbers, further ruling out spleen-derived

macrophages as source of CLECs (Extended Data Fig. 4).

CLECs regulate vascular patterning and support glioma growth.

In late-stage gliomas, CLECs preferentially localized to the peripheral tumor area,

correlating with the zone of most active angiogenesis (Extended Data Fig. 5). Real-time

intravital imaging of mouse gliomas also revealed that CLECs incorporation into vessels

is very dynamic. This implied that CLECs might have unique and potentially transient

functions during tumour vessel formation. To gain a better understanding of CLECs

behaviour during tumour growth, we developed dual-recombination combinatorial

genetics as a tool for CLECs tracing [23,24]. To achieve selective recombination in CLECs,

we used the *Cdh5* endothelial driver for expression of the Dre recombinase, and the same

transgenic mouse line as above expressing the tamoxifen inducible form of the Cre

recombinase under control of the *Csf1r* transgene. The combined action of these distinct

recombinases, Dre and Cre, allowed for deletion of two stop cassettes flanked by RoxP

and LoxP sites, respectively, located upstream of a tdTomato expression cassette (Fig.

4a). Using this strategy, we could selectively trace tdTomato-positive CLECs in mouse

gliomas following tamoxifen exposure (Fig. 4b). Note that the appearance of labelled

CLECs and blood vessel labelling differs from the images of the mTmG line, as the dual

Dre/Cre tdTomato reporter used lacks a general label of unrecombined cells. Fluorescent

dextran injection instead was used to label vascular lumen, highlighting dtTomato

positive CLECs that line the vessel lumen. As expected, counterstaining on fixed samples

demonstrated that CLECs express the prototypical endothelial marker, ERG (ETS-related

gene) (Fig. 4c). In order to assess whether CLECs contribute functionally to tumour

growth and vascular patterning in glioma, we used the same dual recombination strategy

to selectively ablate CLECs (Fig. 4d). In this model, CLECs expressed both GFP as well

as the diphtheria toxin receptor after tamoxifen induction enabling depletion of GFP-

positive CLECs by administration of diphtheria toxin (Fig. 4e, h). Strikingly, CLECs

ablation significantly decreased glioma growth (Fig. 4g, i). CLECs ablation also

decreased the vascular network length and the number of bifurcations in tumour vessels

(Fig. 4j, k, l), indicating that CLECs may promote tumour growth by supporting vessel

branching.

CLECs express a unique set of markers

To further understand the nature of tumour CLECs (T_CLECs), we performed single-cell

RNA sequencing (scRNA-seq) on isolated T_CLECs in the mouse glioma model. Single-

cell transcriptome analysis using Smart-seq2 [25] identified that the gene expression pattern

of T_CLECs was similar to lineage-negative tumour endothelial cells (T_ECs), but not

to tumour macrophages (T_MACs), as determined by UMAP (Uniform Manifold

Approximation and Projection) analysis (Fig. 5a). Single-cell differential expression

(SCDE) analysis [26] confirmed that T_CLECs expressed typical endothelial markers, but

lacked macrophage and myeloid cell markers (Fig. 5b). Unexpectedly, however,

T_CLECs showed no *Csf1r* expression, despite carrying the *Csf1r*-lineage marker (Fig.

5c). Beyond the overlapping gene expression pattern between T_CLECs and T_ECs,

SCDE analysis identified a unique signature that can distinctly define the T_CLECs

population; RNAs coding for 24 genes including three types of cell surface proteins

(*Aqp1: Aquaporin 1, Fabp4: Fatty acid binding protein 4, Kcnj8: Potassium voltage-*

gated channel subfamily j member 8) were significantly enriched in T_CLECs (Fig. 5d;

Extended Data Fig. 6; Supplementary Table 1). *Aqp1* is highly expressed in vascular

endothelial cells and was reported to contribute to tumour growth and angiogenesis [27].

Staining of lineage traced T_CLECs on fixed mouse glioma samples confirmed AQP1

expression (Extended Data Fig. 7).

The T_CLEC state is enriched for "Tip-like" and a lymphatic gene signature

Gene set enrichment analysis (GSEA) identified that the T_CLEC state shows significant

enrichment for a Tip-cell-like associated gene expression program (Fig. 5e), thus

supporting the morphological observations and their functional importance for branching

and tumour growth. Intriguingly, GSEA identified that T_CLECs were also enriched for

lymphatic gene signatures (Fig. 5e). AUCell analysis [28] of scRNA-seq data from multiple

human primary tumours identified subpopulations of endothelial cells that express the

T_CLECs marker gene set (Fig. 5f-k, 442 out of 7314 ECs are candidates: 6.0%;

Supplementary Table 2). These *bona-fide* T_CLECs were highly enriched in tip cell,

neophalanx cell, lymphatic endothelial cell subpopulations and particularly abundant in

human pancreatic cancer (Fig. 5k). Neophalanx endothelial cells, which express two pro-

angiogenesis markers, *Glul (Glutamine synthetase)* and *Cxcl12 (C-X-C motif chemokine*

ligand 12) [29,30] have recently been identified in lung tumour-associated endothelial cell

studies [31,32], and were proposed to line neo-vessels after termination of vessel sprouting.

Together this data suggests that based on the transcriptional state, T_CLECs might have

a supportive role in tumor vessel sprouting and remodeling. Detailed comparison with

recent single cell RNA seq data on tumour endothelial cells in mouse and human tumours

[31] confirmed the Tip-like gene signature of T_CLECs, but also identified significant

similarities with lymphatics (Fig. 5k). The latter is particularly surprising given that our

perfusion experiments with dextran clearly demonstrated that T_CLECs in the glioma

model line blood vessels, not lymphatics.

Acute recombination identifies a CLEC lineage in lymph nodes

The transcriptional similarities between T_CLECs and lymphatic endothelial cells are

noteworthy as blood endothelial cells, in particular in the brain environment, and

lymphatic EC represent highly differentiated cell types. During vertebrate embryogenesis,

lymphovenous specification sees common progenitor cells diverge in their differentiation

to establish mature blood endothelium and lymphatic endothelium[33]. The apparent

similarities could therefore suggest that T_CLECs represent an immature endothelial cell

state with partially overlapping profiles of both immature blood and lymphatic cells. In

this case, the expression of lymphatic genes would signify a return to an earlier cell state

or potentially some level of transdifferentiation between the two endothelial cell states.

The latter would imply a distant source of lymphatic endothelium as possible origin of

CLECs. Indeed, when searching for CLECs in all organs, we observed selective

endothelial labelling in lymph nodes, notably restricted to the periphery where specialized

lymphatic endothelial cells form the floor of the subcapsular sinus (SCS, Fig. 6a). Bone-

marrow transplantation to eliminate the abundant Csfr1-lineage macrophage population

highlighted the specific labelling of CLECs in lymph nodes (Fig. 6b).

To gain more insight into the emergence of T_CLECs, we asked whether acute

recombination could indicate their possible origin. Acute induction experiments (24h

after single tamoxifen injection) illustrated that the Csf1r-expressing progenitors of this

tumour endothelial cell population do not reside locally in pre-existing brain or the

associated brain tumour vasculature (Extended Fig. 8). Close inspection however

revealed GFP-positive endothelial cells with significant labelling 24h post-injection in

the floor endothelial cells of the SCS of lymph nodes (Fig. 6c). The mTmG Cre-reporter

switches from red fluorescent protein expression to mEGFP expression upon

recombination. Therefore, cells that acutely recombine, carry both GFP and Tomato

expression until the latter is fully degraded. Strikingly, floor SCS endothelial cells

exhibited both GFP and simultaneous Tomato expression, demonstrating recent

recombination. FACS analysis confirmed the presence of double positive EC in lymph

nodes, but not in brain or brain tumours. In fact, double positive cells for GFP and Tomato

were not observed in the brain tumours at any stage, excluding the possibility of any acute

or continuous local recombination of T_CLECs within the tumour microenvironment.

Prox-1 staining confirmed the lymphatic nature of CLECs in the lymphnode, and

demonstrated that only a subpopulation of lymphatic endothelial cells in the lymph node

carry the Csfr1-lineage trace, the cells in the floor of the SCS (Fig. 6d).

We decided to perform scRNA seq also on endothelial cells isolated from lymph nodes,

assess gene expression and perform pseudotime analysis to understand differentiation of

T_CLECs, T_EC, LN_CLECs (GFP positive) and LN_ECs (GFP negative) (Fig. 7a-f).

We used the same Smart-seq2 platform for comparison to sequence another 384 cells of

lymph node origin. After quality control (Fig. 7a), 89 cells EC, not Csfr1 lineage EC, and

104 cells expressing GFP (Fig.7b) in the lymph nodes were selected for further analysis.

Comparative analysis with recent mouse lymph node scRNA data[34] confirmed that the

LN_CLEC co-register with identified SCS floor cells (Extended Data Fig. 9a).

Unlike T_CLECs, LN_CLECs also expressed Csf1r, in line with our finding of acute

recombination in the SCS floor cells.

UMAP based dimension reduction placed T_CLECs and T_EC closely together, as before,

and also LN_CLEC and LN_EC co-cluster, although two additional distinct LN_EC

clusters appear highly distinct and thus unrelated to any of the other cell types (Fig. 7c).

Interestingly, some T_CLEC cluster with the LN_CLEC, suggesting that there is at least

some overlap in gene expression between the Csf1r lineage cells in the brain tumour and

in the lymph nodes. Looking at relative expression levels of all genes in a heatmap

comparing the lineage positive and negative EC from lymph nodes and tumours

highlighted that differential expression is more prevalent than any overlap between

tumour EC and LN EC (Fig. 7d). However, there are small number of genes like Pde4d,

Robo1 and Itga9 that show higher expression in the two lineage positive populations in

LN and tumour, compared to the lineage negative cells (Extended Data Fig. 9b). Itga9 is

well known to be important in lymphatic development[35], confirming again that T_CLECs

carry some gene expression patterns normally associated with lymphatic EC.

Nevertheless, despite the fact that only LN_CLEC appear to acutely recombine,

differential gene expression analysis does not provide clear evidence for a lineage

relationship between LN_CLEC and T_CLEC.

T_EC and T_CLECs "co-evolve" during tumour growth, suggesting distinct origin

To gain more insight into the differentiation state and possible route of differentiation,

we next performed pseudotemporal ordering using Monocle[36] (Fig. 7f). The Monocle

inferred trajectory suggested equally distributed pseudotime values for T_CLECs and

T_EC then LN_EC and finally LN_CLECs. When measuring single-cell gene expression

diversity as surrogate for differentiation potential with the CytoTRACE algorithm[37]

(higher score means more undifferentiated, stem-like), we observed again equal scores

between T_CLECs and T_EC but lower cytoTRACE values for LN_EC and LN_CLECs

(Fig. 7e-f). The equal scores between T_CLECs and T_ECs argue against a role of T_ECs

as precursors for T_CLECs, independently confirming that T_CLECs are unlikely

emerging through differentiation from the pre-existing brain tumour vasculature.

However, the LN_CLECs display the lowest cytoTRACE scores (most differentiated).

Based on the cytoTRACE analysis, and its logic of developmental potential, it would

seem unlikely that LN_CLEC are the origin of T_CLEC (Fig. 7e-f). Moreover, it appears

that both lineage positive and negative EC in the lymph nodes are more differentiated

than endothelial cells in the tumour. It is noteworthy that Monocle was designed for

developmental questions, which are strictly unidirectional. Interestingly, when plotting

CytoTRACE scores onto the Monocle trajectory (Fig. 7f), the right side shows higher

scores (T_CLEC and T_EC) and the left side (darker blue) lower scores (LN_CLECs

lowest), being in line with pseudotime. This again predicts that LN_CLECs represent an

end state rather than the origin. In summary, these data predict that during tumour growth,

the T_EC and T_CLECs coevolve from a distinct origin to then share many aspects of

their gene expression and biology, yet with a few very distinct gene profiles.

DISCUSSION

In this study, using inducible *Csf1r* lineage tracing in adult mice intended to study

macrophage recruitment during glioma development, we identify a distinct endothelial

cell lineage that forms up to 10% of the tumour vasculature in adult mice. Selective

experimental depletion of this *Csf1r* endothelial cell lineage demonstrated a role in

promoting vascular branching and tumour growth. Through a combination of genetic

lineage tracing, longitudinal live two-photon imaging, and single cell analysis, we find

that these cells arise during tumour progression, adopt dynamic endothelial phenotypes

including tip cell characteristics, and express common endothelial as well as unique

markers that can also be found in patient-derived tumour endothelium.

Csf1r is a well-known regulator of monocyte/ macrophage differentiation [38,39], but is

also important for yolk sac hematopoiesis and progenitor cell differentiation. Klotz and

coworkers reported that *Csf1r*-expressing cells in the mouse embryo contribute to *Prox1-*

positive lymphatic endothelial cells in the cardiac lymphatic vasculature [40]. Moreover, a

recent report identified that *Csf1r*-expressing erythro-myeloid progenitors from the early

embryonic yolk-sac could give rise to intraembryonic endothelial cells in mouse

development[41]. *Csf1r* is also expressed in EPCs, such as colony forming unit-Hill cells

and circulating angiogenic cells [42]. In the adult however, *Csf1r* is not expressed in the

mature endothelium of blood and lymphatic vessels, but continues to drive recruitment

and expansion of myeloid cells in inflammation, ischemia and tumour growth [43].

Intriguingly, *Csf1r* itself is no longer expressed in the Csf1r lineage cells within the

tumour and acute recombination of endothelial cells 24h after tamoxifen injection fails to

label cells in the tumour vasculature. These results raise the possibility that a *Csf1r-*

expressing cell population of endothelial nature or potential is present elsewhere in the

body, recombines upon tamoxifen injection and is recruited to the tumour vasculature

where it differentiates into blood endothelium to contribute to tumour angiogenesis.

Alternatively, the fact that Csf1r expression is not detected in isolated tumour CLECs

could also indicate that the Cre-line is leaky and gives rise to labelled cells through

stochastic recombination. In that case, one would expect the labelled cells to populate any

part of the vasculature and be otherwise indistinguishable from the unlabelled tumour EC.

Single cell analysis however identifies a unique gene expression signature beyond the

large number of genes that are co-expressed in T_CLEC and T_EC, and demonstrates

that T_CLEC are distinctly enriched in tip-cell and neo-phalanx cell populations. Thus,

labelled cells show distinct phenotypes, location and expression patterns, ruling out

purely stochastic recombination events. Furthermore, such stochastic recombination

within the tumour vasculature would necessitate that we identify acutely labelled cells

marked by co-expression of Tomato and GFP. As we failed to identify such cells at any

timepoint during tumour growth, and also could detect the emergence of T_CLECs in

mice only injected with tamoxifen before tumour implantation, the cumulative evidence

would suggest that T_CLECs originate outside of the tumour and also that the progenitor

population resides outside of the brain. Despite all our efforts, the present work fails to

provide ultimate and definitive proof of the true origin of T_CLECs, but arrives at a clear

direction and working hypothesis for future studies. In search of their origin and a

potential progenitor population, we imaged and analysed all organs by FACS. We

performed bone-marrow transplantation, performed splenectomy and examined the entire

mouse by fluorescence stereomicroscopy after clearing to identify regions of

recombination. The results of bone-marrow transplantation effectively ruled out EPC or

any other known progenitors that may be found in circulation, including

transdifferentation from macrophages. Four independent observations placed a spotlight

on lymphatics, and the lymph nodes in particular. First, the only *bonafide* Csf1r lineage

endothelial cell population that we could identify outside of the tumours, including after

complete reconstitution of the bone marrow by wildtype bone marrow, was found in the

subcapsular sinus of the lymph nodes. Here the floor cells, specialized lymphatic

endothelial cells that we identify as GFP and Csf1r positive line the capsule in lymph

nodes in close apposition to the Csf1 expressing ceiling cells. Second, after acute

tamoxifen injection the cells in the floor of the subcapsular sinus are uniquely double

positive for Tomato and GFP, demonstrating acute and likely local recombination. Third,

single cell RNAseq of T-CLECs identified lymphatic markers including the lymphatic

master regulator Prox1 co-expressed with blood endothelial markers, and SCDE placed

T_CLECs close to lymphatic endothelial cells from various mouse tumour models in

recently published datasets. And finally, CLECs can also be found in lymphatics in

subcutaneous tumour models and in skin. These independent observations clearly make

the SCS floor cells prime candidates for the origin of T_CLECs, but future work will be

necessary for a possible direct demonstration of this link.

In developmental studies, lineage relationships have recently been studied by scRNAseq

data sets through pseudotime analysis. In particular the CytoTRACE analysis can be

useful to estimate the degree of differentiation of cell populations and thereby also predict

their stemness [37]. Our comparative analysis of endothelial cells isolated from tumour and

lymph nodes assigned the highest differentiation scores to cells in the lymph node, and

lower values to the EC in tumour, including T_CLECs. Accordingly, the cells in the floor

of the SCS that carry the lineage label would rather be assumed at the end of a

differentiation process rather than progenitors for other populations such as the T_CLECs.

Whether however such algorithms are suitable to determine the potential of endothelial

cells to become activated and possibly transdifferentiate is unclear. In particular given the

altered environmental context of EC within a growing tumour compared to a vascular bed

in homeostasis. The combined monocle[36] and cytoTRACE analysis would fit with a

general level of dedifferentiation of endothelial cells within the tumour microenvironment,

and therefore a co-evolution of the cell states of both T_EC and T_CLEC. It is likely that

pseudotemporal ordering of endothelial cells in a reactivation state will always show the

reactivated cells, like in the tumour environment as less differentiated. Indeed, recent

work studying the effect of chronic hypoxia on brain endothelium illustrated the wide

range of reactivation, in particular noting a strong increase in tip cell population amongst

the brain endothelial cells[44]. The current assumption states that these new tip cells arise

under hypoxia from pre-existing brain endothelial cells, and possibly emerge from

particular endothelial subpopulations. In the absence of lineage tracing, an alternative

possibility, raised by our current observation in tumour angiogenesis, could be that new

tip cells arise at least in part elsewhere, and are recruited to the hypoxic vessel areas.

More work on vascular single cell analysis will be required to establish the fundamental

principles of endothelial activation and their interpretation for the different tissue

challenges that require vascular adaptations.

Angiogenesis is essential for tumour growth, and contributes to metastasis [45,46]. Early

hopes that targeting tumour angiogenesis may prove uniquely selective and effective as

anti-tumour treatment have, however, been disappointing. Intense research efforts

therefore currently focus on understanding reasons and mechanisms for resistance to anti-

angiogenic therapy, and identifying new cellular and molecular targets to effectively

modify the tumour vasculature [47]. Intravital imaging shows that T_CLECs arise during

tumour progression, adopt endothelial phenotypes with tip cell characteristics and line

perfused vascular tubes. Consistent with this observation, AUCell analysis of scRNA-seq

data in human samples identifies that T_CLECs signatures are highly enriched in tip cell

subpopulation. A recent study using single-cell transcriptome analysis revealed that anti-

VEGF (vascular endothelial growth factor) treatment reduces the proportion of

endothelial tip-like cells in a tumour xenografts [48]. How CLECs affect the outcome of

anti-angiogenic treatment will be important to investigate in the future. Considering that

T_CLECs prominently express *Vegfr3*, even more so than regular endothelial cells in the

tumour, we can speculate that T_CLECs are resistant to anti-VEGFR2 treatment, and

possibly escape treatment through the activity of VEGFR3. If so CLECs may adversely

affect anti-angiogenic therapy targeting VEGFR2 selectively. Tumour blood vessels are

generally highly tortuous and the branching pattern is greatly different from normal blood

vessels. Our results from the selective targeting of CLECs in the mouse glioma model

show that CLECs ablation decreases both vessel branching and tumour growth. Thus, in

the absence of CLEC, the dysmorphic vascular phenotype in tumours appears exacerbated,

suggesting that CLECs support the formation of a highly branched vascular network,

which in turn better promotes tumour growth. Future work will be required to identify the

true origin of T_CLECs and to extend this functional analysis to fully appreciate the

consequence and opportunities to target or utilize CLECs for therapeutic approaches.

METHODS

Mice

Mice expressing both the *Csf1r-Mer-iCre-Mer* transgene (*Csf1r-Mer2.Cre*) [17] and the

Rosa^mTmG Cre recombination reporter [18] were generated by breeding. Mice carrying the.

Endothelial specific *Cdh5-Dre* mice [24] were mated with *Csf1r-Mer2.Cre* mice and

Rosa26-RSR-LSL-tdTomato mice [23] to obtain *Cdh5-Dre*:: *Csf1r-Mer2.Cre*:: *Rosa26-RSR-*

LSL-tdTomato mice (*DreMer2.Cre^tdTomato*) that allows combinatorial genetics as a tool for

CLECs tracing.   Endothelial specific *Cdh5-Dre* mice were mated with *Csf1r-Mer2.Cre*

mice and *Rosa26-RSR-mTmG-DTR* mice to obtain *Cdh5-Dre*:: *Csf1r-Mer2.Cre*:: *Rosa26-*

RSR-mTmG-DTR mice (*DreMer2.Cre^mTmG-DTR*) that allows for selective tracing and

ablation of CLECs. *Rosa26-RSR-mTmG-DTR* mice were designed by Dr. Fabio Stanchi

and generated by the team of Dr. Ralf Kühn at the Max Delbrück Center in Germany. All

mouse strains were maintained on C57BL/6J background. Animal housing and all

experimental procedures were approved by the Institutional Animal Care and Research

Advisory Committee of the KU Leuven (085/2016).

Cell lines

CT2A, GL261 glioma cell lines and B16F1 melanoma cell line were cultured in DMEM

(Life Technologies) supplemented with 10% FBS (Life Technologies), 1% penicillin/

streptomycin (Life Technologies) and 1% glutamine (Life Technologies). Spheroids were

obtained by seeding the glioma cells for 48h on non-adherent culture dishes. Spheroids

of 200- 250μm were selected for implantation.

Intravital imaging on mouse gliomas

Surgery for tumor implantation, installation of cranial windows and *in vivo* imaging were

performed as described previously [16]. Briefly, 8– 12-weeks-old mice were anesthetized

with ketamine/ xylazine and a craniotomy was performed on the parietal bone. Spheroids

of CT2A or GL261 cells modified to express the blue fluorescent protein TagBFP [50] were

injected in the exposed parietal brain cortex, which was then sealed by cementing a glass

coverslip to the bone surrounding the craniotomy. After surgery, mice were injected

intraperitoneally with tamoxifen (65 μg/g body weight) every 5 days. For CLECs

depletion, mice were injected intraperitoneally with diphtheria toxin (4 ng/g body weight)

the next day after tamoxifen induction. Longitudinal *in vivo* imaging of growing tumors

was performed in mice under isoflurane anesthesia using a SP8 upright microscope (Leica

Microsystems) equipped with a HCX IRAPO L25x/0.95 water objective and a Titanium:

Sapphire laser (Vision II, Coherent Inc.) tuned at 925 nm. Humane end-point of

experiments was applied if animals lost 15-20% of their original weight or shown evident

signs of distress.

Immunofluorescence imaging

Mice anesthetized with ketamine/ xylazine were perfused through the heart with 15 ml of

ice-cold PBS, followed by 10 ml of 2% PFA in PBS. Tumour tissues or organs were

harvested and fixed overnight in 4% PFA. For tumour tissues, sections (200 μm-thick)

were prepared with a vibratome 650V (Thermo Scientific). For organ samples, frozen

sections (10 μm-thick) were prepared with a cryostat NX70 (Thermo Scientific). For

immunofluorescence imaging studies, the PFA-fixed sections were blocked and

permeabilized in TNBT (0.1 M Tris, pH 7.4, 150 mM NaCl, 0.5% blocking reagent from

Perkin Elmer, 0.5% Triton X-100) for 4h at room temperature. Sections were incubated

with antibodies against CDH5 (1:25, AF1002, R&D), CD31 (1:100, ab28364, Abcam),

F4/80 (1:100, MF48000, Invitrogen), CD45 (1:100, 550539, BD Pharmingen),   CD169

(1:100, 142401, Biolegend), PROX1 (1:200, 11-002, Angiobio), AQP1 (1:4000, AB2219,

Millipore), LYVE1 (1:100, 50-0433-80, eBioscience) diluted in TNBT buffer overnight

at 4°C, washed in TNT buffer (0.1 M Tris pH 7.4; 150 mM NaCl, 0.5% Triton X-100)

and incubated with an Alexa Fluor conjugated antibodies (1:200, ThermoFisher

Scientific). Sections were washed and mounted in fluorescent mounting medium (Dako).

Images were obtained with a Leica TCS SP8 confocal microscope.

Flow cytometric analysis

Mice were anesthetized with ketamine/ xylazine, then tumour tissues or organs were

harvested and incubated with PBS containing 1mg/ml collagenase I (Gibco), 2 mg/ ml

Dispase I (Gibco), 100ug/ml DNase I (Roche) and 2 mM $CaCl_2$ for 30 min at 37 °C. After

incubation, the digested tissue was passed through a cell strainer and then washed by PBS

including 2% FBS. For red cell exclusion, we incubated samples with red cell lysis buffer

(Sigma) for 5 min at 37 °C and then washed by PBS including 2% FBS and 2 mM EDTA.

Cells were stained with the following monoclonal antibodies: PE/ Cy7 anti-CD45

(552848, BD), APC anti-CD31 (561814, BD). Data acquisition was performed with BD

FACSVerse and analysis was performed with BD FACSuite software, and FlowJo

software.

Mouse melanoma model

Mice (8–12-weeks-old) were anesthetized with ketamine/ xylazine and then 1×10^6 B16F1

melanoma cells in PBS were implanted into the right flank region of the mice. B16F1

melanomas were removed from mice 12 days after implantation for immunofluorescent

imaging studies and flow cytometric analysis. Mice were injected intraperitoneally with

tamoxifen (65 µg/g body weight) 4 time/ 2 weeks before tumour implantation and then

intraperitoneally with tamoxifen (65 μg/g body weight) every 5 days after tumour

implantation.

Single cell isolation and RNA-sequencing

Single cells were sorted (BD FACSAriaIII) in 96 well plates (VWR) containing 2 μL of

PBS including 0.2% Triton X-100 and 4U of RNase inhibitor (Takara) per well. Plates

were properly sealed and spun down at 2000 g for 1 min before storing at −80°C. Whole

transcriptome amplification was performed with a modified SMART-seq2 protocol as

described previously [25], using 20 instead of 18 cycles of cDNA amplification. PCR

purification was realized with a 0.8:1 ratio (ampureXP beads:DNA). Amplified cDNA

quality was monitored with a high sensitivity DNA chip (Agilent) using the Bioanalyzer

(Agilent). Sequencing libraries were performed using the Nextera XT kit (Illumina) as

described previously [25], using 1/4th of the recommended reagent volumes and 1/5th of

input DNA with a tagmentation time of 9 min. Library quality was monitored with a high

sensitivity DNA chip (Agilent) using the Bioanalyzer (Agilent). Indexing was performed

with the Nextera XT index Kit V2 (A-D). Up to 4 x 96 single cells were pooled per

sequencing lane. Samples were sequenced on the Illumina NextSeq 500 platform using

75bp single-end reads.

scRNA-sequencing data analysis

BAM files were converted to merged, demultiplexed FASTQ files, cleaned using fastq-

mcf (ea-utils r819), and QC checked with FastQC (0.11.4). Reads were then mapped to

the mouse genome (mm18) using STAR (2.4.1b) and quantified with Subread (1.4.6-p2).

Cells with less than 100,000 reads and/or 500 genes expressed, more than 20%

mitochondrial reads, and less than an average expression level of 3.0 of about 80

housekeeping genes [51] were discarded. 469 cells passed these stringent quality criteria.

Winsorized Highly Variable genes (HVGs) were identified for tumor derived

macrophages (T_MACs), endothelial cell (T_ECs) and *Csf1r* lineage endothelial cells (T_CLECs) [26]. These cells were then clustered based on HVG expression using non-negative matrix factorization as dimension reduction approach (run=40, rank=10, in MeV 4.8.1). The "best fit" (numbers of clusters) was chosen based on the highest cophenetic correlation coefficient. Next, Single-cell Differential Expression analysis (SCDE) was performed between the different NMF-clusters using the global gene expression matrix [26]. Expression values were library-size-factor-normalized (DESeq) and log2 transformed (log2+1). T_MAC, T_EC, T_CLEC and LN_CLECS, LN_EC, T_CLEC, T_EC cells were projected into a two-dimensional space using Uniform Manifold Approximation and Projection (dims=15, resolution = 0.4) based on 5k most variable features (Seurat pipeline) [52]. Monocle (2.14) based trajectory analysis was performed on T_CLEC, T_EC, LN_CELC and LN_EC (total=454 cells) taking into account n=790 m3Drop genes. The same data set was subjected to CytoTRACE analysis using the online tool (https://cytotrace.stanford.edu/) [37].

AUCell analysis of human scRNA-seq data in multiple tumours

The droplet-based scRNA-seq data of 4 human tumour types were downloaded from publicly available sources including lung cancer (ArrayExpress: E-MTAB-6149 and E-MTAB-6653)[53], PDAC (GSA: CRA001160)[54], liver cancer (GEO: GSE125449)[55], and BCC (GEO: GSE123814)[56]. These datasets were processed and clustered per tumour type using Seurat (v3.0.4) package, and the endothelial cell clusters were identified using marker genes (*CLDN5*, *VWF*, *PECAM1*). The resulting endothelial cell data were pooled together and aligned using anchor-based canonical correlation analysis (CCA) to mitigate the difference related to distinct tumor types and scRNA-seq technologies. Initial graph-based clustering identified 10 clusters including 3 potential doublet clusters as predicted

by DoubleFinder (v2), which were subsequently removed before second round of

clustering. The resulting 8 clusters consisting of 7314 endothelial cells were annotated

based on known marker genes, and their CLECs signature were calculated using AUCell

package (v1.6.1). The threshold of AUC score was chosen at the peak of the right modal,

and the percentage of putative CLECs were calculated for different subtypes and their

tumour type of origin.

Generation of bone marrow chimeras

Recipient 8-10-weeks-old mice were lethally irradiated (9.5 Gy) and then intravenously

injected with 1×10^7 bone marrow cells from donor mice 16 h later. Tumour implantation

experiments were initiated 8 weeks after bone marrow reconstitution. Reconstitution rate

was determined using flow cytometric analysis in the peripheral blood.

Surgical removal of the spleen

Mice (8–12-weeks-old) were anesthetized with ketamine/ xylazine and then the spleen

was gently taken out from the connective tissue, while cauterizing associated vessels. The

incision was thereafter sutured. These procedures were performed as described previously

[57].

Vessel morphology analysis

Confocal images of the tumour vasculature were analyzed for two-dimensional vessel

area, network length and number of network bifurcation points. The tdTomato and GFP

channels were merged and a Gaussian blur filter as well as Otsu thresholding was applied

to obtain binary vessel mask for sectional area estimation. To obtain the vessel network,

a semi-automatic multicut workflow from ilastik [58] was applied to all images. The

segmented vessels were skeletonized and converted into a network graph with nodes and

edges by using the Python package ImagePy [59]. The lengths of network edges were

summed up to obtain the total network length for each image, all nodes with a degree of

three or more were counted as bifurcations.

Statistics

Statistical analyses of data were performed with GraphPad Prism 7.0 software. All data

are shown as the means ± standard deviations. Differences were assessed using a two-

tailed unpaired Mann-Whitney's U test or two-tailed unpaired Welch's correction.

AUTHOR CONTRIBUTIONS

KM led and conceived the project, designed and performed experiments, analysed and

interpreted data. FR, FS, TM, WG, LH, JQ performed experiments and analyzed data.

DL, BZ, CB, JM and HG analyzed data. HG conceived and supervised the project. KM

and HG wrote and edited the manuscript, with contribution by all authors.

ACKNOWLEDGEMENT

This work was supported by the Belgian Cancer Foundation (Stichting Tegen Kanker,

grant 2012-181, 2018-074) and a Hercules type 2 grant (Herculesstichting: AKUL11033).

K.M. is a recipient of a Japan Society for the Promotion of Science Overseas Research

Fellowship. We would like to thank Dr. Till Acker (Institute of Neuropathology,

University of Giessen, Germany) and Dr. Thomas N. Seyfried (Biology department,

Boston College, USA) for the gift of GL261 and CT2A cells, respectively. We are grateful

for outstanding advice and service by Dr. Ralf Kühn heading the Transgenics Core

Facility at the Max-Delbrueck-Center, Berlin, Germany. We would like to thank our

colleagues Marly Balcer, Lisse Decraecker, Greet Bervoets and Irene Hollfinger for

excellent technical assistance.

CODE AVAILABILITY

The computational code (R) used for scRNA-seq analysis is based on already published

pipelines (see Material and Methods) and available upon request.

DATA AVAILABILITY

All scRNA-seq data are available at Gene Expression Omnibus under accession number

GSE157507.

STATISTICS

Differences were assessed using either a two-tailed unpaired Mann-Whitney's or two-

tailed unpaired t-test with Welch correction. All planned comparisons were considered

stand-alone and were not corrected for multiple testing.

REFERENCES

1　Asahara, T. *et al.* Isolation of putative progenitor endothelial cells for

angiogenesis. *Science* **275**, 964-967, doi:10.1126/science.275.5302.964 (1997).

2　Shi, Q. *et al.* Evidence for circulating bone marrow-derived endothelial cells.

Blood **92**, 362-367 (1998).

3　Asahara, T. *et al.* Bone marrow origin of endothelial progenitor cells responsible

for postnatal vasculogenesis in physiological and pathological neovascularization.

Circ Res **85**, 221-228, doi:10.1161/01.res.85.3.221 (1999).

4　Peters, B. A. *et al.* Contribution of bone marrow-derived endothelial cells to

human tumor vasculature. *Nat Med* **11**, 261-262, doi:10.1038/nm1200 (2005).

5　Rafii, S., Lyden, D., Benezra, R., Hattori, K. & Heissig, B. Vascular and

haematopoietic stem cells: novel targets for anti-angiogenesis therapy? *Nat Rev*

Cancer **2**, 826-835, doi:10.1038/nrc925 (2002).

6   Rafii, S. & Lyden, D. Therapeutic stem and progenitor cell transplantation for

organ vascularization and regeneration. *Nat Med* **9**, 702-712,

doi:10.1038/nm0603-702 (2003).

7   Religa, P. *et al.* Presence of bone marrow-derived circulating progenitor

endothelial cells in the newly formed lymphatic vessels. *Blood* **106**, 4184-4190,

doi:10.1182/blood-2005-01-0226 (2005).

8   Kerjaschki, D. *et al.* Lymphatic endothelial progenitor cells contribute to de novo

lymphangiogenesis in human renal transplants. *Nat Med* **12**, 230-234,

doi:10.1038/nm1340 (2006).

9   Purhonen, S. *et al.* Bone marrow-derived circulating endothelial precursors do not

contribute to vascular endothelium and are not needed for tumor growth. *Proc*

Natl Acad Sci U S A **105**, 6620-6625, doi:10.1073/pnas.0710516105 (2008).

10   Kerbel, R. S. *et al.* Endothelial progenitor cells are cellular hubs essential for

neoangiogenesis of certain aggressive adenocarcinomas and metastatic transition

but not adenomas. *Proc Natl Acad Sci U S A* **105**, E54; author reply E55,

doi:10.1073/pnas.0804876105 (2008).

11   Basile, D. P. & Yoder, M. C. Circulating and tissue resident endothelial progenitor

cells. *J Cell Physiol* **229**, 10-16, doi:10.1002/jcp.24423 (2014).

12   Psaltis, P. J. & Simari, R. D. Vascular wall progenitor cells in health and disease.

Circ Res **116**, 1392-1412, doi:10.1161/CIRCRESAHA.116.305368 (2015).

13   Fujisawa, T. *et al.* Endothelial Progenitor Cells Do Not Originate From the Bone

Marrow. *Circulation* **140**, 1524-1526,

doi:10.1161/CIRCULATIONAHA.119.042351 (2019).

14   Martinez-Murillo, R. & Martinez, A. Standardization of an orthotopic mouse

brain tumor model following transplantation of CT-2A astrocytoma cells. *Histol*

Histopathol **22**, 1309-1326, doi:10.14670/HH-22.1309 (2007).

15   Zagzag, D. *et al.* Expression of hypoxia-inducible factor 1alpha in brain tumors:

association with angiogenesis, invasion, and progression. *Cancer* **88**, 2606-2618

(2000).

16   Stanchi, F., Matsumoto, K. & Gerhardt, H. Imaging Glioma Progression by

Intravital Microscopy. *Methods Mol Biol* **1862**, 227-243, doi:10.1007/978-1-

4939-8769-6_16 (2019).

17   Qian, B. Z. *et al.* CCL2 recruits inflammatory monocytes to facilitate breast-

tumour metastasis. *Nature* **475**, 222-225, doi:10.1038/nature10138 (2011).

18   Muzumdar, M. D., Tasic, B., Miyamichi, K., Li, L. & Luo, L. A global double-

fluorescent Cre reporter mouse. *Genesis* **45**, 593-605, doi:10.1002/dvg.20335

(2007).

19   Mathivet, T. *et al.* Dynamic stroma reorganization drives blood vessel dysmorphia

during glioma growth. *EMBO Mol Med* **9**, 1629-1645,

doi:10.15252/emmm.201607445 (2017).

20   Raivich, G. *et al.* Regulation of MCSF receptors on microglia in the normal and

injured mouse central nervous system: a quantitative immunofluorescence study

using confocal laser microscopy. *J Comp Neurol* **395**, 342-358,

doi:10.1002/(sici)1096-9861(19980808)395:3<342::aid-cne6>3.0.co;2-2 (1998).

21   Jin, S. *et al.* Interleukin-34 restores blood-brain barrier integrity by upregulating

tight junction proteins in endothelial cells. *PLoS One* **9**, e115981,

doi:10.1371/journal.pone.0115981 (2014).

22   Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain

vasculature. *Nature* **554**, 475-480, doi:10.1038/nature25739 (2018).

23   Liu, Q. *et al.* Lung regeneration by multipotent stem cells residing at the

bronchioalveolar-duct junction. *Nat Genet* **51**, 728-738, doi:10.1038/s41588-019-

0346-6 (2019).

24   Tang, M. *et al.* Simultaneous quantitative assessment of two distinct cell lineages

with a nuclear-localized dual genetic reporter. *J Mol Cell Cardiol* **146**, 60-68,

doi:10.1016/j.yjmcc.2020.07.002 (2020).

25   Picelli, S. *et al.* Full-length RNA-seq from single cells using Smart-seq2. *Nat*

Protoc **9**, 171-181, doi:10.1038/nprot.2014.006 (2014).

26   Kharchenko, P. V., Silberstein, L. & Scadden, D. T. Bayesian approach to single-

cell differential expression analysis. *Nat Methods* **11**, 740-742,

doi:10.1038/nmeth.2967 (2014).

27   Saadoun, S., Papadopoulos, M. C., Hara-Chikuma, M. & Verkman, A. S.

Impairment of angiogenesis and cell migration by targeted aquaporin-1 gene

disruption. *Nature* **434**, 786-792, doi:10.1038/nature03460 (2005).

28   Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering.

Nat Methods **14**, 1083-1086, doi:10.1038/nmeth.4463 (2017).

29   Eelen, G. *et al.* Role of glutamine synthetase in angiogenesis beyond glutamine

synthesis. *Nature* **561**, 63-69, doi:10.1038/s41586-018-0466-7 (2018).

30   Zhang, M. *et al.* CXCL12 enhances angiogenesis through CXCR7 activation in

human umbilical vein endothelial cells. *Sci Rep* **7**, 8289, doi:10.1038/s41598-017-

08840-y (2017).

31   Goveia, J. *et al.* An Integrated Gene Expression Landscape Profiling Approach to

Identify Lung Tumor Endothelial Cell Heterogeneity and Angiogenic Candidates.

Cancer cell **37**, 21-36 e13, doi:10.1016/j.ccell.2019.12.001 (2020).

32   Rohlenova, K. *et al.* Single-Cell RNA Sequencing Maps Endothelial Metabolic

Plasticity in Pathological Angiogenesis. *Cell Metab* **31**, 862-877 e814,

699   doi:10.1016/j.cmet.2020.03.009 (2020).

700   33   Yang, Y. & Oliver, G. Development of the mammalian lymphatic vasculature.

701   *The Journal of clinical investigation* **124**, 888-897, doi:10.1172/JCI71609 (2014).

702   34   Fujimoto, N. *et al.* Single-cell mapping reveals new markers and functions of

703   lymphatic endothelial cells in lymph nodes. *PLoS biology* **18**, e3000704,

704   doi:10.1371/journal.pbio.3000704 (2020).

705   35   Bazigou, E. *et al.* Integrin-alpha9 is required for fibronectin matrix assembly

706   during lymphatic valve morphogenesis. *Developmental cell* **17**, 175-186,

707   doi:10.1016/j.devcel.2009.06.017 (2009).

708   36   Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed

709   by pseudotemporal ordering of single cells. *Nature biotechnology* **32**, 381-386,

710   doi:10.1038/nbt.2859 (2014).

711   37   Gulati, G. S. *et al.* Single-cell transcriptional diversity is a hallmark of

712   developmental potential. *Science* **367**, 405-411, doi:10.1126/science.aax0249

713   (2020).

714   38   Tushinski, R. J. *et al.* Survival of mononuclear phagocytes depends on a lineage-

715   specific growth factor that the differentiated cells selectively destroy. *Cell* **28**, 71-

716   81, doi:10.1016/0092-8674(82)90376-2 (1982).

717   39   Dai, X. M. *et al.* Targeted disruption of the mouse colony-stimulating factor 1

718   receptor gene results in osteopetrosis, mononuclear phagocyte deficiency,

719   increased primitive progenitor cell frequencies, and reproductive defects. *Blood*

720   **99**, 111-120, doi:10.1182/blood.v99.1.111 (2002).

721   40   Klotz, L. *et al.* Cardiac lymphatics are heterogeneous in origin and respond to

722   injury. *Nature* **522**, 62-67, doi:10.1038/nature14483 (2015).

41   Plein, A., Fantin, A., Denti, L., Pollard, J. W. & Ruhrberg, C. Erythro-myeloid

progenitors contribute endothelial cells to blood vessels. *Nature* **562**, 223-228,

doi:10.1038/s41586-018-0552-x (2018).

42   Hirschi, K. K., Ingram, D. A. & Yoder, M. C. Assessing identity, phenotype, and

fate of endothelial progenitor cells. *Arterioscler Thromb Vasc Biol* **28**, 1584-1595,

doi:10.1161/ATVBAHA.107.155960 (2008).

43   Mantovani, A., Marchesi, F., Malesci, A., Laghi, L. & Allavena, P. Tumour-

associated macrophages as treatment targets in oncology. *Nat Rev Clin Oncol* **14**,

399-416, doi:10.1038/nrclinonc.2016.217 (2017).

44   Heng, J. S. *et al.* Hypoxia tolerance in the Norrin-deficient retina and the

chronically hypoxic brain studied at single-cell resolution. *Proceedings of the*

National Academy of Sciences of the United States of America **116**, 9103-9114,

doi:10.1073/pnas.1821122116 (2019).

45   Folkman, J. Tumor angiogenesis: therapeutic implications. *N Engl J Med* **285**,

1182-1186 (1971).

46   Ferrara, N. & Kerbel, R. S. Angiogenesis as a therapeutic target. *Nature* **438**, 967-

974, doi:10.1038/nature04483 (2005).

47   Welti, J., Loges, S., Dimmeler, S. & Carmeliet, P. Recent molecular discoveries

in angiogenesis and antiangiogenic therapies in cancer. *J Clin Invest* **123**, 3190-

3200, doi:10.1172/JCI70212 (2013).

48   Zhao, Q. *et al.* Single-Cell Transcriptome Analyses Reveal Endothelial Cell

Heterogeneity in Tumors and Changes following Antiangiogenic Treatment.

Cancer Res **78**, 2370-2382, doi:10.1158/0008-5472.CAN-17-2728 (2018).

49   Bazigou, E. *et al.* Genes regulating lymphangiogenesis control venous valve

formation and maintenance in mice. *The Journal of clinical investigation* **121**,

2984-2992, doi:10.1172/JCI58050 (2011).

50   Subach, O. M. *et al.* Conversion of red fluorescent protein into a bright blue probe.

Chem Biol **15**, 1116-1124, doi:10.1016/j.chembiol.2008.08.006 (2008).

51   Tirosh, I. *et al.* Dissecting the multicellular ecosystem of metastatic melanoma by

single-cell RNA-seq. *Science* **352**, 189-196, doi:10.1126/science.aad0501 (2016).

52   Stuart, T. *et al.* Comprehensive Integration of Single-Cell Data. *Cell* **177**, 1888-

1902 e1821, doi:10.1016/j.cell.2019.05.031 (2019).

53   Lambrechts, D. *et al.* Phenotype molding of stromal cells in the lung tumor

microenvironment. *Nat Med* **24**, 1277-1289, doi:10.1038/s41591-018-0096-5

(2018).

54   Peng, J. *et al.* Single-cell RNA-seq highlights intra-tumoral heterogeneity and

malignant progression in pancreatic ductal adenocarcinoma. *Cell Res* **29**, 725-738,

doi:10.1038/s41422-019-0195-y (2019).

55   Ma, L. *et al.* Tumor Cell Biodiversity Drives Microenvironmental

Reprogramming in Liver Cancer. *Cancer Cell* **36**, 418-430 e416,

doi:10.1016/j.ccell.2019.08.007 (2019).

56   Yost, K. E. *et al.* Clonal replacement of tumor-specific T cells following PD-1

blockade. *Nat Med* **25**, 1251-1259, doi:10.1038/s41591-019-0522-3 (2019).

57   Cortez-Retamozo, V. *et al.* Origins of tumor-associated macrophages and

neutrophils. *Proc Natl Acad Sci U S A* **109**, 2491-2496,

doi:10.1073/pnas.1113744109 (2012).

58   Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nat*

Methods **16**, 1226-1232, doi:10.1038/s41592-019-0582-9 (2019).

59   Wang, A., Yan, X. & Wei, Z. ImagePy: an open-source, Python-based and

platform-independent software package for bioimage analysis. *Bioinformatics* **34**,

3238-3240, doi:10.1093/bioinformatics/bty313 (2018).

FIGURE LEGENDS

Fig. 1: Csf1r lineage endothelial cells (CLECs) contribute to perfused, dynamically

sprouting and remodeling blood vessels in mouse glioma. a, Csf1r-Mer2.CremTmG:

a mouse model for Csf1r-lineage tracing combining the tamoxifen-inducible Cre-driver

under control of the Csf1r-promoter and the mTmG Cre-reporter. b, Timing schedule of

glioma implantation, tamoxifen induction and in vivo imaging. c-f, Longitudinal

intravital imaging of the tumour vasculature in CT2A mouse glioma of Csf1r-

Mer2.CremTmG by cranial window. Scale bar: 50 μm. c, In vivo imaging in 3 weeks

glioma. Endothelial cells in red (tdTomato), Csf1r-labeled cells in green (GFP) and

glioma cells in blue (BFP). d, Detail image of outlined square in Fig. 1c illustrating tracing

of GFP-positive cells. White arrowheads, macrophages; blue arrowhead, CLECs at tip

position. e, In vivo imaging of CLECs in 2 weeks glioma. Blue arrowheads, tip of

endothelial cells. f, Visualization of blood vessels in 3 weeks glioma by Intravenous

injection of dextran with texas red dye. Yellow arrowhead, bridging CLECs; orange

arrowhead, tube formation by CLECs; blue arrowhead, CLECs at tip position. g-h,

Counterstaining on PFA-fixed 4 weeks CT2A glioma sections of Csf1r-Mer2.CremTmG

with indicated antibodies. Yellow arrows, CLECs. Scale bar: 25 μm.

Fig. 2: Up to 10% of intra-tumour endothelium is comprised of CLECs not only in

glioma, but also in other tumour types and locations. a-i, Quantification of the

population of CLECs in glioma of Csf1r-Mer2.CremTmG mice. a, Gross appearance of

sham control and 4 weeks CT2A glioma (Upper panel). Purple arrowhead, the area of

capillary injection; blue arrowhead, glioma. Scale bar: 5 mm. Flow cytometric analysis

of CLECs in sham control and 4 weeks CT2A glioma (Lower panel). b, In 3 weeks CT2A

glioma (n= 9), CLECs constitute $2.1 \pm 1.0\%$ of total endothelial cells (CD45-CD31+); in

4 weeks CT2A glioma (n= 5), they make up $7.5 \pm 1.6\%$ of total endothelial cells. c, In 3

weeks GL261 glioma (n= 5), CLECs constitute $1.5 \pm 1.6\%$ of total endothelial cells. d,

Gross appearance of normal brain, sham control and 3 weeks CT2A glioma. Purple

arrowhead, area of capillary injection. Scale bar: 5 mm. e-f, Quantification of the

population of CLECs by flow cytometric analysis in brain and tumour samples. Bars

represent mean $\pm$ s.d. $*p<0.05$, $**p<0.01$, $***p<0.001$, $****P<0.0001$. Two-tailed

unpaired Mann-Whitney' s U test. i-j, Analysis of CLECs in B16F1 melanoma in Csf1r-

Mer2.CremTmG mice. g, Counterstaining on PFA-fixed section of day12 melanoma with

CDH5 antibody. Yellow arrows, CLECs. Scale bar: 50 μm. h, CLECs constitute $2.8 \pm$

0.75% of total endothelial cells in day 12 melanoma (n=4).

Fig. 3: CLECs do not originate from the bone marrow. a-b, Tamoxifen induction

timing. b, Quantification of the population of tumour macrophagens (T_MACs:

CD45+GFP+) and tumour CLECs (T_CLECs: CD45-CD31+GFP+) by flow cytometric

analysis in tumour samples (n=4). Bars represent mean $\pm$ s.d. c, Fluorescent images of

PFA-fixed 3 weeks CT2A glioma de of Csf1r-Mer2.CremTmG mice within 24 hours after

tamoxifen induction. Scale bar: 100 μm. d-e, Flow cytometric analysis of CT2A-glioma

in bone marrow chimeras. d, BMCsf1r-mer2.CremTmG: C57BL/6J mice are used as

recipients and Csf1r-Mer2.CremTmG mice used as donors. Within the sorted myeloid

cell population (BFP-CD45+), $32.3 \pm 12.1\%$ are GFP-positive in Csf1r-Mer2.CremTmG

mice (n= 10), and $33.8 \pm 6.9\%$ are GFP-positive in bone-marrow transplanted BMCsf1r-

Mer2.CremTmG mice (n= 12). Among endothelial cell population (BFP-CD45-CD31+),

$4.3 \pm 1.9\%$ are GFP-positive in Csf1r-Mer2.CremTmG mice (n= 10), whereas no GFP-

positive endothelial cells are found in BMCsf1r-Mer2.CremTmG mice (n= 12). e, Csf1r-

Mer2.CremTmG:: BMWT: Csf1r-Mer2.CremTmG are used as recipients and C57BL/6J

as donors. Of the sorted myeloid cell population, $43.4 \pm 11.3\%$ GFP-positive cells are

found in Csf1r-Mer2.CremTmG mice (n= 6), whereas $0.1 \pm 0.1\%$ GFP-positive cells are

found in Csf1r-Mer2.CremTmG:: BMWT mice (n= 4). Of the endothelial cell population,

$10.5 \pm 9.7\%$ GFP-positive cells are found in Csf1r-Mer2.CremTmG mice (n= 6), and 9.3

$\pm 3.7\%$ GFP-positive cells are found in Csf1r-Mer2.CremTmG:: BMWT mice (n= 4).

Bars represent mean $\pm$ s.d. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Two-tailed

unpaired Mann-Whitney' s U test. e, Longitudinal intravial imaging of 2 weeks CT2A

glioma of Csf1r-Mer2.CremTmG and Csf1r-Mer2.CremTmG :: BMWT mice. White

arrows, CLECs. Scale bar: 50 μm.

Fig. 4: Dual recombination lineage tracing and inducible depletion model identifies

a role for CLECs in tumour blood vessel patterning supporting glioma growth. a,

Cdh5-Dre: Csf1r-Mer2.Cre: RSR-LSL-tdTomato (DreMer2.CretdTomato): a genetic

mouse model for CLEC selective tracing by dual recombination. b-c, Analysis in

DreMer2.CretdTomato. b, Longitudinal intravital imaging of the tumour vasculature in 3

weeks CT2A mouse glioma by Intravenous injection of FITC dextran. Yellow

arrowheads, CLECs. Scale bar: 50 μm. c, ERG antibody counterstaining on PFA-fixed

sections of 3 weeks CT2A glioma. Yellow arrows, CLECs. Scale bar: 50 μm. d, Cdh5-

Dre: Csf1r-Mer2.Cre: RSR-mTmG-DTR (DreMer2.CremTmG-DTR): a genetic mouse

model for CLEC selective tracing and inducible ablation. DTR, diphtheria toxin receptor.

e-m, Analysis in DreMer2.CremTmG-DTR. e, Longitudinal intravital imaging of the

tumour vasculature in 3 weeks CT2A mouse glioma. Mer2.CremTmG, Csf1r-

Mer2.CremTmG. Yellow arrowheads, CLECs. f, Timing schedule of glioma implantation,

tamoxifen induction and diphtheria toxin (DT) treatment. g, Gross appearance of 3 weeks

CT2A gliomas. Scale bar: 5 mm. h, Confocal images of PFA-fixed 3 weeks CT2A glioma

sections. Yellow arrows, CLECs. Scale bar: 50 μm. i, Tumour weight at 3 weeks CT2A

glioma. j-l, Analysis of PFA-fixed 3 weeks CT2A glioma sections. j, Tumour blood

vessels area. k, Network length of tumour blood vessels. l, Number of bifurcations of

tumour blood vessels. Bars represent mean ± s.d. *p<0.05, **p<0.01. Two-tailed unpaired

Mann-Whitney' s U test.

Fig. 5: Single-cell analysis identifies a unique signature that defines the tumour

CLECs. a-d, Single-cell RNA sequencing (scRNA-seq) analysis of tumour macrophages

(T_MACs: CD45+GFP+, 79 cells), tumour endothelial cell (T_ECs: CD45-CD31+GFP-,

78 cells), tumour CLECs (T_CLECs: CD45-CD31+GFP+, 141 cells) in 4 weeks CT2A

glioma of Csf1r-Mer2.CremTmG mice. b, Gene expression of typical endothelial cell

markers, macrophages and myeloid cell markers. c, Csf1r gene expression. d, Three

characteristic genes expressed in T_CLECs. e, Enrichment of Tip-cell signature in

T_CLECs compared to T_ECs as shown by GSEA. f-k, Analysis of scRNA-seq data of

human endothelial cells from multiple primary tumours. f, UMAP plot of human

endothelial cell subtypes. g, UMAP plot of human endothelial cells colour-coded for the

tumour type of origins. h, UMAP plots with marker gene expression for each EC subtype.

i, Heatmap of marker gene expression for each EC phenotype (each cell as a column and

each gene as a row). j-k, AUCell analysis of 24 T_CLECs markers. j, T_CLECs

subpopulation indicated (AUC>0.165) by red colored dots. k, Cell type-specific (left) and

tumour type-specific (right) populations of T_CLECs. BCC, basal cell carninoma: PDAC,

pancreatic ductal adenocarcinoma.

Fig. 6: CLECs are linked to Prox1-positive lymphatic endothelium of the

subcapsular sinus floor in lymph nodes. a, Fluorescent images of the indicated organs

of Csf1r-Mer2.CremTmG mice after multiple tamoxifen induction. Scale bar: 50 μm. b,

Quantification of the population of CLECs in Csf1r-Mer2.CremTmG mice by flow

cytometric analysis. Of the endothelial cell population (CD45-CD31+), 9.8 ± 2.1% GFP-

positive cells are found in the axillary lymph node of Csf1r-Mer2.CremTmG mice (n=

10). c, Counterstaining on PFA-fixed Inguinal lymph node of Csf1r-Mer2.CremTmG::

BMWT mice using PROX1 antibody within 24 hours after tamoxifen induction. Yellow

arrowheads, GFP+tdTomato+PROX1+ cells in the floor of the subcapsular sinus. Scale

bar 25 μm. d, Fluorescent images of PROX1 antibody counterstaining on PFA-fixed

inguinal lymph node of Csf1r-Mer2.CremTmG::BMWT mice after multiple rounds of

tamoxifen induction. Scale bar: 100 μm. Detailed image of square outlined in d. Yellow

arrowheads, GFP+Prox1+ cells; blue arrowheads, GFP+Prox1- cells. h-k, Analysis of

CT2A glioma in Prox1-CreERT2mTmG mice.

Fig. 7: CLEC and EC heterogeneity in tumor and lymph node. a, Quality measures

of 4 SMARTseq2 scRNA-seq. libraries: Csf1r-lineage endothelial cells (CLECs) and

endothelial cells (ECs) isolated from tumor (T_) and lymph node (LN_). b, Violin plots

show GFP expression per single cell. c, Uniform manifold approximation and projection

(UMAP) of T_CLECs, T_ECs and LN_CLECs, LN_ECs. d, Heatmap shows the 20 most

characteristic markers per cell population. e, UMAP colored by gene expression diversity

(CytoTRACE score) and predicted ordering by CytoTRACE. f, Monocle based trajectory

analysis of four cell populations, colored by pseudotime and CytoTRACE score.

Extended Data Fig. 1: CLECs in mouse glioma express endothelial cell markers, but

lack macrophage and myeloid cell markers.   Counterstaining on PFA-fixed 4 weeks

CT2A glioma sections of Csf1r-Mer2.CremTmG with indicated antibodies. Lower panels

show the detail images of outlined squares in upper panels. Yellow arrowheads, CLECs;

blue arrowheads, macrophages. Scale bar: 25 μm.

Extended Data Fig. 2: CLECs contribute to lymphatic endothelium in the tumour

environment. Fluorescent images of counterstaining on PFA-fixed section of day 12

B16F1 melanoma under ear skin of Csf1r-Mer2.CremTmG mice using LYVE1 antibody.

Lower panel indicates detail image of yellow outlined square in upper panel. Yellow

arrow heads, CLECs. Scale bar: 100 μm.

Extended Data Fig. 3: Reconstitution rate in bone marrow chimeras. Reconstitution

rate was determined in peripheral blood 8 weeks after bone marrow transplantation.

TdTomato-positive cells in blood of: C57BL/6J (0.3 ± 0.1%, n= 6), Csf1r-

Mer2.CremTmG (99.2 ± 0.5%, n= 7), BMCsf1r-Mer2.CremTmG (97.3 ± 1.6%, n= 13).

Bars represent mean ± s.d. ****$p<0.0001$. Two-tailed unpaired Mann-Whitney' s U test.

Extended Data Fig. 4: CLECs do not originate from spleen. Surgical removal of

spleen (splenectomy) was performed 3 weeks before glioma injection. CLECs isolated

from CT2A glioma of sham control Csf1r-Mer2.CremTmG mice (n= 5) constitute 1.5 ±

0.8 % of total tumour endothelial cell population, compared to 1.5 ± 0.5% found in Csf1r-

Mer2.CremTmG with splenectomy (n= 4). Bars represent mean ± s.d. Two-tailed

unpaired Mann-Whitney' s U test.

Extended Data Fig. 5: CLECs locate at peripheral area of mouse glioma. a,

Vibratome section (200 μm-thick) of 4 weeks CT2A glioma of Csf1r-Mer2.CremTmG::

BMWT mice. Scale bar: 1 mm. b, Fluorescent images of area outlined in a. Yellow

arrowheads, CLECs. Scale bar: 100 μm.

Extended Data Fig. 6: 24 tumour CLECs markers in mouse glioma. SCDE analysis of scRNA-seq data of tumour macrophages (T_MACs: CD45+GFP+, n= 79), tumour endothelial cells (T_ECs: CD45-CD31+GFP-, n= 78), tumour CLECs (T_CLECs: CD45-CD31+GFP+, n= 141) in 4 weeks CT2A glioma in Csf1r-Mer2.CremTmG mice. Bars represent mean ± s.d. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001. Two-tailed unpaired t-test with Welch correction.

Extended Data Fig. 7: CLECs in glioma express AQP1. Counterstaining on PFA-fixed CT2A glioma of Csf1r-Mer2.CremTmG::BMWT mice using AQP1 antibody. Yellow arrowheads, T_CLECs; blue arrowheads, T_ECs. Scale bar: 25 μm.

Extended Data Fig. 8: CLECs in glioma cannot be labelled by GFP within 24 hours after tamoxifen induction. a-b, Tamoxifen induction timing. b, Quantification of the population of tumour macrophagens (T_MACs: CD45+GFP+) and tumour CLECs (T_CLECs: CD45-CD31+GFP+) by flow cytometric analysis in tumour samples (n=4). Bars represent mean ± s.d. c, Fluorescent images of PFA-fixed 3 weeks CT2A glioma de of Csf1r-Mer2.CremTmG mice within 24 hours after tamoxifen induction. Scale bar: 100 μm.

Extended Data Fig. 9: CLEC and EC heterogeneity in tumor and lymph node .
a, Heatmap shows the expression of n=91 fLECs markers in LN_CLEC, LN_EC, T_CLEC and T_EC single cells. b, Violin plots show the expression of 3 CLEC markers: Pde4d, Robo1 and Itga9.

Supplemental Movie 1: Time-lapse *in vivo* imaging of Csf1r lineage endothelial tip cell in CT2A glioma of *Csf1r-Mer2.Cre^{mTmG}* mice. Imaging was performed over 4 hours (1 frame/ 10 min).

Supplemental Movie 2: Time-lapse *in vivo* imaging of *Csf1r* lineage endothelial tip cell in CT2A glioma of *Csf1r-Mer2.Cre^{mTmG}*::BM^{WT}. Imaging was performed over 50 minutes (1 frame/ 2.5 min).

Supplemental Table 1:   Details of gene expression analysis. Seurat based differential analysis (non-parameteric Wilcoxon rank sum test) was performed by comparing cell types amongst each other (green columns) and the different Seurat clusters (yellow columns). p_val: p_val (unadjusted). avg_logFC: log fold-change of the average expression between the two groups. Positive values indicate that the feature is more highly expressed in the first group. pct.1: The percentage of cells where the feature is detected in the first group. pct.2: The percentage of cells where the feature is detected in the second group. p_val_adj: Adjusted p-value, based on Bonferroni correction using all features in the dataset.

Supplemental Table 2:   Marker genes for EC subtypes of multiple tumour types.
p_val: p_val (unadjusted). avg_logFC: Average log (fold change). pct.1: Percentage of expressed cells in current cluster. pct.2: Percentage of expressed cells in other cluster. p_val_adj: Adjusted p-value. cluster: Cluster name. gene: Gene name. log2fc: log2 (fold change). Log2pct: log2 ((pct+0.005)/(pct2+0.005)).

Fig. 1: ***Csf1r*** **lineage endothelial cells (CLECs) contribute to perfused, dynamically sprouting and remodeling blood vessels in mouse glioma. a,** *Csf1r-Mer2.Cre*[mTmG]: a mouse model for *Csf1r*-lineage tracing combining the tamoxifen-inducible Cre-driver under control of the *Csf1r*-promoter and the *mTmG* Cre-reporter. **b,** Timing schedule of glioma implantation, tamoxifen induction and in vivo imaging. **c-f,** Longitudinal intravital imaging of the tumour vasculature in CT2A mouse glioma of *Csf1r-Mer2.Cre*[mTmG] by cranial window. Scale bar: 50 μm. **c,** In vivo imaging in 3 weeks glioma. Endothelial cells in red (tdTomato), *Csf1r*-labeled cells in green (GFP) and glioma cells in blue (BFP). **d,** Detail image of outlined square in Fig. 1c illustrating tracing of GFP-positive cells. White arrowheads, macrophages; blue arrowhead, CLECs at tip position. **e,** In vivo imaging of CLECs in 2 weeks glioma. Blue arrowheads, tip of endothelial cells. **f,** Visualization of blood vessels in 3 weeks glioma by Intravenous injection of dextran with texas red dye. Yellow arrowhead, bridging CLECs; orange arrowhead, tube formation by CLECs; blue arrowhead, CLECs at tip position. **g-h,** Counterstaining on PFA-fixed 4 weeks CT2A glioma sections of *Csf1r-Mer2.Cre*[mTmG] with indicated antibodies. Yellow arrows, CLECs. Scale bar: 25 μm.

Matsumoto K et al., Figure1

Fig. 2: Up to 10% of intra-tumour endothelium is comprised of CLECs not only in glioma, but also in other tumour types and locations. a-f, Quantification of the population of CLECs in glioma of Csf1r-Mer2.Cre^mTmG mice. a, Gross appearance of sham control and 4 weeks CT2A glioma (Upper panel). Purple arrowhead, the area of capillary injection; blue arrowhead, glioma. Scale bar: 5 mm. Flow cytometric analysis of CLECs in sham control and 4 weeks CT2A glioma (Lower panel). b, In 3 weeks CT2A glioma (n= 9), CLECs constitute 2.1 ± 1.0% of total endothelial cells (CD45-CD31+); in 4 weeks CT2A glioma (n= 5), they make up 7.5 ± 1.6% of total endothelial cells. c, In 3 weeks GL261 glioma (n=5), CLECs constitute 1.5 ± 1.6% of total endothelial cells. d, Gross appearance of normal brain, sham control and 3 weeks CT2A glioma. Purple arrowhead, area of capillary injection. Scale bar: 5 mm. e-f, Quantification of the population of CLECs by flow cytometric analysis in brain and tumour samples. Bars represent mean ± s.d. *P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. Two-tailed unpaired Mann-Whitney's U test. g-h, Analysis of CLECs in B16F1 melanoma in Csf1r-Mer2.CremTmG mice. g, Counterstaining on PFA-fixed section of day12 melanoma with CDH5 antibody. Yellow arrows, CLECs. Scale bar: 50 μm. h, CLECs constitute 2.8 ± 0.75% of total endothelial cells in day 12 melanoma (n=4).

Matsumoto K et al., Figure 2

Fig. 3: CLECs do not originate from bone marrow. a-b, Flow cytometric analysis of CT2A-glioma in bone marrow chimeras.

a, BM*Csf1r-mer2.CremTmG*: C57BL/6J mice are used as recipients and *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ mice used as donors. Within the sorted myeloid cell population (BFP⁻CD45⁺), 32.3 ± 12.1% are GFP-positive in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ mice (n= 10), and 33.8 ± 6.9% are GFP-positive in bone-marrow transplanted BM*Csf1r-Mer2.CremTmG* mice (n= 12). Among endothelial cell population (BFP⁻CD45⁻CD31⁺), 4.3 ± 1.9% are GFP-positive in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ mice (n= 10), whereas no GFP-positive endothelial cells are found in BM*Csf1r-Mer2.CremTmG* mice (n= 12). b, *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ:: BMᵂᵀ: *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ are used as recipients and C57BL/6J as donors. Of the sorted myeloid cell population, 43.4 ± 11.3% GFP-positive cells are found in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ mice (n= 6), whereas 0.1 ± 0.1% GFP-positive cells are found in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ:: BMᵂᵀ mice (n= 4). Of the endothelial cell population, 10.5 ± 9.7% GFP-positive cells are found in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ mice (n= 6), and 9.3 ± 3.7% GFP-positive cells are found in *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ:: BMᵂᵀ mice (n= 4). Bars represent mean ± s.d. *P<0.05,**P<0.01,***P<0.001, ****P<0.0001. Two-tailed unpaired Mann-Whitney' s U test. c, Longitudinal intravial imaging of 2 weeks CT2A glioma of *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ and *Csf1r-Mer2.Cre*ᵐᵀᵐᴳ :: BMᵂᵀ mice. White arrows, CLECs. Scale bar: 50 μm.

Matsumoto K et al., Figure 3

Fig. 4: Dual recombination lineage tracing and inducible depletion model identifies a role for CLECs in tumour blood vessel patterning supporting glioma growth. a, *Cdh5-Dre*: *Csf1r-Mer2.Cre*: *RSR-LSL-tdTomato* (*DreMer2.Cre^tdTomato*): a genetic mouse model for CLEC selective tracing by dual recombination. b-c, Analysis in *DreMer2.Cre^tdTomato*. b, Longitudinal intravital imaging of the tumour vasculature in 3 weeks CT2A mouse glioma by Intravenous injection of FITC dextran. Yellow arrowheads, CLECs. Scale bar: 50 *μ*m. c, ERG antibody counterstaining on PFA-fixed sections of 3 weeks CT2A glioma. Yellow arrows, CLECs. Scale bar: 50 *μ*m. d, *Cdh5-Dre*: *Csf1r-Mer2.Cre*: *RSR-mTmG-DTR* (*DreMer2.Cre^mTmG-DTR*): a genetic mouse model for CLEC selective tracing and inducible ablation. DTR, diphtheria toxin receptor. e-l, Analysis in *DreMer2.Cre^mTmG-DTR*. e, Longitudinal intravital imaging of the tumour vasculature in 3 weeks CT2A mouse glioma. *Mer2.Cre^mTmG*, *Csf1r-Mer2.Cre^mTmG*. Yellow arrowheads, CLECs. f, Timing schedule of glioma implantation, tamoxifen induction and diphtheria toxin (DT) treatment. g, Gross appearance of 3 weeks CT2A gliomas. Scale bar: 5 mm. h, Confocal images of PFA-fixed 3 weeks CT2A glioma sections. Yellow arrows, CLECs. Scale bar: 50 *μ*m. i, Tumour weight at 3 weeks CT2A glioma. j-l, Analysis of PFA-fixed 3 weeks CT2A glioma sections. j, Tumour blood vessels area. k, Network length of tumour blood vessels. l, Number of bifurcations of tumour blood vessels. Bars represent mean ± s.d. *P<0.05, **P<0.01. Two-tailed unpaired Mann-Whitney's U test.

Matsumoto K et al., Figure 4

Fig. 5: Single-cell analysis identifies a unique signature that defines the tumour CLECs. a-d, Single-cell RNA sequencing (scRNA-seq) analysis of tumour macrophages (T_MACs: CD45+GFP+, 79 cells), tumour endothelial cell (T_ECs: CD45-CD31+GFP-, 78 cells), tumour CLECs (T_CLECs: CD45-CD31+GFP+, 141 cells) in 4 weeks CT2A glioma of *Csf1r-Mer2.Cre*[mTmG] mice. b, Gene expression of typical endothelial cell markers, macrophages and myeloid cell markers. c, *Csf1r* gene expression. d, Three characteristic genes expressed in T_CLECs. e, Enrichment of Tip-cell and lymphatic signature in T_CLECs compared to T_ECs as shown by GSEA. f-k, Analysis of scRNA-seq data of human endothelial cells from multiple primary tumours. f, UMAP plot of human endothelial cell subtypes. g, UMAP plot of human endothelial cells colour-coded for the tumour type of origins. h, UMAP plots with marker gene expression for each EC subtype. i, Heatmap of marker gene expression for each EC phenotype (each cell as a column and each gene as a row). j-k, AUCell analysis of 24 T_CLECs markers. j, T_CLECs subpopulation indicated (AUC>0.165) by red colored dots. k, Cell type-specific (left) and tumour type-specific (right) populations of T_CLECs. BCC, basal cell carninoma: PDAC, pancreatic ductal adenocarcinoma.

Matsumoto K et al., Figure 5

Fig. 6: CLECs are linked to *Prox1*-positive lymphatic endothelium of the subcapsular sinus floor in lymph nodes. a, Fluorescent images of the indicated organs of *Csf1r-Mer2.Cre^mTmG* mice after multiple tamoxifen induction. Scale bar: 50 *μ*m. b, Quantification of the population of CLECs in *Csf1r-Mer2.Cre^mTmG* mice by flow cytmetric analysis. Of the endothelial cell population (CD45⁻CD31⁺), 9.8 ± 2.1% GFP-positive cells are found in the axillary lymph node of *Csf1r-Mer2.Cre^mTmG* mice (n= 10). c, Counterstaining on PFA-fixed Inguinal lymph node of *Csf1r-Mer2.Cre^mTmG*:: BM^WT mice using PROX1 antibody within 24 hours after tamoxifen induction. Yellow arrowheads, GFP⁺tdTomato⁺PROX1⁺ cells in the floor of the subcapsular sinus. Scale bar 25 *μ*m. d, Fluorescent images of PROX1 antibody counterstaining on PFA-fixed inguinal lymph node of *Csf1r-Mer2.Cre^mTmG*::BM^WT mice after multiple rounds of tamoxifen induction. Scale bar: 100 *μ*m. Detailed image of square outlined in d. Yellow arrowheads, GFP⁺Prox1⁺ cells; blue arrowheads, GFP⁺Prox1⁻ cells.

Fig. 7: CLEC and EC heterogeneity in tumor and lymph node .

a, Quality measures of 4 SMARTseq2 scRNA-seq. libraries: Csf1r-lineage endothelial cells (CLECs) and endothelial cells (ECs) isolated from tumor (T_) and lymph node (LN_). b, Violin plots show GFP expression per single cell. c, Uniform manifold approximation and projection (UMAP) of T_CLECs, T_ECs and LN_CLECs, LN_ECs. d, Heatmap shows the 20 most characteristic markers per cell population. e, UMAP colored by gene expression diversity (CytoTRACE score) and predicted ordering by CytoTRACE. f, Monocle based trajectory analysis of four cell populations, colored by pseudotime and CytoTRACE score.

Matsumoto K et al., Figure 7